



HAL
open science

Cauchy Problem for the Kuznetsov Equation

Adrien Dekkers, Anna Rozanova-Pierrat

► **To cite this version:**

Adrien Dekkers, Anna Rozanova-Pierrat. Cauchy Problem for the Kuznetsov Equation. *Discrete and Continuous Dynamical Systems - Series A*, In press, 39 (1). hal-01648010v1

HAL Id: hal-01648010

<https://hal.science/hal-01648010v1>

Submitted on 24 Nov 2017 (v1), last revised 8 Oct 2018 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Cauchy Problem for the Kuznetsov Equation

Adrien Dekkers and Anna Rozanova-Pierrat*

November 24, 2017

Abstract

We consider the Cauchy problem for a model of non-linear acoustics, named the Kuznetsov equation, describing sound propagation in thermo-viscous elastic media. For the viscous case, it is a weakly quasi-linear strongly damped wave equation, for which we prove the global existence in time of regular solutions for sufficiently small initial data, the size of which is specified, and give the corresponding energy estimates. In the non-viscous case, we update the known results of John for quasi-linear wave equations, obtaining the well-posedness results for less regular initial data. We obtain, using a priori estimates and a Klainerman inequality, the estimations of the maximal existence time, depending on the space dimension, which are optimal, thanks to the blow-up results of Alinhac. Alinhac's blow-up results are also confirmed by a L^2 -stability estimate, obtained between a regular and a less regular solutions.

1 Introduction

The Kuznetsov equation [16] models a propagation of non-linear acoustical waves in thermo-viscous elastic media. This equation describes the evolution of the velocity potential and can be derived, as in [18], from a compressible isentropic Navier-Stokes system, for the viscous case, or the Euler system for the non-viscous case, using small perturbations of the density and of the velocity characterized by a small dimensionless parameter $\varepsilon > 0$. The Cauchy problem for the Kuznetsov equation reads for $\alpha = \frac{\gamma-1}{c^2}$, $\beta = 2$ and $\nu = \frac{\delta}{\rho_0}$ as

$$u_{tt} - c^2 \Delta u - \nu \varepsilon \Delta u_t = \alpha \varepsilon u_t u_{tt} + \beta \varepsilon \nabla u \cdot \nabla u_t, \quad x \in \mathbb{R}^n, \quad (1)$$

$$u(x, 0) = u_0(x), \quad u_t(x, 0) = u_1(x), \quad x \in \mathbb{R}^n, \quad (2)$$

where c , ρ_0 , γ , δ are the velocity of the sound, the density, the ratio of the specific heats and the viscosity of the medium respectively. In what follows, we can just suppose that α and β are some positive constants. Eq. (1) is a weakly quasi-linear damped

*Laboratory Mathématiques et Informatique pour la Complexité et les Systèmes, CentraleSupélec, Université Paris-Saclay, Campus de Gif-sur-Yvette, Plateau de Moulon, 3 rue Joliot Curie, 91190 Gif-sur-Yvette, France

wave equation, that describes a propagation of a high amplitude wave in fluids. The Kuznetsov equation is one of the models derived from the Navier-Stokes system, and it is well suited for the plane, cylindrical and spherical waves in a fluid [7]. Most of the works on the Kuznetsov equation (1) are treated in the one space dimension [11] or in a bounded spatial domain of \mathbb{R}^n [13, 12, 17]. For the viscous case Kaltenbacher and Lasiecka [13] have considered the Dirichlet boundary valued problem and proved for sufficiently small initial data the global well-posedness for $n \leq 3$. Meyer and Wilke [17] have proved it for all n . In [12] it was proven a local well-posedness of the Neumann boundary valued problem for $n \leq 3$.

In this article we study the well-posedness properties of the Cauchy problem (1)–(2). In the non-viscous case for $\nu = 0$, the Cauchy problem for the Kuznetsov equation is a particular case of a general quasi-linear hyperbolic system of the second order considered by Hughes, Kato and Marsden [8] (see Theorem 1 Points 1 and 2 for the application of their results to the Kuznetsov equation). The local well-posedness result, proved in [8], does not use a priori estimate techniques and is based on the semi-group theory. Hence, thanks to [8], we have the well-posedness of (1)–(2) in the Sobolev spaces H^s with a real $s > \frac{n}{2} + 1$. Therefore, actually, to extend the local well-posedness to a global one (for $n \geq 4$) and to estimate the maximal time interval on which there exists a regular solution, John [10] has developed a priori estimates for the Cauchy problem for a general quasi-linear wave equation. This time, due to the non-linearities $u_t u_{tt}$ and $\nabla u \nabla u_t$, including the time derivatives, to have an a priori estimate for the Kuznetsov equation we need to work with Sobolev spaces with a natural s , thus denoted in what follows by m . If we directly apply general results of Ref. [10] to our case of the Kuznetsov equation, we obtain a well-posedness result with a high regularity of the initial data. We improve it in Theorem 3 and show John's results for the Kuznetsov equation with the minimal regularity on the initial data corresponding to the regularity obtained by Hughes, Kato and Marsden [8]. For instance, we prove the analogous energy estimates in H^m with $m \geq [\frac{n}{2} + 2]$ instead of John's $m \geq \frac{3}{2}n + 4$ (see Eq. (20) in Proposition 1) and its slight modified version in H^m with $m \geq [\frac{n}{2} + 3]$ instead of $m \geq \frac{3}{2}n + 7$ (see Eq. (36) in Proposition 2). The energy estimates allow us to evaluate the maximal existence time interval (see Theorem 1 Point 5 and Theorem 4 for more details). In \mathbb{R}^2 and \mathbb{R}^3 the optimality of obtained estimations for the maximal existence time is ensured by the results of Alinhac [2]. In Ref. [2] a geometric blow-up for small data is proved for $\partial_t^2 u$ and Δu at a finite time of the same order as predicted by our a priori estimates (see Theorem 1). From the other hand, the blow-up of $\partial_t^2 u$ and Δu is also confirmed by the stability estimate (8) in Theorem 1: if the maximal existence time interval is finite and limited by T^* , by Eq. (8), we have the divergence

$$\int_0^{T^*} (\|u_{tt}\|_{L^\infty(\mathbb{R}^n)} + \|\Delta u\|_{L^\infty(\mathbb{R}^n)}) ds = +\infty. \quad (3)$$

For $n \geq 4$ and $\nu = 0$, we also improve the results of John [10] and show the global existence for sufficiently small initial data $u_0 \in H^{m+1}(\mathbb{R}^n)$ and $u_1 \in H^m(\mathbb{R}^n)$ with $m \geq n + 2$ instead of $m \geq \frac{3}{2}n + 7$ (see Proposition 4 and Theorem 4). The smallness of the initial data here directly ensures the hyperbolicity of the Kuznetsov equation for all time, *i.e.* it ensures that $1 - \alpha \varepsilon u_t$ is strictly positive and bounded for all time. The proof

uses the generalized derivatives for the wave type equations [10] and a priori estimate of Klainerman [14, 15] (see Section 3.2).

Let us now formulate our main well-posedness result for the non-viscous case:

Theorem 1 (Non-viscous case) *Let $\nu = 0$, $n \in \mathbb{N}^*$ and $s > \frac{n}{2} + 1$. For all $u_0 \in H^{s+1}(\mathbb{R}^n)$ and $u_1 \in H^s(\mathbb{R}^n)$ such that $\|u_1\|_{L^\infty(\mathbb{R}^n)} < \frac{1}{2\alpha\varepsilon}$, the following results hold:*

1. *There exists $T > 0$ such that there exists a unique solution u of (1)–(2) with the following regularity*

$$u \in C^r([0, T]; H^{s+1-r}(\mathbb{R}^n)) \text{ for } 0 \leq r \leq s, \quad (4)$$

$$\forall t \in [0, T], \quad \|u_t(t)\|_{L^\infty(\mathbb{R}^n)} < \frac{1}{2\alpha\varepsilon}. \quad (5)$$

2. *The map $(u_0, u_1) \mapsto (u(t, \cdot), \partial_t u(t, \cdot))$ is continuous in the topology of $H^{s+1} \times H^s$ uniformly in $t \in [0, T]$.*
3. *Let T^* be the largest time on which such a solution is defined, and in addition $s \in \mathbb{N}$, i.e. $s = m \geq m_0 = \lceil \frac{n}{2} + 2 \rceil$. With the notation*

$$E_m[u](t) = \|\nabla u(t)\|_{H^m}^2 + \sum_{i=1}^{m+1} \|\partial_t^i u(t)\|_{H^{m+1-i}}^2, \quad (6)$$

there exist constants $C(n, c, \alpha) > 0$ and $\hat{C}(n, c, \alpha, \beta) > 0$ (see Theorem 3) such that if the initial data satisfies $\sqrt{E_{m_0}[u]}(0) \leq \frac{1}{C(n, c, \alpha)\varepsilon}$, then

$$T^* \geq \frac{1}{\varepsilon \hat{C}(n, c, \alpha, \beta) \sqrt{E_{m_0}[u]}(0)}, \text{ such that it holds (3)}. \quad (7)$$

4. *For two solutions u and v of the Kuznetsov equation for $\nu = 0$ defined on $[0, T^*[$ assume that u be regular as in (4)–(5), i.e. $u \in L^\infty([0, T^*]; H^{m+1}(\mathbb{R}^n))$, $u_t \in L^\infty([0, T^*]; H^m(\mathbb{R}^n))$ ($s = m$ as in Point 3), and*

$$v \in L^\infty([0, T^*]; H^1(\mathbb{R}^n)), \quad v_t \in L^\infty([0, T^*]; L^2(\mathbb{R}^n)) \text{ with } \|v\|_{L^\infty(\mathbb{R}^n)} < \frac{1}{2\alpha\varepsilon}$$

and with a bounded $\|\nabla v_t\|_{L^\infty(\mathbb{R}^n)}$ norm on $[0, T^[$. Then it holds the following stability uniqueness result: there exist constants $C_1 > 0$ and $C_2 > 0$, independent on time, such that*

$$\begin{aligned} (\|(u-v)_t\|_{L^2}^2 + \|\nabla(u-v)\|_{L^2}^2)(t) &\leq C_1 \exp(C_2\varepsilon \int_0^t \sup(\|u_{tt}\|_{L^\infty(\mathbb{R}^n)}, \|\Delta u\|_{L^\infty(\mathbb{R}^n)}) ds) \\ &\cdot (\|u_1 - v_1\|_{L^2}^2 + \|\nabla(u_0 - v_0)\|_{L^2}^2). \end{aligned} \quad (8)$$

5. *If $s = m \geq n + 2$, then for sufficiently small initial data (see Theorem 4 in Section 3.2)*

- (a) $\liminf_{\varepsilon \rightarrow 0} \varepsilon^2 T^* > 0$ for $n = 2$,
- (b) $\liminf_{\varepsilon \rightarrow 0} \varepsilon \log(T^*) > 0$ for $n = 3$,
- (c) $T^* = +\infty$ for $n \geq 4$.

Theorem 1 is principally based on the a priori estimates proved in Sections 3.1 (for Point 3) and 3.2 (for Point 5) and on the local existence result updated from Ref. [8] (Points 1 and 2). Point 4 is proved in Section 3.3. Analyzing the structure of the Kuznetsov equation and the difficulties involving by its non-linear terms, we start in Section 2 with preliminary remarks on the L^2 -energy properties for the Kuznetsov equation to compare with its simplified versions. Developing the energy estimates in the Sobolev spaces, we however recognize the structure of the L^2 -energy of the wave equation which keeps unchanged.

In the presence of the term Δu_t for the viscous case $\nu > 0$, the regularity of the higher order time derivatives of u is different (to compare to the non-viscous case), and the way to control the non-linearities in the a priori estimates becomes different. As it was shown in [19], this dissipative term changes a finite speed of propagation of the wave equation to the infinite one. Indeed, the linear part of Eq. (1) can be viewed as two compositions of the heat operator $\partial_t - \Delta$ in the following way:

$$u_{tt} - c^2 \Delta u - \nu \varepsilon \Delta u_t = \partial_t(\partial_t u - \varepsilon \nu \Delta u) - c^2 \Delta u.$$

For the viscous case we prove the global in time well-posedness results in \mathbb{R}^n (see Section 4) for small enough initial data, the size of which we specify (see Point 1 of Theorem 2 and Subsection 4.1 for its proof). In Subsection 4.2 for $n \geq 3$ (see Point 2 of Theorem 2) we establish an a priori estimate which gives also a sufficient condition of the existence of a global solution for a sufficiently small initial energy of the same order on ε as in Point 1 of Theorem 2. The same results (see Point 3 of Theorem 2) hold in $(\mathbb{R}/L\mathbb{Z}) \times \mathbb{R}^{n-1}$ for $n \geq 2$ (with a periodicity and mean value zero on one variable).

Theorem 2 (Viscous case) *Let $\nu > 0$, $n \in \mathbb{N}^*$, $s > \frac{n}{2}$ and $\mathbb{R}_t^+ = [0, +\infty[$. Considering the Cauchy problem for the Kuznetsov equation (1)–(2), the following results hold:*

1. *Let*

$$X := H^2(\mathbb{R}_t^+; H^s(\mathbb{R}^n)) \cap H^1(\mathbb{R}_t^+; H^{s+2}(\mathbb{R}^n)),$$

the initial data

$$u_0 \in H^{s+2}(\mathbb{R}^n) \quad \text{and} \quad u_1 \in H^{s+1}(\mathbb{R}^n),$$

$r_ = O(1)$ be the positive constant defined in Eq. (53) and $C_1 = O(1)$ be the minimal constant such that the solution u^* of the corresponding linear Cauchy problem (50) satisfies*

$$\|u^*\|_X \leq \frac{C_1}{\sqrt{\nu \varepsilon}} (\|u_0\|_{H^{s+2}} + \|u_1\|_{H^{s+1}}).$$

Then for all $r \in [0, r_[$ and all initial data satisfying*

$$\|u_0\|_{H^{s+2}} + \|u_1\|_{H^{s+1}} \leq \frac{\sqrt{\nu \varepsilon}}{C_1} r, \tag{9}$$

there exists the unique solution $u \in X$ of the Cauchy problem for the Kuznetsov equation and $\|u\|_X \leq 2r$.

2. Let $n \geq 3$, $s = m \in \mathbb{N}$ be even and $m \geq [\frac{n}{2} + 3]$. With the notation

$$E_{\frac{m}{2}}[u](t) = \|\nabla u(t)\|_{H^m}^2 + \sum_{i=1}^{\frac{m}{2}+1} \|\partial_t^i u(t)\|_{H^{m-2(i-1)}}^2, \quad (10)$$

there exists a constant $\rho = O(1) > 0$ (see Theorem 7 Point 2), independent on time, such that for all initial data $u_0 \in H^{m+1}(\mathbb{R}^n)$ and $u_1 \in H^m(\mathbb{R}^n)$ satisfying

$$E_{\frac{m}{2}}[u](0) < \rho\varepsilon, \quad (11)$$

there exists a unique $u \in C^0(\mathbb{R}_t^+; H^{m+1}(\mathbb{R}^n)) \cap C^i(\mathbb{R}_t^+; H^{m+2-2i}(\mathbb{R}^n))$, for $i = 1, \dots, \frac{m}{2} + 1$ with the bounded energy

$$\forall t \in \mathbb{R}^+, \quad E_{\frac{m}{2}}[u](t) \leq C(c) E_{\frac{m}{2}}[u](0) < +\infty.$$

Here the constant $C(c) > 0$ is independent on time, on u and on m .

3. For $n \in \mathbb{N}^*$ in $\Omega = (\mathbb{R}/L\mathbb{Z}) \times \mathbb{R}^{n-1}$ with $s = m \in \mathbb{N}$ even and $m \geq [\frac{n}{2} + 3]$ there hold Points 1 and 2 in the class of periodic in one direction functions with the mean value zero

$$\int_{\mathbb{R}/L\mathbb{Z}} u(t, x, y) dx = 0. \quad (12)$$

Let us notice that the hyperbolicity condition (5) is also satisfied if we require conditions (9) and (11). For $\nu > 0$ Point 4 of Theorem 1 obviously holds for all $n \in \mathbb{N}^*$. Point 1 of Theorem 2 is proved in Subsection 4.1 using a theorem of a non-linear analysis [20] (see Theorem 6) and regularity results for the strongly damped wave equation following [6], which can also be used for $\Omega = (\mathbb{R}/L\mathbb{Z}) \times \mathbb{R}^{n-1}$ in point 3. Point 2 of Theorem 2 is proved in Subsection 4.2, using a priori estimates given in Proposition 1, see also Theorem 7. The last point of Theorem 2 is a direct corollary of the Poincaré inequality

$$\|u\|_{L^2((\mathbb{R}/L\mathbb{Z}) \times \mathbb{R}^{n-1})} \leq C \|\partial_x u\|_{L^2((\mathbb{R}/L\mathbb{Z}) \times \mathbb{R}^{n-1})}, \quad (13)$$

which holds in the class of periodic functions with the mean value zero. Estimate (13) allows to have the same estimate as in Proposition 1 (see Section 4) for $n = 2$, which fails in \mathbb{R}^2 . Thus, it also gives the global existence for rather small initial data detailed in Point 2.

2 Preliminary remarks on L_2 -energies

We can notice that Eq. (1) is a wave equation containing a dissipative term Δu_t and two non-linear terms: $\nabla u \nabla u_t$ describing local non-linear effects and $u_t u_{tt}$ describing

global or cumulative effects. Actually, the linear wave equation appears from Eq. (1) if we consider only the terms of the zero order on ε :

$$u_{tt} - c^2 \Delta u = 0. \quad (14)$$

The semi-group theory permits in the usual way to show that for $u_0 \in H^1(\mathbb{R}^n)$ and $u_1 \in L^2(\mathbb{R}^n)$ there exists a unique solution of the Cauchy problem (14), (2)

$$u \in C^0(\mathbb{R}_t^+; H^1(\mathbb{R}^n)) \cap C^1(\mathbb{R}_t^+; L^2(\mathbb{R}^n)).$$

So the energy of the wave equation (14)

$$E(t) = \int_{\mathbb{R}^n} [(u_t)^2 + c^2(\nabla u)^2](t, x) dx, \quad (15)$$

is well defined and conserved

$$\frac{d}{dt} E(t) = 0.$$

For $\nu > 0$ and without non-linear terms, the Kuznetsov equation (1) becomes the known strongly damped wave equation:

$$u_{tt} - c^2 \Delta u - \frac{\nu}{\rho_0} \varepsilon \Delta u_t = 0, \quad (16)$$

which is well-posed [9]: for $m \in \mathbb{N}$, $u_0 \in H^{m+1}(\mathbb{R}^n)$ and $u_1 \in H^m(\mathbb{R}^n)$ there exists a unique solution of the Cauchy problem (16), (2)

$$u \in C^0(\mathbb{R}_t^+; H^{m+1}(\mathbb{R}^n)) \cap C^1(\mathbb{R}_t^+; H^m(\mathbb{R}^n)).$$

Multiplying Eq. (16) by u_t in $L^2(\mathbb{R}^n)$, we obtain for the energy of the wave equation (15)

$$\frac{d}{dt} E(t) = -2 \frac{\nu}{\rho_0} \varepsilon \int_{\mathbb{R}^n} (\nabla u_t)^2(t, s) ds \leq 0,$$

what means that the energy $E(t)$ decreases in time, thanks to the viscosity term with $\nu > 0$. The rate of the decreasing is found for more regular energies in [19] in accordance with the regularity of the initial conditions. Without the term $\nabla u \nabla u_t$ (local non-linear effects), the Kuznetsov equation becomes similar to the Westervelt equation, initially derived by Westervelt [21] before Kuznetsov. The Westervelt equation, historically derived [21] for the acoustic pressure fluctuation, has the following form

$$p_{tt} - c^2 \Delta p - \frac{\nu}{\rho_0} \varepsilon \Delta p_t = \frac{\gamma + 1}{c^2} \varepsilon p_t p_{tt}, \quad (17)$$

and can also be seen as an approximation of an isentropic Navier-Stokes system.

In the sequel we conveniently denote p by u . We multiply Eq. (17) by u_t and integrate over \mathbb{R}^n to obtain

$$\frac{1}{2} \frac{d}{dt} \left(\int_{\mathbb{R}^n} [(u_t)^2 + c^2(\nabla u)^2] dx \right) + \frac{\nu}{\rho_0} \varepsilon \int_{\mathbb{R}^n} (\nabla u_t)^2 dx = \frac{1}{3} \frac{\gamma + 1}{c^2} \varepsilon \frac{d}{dt} \left(\int_{\mathbb{R}^n} (u_t)^3 dx \right).$$

Then we have

$$\frac{1}{2} \frac{d}{dt} \left(\int_{\mathbb{R}^n} \left[\left(1 - \frac{2\gamma+1}{3} \frac{\varepsilon u_t}{c^2} \right) (u_t)^2 + c^2 (\nabla u)^2 \right] dx \right) + \frac{\nu}{\rho_0} \varepsilon \int_{\mathbb{R}^n} (\nabla u_t)^2 dx = 0.$$

For $\alpha = \frac{2\gamma+1}{3} \frac{\varepsilon}{c^2}$ we consider the energy

$$E_{west}(t) = \int_{\mathbb{R}^n} [(1 - \alpha \varepsilon u_t) (u_t)^2 + c^2 (\nabla u)^2] dx, \quad (18)$$

which is monotonous decreasing for $\nu > 0$ and is conserved for $\nu = 0$. Let us also notice that, taking the same initial data for different values of ν , energy (18) is a decreasing function of ν :

$$\text{for all } 0 < \nu_1 < \nu_2 \text{ and } t > 0 \quad E_{west}(t, \nu = 0) > E_{west}(t, \nu_1) > E_{west}(t, \nu_2) \geq 0,$$

in the assumption that $1 - \alpha \varepsilon u_t \geq 0$ almost everywhere.

While $\frac{1}{2} \leq 1 - \alpha \varepsilon u_t \leq \frac{3}{2}$, that is to say $\|u_t(t)\|_{L^\infty(\mathbb{R}^n)}$ remains small enough in time, then we can compare E_{west} to the energy of the wave equation

$$\frac{1}{2} E(t) \leq E_{west}(t) \leq \frac{3}{2} E(t).$$

Then a sufficiently regular solution of the Cauchy problem for the Westervelt equation has the energy E controlled by a decreasing in time function:

$$E(t) \leq 3E(0) - 4 \frac{\nu}{\rho_0} \varepsilon \int_0^t \int_{\mathbb{R}^n} (\nabla u_t(s, x))^2 dx ds.$$

Now, let us consider the Kuznetsov equation (1). We multiply it by u_t and integrate on \mathbb{R}^n to obtain

$$\frac{1}{2} \frac{d}{dt} E_{nonl}(t) + \frac{\nu}{\rho_0} \varepsilon \int_{\mathbb{R}^n} (\nabla u_t)^2 dx = 2\varepsilon \int_{\mathbb{R}^n} \nabla u \nabla u_t u_t dx,$$

where $E_{nonl}(t)$ is given by Eq. (18) with $\alpha = \frac{2\gamma-1}{3} \frac{\varepsilon}{c^2}$. As

$$2\varepsilon \int_{\mathbb{R}^n} \nabla u \nabla u_t u_t dx = \varepsilon \frac{d}{dt} \int_{\mathbb{R}^n} u_t (\nabla u)^2 dx - \varepsilon \int_{\mathbb{R}^n} u_{tt} (\nabla u)^2 dx,$$

we find

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \left(\int_{\mathbb{R}^n} \left[\left(1 - \frac{2\gamma-1}{3} \frac{\varepsilon u_t}{c^2} \right) (u_t)^2 + (c^2 - 2\varepsilon u_t) (\nabla u)^2 \right] dx \right. \\ \left. + 2\varepsilon \int_0^t \int_{\mathbb{R}^n} u_{tt} |\nabla u|^2 dx ds \right) + \frac{\nu}{\rho_0} \varepsilon \int_{\mathbb{R}^n} (\nabla u_t)^2 dx = 0. \quad (19) \end{aligned}$$

Thus, for $\alpha = \frac{2\gamma-1}{3} \frac{\varepsilon}{c^2}$, the function

$$F_\nu(t) = \int_{\mathbb{R}^n} [(1 - \alpha \varepsilon u_t) (u_t)^2 + (c^2 - 2\varepsilon u_t) (\nabla u)^2] dx + 2\varepsilon \int_0^t \int_{\mathbb{R}^n} u_{tt} |\nabla u|^2 dx ds$$

is constant if $\nu = 0$ and decreases if $\nu > 0$. Let us notice that while $\frac{1}{2} \leq 1 - \alpha \varepsilon u_t \leq \frac{3}{2}$, the coefficient $c^2 - 2\varepsilon u_t$ is always positive (since c is the sound speed in the chosen medium, $c^2 \gg 1$), hence the first integral in $F_\nu(t)$ is positive, but we a priori don't know the sign of the second integral, *i.e.* the sign of u_{tt} . However, for $\nu = 0$, $F_{\nu=0}(t)$ is positive, as soon as $0 \leq 1 - \alpha \varepsilon u_1$:

$$F_{\nu=0}(t) = F_{\nu=0}(0) = \int_{\mathbb{R}^n} [(1 - \alpha \varepsilon u_1) (u_1)^2 + (c^2 - 2\varepsilon u_1) (\nabla u_0)^2] dx \geq 0,$$

and, if we take the same initial data for the Cauchy problems with $\nu = 0$ and $\nu > 0$, for all $t > 0$ (for all time where $F_{\nu=0}$ exists) it holds $F_{\nu=0}(t) = F_{\nu=0}(0) > F_{\nu>0}(t)$.

For $n \geq 3$, we can control the term $2\varepsilon \int_{\mathbb{R}^n} \nabla u \nabla u_t u_t dx$ using the Hölder inequality and the Sobolev embeddings (which fails in \mathbb{R}^2):

$$\left| \int_{\mathbb{R}^n} \nabla u \nabla u_t u_t dx \right| \leq \|\nabla u\|_{L^n} \|\nabla u_t\|_{L^2} \|u_t\|_{L^{\frac{2n}{n-2}}} \leq C \|\nabla u\|_{L^n} \|\nabla u_t\|_{L^2}^2.$$

Then, instead of Eq. (19) for F_ν , we have the relation for E_{nonl} :

$$\frac{1}{2} \frac{d}{dt} E_{nonl}(t) + \left(\frac{\nu}{\rho_0} \varepsilon - 2\varepsilon C \|\nabla u\|_{L^n} \right) \int_{\mathbb{R}^n} (\nabla u_t)^2 dx \leq 0.$$

So, if a solution of the Kuznetsov equation u is such that $\|\nabla u(t)\|_{L^n}$ and $\|u_t(t)\|_{L^\infty}$ stay small enough for all time, then E_{nonl} decreases and, as previously for the Westervelt equation, thanks to $\frac{1}{2}E(t) \leq E_{nonl}(t) \leq \frac{3}{2}E(t)$, the energy E has for upper bound a decreasing function.

This fact leads us to look for global well-posedness results for the Cauchy problem for the Kuznetsov equation in the viscous case.

3 Well-posedness for the non-viscous case

3.1 Proof of Point 3 of Theorem 1

Let us give an estimation of the maximum existence time for a solution of problem (1)–(2) with $\nu = 0$. For this we are inspiring of the work of John [10] with the use of a priori estimate. However we don't directly apply the general results of John, but we improve them for our specific problem as we can take less regular initial conditions in order to have suitable a priori estimates.

Proposition 1 *For a fixed $m \in \mathbb{N}$ with $m \geq m_0 = \lceil \frac{n}{2} + 2 \rceil$, let u be a local solution of problem (1)–(2) with $\nu = 0$ on $[0, T]$ satisfying (4) and (5) for $s = m$. For $t \in [0, T]$ we have for $E_m[u](t)$, defined in Eq. (6),*

$$E_m[u](t) \leq B E_m[u](0) + C_m \max(\alpha, \beta) \varepsilon \int_0^t E_m[u](s)^{\frac{3}{2}} ds, \quad (20)$$

with constants $B = \frac{(3+2c^2)}{\min(1/2, c^2)} > 0$, depending only on c , and $C_m > 0$, depending only on m , on the dimension n and on c (only if $\min(1/2, c^2) = c^2$).

Proof : Following [10], let us consider

$$L_u v = v_{tt} - c^2 \Delta v - \alpha \varepsilon u_t v_{tt} - \beta \varepsilon \nabla u \nabla v_t, \quad (21)$$

where u is a local solution on $[0, T]$ of the problem (1)–(2) with $\nu = 0$, satisfying (4) and (5) for $s = m$. We multiply Eq. (21) by v_t and integrate over \mathbb{R}^n

$$\begin{aligned} \int_{\mathbb{R}^n} L_u v v_t &= \frac{1}{2} \frac{d}{dt} \left(\int_{\mathbb{R}^n} v_t^2 + c^2 (\nabla v)^2 dx \right) - \alpha \varepsilon \int_{\mathbb{R}^n} u_t v_{tt} v_t dx - \beta \varepsilon \int_{\mathbb{R}^n} \nabla u \nabla v_t v_t dx \\ &= \frac{1}{2} \frac{d}{dt} \left(\int_{\mathbb{R}^n} v_t^2 + c^2 (\nabla v)^2 dx \right) - \frac{\alpha}{2} \varepsilon \left[\frac{d}{dt} \left(\int_{\mathbb{R}^n} u_t v_t^2 dx \right) - \int_{\mathbb{R}^n} u_{tt} v_t^2 dx \right] \\ &\quad - \frac{\beta}{2} \varepsilon \int_{\mathbb{R}^n} \nabla u \nabla (v_t)^2 dx \\ &= \frac{1}{2} \frac{d}{dt} \left(\int_{\mathbb{R}^n} v_t^2 + c^2 (\nabla v)^2 dx \right) - \frac{\alpha}{2} \varepsilon \left[\frac{d}{dt} \left(\int_{\mathbb{R}^n} u_t v_t^2 dx \right) - \int_{\mathbb{R}^n} u_{tt} v_t^2 dx \right] \\ &\quad + \frac{\beta}{2} \varepsilon \int_{\mathbb{R}^n} \Delta u (v_t)^2 dx. \end{aligned}$$

Hence, denoting by

$$I[v] = v_t^2 + c^2 (\nabla v)^2 - \alpha \varepsilon u_t v_t^2, \quad (22)$$

$$J[v] = 2L_u v v_t - [\alpha \varepsilon u_{tt} + \beta \varepsilon \Delta u] (v_t)^2, \quad (23)$$

we have the following equation

$$\frac{d}{dt} \int_{\mathbb{R}^n} I[v](t, x) dx = \int_{\mathbb{R}^n} J[v](t, x) dx. \quad (24)$$

Let $A = (A_0, A_1, \dots, A_n)$ be a multi-index, and $D^A v = \partial_t^{A_0} \partial_{x_1}^{A_1} \dots \partial_{x_n}^{A_n}$. To prove estimate (20), we study $|\int_{\mathbb{R}^n} J[v](t, x) dx|$ for $v = D^A u$ with $|A| = A_0 + A_1 + \dots + A_n \leq m$.

For $m \geq [\frac{n}{2} + 2]$ and a multi-index A with $|A| \leq m$ we estimate, thanks to the definition of $E_m[u]$ by Eq. (6),

$$\int_{\mathbb{R}^n} |u_{tt} (D^A u_t)^2| dx \leq \|u_{tt}\|_{L^\infty} \|D^A u_t\|_{L^2}^2 \leq C \|u_{tt}\|_{H^{[\frac{n}{2}+1]}} E_m[u] \leq C E_m[u]^{\frac{3}{2}}, \quad (25)$$

with a constant $C > 0$, depending only of n by the Sobolev embedding [1] Theorem 7.57 p. 228

$$H^s(\mathbb{R}^n) \hookrightarrow L^\infty(\mathbb{R}^n) \quad \text{for } s > \frac{n}{2}. \quad (26)$$

In the same way, using the Sobolev embedding (26), we obtain

$$\begin{aligned} \int_{\mathbb{R}^n} |\Delta u (D^A u_t)^2| dx &\leq \|\Delta u\|_{L^\infty} \|D^A u_t\|_{L^2}^2 \leq C \|\Delta u\|_{H^{[\frac{n}{2}+1]}} E_m[u] \\ &\leq C \|\nabla u\|_{H^m} E_m[u] \leq C E_m[u]^{\frac{3}{2}}. \end{aligned} \quad (27)$$

To calculate $L_u D^A u$ we apply the chain rule of differentiation to $D^A L_u u = 0$. As $L_u u = 0$ we suppose $|A| \geq 1$. By developing $D^A(u_t u_{tt})$ and $D^A(\nabla u \nabla u_t) = \sum_{i=1}^n D^A(\partial_{x_i} u \partial_{x_i} u_t)$, we have

$$L_u D^A u = \varepsilon \sum_j \left(C_j \alpha D^{A^{j1}} u_t D^{A^{j2}} u_t + \sum_{i=1}^n E_{ij} \beta D^{A^{j1}} \partial_{x_i} u D^{A^{j2}} \partial_{x_i} u \right), \quad (28)$$

where \sum_j is a finite sum, with C_j and E_{ij} depending only of $|A| \leq m$, and A^{j1} and A^{j2} are multi-index such that

$$\begin{cases} |A^{j1}| + |A^{j2}| = |A| + 1, \\ |A^{j1}| \geq 1, \quad |A^{j2}| \geq 1, \\ A_0^{j1} + A_0^{j2} = A_0 + 1, \quad A_i^{j1} + A_i^{j2} = A_i \text{ for } 1 \leq i \leq n. \end{cases} \quad (29)$$

Let us show for $m \geq \lfloor \frac{n}{2} + 2 \rfloor$ the estimate

$$\left| \int_{\mathbb{R}^n} L_u D^A u D^A u_t \, dx \right| \leq C \varepsilon \max(\alpha, \beta) E_m[u]^{\frac{3}{2}}. \quad (30)$$

Without loss of generality, we consider two multi-indexes A^1 and A^2 satisfying (29) and divide the proof of (30) in two parts: firstly we estimate $\int_{\mathbb{R}^n} |D^{A^1} u_t D^{A^2} u_t D^A u_t| \, dx$, and secondary $\int_{\mathbb{R}^n} |D^{A^1} \partial_{x_i} u D^{A^2} \partial_{x_i} u D^A u_t| \, dx$. As the proof of each part is very similar, we give the details only for the first one.

To estimate $\int_{\mathbb{R}^n} |D^{A^1} u_t D^{A^2} u_t D^A u_t| \, dx$, we consider three cases:

Case 1: $1 < |A^1| < m$ and $1 < |A^2| < m$,

Case 2: $|A^1| \leq m$ and $|A^2| = 1$,

Case 3: $|A^2| \leq m$ and $|A^1| = 1$.

Case 1: For $2 \leq |A^1| \leq m - 1$ and $2 \leq |A^2| \leq m - 1$, it holds

$$\int_{\mathbb{R}^n} |D^{A^1} u_t D^{A^2} u_t D^A u_t| \, dx \leq \|D^{A^1} u_t\|_{L^p} \|D^{A^2} u_t\|_{L^q} \|D^A u_t\|_{L^2},$$

with $\frac{1}{p} + \frac{1}{q} = \frac{1}{2}$ by the general Hölder inequality [4]. Hence, using the Sobolev embedding [1]

$$H^{m_1}(\mathbb{R}^n) \hookrightarrow L^p(\mathbb{R}^n) \quad \text{with } \frac{1}{p} = \frac{1}{2} - \frac{m_1}{n} \text{ and } 0 < m_1 < \frac{n}{2}, \quad (31)$$

we find

$$\int_{\mathbb{R}^n} |D^{A^1} u_t D^{A^2} u_t D^A u_t| \, dx \leq C \|D^{A^1} u_t\|_{H^{m_1}} \|D^{A^2} u_t\|_{H^{\frac{n}{2} - m_1}} \|D^A u_t\|_{L^2}.$$

In what follows by $C > 0$ is denoted an arbitrary constant depending only on m and on n .

We have

$$\|D^{A^1} u_t\|_{H^{m_1}} \leq \|\partial_t^{A_0^1} u_t\|_{H^{m_1+|A^1|-A_0^1}}, \quad \|D^{A^2} u_t\|_{H^{\frac{n}{2}-m_1}} \leq \|\partial_t^{A_0^2} u_t\|_{H^{\frac{n}{2}-m_1+|A^2|-A_0^2}}.$$

We need to find m for which there exists m_1 with $0 < m_1 < \frac{n}{2}$, such that

$$\begin{cases} m_1 + |A^1| - A_0^1 \leq m + 1 - (A_0^1 + 1), \\ \frac{n}{2} - m_1 + |A^2| - A_0^2 \leq m + 1 - (A_0^2 + 1), \end{cases} \quad (32)$$

or equivalently, by (29) $|A^2| = |A| + 1 - |A^1|$,

$$\begin{cases} m_1 + |A^1| \leq m, \\ \frac{n}{2} - m_1 + |A| + 1 - |A^1| \leq m. \end{cases}$$

As $m - |A| \geq 0$ it is sufficient to find m_1 , such that

$$\frac{n}{2} + 1 \leq m_1 + |A^1| \leq m,$$

with $2 \leq |A^1| \leq m - 1$ and $0 < m_1 < \frac{n}{2}$. In particular, the last three inequalities imply that $m \geq [2 + \frac{n}{2}]$. For the existence of m_1 , we see that, for instance,

if $|A_1| = 2$: we can take $m_1 = \frac{n}{2} - \frac{1}{4}$,

if $2 < |A_1| < \frac{n}{2} + 1$: we can take $m_1 = \frac{n}{2} + 1 - |A_1|$,

if $\frac{n}{2} + 1 \leq |A_1| \leq m - 1$: we can take $m_1 = \frac{1}{4}$.

Moreover,

$$\|D^A u_t\|_{L^2} \leq \|\partial_t^{A_0} u_t\|_{H^{|A|-A_0}} \leq \|\partial_t^{A_0} u_t\|_{H^{m-A_0}}.$$

Then, thanks to relations (32), we conclude

$$\begin{aligned} \int_{\mathbb{R}^n} |D^{A^1} u_t D^{A^2} u_t D^A u_t| &\leq C \|\partial_t^{A_0^1} u_t\|_{H^{m-A_0^1}} \|\partial_t^{A_0^2} u_t\|_{H^{m-A_0^2}} \|\partial_t^{A_0} u_t\|_{H^{m-A_0}} \\ &\leq C E_m[u]^{\frac{3}{2}}. \end{aligned}$$

Case 2: Now, if $|A^1| \leq m$ and $|A^2| = 1$, it holds

$$\int_{\mathbb{R}^n} |D^{A^1} u_t D^{A^2} u_t D^A u_t| \leq \|D^{A^1} u_t\|_{L^2} \|D^{A^2} u_t\|_{L^\infty} \|D^A u_t\|_{L^2}.$$

Since $|A^1| \leq m$ and $|A| \leq m$, we have

$$\|D^{A^1} u_t\|_{L^2} \leq \|\partial_t^{A_0^1} u_t\|_{H^{m-A_0^1}} \leq \sqrt{E_m[u]}, \quad \|D^A u_t\|_{L^2} \leq \|\partial_t^{A_0} u_t\|_{H^{m-A_0}} \leq \sqrt{E_m[u]}.$$

In addition, since $m \geq [\frac{n}{2} + 2]$ and $A_0^2 \leq 1$, by the Sobolev embeddings (26) it holds

$$\|D^{A^2} u_t\|_{L^\infty} \leq C \|D^{A^2} u_t\|_{H^{[\frac{n}{2}+1]}} \leq C \|\partial_t^{A_0^2} u_t\|_{H^{[\frac{n}{2}+2]-A_0^2}} \leq C \sqrt{E_m[u]}.$$

Case 3: The case 3, *i.e.* when $|A^2| \leq m$ and $|A^1| = 1$, is the same as Case 2.

Consequently, for $m \geq [\frac{n}{2} + 2]$, and A^1 and A^2 , satisfying properties (29), it holds

$$\int_{\mathbb{R}^n} |D^{A^1} u_t D^{A^2} u_t D^A u_t| dx \leq C E_m[u]^{\frac{3}{2}}. \quad (33)$$

By the same argument, for $m \geq [\frac{n}{2} + 2]$ and A^1 and A^2 , satisfying properties (29), we control the terms of the form $\int_{\mathbb{R}^n} |D^{A^1} \partial_{x_i} u D^{A^2} \partial_{x_i} u D^A u_t| dx$:

$$\int_{\mathbb{R}^n} |D^{A^1} \partial_{x_i} u D^{A^2} \partial_{x_i} u D^A u_t| \leq C E_m[u]^{\frac{3}{2}}. \quad (34)$$

Thus, considering (28), (33) and (34) for $m \geq [\frac{n}{2} + 2]$ and for a multi-index A with $|A| \leq m$, we have estimate (30).

Thanks to estimations (25), (27) and (30), we are able to control each term of $J[D^A u]$ from Eq. (23):

$$\left| \int_{\mathbb{R}^n} J[D^A u](t, x) dx \right| \leq C \max(\alpha, \beta) \varepsilon E_m[u](t)^{\frac{3}{2}}. \quad (35)$$

By the hypothesis that u is a local solution of the non-viscous Kuznetsov equation, u satisfies Eq. (5), *i.e.* $\|u_t(t)\|_{L^\infty} \leq \frac{1}{2\alpha\varepsilon}$ on $[0, T]$, which implies the equivalence of energies

$$\int_{\mathbb{R}^n} \frac{1}{2} (D^A u_t)^2 + c^2 (\nabla D^A u)^2 dx \leq \int_{\mathbb{R}^n} I[D^A u](t, x) dx \leq \int_{\mathbb{R}^n} \frac{3}{2} (D^A u_t)^2 + c^2 (\nabla D^A u)^2 dx.$$

We integrate relation (24) over $[0, t]$ with $t \leq T$ to obtain

$$\begin{aligned} \|D^A u_t(t)\|_{L^2}^2 + \|\nabla D^A u(t)\|_{L^2}^2 &\leq \frac{(\frac{3}{2} + c^2)}{\min(1/2, c^2)} (\|D^A u_t(0)\|_{L^2}^2 + \|\nabla D^A u(0)\|_{L^2}^2) \\ &\quad + \frac{1}{\min(1/2, c^2)} \int_0^t \int_{\mathbb{R}^n} J(s, x) dx ds. \end{aligned}$$

Then, using estimate (35), we find

$$\begin{aligned} \|D^A u_t(t)\|_{L^2}^2 + \|\nabla D^A u(t)\|_{L^2}^2 &\leq \frac{(\frac{3}{2} + c^2)}{\min(1/2, c^2)} (\|D^A u_t(0)\|_{L^2}^2 + \|\nabla D^A u(0)\|_{L^2}^2) \\ &\quad + \frac{1}{\min(1/2, c^2)} C \max(\alpha, \beta) \varepsilon \int_0^t E_m[u](s)^{\frac{3}{2}} ds. \end{aligned}$$

As we have this for all multi-index A with $|A| \leq m$, by summing, we obtain

$$E_m[u](t) \leq \frac{(3 + 2c^2)}{\min(1/2, c^2)} E_m[u](0) + \frac{1}{\min(1/2, c^2)} C \max(\alpha, \beta) \varepsilon \int_0^t E_m[u](s)^{\frac{3}{2}} ds$$

with a constant $C > 0$, depending only of n and m . This gives estimate (20). \square

Inequality (20), proved in Proposition 1, gives us an a priori estimate in order to have, with the help of the Gronwall Lemma, an estimation of the maximum existence time T^* . However, when m increases, C_m increases, and the maximum existence time, given by estimate (20), decreases whereas the initial conditions become more regular. Therefore we prove the second a priori estimate (see Eq. (36)), playing a key role in order to avoid this problem:

Proposition 2 *Let conditions of Proposition 1 be satisfied. Then for $t \in [0, T]$ and $m \geq \lceil \frac{n}{2} + 3 \rceil$ we have*

$$E_m[u](t) \leq B E_m[u](0) + D_m \max(\alpha, \beta) \varepsilon \int_0^t E_{m-1}[u](s)^{\frac{1}{2}} E_m[u](s) ds, \quad (36)$$

with a constant $D_m > 0$, depending only on m , on n and on c and the same constant B as in Proposition 1.

Proof :

Using the proof of Proposition 1 for A a multi-index with $|A| \leq m$ and $J[D^A u]$ from Eq. (23), we just have to show that for $m \geq \lceil \frac{n}{2} + 3 \rceil$

$$\left| \int_{\mathbb{R}^n} J[D^A u](s, x) dx \right| \leq C \varepsilon \sqrt{E_{m-1}[u]} E_m[u].$$

Indeed, for $m \geq \lceil \frac{n}{2} + 3 \rceil$ and a multi-indexes A with $|A| \leq m$ we have

$$\int_{\mathbb{R}^n} |u_{tt}(D^A u_t)^2| dx \leq \|u_{tt}\|_{L^\infty} \|D^A u_t\|_{L^2}^2 \leq C \|u_{tt}\|_{H^{m-2}} E_m[u] \leq C \sqrt{E_{m-1}[u]} E_m[u],$$

and

$$\int_{\mathbb{R}^n} |\Delta u(D^A u_t)^2| dx \leq \|\Delta u\|_{L^\infty} \|D^A u_t\|_{L^2}^2 \leq C \|\nabla u\|_{H^{m-1}} E_m[u] \leq C \sqrt{E_{m-1}[u]} E_m[u].$$

Now we consider two multi-indexes A^1 and A^2 with properties (29). As previously, we have to distinguish three cases:

Case 1: $1 < |A^1| < m$ and $1 < |A^2| < m$,

Case 2: $|A^1| \leq m$ and $|A^2| = 1$,

Case 3: $|A^2| \leq m$ and $|A^1| = 1$.

First, if $1 < |A^1| < m$ and $1 < |A^2| < m$, in Case 1 we have

$$\begin{aligned} \int_{\mathbb{R}^n} |D^{A^1} u_t D^{A^2} u_t D^A u_t| dx &\leq C \|\partial_t^{A_0^1} u_t\|_{H^{m_1+|A^1|-A_0^1}} \|\partial_t^{A_0^2} u_t\|_{H^{\frac{n}{2}-m_1+|A^2|-A_0^2}} \|D^A u_t\|_{L^2} \\ &\leq \|\partial_t^{A_0^1} u_t\|_{H^{m_1+|A^1|-A_0^1}} \|\partial_t^{A_0^2} u_t\|_{H^{\frac{n}{2}-m_1+|A^2|-A_0^2}} \sqrt{E_m[u]}. \end{aligned}$$

By symmetry of their roles we can take $|A_1| \leq \lceil \frac{m+1}{2} \rceil \leq |A_2|$. We look for m for which there exists m_1 , such that

$$\begin{cases} m_1 + |A^1| - A_0^1 \leq m - (A_0^1 + 1), \\ \frac{n}{2} - m_1 + |A^2| - A_0^2 \leq m + 1 - (A_0^2 + 1), \end{cases}$$

in order to have

$$\|\partial_t^{A_0^1} u_t\|_{H^{m_1+|A^1|-A_0^1}} \leq \sqrt{E_{m-1}[u]} \text{ and } \|\partial_t^{A_0^2} u_t\|_{H^{\frac{n}{2}-m_1+|A^2|-A_0^2}} \leq \sqrt{E_m[u]}.$$

As $|A_2| = |A| + 1 - |A^1|$ and $m \geq |A|$, it is sufficient to find m_1 , such that

$$\frac{n}{2} + 1 \leq m_1 + |A^1| \leq m - 1,$$

with $2 \leq |A^1| \leq m - 1$, $|A_1| \leq \lceil \frac{m+1}{2} \rceil$ and $0 < m_1 < \frac{n}{2}$. This directly implies that $m \geq \lceil \frac{n}{2} + 3 \rceil$. In addition, this also implies, except if $m \leq 3$, that $|A^1| \leq m - 2$. Thus the existence of a required m_1 is justified in Case 1 of the proof of estimate (20). If $m \leq 3$, as $m \geq \lceil \frac{n}{2} + 3 \rceil$, we have only the case $m = 3$, $n = 1$ and $|A^1| = |A^2| = 2$, for which it holds

$$\begin{aligned} \int_{\mathbb{R}^n} |D^{A^1} u_t D^{A^2} u_t D^A u_t| dx &\leq \|D^{A^1} u_t\|_{L^2} \|D^{A^2} u_t\|_{L^\infty} \|D^A u_t\|_{L^2} \\ &\leq C \|D^{A^1} u_t\|_{L^2} \|D^{A^2} u_t\|_{H^1} \|D^A u_t\|_{L^2} \leq C \sqrt{E_{m-1}[u]} E_m[u], \end{aligned}$$

by the Sobolev embedding (26).

For $|A^1| \leq m$ and $|A^2| = 1$, in Case 2, we find

$$\int_{\mathbb{R}^n} |D^{A^1} u_t D^{A^2} u_t D^A u_t| dx \leq \|D^{A^1} u_t\|_{L^2} \|D^{A^2} u_t\|_{L^\infty} \|D^A u_t\|_{L^2},$$

where the two L^2 -norms are controlled by $\sqrt{E_m[u]}$:

$$\|D^{A^1} u_t\|_{L^2} \leq \|\partial_t^{A_0^1} u_t\|_{H^{m-A_0^1}} \leq \sqrt{E_m[u]}, \quad \|D^A u_t\|_{L^2} \leq \|\partial_t^{A_0} u_t\|_{H^{m-A_0}} \leq \sqrt{E_m[u]}.$$

As $A_0^2 \leq 1$, for the L^∞ -norm, for $m \geq \lceil \frac{n}{2} + 3 \rceil$, we have

$$\|D^{A^2} u_t\|_{L^\infty} \leq C \|D^{A^2} u_t\|_{H^{\lceil \frac{n}{2} + 1 \rceil}} \leq C \|\partial_t^{A_0^2} u_t\|_{H^{\lceil \frac{n}{2} + 2 \rceil - A_0^2}} \leq C \sqrt{E_{m-1}[u]}.$$

Case 3, *i.e.* for $|A^2| \leq m$ and $|A^1| = 1$, can be treated in the same way.

So, if $m \geq \lceil \frac{n}{2} + 3 \rceil$, and A^1 and A^2 satisfy properties (29), we obtain

$$\int_{\mathbb{R}^n} |D^{A^1} u_t D^{A^2} u_t D^A u_t| dx \leq C \sqrt{E_{m-1}[u]} E_m[u].$$

Using similar arguments, we can show with the same restrictions on m , A^1 and A^2 that

$$\int_{\mathbb{R}^n} |D^{A^1} \partial_{x_i} u D^{A^2} \partial_{x_i} u D^A u_t| dx \leq C \sqrt{E_{m-1}[u]} E_m[u].$$

Consequently,

$$\int_{\mathbb{R}^n} |L_u D^A u D^A u_t| dx \leq C \max(\alpha, \beta) \sqrt{E_{m-1}[u]} E_m[u],$$

from where follows estimate (36). □

We are now going to give a first estimation of the lifespan T^* of a local solution of problem (1)–(2) with $\nu = 0$.

Theorem 3 Let $m \geq m_0 = \lfloor \frac{n}{2} + 2 \rfloor$ and let u be the unique solution on $[0, T^*[$ of the problem (1)–(2) with $\nu = 0$ for

$$u_0 \in H^{m+1}(\mathbb{R}^n), \quad u_1 \in H^m(\mathbb{R}^n) \quad \text{and} \quad \|u_1\|_{L^\infty} < \frac{1}{2\alpha\varepsilon}.$$

If $\sqrt{E_{m_0}[u](0)} \leq \frac{1}{4\sqrt{B}C_\infty\alpha\varepsilon}$, then

$$T^* > T_0 = \frac{1}{C_{m_0} \max(\alpha, \beta)\varepsilon\sqrt{B}E_{m_0}[u](0)} \quad (37)$$

and

$$u \in C^r([0, T_0]; H^{m+1-r}) \text{ for } 0 \leq r \leq m+1,$$

with

$$\forall t \in [0, T_0], \quad E_m[u](t) \leq C < +\infty.$$

Here B and C_{m_0} are the constants from estimate (20) and C_∞ is the embedding constant from the embedding of the Sobolev space $H^{\lfloor \frac{n}{2} + 1 \rfloor}(\mathbb{R}^n)$ in $L^\infty(\mathbb{R}^n)$.

Proof : Thanks to Point 1 of Theorem 1, for $u_0 \in H^{m+1}(\mathbb{R}^n)$, $u_1 \in H^m(\mathbb{R}^n)$ and $\|u_1\|_{L^\infty} < \frac{1}{2\alpha\varepsilon}$ there exists a unique solution u on a sufficiently small interval $[0, T]$ of problem (1)–(2) with $\nu = 0$, satisfying (4) and (5) for $s = m$. Moreover it implies that $E_m[u](0)$ is finite. Hence, we can add the hypothesis

$$\sqrt{E_{m_0}[u](0)} \leq \frac{1}{4\sqrt{B}C_\infty\alpha\varepsilon}$$

without adding further conditions of regularity on u_0 and u_1 as it can be reduced on a smallness condition on $\|u_0\|_{H^{m+1}} + \|u_1\|_{H^m}$.

Let us take T_0 , as defined in Eq. (37), and show by induction on $j \in \mathbb{N}$ with $m_0 \leq j \leq m$ that

$$\forall j \in \mathbb{N}, \text{ with } m_0 \leq j \leq m \quad \sup_{t \in [0, T_0]} E_j[u](t) < \infty.$$

For $j = m_0$, $u_0 \in H^{m+1}(\mathbb{R}^n) \subseteq H^{m_0+1}(\mathbb{R}^n)$ and $u_1 \in H^m(\mathbb{R}^n) \subseteq H^{m_0}(\mathbb{R}^n)$, and consequently

$$E_{m_0}[u](0) \leq E_m[u](0) < \infty.$$

For $t \geq 0$, while $\|u_t(t)\|_{L^\infty} \leq \frac{1}{2\alpha\varepsilon}$, it holds estimate (20) with $m = m_0$. According to the Gronwall Lemma, applied to (20) with $m = m_0$, we have

$$E_{m_0}[u](t) \leq z(t),$$

where $z(t)$ is the solution of the Cauchy problem for an ordinary differential equation

$$z(t) = z_0 + C_{m_0} \max(\alpha, \beta)\varepsilon \int_0^t (z(s))^{3/2} ds \quad \text{with } z_0 = B E_{m_0}[u](0).$$

This problem can be solved explicitly:

$$z(t) = \frac{z_0}{(1 - \frac{1}{2}z_0^{1/2}C_{m_0} \max(\alpha, \beta)\varepsilon t)^2}.$$

We can see that, as long as $0 \leq t \leq T_0$, the function $z(t)$ has the finite upper bound $z(t) \leq 4z_0$. It implies the upper boundness of $E_{m_0}[u]$:

$$E_{m_0}[u](t) \leq 4B E_{m_0}[u](0). \quad (38)$$

Moreover, thanks to our notations,

$$\frac{\|u_t(t)\|_{L^\infty}}{C_\infty} \leq \|u_t(t)\|_{H^{[\frac{n}{2}+1]}} \leq \sqrt{E_{m_0}[u](t)},$$

from where, using inequality (38), we find

$$\|u_t(t)\|_{L^\infty} \leq 2C_\infty \sqrt{B E_{m_0}[u](0)} \leq \frac{1}{2\alpha\varepsilon},$$

since $\sqrt{E_{m_0}[u](0)} \leq \frac{1}{4\sqrt{B}C_\infty\alpha\varepsilon}$. Thus Eq. (5) holds on all interval $[0, T_0]$ and $\sup_{t \in [0, T_0]} E_{m_0}[u](t)$ is finite.

Let $j \in \mathbb{N}$, $m_0 \leq j \leq m-1$ be such that $\sup_{t \in [0, T_0]} E_j[u](t) < \infty$.

Since Eq. (5) holds on all interval $[0, T_0]$, we can use the a priori estimate (36) and write that for all $t \in [0, T_0]$

$$E_{j+1}[u](t) \leq B E_{j+1}[u](0) + D_{j+1} \max(\alpha, \beta) \varepsilon \int_0^t \sqrt{E_j[u](s)} E_{j+1}[u](s) ds.$$

By the induction hypothesis $\sup_{t \in [0, T_0]} E_j[u](t)$ is bounded by a constant, denoted here by E^2 , and hence on $[0, T_0]$ it holds

$$E_{j+1}[u](t) \leq B E_{j+1}[u](0) + D_{j+1} \max(\alpha, \beta) E \varepsilon \int_0^t E_{j+1}[u](s) ds.$$

Applying the Gronwall Lemma, we obtain for $t \in [0, T_0]$

$$E_{j+1}[u](t) \leq B E_{j+1}[u](0) e^{D_{j+1} \max(\alpha, \beta) E \varepsilon t} \leq B E_{j+1}[u](0) e^{D_{j+1} \max(\alpha, \beta) E \varepsilon T_0}.$$

This means, as $E_{j+1}[u](0) \leq E_m[u](0) < +\infty$, that $\sup_{t \in [0, T_0]} E_{j+1}[u](t) < \infty$ and finishes the proof. \square

Theorem 3 estimates the lifespan T^* as at least of the order $\frac{1}{\varepsilon}$, or more precisely, implies that

$$\liminf_{\varepsilon \rightarrow 0} \varepsilon T^* > 0.$$

This result is independent of the dimension n . However, much better estimations for the lifespan can be obtained, if we use an inequality that takes into account the time decay of the solutions for $n > 1$, what we do in the next Section.

3.2 Proof of Point 5 of Theorem 1. Optimal estimations of the existence time

In [10] John uses the group of linear transformations preserving the equation $u_{tt} - \Delta u = 0$. The generators of this group (the derivatives with respect to group parameters taken at the identity), here called generalized derivatives, include in addition to the derivatives $\partial_t, \partial_{x_1}, \dots, \partial_{x_n}$ certain first-order differential operators L_α with $\alpha = 0, \dots, n$ and Ω_{ik} with $1 \leq i < k \leq n$:

Definition 1 (Generalized derivatives [10]) *The following operators*

$$L_0 = t\partial_t + \sum_i x_i \partial_{x_i}, \quad L_i = x_i \partial_t + t\partial_{x_i} \text{ for } i = 1, \dots, n,$$

$$\Omega_{ik} = x_i \partial_{x_k} - x_k \partial_{x_i} \text{ for } 1 \leq i < k \leq n, \text{ and } \partial_t, \partial_{x_i} \text{ for } i = 1, \dots, n$$

are called the generalized derivatives. The operators

$$L_0, \dots, L_n, \Omega_{12}, \Omega_{13}, \dots, \Omega_{n-1n}, \partial_t, \partial_{x_1}, \dots, \partial_{x_n},$$

(taken in this order) are denoted respectively by $\Gamma_0, \dots, \Gamma_\mu$ with $\mu = \frac{1}{2}(n^2 + 3n + 2)$. For a multi-index $A = (A_0, \dots, A_\mu)$ we write in the usual way

$$|A| = A_0 + \dots + A_\mu, \quad \Gamma^A = (\Gamma_0)^{A_0} (\Gamma_1)^{A_1} \dots (\Gamma_\mu)^{A_\mu}.$$

Therefore, in the framework of the general derivatives, we define for $m \in \mathbb{N}$

$$E_{\infty, m}[u](t) = \sup_x \left| \sup_{|A| \leq m} [(\Gamma^A \partial_t u(t, x))^2 + (\Gamma^A \nabla u(x, t))^2] \right|, \quad (39)$$

$$E_{1, m}[u](t) = \sum_{|A| \leq m} (\|\Gamma^A \partial_t u\|_{L^2}^2 + \|\Gamma^A \nabla u\|_{L^2}^2)(t). \quad (40)$$

Let us give a remarkable estimation proved in Ref. [15] by Klainerman:

Proposition 3 (Klainerman 1987) *For $n^* = [\frac{n}{2} + 1]$, $m \in \mathbb{N}$, and $t > 0$, as soon as u is such that $E_{1, m+n^*}[u](t)$ is finite, it holds*

$$\sqrt{E_{\infty, m}[u](t)} \leq C_n (1+t)^{\frac{1-n}{2}} \sqrt{E_{1, m+n^*}[u](t)}. \quad (41)$$

Thanks to Proposition 3, we improve the results of John [10] for the case of the Kuznetsov equation and state:

Proposition 4 *For n and m in \mathbb{N}^* , $m \geq n+2$, let u be a local solution on an interval $[0, T]$ of the problem (1)–(2) with $\nu = 0$, satisfying (4) and (5) with $s = m$. Then for all $t \in [0, T]$, it holds*

$$E_{1, m}[u](t) \leq B E_{1, m}[u](0) + C_m \max(\alpha, \beta) \varepsilon \int_0^t (1+s)^{(1-n)/2} E_{1, m}[u](s)^{\frac{3}{2}} ds, \quad (42)$$

with a positive constant $B > 0$, depending only on c , on α and on β , and with a positive constant $C_m > 0$, depending only on m , on n and on c .

Proof : The proof follows identically the proof of Proposition 1 up to Eq. (28) replacing everywhere D^A by Γ^A . This time Eq. (28) becomes

$$L_u \Gamma^A u = \varepsilon \sum_{j=0}^{\mu} \left(\alpha C_j \Gamma^{A^{j1}} u_t \Gamma^{A^{j2}} u_t + \sum_{i=1}^n \beta E_{ij} \Gamma^{A^{j1}} \partial_{x_i} u \Gamma^{A^{j2}} \partial_{x_i} u \right), \quad (43)$$

where μ is defined in Definition 1, C_j and E_{ij} depend only on $|A| \leq m$, and A^{j1} and A^{j2} are multi-indexes, such that

$$|A^{j1}| + |A^{j2}| \leq m + 1.$$

It follows that $|A^{j1}| \leq [\frac{m+1}{2}]$ or $|A^{j2}| \leq [\frac{m+1}{2}]$. Therefore, if we set $m' = [\frac{m+1}{2}]$, we obtain

$$\begin{aligned} |J[\Gamma^A u](s, x)| &\leq C_m \max(\alpha, \beta) \varepsilon \sqrt{\sup_{|B| \leq m'} ((\Gamma^B \partial_t u(t, x))^2 + (\Gamma^B \nabla u(x, t))^2)} \\ &\quad \cdot \sup_{|B| \leq m} ((\Gamma^B \partial_t u(t, x))^2 + (\Gamma^B \nabla u(x, t))^2) \\ &\leq C_m \max(\alpha, \beta) \varepsilon \sqrt{E_{\infty, m'}[u](s)} \sum_{|B| \leq m} ((\Gamma^B \partial_t u(t, x))^2 + (\Gamma^B \nabla u(x, s))^2), \end{aligned}$$

and thus

$$\left| \int_{\mathbb{R}^n} J[\Gamma^A u](s, x) dx \right| \leq C_m \max(\alpha, \beta) \varepsilon \sqrt{E_{\infty, m'}[u](s)} E_{1, m}[u](s).$$

By hypothesis on u ,

$$\|u_t(t)\|_{L^\infty} \leq \frac{1}{2\alpha\varepsilon} \text{ on } [0, T],$$

and then, by integrating of Eq. (24) on $[0, t]$ with $t \in [0, T]$, we have

$$\begin{aligned} \frac{1}{2} \|\partial_t \Gamma^A u(t)\|_{L^2}^2 + c^2 \|\nabla \Gamma^A u(t)\|_{L^2}^2 &\leq \frac{3}{2} \|\partial_t \Gamma^A u(0)\|_{L^2}^2 + c^2 \|\nabla \Gamma^A u(0)\|_{L^2}^2 \\ &\quad + C_m \max(\alpha, \beta) \varepsilon \int_0^t \sqrt{E_{\infty, m'}[u](s)} E_{1, m}[u](s) ds. \end{aligned}$$

By summing for $|A| \leq m$, we obtain

$$E_{1, m}[u](t) \leq B E_{1, m}[u](0) + C_m \max(\alpha, \beta) \varepsilon \int_0^t \sqrt{E_{\infty, m'}[u](s)} E_{1, m}[u](s) ds.$$

Now we use the Klainerman inequality (41), noticing that, if we take $m \geq n + 2$, we have

$$m' + n^* = \left[\frac{m+1}{2} \right] + \left[\frac{n}{2} + 1 \right] \leq m.$$

This finishes the proof. \square

We use the a priori estimate (42) to improve our estimation of the lifespan T^* as a function of n .

Theorem 4 Let $m \geq n+2$. For $u_0 \in H^{m+1}(\mathbb{R}^n)$ and $u_1 \in H^m(\mathbb{R}^n)$ with $\|u_1\|_{L^\infty} \leq \frac{1}{2\alpha\varepsilon}$, we consider the local solution u of problem (1)–(2) with $\nu = 0$ on an interval $[0, T]$, satisfying (4) and (5) for $s = m$ as in Point 1 of Theorem 1. If $\sqrt{E_{1,m}[u](0)} \leq \frac{1}{4\sqrt{BC_\infty\alpha\varepsilon}}$, then

$$E_{1,m}[u](t) \leq 4B E_{1,m}[u](0),$$

as long as

$$\begin{aligned} t &\leq \left(2C_m \max(\alpha, \beta)\varepsilon \sqrt{B E_{1,m}[u](0)}\right)^{-2} \quad \text{when } n = 2, \\ t &\leq 2\exp\left(\frac{1}{C_m \max(\alpha, \beta)\varepsilon \sqrt{B E_{1,m}[u](0)}}\right) \quad \text{when } n = 3, \\ 1 &\leq \left(2C_m \max(\alpha, \beta)\varepsilon \sqrt{B E_{1,m}[u](0)}\right)^{-1} \quad \text{when } n \geq 4. \end{aligned}$$

Consequently,

$$\begin{aligned} \liminf_{\varepsilon \rightarrow 0} \varepsilon^2 T^* &> 0 \quad \text{when } n = 2, \\ \liminf_{\varepsilon \rightarrow 0} \varepsilon \log(T^*) &> 0 \quad \text{when } n = 3, \end{aligned}$$

and, for a small enough ε , $T^* = +\infty$ when $n \geq 4$, i.e. the solution u is global.

Proof : This is a direct consequence of the Gronwall lemma, used with the a priori estimate (42), as it is done by John in [10]. \square

Remark 1 The estimations, given for T^* in the case $n = 1, 2, 3$, are optimal as the blow-up times, found by Alinhac in [2], have the same order.

3.3 Proof of Point 4 of Theorem 1. Stability and uniqueness result

By definition of u and v we have

$$(u - v)_{tt} - c^2 \Delta(u - v) = \alpha\varepsilon(u_t u_{tt} - v_t v_{tt}) + \beta\varepsilon(\nabla u \nabla u_t - \nabla v \nabla v_t). \quad (44)$$

We multiply this equation by $(u - v)_t$ and integrate on \mathbb{R}^n . By integration by parts we obtain

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} (\|(u - v)_t\|_{L^2}^2 + c^2 \|\nabla(u - v)\|_{L^2}^2) &= \alpha\varepsilon \int_{\mathbb{R}^n} (u_t u_{tt} - v_t v_{tt})(u - v)_t dx \\ &+ \beta\varepsilon \int_{\mathbb{R}^n} (\nabla u \nabla u_t - \nabla v \nabla v_t)(u - v)_t dx. \end{aligned} \quad (45)$$

For the first right hand side term in Eq. (45) we find

$$\begin{aligned}
\int_{\mathbb{R}^n} (u_t u_{tt} - v_t v_{tt})(u - v)_t dx &= \int_{\mathbb{R}^n} u_{tt}(u - v)_t^2 dx + \int_{\mathbb{R}^n} v_t(u - v)_{tt}(u - v)_t dx \\
&= \int_{\mathbb{R}^n} u_{tt}(u - v)_t^2 dx + \int_{\mathbb{R}^n} v_t \frac{1}{2} \frac{d}{dt} [(u - v)_t^2] dx \\
&= \int_{\mathbb{R}^n} u_{tt}(u - v)_t^2 dx + \frac{1}{2} \frac{d}{dt} \left[\int_{\mathbb{R}^n} v_t(u - v)_t^2 dx \right] - \frac{1}{2} \int_{\mathbb{R}^n} v_{tt}(u - v)_t^2 dx \\
&= \int_{\mathbb{R}^n} u_{tt}(u - v)_t^2 dx + \frac{1}{2} \frac{d}{dt} \left[\int_{\mathbb{R}^n} v_t(u - v)_t^2 dx \right] \\
&\quad + \frac{1}{2} \int_{\mathbb{R}^n} (u_{tt} - v_{tt})(u - v)_t^2 dx - \frac{1}{2} \int_{\mathbb{R}^n} u_{tt}(u - v)_t^2 dx \\
&= \frac{1}{2} \int_{\mathbb{R}^n} u_{tt}(u - v)_t^2 dx + \frac{1}{2} \frac{d}{dt} \left[\int_{\mathbb{R}^n} v_t(u - v)_t^2 dx \right] + \frac{1}{6} \frac{d}{dt} \left[\int_{\mathbb{R}^n} (u - v)_t^3 dx \right] \\
&= \frac{1}{2} \int_{\mathbb{R}^n} u_{tt}(u - v)_t^2 dx + \frac{d}{dt} \left[\int_{\mathbb{R}^n} \left(\frac{1}{6} u_t + \frac{1}{3} v_t \right) (u - v)_t^2 dx \right].
\end{aligned}$$

On one hand,

$$\left| \int_{\mathbb{R}^n} u_{tt}(u - v)_t^2 dx \right| \leq \|u_{tt}\|_{L^\infty} \|(u - v)_t\|_{L^2}^2,$$

and on the other hand, we can put the term $\alpha \varepsilon \frac{d}{dt} \left[\int_{\mathbb{R}^n} \left(\frac{1}{6} u_t + \frac{1}{3} v_t \right) (u - v)_t^2 dx \right]$ on the left hand side of Eq. (44) and combine it with $\frac{1}{2} \frac{d}{dt} (\|(u - v)_t\|_{L^2}^2)$, so that we obtain the term

$$\frac{1}{2} \frac{d}{dt} \int_{\mathbb{R}^n} A(u - v)_t^2 dx, \quad \text{with } A = 1 - \alpha \varepsilon \left(\frac{1}{3} u_t + \frac{2}{3} v_t \right).$$

We notice that with condition (5) on u_t and v_t (which keeps our model hyperbolic) we have $\frac{1}{2} \leq A(t) \leq \frac{3}{2}$ for $t < T^*$. Therefore, for the second term at the right hand side we find

$$\begin{aligned}
\int_{\mathbb{R}^n} (\nabla u \nabla u_t - \nabla v \nabla v_t)(u - v)_t dx &= \int_{\mathbb{R}^n} \nabla v_t (\nabla u - \nabla v)(u - v)_t dx \\
&\quad + \int_{\mathbb{R}^n} \nabla u (\nabla u - \nabla v)_t (u - v)_t dx \\
&= \int_{\mathbb{R}^n} \nabla v_t (\nabla u - \nabla v)(u - v)_t dx - \frac{1}{2} \int_{\mathbb{R}^n} \Delta u (u - v)_t^2 dx.
\end{aligned}$$

We estimate the obtained two terms:

$$\begin{aligned}
\left| \int_{\mathbb{R}^n} \nabla v_t (\nabla u - \nabla v)(u - v)_t dx \right| &\leq C \|\nabla v_t\|_{L^\infty} (\|\nabla(u - v)\|_{L^2}^2 + \|(u - v)_t\|_{L^2}^2), \\
\left| \int_{\mathbb{R}^n} \Delta u (u - v)_t^2 dx \right| &\leq \|\Delta u\|_{L^\infty} \|(u - v)_t\|_{L^2}^2.
\end{aligned}$$

Thus, we find the following estimate

$$\frac{1}{2} \frac{d}{dt} \left(\int_{\mathbb{R}^n} A(u-v)_t^2 + c^2 \|\nabla(u-v)\|_{L^2}^2 dx \right) \leq C\varepsilon \sup(\|u_{tt}\|_{L^\infty}, \|\Delta u\|_{L^\infty}, \|\nabla v_t\|_{L^\infty}) \left(\|(u-v)_t\|_{L^2}^2 + \|\nabla(u-v)\|_{L^2}^2 \right).$$

Applying the Gronwall Lemma, as $\frac{1}{2} \leq A(t) \leq \frac{3}{2}$ for $t < T^*$, from the last estimate we have

$$\left(\|(u-v)_t\|_{L^2}^2 + \|\nabla(u-v)\|_{L^2}^2 \right)(t) \leq C_1 \exp\left(C_2\varepsilon \int_0^t \sup(\|u_{tt}\|_{L^\infty}, \|\Delta u\|_{L^\infty}, \|\nabla v_t\|_{L^\infty}) ds\right) \cdot \left(\|u_1 - v_1\|_{L^2}^2 + \|\nabla(u_0 - v_0)\|_{L^2}^2 \right).$$

With the hypothesis that $\|\nabla v_t\|_{L^\infty}$ is bounded on $[0, T^*[$, we obtain the result of Point 4 in Theorem 1 for all $t < T^*$.

Remark 2 *It is easy to verify that the same stability estimate also holds for $\nu > 0$: by the adding of the term $-\nu\Delta(u-v)_t$ to the equation, it gives a positive term $+\nu \int_0^t |\nabla(u-v)_t|^2 dx$ in the right hand side of the estimate and can be simply omitted.*

4 Well-posedness for the viscous case

4.1 Proof of Point 1 of Theorem 2

Let us show the global well-posedness, of the solution of the Cauchy problem (1)-(2). We start with the study of the linear problem, associated to the Kuznetsov equation.

Theorem 5 *Let $s \geq 0$ and X be the space defined in Point 1 of Theorem 2. Then the system*

$$\begin{cases} u_{tt} + c^2\Delta u + \varepsilon\nu\Delta u_t = f, \\ u(0) = u_0, \quad u_t(0) = u_1 \end{cases} \quad (46)$$

has a unique solution $u \in X$, if and only if $f \in L^2(\mathbb{R}_t^+; H^s)$, $u_0 \in H^{s+2}$ and $u_1 \in H^{s+1}$. Moreover it holds the following a priori estimate

$$\|u\|_X \leq C \left(\|f\|_{L^2(\mathbb{R}_t^+; H^s)} + \|u_0\|_{H^{s+2}} + \|u_1\|_{H^{s+1}} \right) \quad (47)$$

with $\|u\|_X := \|u\|_{H^2(\mathbb{R}_t^+; H^s)} + \|u\|_{L^2(\mathbb{R}_t^+; H^{s+2})} + \|u_t\|_{L^2(\mathbb{R}_t^+; H^{s+2})}$.

Proof : First we take $f \in L^2(\mathbb{R}_t^+; H^s)$, $u_0 \in H^{s+2}$ and $u_1 \in H^{s+1}$. We use the ideas of [6] (see Eq. (4.26)). For the sake of clarity, let us take $s = 0$. We take the inner product in $L^2(\mathbb{R}^n)$ of the equation with $-\Delta u_t$ and integrate by parts:

$$\frac{1}{2} \frac{d}{dt} \left(\|\nabla u_t\|_{L^2(\mathbb{R}^n)}^2 + c^2 \|\Delta u\|_{L^2(\mathbb{R}^n)}^2 \right) + \varepsilon\nu \|\Delta u_t\|_{L^2(\mathbb{R}^n)}^2 = - \int_{\mathbb{R}^n} f \Delta u_t dx.$$

Using the Young inequality and integrating over $[0, t]$, we find

$$\begin{aligned} \frac{1}{2} \left(\|\nabla u_t\|_{L^2(\mathbb{R}^n)}^2 + c^2 \|\Delta u\|_{L^2(\mathbb{R}^n)}^2 \right) + \frac{\varepsilon \nu}{2} \int_0^t \|\Delta u_s\|_{L^2(\mathbb{R}^n)}^2 ds \\ \leq \frac{1}{2} \|\nabla u_1\|_{L^2(\mathbb{R}^n)}^2 + \frac{1}{2} \|\Delta u_0\|_{L^2(\mathbb{R}^n)}^2 + \frac{1}{2\varepsilon \nu} \int_0^t \int_{\mathbb{R}^n} |f|^2 dx ds. \end{aligned} \quad (48)$$

Since $f \in L^2(\mathbb{R}_t^+ \times \mathbb{R}^n)$ and $(u_0, u_1) \in H^2(\mathbb{R}^n) \times H^1(\mathbb{R}^n)$, the last estimate implies that

$$\int_0^{+\infty} \int_{\mathbb{R}^n} |\Delta u_s|^2 dx ds < +\infty.$$

Since the domain of $-\Delta$ is H^2 , we obtain that

$$u, u_t \in L^2(\mathbb{R}_t^+; H^2), \quad \text{and} \quad u_{tt} \in L^2(\mathbb{R}_t^+ \times \mathbb{R}^n),$$

and hence, $u \in X$ for $s = 0$. For $s > 0$, as the equation is linear, we perform the same proof, using the fact that, the operator $\Lambda = (1 - \Delta)^{\frac{1}{2}}$, defined by its Fourier transform by the formula $\widehat{(\Lambda u)}(\zeta) = (1 + |\zeta|^2)^{\frac{1}{2}} \hat{u}(\zeta)$, relies the norm of H^s with the L^2 -norm:

$$\Lambda^s = (1 - \Delta)^{\frac{s}{2}}, \quad \|u\|_{H^s} = \|\Lambda^s u\|_{L^2}. \quad (49)$$

The unicity of u follows from the linearity of the operator and the unicity of the solution of system (46) in the case $f = 0$ (see [9]).

Conversely, if $u \in X$ solution of system (46), this implies that

$$u \in C(\mathbb{R}_t^+; H^{s+2}(\mathbb{R}^n)) \quad \text{and} \quad u_t \in H^1(\mathbb{R}_t^+; H^s(\mathbb{R}^n)) \cap L^2(\mathbb{R}_t^+; H^{s+2}(\mathbb{R}^n)).$$

Thanks to Theorem III.4.10.2 in [3], it follows that $u_t \in C(\mathbb{R}_t^+; H^{s+1}(\mathbb{R}^n))$. Then we have $u(0) \in H^{s+2}$, $u_t(0) \in H^{s+1}$. Moreover, it reads directly from the definition of X , that $f \in L^2(\mathbb{R}_t^+; H^s)$ for $u \in X$.

The a priori estimate follows from the closed graph theorem. \square

Let us notice that Theorem 5 states that problem (46) has L^2 -maximal regularity (see [5] Definition 2.1) on \mathbb{R}_t^+ . To prove the main result of the global well-posedness of the Cauchy problem for the Kuznetsov equation we use the following theorem from [20]:

Theorem 6 *Let X be a Banach space, let Y be a separable topological vector space, let $L : X \rightarrow Y$ be a linear continuous operator, let U be the open unit ball in X , let $P_{LU} : LX \rightarrow [0, \infty[$ be the Minkowski functional of the set LU , and let $\Phi : X \rightarrow LX$ be a mapping satisfying the condition*

$$P_{LU}(\Phi(x) - \Phi(\bar{x})) \leq \Theta(r) \|x - \bar{x}\| \quad \text{for} \quad \|x - x_0\| \leq r, \quad \|\bar{x} - x_0\| \leq r$$

for some $x_0 \in X$, where $\Theta : [0, \infty[\rightarrow [0, \infty[$ is a monotone non-decreasing function. Set $b(r) = \max(1 - \Theta(r), 0)$ for $r \geq 0$.

Suppose that

$$w = \int_0^\infty b(r) dr \in]0, \infty], \quad r_* = \sup\{r \geq 0 \mid b(r) > 0\},$$

$$w(r) = \int_0^r b(t)dt \quad (r \geq 0) \quad \text{and} \quad f(x) = Lx + \Phi(x) \quad \text{for} \quad x \in X.$$

Then for any $r \in [0, r_*[$ and $y \in f(x_0) + w(r)LU$, there exists an $x \in x_0 + rU$ such that $f(x) = y$.

Remark 3 *If either L is injective or $\text{Ker}L$ has a topological complement E in X such that $L(E \cap U) = LU$, then the assertion of Theorem 6 follows from the contraction mapping principle [20]. In particular, if L is injective, then the solution is unique.*

Now, we have all elements to prove Point 1 of Theorem 2:

Proof :

For $u_0 \in H^{s+2}(\mathbb{R}^n)$ and $u_1 \in H^{s+1}(\mathbb{R}^n)$ let us denote by $u^* \in X$ the unique solution of the linear problem

$$\begin{cases} u_{tt}^* - c^2 \Delta u^* - \nu \varepsilon \Delta u_t^* = 0, \\ u^*(0) = u_0 \in H^{s+2}(\mathbb{R}^n), \quad u_t^*(0) = u_1 \in H^{s+1}(\mathbb{R}^n). \end{cases} \quad (50)$$

In addition, according to Theorem 5, we take

$$X := H^2(\mathbb{R}_t^+; H^s(\mathbb{R}^n)) \cap H^1(\mathbb{R}_t^+; H^{s+2}(\mathbb{R}^n)),$$

this time for $s > \frac{n}{2}$ (we need it to control the non-linear terms), and introduce the Banach spaces

$$X_0 := \{u \in X \mid u(0) = u_t(0) = 0\} \quad (51)$$

and $Y = L^2(\mathbb{R}_t^+; H^s(\mathbb{R}^n))$. Then by Theorem 5, the linear operator

$$L : X_0 \rightarrow Y, \quad u \in X_0 \mapsto L(u) := u_{tt} - c^2 \Delta u - \nu \varepsilon \Delta u_t \in Y,$$

is a bi-continuous isomorphism.

Let us now notice that if v is the unique solution of the non-linear Cauchy problem

$$\begin{cases} v_{tt} - c^2 \Delta v - \nu \varepsilon \Delta v_t - \alpha \varepsilon (v + u^*)_t (v + u^*)_{tt} - \beta \varepsilon \nabla(v + u^*) \cdot \nabla(v + u^*)_t = 0, \\ v(0) = 0, \quad v_t(0) = 0, \end{cases} \quad (52)$$

then $u = v + u^*$ is the unique solution of the Cauchy problem for the Kuznetsov equation (1)–(2). Let us prove the existence of a such v , using Theorem 6.

We suppose that $\|u^*\|_X \leq r$ and define for $v \in X_0$

$$\Phi(v) := \alpha \varepsilon (v + u^*)_t (v + u^*)_{tt} + \beta \varepsilon \nabla(v + u^*) \cdot \nabla(v + u^*)_t.$$

For w and z in X_0 such that $\|w\|_X \leq r$ and $\|z\|_X \leq r$, we estimate

$$\begin{aligned} \|\Phi(w) - \Phi(z)\|_Y &= \|\alpha \varepsilon (u_t^*(w - z))_{tt} + (w - z)_t u_{tt}^* + w_t w_{tt} - z_t z_{tt} \\ &\quad + \beta \varepsilon (\nabla u^* \nabla(w - z)_t + \nabla(w - z) \nabla u_t^* + \nabla w \nabla w_t - \nabla z \nabla z_t)\|_Y \\ &= \|\alpha \varepsilon (u_t^*(w - z))_{tt} + (w - z)_t u_{tt}^* + w_t (w - z)_{tt} + (w - z)_t z_{tt} \\ &\quad + \beta \varepsilon (\nabla u^* \nabla(w - z)_t + \nabla(w - z) \nabla u_t^* + \nabla w \nabla(w - z)_t + \nabla(w - z) \nabla z_t)\|_Y \end{aligned}$$

by applying the triangular inequality

$$\begin{aligned} \|\Phi(w) - \Phi(z)\|_Y &\leq \alpha\varepsilon \left(\|u_t^*(w-z)_{tt}\|_Y + \|(w-z)_t u_{tt}^*\|_Y \right. \\ &\quad \left. + \|w_t(w-z)_{tt}\|_Y + \|(w-z)_t z_{tt}\|_Y \right) \\ &\quad + \beta\varepsilon \left(\|\nabla u^* \nabla(w-z)_t\|_Y + \|\nabla(w-z) \nabla u_t^*\|_Y \right. \\ &\quad \left. + \|\nabla w \nabla(w-z)_t\|_Y + \|\nabla(w-z) \nabla z_t\|_Y \right). \end{aligned}$$

Now, for all a and b in X with $s \geq s_0 > \frac{n}{2}$ it holds

$$\begin{aligned} \|a_t b_{tt}\|_Y &\leq \|a_t\|_{L^\infty(\mathbb{R}_t^+ \times \mathbb{R}^n)} \|b_{tt}\|_Y \\ &\leq C_{H^1(\mathbb{R}_t^+; H^{s_0}) \rightarrow L^\infty(\mathbb{R}_t^+ \times \mathbb{R}^n)} \|a_t\|_{H^1(0, +\infty; H^s)} \|b\|_X \\ &\leq C_{H^1(\mathbb{R}_t^+; H^{s_0}) \rightarrow L^\infty(\mathbb{R}_t^+ \times \mathbb{R}^n)} \|a\|_X \|b\|_X, \end{aligned}$$

where $C_{H^1(\mathbb{R}_t^+; H^{s_0}) \rightarrow L^\infty(\mathbb{R}_t^+ \times \mathbb{R}^n)}$ is the embedding constant of $H^1(\mathbb{R}_t^+; H^{s_0})$ in $L^\infty(\mathbb{R}_t^+ \times \mathbb{R}^n)$, independent on s , but depending only on the dimension n . In the same way, for all a and b in X it holds

$$\|\nabla a \nabla b_t\|_Y \leq C_{H^1(\mathbb{R}_t^+; H^{s_0}) \rightarrow L^\infty(\mathbb{R}_t^+ \times \mathbb{R}^n)} \|a\|_X \|b\|_X.$$

Taking a and b equal to u^* , w , z or $w-z$, as $\|u^*\|_X \leq r$, $\|w\|_X \leq r$ and $\|z\|_X \leq r$, we obtain

$$\|\Phi(w) - \Phi(z)\|_Y \leq 4(\alpha + \beta) C_{H^1(\mathbb{R}_t^+; H^{s_0}) \rightarrow L^\infty(\mathbb{R}_t^+ \times \mathbb{R}^n)} \varepsilon r \|w - z\|_X.$$

By the the fact that L is a bi-continuous isomorphism, there exists a minimal constant $C_\varepsilon = O\left(\frac{1}{\varepsilon^\nu}\right) > 0$ (coming from the inequality $C_0 \varepsilon^\nu \|u\|_X^2 \leq \|f\|_Y \|u\|_X$ for u , a solution of the linear problem (46) with homogeneous initial data [for a maximal constant $C_0 = O(1) > 0$]) such that

$$\forall u \in X_0 \quad \|u\|_X \leq C_\varepsilon \|Lu\|_Y.$$

Hence, for all $f \in Y$

$$P_{LU_{X_0}}(f) \leq C_\varepsilon P_{U_Y}(f) = C_\varepsilon \|f\|_Y.$$

Then we find for w and z in X_0 , such that $\|w\|_X \leq r$, $\|z\|_X \leq r$, and also with $\|u^*\|_X \leq r$, that

$$P_{LU_{X_0}}(\Phi(w) - \Phi(z)) \leq \Theta(r) \|w - z\|_X$$

with $\Theta(r) := 4C_\varepsilon(\alpha + \beta) C_{H^1(\mathbb{R}_t^+; H^{s_0}) \rightarrow L^\infty(\mathbb{R}_t^+ \times \mathbb{R}^n)} \varepsilon r$. Thus we apply Theorem 6 with $f(x) = L(x) - \Phi(x)$ and $x_0 = 0$. Therefore, knowing that $C_\varepsilon = \frac{C_0}{\varepsilon^\nu}$, we have, that for all $r \in [0, r_*[$ with

$$r_* = \frac{\nu}{4C_0(\alpha + \beta) C_{H^1(\mathbb{R}_t^+; H^{s_0}) \rightarrow L^\infty(\mathbb{R}_t^+ \times \mathbb{R}^n)}} = O(1), \quad (53)$$

for all $y \in \Phi(0) + w(r)LU_{X_0} \subset Y$ with

$$w(r) = r - 2\frac{C_0}{\nu} C_{H^1(\mathbb{R}_t^+; H^{s_0}) \rightarrow L^\infty(\mathbb{R}_t^+ \times \mathbb{R}^n)} (\alpha + \beta) r^2,$$

there exists a unique $v \in 0 + rU_{X_0}$ such that $L(v) - \Phi(v) = y$. But, if we want that v be the solution of the non-linear Cauchy problem (52), then we need to impose $y = 0$ and thus, to ensure that $0 \in \Phi(0) + w(r)LU_{X_0}$. Since $-\frac{1}{w(r)}\Phi(0)$ is an element of Y and $LU_{X_0} = Y$, there exists a unique $z \in X_0$ such that

$$Lz = -\frac{1}{w(r)}\Phi(0). \quad (54)$$

Let us show that $\|z\|_X \leq 1$, what will implies that $0 \in \Phi(0) + w(r)LU_{X_0}$. Noticing that

$$\begin{aligned} \|\Phi(0)\|_Y &\leq \alpha\varepsilon\|v_t v_{tt}\|_Y + \beta\varepsilon\|\nabla v \nabla v_t\|_Y \\ &\leq (\alpha + \beta)\varepsilon C_{H^1(\mathbb{R}_t^+; H^{s_0}) \rightarrow L^\infty(\mathbb{R}_t^+ \times \mathbb{R}^n)} \|v\|_X^2 \\ &\leq (\alpha + \beta)\varepsilon C_{H^1(\mathbb{R}_t^+; H^{s_0}) \rightarrow L^\infty(\mathbb{R}_t^+ \times \mathbb{R}^n)} r^2 \end{aligned}$$

and using (54), we find

$$\begin{aligned} \|z\|_X &\leq C_\varepsilon \|Lz\|_Y = C_\varepsilon \frac{\|\Phi(0)\|_Y}{w(r)} \\ &\leq \frac{C_\varepsilon C_{H^1(\mathbb{R}_t^+; H^{s_0}) \rightarrow L^\infty(\mathbb{R}_t^+ \times \mathbb{R}^n)} (\alpha + \beta)\varepsilon r}{(1 - 2C_\varepsilon C_{H^1(\mathbb{R}_t^+; H^{s_0}) \rightarrow L^\infty(\mathbb{R}_t^+ \times \mathbb{R}^n)} (\alpha + \beta)\varepsilon r)} < \frac{1}{2}, \end{aligned}$$

as soon as $r < r^*$.

Consequently, $z \in U_{X_0}$ and $\Phi(0) + w(r)Lz = 0$.

Then we conclude that for all $r \in [0, r_*[$, if $\|u^*\|_X \leq r$, there exists a unique $v \in rU_{X_0}$ such that $L(v) - \Phi(v) = 0$, *i.e.* the solution of the non-linear Cauchy problem (52). Thanks to the maximal regularity and a priori estimate following from inequality (48) with $f = 0$, there exists a constant $C_1 = O(\varepsilon^0) > 0$, such that

$$\|u^*\|_X \leq \frac{C_1}{\sqrt{\nu\varepsilon}} (\|u_0\|_{H^{s+2}} + \|u_1\|_{H^{s+1}}).$$

Thus, for all $r \in [0, r_*[$ and $\|u_0\|_{H^{s+2}} + \|u_1\|_{H^{s+1}} \leq \frac{\sqrt{\nu\varepsilon}}{C_1} r$, the function $u = u^* + v \in X$ is the unique solution of the Cauchy problem for the Kuznetsov equation and $\|u\|_X \leq 2r$. \square

4.2 Proof of Point 2 of Theorem 2: case $n \geq 3$

Knowing the existence of a solution u of the Kuznetsov equation in

$$X = H^2(\mathbb{R}_t^+; H^s(\mathbb{R}^n)) \cap H^1(\mathbb{R}_t^+; H^{s+2}(\mathbb{R}^n)),$$

we notice that this directly implies that

$$u \in C(\mathbb{R}_t^+; H^{s+2}(\mathbb{R}^n)) \quad \text{and} \quad u_t \in H^1(\mathbb{R}_t^+; H^s(\mathbb{R}^n)) \cap L^2(\mathbb{R}_t^+; H^{s+2}(\mathbb{R}^n)).$$

By Theorem III.4.10.2 in [3], it implies that $u_t \in C(\mathbb{R}_t^+; H^{s+1}(\mathbb{R}^n))$, which gives that

$$u \in C^1(\mathbb{R}_t^+; H^{s+1}(\mathbb{R}^n)) \cap C(\mathbb{R}_t^+; H^{s+2}(\mathbb{R}^n))$$

with, this time with the help of the Kuznetsov equation, $u_{tt} \in C(\mathbb{R}_t^+; H^{s-1}(\mathbb{R}^n))$. Consequently, in the viscous case the regularity of the time derivatives of the order greater than two of the solutions differs from the regularity, obtained in Section 3 for the non-viscous case. Thus we have to consider estimates with different energies: the energy $E_{\frac{m}{2}}[u](t)$, defined in Eq. (10), and the energy

$$S_{\frac{m}{2}}[u](t) = \sum_{i=1}^{\frac{m}{2}+1} \|\nabla \partial_t^i u(t)\|_{H^{m-2(i-1)}}^2, \quad (55)$$

defined, as $E_{\frac{m}{2}}[u](t)$, for $m \in \mathbb{N}$ and m even, which respect to the obtained regularity of u and its derivatives.

Lemma 1 *Let $n \in \mathbb{N}^*$, $n \geq 3$, $m \in \mathbb{N}$, and u be the solution of problem (1)-(2). Then for $m \geq [\frac{n}{2} + 3]$, m even, and all multi-index $A = (A_0, A_1, \dots, A_n)$ with $|A| - A_0 \leq m - 2A_0$ it holds*

$$\begin{aligned} \frac{d}{dt} \left(\int_{\mathbb{R}^n} ((1 - \alpha \varepsilon u_t)(D^A u_t)^2 + c^2 (\nabla D^A u)^2)(s, x) dx \right) \\ + 2\nu \varepsilon \int_{\mathbb{R}^n} (\nabla D^A u_t)^2(s, x) dx \\ \leq C_m \max(\alpha, \beta) \varepsilon \sqrt{E_{\frac{m}{2}}[u](s)} S_{\frac{m}{2}}[u](s) \end{aligned} \quad (56)$$

with a constant $C_m > 0$, depending only on m and on the dimension n .

Proof : Following notations of Proposition 1, we redefine

$$L_u v := v_{tt} - c^2 \Delta v - \nu \varepsilon \Delta v_t - \alpha \varepsilon u_t v_{tt} - \beta \varepsilon \nabla u \nabla v_t,$$

where u is the solution of problem (1). For this new $L_u v$ with the additional term $\nu \varepsilon \Delta v_t$, we have a modified version of relation (24)

$$\frac{d}{dt} \int_{\mathbb{R}^n} I[v](t, x) dx + 2\nu \varepsilon \int_{\mathbb{R}^n} (\nabla v_t)^2 = \int_{\mathbb{R}^n} J[v](t, x) dx, \quad (57)$$

where $I[v]$ and $J[v]$ are defined in Eqs. (22)–(23). We still take $v = D^A u$ with $A = (A_0, A_1, \dots, A_n)$, but this time $|A| - A_0 \leq m - 2A_0$ and m is even. Then we just need to show

$$\left| \int_{\mathbb{R}^n} J[D^A u](t, x) dx \right| \leq \varepsilon C_m \max(\alpha, \beta) \sqrt{E_{\frac{m}{2}}[u](t)} S_{\frac{m}{2}}[u](t). \quad (58)$$

For $n \geq 3$, $m \geq [\frac{n}{2} + 3]$ and m even, we have, thanks to the Hölder inequality,

$$\int_{\mathbb{R}^n} |u_{tt} (D^A u_t)^2| dx \leq \|u_{tt}\|_{L^{\frac{n}{2}}} \|D^A u_t\|_{L^{\frac{2n}{n-2}}}^2.$$

Noticing, that, thanks to Ref. [1] Theorem 7.57 p. 228, for $s > \frac{n}{2}$ there hold the continuous embeddings $H^s(\mathbb{R}^n) \subset C_B^0(\mathbb{R}^n) \subset L^{\frac{n}{2}}(\mathbb{R}^n)$ (where C_B^0 is the Banach space of bounded continuous functions equal to zero at the infinity), we can write for $m \geq [\frac{n}{2} + 3]$

$$\|u_{tt}\|_{L^{\frac{n}{2}}(\mathbb{R}^n)} \leq C \|u_{tt}\|_{H^{[\frac{m}{2}+1]}(\mathbb{R}^n)} \leq C \|u_{tt}\|_{H^{m-2}(\mathbb{R}^n)} \leq C \sqrt{E_{\frac{m}{2}}[u]}. \quad (59)$$

In addition, with the help of the Gagliardo-Nirenberg-Sobolev inequality

$$\|v\|_{L^{\frac{2n}{n-2}}(\mathbb{R}^n)} \leq C\|\nabla v\|_{L^2(\mathbb{R}^n)}, \quad (60)$$

we also have

$$\|D^A u_t\|_{L^{\frac{2n}{n-2}}} \leq C\|\nabla D^A u_t\|_{L^2} \leq C\|\nabla D_t^{A_0+1} u\|_{H^{|A|-A_0}}.$$

With the hypothesis that $|A| - A_0 \leq m - 2A_0$, there hold $2A_0 \leq m$ and

$$\|\nabla D_t^{A_0+1} u\|_{H^{|A|-A_0}} \leq \|\nabla D_t^{A_0+1} u\|_{H^{m-2A_0}}.$$

Therefore, all norms $\|\nabla D_t^{A_0+1} u\|_{H^{m-2A_0}}^2$, for the chosen n , m and A_0 , are present in $S_{\frac{m}{2}}$. Hence, we find

$$\int_{\mathbb{R}^n} |u_{tt}(D^A u_t)^2| dx \leq C\|u_{tt}\|_{H^{m-2}} \|\nabla D^A u_t\|_{L^2}^2 \leq C\sqrt{E_{\frac{m}{2}}[u]} S_{\frac{m}{2}}[u], \quad (61)$$

and in the same way,

$$\begin{aligned} \int_{\mathbb{R}^n} |\Delta u(D^A u_t)^2| dx &\leq \|\Delta u\|_{L^{\frac{n}{2}}} \|D^A u_t\|_{L^{\frac{2n}{n-2}}}^2 \leq C\|\Delta u\|_{H^{[\frac{n}{2}+1]}} \|\nabla D^A u_t\|_{L^2}^2 \\ &\leq C\sqrt{E_{\frac{m}{2}}[u]} S_{\frac{m}{2}}[u]. \end{aligned}$$

To calculate $L_u D^A u$ we use expression (28) with multi-indexes A^{j1} and A^{j2} satisfying (29). As in Proposition 1, without loss of generality, we consider two multi-indexes A^1 and A^2 with the same properties (29). We perform two steps:

Step 1: we prove

$$\int_{\mathbb{R}^n} |D^{A^1} u_t D^{A^2} u_t D^A u_t| dx \leq C\sqrt{E_{\frac{m}{2}}[u]} S_{\frac{m}{2}}[u], \quad (62)$$

Step 2: we prove

$$\int_{\mathbb{R}^n} |D^{A^1} \partial_{x_i} u D^{A^2} \partial_{x_i} u D^A u_t| dx \leq C\sqrt{E_{\frac{m}{2}}[u]} S_{\frac{m}{2}}[u]. \quad (63)$$

Step 1. Thanks to properties (29) of A_1 and A_2 and to the symmetry of the general case

$$\int_{\mathbb{R}^n} |(D_t^{A_0^1} D_x^{(A_1^1, \dots, A_n^1)} u_t)(D_t^{A_0^2} D_x^{(A_1^2, \dots, A_n^2)} u_t)(D^A u_t)| dx,$$

we divide our proof on three typical cases:

Case 1: $|A^1| - A_0^1 \geq 0$, $A_0^1 \geq 0$, $|A^2| - A_0^2 > 0$ and $A_0^2 > 0$, *i.e.* a non trivial presence of $D_t^{A_0^2}$ and $D_x^{(A_1^2, \dots, A_n^2)}$ is imposed,

Case 2: $|A^1| - A_0^1 = 0$, $A_0^1 > 0$, $|A^2| - A_0^2 > 0$ and $A_0^2 = 0$, *i.e.* we consider the integrals of the form $\int_{\mathbb{R}^n} |(D_t^{A_0^1} u_t)(D_x^{(A_1^2, \dots, A_n^2)} u_t)(D^A u_t)| dx$,

Case 3: $|A^1| - A_0^1 = 0$, $A_0^1 > 0$, $|A^2| - A_0^2 = 0$ and $A_0^2 > 0$, *i.e.* we consider only non-trivial time derivatives $\int_{\mathbb{R}^n} |(D_t^{A_0^1} u_t)(D_t^{A_0^2} u_t)(D^A u_t)| dx$.

Step 1, Case 1. By the generalized Hölder inequality with $\frac{1}{p} + \frac{1}{q} = \frac{n+2}{2n}$, we have

$$\int_{\mathbb{R}^n} |D^{A^1} u_t D^{A^2} u_t D^A u_t| dx \leq \|D^{A^1} u_t\|_{L^p} \|D^{A^2} u_t\|_{L^q} \|D^A u_t\|_{L^{\frac{2n}{n-2}}}.$$

By the Sobolev embeddings (31) of $H^{m_1} \subset L^p$ and $H^{m_2} \subset L^q$ with $m_1 + m_2 = \frac{n}{2} - 1$ and $0 < m_1 < \frac{n}{2} - 1$, we find

$$\int_{\mathbb{R}^n} |D^{A^1} u_t D^{A^2} u_t D^A u_t| dx \leq C \|D^{A^1} u_t\|_{H^{m_1}} \|D^{A^2} u_t\|_{H^{m_2}} \|\nabla D^A u_t\|_{L^2},$$

where we have also applied the Gagliardo-Nirenberg-Sobolev inequality (60). Hence,

$$\int_{\mathbb{R}^n} |D^{A^1} u_t D^{A^2} u_t D^A u_t| dx \leq C \|\partial_t^{A_0^1} u_t\|_{H^{m_1+|A^1|-A_0^1}} \|\nabla \partial_t^{A_0^2} u_t\|_{H^{m_2+|A^2|-A_0^2-1}} S_{\frac{m}{2}}[u]^{\frac{1}{2}}. \quad (64)$$

Now we are looking for $0 < m_1 < \frac{n}{2} - 1$, such that

$$\begin{cases} m_1 + |A^1| - A_0^1 \leq m - 2A_0^1, \\ m_2 + |A^2| - A_0^2 - 1 \leq m - 2A_0^2, \end{cases} \quad (65)$$

in order to have

$$\|\partial_t^{A_0^1} u_t\|_{H^{m_1+|A^1|-A_0^1}} \leq \sqrt{E_{\frac{m}{2}}[u]} \quad \text{and} \quad \|\nabla \partial_t^{A_0^2} u_t\|_{H^{m_2+|A^2|-A_0^2-1}} \leq \sqrt{S_{\frac{m}{2}}[u]}. \quad (66)$$

Since $m_2 = \frac{n}{2} - 1 - m_1$, and by (29), $|A^2| = |A| + 1 - |A^1|$ and $A_0^2 = A_0 + 1 - A_0^1$, system (65) is equivalent to

$$\begin{cases} m_1 + |A^1| + A_0^1 \leq m, \\ \frac{n}{2} - 1 - m_1 + |A| + 1 - |A^1| + A_0 + 1 - A_0^1 - 1 \leq m. \end{cases}$$

The last system, thanks to $|A| + A_0 \leq m$, corresponding to the assumptions of the Proposition, is satisfied if

$$\frac{n}{2} \leq m_1 + |A^1| + A_0^1 \leq m.$$

Using (29), we find that

$$|A^1| + A_0^1 = |A| + A_0 + 2 - (|A^2| + A_0^2).$$

Therefore, since for Case 1 $|A^2| \geq 2$ and $A_0^2 \geq 1$, recalling that (again by (29)) $|A| + A_0 \leq m$, we obtain

$$1 \leq |A^1| + A_0^1 \leq m - 1.$$

Thus, we distinguish three sub-cases:

For $n \geq 3$, $\frac{n}{2} \leq |A^1| + A_0^1 \leq m - 1$: taking $m_1 = \frac{1}{4}$, we obtain (66).

For $n \geq 5$, $2 \leq |A^1| + A_0^1 < \frac{n}{2}$: as $m \geq [\frac{n}{2} + 3]$, it is sufficient to take $m_1 = \frac{n}{2} - (|A^1| + A_0^1)$.

For $n \geq 3$, $|A^1| + A_0^1 = 1$: instead of finding m_1 , we notice, that we have only two possibility: either $D^{A^1} = \partial_t$ and $A^2 = A$, which gives estimate (61), or $D^{A^1} = \partial_{x_i}$ with $A_0^2 = A_0 + 1$ and $|A^2| - A_0^2 = |A| - A_0 - 1 > 0$. For the last case, by the generalized Hölder inequality, we have

$$\int_{\mathbb{R}^n} |\partial_{x_i} u_t D^{A^2} u_t D^A u_t| dx \leq \|\partial_{x_i} u_t\|_{L^n} \|D^{A^2} u_t\|_{L^2} \|D^A u_t\|_{L^{\frac{2n}{n-2}}}. \quad (67)$$

For $m \geq [\frac{n}{2} + 3]$ the first norm in Eq. (67) can be estimated using the continuous embedding $H^s(\mathbb{R}^n) \subset L^n(\mathbb{R}^n)$ holding for $s > \frac{n}{2}$:

$$\|\partial_{x_i} u_t\|_{L^n} \leq C \|\partial_{x_i} u_t\|_{H^{[\frac{n}{2}+1]}} \leq C \|u_t\|_{H^{m-1}} \leq C \sqrt{E_{\frac{m}{2}}[u]}.$$

With the help of the Gagliardo-Nirenberg-Sobolev inequality (60), we also estimate the second norm in (67)

$$\|D^A u_t\|_{L^{\frac{2n}{n-2}}} \leq C \|\nabla D^A u_t\|_{L^2} \leq C \sqrt{S_{\frac{m}{2}}[u]}, \quad (68)$$

and for the last one we directly have

$$\|D^{A^2} u_t\|_{L^2} \leq \|\nabla \partial_t^{A_0+2} u\|_{H^{|A|-A_0-2}} \leq \|\nabla \partial_t^{A_0+2} u\|_{H^{m-2A_0-2}} \leq \sqrt{S_{\frac{m}{2}}[u]}.$$

Thus we obtain as previously estimate (62) of Step 1.

This permits to conclude Case 1 of Step 1.

Step 1, Case 2. We have $|A^1| - A_0^1 = 0$, $A_0^1 > 0$, $|A^2| - A_0^2 > 0$ and $A_0^2 = 0$. Therefore, by (29), $A_0^1 = 1 + A_0$, and, updating (64), we directly have

$$\int_{\mathbb{R}^n} |D_t^{A_0^1} u_t D_x^{(A_1^2, \dots, A_n^2)} u_t D^A u_t| dx \leq C \|\partial_t^{A_0+1} u_t\|_{H^{m_1}} \|\nabla u_t\|_{H^{m_2+|A^2|-1}} S_{\frac{m}{2}}[u]^{\frac{1}{2}}$$

with $m_1 + m_2 = \frac{n}{2} - 1$, $0 < m_1 < \frac{n}{2} - 1$. Now we need to find m_1 , belonging to $]0, \frac{n}{2} - 1[$, such that

$$\begin{cases} m_1 \leq m - 2(A_0 + 1), \\ m_2 + |A^2| - 1 \leq m, \end{cases} \quad (69)$$

in order to have

$$\|\partial_t^{A_0+1} u_t\|_{H^{m_1}} \leq \sqrt{E_{\frac{m}{2}}[u]} \quad \text{and} \quad \|\nabla u_t\|_{H^{m_2+|A^2|-1}} \leq \sqrt{S_{\frac{m}{2}}[u]}.$$

From $1 + |A| = |A^1| + |A^2|$, by (29), with the relation $|A^1| = A_0^1 = 1 + A_0$ it follows that

$$|A^2| = |A| - A_0. \quad (70)$$

Therefore, as $m_2 = \frac{n}{2} - m_1 - 1$, system (69) is equivalent to

$$\begin{cases} m_1 + 2A_0 \leq m - 2, \\ \frac{n}{2} - 2 \leq m_1 + m - |A| + A_0. \end{cases}$$

By the assumption of the proposition

$$m - |A| + A_0 \geq 2A_0, \quad (71)$$

hence the last system is satisfied if we have m_1 such that

$$\frac{n}{2} - 2 \leq m_1 + 2A_0 \leq m - 2.$$

Knowing that $|A^2| > 0$ (by the assumption of Case 2), Eq. (70) implies that $|A| - A_0 > 0$. Thus, relation (71) gives $2A_0 \leq m - 1$, or more precisely

$$2A_0 \leq m - 2,$$

since m is even. So, a m_1 with $0 < m_1 < \frac{n}{2} - 1$ exists if $m - 2A_0 > 2$. Indeed, if $2A_0 < \frac{n}{2} - 2$ we can take $m_1 = \frac{n}{2} - 2 - 2A_0$, and if $m - 3 \geq 2A_0 \geq \frac{n}{2} - 2$ we can take $m_1 = \frac{1}{2}$.

Let us now consider the limit case $2A_0 = m - 2$. Then we have $|A^1| = A_0^1 = \frac{m}{2}$. Moreover, from (71) viewed, thanks to Eq. (70), as $|A^2| + 2A_0 \leq m$, follows that $1 \leq |A^2| \leq 2$. We apply the generalized Hölder inequality and estimate (60) to obtain

$$\begin{aligned} \int_{\mathbb{R}^n} |\partial_t^{\frac{m}{2}} u_t D_x^{(A_1^2, \dots, A_n^2)} u_t D^A u_t| dx &\leq \|\partial_t^{\frac{m}{2}} u_t\|_{L^2} \|D_x^{(A_1^2, \dots, A_n^2)} u_t\|_{L^n} \|D^A u_t\|_{L^{\frac{2n}{n-2}}} \\ &\leq C \|\partial_t^{\frac{m}{2}} u_t\|_{L^2} \|D_x^{(A_1^2, \dots, A_n^2)} u_t\|_{L^n} \sqrt{S_{\frac{m}{2}}[u]}. \end{aligned}$$

Moreover,

$$\|\partial_t^{\frac{m}{2}} u_t\|_{L^2} \leq \sqrt{E_{\frac{m}{2}}[u]}.$$

Using the continuity of the embedding $H^s(\mathbb{R}^n) \subset L^n(\mathbb{R}^n)$ for $s > \frac{n}{2}$, we also find for $m \geq [\frac{n}{2} + 3]$

$$\|D^{A^2} u_t\|_{L^n} \leq C \|D^{A^2} u_t\|_{H^{[\frac{n}{2}+1]}} \leq C \|\nabla u_t\|_{H^{[\frac{n}{2}+2]}} \leq C \|\nabla u_t\|_{H^m} \leq C \sqrt{S_{\frac{m}{2}}[u]}.$$

Hence, estimate (62) of Step 1 is also proved for Case 2.

Step 1, Case 3. Let us notice that thanks to relations (29), from $|A^1| = A_0^1$ and $|A^2| = A_0^2$ it follows $|A| = A_0$. We start as usual with the generalized Hölder inequality

$$\int_{\mathbb{R}^n} |D_t^{A_0^1} u_t D_t^{A_0^2} u_t D_t^{A_0} u_t| dx \leq \|D_t^{A_0^1} u_t\|_{L^p} \|D_t^{A_0^2} u_t\|_{L^q} \|D_t^{A_0} u_t\|_{L^{\frac{2n}{n-2}}}$$

with $\frac{1}{p} + \frac{1}{q} = \frac{n+2}{2n}$. Then we apply the Gagliardo-Nirenberg-Sobolev inequality (60) and its more general version, which can be viewed as the embedding of the Sobolev space $W_{q^*}^1(\mathbb{R}^n)$ in the Lebesgue space $L^q(\mathbb{R}^n)$ with $\frac{1}{q} = \frac{1}{q^*} - \frac{1}{n}$ and $1 \leq q^* < n$:

$$\int_{\mathbb{R}^n} |D_t^{A_0^1} u_t D_t^{A_0^2} u_t D_t^{A_0} u_t| dx \leq C \|D_t^{A_0^1} u_t\|_{L^p} \|\nabla D_t^{A_0^2} u_t\|_{L^{q^*}} \|\nabla D_t^{A_0} u_t\|_{L^2}$$

with $\frac{1}{p} + \frac{1}{q^*} = \frac{n+4}{2n}$. We notice that if we want to use the Sobolev embeddings (31) to L^p and to L^{q^*} , it is only possible if $\frac{1}{p}$ and $\frac{1}{q^*}$ are smaller than $\frac{1}{2}$, or equivalently, if $\frac{1}{p} + \frac{1}{q^*} = \frac{n+4}{2n} < 1$. Knowing that $\frac{n+4}{2n} < 1$ for $n \geq 5$, $\frac{n+4}{2n} > 1$ for $n = 3$ and $\frac{n+4}{2n} = 1$ for $n = 4$, we treat separately two cases: $n \geq 5$ and $n = 3, 4$.

For $n = 3$ or 4 , we choose $p = \frac{n}{2}$ and $q = \frac{2n}{n-2}$, implying $q^* = 2$. Thus, for $n = 3$ we use the continuous embedding $H^2(\mathbb{R}^3) \subset L^{\frac{3}{2}}(\mathbb{R}^3)$ [1] (since $2 > \frac{3}{2}$) and for $n = 4$ we use $H^2(\mathbb{R}^4) \subset L^2(\mathbb{R}^4)$ to obtain

$$\begin{aligned} \int_{\mathbb{R}^n} |D_t^{A_0^1} u_t D_t^{A_0^2} u_t D_t^{A_0} u_t| dx &\leq \|D_t^{A_0^1} u_t\|_{L^{\frac{n}{2}}} \|\nabla D_t^{A_0^2} u_t\|_{L^2} \|\nabla D_t^{A_0} u_t\|_{L^2} \\ &\leq C \|D_t^{A_0^1} u_t\|_{H^2} S_{\frac{m}{2}}[u]. \end{aligned}$$

If $m - 2A_0^1 \geq 2$, then we directly have

$$\|D^{A_0^1} u_t\|_{H^2} \leq \|D^{A_0^1} u_t\|_{H^{m-2A_0^1}} \leq \sqrt{E_{\frac{m}{2}}[u]}.$$

Recalling that m is even, and, by our assumption $|A^1| + A_0^1 \leq m$, $2A_0^1 \leq m$, there is only one additional possibility: $m - 2A_0^1 = 0$, *i.e.* $A_0^1 = \frac{m}{2}$.

For $A_0^1 = \frac{m}{2}$, thanks to (29) and the assumption $2A_0 \leq m$, we necessary have $|A_0^2| = 1$, and consequently, by (68),

$$\int_{\mathbb{R}^n} |\partial_t^{\frac{m}{2}} u_t u_{tt} \partial_t^{\frac{m}{2}} u_t| dx \leq C \|u_{tt}\|_{H^2} \|\partial_t^{\frac{m}{2}} u_t\|_{L^{\frac{2n}{n-2}}}^2 \leq \sqrt{E_{\frac{m}{2}}[u]} S_{\frac{m}{2}}[u].$$

Thus for $n = 3$ and $n = 4$ we find estimate (62).

Now, for $n \geq 5$, when $\frac{1}{p} + \frac{1}{q^*} = \frac{n+4}{2n} < 1$, we have

$$\begin{aligned} \int_{\mathbb{R}^n} |D_t^{A_0^1} u_t D_t^{A_0^2} u_t D_t^{A_0} u_t| dx &\leq C \|D_t^{A_0^1} u_t\|_{L^p} \|\nabla D_t^{A_0^2} u_t\|_{L^{q^*}} \|\nabla D_t^{A_0} u_t\|_{L^2} \\ &\leq C \|D_t^{A_0^1} u_t\|_{H^{m_1}} \|\nabla D_t^{A_0^2} u_t\|_{H^{m_2}} \sqrt{S_{\frac{m}{2}}[u]} \end{aligned}$$

with $m_1 + m_2 = \frac{n}{2} - 2$ and $0 < m_1 < \frac{n}{2} - 2$ by the Sobolev embeddings (31) which give us $H^{m_1} \subset L^p$ and $H^{m_2} \subset L^{q^*}$. We look for m_1 such that

$$m_1 \leq m - 2A_0^1, \quad m_2 \leq m - 2A_0^2 \quad (72)$$

in order to have

$$\|D_t^{A_0^1} u_t\|_{H^{m_1}} \leq \sqrt{E_{\frac{m}{2}}[u]} \quad \text{and} \quad \|\nabla D_t^{A_0^2} u_t\|_{H^{m_2}} \leq \sqrt{S_{\frac{m}{2}}[u]}.$$

As $m_2 = \frac{n}{2} - 2 - m_1$ and $A_0^2 = A_0 + 1 - A_0^1$, system (72) is equivalent to

$$\begin{cases} m_1 + 2A_0^1 \leq m, \\ \frac{n}{2} - 2 \leq m - 2A_0 + m_1 + 2A_0^1 - 2. \end{cases}$$

As $m - 2A_0 \geq 0$, it is sufficient to have m_1 such that

$$\frac{n}{2} \leq m_1 + 2A_0^1 \leq m$$

with $0 < m_1 < \frac{n}{2} - 2$ and $1 \leq A_0^1 \leq \frac{m}{2}$. If $2 \leq A_0^1 < \frac{n}{4}$ we can take $m_1 = \frac{n}{2} - 2A_0^1$. And if $\frac{n}{4} \leq A_0^1 \leq \frac{m}{2} - 1$ we can take $m_1 = \frac{1}{4}$.

If $A_0^1 = 1$, then necessary $A_0^2 = A_0$, and using estimates (59) and (68) we directly find

$$\int_{\mathbb{R}^n} |u_{tt} (D_t^{A_0} u_t)^2| dx \leq C \|u_{tt}\|_{L^{\frac{n}{2}}} \|D_t^{A_0} u_t\|_{L^{\frac{2n}{n-2}}}^2 \leq C \sqrt{E_{\frac{m}{2}}[u]} S_{\frac{m}{2}}[u].$$

If $A_0^1 = \frac{m}{2}$ we are in a symmetric case as $A_0^2 = 1$. This conclude the proof of Case 3 and of Step 1, *i.e.* of estimate (62).

Step 2. Let us show estimate (63). Thanks to properties (29) of A_1 and A_2 and to the symmetry of the general case

$$\int_{\mathbb{R}^n} |(D_t^{A_0} D_x^{(A_1^1, \dots, A_n^1)} u_{x_i})(D_t^{A_0^2} D_x^{(A_1^2, \dots, A_n^2)} u_{x_i})(D^A u_t)| dx,$$

we divide our proof on two typical cases:

Case 1: $|A^1| - A_0^1 \geq 0$, $A_0^1 > 0$, $|A^2| - A_0^2 \geq 0$ and $A_0^2 > 0$, *i.e.* a non trivial presence of $D_t^{A_0^1}$ and $D_t^{A_0^2}$ is imposed,

Case 2: $|A^1| - A_0^1 > 0$, $A_0^1 = 0$, $|A^2| - A_0^2 \geq 0$ and $A_0^2 > 0$, *i.e.* we consider the integrals of the form $\int_{\mathbb{R}^n} |(D_x^{A_1^1 + \dots + A_n^1} u_{x_i})(D_t^{A_0^2} D_x^{A_1^2 + \dots + A_n^2} u_{x_i})(D^A u_t)| dx$ with a non-trivial $D_t^{A_0^2}$.

Case 1. Using estimate $\|D^A u_t\|_{L^2} \leq \sqrt{E_{\frac{m}{2}}[u]}$, we have

$$\begin{aligned} & \int_{\mathbb{R}^n} |(D_t^{A_0^1} D_x^{(A_1^1, \dots, A_n^1)} u_{x_i})(D_t^{A_0^2} D_x^{(A_1^2, \dots, A_n^2)} u_{x_i})(D^A u_t)| dx \\ & \leq C \|\nabla \partial_t^{A_0^1} u\|_{H^{m_1 + |A^1| - A_0^1}} \|\nabla \partial_t^{A_0^2} u\|_{H^{m_2 + |A^2| - A_0^2}} \sqrt{E_{\frac{m}{2}}[u]} \end{aligned}$$

with $m_1 + m_2 = \frac{n}{2}$ and $0 < m_1 < \frac{n}{2}$.

Let us find m_1 with $0 < m_1 < \frac{n}{2}$ such that

$$\begin{cases} m_1 + |A^1| - A_0^1 \leq m - 2(A_0^1 - 1), \\ m_2 + |A^2| - A_0^2 \leq N - 2(A_0^2 - 1) \end{cases} \quad (73)$$

in order to have

$$\|\nabla \partial_t^{A_0^1} u\|_{H^{m_1+|A^1|-A_0^1}} \leq \sqrt{S_{\frac{m}{2}}[u]} \quad \text{and} \quad \|\nabla \partial_t^{A_0^2} u\|_{H^{m_2+|A^2|-A_0^2}} \leq \sqrt{S_{\frac{m}{2}}[u]}.$$

As $m_2 = \frac{n}{2} - m_1$, $|A^1| + |A^2| = |A| + 1$, and $A_0^1 + A_0^2 = A_0 + 1$, system (73) is equivalent to

$$\begin{cases} m_1 + |A^1| + A_0^1 \leq m + 2, \\ \frac{n}{2} + |A| + A_0 + 2 \leq m + 2 + m_1 + |A^1| + A_0^1. \end{cases}$$

By our assumption $|A| + A_0 \leq m$, and hence the last system is satisfied if m_1 verifies

$$\frac{n}{2} \leq m_1 + |A^1| + A_0^1 \leq m + 2.$$

In our case $A_0^1 > 0$, thus $2 \leq |A^1| + A_0^1 \leq m$, which implies the existence of a such m_1 with $0 < m_1 < \frac{n}{2}$. Indeed, if $m \geq |A^1| + A_0^1 \geq \frac{n}{2}$ we can take $m_1 = 1$, else if $2 \leq |A^1| + A_0^1 < \frac{n}{2}$ it is possible to take $m_1 = \frac{n}{2} - (|A^1| + A_0^1)$. This concludes Case 1 of Step 2.

Case 2. Thanks to (29), the conditions $|A^1| > 0$ with $A_0^1 = 0$ imply that $|A| - A_0 > 0$. Consequently, with $m_1 + m_2 = \frac{n}{2}$ and $0 < m_1 < \frac{n}{2}$ as in the previous case, we obtain

$$\begin{aligned} \int_{\mathbb{R}^n} |D_x^{A^1} \partial_{x_i} u D^{A^2} \partial_{x_i} u D^A u_t| dx &\leq C \|\nabla u\|_{H^{m_1+|A^1|}} \|\nabla \partial_t^{A_0^2} u\|_{H^{m_2+|A^2|-A_0^2}} \\ &\cdot \|\nabla \partial_t^{A_0} u_t\|_{H^{|A|-A_0-1}} \leq C \|\nabla u\|_{H^{m_1+|A^1|}} \|\nabla \partial_t^{A_0^2} u\|_{H^{m_2+|A^2|-A_0^2}} \sqrt{S_{\frac{m}{2}}[u]}. \end{aligned}$$

In the aim to have

$$\|\nabla u\|_{H^{m_1+|A^1|}} \leq \sqrt{E_{\frac{m}{2}}[u]} \quad \text{and} \quad \|\nabla \partial_t^{A_0^2} u\|_{H^{m_2+|A^2|-A_0^2}} \leq \sqrt{S_{\frac{m}{2}}[u]},$$

we need to find m_1 with $0 < m_1 < \frac{n}{2}$, such that

$$\begin{cases} m_1 + |A^1| \leq m, \\ m_2 + |A^2| - A_0^2 \leq m - 2(A_0^2 - 1). \end{cases}$$

As $m_2 = \frac{n}{2} - m_1$, $|A^2| = |A| + 1 - |A^1|$ and $A_0^2 = A_0 + 1$ it is equivalent to solve

$$\begin{cases} m_1 + |A^1| \leq m, \\ \frac{n}{2} - m_1 + |A| + 1 - |A^1| + A_0 + 1 - 2 \leq m. \end{cases}$$

As $m - |A| - A_0 \geq 0$, the last system is satisfied if m_1 verifies

$$\frac{n}{2} \leq m_1 + |A^1| \leq m.$$

By assumptions of this case $1 \leq |A^1| \leq m$, what guarantees the existence of such m_1 with $0 < m_1 < \frac{n}{2}$. Indeed, if $1 \leq |A^1| < \frac{n}{2}$, then we can take $m_1 = \frac{n}{2} - |A^1|$, and

if $|A^1| \leq m - 1$, then we can take $m_1 = \frac{1}{2}$. In the case $|A^1| = m$, corresponding to $D^{A^2} = \partial_t$, we directly obtain

$$\begin{aligned} \int_{\mathbb{R}^n} |D_x^{A^1} \partial_{x_i} u \partial_{x_i} u_t D^A u_t| dx &\leq C \|D_x^{A^1} \partial_{x_i} u\|_{L^2} \|\partial_{x_i} u_t\|_{L^n} \|D^A u_t\|_{L^{\frac{2n}{n-2}}} \\ &\leq C \|\nabla u\|_{H^m} \|\nabla u_t\|_{H^m} \|\nabla D^A u_t\|_{L^2} \leq C \sqrt{E_{\frac{m}{2}}[u]} S_{\frac{m}{2}}[u]. \end{aligned}$$

This completes the proof of Step 2 and hence the proof of estimate (63).

Thus, estimates (62) and (63) imply

$$\left| \int_{\mathbb{R}^n} L_u D^A u D^A u_t dx \right| \leq C \max(\alpha, \beta) \varepsilon \sqrt{E_{\frac{m}{2}}[u]} S_{\frac{m}{2}}[u],$$

from where follows (58). □

Thanks to Lemma 1, we have the following energy decreasing result:

Theorem 7 *Let $n \geq 3$, $m \in \mathbb{N}$ be even and $m \geq [\frac{n}{2} + 3]$. For $u_0 \in H^{m+1}(\mathbb{R}^n)$ and $u_1 \in H^m(\mathbb{R}^n)$, satisfying the smallness condition according to Point 1 of Theorem 2, there exists a unique global solution $u \in C^1(\mathbb{R}_t^+; H^{m-1}(\mathbb{R}^n)) \cap C(\mathbb{R}_t^+; H^m(\mathbb{R}^n))$ of problem (1)–(2) and the energy $E_{\frac{m}{2}}[u](0) < \infty$ is well-defined. Then*

1. *it holds the a priori estimate*

$$\frac{d}{dt} E(t) + \sqrt{2} \varepsilon S_{\frac{m}{2}}[u](t) \left(\sqrt{2} \nu - C_m \max(\alpha, \beta) \sqrt{E(t)} \right) \leq 0, \quad (74)$$

where, denoting by V the set of all multi-indexes $A = (A_0, A_1, \dots, A_n)$ with $|A| - A_0 \leq m - 2A_0$,

$$E(t) = \sum_{A \in V} \int_{\mathbb{R}^n} (1 - \alpha \varepsilon u_t) (D^A u_t)^2 + c^2 (\nabla D^A u)^2(t, x) dx.$$

2. *if in addition $\sqrt{E_{\frac{m}{2}}[u](0)} \leq \frac{\sqrt{2} \nu}{\sqrt{\frac{3}{2} + c^2} C_m \max(\alpha, \beta)} = O(\sqrt{\varepsilon})$, then*

$$\forall t \in \mathbb{R}^+, \quad E_{\frac{m}{2}}[u](t) \leq (3 + 2c^2) E_{\frac{m}{2}}[u](0) < +\infty. \quad (75)$$

Proof : We sum (56) on all $A \in V$ to obtain

$$\frac{d}{dt} E(t) + 2\nu \varepsilon S_{\frac{m}{2}}[u] \leq C_m \max(\alpha, \beta) \varepsilon \sqrt{E_{\frac{m}{2}}[u]} S_{\frac{m}{2}}[u].$$

While $\|u_t(t)\|_{L^\infty} \leq \frac{1}{2\alpha\varepsilon}$ it holds

$$\frac{1}{2} E_{\frac{m}{2}}[u](t) \leq E(t) \leq \left(\frac{3}{2} + c^2\right) E_{\frac{m}{2}}[u](t),$$

and consequently,

$$\frac{d}{dt}E(t) + 2\nu\varepsilon S_{\frac{m}{2}}[u](t) \leq \sqrt{2}C_m \max(\alpha, \beta)\varepsilon\sqrt{E(t)}S_{\frac{m}{2}}[u](t).$$

Thus, if for all time $\sqrt{E(t)} < \frac{\sqrt{2}\nu}{\max(\alpha, \beta)C_m}$, and in particular,

$$E(0) \leq \left(\frac{3}{2} + c^2\right)E_{\frac{m}{2}}[u](0) < 2\left(\frac{\nu}{C_m \max(\alpha, \beta)}\right)^2, \quad (76)$$

then we have the decreasing of E in time:

$$\frac{d}{dt}E(t) < 0 \quad \text{and} \quad E(t) \leq E(0).$$

Moreover, for all time $t \geq 0$

$$\begin{aligned} \|u_t(t)\|_{L^\infty} &\leq C_\infty \sqrt{E_{\frac{m}{2}}[u](t)} \leq C_\infty \sqrt{2}\sqrt{E(t)} \leq C_\infty \sqrt{2}\sqrt{E(0)} \\ &< 2C_\infty \frac{\nu}{C_m \max(\alpha, \beta)} < \frac{1}{2\alpha\varepsilon}. \end{aligned}$$

To be able to write $2C_\infty \frac{\nu}{C_m \max(\alpha, \beta)} < \frac{1}{2\alpha\varepsilon}$, we recall that, using the physical values of coefficients, $\varepsilon \ll 1$, $c^2 = O(\frac{1}{\varepsilon})$, $\alpha = \frac{\gamma-1}{c^2} < \beta = 2$, and consequently, as $\nu = O(1)$, the last inequality becomes

$$\frac{C_\infty}{C_m}\nu < \frac{1}{2\alpha\varepsilon},$$

which is obviously true in the case of $\varepsilon \ll 1$ (and, for instance, taking $C_m = 2C_\infty$). Hence, if Eq. (76) holds, then for all time $\|u_t(t)\|_{L^\infty} < \frac{1}{2\alpha\varepsilon}$ and the well-posedness of the Cauchy problem is ensured with the following energy estimate

$$E_{\frac{m}{2}}[u](t) \leq 2E(0) \leq (3 + 2c^2)E_{\frac{m}{2}}[u](0).$$

□

References

- [1] R. A. ADAMS, *Sobolev spaces*, vol. 65 of Pure and Applied Mathematics, Academic Press, New York, 1975.
- [2] S. ALINHAC, *A minicourse on global existence and blowup of classical solutions to multidimensional quasilinear wave equations*, in Journées “Équations aux Dérivées Partielles” (Forges-les-Eaux, 2002), Univ. Nantes, Nantes, 2002, pp. Exp. No. I, 33.
- [3] H. AMANN, *Linear and quasilinear parabolic problems. Vol. I*, vol. 89 of Monographs in Mathematics, Birkhäuser Boston, Inc., Boston, MA, 1995. Abstract linear theory.

- [4] H. BREZIS, *Analyse fonctionnelle : Théorie et applications*, Sciences SUP, 2005.
- [5] R. CHILL AND S. SRIVASTAVA, *L^p -maximal regularity for second order Cauchy problems*, Math. Z., 251 (2005), pp. 751–781.
- [6] M. GHISI, M. GOBBINO, AND A. HARAUX, *Local and global smoothing effects for some linear hyperbolic equations with a strong dissipation*, Trans. Amer. Math. Soc., 368 (2016), pp. 2039–2079.
- [7] M. HAMILTON AND D. BLACKSTOCK, *Nonlinear Acoustics*, Academic Press, 1998.
- [8] T. J. R. HUGHES, T. KATO, AND J. E. MARSDEN, *Well-posed quasi-linear second-order hyperbolic systems with applications to nonlinear elastodynamics and general relativity*, Arch. Rational Mech. Anal., 63 (1976), pp. 273–294 (1977).
- [9] R. IKEHATA, G. TODOROVA, AND B. YORDANOV, *Wave equations with strong damping in Hilbert spaces*, J. Differential Equations, 254 (2013), pp. 3352–3368.
- [10] F. JOHN, *Nonlinear wave equations, formation of singularities*, vol. 2 of University Lecture Series, American Mathematical Society, Providence, RI, 1990. Seventh Annual Pitcher Lectures delivered at Lehigh University, Bethlehem, Pennsylvania, April 1989.
- [11] P. M. JORDAN, *An analytical study of Kuznetsov's equation: diffusive solitons, shock formation, and solution bifurcation*, Phys. Lett. A, 326 (2004), pp. 77–84.
- [12] B. KALTENBACHER AND I. LASIECKA, *Well-posedness of the Westervelt and the Kuznetsov equation with nonhomogeneous Neumann boundary conditions*, Discrete Contin. Dyn. Syst., (2011), pp. 763–773.
- [13] —, *An analysis of nonhomogeneous Kuznetsov's equation: local and global well-posedness; exponential decay*, Math. Nachr., 285 (2012), pp. 295–321.
- [14] S. KLAINERMAN, *Uniform decay estimates and the Lorentz invariance of the classical wave equation*, Comm. Pure Appl. Math., 38 (1985), pp. 321–332.
- [15] —, *Remarks on the global sobolev inequalities in the minkowski space \mathbb{R}^{n+1}* , Comm. Pure Appl. Math., 40 (1987), pp. 111–117.
- [16] V. KUZNETSOV, *Equations of nonlinear acoustics*, Soviet Phys. Acoust., 16 (1971), pp. 467–470.
- [17] S. MEYER AND M. WILKE, *Global well-posedness and exponential stability for Kuznetsov's equation in L_p -spaces*, Evol. Equ. Control Theory, 2 (2013), pp. 365–378.
- [18] A. ROZANOVA-PIERRAT, *Approximation of a compressible Navier-Stokes system by non-linear acoustical models*, Proceedings of the International Conference "Days on Diffraction 2015", St. Petersburg, Russia, IEEE , St. Petersburg, Russia, 2015.

- [19] Y. SHIBATA, *On the rate of decay of solutions to linear viscoelastic equation*, Math. Methods Appl. Sci., 23 (2000), pp. 203–226.
- [20] M. F. SUKHININ, *On the solvability of the nonlinear stationary transport equation*, Teoret. Mat. Fiz., 103 (1995), pp. 23–31.
- [21] P. J. WESTERVELT, *Parametric acoustic array*, The Journal of the Acoustical Society of America, 35 (1963), pp. 535–537.