



HAL
open science

The ALMA-PILS survey: Formaldehyde deuteration in warm gas on small scales toward IRAS 16293-2422 B

M. V. Persson, J. K. Jørgensen, H. S. P. Müller, Audrey Coutens, E. F. van Dishoeck, V. Taquet, H. Calcutt, M. H. D. van Der Wiel, T. L. Bourke, S. Wampfler

► To cite this version:

M. V. Persson, J. K. Jørgensen, H. S. P. Müller, Audrey Coutens, E. F. van Dishoeck, et al.. The ALMA-PILS survey: Formaldehyde deuteration in warm gas on small scales toward IRAS 16293-2422 B. *Astronomy and Astrophysics - A&A*, 2018, 610, pp.A54. 10.1051/0004-6361/201731684 . hal-01650801v2

HAL Id: hal-01650801

<https://hal.science/hal-01650801v2>

Submitted on 1 Jun 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

The ALMA-PILS survey: Formaldehyde deuteration in warm gas on small scales toward IRAS 16293–2422 B

M. V. Persson^{1,2}, J. K. Jørgensen³, H. S. P. Müller⁴, A. Coutens⁵, E. F. van Dishoeck^{2,6}, V. Taquet⁷, H. Calcutt²,
M. H. D. van der Wiel⁸, T. L. Bourke⁹, and S. Wampfler¹⁰

¹ Department of Space, Earth and Environment, Chalmers University of Technology, Onsala Space Observatory, 439 92, Onsala, Sweden, e-mail: magnpe@chalmers.se

² Leiden Observatory, Leiden University, P.O. Box 9513, NL-2300 RA Leiden, The Netherlands

³ Centre for Star and Planet Formation, Niels Bohr Institute & Natural History Museum of Denmark, University of Copenhagen, Øster Voldgade 5–7, DK-1350, Copenhagen K, Denmark

⁴ I. Physikalisches Institut, Universität zu Köln, Zùlpicher Str. 77, 50937 Köln, Germany

⁵ Laboratoire d’astrophysique de Bordeaux, Univ. Bordeaux, CNRS, B18N, allée Geoffroy Saint-Hilaire, 33615 Pessac, France

⁶ Max-Planck Institute für extraterrestrische Physik (MPE), Giessenbachstrasse, 85748 Garching, Germany

⁷ INAF-Osservatorio Astrofisico di Arcetri, Largo E. Fermi 5, I-50125 Firenze, Italy

⁸ ASTRON Netherlands Institute for Radio Astronomy, PO Box 2, 7990 AA Dwingeloo, The Netherlands

⁹ SKA Organization, Jodrell Bank Observatory, Lower Withington, Macclesfield, Cheshire SK11 9DL, UK

¹⁰ Center for Space and Habitability (CSH) University of Bern Sidlerstrasse 5, CH-3012 Bern, Switzerland

Received XX July, 2017; accepted XX YY, 201Z

ABSTRACT

Context. The enhanced degrees of deuterium fractionation observed in envelopes around protostars demonstrate the importance of chemistry at low temperatures, relevant in pre- and protostellar cores. Formaldehyde is an important species in the formation of methanol and more complex molecules.

Aims. Here, we present the first study of formaldehyde deuteration on small scales around the prototypical low-mass protostar IRAS 16293–2422 using high spatial and spectral resolution Atacama Large Millimeter/submillimeter Array (ALMA) observations. We determine the excitation temperature, abundances and fractionation level of several formaldehyde isotopologues, including its deuterated forms.

Methods. Excitation temperature and column densities of formaldehyde in the gas close to one of the components of the binary are constrained through modeling of optically thin lines assuming local thermodynamical equilibrium. The abundance ratios are compared to results from previous single dish observations, astrochemical models and local ISM values.

Results. Numerous isotopologues of formaldehyde are detected, among them H₂C¹⁷O, and D₂¹³CO for the first time in the ISM. The large range of upper energy levels covered by the HDCO lines help constrain the excitation temperature to 106±13 K. Using the derived column densities, formaldehyde shows a deuterium fractionation of HDCO/H₂CO=6.5±1%, D₂CO/HDCO=12.8^{+3.3}_{-4.1}%, and D₂CO/H₂CO=0.6(4)±0.1%. The isotopic ratios derived are ¹⁶O/¹⁸O=805⁺⁴³₋₇₉, ¹⁸O/¹⁷O=3.2^{+0.2}_{-0.3} and ¹²C/¹³C=56⁺⁸₋₁₁.

Conclusions. The HDCO/H₂CO ratio is lower than found in previous studies, highlighting the uncertainties involved in interpreting single dish observations of the inner warm regions. The D₂CO/HDCO ratio is only slightly larger than the HDCO/H₂CO ratio. This is consistent with formaldehyde forming in the ice as soon as CO has frozen onto the grains, with most of the deuteration happening towards the end of the prestellar core phase. A comparison with available time-dependent chemical models indicates that the source is in the early Class 0 stage.

Key words. astrochemistry — stars: formation — stars: protostars — ISM: molecules — ISM: individual objects: IRAS 16293–2422

1. Introduction

Among molecular abundance ratios, the deuterium fractionation (D/H ratio) is commonly used to infer the formation conditions (e.g. Roberts & Millar 2000). In general, a high D/H ratio indicates a low temperature, and a low D/H ratio indicates a high temperature during formation. The deuterium-to-hydrogen ratio in the local interstellar medium (ISM) is 2.0±0.1×10⁻⁵ (Prodanović et al. 2010). Many molecules are formed on the surfaces of dust grains, and once it is cold enough for CO to freeze out onto the grains, the deuterium chemistry is enhanced. This means that any molecules formed after CO freeze-out are expected to have high levels of deuterium fractionation (e.g. Taquet et al. 2014). Single dish observations of various molecules to-

ward deeply embedded protostars have shown high deuterium fractions, above 10% (e.g. van Dishoeck et al. 1995; Parise et al. 2006). Optical depth and source size effects (beam dilution) are significant for single dish observations of lines of the main isotopologue which makes the uncertainty in the fractions large. High sensitivity and high resolution interferometric observations have the possibility to circumvent these problems since they probe smaller scales, close to the forming star where the temperature is high enough to sublimate the grain surface ices completely (i.e. $T \gtrsim 100$ K). The higher sensitivity makes it possible to detect weaker lines of minor isotopologues where optical depth effects are less important.

Observations of water have revealed a low deuterium fractionation (e.g. HDO/H₂O ratio) on small scales (warm gas)

in young protostellar envelopes. While the cold gas shows a HDO/H₂O ratio of a few %, the warm gas has a ratio closer to 0.1% (e.g. Persson et al. 2014; Coutens et al. 2012). Furthermore, the water deuterium fractionation of the warm gas in the inner region toward the protostar NGC 1333 IRAS 2A shows a D₂O/HDO ratio significantly higher than the HDO/H₂O ratio (seven times higher, Coutens et al. 2014). Furuya et al. (2016) explain the observed ratios by modeling the pre-stellar core physical and chemical evolution. The gas-phase deuteration processes are inefficient early on due to the higher H₂ o/p-ratio raising the destruction rate of the main seed molecule for deuteration processes, H₂D⁺ (Pagani et al. 1992). However, the deuteration processes through H₂D⁺ increase as soon as the o/p-ratio decreases and other molecules that destroy it freezes out (e.g. CO, Furuya et al. 2015). This and the low temperature results in higher gas-phase atomic D/H ratio which in turn causes more D atoms to accrete (mainly through H₂D⁺) on the grains (first pointed out by Tielens 1983). This drives the hydrogen-surface deuterium chemistry, however the total production rate of water is significantly decreased at this point, thus for water the deuteration is low. This explains both the low absolute deuteration that water has, and the relative ratios between the deuterated forms. Whether other molecules show similar trends is not yet clear. However many molecules are thought to form later, once CO has frozen onto the grains and could thus show different ratios in this model.

Formaldehyde is an important molecule, not only for constraining physical conditions in the gas of star-forming regions, but also as an intermediary in the formation path to complex organic molecules. H₂CO formation in ices was shown experimentally and models highlighted that it proceeds at a significantly higher rate in ices than in gas-phase reactions (Roberts et al. 2004). Thus it is thought that formaldehyde is mainly formed on the surface of dust grains once CO has frozen onto the grains through the hydrogen addition reactions CO + H → HCO and then HCO + H → H₂CO seen in the laboratory (Watanabe & Kouchi 2002; Fuchs et al. 2009). The deuterated forms of H₂CO can be obtained through subsequent substitution reactions involving D-atoms, i.e. H₂CO + D → HDCO+H and HDCO + D → D₂CO+H. They can also be obtained through abstraction and addition reactions through the HCO and DCO radical (e.g. Tielens 1983), which has been shown to be efficient at low temperature in ices (Hidaka et al. 2009). Naturally, just as with H₂CO there is also a direct channel from CO through DCO, i.e. the tunneling reaction CO + D → DCO, with subsequent addition reactions to form D₂CO and HDCO.

Assuming a grain surface formation route, Turner (1990) and later Charnley et al. (1997) estimated the relation between abundance ratios of the deuterated isotopologues of formaldehyde, assuming equal transmission probabilities for the CO+H and CO+D reaction channels and neglecting abstraction reactions, to be

$$\frac{D_2CO}{HDCO} = \frac{1}{4} \frac{HDCO}{H_2CO}. \quad (1)$$

While this was enough to explain the observed deuterium fractionation of formaldehyde at the time, it seems the situation is slightly more complex for deeply-embedded protostars, and that substitution and abstraction (with subsequent addition) reactions are important.

In this paper we study the deuterium fractionation of formaldehyde (H₂CO) on small solar-system scales toward one of the sources in the Class 0 multiple system IRAS 16293–2422, located in the ρ Ophiuchus star forming region at a distance of

120 pc (Loinard et al. 2008). The deuterium fractionation is determined through line analysis of various deuterated and non-deuterated isotopologues of formaldehyde. IRAS 16293–2422 is a protostellar binary with a separation of about 5'' (600 AU), where the SE source is referred to as "A" and the NW source as "B". Several studies have shown IRAS 16293–2422 to be a chemically rich source, with a wealth of complex organic molecular species (e.g. Bottinelli et al. 2004; Kuan et al. 2004; Bisschop et al. 2008; Jørgensen et al. 2011) and with thermal water (Persson et al. 2013) associated with both sources. Recent detections include the prebiotic molecule glycolaldehyde (Jørgensen et al. 2012), its deuteration (Jørgensen et al. 2016), and the detection of ethylene oxide, acetone, propanal, methyl isocyanate and formamide (Coutens et al. 2016; Lykke et al. 2017; Martín-Doménech et al. 2017; Ligterink et al. 2017). For an extended overview of the literature on IRAS 16293–2422 and a review of the ALMA-PILS survey see Jørgensen et al. (2016).

Measurements of the deuterium fractionation of formaldehyde toward IRAS 16293–2422 have previously been limited to single dish observations. Using the JCMT and the CSO van Dishoeck et al. (1995) derived a HDCO/H₂CO ratio of 14±7% in a 20'' beam (2400 AU), taking non-LTE effects into account. Assuming the same excitation temperature for both HDCO and H₂CO an even higher ratio of HDCO/H₂CO=33⁺⁴²₋₁₈% was obtained with the same observations by Parise et al. (2006). The D₂CO/H₂CO ratio was mapped around the source and constrained to 3% toward the edge of the envelope and peaking at 16% toward a position roughly one single dish beam south (Ceccarelli et al. 1998, 2001). Further single dish observations, also with signatures of self-absorption, constrain the ratios to HDCO/H₂CO=13–16%, D₂CO/H₂CO = 5–6%, and D₂CO/HDCO= 33–40% (Loinard et al. 2000). However, for these studies opacity effects and multiple contributions to the emission within the beam might affect the column densities by factors of a couple to a few (Parise et al. 2006), thus the abundances in cold gas need to be more accurately constrained for a proper comparison with ratios derived here for the warm gas on small scales.

Formaldehyde was mapped at high angular resolution toward IRAS 16293–2422 by Schöier et al. (2004), showing some of the uncertainties involved in deriving abundances of formaldehyde with single dish telescopes, and highlighting the importance of higher resolution.

The formation and deuteration of formaldehyde, methanol and other species were studied through chemical modeling by Taquet et al. (2012) and more recently by Taquet et al. (2014). The more recent models trace the deuteration with time from the prestellar core stage to the end of the deeply-embedded stage. The models cannot reproduce the high levels of formaldehyde deuteration previously observed. To further test the chemical models and assess the importance of grain surface formation as well as the various formation paths it is important to accurately constrain the relative abundances of the different deuterated forms of formaldehyde. Interferometric observations of the inner warm region have the possibility to constrain the abundances where the grain surface ice has just been completely sublimated, without the pitfalls of single dish observations.

This paper is laid out as follows. In Sect. 2 details of the observations and spectroscopic data are given together with a description of the analysis method. Section 3 gives the result of the analysis; the measured excitation temperature and the column densities of the various isotopologues of formaldehyde. In Section 4 the derived abundances and isotopologue ratios are discussed with previous measurements and chemical models in

mind. This is followed by a short summary and outlook on future prospects in Sect. 5.

2. Observations and Analysis

This study is based on observations from the Protostellar Interferometric Line Survey (PILS¹), an ALMA band 7 unbiased spectral line survey with complete coverage between 329.15 and 362.90 GHz at 0.244 MHz resolution. For details, including observing conditions, calibration and imaging, see Jørgensen et al. (2016). The observations include both the main array of 12 m dishes, and the 7 m dishes of the Atacama Compact Array (ACA). The observations cover both sources in the IRAS 16293–2422 multiple system. The PILS spectrum toward source A shows lines with larger line widths (2–8 km s⁻¹ FWHM) than for source B (around 1 km s⁻¹ FWHM) making line identification much easier toward the B source, also when comparing to the Galactic Center and high-mass protostars previously observed in the same fashion. The inclusion of the ACA observations fills in the shorter baselines and thus the extended emission encapsulating the sources with a maximum recoverable scale of $\sim 13''$. The final spectral image cubes have a RMS of 4–8 mJy beam⁻¹ km s⁻¹, a synthesized beam with a FWHM of 0''.5 and a flux calibration uncertainty of $\sim 5\%$. The phase center is located half-way between the large scale binary A and B ($\alpha_{J2000} = 16^{\text{h}}32^{\text{m}}22^{\text{s}}.72$; $\delta_{J2000} = -24^{\circ}28'34''.3$) and the field of view covers roughly 15''.

The observed spectra toward the two main sources show a large number of lines. The unprecedented sensitivity and richness of the observed spectrum have so far facilitated several new detections and constraints on isotopologue ratios. So far the abundances and excitation conditions for acetone, propanal and methyl isocyanate (Lykke et al. 2017; Ligterink et al. 2017), and also the deuterium fraction of formamide, glycolaldehyde, ketene and other oxygen-bearing (complex) organic molecules have been presented (Coutens et al. 2016; Jørgensen et al. 2016, 2017).

Similar to the analysis presented in Lykke et al. (2017), Coutens et al. (2016) and Ligterink et al. (2017), the spectrum from an offset position toward source B was extracted. Located at $\alpha_{J2000} = 16^{\text{h}}32^{\text{m}}22^{\text{s}}.58$ $\delta_{J2000} = -24^{\circ}28'32''.8$ it is $\sim 0''.5$ offset (i.e. one beam) in the Southwest direction relative to the continuum peak of source B (see Fig. 1). The main reason for this is that toward the continuum peak, lines are affected by varying degrees of absorption and continuum optical depth, complicating the line identification and analysis of the abundances (see contours in the right panel of Fig. 1).

2.1. Laboratory spectroscopy data

The transition frequencies and other data (including partition function values) of all forms of formaldehyde were taken from the CDMS database (Müller et al. 2001, 2005).

The H₂CO, H₂C¹⁷O and H₂C¹⁸O entries are based on Müller & Lewen (2017), and the H₂¹³CO entry on Müller et al. (2000c). Important additional data are from Brünken et al. (2003); Bocquet et al. (1996) in the case of H₂CO along with ground state combination differences employed in Müller et al. (2000a). Müller et al. (2000b); Cornet & Winnewisser (1980a) contributed additional data to the H₂C¹⁸O entry, the latter also to those of H₂CO and H₂¹³CO. The H₂C¹⁷O entry employed previ-

ous data from Cornet et al. (1980b) and from Flygare & Lowe (1965).

Entries of the deuterated isotopologues, HDCO, D₂CO, HD¹³CO, D₂¹³CO, HDC¹⁸O and D₂C¹⁸O, were based on the very recent analyses by Zakharenko et al. (2015). The analyses took into account earlier data for five of these six isotopologues (no earlier data for HDC¹⁸O) from Dangoisse et al. (1978) as well as extensive data for HDCO and D₂CO from Bocquet et al. (1999). Far-infrared data of D₂CO (Lohilahti & Horneman 2004) and D₂¹³CO (Lohilahti et al. 2005) were also employed for the CDMS entries.

2.2. Abundances

The abundances are estimated by computing a synthetic spectrum assuming Local Thermal Equilibrium (LTE) and comparing it with the observed spectrum. In addition to this, CASSIS² was used as a support tool to e.g. check for possible line blending from other species, which is the same method as in other PILS survey studies e.g. Coutens et al. (2016); Ligterink et al. (2017); Lykke et al. (2017). For HDCO a full grid in excitation temperature and column density is calculated and the coordinates of the minimum χ^2 are taken as the best estimates. Because the excitation temperature is better constrained using HDCO than through the other isotopologues of formaldehyde, its resulting temperature is used to constrain the column density for all forms. This is done by varying the column density, calculating the integrated line flux for each line not affected by optical depth effects or significant blending by other species and comparing to the observed line flux by calculating the χ^2 estimate. The uncertainty in abundance is estimated by finding the minimum χ^2 for the lower and upper bound excitation temperature (i.e. 93 and 119 K, Fig. 2). The column densities are corrected for the continuum emission from the surrounding dense dust (by multiplying with a factor of 1.1658, Jørgensen et al. 2016). A line width of 1.15 km s⁻¹ (FWHM) and system velocity of 2.7 km s⁻¹ were used (relevant for source B). The assumed line width agrees well with the observed line profiles for formaldehyde and is similar to that of other species (Coutens et al. 2016; Lykke et al. 2017).




Fig. 1. PILS ALMA continuum map of the IRAS 16293–2422 system at 850 μm . The background image shows the continuum emission, the contours indicate the H₂C¹⁷O 5_{2,3}–4_{2,2} line ($E_u = 98.5$ K) integrated emission, the star is the position of the continuum peak, and the plus sign the position of the extracted spectrum (one beam, i.e. 0''.5, away from the continuum). The continuum image is cut off at 15 mJy (3σ) and the line contours start at 25 mJy km s⁻¹ (5σ) in steps of 120 mJy km s⁻¹.

The detected transitions of the various isotopologues, deuterated and non-deuterated are listed in Table A.1 in the Appendix. The analysis focuses on H₂CO (3 lines detected) and several deuterated and non-deuterated isotopologues (number of de-

¹ <http://youngstars.nbi.dk/PILS>

² Developed by IRAP-UPS/CNRS <http://cassis.irap.omp.eu>

ected lines in parenthesis): H_2^{13}CO (9), $\text{H}_2\text{C}^{18}\text{O}$ (9), $\text{H}_2\text{C}^{17}\text{O}$ (8), HDCO (6), HDC^{18}O (1 tentative), D_2CO (11), and D_2^{13}CO (10).




Fig. 2. Grid constraining the excitation temperature and column density of HDCO. The cross marks the best fit values, $1.3 \pm 0.2 \times 10^{17} \text{ cm}^{-2}$ and $106 \pm 13 \text{ K}$ from interpolation of the values within the marked area. The contour line marks the $\Delta\chi^2 = 2.3$ region. Note that the column density value has been corrected for the continuum. The dots show the evaluated column densities and excitation temperatures.

3. Results

In the following sections we present the results of a detailed analysis of the column densities, the various isotopic ratios and the constraints on the excitation temperature. Only optically thin lines are included in the analysis to minimize any opacity effects.

3.1. Excitation temperature

The excitation temperature was determined to be $106 \pm 13 \text{ K}$ by fitting the spectrum of HDCO, and the resulting temperature is used for all forms of formaldehyde to constrain their column density. The χ^2 in a grid around the best fitting column density and excitation temperature is shown in Fig. 2. The region with an added $\Delta\chi^2 = 2.3$ is taken as the 1σ error in T_{ex} and N_{tot} for HDCO (Avni 1976; Press et al. 2002). This temperature, together with the constrained column densities reproduce the lines for all isotopologues well. The figures in Appendix B show the LTE model for the best fitting column densities and the observed spectrum for all lines of all species, including non-detections.

In some studies of other sources a second temperature component is needed to reproduce the spectra of lines of both lower and higher upper energy levels (e.g. Isokoski et al. 2013). If a second temperature component were relevant for this study, e.g. the low E_u lines would be systematically under/over-produced in the synthetic spectrum. Since this is not the case it is likely that only one temperature component is present. The other isotopologues were not as suitable to use for constraining the excitation temperature, the resulting parameter space of HDCO shows a clearer minimum than any of the other isotopologues investigated (Fig. 2). The spread in E_u of the detected and optically thin lines of HDCO makes it better at constraining the temperature. Using selected lines of $\text{H}_2\text{C}^{18}\text{O}$ a less constrained excitation temperature is obtained ($90 \sim 200 \text{ K}$). For the other isotopo-

Table 1. Best-fit continuum corrected column densities at $T_{\text{ex}} = 106 \pm 13 \text{ K}$ for all the isotopologues of formaldehyde.

Species	$N_{\text{tot}} (\text{cm}^{-2})$
H_2CO	$1.9^{+0.1}_{-0.2} \times 10^{18}$
H_2^{13}CO	$3.6^{+0.7}_{-0.7} \times 10^{16}$
$\text{H}_2\text{C}^{17}\text{O}$	$7.2^{+0.1}_{-0.1} \times 10^{14}$
$\text{H}_2\text{C}^{18}\text{O}$	$2.3 \pm 0.1 \times 10^{15}$
HDCO	$1.3 \pm 0.2 \times 10^{17}$
HDC^{18}O	$\leq 1.4 \pm 0.4 \times 10^{14}$
D_2CO	$1.6^{+0.4}_{-0.5} \times 10^{16}$
D_2^{13}CO	$2.2^{+0.2}_{-0.4} \times 10^{14}$

Table 2. Derived fractionation ratios for the various forms of formaldehyde. The errors are propagated from the uncertainty in the determined column densities.

Species	Ratio	Comb. ^a	D/H
HDCO / H_2CO	$6.8^{+1.1}_{-1.3} \%$	$6.5 \pm 1 \%$	$3.25 \pm 1 \%$
$\text{HDC}^{18}\text{O}/\text{H}_2\text{C}^{18}\text{O}$	$6.0 \pm 1.5 \%$	–	–
$\text{D}_2\text{CO} / \text{HDCO}$	$12.8^{+3.3}_{-2.1} \%$	–	$25.6^{+3.3}_{-4.1} \%$
$\text{D}_2\text{CO} / \text{H}_2\text{CO}$	$0.9^{+0.2}_{-0.3} \%$	$0.6(4) \pm 0.1 \%$	$8.0 \pm 0.1 \%$
$\text{D}_2^{13}\text{CO}/\text{H}_2^{13}\text{CO}$	$0.6^{+0.1}_{-0.2} \%$	–	–
$\text{H}_2\text{CO} / \text{H}_2^{13}\text{CO}$	52^{+10}_{-16}	56^{+8}_{-11}	–
$\text{D}_2\text{CO} / \text{D}_2^{13}\text{CO}$	74^{+18}_{-24}	–	–
$\text{H}_2\text{CO} / \text{H}_2\text{C}^{18}\text{O}$	800^{+47}_{-98}	805^{+43}_{-79}	–
HDCO / HDC^{18}O	908 ± 263	–	–
$\text{H}_2\text{CO} / \text{H}_2\text{C}^{17}\text{O}$	2596^{+639}_{-395}	–	–
$\text{H}_2\text{C}^{18}\text{O} / \text{H}_2\text{C}^{17}\text{O}$	$3.2^{+0.2}_{-0.3}$	–	–

Notes. ^(a) Combined isotopologue ratio (following Barlow 2003).

logues, it was possible to constrain the column density to high accuracy with the given excitation temperature.

In addition to the detected, optically thin HDCO lines (Table A.1) we also used upper limits of one non-detected HDCO transition at 348.965 GHz ($E_u = 1017.78 \text{ K}$, $\log_{10}(A_{ij}) = -4.5321$) to constrain the excitation temperature (see Table A.2). Since the LTE model should not overproduce this line, it increases the constraint. Thus, a wide spread in E_u was used to constrain the T_{ex} , but avoiding the $11_{1,10} - 11_{1,11}$ line at 346.74 GHz which is optically thick.

3.2. Column densities and ratios

The best fit column densities at $T_{\text{ex}} = 106 \pm 13 \text{ K}$ for the species are listed in Table 1. The uncertainty in abundance is dominated by the uncertainty in excitation temperature. In Table 2 the various column density ratios for the studied isotopologues are shown. Ratios where multiple estimates exist using isotopologues, i.e. HDCO / H_2CO and $\text{HDC}^{18}\text{O}/\text{H}_2\text{C}^{18}\text{O}$ for the HDCO/ H_2CO ratio, are combined. When averaging/combining values with asymmetric uncertainties care has to be taken when calculating the best combined estimate (for a detailed discussion and solution see Section 3 in Barlow 2003). To constrain the column density of the main isotopologue H_2CO only one detection and one upper-limit (non-detection) are used in the fit. The other H_2CO transitions are optically thick, by a large margin, and they also show some extended emission (van der Wiel, in prep.).

4. Discussion

4.1. Column densities and excitation temperature

The derived excitation temperature of 106 ± 13 K agrees with the results for some of the other molecules studied within the PILS survey. Acetaldehyde, ketene, ethylene oxide, acetone, propanal, and also dimethyl ether show a similar excitation temperature, on the order of 125 K (Lykke et al. 2017; Jørgensen et al. 2017). Interestingly van Dishoeck et al. (1995) derived, after subtracting the extended contribution from the central emission, a kinetic temperature of 100^{+40}_{-20} K from the $\text{H}_2\text{CO } 3_{22}\text{--}3_{21}/3_{03}\text{--}2_{02}$ ratio toward IRAS 16293–2422.

The derived column densities fit the observed spectrum well using the assumed T_{ex} of 106 K from HDCO. The H_2^{13}CO line at 355.04 GHz is partly blended with the weaker counterpart of $\text{H}_2\text{C}^{17}\text{O}$. This means that the H_2^{13}CO column density might be overestimated by a small amount, however the effect is within the uncertainties. A $\text{H}_2\text{C}^{17}\text{O}$ column density and excitation temperature taken at the lower uncertainty limit will make its blending effect on H_2^{13}CO insignificant.

4.2. Deuterium fractionation

In this section we present the deuterium fractionation calculated from the observations and then compare these to the model of Taquet et al. (2014).

4.2.1. Measured D/H ratios

Together with the column density ratios for the studied isotopologues in Table 2 are the best estimated D/H ratios as well (combined measurements where applicable), where D/H is given by $\text{HDCO}/\text{H}_2\text{CO} \times 0.5$, $\text{D}_2\text{CO}/\text{HDCO} \times 2$ and $\sqrt{\text{D}_2\text{CO}/\text{H}_2\text{CO}}$. While the errors quoted for the column densities are derived by fitting spectra with the column density for the highest and lowest excitation temperature, the errors in the ratios were derived with error propagation. Thus the uncertainties of the column densities and in extension the fractionation ratios, reflect also the uncertainty in excitation temperature.

The $\text{HDC}^{18}\text{O}/\text{H}_2\text{C}^{18}\text{O}$ ratio is 6.0% (D/H ratio of 3.0%), and it agrees with the main isotopologues $\text{HDCO}/\text{H}_2\text{CO}$ ratio of 6.8% (D/H ratio of 3.4%), confirming the method used for constraining the column densities. Their combined ratio 6.5% (D/H ratio 3.25%) is taken as the best estimate. The $\text{D}_2^{13}\text{CO}/\text{H}_2^{13}\text{CO}$ ratio (0.6%) is also similar to the main form $\text{D}_2\text{CO}/\text{H}_2\text{CO}$ ratio (0.9%), their combined ratio being 0.6(4)% (D/H ratio 8.0%). Finally, the $\text{D}_2\text{CO}/\text{HDCO}$ ratio is 12.8% (D/H ratio is 25.6%).

These numbers show that the $\text{D}_2\text{CO}/\text{HDCO}$ ratio is two times higher than the $\text{HDCO}/\text{H}_2\text{CO}$ ratio, significantly more than the $\frac{1}{4}$ expected from statistical arguments (see Equation 1 and related text). Furthermore, other organics show D/H ratios similar to this (Jørgensen et al. 2017). This shows that for a full understanding a more complex interpretation than simple statistical arguments is needed. Whether other protostars show similar deuterium fractionation ratios for H_2CO as deduced here is not yet clear, observations of more protostars are needed to investigate what role initial environment and evolution might play in determining the ratios.

4.2.2. Models

Taquet et al. (2012) and more recently Taquet et al. (2014) studied the evolution of deuterated ices during the earliest stages of

star formation, including the prestellar and protostellar (Class 0) phase. In the 2012 study the importance of abstraction and substitution reactions is shown, as also supported by lab data (Hidaka et al. 2009). Furthermore, the $\text{HDCO}/\text{H}_2\text{CO}$ ratio is not affected by the inclusion of abstraction reactions due to the efficiency of the deuterium abstraction on HDCO, forming back H_2CO . Thus, constraining the deuterium fractionation in all forms of formaldehyde, especially the doubly deuterated form is crucial for constraining the formation conditions and the chemical models. The results show that a reaction network without abstraction and substitution reactions cannot reproduce the $\text{HDCO}/\text{H}_2\text{CO}$ and $\text{D}_2\text{CO}/\text{HDCO}$ ratios simultaneously. Including abstraction reactions shows that the ratios are reproducible at moderate densities ($n_{\text{H}} = 10^6 \text{ cm}^{-3}$). The model in the 2014 study traces the density and temperature evolution from a (inside-out) collapsing core to the end of the deeply-embedded stage (Class 0). The observed (combined, best estimate) ratios of $\text{HDCO}/\text{H}_2\text{CO}$ of $6.5 \pm 1\%$ and $\text{D}_2\text{CO}/\text{H}_2\text{CO}$ of $0.6(4) \pm 0.1\%$ are best reproduced by the model at $t = 1.1 \times 10^5$ years. This timescale represents the beginning of the Class 0 stage, where the D/H ratio in HDCO is 4.1% and 0.18% for D_2CO (Table 7 in Taquet et al.). While the density and temperature profile at this time step from Taquet et al. is similar to the physical model of Crimier et al. (2010) for IRAS 16293–2422 specifically, the modeled deuterium fractionation observed in the hot corino gas is mostly due to the deuteration processes of formaldehyde in ices in the previous dense cloud stage and thus less dependent on the current physical structure. In Taquet et al. the assumed size of the beam in the observations of warm gas with interferometers is 0.4 (or 50 AU) which is very similar to the beam in this study (0.5). The excitation temperature derived shows that we are probing the warm gas of the hot corino. The ratios derived using single dish telescopes, which may apply to larger scales, are more difficult to compare with the model results. It is not clear to which extent opacity effects, assumed size of emitting region and contributions of various emission components play a role in the previous single dish observations, thus we refrain from any extensive comparison with those results.

The main production phase of deuterated molecules comes towards the end of the prestellar core stage (Furuya et al. 2016). This is in agreement with previous studies of starless cores using N_2H^+ and N_2D^+ (Crapsi et al. 2005). Taquet et al. (2014) also trace the deuteration through these early stages, with similar conclusions. The model reproduces the large differences between singly and doubly deuterated forms with the $\text{D}_2\text{O}/\text{H}_2\text{O}$ ratio being about 7× higher than the $\text{HDO}/\text{H}_2\text{O}$ ratio. In formaldehyde toward IRAS 16293–2422 the $\text{D}_2\text{CO}/\text{HDCO}$ ratio is about $2 \times \text{HDCO}/\text{H}_2\text{CO}$, a similar but much less pronounced effect than for water.

4.2.3. Formaldehyde and methanol

Methanol (CH_3OH) was studied in the framework of the ALMA-PILS survey (Jørgensen et al. 2017). The deuterium fractionation (D/H) derived toward IRAS 16293–2422 B in methanol is 2%. This is comparable to singly deuterated formaldehyde. Interesting enough, the time scale of the astrochemical model presented by Taquet et al. (2014) that best reproduces this level of fractionation in methanol is also $t = 1.1 \times 10^5$ years. This could indicate that IRAS 16293–2422 B is indeed a very young Class 0 source. Using the column density derived for methanol by Jørgensen et al. the $\text{H}_2\text{CO}/\text{CH}_3\text{OH}$ ratio is 0.19 toward IRAS 16293–2422 B. A ratio of 1 was inferred toward the inner regions of the same source by Maret et al. (2005) using single dish

observations and radiative transfer modeling, clearly different from the ratio presented in this study. However, the large uncertainties inherent in using single dish measurements to constrain inner envelope abundances makes a comparison difficult. The measured $\text{H}_2\text{CO}/\text{CH}_3\text{OH}$ ratio of 0.19 is close to what has been constrained in ices toward protostars, ranging from 0.1 to 0.67 (Keane et al. 2001). This strengthens the view that warm gas in the innermost regions of protostars represent the bulk ice composition. The measured $\text{H}_2\text{CO}/\text{CH}_3\text{OH}$ ratio is in agreement with what was measured in hot cores by Bisschop et al. (2007), where the mean and 1σ standard deviation presented toward seven hot cores was 0.22 ± 0.05 .

4.3. Other isotopic ratios

While the $^{16}\text{O}/^{18}\text{O}$ ratio is 500 at this distance from the Galactic center (Wilson & Rood 1994; Wilson 1999), the ratio measured in formaldehyde for IRAS 16293–2422 B is 800 for H_2CO and 908 for HDCO, both slightly higher than expected at these distances from the Galactic center. The combined ratio for the two measurements is 805^{+43}_{-79} . Given the uncertainties involved, including the Galactic gradient, $^{16}\text{O}/^{18}\text{O}=400\text{--}600$ at 8 kpc, it is difficult to draw any conclusions, although the combined best estimate ratio is higher than the range in the local ISM.

The $^{18}\text{O}/^{17}\text{O}$ ratio in formaldehyde toward IRAS 16293–2422 B is $3.2^{+0.2}_{-0.3}$ ($\text{H}_2\text{C}^{18}\text{O}/\text{H}_2\text{C}^{17}\text{O}$). Upper limits of the isotopic variant HDC^{17}O derived using the $6_{1,6} - 5_{1,5}$ transition at 360.962 GHz gives a $^{18}\text{O}/^{17}\text{O}$ ratio of ≥ 2 , compatible with the local ISM value and $\text{H}_2\text{C}^{18}\text{O}/\text{H}_2\text{C}^{17}\text{O}$ ratios. Jørgensen et al. (2002) measured the $\text{C}^{18}\text{O}/\text{C}^{17}\text{O}$ ratio toward the envelopes of 19 protostars, including Class 0 and I sources. Toward IRAS 16293–2422 the measured ratio was 3.9, in agreement with the ratio measured for formaldehyde, and roughly 74% of all the measured sources showed a ratio 2–4. The $\text{H}_2\text{C}^{18}\text{O}/\text{H}_2\text{C}^{17}\text{O}$ ratio was first measured toward the Galactic center, 4 kpc molecular ring and the local ISM to be 3.2 ± 0.2 (Penzias 1981). Wouterloot et al. (2008) discovered a tentative gradient in the $^{18}\text{O}/^{17}\text{O}$ ratio with galactocentric distance ranging from 3 in the center to 5 in the outer galaxy, and more recently Li et al. (2016, e.g. Fig. 3) strengthened this conclusion. The ratio in the outermost galaxy, beyond 11 kpc is still uncertain. The observed gradient is consistent with an inside-out formation of the galaxy. The observed value toward IRAS 16293–2422 fits in the galactic $^{18}\text{O}/^{17}\text{O}$ gradient and could indicate a fairly normal formation environment at this distance from the Galactic center. The deviating $^{16}\text{O}/^{18}\text{O}$ ratio (see previous paragraph) is not fully consistent with this conclusion.

The $^{12}\text{C}/^{13}\text{C}$ ratio for H_2CO and D_2CO and their combined ratio of 56^{+8}_{-11} are consistent with the ratio of 68 ± 30 , which is relevant for these galactocentric distances (Wilson & Rood 1994; Wilson 1999; Milam et al. 2005). While lines of the isotopic variant HD^{13}CO are too weak to detect in our survey, its upper limit is compatible with a $^{12}\text{C}/^{13}\text{C}$ ratio of ≥ 55 (using mainly the $11_{1,10} - 11_{1,11}$ transition). Recent results from the ALMA-PILS survey presented in Jørgensen et al. (2017) show that the $^{12}\text{C}/^{13}\text{C}$ ratio in glycolaldehyde, ethanol, methyl formate, and dimethyl ether is around 20–40, lower than for formaldehyde, methanol and the surrounding ISM. This is attributed to a later formation of the low $^{12}\text{C}/^{13}\text{C}$ ratio species, when the availability of ^{13}C is higher.

4.4. The path to more complex molecules

Since formaldehyde is thought to play a role in the formation of more complex molecules, it is interesting to compare the deuteration between formaldehyde and its potential daughter species. Formaldehyde may also lead to the formation of formamide (NH_2CHO) on grain surfaces as shown by some laboratory experiments (Fedoseev et al. 2016). Barone et al. (2015) proposed that formamide could form due to the reaction between H_2CO and NH_2 in the gas phase. Recently Coutens et al. (2016) detected the mono-deuterated forms of formamide using the ALMA-PILS survey and reported a deuteration level of $\sim 2\%$ assuming a standard $^{12}\text{C}/^{13}\text{C}$ ratio of 68. Skouteris et al. (2017) also calculated the rate coefficients of the reactions producing deuterated formamide. Consequently, a comparison of the deuteration of formamide and formaldehyde can help determine if this gas phase pathway (for formamide) is possible or if it has to be ruled out in this source. According to the respective rate coefficients determined for NH_2CHO and NH_2CDO by Skouteris et al., the $\text{HDCO}/\text{H}_2\text{CO}$ ratio should be three times higher than the $\text{NH}_2\text{CDO}/\text{NH}_2\text{CHO}$ ratio if this gas-phase reaction occurs. At similar spatial scales, the best estimate $\text{HDCO}/\text{H}_2\text{CO}$ ratio is $6.5 \pm 1\%$, i.e. approximately three times higher than the $\text{NH}_2\text{CDO}/\text{NH}_2\text{CHO}$ ratio. Thus, a gas phase formation pathway for formamide cannot be ruled out. Since the expected deuteration levels for the case of a grain surface formation pathway has not been explored, none of the scenarios for formamide formation can be ruled out from this comparison at this point.

5. Summary and Outlook

Measuring the deuterium fractionation is an important tool in understanding the early chemical evolution of forming protostars. The evolving picture is that the gas phase deuteration of molecules in the inner warm regions, representative of the bulk ice, seems to be lower than previously thought based on single dish observations. This highlights that more high-sensitivity and -resolution radio interferometric observations are needed to probe this gas reservoir and unlock the chemical history and future of deeply embedded low-mass protostars.

In this study we have detected several lines of various isotopologues of formaldehyde toward the deeply embedded low-mass protostar IRAS 16293–2422 B as part of the ALMA-PILS survey. Both $\text{H}_2\text{C}^{17}\text{O}$ and D_2^{13}CO are detected for the first time in the ISM. Many of the lines are optically thick due to the high densities probed. The determined excitation temperature for HDCO is 106 ± 13 K, similar to several other molecules in the same survey, and previous measurements.

Assuming the same excitation temperature constrained from HDCO for all the forms of formaldehyde to determine the column densities shows that many of the transitions are optically thick and abundances high. The detected forms of formaldehyde are: H_2CO , H_2^{13}CO , $\text{H}_2\text{C}^{17}\text{O}$, $\text{H}_2\text{C}^{18}\text{O}$, HDCO, HDC^{18}O , D_2CO and D_2^{13}CO . The large number of lines covered by the unbiased spectral survey makes it possible to constrain the column density by focusing on the optically thin, unblended lines.

The measured $\text{HDCO}/\text{H}_2\text{CO}$ ratio is $6.5 \pm 1\%$. This is significantly lower than what was previously estimated using single dish telescopes (14–33%). This is also true for doubly deuterated formaldehyde, where the $\text{D}_2\text{CO}/\text{HDCO}$ ratio is $12.8^{+3.3}_{-4.1}\%$, and $\text{D}_2\text{CO}/\text{H}_2\text{CO}$ ratio is $0.6 \pm 0.1\%$. The lines observed with single dish telescopes in previous studies also covered by the ALMA-PILS survey are significantly optically thick. The source size is

better constrained here by spatially resolving the emitting region. These effects could be the reason for the different ratio(s).

These levels of deuterium fractionation are in line with formaldehyde forming on the grains once CO has frozen onto the surface as soon as the temperature drops low enough. The astrochemical model of Taquet et al. (2014) can approximately reproduce the deuterium fractionation observed in formaldehyde with a grain surface formation pathway dominating the production. However, the role of gas phase formation routes is still not clear.

The effect seen for water (Persson et al. 2013; Coutens et al. 2012; Furuya et al. 2016), where D_2O/HDO is significantly higher than the HDO/H_2O ratio is only minor for formaldehyde due to the even later start of formation toward the end of the prestellar core phase in comparison to water. This also implies higher absolute deuterium fractionation ($D/H_{H_2CO} \sim 3\%$, $D/H_{H_2O} \sim 0.1\%$), due to the more favorable conditions (low o/p -ratio, low T , higher accretion of D atoms onto grains). The deuterium fractionation presented for formaldehyde is similar to that of other molecules with similar complexity such as methanol (CH_3OH Jørgensen et al. 2017) and formamide (NH_2CHO , Coutens et al. 2016), at around 2–3%. This formation-time dependence on deuteration implies that more complex molecules which form even later should have even higher levels of deuterium fractionation.

The $^{12}C/^{13}C$ and $^{18}O/^{17}O$ ratios in formaldehyde are consistent with the values measured for the surrounding ISM at the relevant galactocentric distances. This could indicate a relatively common formation environment in the Galaxy. However, the $^{16}O/^{18}O$ ratio is slightly higher than the local ISM value thus it is not possible to draw any strong conclusions based on this.

Future high resolution interferometric observations constraining the deuterium fractionation in formaldehyde of other sources, could shed light on any possible dependence on environment and mass. In addition to this, multi transition single dish observations unambiguously characterizing the cold gas would improve the global picture of formaldehyde deuterium fractionation.

Acknowledgements. This research made use of Astropy and Astroquery, community-developed Python packages for Astronomy (Astropy Collaboration et al. 2013; Ginsburg et al. 2017). MVP postdoctoral position is funded by the ERC consolidator grant 614264. JKJ acknowledges support from the European Research Council (ERC) under the European Union’s Horizon 2020 research and innovation programme (grant agreement No 646908) through ERC Consolidator Grant “S4F”. Research at Centre for Star and Planet Formation is funded by the Danish National Research Foundation. AC postdoctoral grant is funded by the ERC Starting Grant 3DICE (grant agreement 336474). VT has received funding from the European Union’s Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement n. 664931. EvD acknowledges EU A-ERC grant 291141 CHEMPLAN and a KNAW professorship prize. This paper makes use of the following ALMA data: ADS/JAO.ALMA#2013.1.00278.S. ALMA is a partnership of ESO (representing its member states), NSF (USA) and NINS (Japan), together with NRC (Canada) and NSC and ASIAA (Taiwan), in cooperation with the Republic of Chile. The Joint ALMA Observatory is operated by ESO, AUI/NRAO and NAOJ.

References

Astropy Collaboration, Robitaille, T. P., Tollerud, E. J., et al. 2013, *A&A*, 558, A33
 Avni, Y. 1976, *ApJ*, 210, 642
 Barlow, R. 2003, in *Statistical Problems in Particle Physics, Astrophysics, and Cosmology*, ed. L. Lyons, R. Mount, & R. Reitmeyer, 250
 Barone, V., Latouche, C., Skouteris, D., et al. 2015, *MNRAS*, 453, L31
 Bisschop, S. E., Jørgensen, J. K., Bourke, T. L., Bottinelli, S., & van Dishoeck, E. F. 2008, *A&A*, 488, 959

Bisschop, S. E., Jørgensen, J. K., van Dishoeck, E. F., & de Wachter, E. B. M. 2007, *A&A*, 465, 913
 Bocquet, R., Demaison, J., Cosléou, J., et al. 1999, *Journal of Molecular Spectroscopy*, 195, 345
 Bocquet, R., Demaison, J., Poteau, L., et al. 1996, *Journal of Molecular Spectroscopy*, 177, 154
 Bottinelli, S., Ceccarelli, C., Neri, R., et al. 2004, *ApJ*, 617, L69
 Brünken, S., Müller, H. S. P., Lewen, F., & Winnewisser, G. 2003, *Physical Chemistry Chemical Physics (Incorporating Faraday Transactions)*, 5
 Ceccarelli, C., Castets, A., Loinard, L., Caux, E., & Tielens, A. G. G. M. 1998, *A&A*, 338, L43
 Ceccarelli, C., Loinard, L., Castets, A., et al. 2001, *A&A*, 372, 998
 Charnley, S. B., Tielens, A. G. G. M., & Rodgers, S. D. 1997, *ApJ*, 482, L203
 Cornet, R. A., Landsberg, B. M., & Winnewisser, G. 1980b, *Journal of Molecular Spectroscopy*, 82, 253
 Cornet, R. A. & Winnewisser, G. 1980a, *Journal of Molecular Spectroscopy*, 80, 438
 Coutens, A., Jørgensen, J. K., Persson, M. V., et al. 2014, *ApJ*, 792, L5
 Coutens, A., Jørgensen, J. K., van der Wiel, M. H. D., et al. 2016, *A&A*, 590, L6
 Coutens, A., Vastel, C., Caux, E., et al. 2012, *A&A*, 539, A132
 Crapsi, A., Caselli, P., Walmsley, C. M., et al. 2005, *ApJ*, 619, 379
 Crimier, N., Ceccarelli, C., Maret, S., et al. 2010, *A&A*, 519, A65
 Dangoisse, D., Willemot, E., & Bellet, J. 1978, *Journal of Molecular Spectroscopy*, 71, 414
 Fedoseev, G., Chuang, K.-J., van Dishoeck, E. F., Ioppolo, S., & Linnartz, H. 2016, *MNRAS*, 460, 4297
 Flygare, W. H. & Lowe, J. T. 1965, *J. Chem. Phys.*, 43, 3645
 Fuchs, G. W., Cuppen, H. M., Ioppolo, S., et al. 2009, *A&A*, 505, 629
 Furuya, K., Aikawa, Y., Hincelin, U., et al. 2015, *A&A*, 584, A124
 Furuya, K., van Dishoeck, E. F., & Aikawa, Y. 2016, *A&A*, 586, A127
 Ginsburg, A., Parikh, M., Woillez, J., et al. 2017, *Astroquery: Access to online data resources*, *Astrophysics Source Code Library*
 Hidaka, H., Watanabe, M., Kouchi, A., & Watanabe, N. 2009, *The Astrophysical Journal*, 702, 291
 Isokoski, K., Bottinelli, S., & van Dishoeck, E. F. 2013, *A&A*, 554, A100
 Jørgensen, J. K., Bourke, T. L., Nguyen Luong, Q., & Takakuwa, S. 2011, *A&A*, 534, A100
 Jørgensen, J. K., Favre, C., Bisschop, S. E., et al. 2012, *ApJ*, 757, L4
 Jørgensen, J. K., Müller, H. S. P., Calcutt, H., et al. 2017, *submit*
 Jørgensen, J. K., Schöier, F. L., & van Dishoeck, E. F. 2002, *A&A*, 389, 908
 Jørgensen, J. K., van der Wiel, M. H. D., Coutens, A., et al. 2016, *A&A*, 595, A117
 Keane, J. V., Tielens, A. G. G. M., Boogert, A. C. A., Schutte, W. A., & Whittet, D. C. B. 2001, *A&A*, 376, 254
 Kuan, Y.-J., Huang, H.-C., Charnley, S. B., et al. 2004, *ApJ*, 616, L27
 Li, H.-K., Zhang, J.-S., Liu, Z.-W., et al. 2016, *Research in Astronomy and Astrophysics*, 16, 47
 Ligterink, N. F. W., Coutens, A., Kofman, V., et al. 2017, *MNRAS*, 469, 2219
 Lohilahti, J. & Horneman, V.-M. 2004, *Journal of Molecular Spectroscopy*, 228, 1
 Lohilahti, J., Mattila, H., Horneman, V.-M., & Pawłowski, F. 2005, *Journal of Molecular Spectroscopy*, 234, 279
 Loinard, L., Castets, A., Ceccarelli, C., et al. 2000, *A&A*, 359, 1169
 Loinard, L., Torres, R. M., Mioduszewski, A. J., & Rodríguez, L. F. 2008, *ApJ*, 675, L29
 Lykke, J. M., Coutens, A., Jørgensen, J. K., et al. 2017, *A&A*, 597, A53
 Maret, S., Ceccarelli, C., Tielens, A. G. G. M., et al. 2005, *A&A*, 442, 527
 Martín-Doménech, R., Rivilla, V. M., Jiménez-Serra, I., et al. 2017, *MNRAS*, 469, 2230
 Milam, S. N., Savage, C., Brewster, M. A., Ziurys, L. M., & Wyckoff, S. 2005, *ApJ*, 634, 1126
 Müller, H. S. P., Gendriesch, R., Lewen, F., & Winnewisser, G. 2000b, *Zeitschrift Naturforschung Teil A*, 55, 486
 Müller, H. S. P., Gendriesch, R., Margulès, L., et al. 2000c, *Physical Chemistry Chemical Physics (Incorporating Faraday Transactions)*, 2
 Müller, H. S. P. & Lewen, F. 2017, *Journal of Molecular Spectroscopy*, 331, 28
 Müller, H. S. P., Schlöder, F., Stutzki, J., & Winnewisser, G. 2005, *Journal of Molecular Structure*, 742, 215
 Müller, H. S. P., Thorwirth, S., Roth, D. A., & Winnewisser, G. 2001, *A&A*, 370, L49
 Müller, H. S. P., Winnewisser, G., Demaison, J., Perrin, A., & Valentin, A. 2000a, *Journal of Molecular Spectroscopy*, 200, 143
 Pagani, L., Salez, M., & Wannier, P. G. 1992, *A&A*, 258, 479
 Parise, B., Ceccarelli, C., Tielens, A. G. G. M., et al. 2006, *A&A*, 453, 949
 Penzias, A. A. 1981, *ApJ*, 249, 518
 Persson, M. V., Jørgensen, J. K., & van Dishoeck, E. F. 2013, *A&A*, 549, L3
 Persson, M. V., Jørgensen, J. K., van Dishoeck, E. F., & Harsono, D. 2014, *A&A*, 563, A74
 Press, W. H., Teukolsky, S. A., Vetterling, W. T., & Flannery, B. P. 2002, *Numerical recipes in C++: the art of scientific computing*

- Prodanović, T., Steigman, G., & Fields, B. D. 2010, MNRAS, 406, 1108
- Roberts, H., Herbst, E., & Millar, T. J. 2004, A&A, 424, 905
- Roberts, H. & Millar, T. J. 2000, A&A, 361, 388
- Schöier, F. L., Jørgensen, J. K., van Dishoeck, E. F., & Blake, G. A. 2004, A&A, 418, 185
- Skouteris, D., Vazart, F., Ceccarelli, C., et al. 2017, MNRAS in press [arXiv:1701.06138]
- Taquet, V., Ceccarelli, C., & Kahane, C. 2012, ApJ, 748, L3
- Taquet, V., Charnley, S. B., & Sipilä, O. 2014, ApJ, 791, 1
- Tielens, A. G. G. M. 1983, A&A, 119, 177
- Turner, B. E. 1990, ApJ, 362, L29
- van Dishoeck, E. F., Blake, G. A., Jansen, D. J., & Groesbeck, T. D. 1995, ApJ, 447, 760
- Watanabe, N. & Kouchi, A. 2002, ApJ, 571, L173
- Wilson, T. L. 1999, Reports on Progress in Physics, 62, 143
- Wilson, T. L. & Rood, R. 1994, ARA&A, 32, 191
- Wouterloot, J. G. A., Henkel, C., Brand, J., & Davis, G. R. 2008, A&A, 487, 237
- Zakharenko, O., Motiyenko, R. A., Margulès, L., & Huet, T. R. 2015, Journal of Molecular Spectroscopy, 317, 41

Appendix A: Formaldehyde lines used in the analysis

Appendix B: Observed spectrum of lines

Table A.1. Lines detected in the ALMA-PILS spectrum of IRAS 16293–2422. The stars in the left-most column indicates the lines that are blended with each other. The square in the right-most column identifies the lines used in constraining the column density (and for HDCO also the temperature).

Species	Q ₁ <i>J_{K_AK_B}</i>	Q ₂	ν GHz	A_{ij} $\log_{10}()$	E_u K	Fit?	
H ₂ CO	5 _{1,5}	4 _{1,4}	351.769	-2.9201	62.5		
	8 _{2,6}	9 _{0,9}	361.968	-5.9856	173.5	■	
	5 _{0,5}	4 _{0,4}	362.736	-2.8626	52.3		
H ₂ C ¹⁷ O	5 _{1,5}	4 _{1,4}	343.333	-2.9518	61.3		
	5 _{0,5}	4 _{0,4}	353.820	-2.8950	51.0	■	
	5 _{2,4}	4 _{2,3}	354.908	-2.9665	98.4	■	
*	5 _{4,2}	4 _{4,1}	355.042	-3.3340	240.1		
*	5 _{4,1}	4 _{4,0}	355.042	-3.3340	240.1		
**	5 _{3,3}	4 _{3,2}	355.202	-3.0835	157.5		
	5 _{3,2}	4 _{3,1}	355.214	-3.0834	157.5	■	
	5 _{2,3}	4 _{2,2}	356.187	-2.9619	98.5	■	
H ₂ C ¹⁸ O	5 _{1,5}	4 _{1,4}	335.816	-2.9805	60.2		
	5 _{0,5}	4 _{0,4}	345.881	-2.9246	49.9		
	5 _{2,4}	4 _{2,3}	346.869	-2.9963	97.4	■	
	5 _{4,2}	4 _{4,1}	346.984	-3.3639	239.6	■	
	5 _{4,1}	4 _{4,0}	346.984	-3.3639	239.6	■	
	5 _{3,3}	4 _{3,2}	347.134	-3.1134	156.7		
	5 _{3,2}	4 _{3,1}	347.144	-3.1134	156.7		
	5 _{2,3}	4 _{2,2}	348.032	-2.9920	97.5	■	
H ₂ ¹³ CO	5 _{1,4}	4 _{1,3}	357.741	-2.8982	63.4		
	5 _{1,5}	4 _{1,4}	343.326	-2.9517	61.3		
	5 _{0,5}	4 _{0,4}	353.812	-2.8949	51.0		
	5 _{2,4}	4 _{2,3}	354.899	-2.9664	98.4		
	5 _{4,2}	4 _{4,1}	355.029	-3.3339	240.1		
	5 _{4,1}	4 _{4,0}	355.029	-3.3339	240.1		
	*	12 _{1,11}	12 _{1,12}	355.042	-4.7277	285.4	■
	5 _{3,3}	4 _{3,2}	355.191	-3.0835	157.5		
**	5 _{3,2}	4 _{3,1}	355.203	-3.0834	157.5		
	5 _{2,3}	4 _{2,2}	356.176	-2.9618	98.5		
HDCO	5 _{1,4}	4 _{1,3}	335.097	-2.9841	56.2		
	17 _{2,15}	17 _{2,16}	340.801	-4.5839	517.6	■	
	3 _{1,3}	2 _{0,2}	343.284	-5.2876	25.8		
	11 _{1,10}	11 _{1,11}	346.739	-4.6725	219.4		
	4 _{2,2}	5 _{0,5}	347.286	-6.1800	62.9	■	
10 _{1,9}	10 _{0,10}	355.075	-5.2163	184.3	■		
HDC ¹⁸ O	6 _{1,6}	5 _{1,5}	353.098	-2.9039	67.4	■	
D ₂ CO	6 _{1,6}	5 _{1,5}	330.674	-2.9892	61.1		
	6 _{0,6}	5 _{0,5}	342.522	-2.9331	58.1		
	15 _{2,13}	15 _{2,14}	345.075	-4.4700	370.5	■	
	6 _{2,5}	5 _{2,4}	349.631	-2.9553	80.4		
	6 _{5,2}	5 _{5,1}	351.196	-3.4129	193.7	■	
	6 _{5,1}	5 _{5,0}	351.196	-3.4129	193.7	■	
	6 _{4,3}	5 _{4,2}	351.487	-3.1522	145.2		
	6 _{4,2}	5 _{4,1}	351.492	-3.1522	145.2		
	6 _{3,4}	5 _{3,3}	351.894	-3.0204	107.6		
	6 _{3,3}	5 _{3,2}	352.244	-3.0192	107.6		
	6 _{2,4}	5 _{2,3}	357.871	-2.9248	81.2		
	D ₂ ¹³ CO	6 _{0,6}	5 _{0,5}	337.552	-2.9461	57.2	
		6 _{2,5}	5 _{2,4}	344.225	-2.9696	79.6	■
		6 _{4,3}	5 _{4,2}	345.957	-3.1670	144.7	■
		6 _{4,2}	5 _{4,1}	345.961	-3.1670	144.7	■
6 _{3,4}		5 _{3,3}	346.346	-3.0352	106.9		
6 _{3,3}		5 _{3,2}	346.662	-3.0340	106.9		
6 _{2,4}		5 _{2,3}	351.961	-2.9406	80.3	■	
6 _{1,5}		5 _{1,4}	360.989	-2.8691	66.2	■	

Table A.2. Lines not detected but used to constrain the column densities.

Species	Q ₁	Q ₂	ν	A _{ij}	E _u
	$J_{K_A K_B}$		GHz	log ₁₀ (s^{-1})	K
H ₂ CO	28 _{3,25}	28 _{3,26}	336.14	-4.6821	1542.1
H ₂ C ¹⁷ O	18 _{3,15}	19 _{1,18}	354.222	-6.0022	691.4
	12 _{1,11}	12 _{1,12}	355.087	-4.7276	285.4
H ₂ C ¹⁸ O	12 _{1,11}	12 _{1,12}	339.498	-4.7872	279.1
H ₂ ¹³ CO	29 _{3,26}	29 _{3,27}	349.48	-4.6636	1606.4
HDCO	24 _{3,21}	24 _{3,22}	348.965	-4.5321	1017.8
HDC ¹⁸ O	2 _{2,0}	3 _{1,3}	332.193	-6.2815	41.0
D ₂ CO	7 _{3,4}	8 _{1,7}	340.380	-6.1185	127.4




Fig. B.1. All lines of (a) H₂CO, (b) H₂C¹⁷O, and (c) H₂C¹⁸O with the best fit LTE model overplotted in blue. Synthetic spectra in red indicate optical depth beginning to play a significant role i.e. $\tau > 0.6$. Numbers in top right corner with red boxes indicate lines used to fit the column density. Note that the two optically thick lines of H₂CO are the ones with a significant amount of extended emission (covered in van der Wiel, in prep.). The number in parenthesis is the estimated optical depth. Lines in figure (b) panel 5 and 8 have stronger H₂¹³CO lines blended (see panel 9 and 11 in Fig. B.2).




Fig. B.2. All lines of H₂¹³CO with the best fit LTE model overplotted. Synthetic spectra in red indicate optical depth beginning to play a significant role i.e. $\tau > 0.6$. Numbers in top right corner with red boxes indicate lines used to fit the column density. The number in parenthesis is the estimated optical depth. The line in panels 9 and 11, at 355.04 and 355.20 GHz could be blended with significantly weaker H₂C¹⁷O lines (see panel 5 and 8 in Fig. B.1).




Fig. B.3. All the lines of HDCO with the best fit LTE model overplotted. Synthetic spectra in red indicate optical depth beginning to play a significant role i.e. $\tau > 0.6$. Numbers in top right corner with red boxes indicate lines used to fit the column density. The number in parenthesis is the estimated optical depth.




Fig. B.4. All the lines of HDC^{18}O with the best fit LTE model overplotted. Synthetic spectra in red indicate optical depth beginning to play a significant role i.e. $\tau > 0.6$. Numbers in top right corner with red boxes indicate lines used to fit the column density. The number in parenthesis is the estimated optical depth. The emission line at 350.10 GHz is probably methanol.




Fig. B.5. All the lines of D_2CO with the best fit LTE model overplotted. Synthetic spectra in red indicate optical depth beginning to play a significant role i.e. $\tau > 0.6$. Numbers in top right corner with red boxes indicate lines used to fit the column density. The emission line at 332.96 GHz is possibly glycolaldehyde and ethylene glycol, and at 357.91 GHz methyl formate.




Fig. B.6. All the lines of $D_2^{13}CO$ with the best fit LTE model overplotted. Synthetic spectra in red indicate optical depth beginning to play a significant role i.e. $\tau > 0.6$. Numbers in top right corner with red boxes indicate lines used to fit the column density. The number in parenthesis is the estimated optical depth.