



About recurrence time for a semi-Lagrangian discontinuous Galerkin Vlasov solver

Laurent Navoret, Michel Mehrenberger

► To cite this version:

Laurent Navoret, Michel Mehrenberger. About recurrence time for a semi-Lagrangian discontinuous Galerkin Vlasov solver. Collisionless Boltzmann (Vlasov) Equation and Modeling of Self-Gravitating Systems and Plasmas, Oct 2017, Marseille, France. hal-01653023

HAL Id: hal-01653023

<https://hal.science/hal-01653023>

Submitted on 30 Nov 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Introduction

Cheng-Knorr, 1976

Comments and objective

The SLDG method for
Vlasov-Poisson

history

The scheme

Free streaming for
Gauss type mesh

Recurrence & quadrature
Trigonometric quadrature

Numerical results

free streaming

linear Landau damping

Plasma echo

Conclusion and
perspectives

About recurrence time for a semi-Lagrangian discontinuous Galerkin Vlasov solver

Michel Mehrenberger

IRMA, Université de Strasbourg & projet TONUS (INRIA), France

CIRM, 31.10.2017

Joint work with Laurent Navoret (IRMA)

Introduction : results from Cheng-Knorr, 1976



Recurrence time for
SLDG

M. Mehrenberger

Recurrence time $T_R = \frac{2\pi}{k\Delta v}$ is analyzed for free streaming case...

$$\partial_t f + v \partial_x f = 0, \rho = \int f dv, f(t=0) = A \cos(kx) f_0(v), f_0(v) = \frac{1}{\sqrt{2\pi}} e^{-v^2/2}$$

$$\rho(t, x) = A \Delta v \sum_{j=-M}^{M-1} f_0(v_j) \cos(k(x - (j + 1/2)\Delta vt)) \text{ is } T_R \text{ periodic in time}$$



Introduction

Cheng-Knorr, 1976

Comments and objective

The SLDG method for
Vlasov-Poisson

history

The scheme

Free streaming for
Gauss type mesh

Recurrence & quadrature

Trigonometric quadrature

Numerical results

free streaming

linear Landau damping

Plasma echo


Conclusion and
perspectives

Introduction : results from Cheng-Knorr 1976

...and observed for the simulation of the linear Landau damping

$$\partial_t f + v \partial_x f + E \partial_v f = 0, \quad \partial_x E = \rho - 1$$

$$f(t=0) = (1 + A \cos(kx)) f_0(v)$$



Recurrence time for
SLDG

M. Mehrenberger

Introduction

Cheng-Knorr, 1976

Comments and objective

The SLDG method for
Vlasov-Poisson

history

The scheme

Free streaming for
Gauss type mesh

Recurrence & quadrature
Trigonometric quadrature

Numerical results

free streaming
linear Landau damping
Plasma echo

Conclusion and
perspectives

Comments and objective

Recurrence time for
SLDG

M. Mehrenberger

Introduction

Cheng-Knorr, 1976

Comments and objective

The SLDG method for Vlasov-Poisson

history

The scheme

Free streaming for Gauss type mesh

Recurrence & quadrature

Trigonometric quadrature

Numerical results

free streaming

linear Landau damping

Plasma echo

Conclusion and perspectives

The SLDG method for Vlasov-Poisson : history

Recurrence time for
SLDG

M. Mehrenberger

Introduction

Cheng-Knorr, 1976

Comments and objective

The SLDG method for Vlasov-Poisson

history

The scheme

Free streaming for Gauss type mesh

Recurrence & quadrature

Trigonometric quadrature

Numerical results

free streaming

linear Landau damping

Plasma echo

Conclusion and perspectives

- ▶ Rossmanith-Seal → Qiu-Shu → Crouseilles-M-Vencil (2010-2011)
- ▶ already ideas of SLDG in Mangeney-Califano-et-al (2002)
- ▶ variant+analysis : Einkemmer-Ostermann (2014)
- ▶ adaptive : Besse-Deriaz-Madaule (2016)
- ▶ no splitting : Cai-Guo-Qiu (2017)
- ▶ other contexts :
 - ▶ general approach : Restelli-Bonaventuro-Sacco (2006)
 - ▶ Guo-Nair-Qiu (2014)
 - ▶ analysis : Bokanowski-Simarmata, Bokanowski-Cheng-Shu (2016)
- ▶ some ref. for DG for Vlasov :
 - ▶ Heath-Gamba-et-al (2012)
 - ▶ Cheng-Christlieb-Zhong, Madaule-Restelli-Sonnendrücker (2014)
 - ▶ analysis : Ayuso-Carillo-Shu (2012)

The SLDG method for Vlasov-Poisson : the scheme

Recurrence time for
SLDG

M. Mehrenberger

► In 1D for free streaming

- ▶ polynomial representation on each cell
- ▶ exact transport (constant velocity)
- ▶ projection on each cell : piecewise polynomial → polynomial
- ▶ can be interpreted as advection of Gauss points¹

$$x_{j,0}, \dots, x_{j,d} \in]x_{\min} + j\Delta x, x_{\min} + (j+1)\Delta x[$$

and written as

$$\begin{pmatrix} f(t + \Delta t, x_{j,0}) \\ \dots \\ f(t + \Delta t, x_{j,d}) \end{pmatrix} = A(\alpha) \begin{pmatrix} f(t, x_{i,0}) \\ \dots \\ f(t, x_{i,d}) \end{pmatrix} + B(\alpha) \begin{pmatrix} f(t, x_{i+1,0}) \\ \dots \\ f(t, x_{i+1,d}) \end{pmatrix}$$

with $x_{j,\cdot} - v\Delta t = x_{i,\cdot} + \alpha\Delta x$ and $A(\alpha), B(\alpha) \in \mathbb{R}^{d+1} \times \mathbb{R}^{d+1}$

► In $1D \times 1D$ phase-space for Vlasov-Poisson

- ▶ polynomial represented by Gauss points on each rectangular cell
- ▶ splitting : succession of
 - ▶ 1D advects of Gauss points in x direction
(each 1D advection updates its horizontal line)
 - ▶ Poisson
 - ▶ 1D advects of Gauss points in v direction
(each 1D advection updates its vertical line)

Introduction

Cheng-Knorr, 1976

Comments and objective

The SLDG method for
Vlasov-Poisson

history

The scheme

Free streaming for
Gauss type mesh

Recurrence & quadrature

Trigonometric quadrature

Numerical results

free streaming

linear Landau damping

Plasma echo

Conclusion and
perspectives

Free streaming for Gauss type mesh

Recurrence time for
SLDG

M. Mehrenberger

The mesh in velocity is given by

$$v_{j,\ell} = (j + \bar{v}_\ell) \Delta v, \ell = 0, \dots, d$$

with $0 < \bar{v}_0 < \dots < \bar{v}_d < 1$ and we solve

$$\partial_t f + v_{j,\ell} \partial_x f = 0$$

- ▶ Framework not only for SLDG : what counts here is the mesh
- ▶ Do we have a recurrence time ?
- ▶ is it formula $T_R = 2\pi/(k\Delta v)$?
 - ▶ OK for $d = 0$ and $\bar{v}_0 = 1/2$.
 - ▶ but looking at uniform cases :
 - ▶ $d = 1$ and $\bar{v}_0 = 1/4, \bar{v}_1 = 3/4, \rightarrow T_R = 2\pi/(k\Delta v/2)$
 - ▶ $d = 2$ and $\bar{v}_0 = 1/6, \bar{v}_1 = 3/6, \bar{v}_2 = 5/6, \rightarrow T_R = 2\pi/(k\Delta v/3)$
 - ▶ ... $\bar{v}_k = (2\ell + 1)/(2d + 2), \ell = 0, \dots, d$
- ▶ we can hope formula $T_R = 2\pi(d + 1)/(k\Delta v)$
- ▶ We will see that formula is between :

$$T_R \simeq 2\pi(\lfloor d/2 \rfloor + 1)/(k\Delta v)$$

Introduction

Cheng-Knorr, 1976

Comments and objective

The SLDG method for
Vlasov-Poisson

history

The scheme

Free streaming for
Gauss type mesh

Recurrence & quadrature

Trigonometric quadrature

Numerical results

free streaming

linear Landau damping

Plasma echo

Conclusion and
perspectives

Recurrence & quadrature

Recurrence time for
SLDG

M. Mehrenberger

We get

$$\rho(t, x) = A\Delta v \sum_{j=-M}^{M-1} \sum_{\ell=0}^d \omega_\ell f_0(v_{j,\ell}) \cos(k(x - (j + \bar{v}_\ell)\Delta vt))$$

With symmetry of discrete velocity distribution, we obtain

$$\rho(t, x) = A \cos(kx) h(t),$$

with

$$h(t) = \Delta v \sum_{j=-M}^{M-1} \sum_{\ell=0}^d \omega_\ell f_0(v_{j,\ell}) \cos(k(j + \bar{v}_\ell)\Delta vt))$$

We look the expression for $t = t_m = \frac{2\pi m}{k\Delta v}$, $m \in \mathbb{N}^*$

$$h(t_m) = \sum_{\ell=0}^d \omega_\ell \cos(2\pi m \bar{v}_\ell) \Delta v \sum_{j=-M}^{M-1} f_0(v_{j,\ell})$$

Introduction

Cheng-Knorr, 1976

Comments and objective

The SLDG method for
Vlasov-Poisson

history

The scheme

Free streaming for
Gauss type mesh

Recurrence & quadrature

Trigonometric quadrature

Numerical results

free streaming

linear Landau damping

Plasma echo

Conclusion and
perspectives

Recurrence & quadrature

As $\Delta v \sum_{j=-M}^{M-1} f_0(v_{j,\ell}) \rightarrow \int f_0(v) dv$ exponentially, we have

$$h(t_m) = h_m \int f_0(v) dv + O(\Delta v^\rho), \quad h_m = \sum_{\ell=0}^d \omega_\ell \cos(2\pi m \bar{v}_\ell)$$

- ▶ Energy restituted at $t_m = \frac{2\pi m}{k\Delta v}$, $m \in \mathbb{N}^*$ is \simeq proportional to h_m .
- ▶ h_m is a quadrature formula for $\int_0^1 \cos(2\pi mv) dv$
- ▶ At continuous level, we have

$$\int_0^1 \cos(2\pi mv) dv = 0 : \text{no recurrence}$$

⇒ link between recurrence effect and quadrature rule

- ▶ For uniform mesh :
- ▶ $\cos(2\pi m \cdot)$ exactly reproduced by standard quadrature for $m = 0, 1, \dots, d$:

$$\frac{1}{d+1} \sum_{\ell=0}^d \cos\left(2\pi m \frac{2\ell+1}{2d+2}\right) = \begin{cases} 0, & \text{if } m = q(d+1) + r, \quad r = 1, \dots, d \\ (-1)^q, & \text{if } m = q(d+1). \end{cases}$$

- ▶ consistent with $T_R = 2\pi(d+1)/(k\Delta v)$.

We look for a quadrature rule such that

$$\sum_{\ell=0}^d \omega_\ell \cos(2\pi m \bar{v}_\ell) = \int_0^1 \cos(2\pi mv) dv,$$

for all $m \leq n$ with n as high as possible.

- ▶ Possible for $n = \lfloor d/2 \rfloor$ and

$$\omega_\ell = \frac{1}{2\pi} \int_{-\pi}^{\pi} \prod_{j=0, j \neq \ell}^{\lfloor d/2 \rfloor} \frac{\cos(v) - \cos(2\pi \bar{v}_j)}{\cos(2\pi \bar{v}_\ell) - \cos(2\pi \bar{v}_j)} dv, \quad \ell = 0, \dots, d.$$

- ▶ Optimal points are uniform and lead to $n = d$ (like Gauss integration)
 - ▶ for Gauss points, weights remain positive until $d = 18$.
 - ▶ recent references : Peherstorfer, 2011 ; Austin, PhD 2016
- ⇒ Instead of using classical *polynomial* integration quadrature rule,
we can use this *trigonometric* integration

Introduction

Cheng-Knorr, 1976

Comments and objective

The SLDG method for
Vlasov-Poisson

history

The scheme

Free streaming for
Gauss type mesh

Recurrence & quadrature

Trigonometric quadrature

Numerical results

free streaming

linear Landau damping

Plasma echo

Conclusion and
perspectives

Numerical results for free streaming

$d = 2$ (gauss points)

	classical	new
h_1	0.02245	0
h_2	0.52996	0.51916
h_3	0.73652	0.73047

$d = 3$ (gauss points)

	classical	new
h_1	$-1.0678 \cdot 10^{-3}$	0
h_2	-0.12573	-0.12664
h_3	-0.74097	-0.74155
h_4	-0.33811	-0.33830

$d = 4$ (gauss points)

	classical	new
h_1	$3.0803 \cdot 10^{-5}$	0
h_2	0.016669	0
h_3	0.30402	0.30657
h_4	0.79870	0.80082
h_5	-0.010458	-0.018835

Introduction

Cheng-Knorr, 1976

Comments and objective

The SLDG method for
Vlasov-Poisson

history

The scheme

Free streaming for
Gauss type mesh

Recurrence & quadrature

Trigonometric quadrature

Numerical results

free streaming

linear Landau damping

Plasma echo

Conclusion and
perspectives

Numerical results for free streaming

Recurrence time for
SLDG

M. Mehrenberger

$d = 5$ (gauss points)

	classical	new
h_1	$-5.9704 \cdot 10^{-7}$	0
h_2	$-1.4223 \cdot 10^{-3}$	0
h_3	-0.071038	-0.073117
h_4	-0.50843	-0.50769
h_5	-0.67454	-0.67455
h_6	0.30953	0.30852

$d = 6$ (gauss points)

	classical	new
h_1	$8.3166 \cdot 10^{-9}$	0
h_2	$8.5037 \cdot 10^{-5}$	0
h_3	0.010947	0
h_4	0.17907	0.18741
h_5	0.66873	0.66483
h_6	0.40512	0.40575
h_7	-0.41364	-0.41456

Introduction

Cheng-Knorr, 1976

Comments and objective

The SLDG method for
Vlasov-Poisson

history

The scheme

Free streaming for
Gauss type mesh

Recurrence & quadrature

Trigonometric quadrature

Numerical results

free streaming

linear Landau damping

Plasma echo

Conclusion and
perspectives

Numerical results for free streaming

Recurrence time for
SLDG

M. Mehrenberger

Introduction

Cheng-Knorr, 1976

Comments and objective

The SLDG method for
Vlasov-Poisson

history

The scheme

Free streaming for
Gauss type mesh

Recurrence & quadrature

Trigonometric quadrature

Numerical results

free streaming

linear Landau damping

Plasma echo

Conclusion and
perspectives

- ▶ Gauss quadrature (classical) already good
- ▶ Gauss quadrature becomes better when d increases
- ▶ small improvement for new trigonometric quadrature
- ▶ do not use when $d > 18$ (negative weights)
- ▶ T_R coherent with result of Madaule-Restelli-Sonnendrücker (2014)

Gauss-Lobatto points

The analysis can be extended to Gauss-Lobatto points

Example : $d = 2$, the mesh is uniform :

$$x_0 = 0, x_1 = 1/2, x_2 = 1$$

- ▶ Trigonometric interpolation corresponds to trapezoidal formula :

$$\omega_0 = \omega_2 = 1/4, \omega_1 = 1/2$$

- ▶ Gauss-Lobatto quadrature is Simpson rule :

$$\omega_0 = \omega_2 = 1/6, \omega_1 = 2/3$$

$d = 2$ (gauss-lobatto points)

	classical	new
h_1	1/3	0
h_2	1	1
h_3	1/3	0
h_4	1	1

Introduction

Cheng-Knorr, 1976

Comments and objective

The SLDG method for
Vlasov-Poisson

history

The scheme

Free streaming for
Gauss type mesh

Recurrence & quadrature

Trigonometric quadrature

Numerical results

free streaming

linear Landau damping

Plasma echo

Conclusion and
perspectives

Numerical results for free streaming

Recurrence time for
SLDG

M. Mehrenberger

$d = 4$ (gauss-lobatto points)

	classical	new
h_1	$1.3174 \cdot 10^{-3}$	0
h_2	0.14855	0
h_3	0.79651	0.93393
h_4	0.25734	0.12776
h_5	-0.10028	-0.11932

$d = 6$ (gauss-lobatto points)

	classical	new
h_1	$6.9354 \cdot 10^{-7}$	0
h_2	$1.6300 \cdot 10^{-3}$	0
h_3	0.079357	0
h_4	0.542841	0.663454
h_5	0.64533	0.51383
h_6	-0.34435	-0.22468
h_7	0.15274	$2.1373 \cdot 10^{-3}$

positive weights until $d = 7$.

Introduction

Cheng-Knorr, 1976

Comments and objective

The SLDG method for
Vlasov-Poisson

history

The scheme

Free streaming for
Gauss type mesh

Recurrence & quadrature

Trigonometric quadrature

Numerical results

free streaming

linear Landau damping

Plasma echo


Conclusion and
perspectives

Numerical results for linear Landau damping

Recurrence time for
SLDG

M. Mehrenberger

$$N = 16, v_{\max} = 4, A = 10^{-2}, k = 0.5, 2\pi/(k\Delta v) \simeq 25.13$$



Introduction

Cheng-Knorr, 1976

Comments and objective

The SLDG method for
Vlasov-Poisson

history

The scheme

Free streaming for
Gauss type mesh

Recurrence & quadrature

Trigonometric quadrature

Numerical results

free streaming



linear Landau damping

Plasma echo

Conclusion and
perspectives

Numerical results for linear Landau damping

$N = 64$, $v_{\max} = 8$, $A = 10^{-6}$, $k = 0.5$, $2\pi/(k\Delta v) \simeq 50.26$



Introduction

Cheng-Knorr, 1976

Comments and objective

The SLDG method for Vlasov-Poisson

history

The scheme

Free streaming for Gauss type mesh

Recurrence & quadrature

Trigonometric quadrature

Numerical results

free streaming



linear Landau damping

Plasma echo

Conclusion and perspectives

Numerical results for plasma echo

$N = 600$, $v_{\max} = 8$, $A = 10^{-3}$, $k_1 = \frac{12\pi}{100}$ perturbation with $k_2 = \frac{24\pi}{100}$ at time $t = 200$ is added. $2\pi/(k_1 \Delta v) = 625$



Conclusion and perspectives

Recurrence time for
SLDG

M. Mehrenberger

Introduction

Cheng-Knorr, 1976

Comments and objective

The SLDG method for Vlasov-Poisson

history

The scheme

Free streaming for Gauss type mesh

Recurrence & quadrature

Trigonometric quadrature

Numerical results

free streaming

linear Landau damping

Plasma echo

Conclusion and perspectives

► Conclusion

- ▶ link between recurrence and quadrature
- ▶ introduction of trigonometric quadrature for non uniform velocity discretization
- ▶ analysis for free streaming
- ▶ numerical results for Landau damping

► Perspectives/possible applications

- ▶ velocity semi-discretization method (N. Pham, PhD 2016)
- ▶ curvilinear grid
- ▶ DG, NUFFT