

# Draft Genome Sequence of Pseudomonas graminis PDD-13b-3, a Model Strain Isolated from Cloud Water

Ludovic Besaury, P. Amato, N. Wirgot, M. Sancelme, A. M. Delort

## ▶ To cite this version:

Ludovic Besaury, P. Amato, N. Wirgot, M. Sancelme, A. M. Delort. Draft Genome Sequence of Pseudomonas graminis PDD-13b-3, a Model Strain Isolated from Cloud Water. Genome Announcements, 2017, 5 (26), pp.e00464-17. 10.1128/genomeA.00464-17. hal-01654897

# HAL Id: hal-01654897 https://hal.science/hal-01654897v1

Submitted on 13 Sep 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

#### PROKARYOTES



# Draft Genome Sequence of *Pseudomonas* graminis PDD-13b-3, a Model Strain Isolated from Cloud Water

gen@meAnnouncements™

### L. Besaury, P. Amato, N. Wirgot, M. Sancelme, A. M. Delort

AMERICAN SOCIETY FOR MICROBIOLOGY

CNRS, Université Clermont Auvergne, Institut de Chimie de Clermont-Ferrand, Clermont-Ferrand, France

**ABSTRACT** The whole genome of *Pseudomonas graminis* PDD-13b-3, a strain of bacteria isolated from cloud water, was sequenced. This showed that this microorganism is equipped with genes that could potentially be involved in its survival in the atmosphere and clouds: those for oxidative stress and carbon starvation responses, DNA repair, and iron uptake.

louds play a major role in the transformation of atmospheric compounds. They also host living microorganisms, despite high oxidative capacity and UV radiation levels, low temperatures, and rapid shifts of abiotic conditions causing physiological shocks (1-4). Among those microorganisms recovered alive from cloud water, bacteria affiliated with the genus Pseudomonas are the most frequently recovered (5). Pseudomonas graminis PDD-13b-3 (1) was recovered from a culture of cloud water sampled in July 2004 from Puy de Dôme Mountain's meteorological station (France), at 1,465 m altitude, on R2A agar medium incubated at 17°C under aerobic conditions. Once purified from the original colony, the strain was identified based on its 16S rRNA gene sequence (GenBank accession number DQ512786). Pseudomonas graminis PDD-13b-3 has the ability to develop at low temperature, degrade some organic compounds present in clouds (6), and produce siderophores to satisfy their iron requirements (7); it thus potentially interferes with cloud chemical reactivity. This strain was found to be ice-nucleation negative (8), but it produces biosurfactants known for facilitating spread on surfaces, and perhaps also participating in cell aerosolization and cloud formation (9). Pseudomonas graminis PDD-13b-3 appears thus to be a model of cloud water microorganisms. To further understand the specificities of Pseudomonas graminis PDD-13b-3, its ability to survive in cloud water, and the implications of its presence in clouds, its whole genome was sequenced and is reported here. Whole-genome shotgun sequencing (2  $\times$  150 bp) was prepared using the Nextera DNA sample preparation kit (Illumina, San Diego, CA, USA), according to the manufacturer's user guide, and sequenced on an Illumina MiSeq sequencer (MR DNA; Molecular Research, Shallowater, TX, USA). Sequence data files were filtered for quality using FastQC (https://www .bioinformatics.babraham.ac.uk/projects/fastqc/), trimmed using Prinseq-Lite (10), and then de novo assembled with SPAdes (11). A total of 102 contigs were generated, with an average coverage of 22.6-fold. The average contig size was 58,931 bp, and the  $N_{50}$ contig size was 152,216 bp. The size of the assembled genome is 5,686,785 bp, with a G+C content of 60.11%, which are within the ranges of known values for Pseudomonas genomes (12).

The draft genome of *Pseudomonas graminis* PDD-13b-3 was annotated using the RAST annotation server (http://rast.nmpdr.org). It contains 61 RNAs and 5,177 proteincoding genes, of which 52% were assigned to a total of 527 SEED subsystems. Among these SEED subsystems, 98 were affiliated with oxidative stress response (encoding NADPH:quinone oxidoreductase, glutathione synthetase, peroxidase, etc.) and 85 with DNA repair processes. Eleven protein-coding genes were related to iron acquisition and

## Received 20 April 2017 Accepted 12 May 2017 Published 29 June 2017

**Citation** Besaury L, Amato P, Wirgot N, Sancelme M, Delort AM. 2017. Draft genome sequence of *Pseudomonas graminis* PDD-13b-3, a model strain isolated from cloud water. Genome Announc 5:e00464-17. https://doi .org/10.1128/genomeA.00464-17.

**Copyright** © 2017 Besaury et al. This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International license.

Address correspondence to L. Besaury, ludovic.besaury@uca.fr.

metabolism, encoding iron siderophore (pyoverdine) and related sensor and receptor systems. Interestingly, 8 protein-coding genes were associated with carbon starvation. All these could have contributed to *Pseudomonas graminis* PDD-13b-3 survival in clouds.

**Accession number(s).** This whole-genome shotgun project has been deposited in the GenBank database under the accession number MTSB00000000.

#### ACKNOWLEDGMENTS

We thank Magali Abrantes for the DNA extraction of *Pseudomonas graminis* PDD-13b-3.

This study was funded by the project BIOCAP supported by the French National Research Agency (ANR).

#### REFERENCES

- Amato P, Parazols M, Sancelme M, Laj P, Mailhot G, Delort AM. 2007. Microorganisms isolated from the water phase of tropospheric clouds at the Puy de Dôme: major groups and growth abilities at low temperatures. FEMS Microbiol Ecol 59:242–254. https://doi.org/10.1111/j.1574 -6941.2006.00199.x.
- Vaïtilingom M, Amato P, Sancelme M, Laj P, Leriche M, Delort AM. 2010. Contribution of microbial activity to carbon chemistry in clouds. Appl Environ Microbiol 76:23–29. https://doi.org/10.1128/AEM.01127-09.
- Vaïtilingom M, Attard E, Gaiani N, Sancelme M, Deguillaume L, Flossmann Al, Amato P, Delort AM. 2012. Long-term features of cloud microbiology at the Puy de Dôme (France). Atmos Environ 56:88–100. https:// doi.org/10.1016/j.atmosenv.2012.03.072.
- Joly M, Amato P, Sancelme M, Vinatier V, Abrantes M, Deguillaume L, Delort AM. 2015. Survival of microbial isolates from clouds toward simulated atmospheric stress factors. Atmos Environ 117:92–98. https:// doi.org/10.1016/j.atmosenv.2015.07.009.
- Amato P, Ménager M, Sancelme M, Laj P, Mailhot G, Delort AM. 2005. Microbial population in cloud water at the Puy de Dôme: implications for the chemistry of clouds. Atmos Environ 39:4143–4153. https://doi.org/ 10.1016/j.atmosenv.2005.04.002.
- Amato P, Demeer F, Melaouhi A, Fontanella S, Martin-Biesse A-S, Sancelme M, Laj P, Delort A-M. 2007. A fate for organic acids, formaldehyde and methanol in cloud water: their biotransformation by microorganisms. Atmos Chem Phys 7:4159–4169. https://doi.org/10.5194/acp -7-4159-2007.

- Vinatier V, Wirgot N, Joly M, Sancelme M, Abrantes M, Deguillaume L, Delort AM. 2016. Siderophores in cloud waters and potential impact on atmospheric chemistry: production by microorganisms isolated at the Puy de Dôme station. Environ Sci Technol 50:9315–9323. https://doi.org/ 10.1021/acs.est.6b02335.
- Joly M, Attard E, Sancelme M, Deguillaume L, Guilbaud C, Morris CE, Amato P, Delort AM. 2013. Ice nucleation activity of bacteria isolated from cloud water. Atmos Environ 70:392–400. https://doi.org/10.1016/j .atmosenv.2013.01.027.
- Renard P, Canet I, Sancelme M, Wirgot N, Deguillaume L, Delort AM. 2016. Screening of cloud microorganisms isolated at the Puy de Dôme (France) station for the production of biosurfactants. Atmos Chem Phys 16:12347–12358. https://doi.org/10.5194/acp-16-12347-2016.
- Schmieder R, Edwards R. 2011. Quality control and preprocessing of metagenomic datasets. Bioinformatics 27:863–864. https://doi.org/10 .1093/bioinformatics/btr026.
- Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Prjibelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. J Comput Biol 19:455–477.
- Silby MW, Winstanley C, Godfrey SA, Levy SB, Jackson RW. 2011. Pseudomonas genomes: diverse and adaptable. FEMS Microbiol Rev 35: 652–680.