



HAL
open science

Draft Genome Sequence of *Rhodococcus enclensis* 23b-28, a Model Strain Isolated from Cloud Water

A. Lallement, Ludovic Besaury, B. Eyheraguibel, P. Amato, M. Sancelme, G. Mailhot, A. M. Delort

► **To cite this version:**

A. Lallement, Ludovic Besaury, B. Eyheraguibel, P. Amato, M. Sancelme, et al.. Draft Genome Sequence of *Rhodococcus enclensis* 23b-28, a Model Strain Isolated from Cloud Water. *Genome Announcements*, 2017, 5 (43), pp.e01199-17. 10.1128/genomeA.01199-17 . hal-01654935

HAL Id: hal-01654935

<https://hal.science/hal-01654935v1>

Submitted on 13 Sep 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Draft Genome Sequence of *Rhodococcus enclensis* 23b-28, a Model Strain Isolated from Cloud Water

A. Lallement, L. Besaury, B. Eyheraguibel, P. Amato, M. Sancelme, G. Mailhot, A. M. Delort

Université Clermont Auvergne, CNRS, Institut de Chimie de Clermont-Ferrand, Clermont-Ferrand, France

ABSTRACT The whole genome of *Rhodococcus enclensis* 23b-28, a bacterial strain isolated from cloud water, was sequenced. This microorganism is equipped with genes able to degrade aromatic compounds and could thus play a role in complex organic matter decomposition in cloud water.

Phenol can be found in all environmental compartments (soil and water), including the atmosphere (1, 2). Due to its high volatility, phenol is present in the gas phase, but this compound can also be transferred to the aqueous phase of the atmosphere (rain, snow, and clouds) thanks to its solubility (defined by Henry's law constant). The degradation of organic matter in cloud water occurs through photochemical reactions due to solar light, but also through enzymatic reactions performed by viable microorganisms (3) that can survive in clouds despite the numerous stresses they encounter (oxidative, radiative, and osmotic stresses) (4). Among the different strains isolated from cloud water from the Puy de Dôme station (1,465 m above sea level, France) the strain *Rhodococcus enclensis* 23b-28 was one of the most efficient for phenol degradation. This strain has been isolated aerobically, has the ability to develop at low temperature (17°C), and can produce biosurfactants (5). To further understand the specificities of this strain, its ability to degrade aromatic compounds, its presence, and the implications for cloud biochemistry, its whole genome was sequenced. Whole-genome shotgun sequencing (2×150 bp) was prepared using the Nextera DNA sample preparation kit (Illumina, San Diego, CA, USA), following the manufacturer's user guide, and sequencing was done on an Illumina MiSeq sequencer (MR DNA, Shallowater, TX, USA). Sequence data files were filtered for quality using FastQC (<https://www.bioinformatics.babraham.ac.uk/projects/fastqc/>), trimmed using Prinseq-Lite (6), and *de novo* assembled with SPAdes (7). A total of 118 contigs were generated, with an average coverage of 23.1-fold. The average contig size was 61,133 bp, and the N_{50} contig size was 158,305 bp. The size of the assembled genome is 7,213,744 bp, with a GC content of 62.3% (this is a low value for *R. enclensis*, for which the GC content is 66.9% [8], but it is in the range of those obtained for *Rhodococcus* species in general, at 62 to 69% [9]). The RAST annotation server was used for annotating the draft genome of *Rhodococcus enclensis* 23b-28 (<http://rast.nmpdr.org>). It contains 59 RNAs and 6,755,000 protein-coding genes, of which 34% were assigned to a total of 2,367 subsystems; among those, 96 were affiliated with aromatic degradation. Genetic sequences affiliated with enzymes involved in benzoate degradation were found, along with the two degradation pathways for catechol (ortho- and meta-). This information allows us to think that *Rhodococcus enclensis* 23b-28 can play a role in the degradation of aromatic compounds in clouds.

Accession number(s). This whole-genome shotgun project has been deposited in the GenBank database under the accession number [NOVD00000000](https://ncbi.nlm.nih.gov/nucl/NOVD00000000).

Received 25 September 2017 Accepted 2 October 2017 Published 26 October 2017

Citation Lallement A, Besaury L, Eyheraguibel B, Amato P, Sancelme M, Mailhot G, Delort AM. 2017. Draft genome sequence of *Rhodococcus enclensis* 23b-28, a model strain isolated from cloud water. *Genome Announc* 5:e01199-17. <https://doi.org/10.1128/genomeA.01199-17>.

Copyright © 2017 Lallement et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to A. M. Delort, a-marie.delort@uca.fr.

ACKNOWLEDGMENT

This study was funded by the project BIOCAP, supported by the French National Research Agency (ANR).

REFERENCES

1. Atkinson R, Aschmann SM, Arey J. 1992. Reactions of OH and NO₃ radicals with phenol, cresols, and 2-nitrophenol at 296 ± 2K. *Environ Sci Technol* 26:1397–1403. <https://doi.org/10.1021/es00031a018>.
2. Rubio MA, Lissi E, Herrera N, Pérez V, Fuentes N. 2012. Phenol and nitrophenols in the air and dew waters of Santiago de Chile. *Chemosphere* 86:1035–1039. <https://doi.org/10.1016/j.chemosphere.2011.11.046>.
3. Vaitilingom M, Attard E, Gaiani N, Sancelme M, Deguillaume L, Flossmann AI, Amato P, Delort A-M. 2012. Long-term features of cloud microbiology at the Puy de Dôme (France). *Atmos Environ* 56:88–100. <https://doi.org/10.1016/j.atmosenv.2012.03.072>.
4. Delort A-M, Vaitilingom M, Amato P, Sancelme M, Parazols M, Mailhot G, Laj P, Deguillaume L. 2010. A short overview of the microbial population in clouds: potential roles in atmospheric chemistry and nucleation processes. *Atmos Res* 98:249–260. <https://doi.org/10.1016/j.atmosres.2010.07.004>.
5. Renard P, Canet I, Sancelme M, Wirgot N, Deguillaume L, Delort A-M. 2016. Screening of cloud microorganisms isolated at the Puy de Dôme (France) station for the production of biosurfactants. *Atmos Chem Phys* 16:12347–12358. <https://doi.org/10.5194/acp-16-12347-2016>.
6. Schmieder R, Edwards R. 2011. Quality control and preprocessing of metagenomic datasets. *Bioinformatics* 27:863–864. <https://doi.org/10.1093/bioinformatics/btr026>.
7. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Prjibelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <https://doi.org/10.1089/cmb.2012.0021>.
8. Dastager SG, Mawlankar R, Tang SK, Krishnamurthi S, Ramana VV, Joseph N, Shouche YS. 2014. *Rhodococcus enclensis* sp. nov., a novel member of the genus *Rhodococcus*. *Int J Syst Evol Microbiol* 64:2693–2699. <https://doi.org/10.1099/ijs.0.061390-0>.
9. Alvarez HM. 2010. Overview of *Rhodococcus* genome, p. 76–82. *In* *Biology of Rhodococcus*. Microbiology Monographs. Springer, Berlin, Germany.