



**HAL**  
open science

# Derepression of Excision of Integrative and Potentially Conjugative Elements from *Streptococcus thermophilus* by DNA Damage Response: Implication of a cI-Related Repressor

Xavier Bellanger, Catherine Morel, Bernard Decaris, G. Guedon

► **To cite this version:**

Xavier Bellanger, Catherine Morel, Bernard Decaris, G. Guedon. Derepression of Excision of Integrative and Potentially Conjugative Elements from *Streptococcus thermophilus* by DNA Damage Response: Implication of a cI-Related Repressor. *Journal of Bacteriology*, 2007, 189 (4), pp.1478 - 1481. 10.1128/JB.01125-06 . hal-01659457

**HAL Id: hal-01659457**

**<https://hal.science/hal-01659457v1>**

Submitted on 8 Dec 2017

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Derepression of Excision of Integrative and Potentially Conjugative Elements from *Streptococcus thermophilus* by DNA Damage Response: Implication of a *cI*-Related Repressor<sup>∇</sup>

Xavier Bellanger, Catherine Morel, Bernard Decaris, and Gérard Guédon\*

Laboratoire de Génétique et Microbiologie UMR1128, INRA, IFR110, Université Henri Poincaré Nancy 1, F-54506 Vandœuvre-lès-Nancy, France

Received 27 July 2006/Accepted 9 November 2006

**A DNA-damaging agent, mitomycin C, derepresses the site-specific excision of two integrative and potentially conjugative elements from *Streptococcus thermophilus*, ICES*t1* and ICES*t3*. The regulation pathway involves a repressor related to phage lambda *cI* repressor. It could also involve a putative regulator related to another type of phage repressors, the “*cI*-like” repressors.**

Whereas *in silico* analyses revealed that numerous genomic islands could be integrative and conjugative elements (ICEs) or elements derived from ICEs (2, 4, 9, 13, 15, 19, 25, 26), only a few ICEs have been described. ICEs excise by site-specific recombination, transfer through conjugation, and integrate into a replicon of the recipient cell (8). Whereas the regulation of the excision of numerous prophages is well known, the regulation of site-specific excision of only a few ICEs, including Tn916, ICEs from *Bacteroides*, pSAM2, and *clc*, has been described (6, 11, 22, 23). These few regulation systems are very different from each other. Recently, DNA-damaging agents were found to derepress the excision and transfer of two other ICE types, ICE*Bs1* from *Bacillus subtilis* (1) and IncJ elements, including SXT from *Vibrio cholerae* (3) and SXT-related elements from enterobacteria (18). Such regulation systems are similar to the derepression of the site-specific excision of numerous prophages by DNA damage.

Two putative ICEs, ICES*t1* and ICES*t3*, are integrated in the 3' end of the *fda* locus of the lactic acid bacteria *Streptococcus thermophilus* CNRZ368 and CNRZ385, respectively (Fig. 1A). These ICEs harbor almost identical recombination and conjugation modules (20). The tyrosine integrase and the excisionase encoded by the ICES*t1* recombination module catalyze its excision by recombination between the *attL* and *attR* flanking sites, leading to an excised circular ICE harboring an *attI* site and to a chromosomal *attB* site (10). Furthermore, ICES*t1* carries an internal recombination site related to *attL*, *attL'* (Fig. 1A) (20). Recombination between *attL'* and *attR* leads to excision of the circular form of a shorter putative ICE, ICES*t2*, carrying an *attI'* site. A genomic island corresponding to the left part of ICES*t1* and flanked by *attL* and *attB'* sites remains integrated in *fda*.

The closely related regulation modules of ICES*t1*/ICES*t2* and ICES*t3* contain three shared open reading frames (ORFs)

(*arp1*, *orfQ*, and *arp2*) (Fig. 1B). The ICES*t3* recombination module also includes three specific ORFs or pseudogenes (*orf385A*, *orf385B*, and  $\Delta$ *orf385C*), and the ICES*t1*/ICES*t2* recombination module contains two specific ORFs (*orfP* and *orfR*). The putative regulatory proteins encoded by *arp1*, *arp2*, and *orf385A* have a helix-turn-helix (HTH) DNA binding domain. The functions of the putative proteins encoded by the other ORFs are unknown.

The 5' part of *arp1* encodes an HTH domain, and its 3' part encodes a region characteristic of the COG2932 protein family, including the *cI* repressor of phage  $\lambda$  (9, 20). This region has two functions, *cI* autoproteolysis and *cI* oligomerization. In the presence of damaged DNA, the RecA protein, activated by single-stranded DNA (RecA\*), induces autoproteolysis of *cI* and related proteins. Cleaved proteins are not able to oligomerize and therefore are not able to repress their target genes (12). *In silico* analysis suggested that DNA damage could derepress excision of ICEs from *S. thermophilus*.

To test this hypothesis, strains harboring ICES*t1*/ICES*t2* (CNRZ368) or ICES*t3* (CNRZ385) were grown at 42°C in HJL medium (24). Exponentially growing cells (optical density at 600 nm for MIC/4 and MIC/2, 0.04; optical density at 600 nm for 2× MIC and 4× MIC, 0.4) were treated with mitomycin C (MC) concentrations close to the MIC for 2.5 h to induce DNA damage. Then the recombination sites resulting from excision, *attI* and *attB* (ICES*t1* and ICES*t3*) or *attI'* (ICES*t2*), were amplified by PCR using 1  $\mu$ g genomic DNA for CNRZ368 and 1 ng genomic DNA for CNRZ385. PCR experiments were performed in 25- $\mu$ l mixtures using 0.5 U of *Taq* DNA polymerase (Biolabs) according to the manufacturer's specifications and primers (melting temperatures, 49.6 to 53.1°C) described in the legend to Fig. 1. After an initial denaturation step consisting of 4 min at 95°C, PCR was performed for 30 cycles of denaturation for 30 s at 95°C, annealing for 30 s at 47.5°C, and extension for 30 s at 72°C, followed by final extension for 7 min at 72°C.

Treatment of CNRZ368 with MC induced an increase in the PCR signal intensity corresponding to the *attB* and *attI* sites, while this treatment did not induce an increase in the PCR signal intensity corresponding to the *fda* gene used as a control (Fig. 2). The most intense signal for both *attB* and *attI* sites was

\* Corresponding author. Mailing address: Laboratoire de Génétique et Microbiologie UMR1128, INRA, Faculté des Sciences et Techniques, Université Henri Poincaré Nancy 1, 1 Bd des Aiguillettes, BP239, F-54506 Vandœuvre-lès-Nancy, France. Phone: (33) 03 83 68 49 72. Fax: (33) 03 83 68 11 76. E-mail: guedon@nancy.inra.fr.

<sup>∇</sup> Published ahead of print on 17 November 2006.




FIG. 1. Comparison of *ICES1/ICES2* and *ICES3* maps. (A) Location of the recombination, conjugation, and regulation modules and the recombination sites on the *ICES1/ICES2* and *ICES3* maps. The recombination sites are indicated by solid rectangles and are magnified. The schematic localizations and orientations of oligonucleotides used for PCRs are indicated by arrowheads. The following primer pairs were used for amplification of fragments containing the recombination sites resulting from ICE excision: primers O131.2 and O132.3 (*attB* from *ICES1* and *ICES3*), primers *attI2* and *attI3* (*attI* from *ICES1*), primers *attI2* and O74.1 (*attI'* from *ICES2*), and primers O131.5 and 385.2 (*attI* from *ICES3*). The *fda1-fda2* primer pair was used for amplification of an *fda* internal region. The dotted rectangle indicates the area shown in panel B. (B) Map of regulation modules of *ICES1/ICES2* and *ICES3*. ORF designations beginning with “orf” are abbreviated with the corresponding letter or name. The locations and orientations of ORFs and truncated ORFs ( $\Delta 385C$ ) belonging to the regulation modules are indicated by arrow boxes. The angled arrows and the lollipops indicate the putative promoters and rho-independent transcription terminators deduced from in silico analyses. The gray areas join closely related modules and *att* sites belonging to the ICES (>65% nucleotide identity); the levels of identity are indicated. The close relationships between short sequences and insertion elements are not shown.

obtained when 0.1  $\mu\text{g/ml}$  MC (MIC/2) was used. The effects of MC were also determined by PCR amplification of fragments harboring *attI* and *attB*, using 1  $\mu\text{g}$  to 1 ng genomic CNRZ368 DNA. The minimum amounts of template DNA that resulted




FIG. 2. Effects of MC treatment of exponentially growing cells on the excision frequency of *ICES1*. Fragments containing the *attB* (361 bp) or *attI* (400 bp) recombination site were amplified using 1  $\mu\text{g}$  template DNA from MC-treated cells and primers described in the legend to Fig. 1. An internal fragment of the *fda* gene (339 bp) was amplified as a control using 1 ng template DNA.

in positive PCR amplification of the *attB* and *attI* sites were 100 ng using CNRZ368 DNA in the absence of MC and 10 ng after treatment with MC (MIC/2), suggesting that DNA damage caused by MC induced at least a 10-fold increase in *ICES1* excision.

In the absence of MC, the following ratios of the excised circular form to the genome were found in the stationary phase of growth:  $10^{-6}$  for *ICES1*,  $<10^{-6}$  for *ICES2*, and  $9 \times 10^{-3}$  for *ICES3* (20). However, treatment of the strains harboring *ICES2* or *ICES3* with MC (MIC/2) was also found to induce excision (data not shown). This suggests that DNA damage induced excision of the three ICES.

In order to examine the *arp1* function, two fragments containing the 5' and 3' ends of *arp1* of *ICES1/ICES2* were cloned in the thermosensitive vector pG<sup>+</sup>host9 (5, 17). The recombinant plasmid, carrying a very short deleted ORF,  $\Delta arp1$ , was introduced into *S. thermophilus* CNRZ368. A strain harboring the  $\Delta arp1$  ORF instead of *arp1* (CNRZ368  $\Delta arp1$ ) was obtained by two successive homologous recombination events.

The effects of the *arp1* deletion were determined by PCR




FIG. 3. Effects of *arp1* deletion on the excision frequency of *ICES1*. Fragments containing the *attB* (361 bp) and *attI* (400 bp) recombination sites from the strains harboring wild-type *ICES1* (*ICES1* WT) or *ICES1*  $\Delta$ *arp1* were amplified by PCR using primers described in the legend to Fig. 1 and amounts of template DNA ranging from 1  $\mu$ g to 1 pg for CNRZ368  $\Delta$ *arp1* and from 1  $\mu$ g to 1 ng for CNRZ368. An internal fragment of the *fda* gene (339 bp) was amplified as a control.

amplification of fragments harboring *attI* and *attB*, using amounts of genomic DNA ranging from 1  $\mu$ g to 1 ng for wild-type *ICES1* and from 1  $\mu$ g to 1 pg for *ICES1*  $\Delta$ *arp1* (Fig. 3). The minimum amounts of DNA that produced a positive result for the *attB* site were 10 ng of CNRZ368 template DNA and 10 pg of CNRZ368  $\Delta$ *arp1* template DNA. Using the same procedure, the minimum amounts of DNA that produced a positive PCR result for the *attI* site were 0.1  $\mu$ g of CNRZ368 DNA and 0.1 ng of CNRZ368  $\Delta$ *arp1* DNA. The *arp1* deletion did not induce an increase in the PCR signal intensity when the *fda* gene was used as a control. Thus, deletion of the *arp1* gene resulted in at least a 1,000-fold increase in the concentration of *ICES1* *attB* and *attI* sites. Furthermore, whereas *ICES1* and *ICES2* were excised at different frequencies, the concentrations of a fragment carrying the *attI'* site from wild-type *ICES2* or *ICES2*  $\Delta$ *arp1* (Fig. 1A) increased by a factor similar to that observed for *ICES1* (data not shown). Therefore, the *arp1* gene repressed the excision of *ICES1* and *ICES2*.

To validate the assumption that Arp1 autoproteolysis is involved in the induction of ICE excision by MC, exponentially growing cells of CNRZ368  $\Delta$ *arp1* were treated with MC at a concentration of 0.1  $\mu$ g/ml (MIC/2). In three replicate experiments, the minimal amount of DNA that allowed detection of *attB* amplification (1 ng) was the same in the presence and in the absence of MC. Furthermore, in these three replicate experiments, the minimal amounts of DNA that allowed detection of *attI* amplification were 0.1 ng or 1 ng in the absence of MC and 0.1 ng after treatment with MC. Therefore, the inducibility of *ICES1*  $\Delta$ *arp1* excision by MC was reduced or suppressed compared to the results for the same MC treatment in the wild-type strain. This suggests that MC treatment alleviates the repression of *ICES1* excision mediated by Arp1. The minimal amount of DNA that allowed detection of *attI'* amplification (1 ng) was the same in the presence and in the absence of MC. This suggests that MC treatment also alleviates the repression of *ICES2* excision mediated by Arp1. In the same way, MC treatment derepressed the excision and transfer of another ICE, SXT from *V. cholerae*, probably by promoting the autocleavage of a cI homologue encoded by the element (3).

The functions of the other genes harbored by the *ICES1*/*ICES2* and *ICES3* regulation modules (Fig. 1B) remain unknown. Nevertheless, the regulation modules of numerous *Firmicutes* prophages, such as TP901-1 from *Lactococcus lactis* (16), and of another ICE, *ICEBs1* from *B. subtilis* (1), encode an OrfQ homologue and a "cI-like" repressor (i.e., a repressor

related to Arp2), while none of these elements encode a genuine homologue of the cI repressor. OrfQ and the OrfQ homologues have a DUF955 domain that has a conserved H-E-X-X-H motif, suggesting that these proteins could be Zn<sup>2+</sup> metalloproteinases (21). However, the activity of these proteins has not been demonstrated. Whereas the "cI-like" proteins encoded by prophages repress their lytic growth, these proteins are shorter than genuine cI homologues. Indeed, the cI-like repressors contain an HTH DNA binding domain but lack a region harboring autoproteolytic and oligomerization functions related to the genuine cI. However, the TP901-1 "cI-like" repressor binds to operators as dimers and higher multimers (14). Furthermore, the excision of prophages with cI-like genes and *orfQ* homologues, such as  $\phi$ Sfi21 from *S. thermophilus* (7) or TP901-1 (16), is inducible by MC, and the prophage TP901-1 cI-like gene is involved in the lytic phase induction pathway. Moreover, the induction of the prophage TP901-1 (16) and of *ICEBs1* by MC (1) is RecA dependent. Thus, *arp2* and *orfQ* might also be involved in the derepression pathway of the excision of the *S. thermophilus* ICEs.

To our knowledge, *ICES1*/*ICES2* and *ICES3* are the only integrative elements (i.e., phages, ICEs, or related elements) that encode a genuine cI homologue and might encode a cI-like homologue. Since MC induces the conjugative transfer of SXT, an ICE from *V. cholerae* coding for a genuine cI (3), and of *ICEBs1*, an ICE from *B. subtilis* coding for a cI-like repressor (1), DNA damage could regulate not only the excision but also the conjugative transfer of the ICEs from *S. thermophilus*.

#### REFERENCES

1. Auchtung, J. M., C. A. Lee, R. E. Monson, A. P. Lehman, and A. D. Grossman. 2005. Regulation of a *Bacillus subtilis* mobile genetic element by intercellular signaling and the global DNA damage response. *Proc. Natl. Acad. Sci. USA* **102**:12554–12559.
2. Bacic, M., A. C. Parker, J. Stagg, H. P. Whitley, W. G. Wells, L. A. Jacob, and C. J. Smith. 2005. Genetic and structural analysis of the *Bacteroides* conjugative transposon CTn341. *J. Bacteriol.* **187**:2858–2869.
3. Beaver, J. W., B. Hochhut, and M. K. Waldor. 2004. SOS response promotes horizontal dissemination of antibiotic resistance genes. *Nature* **427**:72–74.
4. Bentley, S. D., K. F. Chater, A. M. Cerdeno-Tarraga, G. L. Challis, N. R. Thomson, K. D. James, D. E. Harris, M. A. Quail, H. Kieser, D. Harper, A. Bateman, S. Brown, G. Chandra, C. W. Chen, M. Collins, A. Cronin, A. Fraser, A. Goble, J. Hidalgo, T. Hornsby, S. Howarth, C. H. Huang, T. Kieser, L. Larke, L. Murphy, K. Oliver, S. O'Neil, E. Rabinowitz, M. A. Rajandream, K. Rutherford, S. Rutter, K. Seeger, D. Saunders, S. Sharp, R. Squares, S. Squares, K. Taylor, T. Warren, A. Wietzorrek, J. Woodward, B. G. Barrell, J. Parkhill, and D. A. Hopwood. 2002. Complete genome sequence of the model actinomycete *Streptomyces coelicolor* A3(2). *Nature* **417**:141–147.
5. Biswas, I., A. Gruss, S. D. Ehrlich, and E. Maguin. 1993. High-efficiency gene inactivation and replacement system for gram-positive bacteria. *J. Bacteriol.* **175**:3628–3635.

6. **Bonheyo, G. T., B. D. Hund, N. B. Shoemaker, and A. A. Salyers.** 2001. Transfer region of a *Bacteroides* conjugative transposon contains regulatory as well as structural genes. *Plasmid* **46**:202–209.
7. **Brüssow, H., and A. Bruttin.** 1995. Characterization of a temperate *Streptococcus thermophilus* bacteriophage and its genetic relationship with lytic phages. *Virology* **212**:632–640.
8. **Burrus, V., G. Pavlovic, B. Decaris, and G. Guédon.** 2002. Conjugative transposons: the tip of the iceberg. *Mol. Microbiol.* **46**:601–610.
9. **Burrus, V., G. Pavlovic, B. Decaris, and G. Guédon.** 2002. The ICES*II* element of *Streptococcus thermophilus* belongs to a large family of integrative and conjugative elements that exchange modules and change their specificity of integration. *Plasmid* **48**:77.
10. **Burrus, V., Y. Roussel, B. Decaris, and G. Guédon.** 2000. Characterization of a novel integrative element, ICES*II*, in the lactic acid bacterium *Streptococcus thermophilus*. *Appl. Environ. Microbiol.* **66**:1749–1753.
11. **Clewell, D. B., D. D. Jaworski, S. E. Flannagan, L. A. Zitzow, and Y. A. Su.** 1995. The conjugative transposon Tn916 of *Enterococcus faecalis*: structural analysis and some key factors involved in movement. *Dev. Biol. Stand.* **85**:11–17.
12. **Dodd, I. B., K. E. Shearwin, A. J. Perkins, T. Burr, A. Hochschild, and J. B. Egan.** 2004. Cooperativity in long-range gene regulation by the lambda CI repressor. *Genes Dev.* **18**:344–354.
13. **Gaillard, M., T. Vallaey, F. J. Vorholter, M. Minoia, C. Werlen, V. Sentschilo, A. Puhler, and J. R. van der Meer.** 2006. The *clc* element of *Pseudomonas* sp. strain B13, a genomic island with various catabolic properties. *J. Bacteriol.* **188**:1999–2013.
14. **Johansen, A. H., L. Brondsted, and K. Hammer.** 2003. Identification of operator sites of the CI repressor of phage TP901-1: evolutionary link to other phages. *Virology* **311**:144–156.
15. **Larbig, K. D., A. Christmann, A. Johann, J. Klockgether, T. Hartsch, R. Merkl, L. Wiehlmann, H. J. Fritz, and B. Tummler.** 2002. Gene islands integrated into tRNA<sup>Gly</sup> genes confer genome diversity on a *Pseudomonas aeruginosa* clone. *J. Bacteriol.* **184**:6665–6680.
16. **Madsen, P. L., A. H. Johansen, K. Hammer, and L. Brondsted.** 1999. The genetic switch regulating activity of early promoters of the temperate lactococcal bacteriophage TP901-1. *J. Bacteriol.* **181**:7430–7438.
17. **Maguin, E., H. Prévost, S. D. Ehrlich, and A. Gruss.** 1996. Efficient insertional mutagenesis in lactococci and other gram-positive bacteria. *J. Bacteriol.* **178**:931–935.
18. **McGrath, B. M., J. A. O'Halloran, and J. T. Pembroke.** 2005. Pre-exposure to UV irradiation increases the transfer frequency of the IncJ conjugative transposon-like elements R391, R392, R705, R706, R997 and pMERPH and is recA<sup>+</sup> dependent. *FEMS Microbiol. Lett.* **243**:461–465.
19. **Mohd-Zain, Z., S. L. Turner, A. M. Cerdeno-Tarraga, A. K. Lilley, T. J. Inzana, A. J. Duncan, R. M. Harding, D. W. Hood, T. E. Peto, and D. W. Crook.** 2004. Transferable antibiotic resistance elements in *Haemophilus influenzae* share a common evolutionary origin with a diverse family of syntenic genomic islands. *J. Bacteriol.* **186**:8114–8122.
20. **Pavlovic, G., V. Burrus, B. Gintz, B. Decaris, and G. Guédon.** 2004. Evolution of genomic islands by deletion and tandem accretion by site-specific recombination: ICES*II*-related elements from *Streptococcus thermophilus*. *Microbiology* **150**:759–774.
21. **Rawlings, N. D., and A. J. Barrett.** 1995. Evolutionary families of metallo-peptidases. *Methods Enzymol.* **248**:183–228.
22. **Sentschilo, V., R. Ravatn, C. Werlen, A. J. Zehnder, and J. R. van der Meer.** 2003. Unusual integrase gene expression on the *clc* genomic island in *Pseudomonas* sp. strain B13. *J. Bacteriol.* **185**:4530–4538.
23. **Sezonov, G., C. Possoz, A. Friedmann, J. L. Pernodet, and M. Guéroux.** 2000. KorSA from the *Streptomyces* integrative element pSAM2 is a central transcriptional repressor: target genes and binding sites. *J. Bacteriol.* **182**:1243–1250.
24. **Stingle, F., and B. Mollet.** 1996. Disruption of the gene encoding penicillin-binding protein 2b (pbp2b) causes altered cell morphology and cease in exopolysaccharide production in *Streptococcus thermophilus* Sf16. *Mol. Microbiol.* **22**:357–366.
25. **Toussaint, A., C. Merlin, S. Monchy, M. A. Benotmane, R. Leplae, M. Mergeay, and D. Springael.** 2003. The biphenyl- and 4-chlorobiphenyl-catabolic transposon Tn4371, a member of a new family of genomic islands related to IncP and Ti plasmids. *Appl. Environ. Microbiol.* **69**:4837–4845.
26. **van der Meer, J. R., and V. Sentschilo.** 2003. Genomic islands and the evolution of catabolic pathways in bacteria. *Curr. Opin. Biotechnol.* **14**:1–7.