



HAL
open science

Relaxation of nonconvex unbounded integrals with general growth conditions in Cheeger–Sobolev spaces

Omar Anza Hafsa, Jean-Philippe Mandallena

► **To cite this version:**

Omar Anza Hafsa, Jean-Philippe Mandallena. Relaxation of nonconvex unbounded integrals with general growth conditions in Cheeger–Sobolev spaces. *Bulletin des Sciences Mathématiques*, 2018, 142, pp.49-93. 10.1016/j.bulsci.2017.09.002 . hal-01662240

HAL Id: hal-01662240

<https://hal.science/hal-01662240v1>

Submitted on 13 Dec 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Relaxation of nonconvex unbounded integrals with general growth conditions in Cheeger–Sobolev spaces

Omar Anza Hafsa ^{a,b}, Jean-Philippe Mandallena ^{a,*}

^a *Université de Nîmes, Laboratoire MIPA, Site des Carmes, Place Gabriel Péri, 30021 Nîmes, France*

^b *LMGC, UMR-CNRS 5508, Place Eugène Bataillon, 34095 Montpellier, France*

We study relaxation of nonconvex integrals of the calculus of variations in the setting of Cheeger–Sobolev spaces when the integrand does not have polynomial growth and can take infinite values.

Keywords:

Relaxation

Integral representation

Unbounded nonconvex integral

Ru-usc

General growth conditions

Cheeger–Sobolev space

1. Introduction

In this paper we are concerned with relaxation of integrals of type

$$\int_X L(x, \nabla_\mu u(x)) d\mu(x), \tag{1.1}$$

* Corresponding author.

E-mail addresses: omar.anza-hafsa@unimes.fr (O. Anza Hafsa), jean-philippe.mandallena@unimes.fr (J.-P. Mandallena).

where (X, d, μ) is a metric measure space, with (X, d) a length space, i.e., the distance between any two points equals infimum of lengths of curves connecting the points, separable and compact, satisfying a weak $(1, p)$ -Poincaré inequality with $p > 1$ and such that μ is a doubling positive Radon measure on X , $u \in W_\mu^{1,p}(X; \mathbb{R}^m)$ with $m \geq 1$ an integer and $\nabla_\mu u$ is the μ -gradient of u , and $L : X \times \mathbb{M} \rightarrow [0, \infty]$ is a Borel measurable integrand not necessarily convex with respect to $\xi \in \mathbb{M}$, where \mathbb{M} denotes the space of all $m \times N$ matrices with $N \geq 1$ an integer. Such a relaxation problem in such a metric measure setting was studied for the first time in [10] (see also [14,39,2,27,3,40,43,35] and the references therein) when L has p -growth, i.e., there exist $\alpha, \beta > 0$ such that for every $x \in X$ and every $\xi \in \mathbb{M}$,

$$\alpha|\xi|^p \leq L(x, \xi) \leq \beta(1 + |\xi|^p), \quad (1.2)$$

where it is proved (see [10, Theorem 2.21 and Corollary 2.27]) that if (1.2) holds then the relaxation of (1.1) with respect to the norm of $L_\mu^p(X; \mathbb{R}^m)$ is given by

$$\int_X \mathcal{Q}_\mu L(x, \nabla_\mu u(x)) d\mu(x),$$

where $u \in W_\mu^{1,p}(X; \mathbb{R}^m)$ and $\mathcal{Q}_\mu L : X \times \mathbb{M} \rightarrow [0, \infty]$, called the μ -quasiconvexification of L , is defined by

$$\mathcal{Q}_\mu L(x, \xi) := \overline{\lim}_{\rho \rightarrow 0} \inf \left\{ \int_{Q_\rho(x)} L(y, \xi + \nabla_\mu w(y)) d\mu(y) : w \in W_{\mu,0}^{1,p}(Q_\rho(x); \mathbb{R}^m) \right\} \quad (1.3)$$

where $Q_\rho(x)$ is the open ball centered at $x \in X$ with radius $\rho > 0$, and, for each open set A of X , $W_{\mu,0}^{1,p}(A; \mathbb{R}^m)$ is the closure of $\text{Lip}_0(A; \mathbb{R}^m)$ with respect to $W_\mu^{1,p}$ -norm with $\text{Lip}_0(A; \mathbb{R}^m) := \{u \in \text{Lip}(X; \mathbb{R}^m) : u = 0 \text{ on } X \setminus A\}$, where $\text{Lip}(X; \mathbb{R}^m)$ is the class of Lipschitz functions from X to \mathbb{R}^m (see §3.1 for more details).

Our motivation for developing relaxation, and more generally calculus of variations, in the setting of metric measure spaces comes from applications to hyperelasticity. In fact, the interest of considering a general measure is that its support can be interpreted as a hyperelastic structure together with its singularities like for example thin dimensions, corners, junctions, etc. Such mechanical singular objects naturally lead to develop calculus of variations in the setting of metric measure spaces. Indeed, for example, a low multi-dimensional structures can be described by a finite number of smooth compact manifolds S_i of dimension k_i on which a superficial measure $\mu_i = \mathcal{H}^{k_i}|_{S_i}$ is attached. Such a situation leads to deal with the finite union of manifolds S_i , i.e., $X = \cup_i S_i$, together with the finite sum of measures μ_i , i.e., $\mu = \sum_i \mu_i$, whose mathematical framework is that of metric measure spaces (for more examples, we refer the reader to [14, 50,20] and [22, Chapter 2, §10] and the references therein). In this way, having in mind

the two basic conditions of hyperelasticity, i.e., “the non-interpenetration of the matter” and “the necessity of an infinite amount of energy to compress a finite piece of matter into a point”, it is then of interest to study relaxation of nonconvex integrals of type (1.1) when the integrand does not have p -growth and can take infinite values: this is the general purpose of the present paper. Note that although our framework needs some “convexity” assumptions (see especially (2.7) which implies that domain of $L(x, \cdot)$ is convex) it is consistent with the two above conditions of hyperelasticity (see [6, §2.2] and [11, §9]). Nevertheless, this dose of convexity makes our framework not consistent with another condition of hyperelasticity that is “frame-indifference” (see [11, Remark 9.1]). (For more details on the theory of hyperelasticity we refer the reader to [42].)

For related works in the Euclidean case, i.e., when $(X, d, \mu) = (\overline{\Omega}, |\cdot - \cdot|, \mathcal{L}_N)$ where Ω is a bounded open subset of \mathbb{R}^N and \mathcal{L}_N is the Lebesgue measure, we refer the reader to [47, 48, 4, 5, 49, 1, 8, 41, 18, 44, 45] and the references therein.

Generally speaking, in this paper our main contribution (see §2.1, §2.2 and §2.3 for more details and more precisely Theorem 2.7) is to prove that for $p > \kappa$, with $\kappa := \frac{\ln(C_d)}{\ln(2)}$ where $C_d \geq 1$ is the doubling constant, see (2.1), if L is *radially uniformly upper semicontinuous*, i.e., there exists $a \in L^1_\mu(X;]0, \infty])$ such that

$$\overline{\lim}_{t \rightarrow 1^-} \sup_{x \in X} \sup_{\xi \in \mathbb{L}_x} \frac{L(x, t\xi) - L(x, \xi)}{a(x) + L(x, \xi)} \leq 0,$$

where \mathbb{L}_x denotes the effective domain of $L(x, \cdot)$, and if L has G -growth, i.e., there exist $\alpha, \beta > 0$ such that for every $x \in X$ and every $\xi \in \mathbb{M}$,

$$\alpha G(x, \xi) \leq L(x, \xi) \leq \beta(1 + G(x, \xi)), \quad (1.4)$$

where $G : X \times \mathbb{M} \rightarrow [0, \infty]$ is a Borel measurable and p -coercive integrand satisfying some “convexity” assumptions, see (2.6), (2.7), (2.10) and (2.11), then the relaxation of (1.1) with respect to the norm of $L^p_\mu(X; \mathbb{R}^m)$ is given by

$$\int_X \widehat{\mathcal{Q}_\mu L}(x, \nabla_\mu u(x)) d\mu(x),$$

where $\widehat{\mathcal{Q}_\mu L} : X \times \mathbb{M} \rightarrow [0, \infty]$ is defined by

$$\widehat{\mathcal{Q}_\mu L}(x, \xi) := \liminf_{t \rightarrow 1^-} \mathcal{Q}_\mu L(x, t\xi).$$

Moreover, we also prove (see Corollary 2.8) that if, in addition, $\mathcal{Q}_\mu G = G$ and, for each $x \in X$, $\mathcal{Q}_\mu L(x, \cdot)$ is lower semicontinuous on the interior $\text{int}(\mathbb{L}_x)$ of \mathbb{L}_x , then

$$\widehat{\mathcal{Q}_\mu L}(x, \xi) = \begin{cases} \mathcal{Q}_\mu L(x, \xi) & \text{if } x \in X \text{ and } \xi \in \text{int}(\mathbb{L}_x) \\ \liminf_{t \rightarrow 1^-} \mathcal{Q}_\mu L(x, t\xi) & \text{if } x \in X \text{ and } \xi \in \partial\mathbb{L}_x \\ \infty & \text{otherwise.} \end{cases}$$

In the p -growth (and finite) case, i.e., when (1.2) holds, which was already studied in [10], it is not necessary to assume that L is radially uniformly upper semicontinuous, so that the relaxed integrand is simply given by (1.3). The main novelty of the paper is to deal, in the setting of metric measure spaces, with the relaxation of functionals with non-finite integrands satisfying non-homogeneous growth conditions of type (1.4), and to establish an integral representation of corresponding relaxed functionals in this context.

The plan of the paper is as follows. The main result (see Theorem 2.7 and also Corollary 2.8) is given in Section 2. The proof of Theorem 2.7 is established in Section 4, whereas Corollary 2.8, which is a consequence of Theorem 2.7, is proved at the end of Section 2. Section 3 is devoted to several auxiliary results needed for proving Theorem 2.7.

Notation. The open and closed balls centered at $x \in X$ with radius $\rho > 0$ are denoted by:

$$Q_\rho(x) := \left\{ y \in X : d(x, y) < \rho \right\};$$

$$\overline{Q}_\rho(x) := \left\{ y \in X : d(x, y) \leq \rho \right\}.$$

For $x \in X$ and $\rho > 0$ we set

$$\partial Q_\rho(x) := \overline{Q}_\rho(x) \setminus Q_\rho(x) = \left\{ y \in X : d(x, y) = \rho \right\}.$$

For $A \subset X$, the diameter of A (resp. the distance from a point $x \in X$ to the subset A) is defined by $\text{diam}(A) := \sup_{x, y \in A} d(x, y)$ (resp. $\text{dist}(x, A) := \inf_{y \in A} d(x, y)$).

The symbol \int stands for the mean-value integral

$$\int_B f d\mu = \frac{1}{\mu(B)} \int_B f d\mu.$$

2. Main results

2.1. Setting of the problem

Let (X, d, μ) be a metric measure space, where (X, d) is a length space which is separable and compact, and μ is a positive Radon measure on X . In what follows, we assume that μ is doubling, i.e., there exists a constant $C_d \geq 1$ (called doubling constant) such that

$$\mu(Q_\rho(x)) \leq C_d \mu\left(Q_{\frac{\rho}{2}}(x)\right) \tag{2.1}$$

for all $x \in X$ and all $\rho > 0$, and X enjoys a weak $(1, p)$ -Poincaré inequality with $1 < p < \infty$, i.e., there exist $C_P > 0$ and $\sigma \geq 1$ such that for every $x \in X$ and every $\rho > 0$,

$$\int_{Q_\rho(x)} \left| f - \int_{Q_\rho(x)} f d\mu \right| d\mu \leq \rho C_P \left(\int_{Q_{\sigma\rho}(x)} g^p d\mu \right)^{\frac{1}{p}} \quad (2.2)$$

for every $f \in L^p_\mu(X)$ and every p -weak upper gradient $g \in L^p_\mu(X)$ for f . (For the definition of the concept of p -weak upper gradient, see [Definition 3.2](#).) As μ is doubling, for each $Q_r(\bar{x})$ with $r > 0$ and $\bar{x} \in X$ we have

$$\frac{\mu(Q_\rho(x))}{\mu(Q_r(\bar{x}))} \geq 4^{-\kappa} \left(\frac{\rho}{r} \right)^\kappa \quad (2.3)$$

for all $x \in Q_r(\bar{x})$ and all $0 < \rho \leq r$, where $\kappa := \frac{\ln(C_d)}{\ln(2)}$ (see [\[30, Lemma 4.7\]](#)).

From now on, we suppose $p > \kappa$ and we fix an integer $m \geq 1$.

Let $\mathcal{O}(X)$ be the class of open subsets of X and let $E : W^{1,p}_\mu(X; \mathbb{R}^m) \times \mathcal{O}(X) \rightarrow [0, \infty]$ be the variational integral defined by

$$E(u, A) := \int_A L(x, \nabla_\mu u(x)) d\mu(x), \quad (2.4)$$

where $L : X \times \mathbb{M} \rightarrow [0, \infty]$ is a Borel measurable integrand not necessarily convex with respect to $\xi \in \mathbb{M}$, where \mathbb{M} denotes the space of all $m \times N$ matrices with $N \geq 1$ an integer.

The space $W^{1,p}_\mu(X; \mathbb{R}^m)$ denotes the class of p -Cheeger–Sobolev functions from X to \mathbb{R}^m and $\nabla_\mu u$ is the μ -gradient of u (see §3.1 for more details).

Let $\bar{E} : W^{1,p}_\mu(X; \mathbb{R}^m) \times \mathcal{O}(X) \rightarrow [0, \infty]$ be the “relaxed” variational functional of the variational integral E with respect to the strong convergence in $L^p_\mu(X; \mathbb{R}^m)$, i.e.,

$$\bar{E}(u, A) := \inf \left\{ \liminf_{n \rightarrow \infty} E(u_n, A) : u_n \xrightarrow{L^p_k} u \right\}.$$

The object of the paper is to study the problem of finding an integral representation for $\bar{E}(\cdot, X)$ in the case where L does not have p -growth and can take infinite values.

2.2. General growth and ru -usc condition

Let $G : X \times \mathbb{M} \rightarrow [0, \infty]$ be a Borel measurable integrand which is p -coercive, i.e., there exists $c > 0$ such that for every $x \in X$ and every $\xi \in \mathbb{M}$,

$$G(x, \xi) \geq c|\xi|^p, \quad (2.5)$$

and for which there exists $r > 0$ such that

$$\sup_{|\xi| \leq r} G(\cdot, \xi) \in L^1_\mu(X). \quad (2.6)$$

Remark 2.1. If $\sup_{|\xi| \leq r} G(\cdot, \xi) \in L^\infty_\mu(X)$ then (2.6) is satisfied. In particular, this latter condition holds when G depends only on ξ and is convex.

We also assume that there exists $\gamma > 0$ such that for every $x \in X$, every $t \in]0, 1[$ and every $\xi, \zeta \in \mathbb{M}$,

$$G(x, t\xi + (1-t)\zeta) \leq \gamma(1 + G(x, \xi) + G(x, \zeta)). \quad (2.7)$$

Remark 2.2. If (2.7) holds and if $0 \in \text{int}(\{\xi \in \mathbb{M} : G(\cdot, \xi) \in L^1_\mu(X)\})$ then (2.6) is verified, see [8, Lemma 4.1].

Remark 2.3. If (2.7) holds then the effective domain \mathbb{G}_x of $G(x, \cdot)$ is convex.

Let $\mathcal{G}, \overline{\mathcal{G}} : W^{1,p}_\mu(X; \mathbb{R}^m) \rightarrow [0, \infty]$ be the integral functionals defined by:

$$\mathcal{G}(u) := \int_X G(x, \nabla_\mu u(x)) d\mu(x); \quad (2.8)$$

$$\overline{\mathcal{G}}(u) := \inf \left\{ \varliminf_{n \rightarrow \infty} \mathcal{G}(u_n) : u_n \xrightarrow{L^p_\mu} u \right\}. \quad (2.9)$$

Let us denote the effective domains of the functionals \mathcal{G} and $\overline{\mathcal{G}}$ by \mathfrak{G} and $\overline{\mathfrak{G}}$ respectively. We furthermore assume that:

$$\mathfrak{G} = \overline{\mathfrak{G}}; \quad (2.10)$$

$$\text{if } u \in \mathfrak{G} \text{ then } \lim_{r \rightarrow 0} \int_{Q_r(x)} |G(y, \nabla_\mu u(x)) - G(x, \nabla_\mu u(x))| d\mu(y) = 0 \text{ for } \mu\text{-a.a. } x \in X. \quad (2.11)$$

Remark 2.4. The assumptions (2.11) is satisfied in the following cases:

- (1) if $G(\cdot, \xi) \in L^1_\mu(X)$ for all $\xi \in \mathbb{M}$ then (2.11) holds;
- (2) if G only depends on ξ then (2.11) holds;
- (3) if $G(x, \xi) = G_1(x) + G_2(\xi)$ for all $x \in X$ and all $\xi \in \mathbb{M}$ and if $G_1 \in L^1_\mu(X)$ then (2.11) holds;
- (4) if $G(x, \xi) = G_1(x)G_2(\xi)$ for all $x \in X$ and all $\xi \in \mathbb{M}$ and if $G_1 \in L^1_\mu(X)$ then (2.11) holds.

Throughout the paper, we assume that L has G -growth, i.e., there exist $\alpha, \beta > 0$ such that for every $x \in X$ and every $\xi \in \mathbb{M}$,

$$\alpha G(x, \xi) \leq L(x, \xi) \leq \beta(1 + G(x, \xi)). \quad (2.12)$$

Remark 2.5. If (2.7) and (2.12) hold then the effective domain \mathbb{L}_x of $L(x, \cdot)$ is equal to \mathbb{G}_x , and so is convex.

Remark 2.6. If (2.12) is satisfied then the effective domain of the functional $\overline{E}(\cdot, X)$ is equal to $\overline{\mathfrak{S}}$, and so to \mathfrak{S} when (2.10) holds.

When $G(x, \cdot) \equiv |\cdot|^p$ we say that L has p -growth. The p -growth case was already studied in [10]. The object of this paper is to deal with the G -growth case. For this, in addition, we need to suppose that L is radially uniformly upper semicontinuous (ru-usc), i.e., there exists $a \in L_\mu^1(X;]0, \infty])$ such that

$$\overline{\lim}_{t \rightarrow 1^-} \Delta_L^a(t) \leq 0 \quad (2.13)$$

with $\Delta_L^a : [0, 1] \rightarrow]-\infty, \infty]$ given by

$$\Delta_L^a(t) := \sup_{x \in X} \sup_{\xi \in \mathbb{L}_x} \frac{L(x, t\xi) - L(x, \xi)}{a(x) + L(x, \xi)}.$$

(For more details on the concept of ru-usc, see §3.3.)

2.3. Integral representation theorem

In what follows $p > \kappa$, where $\kappa := \frac{\ln(C_d)}{\ln(2)}$ with $C_d \geq 1$ given by the inequality (2.1), and $m \geq 1$. Let $\mathcal{Q}_\mu L : X \times \mathbb{M} \rightarrow [0, \infty]$, called the μ -quasiconvexification of L , be given by

$$\mathcal{Q}_\mu L(x, \xi) := \overline{\lim}_{\rho \rightarrow 0} \inf_{Q_\rho(x)} \left\{ \int L(y, \xi + \nabla_\mu w(y)) d\mu(y) : w \in W_{\mu, 0}^{1,p}(Q_\rho(x); \mathbb{R}^m) \right\}, \quad (2.14)$$

where, for each $A \in \mathcal{O}(X)$, $W_{\mu, 0}^{1,p}(A; \mathbb{R}^m)$ is the closure of $\text{Lip}_0(A; \mathbb{R}^m)$ with respect to $W_\mu^{1,p}$ -norm with $\text{Lip}_0(A; \mathbb{R}^m) := \{u \in \text{Lip}(X; \mathbb{R}^m) : u = 0 \text{ on } X \setminus A\}$, where $\text{Lip}(X; \mathbb{R}^m)$ is the class of Lipschitz functions from X to \mathbb{R}^m (see §3.1 for more details). The main result of the paper is the following.

Theorem 2.7. *If (2.5), (2.6), (2.7), (2.10), (2.11), (2.12) and (2.13) hold then*

$$\overline{E}(u, X) = \begin{cases} \int_X \widehat{\mathcal{Q}}_\mu L(x, \nabla_\mu u(x)) d\mu(x) & \text{if } u \in \mathfrak{G} \\ \infty & \text{if } u \in W_\mu^{1,p}(X; \mathbb{R}^m) \setminus \mathfrak{G}, \end{cases} \quad (2.15)$$

where $\widehat{\mathcal{Q}}_\mu L : X \times \mathbb{M} \rightarrow [0, \infty]$ is given by

$$\widehat{\mathcal{Q}}_\mu L(x, \xi) = \liminf_{t \rightarrow 1^-} \mathcal{Q}_\mu L(x, t\xi).$$

As a consequence of [Theorem 2.7](#), we have

Corollary 2.8. *Under the hypotheses of [Theorem 2.7](#), if $\mathcal{Q}_\mu G = G$ and if, for each $x \in X$, $\mathcal{Q}_\mu L(x, \cdot)$ is lsc on the interior $\text{int}(\mathbb{L}_x)$ of the effective domain \mathbb{L}_x of $L(x, \cdot)$, then (2.15) holds and*

$$\widehat{\mathcal{Q}}_\mu L(x, \xi) = \begin{cases} \mathcal{Q}_\mu L(x, \xi) & \text{if } x \in X \text{ and } \xi \in \text{int}(\mathbb{L}_x) \\ \lim_{t \rightarrow 1^-} \mathcal{Q}_\mu L(x, t\xi) & \text{if } x \in X \text{ and } \xi \in \partial\mathbb{L}_x \\ \infty & \text{otherwise.} \end{cases} \quad (2.16)$$

Proof. As $\mathcal{Q}_\mu G = G$ we have $\mathcal{Q}_\mu \mathbb{L}_x = \mathbb{L}_x$ for all $x \in X$, where $\mathcal{Q}\mathbb{L}_x$ denotes the effective domain of $\mathcal{Q}_\mu L(x, \cdot)$. Moreover, from (2.7) is easily seen that \mathbb{L}_x is convex for all $x \in X$, hence (3.38) holds, and (2.16) follows from [Theorem 3.23](#). \square

3. Auxiliary results

3.1. The p -Cheeger–Sobolev spaces

Let $p > 1$ be a real number, let (X, d, μ) be a metric measure space, where (X, d) is a length space which is separable and compact, and μ is a positive Radon measure on X . We begin with the concept of upper gradient introduced by Heinonen and Koskela (see [\[33\]](#)).

Definition 3.1. A Borel function $g : X \rightarrow [0, \infty]$ is said to be an upper gradient for $f : X \rightarrow \mathbb{R}$ if $|f(c(1)) - f(c(0))| \leq \int_0^1 g(c(s)) ds$ for all continuous rectifiable curves $c : [0, 1] \rightarrow X$.

The concept of upper gradient has been generalized by Cheeger as follows (see [\[19, Definition 2.8\]](#)).

Definition 3.2. A function $g \in L_\mu^p(X)$ is said to be a p -weak upper gradient for $f \in L_\mu^p(X)$ if there exist $\{f_n\}_n \subset L_\mu^p(X)$ and $\{g_n\}_n \subset L_\mu^p(X)$ such that for each $n \geq 1$, g_n is an upper gradient for f_n , $f_n \rightarrow f$ in $L_\mu^p(X)$ and $g_n \rightarrow g$ in $L_\mu^p(X)$.

Denote the algebra of Lipschitz functions from X to \mathbb{R} by $\text{Lip}(X)$. (As X is compact, every Lipschitz function from X to \mathbb{R} is bounded.) From Cheeger and Keith (see [19, Theorem 4.38] and [38, Definition 2.1.1 and Theorem 2.3.1]) we have

Theorem 3.3. *If μ is doubling, i.e., (2.1) holds, and X enjoys a weak $(1, p)$ -Poincaré inequality, i.e., (2.2) holds, then there exists a countable family $\{(X_k, \xi^k)\}_k$ of μ -measurable disjoint subsets X_k of X with $\mu(X \setminus \cup_k X_k) = 0$ and of functions $\xi^k = (\xi_1^k, \dots, \xi_{N(k)}^k) : X \rightarrow \mathbb{R}^{N(k)}$ with $\xi_i^k \in \text{Lip}(X)$ satisfying the following properties:*

- (a) *there exists an integer $N \geq 1$ such that $N(k) \in \{1, \dots, N\}$ for all k ;*
- (b) *for every k and every $f \in \text{Lip}(X)$ there is a unique $D_\mu^k f \in L^\infty(X_k; \mathbb{R}^{N(k)})$ such that for μ -a.e. $x \in X_k$,*

$$\lim_{\rho \rightarrow 0} \frac{1}{\rho} \|f - f_x\|_{L^\infty(Q_\rho(x))} = 0,$$

where $f_x \in \text{Lip}(X)$ is given by $f_x(y) := f(x) + D_\mu^k f(x) \cdot (\xi^k(y) - \xi^k(x))$; in particular

$$D_\mu^k f_x(y) = D_\mu^k f(x) \text{ for } \mu\text{-a.a. } y \in X_k;$$

- (c) *the operator $D_\mu : \text{Lip}(X) \rightarrow L^\infty(X; \mathbb{R}^N)$ given by*

$$D_\mu f := \sum_k \mathbb{1}_{X_k} D_\mu^k f,$$

where $\mathbb{1}_{X_k}$ denotes the characteristic function of X_k , is linear and, for each $f, g \in \text{Lip}(X)$, one has

$$D_\mu(fg) = f D_\mu g + g D_\mu f;$$

- (d) *for every $f \in \text{Lip}(X)$, $D_\mu f = 0$ μ -a.e. on every μ -measurable set where f is constant.*

Remark 3.4. Theorem 3.3 is true without the assumption that (X, d) is a length space.

Let $\text{Lip}(X; \mathbb{R}^m) := [\text{Lip}(X)]^m$ and let $\nabla_\mu : \text{Lip}(X; \mathbb{R}^m) \rightarrow L^\infty(X; \mathbb{M})$ given by

$$\nabla_\mu u := \begin{pmatrix} D_\mu u_1 \\ \vdots \\ D_\mu u_m \end{pmatrix} \text{ with } u = (u_1, \dots, u_m).$$

From Theorem 3.3(c) we see that for every $u \in \text{Lip}(X; \mathbb{R}^m)$ and every $f \in \text{Lip}(X)$, one has

$$\nabla_\mu(fu) = f \nabla_\mu u + D_\mu f \otimes u. \tag{3.1}$$

Definition 3.5. The p -Cheeger–Sobolev space $W_\mu^{1,p}(X; \mathbb{R}^m)$ is defined as the completion of $\text{Lip}(X; \mathbb{R}^m)$ with respect to the norm

$$\|u\|_{W_\mu^{1,p}(X; \mathbb{R}^m)} := \|u\|_{L_\mu^p(X; \mathbb{R}^m)} + \|\nabla_\mu u\|_{L_\mu^p(X; \mathbb{M})}. \quad (3.2)$$

Taking [Proposition 3.7](#)(a) below into account, since $\|\nabla_\mu u\|_{L_\mu^p(X; \mathbb{M})} \leq \|u\|_{W_\mu^{1,p}(X; \mathbb{R}^m)}$ for all $u \in \text{Lip}(X; \mathbb{R}^m)$ the linear map ∇_μ from $\text{Lip}(X; \mathbb{R}^m)$ to $L_\mu^p(X; \mathbb{M})$ has a unique extension to $W_\mu^{1,p}(X; \mathbb{R}^m)$ which will still be denoted by ∇_μ and will be called the μ -gradient.

Remark 3.6. When X is the closure of a bounded open subset Ω of \mathbb{R}^N and μ is the Lebesgue measure on $\overline{\Omega}$, we retrieve the (classical) Sobolev spaces $W^{1,p}(\Omega; \mathbb{R}^m)$. If X is a compact manifold M and if μ is the superficial measure on M , we obtain the (classical) Sobolev spaces $W^{1,p}(M; \mathbb{R}^m)$ on the compact manifold M . For more details on the various possible extensions of the classical theory of the Sobolev spaces to the setting of metric measure spaces, we refer to [\[31, §10-14\]](#) (see also [\[19,46,29,30\]](#)).

The following proposition (whose proof is given below, see also [\[10, Proposition 2.28\]](#)) provides useful properties for dealing with calculus of variations in the metric measure setting.

Proposition 3.7. *Under the hypotheses of [Theorem 3.3](#), we have:*

- (a) *the μ -gradient is closable in $W_\mu^{1,p}(X; \mathbb{R}^m)$, i.e., for every $u \in W_\mu^{1,p}(X; \mathbb{R}^m)$ and every $A \in \mathcal{O}(X)$, if $u(x) = 0$ for μ -a.a. $x \in A$ then $\nabla_\mu u(x) = 0$ for μ -a.a. $x \in A$;*
- (b) *X enjoys a p -Sobolev inequality, i.e., there exist $C_S > 0$ and $\chi \geq 1$ such that*

$$\left(\int_{Q_\rho(x)} |v|^{\chi p} d\mu \right)^{\frac{1}{\chi p}} \leq \rho C_S \left(\int_{Q_\rho(x)} |\nabla_\mu v|^p d\mu \right)^{\frac{1}{p}} \quad (3.3)$$

for all $0 < \rho \leq \rho_0$, with $\rho_0 > 0$, and all $v \in W_{\mu,0}^{1,p}(Q_\rho(x); \mathbb{R}^m)$, where, for each $A \in \mathcal{O}(X)$, $W_{\mu,0}^{1,p}(A; \mathbb{R}^m)$ is the closure of $\text{Lip}_0(A; \mathbb{R}^m)$ with respect to $W_\mu^{1,p}$ -norm defined in [\(3.2\)](#) with

$$\text{Lip}_0(A; \mathbb{R}^m) := \{u \in \text{Lip}(X; \mathbb{R}^m) : u = 0 \text{ on } X \setminus A\};$$

- (c) *X satisfies the Vitali covering theorem, i.e., for every $A \subset X$ and every family \mathcal{F} of closed balls in X , if $\inf\{\rho > 0 : \overline{Q}_\rho(x) \in \mathcal{F}\} = 0$ for all $x \in A$ then there exists a countable disjoint subfamily \mathcal{G} of \mathcal{F} such that $\mu(A \setminus \cup_{Q \in \mathcal{G}} Q) = 0$; in other words, $A \subset (\cup_{Q \in \mathcal{G}} Q) \cup N$ with $\mu(N) = 0$;*
- (d) *for every $u \in W_\mu^{1,p}(X; \mathbb{R}^m)$ and μ -a.e. $x \in X$ there exists $u_x \in W_\mu^{1,p}(X; \mathbb{R}^m)$ such that:*

$$\nabla_\mu u_x(y) = \nabla_\mu u(x) \text{ for } \mu\text{-a.a. } y \in X; \quad (3.4)$$

$$\lim_{\rho \rightarrow 0} \frac{1}{\rho} \|u - u_x\|_{L^\infty(Q_\rho(x); \mathbb{R}^m)} = 0 \text{ if } p > \kappa, \quad (3.5)$$

where $\kappa := \frac{\ln(C_d)}{\ln(2)}$ with $C_d \geq 1$ given by the inequality (2.1);

(e) for every $x \in X$, every $\rho > 0$ and every $\tau \in]0, 1[$ there exists a Urysohn function $\varphi \in \text{Lip}(X)$ for the pair $(X \setminus Q_\rho(x), \overline{Q}_{\tau\rho}(x))$ ¹ such that

$$\|D_\mu \varphi\|_{L^\infty_\mu(X; \mathbb{R}^N)} \leq \frac{\theta}{\rho(1-\tau)}$$

for some $\theta > 0$;

(f) for μ -a.e. $x \in X$,

$$\lim_{\tau \rightarrow 1^-} \lim_{\rho \rightarrow 0} \frac{\mu(Q_{\tau\rho}(x))}{\mu(Q_\rho(x))} = \lim_{\tau \rightarrow 1^-} \overline{\lim}_{\rho \rightarrow 0} \frac{\mu(Q_{\tau\rho}(x))}{\mu(Q_\rho(x))} = 1. \quad (3.6)$$

Remark 3.8. As μ is a Radon measure, if X satisfies the Vitali covering theorem, i.e., Proposition 3.7(c) holds, then for every $A \in \mathcal{O}(X)$ and every $\varepsilon > 0$ there exists a countable family $\{Q_{\rho_i}(x_i)\}_{i \in I}$ of disjoint open balls of A with $x_i \in A$, $\rho_i \in]0, \varepsilon[$ and $\mu(\partial Q_{\rho_i}(x_i)) = 0$ such that $\mu(A \setminus \cup_{i \in I} Q_{\rho_i}(x_i)) = 0$.

Proof of Proposition 3.7. Firstly, X satisfies the Vitali covering theorem, i.e., the property (c) holds, because μ is doubling (see [25, Theorem 2.8.18]). Secondly, the closability of the μ -gradient in $\text{Lip}(X; \mathbb{R}^m)$, given by Theorem 3.3(d), can be extended from $\text{Lip}(X; \mathbb{R}^m)$ to $W_\mu^{1,p}(X; \mathbb{R}^m)$ by using the closability theorem of Franchi, Hajlasz and Koskela (see [26, Theorem 10]). Thus, the property (a) is satisfied. Thirdly, according to Cheeger (see [19, §4, p. 450] and also [32,34]), since μ is doubling and X enjoys a weak $(1, p)$ -Poincaré inequality, we can assert that there exist $c > 0$ and $\chi > 1$ such that for every $0 < \rho \leq \rho_0$, with $\rho_0 \geq 0$, every $v \in W_{\mu,0}^{1,p}(X; \mathbb{R}^m)$ and every p -weak upper gradient $g \in L_\mu^p(X; \mathbb{R}^m)$ for v ,

$$\left(\int_{Q_\rho(x)} |v|^{\chi p} d\mu \right)^{\frac{1}{\chi p}} \leq \rho c \left(\int_{Q_\rho(x)} |g|^p d\mu \right)^{\frac{1}{p}}. \quad (3.7)$$

On the other hand, from Cheeger (see [19, Theorems 2.10 and 2.18]), for each $w \in W_\mu^{1,p}(X)$ there exists a unique p -weak upper gradient for w , denoted by $g_w \in L_\mu^p(X)$ and called the minimal p -weak upper gradient for w , such that for every p -weak upper

¹ Given a metric space (X, d) , by a Urysohn function from X to \mathbb{R} for the pair $(X \setminus V, K)$, where $K \subset V \subset X$ with K compact and V open, we mean a continuous function $\varphi : X \rightarrow \mathbb{R}$ such that $\varphi(x) \in [0, 1]$ for all $x \in X$, $\varphi(x) = 0$ for all $x \in X \setminus V$ and $\varphi(x) = 1$ for all $x \in K$.

gradient $g \in L^p_\mu(X)$ for w , $g_w(x) \leq g(x)$ for μ -a.a. $x \in X$. Moreover (see [19, §4] and also [13, §B.2, p. 363], [16] and [28, Remark 2.15]), there exists $\theta \geq 1$ such that for every $w \in W^{1,p}_\mu(X)$ and μ -a.e. $x \in X$,

$$\frac{1}{\theta}g_w(x) \leq |D_\mu w(x)| \leq \theta g_w(x). \quad (3.8)$$

As for $v = (v_i)_{i=1,\dots,m} \in W^{1,p}_\mu(X; \mathbb{R}^m)$ we have $\nabla_\mu v = (D_\mu v_i)_{i=1,\dots,m}$, it follows that

$$\frac{1}{\theta}|g_v(x)| \leq |\nabla_\mu v(x)| \leq \theta|g_v(x)| \quad (3.9)$$

for μ -a.a. $x \in X$, where $g_v := (g_{v_i})_{i=1,\dots,m}$ is naturally called the minimal p -weak upper gradient for v . Combining (3.7) with (3.9) we obtain the property (b). Fourthly, from Björn (see [16, Corollary 4.6(ii)]) we see that for every k , every $u \in W^{1,p}_\mu(X; \mathbb{R}^m)$ and μ -a.e. $x \in X_k$,

$$\nabla_\mu u_x(y) = \nabla_\mu u(x) \text{ for } \mu\text{-a.a. } y \in X_k,$$

where $u_x \in W^{1,p}_\mu(X; \mathbb{R}^m)$ is given by

$$u_x(y) := u(y) - u(x) - \nabla_\mu u(x) \cdot (\xi^k(y) - \xi^k(x)),$$

and if $p > \kappa$ then u is L^∞_μ -differentiable at x , i.e.,

$$\lim_{\rho \rightarrow 0} \frac{1}{\rho} \|u(y) - u_x(y)\|_{L^\infty(Q_\rho(x); \mathbb{R}^m)} = 0.$$

Hence the property (d) is verified. Fifthly, given $\rho > 0$, $\tau \in]0, 1[$ and $x \in X$, there exists a Urysohn function $\varphi \in \text{Lip}(X)$ for the pair $(X \setminus Q_\rho(x), \overline{Q}_\tau(x))$ such

$$\|\text{Lip}\varphi\|_{L^\infty_\mu(X)} \leq \frac{1}{\rho(1-\tau)},$$

where for every $y \in X$,

$$\text{Lip}\varphi(y) := \overline{\lim}_{d(y,z) \rightarrow 0} \frac{|\varphi(y) - \varphi(z)|}{d(y,z)}.$$

But, since μ is doubling and X enjoys a weak $(1,p)$ -Poincaré inequality, from Cheeger (see [19, Theorem 6.1]) we have $\text{Lip}\varphi(y) = g_\varphi(y)$ for μ -a.a. $y \in X$, where g_φ is the minimal p -weak upper gradient for φ . Hence

$$\|D_\mu \varphi\|_{L^\infty_\mu(X; \mathbb{R}^N)} \leq \frac{\theta}{\rho(1-\tau)}$$

because $|D_\mu \varphi(y)| \leq \theta |g_\varphi(y)|$ for μ -a.a. $y \in X$. Consequently the property (e) holds. Finally, as (X, d) is a length space, from Colding and Minicozzi (see [21] and [19, Proposition 6.12]) we can assert that there exists $\delta > 0$ such that for every $x \in X$, every $\rho > 0$ and every $\tau \in]0, 1[$,

$$\mu(Q_\rho(x) \setminus Q_{\tau\rho}(x)) \leq 2^\delta (1 - \tau)^\delta \mu(Q_\rho(x)),$$

which implies the property (f). \square

The following result comes from Hajlasz and Koskela (see [34, Theorem 5.1] and [30, Theorem 9.7 and Remark 9.8]).

Theorem 3.9. *Assume that the assumptions of Theorem 3.3 hold and let $\kappa := \frac{\ln(C_d)}{\ln(2)}$ where $C_d \geq 1$ is given by the inequality (2.1). If $p > \kappa$ then for every $r > 0$ and every $\bar{x} \in X$ there exists $C(r, \bar{x}) > 0$ such that*

$$|u(y) - u(z)| \leq C(r, \bar{x}) d(y, z)^{1 - \frac{\kappa}{p}} \left(\int_{Q_r(\bar{x})} |\nabla_\mu u|^p d\mu \right)^{\frac{1}{p}}$$

for all $u \in W_\mu^{1,p}(X; \mathbb{R}^m)$ and all $y, z \in Q_r(\bar{x})$.

Proof of Theorem 3.9. Since (X, d, μ) is a compact (and so complete) doubling metric space, (X, d, μ) is proper, i.e., every closed ball is compact (see [36, Lemma 4.1.14]). Moreover, as (X, d) is a length space, from [30, Theorem 9.7 and Remark 9.8] we can assert that there exists $c > 0$ such that

$$|w(y) - w(z)| \leq cr^{\frac{\kappa}{p}} d(y, z)^{1 - \frac{\kappa}{p}} \left(\int_{Q_r(\bar{x})} g_w^p d\mu \right)^{\frac{1}{p}} \quad (3.10)$$

for all $w \in W_\mu^{1,p}(X)$, all $\bar{x} \in X$, all $r > 0$ and all $y, z \in Q_r(\bar{x})$, where $g_w \in L_\mu^p(X)$ denotes the minimal p -weak upper gradient for w . On the other hand, from (2.3) it is easy to see that for every $r > 0$ and every $\bar{x} \in X$ there exists $\theta(r, \bar{x}) > 0$ such that

$$\mu(Q_r(\bar{x})) \geq \theta(r, \bar{x}) r^\kappa.$$

Moreover, by the left inequality in (3.8) we have $\int_{Q_\rho(x)} g_w^p d\mu \leq \alpha^p \int_{Q_\rho(x)} |D_\mu w|^p d\mu$. Thus, for each $r > 0$, each $\bar{x} \in X$ and each $y, z \in Q_r(\bar{x})$, (3.10) can be rewritten as follows

$$|w(y) - w(z)| \leq C(r, \bar{x}) d(y, z)^{1 - \frac{\kappa}{p}} \left(\int_{Q_r(\bar{x})} |D_\mu w|^p d\mu \right)^{\frac{1}{p}}$$

with $C(r, \bar{x}) = \frac{c\alpha}{\theta(r, \bar{x})} > 0$. It follows that for every $r > 0$ and every $\bar{x} \in X$, we have

$$\begin{aligned} |u(y) - u(z)| &\leq C(r, \bar{x})d(y, z)^{1-\frac{\kappa}{p}} \sum_{i=1}^m \left(\int_{Q_r(\bar{x})} |D_\mu u_i|^p d\mu \right)^{\frac{1}{p}} \\ &\leq C(r, \bar{x})d(y, z)^{1-\frac{\kappa}{p}} \left(\int_{Q_r(\bar{x})} \sum_{i=1}^m |D_\mu u_i|^p d\mu \right)^{\frac{1}{p}} \\ &= C(r, \bar{x})d(y, z)^{1-\frac{\kappa}{p}} \left(\int_{Q_r(\bar{x})} |\nabla_\mu u|^p d\mu \right)^{\frac{1}{p}} \end{aligned}$$

for all $u \in W_\mu^{1,p}(X; \mathbb{R}^m)$ and all $y, z \in Q_r(\bar{x})$, and the proof is complete. \square

Denote the space of continuous functions from X to \mathbb{R}^m by $C(X; \mathbb{R}^m)$. As a consequence of [Theorem 3.9](#) we have

Corollary 3.10. *Under the assumptions of [Theorem 3.3](#), if $p > \kappa$ then $W_\mu^{1,p}(X; \mathbb{R}^m)$ continuously embeds into $C(X; \mathbb{R}^m)$, i.e., $W_\mu^{1,p}(X; \mathbb{R}^m) \subset C(X; \mathbb{R}^m)$ and there exists $K_0 > 0$ such that*

$$\|u\|_{C(X; \mathbb{R}^m)} \leq K_0 \|u\|_{W_\mu^{1,p}(X; \mathbb{R}^m)} \quad (3.11)$$

for all $u \in W_\mu^{1,p}(X; \mathbb{R}^m)$. Moreover, there exists $K_1 > 0$ such that

$$|u(y) - u(z)| \leq K_1 d(y, z)^{1-\frac{\kappa}{p}} \|\nabla_\mu u\|_{L_\mu^p(X; \mathbb{M})} \quad (3.12)$$

for all $u \in W_\mu^{1,p}(X; \mathbb{R}^m)$ and all $y, z \in X$.

Proof of [Corollary 3.10](#). Applying [Theorem 3.9](#) with $r = \text{diam}(X)$ and for a fixed $\bar{x} = x_0 \in X$, where $\text{diam}(X) = \sup\{d(y, z) : y, z \in X\} < \infty$ because (X, d) is compact, we see that

$$\begin{aligned} |u(y) - u(z)| &\leq C(\text{diam}(X), x_0) d(y, z)^{1-\frac{\kappa}{p}} \|\nabla_\mu u\|_{L_\mu^p(X; \mathbb{M})} \\ &\leq C(\text{diam}(X), x_0) \text{diam}(X)^{1-\frac{\kappa}{p}} \|\nabla_\mu u\|_{L_\mu^p(X; \mathbb{M})} \end{aligned} \quad (3.13)$$

for all $u \in W_\mu^{1,p}(X; \mathbb{R}^m)$ and all $y, z \in X$. Hence [\(3.12\)](#) holds with $K_1 = C(\text{diam}(X), x_0)$ and every $u \in W_\mu^{1,p}(X; \mathbb{R}^m)$ is $(1 - \frac{\kappa}{p})$ -Hölder continuous. In particular, it follows that $W_\mu^{1,p}(X; \mathbb{R}^m) \subset C(X; \mathbb{R}^m)$. On the other hand, given any $u \in W_\mu^{1,p}(X; \mathbb{R}^m)$ and any $y \in X$, we have $|u(y)|^p \leq 2^p (|u(y) - u(z)|^p + |u(z)|^p)$ for all $z \in X$, and consequently

$$\mu(X)^{\frac{1}{p}}|u(y)| \leq 2^{1+\frac{1}{p}} \left(\int_X |u(y) - u(z)|^p d\mu(z) \right)^{\frac{1}{p}} + 2^{1+\frac{1}{p}} \|u\|_{L_\mu^p(X; \mathbb{R}^m)}. \quad (3.14)$$

But, by (3.13) we have

$$\left(\int_X |u(y) - u(z)|^p d\mu(z) \right)^{\frac{1}{p}} \leq \mu(X)^{\frac{1}{p}} C(\text{diam}(X), x_0) \text{diam}(X)^{1-\frac{\kappa}{p}} \|\nabla_\mu u\|_{L_\mu^p(X; \mathbb{M})}. \quad (3.15)$$

Hence, combining (3.14) and (3.15) we deduce that for every $y \in X$,

$$\begin{aligned} |u(y)| &\leq 2^{1+\frac{1}{p}} C(\text{diam}(X), x_0) \text{diam}(X)^{1-\frac{\kappa}{p}} \|\nabla_\mu u\|_{L_\mu^p(X; \mathbb{M})} + \frac{2^{1+\frac{1}{p}}}{\mu(X)^{\frac{1}{p}}} \|u\|_{L_\mu^p(X; \mathbb{R}^m)} \\ &\leq K_0 \|u\|_{W_\mu^{1,p}(X; \mathbb{R}^m)} \end{aligned}$$

with $K_0 = \sup \left\{ 2^{1+\frac{1}{p}} C(\text{diam}(X), x_0) \text{diam}(X)^{1-\frac{\kappa}{p}}, \frac{2^{1+\frac{1}{p}}}{\mu(X)^{\frac{1}{p}}} \right\}$, and (3.11) follows. \square

As a consequence of Corollary 3.10 we have the following L_μ^∞ -compactness result in the framework of the p -Cheeger–Sobolev spaces with $p > \kappa$.

Corollary 3.11. *Under the assumptions of Theorem 3.3, if $p > \kappa$ and if $u \in W_\mu^{1,p}(X; \mathbb{R}^m)$ and $\{u_n\}_n \subset W_\mu^{1,p}(X; \mathbb{R}^m)$ are such that*

$$\lim_{n \rightarrow \infty} \|u_n - u\|_{L_\mu^p(X; \mathbb{R}^m)} = 0 \text{ and } \sup_{n \geq 1} \|\nabla_\mu u_n\|_{L_\mu^p(X; \mathbb{M})} < \infty, \quad (3.16)$$

then, up to a subsequence,

$$\lim_{n \rightarrow \infty} \|u_n - u\|_{L_\mu^\infty(X; \mathbb{R}^m)} = 0. \quad (3.17)$$

Proof of Corollary 3.11. From (3.16) we deduce that $\sup_{n \geq 1} \|u_n\|_{W_\mu^{1,p}(X; \mathbb{R}^m)} < \infty$, and using Corollary 3.10 we can assert that $\sup_{n \geq 1} \|u_n\|_{C(X; \mathbb{R}^m)} < \infty$, i.e., $\{u_n\}_n$ is bounded in $C(X; \mathbb{R}^m)$ with (X, d) a compact metric space. On the other hand, using (3.12) it is easy to see that $\{u_n\}_n$ is equicontinuous, and (3.17) follows from Arzelà–Ascoli’s theorem. \square

3.2. The De Giorgi–Letta lemma

Let $X = (X, d)$ be a metric space, let $\mathcal{O}(X)$ be the class of open subsets of X and let $\mathcal{B}(X)$ be the class of Borel subsets of X , i.e., the smallest σ -algebra containing the open

(or equivalently the closed) subsets of X . The following result is due to De Giorgi and Letta (see [24] and also [17, Lemma 3.3.6, p. 105]).

Lemma 3.12. *Let $\mathcal{S} : \mathcal{O}(X) \rightarrow [0, \infty]$ be an increasing set function, i.e., $\mathcal{S}(A) \leq \mathcal{S}(B)$ for all $A, B \in \mathcal{O}(X)$ such $A \subset B$, satisfying the following four conditions:*

- (a) $\mathcal{S}(\emptyset) = 0$;
- (b) \mathcal{S} is superadditive, i.e., $\mathcal{S}(A \cup B) \geq \mathcal{S}(A) + \mathcal{S}(B)$ for all $A, B \in \mathcal{O}(X)$ such that $A \cap B = \emptyset$;
- (c) \mathcal{S} is subadditive, i.e., $\mathcal{S}(A \cup B) \leq \mathcal{S}(A) + \mathcal{S}(B)$ for all $A, B \in \mathcal{O}(X)$;
- (d) there exists a finite Radon measure ν on X such that $\mathcal{S}(A) \leq \nu(A)$ for all $A \in \mathcal{O}(X)$.

Then, \mathcal{S} can be uniquely extended to a finite positive Radon measure on X which is absolutely continuous with respect to ν .

3.3. Integral representation of the Vitali envelope of a set function

What follows was first developed in [15,12] (see also [37]). Here we only recall what is needed for proving Theorem 2.7. Let (X, d) be a metric space, let $\mathcal{O}(X)$ be the class of open subsets of X and let μ be a positive finite Radon measure on X . We begin with the concept of differentiability with respect to μ of a set function.

Definition 3.13. We say that a set function $\Theta : \mathcal{O}(X) \rightarrow \mathbb{R}$ is differentiable with respect to μ if

$$d_\mu \Theta(x) := \lim_{\rho \rightarrow 0} \frac{\Theta(Q_\rho(x))}{\mu(Q_\rho(x))} \quad (3.18)$$

exists and is finite for μ -a.e. $x \in X$.

Remark 3.14. It is easy to see that the limit in (3.18) exists and is finite if and only if $-\infty < d_\mu^+ \Theta \leq d_\mu^- \Theta < \infty$, where $d_\mu^- \Theta : X \rightarrow [-\infty, \infty[$ and $d_\mu^+ \Theta : X \rightarrow]-\infty, \infty]$ are given by:

$$d_\mu^- \Theta(x) := \lim_{\rho \rightarrow 0} d_\mu^- \Theta(x, \rho) \text{ with } d_\mu^- \Theta(x, \rho) := \inf \left\{ \frac{\Theta(Q)}{\mu(Q)} : Q \in \text{Ba}(X, x, \rho) \right\}; \quad (3.19)$$

$$d_\mu^+ \Theta(x) := \lim_{\rho \rightarrow 0} d_\mu^+ \Theta(x, \rho) \text{ with } d_\mu^+ \Theta(x, \rho) := \sup \left\{ \frac{\Theta(Q)}{\mu(Q)} : Q \in \text{Ba}(X, x, \rho) \right\}, \quad (3.20)$$

where $\text{Ba}(X, x, \rho)$ denotes the class of open balls Q of X such that $x \in Q$, $\text{diam}(Q) \in]0, \rho[$ and $\mu(\partial Q) = 0$, where $\partial Q := \overline{Q} \setminus Q$. We then have $d_\mu \Theta = d_\mu^- \Theta = d_\mu^+ \Theta$.

Remark 3.15. In (3.19) and (3.20) we can replace $\text{Ba}(X, x, \rho)$ by $\text{Ba}(A, x, \rho)$ whenever $A \in \mathcal{O}(X)$ and $x \in A$.

For each $\varepsilon > 0$ and each $A \in \mathcal{O}(X)$, we denote the class of countable families $\{Q_i := Q_{\rho_i}(x_i)\}_{i \in I}$ of disjoint open balls of A with $x_i \in A$, $\rho_i = \text{diam}(Q_i) \in]0, \varepsilon[$ and $\mu(\partial Q_i) = 0$ such that $\mu(A \setminus \cup_{i \in I} Q_i) = 0$ by $\mathcal{V}_\varepsilon(A)$.

Definition 3.16. Given $\Theta : \mathcal{O}(X) \rightarrow \mathbb{R}$, for each $\varepsilon > 0$ we define $\Theta^\varepsilon : \mathcal{O}(X) \rightarrow [-\infty, \infty]$ by

$$\Theta^\varepsilon(A) := \inf \left\{ \sum_{i \in I} \Theta(Q_i) : \{Q_i\}_{i \in I} \in \mathcal{V}_\varepsilon(A) \right\}. \quad (3.21)$$

By the Vitali envelope of Θ we denote the set function $\Theta^* : \mathcal{O}(X) \rightarrow [-\infty, \infty]$ defined by

$$\Theta^*(A) := \sup_{\varepsilon > 0} \Theta^\varepsilon(A) = \lim_{\varepsilon \rightarrow 0} \Theta^\varepsilon(A). \quad (3.22)$$

The interest of [Definition 3.16](#) comes from the following integral representation result whose proof is given below.

Theorem 3.17. *Let $\Theta : \mathcal{O}(X) \rightarrow \mathbb{R}$ be a set function satisfying the following two conditions:*

- (a) *there exists a finite Radon measure ν on X which is absolutely continuous with respect to μ such that $|\Theta(A)| \leq \nu(A)$ for all $A \in \mathcal{O}(X)$;*
- (b) *Θ is subadditive, i.e., $\Theta(A) \leq \Theta(B) + \Theta(C)$ for all $A, B, C \in \mathcal{O}(X)$ with $B, C \subset A$, $B \cap C = \emptyset$ and $\mu(A \setminus B \cup C) = 0$.*

Then Θ is differentiable with respect to μ , $d_\mu \Theta \in L_\mu^1(X)$ and

$$\Theta^*(A) = \int_A d_\mu \Theta(x) d\mu(x)$$

for all $A \in \mathcal{O}(X)$.

As a direct consequence, we have

Corollary 3.18. *Let $\Theta : \mathcal{O}(X) \rightarrow \mathbb{R}$ be a set function satisfying the assumptions (a) and (b) of [Theorem 3.17](#). Then Θ and Θ^* are differentiable with respect to μ and $d_\mu \Theta^* = d_\mu \Theta$.*

Proof of [Theorem 3.17](#). First of all, from (a) we see that $-d_\mu \nu \leq d_\mu^- \Theta \leq d_\mu^+ \Theta \leq d_\mu \nu$. Hence $d_\mu^- \Theta, d_\mu^+ \Theta \in L_\mu^1(X)$ because ν is a finite Radon measure which is absolutely continuous with respect to the finite Radon measure μ . So $\lambda^-(A), \lambda^+(A) \in \mathbb{R}$ for all $A \in \mathcal{O}(X)$, where $\lambda^-, \lambda^+ : \mathcal{O}(X) \rightarrow \mathbb{R}$ are given by:

$$\lambda^-(A) := \int_A d_\mu^- \Theta(x) d\mu(x);$$

$$\lambda^+(A) := \int_A d_\mu^+ \Theta(x) d\mu(x).$$

In what follows, we consider $\bar{\Theta}^* : \mathcal{O}(X) \rightarrow \mathbb{R}$ defined by

$$\bar{\Theta}^*(A) := \inf_{\varepsilon > 0} \sup \left\{ \sum_{i \in I} \Theta(Q_i) : \{Q_i\}_{i \in I} \in \mathcal{V}_\varepsilon(A) \right\}. \quad (3.23)$$

(It is clear that $\Theta^* \leq \bar{\Theta}^*$. In fact, we are going to prove that under the assumptions (a) and (b) of [Theorem 3.17](#) we have $\Theta^*(A) = \bar{\Theta}^*(A) = \int_A d_\mu \Theta(x) d\mu(x)$ for all $A \in \mathcal{O}(X)$.) We divide the proof into three steps.

Step 1: proving that $\Theta^* = \lambda^-$ and $\bar{\Theta}^* = \lambda^+$. Define $\theta^-, \theta^+ : \mathcal{O}(X) \rightarrow \mathbb{R}$ by:

$$\theta^-(A) := \Theta(A) - \lambda^-(A);$$

$$\theta^+(A) := \Theta(A) - \lambda^+(A).$$

In what follows, θ^* (resp. $\bar{\theta}^*$) is defined by [\(3.22\)](#) (resp. [\(3.23\)](#)) with Θ replaced by θ^- (resp. θ^+).

Substep 1-1: an auxiliary lemma.

Lemma 3.19. *Under the assumption (a) of [Theorem 3.17](#) we have $\theta^* = \bar{\theta}^* = 0$.*

Proof of [Lemma 3.19](#). We only prove that $\theta^* = 0$. (The proof of $\bar{\theta}^* = 0$ follows from similar arguments and is left to the reader.)

First of all, from the assumption (a) it is clear that

$$|\theta^-(A)| \leq \hat{\nu}(A) \quad (3.24)$$

for all $A \in \mathcal{O}(X)$, where $\hat{\nu} := \nu + |\nu|$ is absolutely continuous with respect to μ (with $|\nu|$ denoting the total variation of ν).

Secondly, we can assert that

$$d_\mu^- \theta^- = 0, \quad (3.25)$$

where for any set function $s : \mathcal{O}(X) \rightarrow \mathbb{R}$, the function $d_\mu^- s : X \rightarrow [-\infty, \infty[$ (resp. $d_\mu^+ s : X \rightarrow]-\infty, \infty]$) is defined by [\(3.19\)](#) (resp. [\(3.20\)](#)) with Θ replaced by s . Indeed, for any $x \in X$, it is easily seen that

$$d_\mu^- \Theta(x, \rho) - d_\mu^+ \lambda^-(x, \rho) \leq d_\mu^- \theta^-(x, \rho) \leq d_\mu^- \Theta(x, \rho) - d_\mu^- \lambda^-(x, \rho).$$

for all $\rho > 0$, and letting $\rho \rightarrow 0$, we obtain

$$d_\mu^- \Theta(x) - d_\mu^+ \lambda^-(x) \leq d_\mu^- \theta^-(x) \leq d_\mu^- \Theta(x) - d_\mu^- \lambda^-(x).$$

But $d_\mu^- \lambda^-(x) = d_\mu^+ \lambda^-(x) = d_\mu^- \Theta(x)$, hence $d_\mu^- \theta^-(x) = 0$.

Finally, to conclude we prove that (3.24) and (3.25) imply $\theta^* = 0$. For this, we are going to prove the following two assertions:

$$\text{if } d_\mu^- \theta^- \leq 0 \text{ then } \theta^* \leq 0; \quad (3.26)$$

$$\text{under (3.24), if } d_\mu^- \theta^- \geq 0 \text{ then } \theta^* \geq 0. \quad (3.27)$$

Proof of (3.26). Fix $A \in \mathcal{O}(X)$. Fix any $\varepsilon > 0$. Then $d_\mu^- \theta^- < \varepsilon$, and so in particular $\lim_{\rho \rightarrow 0} d_\mu^- \theta^-(x, \rho) < \varepsilon$ for all $x \in A$. Hence, for each $x \in A$ there exists $\{\rho_{x,n}\}_n \subset]0, \varepsilon[$ with $\rho_{x,n} \rightarrow 0$ as $n \rightarrow \infty$ such that $d_\mu^- \theta^-(x, \rho_{x,n}) < \varepsilon$ for all $n \geq 1$. Taking Remark 3.15 into account, it follows that for each $x \in A$ and each $n \geq 1$ there is $Q_{x,n} \in \text{Ba}(A, x, \rho_{x,n})$ such that for each $x \in A$ and each $n \geq 1$,

$$\frac{\theta^-(Q_{x,n})}{\mu(Q_{x,n})} < \varepsilon. \quad (3.28)$$

Moreover, since $\text{diam}(\overline{Q}_{x,n}) = \text{diam}(Q_{x,n}) \leq \rho_{x,n}$ for all $x \in A$ and all $n \geq 1$, we have $\inf \{\text{diam}(\overline{Q}_{x,n}) : n \geq 1\} = 0$ (where $\overline{Q}_{x,n}$ denotes the closed ball corresponding to the open ball $Q_{x,n}$). Let \mathcal{F}_0 be the family of closed balls of X given by

$$\mathcal{F}_0 := \{\overline{Q}_{x,n} : x \in A \text{ and } n \geq 1\}.$$

As X satisfies the Vitali covering theorem, from the above we deduce that there exists a disjoint countable subfamily $\{\overline{Q}_i\}_{i \in I_0}$ of closed balls of \mathcal{F}_0 (with $Q_i \subset A$, $\mu(\partial Q_i) = 0$ and $\text{diam}(Q_i) \in]0, \varepsilon[$) such that $\mu(A \setminus \cup_{i \in I_0} \overline{Q}_i) = 0$, which means that $\{Q_i\}_{i \in I_0} \in \mathcal{V}_\varepsilon(A)$. From (3.28) we see that $\theta^-(Q_i) < \varepsilon \mu(Q_i)$ for all $i \in I_0$, hence

$$\sum_{i \in I_0} \theta^-(Q_i) \leq \varepsilon \sum_{i \in I_0} \mu(Q_i) = \varepsilon \mu(A).$$

Consequently $\theta^{-,\varepsilon}(A) \leq \varepsilon \mu(A)$ for all $\varepsilon > 0$, where $\theta^{-,\varepsilon}$ is defined by (3.21) with Θ replaced by θ^- , and letting $\varepsilon \rightarrow 0$ we obtain $\theta^*(A) \leq 0$.

Proof of (3.27). Fix $A \in \mathcal{O}(X)$. By Egorov's theorem, there exists a sequence $\{B_n\}_n$ of Borel subsets of A such that:

$$\lim_{n \rightarrow \infty} \mu(A \setminus B_n) = 0; \quad (3.29)$$

$$\lim_{\varepsilon \rightarrow 0} \sup_{x \in B_n} |d_\mu^- \theta^-(x) - d_\mu^- \theta^-(x, \varepsilon)| = 0 \text{ for all } n \geq 1. \quad (3.30)$$

As $\hat{\nu}$ is absolutely continuous with respect to μ , by (3.29) we have

$$\lim_{n \rightarrow \infty} \hat{\nu}(A \setminus B_n) = 0. \quad (3.31)$$

Moreover, as $d_\mu^- \theta^- \geq 0$, from (3.30) we deduce that

$$\lim_{\varepsilon \rightarrow 0} \inf_{x \in B_n} d_\mu^- \theta^-(x, \varepsilon) \geq 0 \text{ for all } n \geq 1. \quad (3.32)$$

Fix any $n \geq 1$ and any $\varepsilon > 0$. By definition of $\theta^{-, \varepsilon}$, there exists $\{Q_i\}_{i \in I} \in \mathcal{V}_\varepsilon(A)$ such that

$$\theta^{-, \varepsilon}(A) > \sum_{i \in I} \theta^-(Q_i) - \varepsilon. \quad (3.33)$$

Set $I_n := \{i \in I : Q_i \cap B_n \neq \emptyset\}$. Using (3.24) we have

$$\begin{aligned} \sum_{i \in I} \theta^-(Q_i) &= \sum_{i \in I_n} \theta^-(Q_i) + \sum_{i \in I \setminus I_n} \theta^-(Q_i) \geq \sum_{i \in I_n} \theta^-(Q_i) - \sum_{i \in I \setminus I_n} \hat{\nu}(Q_i) \\ &\geq \sum_{i \in I_n} \frac{\theta^-(Q_i)}{\mu(Q_i)} \mu(Q_i) - \hat{\nu} \left(\bigcup_{i \in I \setminus I_n} Q_i \right), \end{aligned}$$

and, choosing $x_i \in Q_i \cap B_n$ for each $i \in I_n$ and noticing that $\bigcup_{i \in I \setminus I_n} Q_i \subset A \setminus B_n$, it follows that

$$\begin{aligned} \sum_{i \in I} \theta^-(Q_i) &\geq \sum_{i \in I_n} d_\mu^- \theta^-(x_i, \varepsilon) \mu(Q_i) - \hat{\nu}(A \setminus B_n) \\ &\geq \inf_{x \in B_n} d_\mu^- \theta^-(x, \varepsilon) \sum_{i \in I_n} \mu(Q_i) - \hat{\nu}(A \setminus B_n). \end{aligned}$$

Taking (3.33) into account, we conclude that

$$\theta^{-, \varepsilon}(A) \geq \inf_{x \in B_n} d_\mu^- \theta^-(x, \varepsilon) \sum_{i \in I_n} \mu(Q_i) - \hat{\nu}(A \setminus B_n) - \varepsilon$$

for all $\varepsilon > 0$ and all $n \geq 1$, which gives $\theta^*(A) \geq 0$ by letting $\varepsilon \rightarrow 0$ and using (3.32) and then by letting $n \rightarrow \infty$ and using (3.31). \square

Substep 1-2: using Lemma 3.19. As λ^- and λ^+ are absolutely continuous with respect to μ , it is easy to see that:

$$\begin{aligned} \theta^* &= \Theta^* - \lambda^-; \\ \bar{\theta}^* &= \bar{\Theta}^* - \lambda^+. \end{aligned}$$

Hence $\Theta^* = \lambda^-$ and $\bar{\Theta}^* = \lambda^+$ by Lemma 3.19.

Step 2: proving that $\Theta^* = \overline{\Theta}^*$. We only need to prove that $\overline{\Theta}^* \leq \Theta^*$. For this, it is sufficient to show that for each open ball Q of X with $\mu(\partial Q) = 0$, one has

$$\Theta(Q) \leq \Theta^*(Q). \quad (3.34)$$

Fix any $\varepsilon > 0$. By definition of Θ^ε , there exists $\{Q_i\}_{i \in I} \in \mathcal{V}_\varepsilon(Q)$ such that

$$\sum_{i \in I} \Theta(Q_i) \leq \Theta^\varepsilon(Q) + \varepsilon. \quad (3.35)$$

Since $\mu(Q \setminus \cup_{i \in I} Q_i) = 0$ there is a sequence $\{I_n\}_n$ of finite subsets of I such that

$$\lim_{n \rightarrow \infty} \mu \left(Q \setminus \bigcup_{i \in I_n} Q_i \right) = \lim_{n \rightarrow \infty} \mu \left(\bigcup_{i \in I \setminus I_n} Q_i \right) = 0. \quad (3.36)$$

Fix any $n \geq 1$. As Θ is subadditive by assumption (b), we have

$$\Theta \left(\bigcup_{i \in I_n} Q_i \right) \leq \sum_{i \in I_n} \Theta(Q_i).$$

Moreover, $\mu(Q \setminus [(\cup_{i \in I_n} Q_i) \cup (Q \setminus \overline{\cup_{i \in I_n} Q_i})]) = 0$ because $\mu(\partial Q_i) = 0$ for all $i \in I_n$, so that

$$\Theta(Q) \leq \Theta \left(\bigcup_{i \in I_n} Q_i \right) + \Theta \left(Q \setminus \overline{\bigcup_{i \in I_n} Q_i} \right)$$

by using again the subadditivity of Θ , and consequently

$$\sum_{i \in I_n} \Theta(Q_i) \geq \Theta(Q) - \Theta \left(Q \setminus \overline{\bigcup_{i \in I_n} Q_i} \right).$$

Thus, using the assumption (a), we get

$$\begin{aligned} \sum_{i \in I} \Theta(Q_i) &= \sum_{i \in I \setminus I_n} \Theta(Q_i) + \sum_{i \in I_n} \Theta(Q_i) \\ &\geq \sum_{i \in I \setminus I_n} \Theta(Q_i) + \Theta(Q) - \Theta \left(Q \setminus \overline{\bigcup_{i \in I_n} Q_i} \right) \\ &\geq \Theta(Q) - \nu \left(\bigcup_{i \in I \setminus I_n} Q_i \right) - \nu \left(Q \setminus \overline{\bigcup_{i \in I_n} Q_i} \right). \end{aligned}$$

But, $\nu(\partial Q_i) = 0$ for all $i \in I_n$ because ν is absolutely with respect to μ , so that

$$\nu \left(Q \setminus \overline{\bigcup_{i \in I_n} Q_i} \right) = \nu \left(Q \setminus \bigcup_{i \in I_n} Q_i \right) = \nu \left(\bigcup_{i \in I \setminus I_n} Q_i \right),$$

and thus

$$\sum_{i \in I} \Theta(Q_i) \geq \Theta(Q) - 2\nu \left(\bigcup_{i \in I \setminus I_n} Q_i \right). \quad (3.37)$$

Combining (3.35) with (3.37) we conclude that

$$\Theta(Q) \leq \Theta^\varepsilon(Q) + 2\nu \left(\bigcup_{i \in I \setminus I_n} Q_i \right) + \varepsilon,$$

and (3.34) follows by letting $n \rightarrow \infty$ and using (3.36) and then by letting $\varepsilon \rightarrow 0$.

Step 3: end of the proof. From steps 1 and 2 we have

$$\int_X d_\mu^- \Theta(x) d\mu(x) = \Theta^*(X) = \bar{\Theta}^*(X) = \int_X d_\mu^+ \Theta(x) d\mu(x).$$

Thus $\int_X (d_\mu^+ \Theta(x) - d_\mu^- \Theta(x)) d\mu(x) = 0$. But $d_\mu^+ \Theta \geq d_\mu^- \Theta$, i.e., $d_\mu^+ \Theta - d_\mu^- \Theta \geq 0$, hence $d_\mu^+ \Theta - d_\mu^- \Theta = 0$, i.e., $d_\mu^+ \Theta = d_\mu^- \Theta$, and the proof of [Theorem 3.17](#) is complete. \square

3.4. Ru-usc integrands

Let (X, d, μ) be a metric space measure, where μ is a positive Radon measure on X , and let $L : X \times \mathbb{M} \rightarrow [0, \infty]$ be a Borel measurable integrand. For each $x \in X$, we denote the effective domain of $L(x, \cdot)$ by \mathbb{L}_x and, for each $a \in L_{\text{loc}, \mu}^1(X;]0, \infty])$, we define $\Delta_L^a : [0, 1] \rightarrow]-\infty, \infty]$ by

$$\Delta_L^a(t) := \sup_{x \in X} \sup_{\xi \in \mathbb{L}_x} \frac{L(x, t\xi) - L(x, \xi)}{a(x) + L(x, \xi)}.$$

(When (X, d) is compact, $L_{\text{loc}, \mu}^1(X;]0, \infty])$ can be replaced by $L_\mu^1(X;]0, \infty])$.)

Definition 3.20. We say that L is radially uniformly upper semicontinuous (ru-usc) if there exists $a \in L_{\text{loc}, \mu}^1(X;]0, \infty])$ such that

$$\overline{\lim}_{t \rightarrow 1^-} \Delta_L^a(t) \leq 0.$$

The concept of ru-usc integrand was introduced in [1] and then developed in [6–8, 41, 9, 11]. (Recently, this concept was used by Duerinckx and Gloria in [23] to deal with stochastic homogenization of unbounded nonconvex integrals with convex growth.)

Remark 3.21. If L is ru-usc then $\overline{\lim}_{t \rightarrow 1^-} L(x, t\xi) \leq L(x, \xi)$ for all $x \in X$ and all $\xi \in \mathbb{L}_x$. On the other hand, if there exist $x \in X$ and $\xi \in \mathbb{L}_x$ such that $L(x, \cdot)$ is lsc at ξ then, for each $a \in L_{\text{loc}}^1(U;]0, \infty])$, $\underline{\lim}_{t \rightarrow 1^-} \Delta_L^a(t) \geq 0$, and so if in addition L is ru-usc then $\lim_{t \rightarrow 1^-} \Delta_L^a(t) = 0$ for some $a \in L_{\text{loc}, \mu}^1(X;]0, \infty])$.

Remark 3.22. If, for every $x \in X$, $L(x, \cdot)$ is convex and $0 \in \mathbb{L}_x$, then L is ru-usc.

The interest of [Definition 3.20](#) comes from the following theorem. (For a proof we refer to [\[6, Theorem 3.5\]](#).) Recall that $\widehat{L} : X \times \mathbb{M} \rightarrow [0, \infty]$ is defined by

$$\widehat{L}(x, \xi) := \underline{\lim}_{t \rightarrow 1^-} L(x, t\xi).$$

Theorem 3.23. *If L is ru-usc and if for every $x \in X$,*

$$t\overline{\mathbb{L}}_x \subset \text{int}(\mathbb{L}_x) \text{ for all } t \in]0, 1[\quad (3.38)$$

and $L(x, \cdot)$ is lsc on $\text{int}(\mathbb{L}_x)$, where $\text{int}(\mathbb{L}_x)$ denotes the interior of \mathbb{L}_x , then:

- (a) $\widehat{L}(x, \xi) = \begin{cases} L(x, \xi) & \text{if } \xi \in \text{int}(\mathbb{L}_x) \\ \lim_{t \rightarrow 1^-} L(x, t\xi) & \text{if } \xi \in \partial\mathbb{L}_x \\ \infty & \text{otherwise;} \end{cases}$
- (b) \widehat{L} is ru-usc;
- (c) for every $x \in X$, $\widehat{L}(x, \cdot)$ is the lsc envelope of $L(x, \cdot)$.

For each $\rho > 0$, let $\mathcal{H}_\mu^\rho L : X \times \mathbb{M} \rightarrow [0, \infty]$ be given by

$$\mathcal{H}_\mu^\rho L(x, \xi) := \inf \left\{ \int_{Q_\rho(x)} L(y, \xi + \nabla_\mu w(y)) d\mu(y) : w \in W_{\mu,0}^{1,p}(Q_\rho(x); \mathbb{R}^m) \right\}.$$

Then, we have

$$\mathcal{Q}_\mu L(x, \xi) := \overline{\lim}_{\rho \rightarrow 0} \mathcal{H}_\mu^\rho L(x, \xi)$$

for all $x \in X$ and all $\xi \in \mathbb{M}$. The following proposition shows that ru-usc is conserved under μ -quasiconvexification.

Proposition 3.24. *If L is ru-usc then $\mathcal{Q}_\mu L$ is ru-usc.*

Proof of Proposition 3.24. Fix any $t \in [0, 1]$, any $x \in X$ and any $\xi \in \mathcal{Q}_\mu \mathbb{L}_x$ where $\mathcal{Q}_\mu \mathbb{L}_x$ denotes the effective domain of $\mathcal{Q}_\mu L(x, \cdot)$. Then $\mathcal{Q}_\mu L(x, \xi) = \overline{\lim}_{\rho \rightarrow 0} \mathcal{H}_\mu^\rho L(x, \xi) < \infty$ and without loss of generality we can suppose that $\mathcal{H}_\mu^\rho L(x, \xi) < \infty$ for all $\rho > 0$.

Fix any $\rho > 0$. By definition, there exists $\{w_n\}_n \subset W_{\mu,0}^{1,p}(Q_\rho(x); \mathbb{R}^m)$ such that:

$$\mathcal{H}_\mu^\rho L(x, \xi) = \lim_{n \rightarrow \infty} \int_{Q_\rho(x)} L(y, \xi + \nabla_\mu w_n(y)) d\mu(y); \quad (3.39)$$

$$\xi + \nabla_\mu w_n(y) \in \mathbb{L}_y \text{ for all } n \geq 1 \text{ and } \mu\text{-a.a. } y \in Q_\rho(x). \quad (3.40)$$

Moreover, for every $n \geq 1$,

$$\mathcal{H}_\mu^\rho L(x, t\xi) \leq \int_{Q_\rho(x)} L(y, t(\xi + \nabla_\mu w_n(y))) d\mu(y)$$

since $tw_n \in W_{\mu,0}^{1,p}(Q_\rho(x); \mathbb{R}^m)$, and so

$$\begin{aligned} & \mathcal{H}_\mu^\rho L(x, t\xi) - \mathcal{H}_\mu^\rho L(x, \xi) \\ & \leq \lim_{n \rightarrow \infty} \int_{Q_\rho(x)} (L(y, t(\xi + \nabla_\mu w_n(y))) - L(y, \xi + \nabla_\mu w_n(y))) d\mu(y). \end{aligned} \quad (3.41)$$

But L is ru-usc, i.e., there exists $a \in L_{\text{loc},\mu}^1(X;]0, \infty])$ such that $\overline{\lim}_{t \rightarrow 1^-} \Delta_L^a(t) \leq 0$ with $\Delta_L^a : [0, 1] \rightarrow]-\infty, \infty]$ defined by $\Delta_L^a(t) := \sup_{z \in X} \sup_{\xi \in \mathbb{L}_z} \frac{L(z, t\xi) - L(z, \xi)}{a(z) + L(z, \xi)}$. So, taking (3.40) into account, for every $n \geq 1$ and μ -a.e. $y \in Q_\rho(x)$,

$$L(y, t(\xi + \nabla_\mu w_n(y))) - L(y, \xi + \nabla_\mu w_n(y)) \leq \Delta_L^a(t)(a(y) + L(y, \xi + \nabla_\mu w_n^\rho(y))).$$

Hence

$$\begin{aligned} & \int_{Q_\rho(x)} (L(y, t(\xi + \nabla_\mu w_n(y))) - L(y, \xi + \nabla_\mu w_n(y))) d\mu \\ & \leq \Delta_L^a(t) \left(\int_{Q_\rho(x)} a(y) d\mu + \int_{Q_\rho(x)} L(y, \xi + \nabla_\mu w_n(y)) d\mu \right) \end{aligned}$$

for all $n \geq 1$. Letting $n \rightarrow \infty$ and using (3.39) and (3.41), it follows that

$$\mathcal{H}_\mu^\rho L(x, t\xi) - \mathcal{H}_\mu^\rho L(x, \xi) \leq \Delta_L^a(t) \left(\int_{Q_\rho(x)} a(y) d\mu(y) + \mathcal{H}_\mu^\rho L(x, \xi) \right) \quad (3.42)$$

for all $\rho > 0$, and so

$$\mathcal{Q}_\mu L(x, t\xi) - \mathcal{Q}_\mu L(x, \xi) \leq \Delta_L^a(t)(a(x) + \mathcal{Q}_\mu L(x, \xi))$$

by letting $\rho \rightarrow 0$ in (3.42), which implies that $\Delta_{\mathcal{Q}_\mu L}^a(t) \leq \Delta_L^a(t)$ for all $t \in [0, 1]$, and the proof is complete. \square

4. Proof of the integral representation theorem

This section is devoted to the proof of [Theorem 2.7](#) which is divided into six steps.

Step 1: integral representation of the \overline{E} . For each $u \in W_{\mu}^{1,p}(X; \mathbb{R}^m)$ we consider the set function $\mathcal{S}_u : \mathcal{O}(X) \rightarrow [0, \infty]$ given by

$$\mathcal{S}_u(A) := \overline{E}(u, A).$$

Recall that \mathfrak{G} is the effective domain of the functional $u \mapsto \int_X G(x, \nabla_{\mu} u(x)) d\mu(x)$.

Step 1 consists of proving the following lemma.

Lemma 4.1. *If (2.5), (2.6), (2.7), (2.12) and (2.13) hold then*

$$\mathcal{S}_u(A) = \int_A \lambda_u(x) d\mu(x)$$

for all $u \in \mathfrak{G}$ and all $A \in \mathcal{O}(X)$ with $\lambda_u \in L_{\mu}^1(X)$ given by

$$\lambda_u(x) = \lim_{\rho \rightarrow 0} \frac{\mathcal{S}_u(Q_{\rho}(x))}{\mu(Q_{\rho}(x))}.$$

Proof of Lemma 4.1. Fix $u \in \mathfrak{G}$. Using the right inequality in (2.12) we see that

$$\mathcal{S}_u(A) \leq \beta \mu(A) + \beta \int_A G(x, \nabla_{\mu} u(x)) d\mu(x) < \infty \quad (4.1)$$

for all $A \in \mathcal{O}(X)$. Thus, the condition (d) of Lemma 3.12 is satisfied with $\nu = \beta(1 + G(x, \nabla_{\mu} u(x))) d\mu$ (which is absolutely continuous with respect to μ). On the other hand, it is easily seen that the conditions (a) and (b) of Lemma 3.12 are satisfied. Hence, the proof is completed provided we prove the condition (c) of Lemma 3.12, i.e.,

$$\mathcal{S}_u(A \cup B) \leq \mathcal{S}_u(A) + \mathcal{S}_u(B) \text{ for all } A, B \in \mathcal{O}(X). \quad (4.2)$$

Indeed, by Lemma 3.12, the set function \mathcal{S}_u can be (uniquely) extended to a (finite) positive Radon measure which is absolutely continuous with respect to μ , and the theorem follows by using Radon–Nikodym’s theorem and then Lebesgue’s differentiation theorem.

Remark 4.2. In fact, Lemma 4.1 establishes that for every $u \in \mathfrak{G}$, $E(u, \cdot)$ can be uniquely extended to a finite positive Radon measure on X which is absolutely continuous with respect to μ .

Substep 1-2: an auxiliary result for proving Lemma 4.1. To show (4.2) we need the following lemma.

Lemma 4.3. *If $U, V, Z, T \in \mathcal{O}(X)$ are such that $\overline{Z} \subset U$ and $T \subset V$, then*

$$\mathcal{S}_u(Z \cup T) \leq \mathcal{S}_u(U) + \mathcal{S}_u(V). \quad (4.3)$$

Proof of Lemma 4.3. Let $\{u_n\}_n$ and $\{v_n\}_n$ be two sequences in $W_\mu^{1,p}(X; \mathbb{R}^m)$ such that:

$$\|u_n - u\|_{L_\mu^p(X; \mathbb{R}^m)} \rightarrow 0; \quad (4.4)$$

$$\|v_n - u\|_{L_\mu^p(X; \mathbb{R}^m)} \rightarrow 0; \quad (4.5)$$

$$\lim_{n \rightarrow \infty} \int_U L(x, \nabla_\mu u_n(x)) d\mu(x) = \mathcal{S}_u(U) < \infty; \quad (4.6)$$

$$\lim_{n \rightarrow \infty} \int_V L(x, \nabla_\mu v_n(x)) d\mu(x) = \mathcal{S}_u(V) < \infty. \quad (4.7)$$

Since L is p -coercive (see (2.5) and the left inequality in (2.12)), from (4.6) and (4.7) we see that $\sup_n \|\nabla_\mu u_n\|_{L_\mu^p(X; \mathbb{M})} < \infty$ and $\sup_n \|\nabla_\mu v_n\|_{L_\mu^p(X; \mathbb{M})} < \infty$. As $p > \kappa$, taking (4.4) and (4.5) into account, by Corollary 3.11 we can assert, up to a subsequence, that:

$$\|u_n - u\|_{L_\mu^\infty(X; \mathbb{R}^m)} \rightarrow 0; \quad (4.8)$$

$$\|v_n - u\|_{L_\mu^\infty(X; \mathbb{R}^m)} \rightarrow 0. \quad (4.9)$$

Fix $\delta \in]0, \text{dist}(Z, \partial U)[$ with $\partial U := \overline{U} \setminus U$, fix any $q \geq 1$ and consider $W_i^-, W_i^+ \subset X$ given by:

$$W_i^- := \left\{ x \in X : \text{dist}(x, Z) \leq \frac{\delta}{3} + \frac{(i-1)\delta}{3q} \right\};$$

$$W_i^+ := \left\{ x \in X : \frac{\delta}{3} + \frac{i\delta}{3q} \leq \text{dist}(x, Z) \right\},$$

where $i \in \{1, \dots, q\}$. For every $i \in \{1, \dots, q\}$ there exists a Urysohn function $\varphi_i \in \text{Lip}(X)$ for the pair (W_i^+, W_i^-) . Fix any $n \geq 1$ and define $w_n^i \in W_\mu^{1,p}(X; \mathbb{R}^m)$ by

$$w_n^i := \varphi_i u_n + (1 - \varphi_i) v_n. \quad (4.10)$$

Fix any $t \in]0, 1[$. Setting $W_i := X \setminus (W_i^- \cup W_i^+)$ and using Theorem 3.3(d) and (3.1) we have

$$\begin{aligned} \nabla_\mu(tw_n^i) &= t\nabla_\mu w_n^i \\ &= \begin{cases} t\nabla_\mu u_n & \text{in } W_i^- \\ (1-t)\frac{t}{1-t}D_\mu\varphi_i \otimes (u_n - v_n) + t(\varphi_i\nabla_\mu u_n + (1-\varphi_i)\nabla_\mu v_n) & \text{in } W_i \\ t\nabla_\mu v_n & \text{in } W_i^+. \end{cases} \end{aligned}$$

Noticing that $Z \cup T = ((Z \cup T) \cap W_i^-) \cup (W \cap W_i) \cup (T \cap W_i^+)$ with $(Z \cup T) \cap W_i^- \subset U$, $T \cap W_i^+ \subset V$ and $W := T \cap \{x \in U : \frac{\delta}{3} < \text{dist}(x, Z) < \frac{2\delta}{3}\}$ we deduce that for every $i \in \{1, \dots, q\}$,

$$\begin{aligned} \int_{Z \cup T} L(x, t\nabla_\mu w_n^i) d\mu &\leq \int_U L(x, t\nabla_\mu u_n) d\mu + \int_V L(x, t\nabla_\mu v_n) d\mu \\ &\quad + \int_{W \cap W_i} L(x, t\nabla_\mu w_n^i) d\mu. \end{aligned} \quad (4.11)$$

Fix any $i \in \{1, \dots, q\}$. From the right inequality in (2.12) and the inequality (2.7) we see that

$$\begin{aligned} \int_{W \cap W_i} L(x, t\nabla_\mu w_n^i) d\mu &\leq \beta\mu(W \cap W_i) + \beta \int_{W \cap W_i} G(x, t\nabla_\mu w_n^i) d\mu \\ &\leq \beta(1 + \gamma)\mu(W \cap W_i) \\ &\quad + \beta\gamma \int_{W \cap W_i} G(x, \varphi_i \nabla_\mu u_n + (1 - \varphi_i) \nabla_\mu v_n) d\mu \\ &\quad + \beta\gamma \int_{W \cap W_i} G\left(x, \frac{t}{1-t} D_\mu \varphi_i \otimes (u_n - v_n)\right) d\mu, \end{aligned}$$

and by using again the inequality (2.7) and the left inequality in (2.12) we obtain

$$\begin{aligned} \int_{W \cap W_i} L(x, t\nabla_\mu w_n^i) d\mu &\leq \beta(1 + \gamma + \gamma^2)\mu(W \cap W_i) \\ &\quad + \frac{\beta\gamma^2}{\alpha} \left(\int_{W \cap W_i} L(x, \nabla_\mu u_n) d\mu + \int_{W \cap W_i} L(x, \nabla_\mu v_n) d\mu \right) \\ &\quad + \beta\gamma \int_{W \cap W_i} G\left(x, \frac{t}{1-t} D_\mu \varphi_i \otimes (u_n - v_n)\right) d\mu. \end{aligned} \quad (4.12)$$

On the other hand, we have

$$\left| \frac{t}{1-t} D_\mu \varphi_i(x) \otimes (u_n(x) - v_n(x)) \right| \leq \frac{t}{1-t} \|D_\mu \varphi_i\|_{L_\mu^\infty(X)} \|u_n - v_n\|_{L_\mu^\infty(X; \mathbb{R}^m)}$$

for μ -a.a. $x \in X$. But $\lim_{n \rightarrow \infty} \|u_n - v_n\|_{L_\mu^\infty(X; \mathbb{R}^m)} = 0$ by (4.8) and (4.9), hence for each $t \in]0, 1[$ and each $i \in \{1, \dots, q\}$ there exists $n_{t,i} \geq 1$ such that

$$\left| \frac{t}{1-t} D_\mu \varphi_i(x) \otimes (u_n(x) - v_n(x)) \right| \leq r$$

for μ -a.a. $x \in X$ and all $n \geq n_{t,i}$ with $r > 0$ given by (2.6). Hence

$$\int_{W \cap W_i} G\left(x, \frac{t}{1-t} D_\mu \varphi_i \otimes (u_n - v_n)\right) d\mu \leq \int_{W \cap W_i} \sup_{|\xi| \leq r} G(x, \xi) d\mu(x) \quad (4.13)$$

for all $n \geq N_{t,q}$ with $N_{t,q} = \max\{n_{t,i} : i \in \{1, \dots, q\}\}$. Moreover, we have:

$$\int_U L(x, t \nabla_\mu u_n) d\mu \leq \int_U L(x, \nabla_\mu u_n) d\mu + \Delta_L^a(t) \left(\int_U a(x) d\mu(x) + \int_U L(x, \nabla_\mu u_n) d\mu \right); \quad (4.14)$$

$$\int_V L(x, t \nabla_\mu v_n) d\mu \leq \int_V L(x, \nabla_\mu v_n) d\mu + \Delta_L^a(t) \left(\int_V a(x) d\mu(x) + \int_V L(x, \nabla_\mu v_n) d\mu \right), \quad (4.15)$$

where $a \in L^1(X;]0, \infty])$ is given by (2.13) (and $\overline{\lim}_{t \rightarrow 1^-} \Delta_L^a(t) \leq 0$ because L is ru-use).

Taking (4.13) into account and substituting (4.12), (4.14) and (4.15) into (4.11) and then averaging these inequalities, it follows that for every $q \geq 1$, every $t \in]0, 1[$ and every $n \geq N_{t,q}$, there exists $i_{n,t,q} \in \{1, \dots, q\}$ such that

$$\begin{aligned} & \int_{Z \cup T} L(x, \nabla_\mu (t w_n^{i_{n,t,q}})) d\mu \\ & \leq \int_U L(x, \nabla_\mu u_n) d\mu + \Delta_L^a(t) \left(\int_U a(x) d\mu(x) + \int_U L(x, \nabla_\mu u_n) d\mu \right) \\ & \quad + \int_V L(x, \nabla_\mu v_n) d\mu + \Delta_L^a(t) \left(\int_V a(x) d\mu(x) + \int_V L(x, \nabla_\mu v_n) d\mu \right) \\ & \quad + \frac{c}{q} \left(\int_X \sup_{|\xi| \leq r} G(x, \xi) d\mu + \int_U L(x, \nabla_\mu u_n) d\mu + \int_V L(x, \nabla_\mu v_n) d\mu \right) \end{aligned}$$

with $c = \max\{\beta(1 + \gamma + \gamma^2) + 1, \frac{\beta \gamma^2}{\alpha}\}$, where $\int_X \sup_{|\xi| \leq r} G(x, \xi) d\mu < \infty$ by (2.6). Thus, letting $n \rightarrow \infty$, $t \rightarrow 1^-$ and $q \rightarrow \infty$ and using (4.6) and (4.7), we get

$$\overline{\lim}_{q \rightarrow \infty} \overline{\lim}_{t \rightarrow 1^-} \overline{\lim}_{n \rightarrow \infty} \int_{Z \cup T} L(x, \nabla_\mu (t w_n^{i_{n,t,q}})) d\mu \leq \mathcal{S}_u(U) + \mathcal{S}_u(V). \quad (4.16)$$

On the other hand, taking (4.10) into account and using (4.4) and (4.5) we see that

$$\lim_{q \rightarrow \infty} \lim_{t \rightarrow 1^-} \lim_{n \rightarrow \infty} \|t w_n^{i_{n,t,q}} - u\|_{L_\mu^p(X; \mathbb{R}^m)} = 0.$$

By diagonalization, there exist increasing mappings $n \mapsto t_n$ and $n \mapsto q_n$ with $t_n \rightarrow 1^-$ and $q_n \rightarrow \infty$ such that:

$$\begin{aligned} \varliminf_{n \rightarrow \infty} \int_{Z \cup T} L(x, \nabla_\mu \hat{w}_n) d\mu &\leq \overline{\lim}_{n \rightarrow \infty} \int_{Z \cup T} L(x, \nabla_\mu \hat{w}_n) d\mu \\ &\leq \overline{\lim}_{q \rightarrow \infty} \overline{\lim}_{t \rightarrow 1^-} \overline{\lim}_{n \rightarrow \infty} \int_{Z \cup T} L(x, \nabla_\mu (t w_n^{i_n, t, q})) d\mu; \\ \lim_{n \rightarrow \infty} \|\hat{w}_n - u\|_{L_\mu^p(X; \mathbb{R}^m)} &= 0, \end{aligned}$$

where $\hat{w}_n := t_n w_n^{i_n, t_n, q_n}$. Hence

$$\mathcal{S}_u(Z \cup T) \leq \overline{\lim}_{q \rightarrow \infty} \overline{\lim}_{t \rightarrow 1^-} \overline{\lim}_{n \rightarrow \infty} \int_{Z \cup T} L(x, \nabla_\mu (t w_n^{i_n, t, q})) d\mu,$$

and (4.3) follows from (4.16). \square

Substep 1-2: end of the proof of Lemma 4.1. We now prove (4.2). Fix $A, B \in \mathcal{O}(X)$. Fix any $\varepsilon > 0$ and consider $C, D \in \mathcal{O}(X)$ such that $\overline{C} \subset A$, $\overline{D} \subset B$ and

$$\beta \mu(E) + \beta \int_E g_u(x) d\mu(x) < \varepsilon$$

with $E := A \cup B \setminus \overline{C \cup D}$. Then $\mathcal{S}_u(E) \leq \varepsilon$ by (4.1). Let $\hat{C}, \hat{D} \in \mathcal{O}(X)$ be such that $\overline{C} \subset \hat{C}$, $\hat{C} \subset A$, $\overline{D} \subset \hat{D}$ and $\hat{D} \subset B$. Applying Lemma 4.3 with $U = \hat{C} \cup \hat{D}$, $V = T = E$ and $Z = C \cup D$ (resp. $U = A$, $V = B$, $Z = \hat{C}$ and $T = \hat{D}$) we obtain

$$\mathcal{S}_u(A \cup B) \leq \mathcal{S}_u(\hat{C} \cup \hat{D}) + \varepsilon \quad (\text{resp. } \mathcal{S}_u(\hat{C} \cup \hat{D}) \leq \mathcal{S}_u(A) + \mathcal{S}_u(B)),$$

and (4.2) follows by letting $\varepsilon \rightarrow 0$. \square

Step 2: another formula for \overline{E} . Let $\overline{E}_0 : W_{\mu,0}^{1,p}(X; \mathbb{R}^m) \times \mathcal{O}(X) \rightarrow [0, \infty]$ given by

$$\overline{E}_0(u, A) := \inf \left\{ \varliminf_{n \rightarrow \infty} E(u_n, A) : W_{\mu,0}^{1,p}(A; \mathbb{R}^m) \ni u_n - u \xrightarrow{L_\mu^p} 0 \right\}.$$

Step 2 consists of proving the following lemma.

Lemma 4.4. *If (2.5), (2.6), (2.7), (2.12) and (2.13) hold then for every $u \in \mathfrak{G}$ and every $A \in \mathcal{O}(X)$, one has:*

$$\overline{E}(u, A) \leq \overline{E}_0(u, A); \tag{4.17}$$

$$\overline{E}_0(tu, A) \leq (1 + \Delta_L^a(t)) \overline{E}(u, A) + \Delta_L^a(t) \|a\|_{L_\mu^1(A)} \quad \text{for all } t \in]0, 1[, \tag{4.18}$$

where $a \in L^1_\mu(X;]0, \infty])$ is given by (2.13). As a direct consequence we have

$$\overline{E}(u, A) = \lim_{t \rightarrow 1^-} \overline{E}_0(tu, A) \quad (4.19)$$

for all $u \in \mathfrak{G}$ and all $A \in \mathcal{O}(X)$.

Proof of Lemma 4.4. Fix $u \in \mathfrak{G}$ and $A \in \mathcal{O}(X)$. Clearly, $\overline{E}_0(u; A) \geq \overline{E}(u, A)$ because $W_{\mu,0}^{1,p}(A; \mathbb{R}^m) \subset W_\mu^{1,p}(X; \mathbb{R}^m)$. So (4.17) is satisfied. Thus, it remains to prove (4.18). Let $\{u_n\}_n \subset W_\mu^{1,p}(X; \mathbb{R}^m)$ be such that:

$$u_n \rightarrow u \text{ in } L^p_\mu(X; \mathbb{R}^m); \quad (4.20)$$

$$\lim_{n \rightarrow \infty} \int_A L(x, \nabla_\mu u_n(x)) d\mu(x) = \overline{E}(u, A) < \infty. \quad (4.21)$$

Remark 4.5. Without loss of generality we can always suppose that $\overline{E}(u, A) < \infty$. So, in fact, if we furthermore assume that (2.10) is satisfied, Lemma 4.4 remains true if we replace “ $u \in \mathfrak{G}$ ” by “ $u \in W_\mu^{1,p}(X; \mathbb{R}^m)$ ”.

Since L is p -coercive (see (2.5) and the left inequality in (2.12)), from (4.21) we see that $\sup_n \|\nabla_\mu u_n\|_{L^p_\mu(X; \mathbb{M})} < \infty$. As $p > \kappa$, taking (4.20) into account, by Corollary 3.11 we can assert, up to a subsequence, that:

$$\|u_n - u\|_{L^\infty(X; \mathbb{R}^m)} \rightarrow 0. \quad (4.22)$$

Fix $\delta > 0$ and set $A_\delta := \{x \in A : \text{dist}(x, \partial A) > \delta\}$ with $\partial A := \overline{A} \setminus A$. Fix any $n \geq 1$ and any $q \geq 1$ and consider $W_i^-, W_i^+ \subset X$ given by

$$W_i^- := \left\{ x \in X : \text{dist}(x, A_\delta) \leq \frac{\delta}{3} + \frac{(i-1)\delta}{3q} \right\};$$

$$W_i^+ := \left\{ x \in X : \frac{\delta}{3} + \frac{i\delta}{3q} \leq \text{dist}(x, A_\delta) \right\},$$

where $i \in \{1, \dots, q\}$. (Note that $W_i^- \subset A$.) For every $i \in \{1, \dots, q\}$ there exists a Urysohn function $\varphi_i \in \text{Lip}(X)$ for the pair (W_i^+, W_i^-) . Define $w_n^i : X \rightarrow \mathbb{R}^m$ by

$$w_n^i := \varphi_i u_n + (1 - \varphi_i)u. \quad (4.23)$$

Then $w_n^i - u \in W_{\mu,0}^{1,p}(A; \mathbb{R}^m)$. Fix any $t \in]0, 1[$. Setting $W_i := X \setminus (W_i^- \cup W_i^+) \subset A$ and using Theorem 3.3(d) and (3.1) we have

$$\begin{aligned} \nabla_\mu(tw_n^i) &= t\nabla_\mu w_n^i \\ &= \begin{cases} t\nabla_\mu u_n & \text{in } W_i^- \\ (1-t)\frac{t}{1-t}D_\mu\varphi_i \otimes (u_n - u) + t(\varphi_i\nabla_\mu u_n + (1-\varphi_i)\nabla_\mu u) & \text{in } W_i \\ t\nabla_\mu u & \text{in } W_i^+. \end{cases} \end{aligned}$$

Noticing that $A = W_i^- \cup W_i \cup (A \cap W_i^+)$ we deduce that for every $i \in \{1, \dots, q\}$,

$$\begin{aligned} \int_A L(x, t\nabla_\mu w_n^i) d\mu &\leq \int_A L(x, t\nabla_\mu u_n) d\mu + \int_{A \cap W_i^+} L(x, t\nabla_\mu u) d\mu \\ &\quad + \int_{W_i} L(x, t\nabla_\mu w_n^i) d\mu. \end{aligned} \quad (4.24)$$

Fix any $q \in \{1, \dots, q\}$. From the right inequality in (2.12) and the inequality (2.7) we see that

$$\begin{aligned} \int_{W_i} L(x, t\nabla_\mu w_n^i) d\mu &\leq \beta\mu(W_i) + \beta \int_{W_i} G(x, t\nabla_\mu w_n^i) d\mu \\ &\leq \beta(1 + \gamma)\mu(W_i) \\ &\quad + \beta\gamma \int_{W_i} G(x, \varphi_i \nabla_\mu u_n + (1 - \varphi_i) \nabla_\mu u) d\mu \\ &\quad + \beta\gamma \int_{W_i} G\left(x, \frac{t}{1-t} D_\mu \varphi_i \otimes (u_n - u)\right) d\mu, \end{aligned}$$

and by using again the inequality (2.7) and the left inequality in (2.12) we obtain

$$\begin{aligned} \int_{W_i} L(x, t\nabla_\mu w_n^i) d\mu &\leq \beta(1 + \gamma + \gamma^2)\mu(W_i) \\ &\quad + \frac{\beta\gamma^2}{\alpha} \left(\int_{W_i} L(x, \nabla_\mu u_n) d\mu + \int_{W_i} L(x, \nabla_\mu u) d\mu \right) \\ &\quad + \beta\gamma \int_{W_i} G\left(x, \frac{t}{1-t} D_\mu \varphi_i \otimes (u_n - u)\right) d\mu. \end{aligned} \quad (4.25)$$

Remark 4.6. As $u \in \mathfrak{G}$ and (2.12) holds, we have $\int_E L(x, \nabla_\mu u) d\mu < \infty$ for all $E \in \mathcal{O}(X)$.

On the other hand, we have

$$\left| \frac{t}{1-t} D_\mu \varphi_i(x) \otimes (u_n(x) - u(x)) \right| \leq \frac{t}{1-t} \|D_\mu \varphi_i\|_{L_\mu^\infty(X)} \|u_n - u\|_{L_\mu^\infty(X; \mathbb{R}^m)}$$

for μ -a.a. $x \in X$. But $\lim_{n \rightarrow \infty} \|u_n - u\|_{L_\mu^\infty(X; \mathbb{R}^m)} = 0$ by (4.22), hence for each $i \in \{1, \dots, q\}$ there exists $n_i \geq 1$ such that

$$\left| \frac{t}{1-t} D_\mu \varphi_i(x) \otimes (u_n(x) - u(x)) \right| \leq r$$

for μ -a.a. $x \in X$ and all $n \geq n_i$ with $r > 0$ given by (2.6). Hence

$$\int_{W_i} G\left(x, \frac{t}{1-t} D_\mu \varphi_i \otimes (u_n - u)\right) d\mu \leq \int_{W_i} \sup_{|\xi| \leq r} G(x, \xi) d\mu(x) \quad (4.26)$$

for all $n \geq N_q$ with $N_q = \max\{n_i : i \in \{1, \dots, q\}\}$. Moreover, we have:

$$\int_A L(x, t \nabla_\mu u_n) d\mu \leq \int_A L(x, \nabla_\mu u_n) d\mu + \Delta_L^a(t) \left(\int_A a(x) d\mu(x) + \int_A L(x, \nabla_\mu u_n) d\mu \right); \quad (4.27)$$

$$\begin{aligned} \int_{A \cap W_i^+} L(x, t \nabla_\mu u) d\mu &\leq \int_{A \cap W_i^+} L(x, \nabla_\mu u) d\mu \\ &+ \Delta_L^a(t) \left(\int_{A \cap W_i^+} a(x) d\mu(x) + \int_{A \cap W_i^+} L(x, \nabla_\mu u) d\mu \right), \end{aligned} \quad (4.28)$$

where $a \in L_\mu^1(X;]0, \infty])$ is given by (2.13) (and $\overline{\lim}_{t \rightarrow 1^-} \Delta_L^a(t) \leq 0$ because L is ru-usc).

Taking (4.26) into account and substituting (4.25), (4.27) and (4.28) into (4.24) and then averaging these inequalities, it follows that for every $q \geq 1$ and every $n \geq N_q$, there exists $i_{n,q} \in \{1, \dots, q\}$ such that

$$\begin{aligned} &\int_A L(x, \nabla_\mu (t w_n^{i_{n,q}})) d\mu \\ &\leq \int_A L(x, \nabla_\mu u_n) d\mu + \Delta_L^a(t) \left(\int_A a(x) d\mu(x) + \int_A L(x, \nabla_\mu u_n) d\mu \right) \\ &+ \frac{1}{q} \left[\int_A L(x, \nabla_\mu u) d\mu + \Delta_L^a(t) \left(\int_A a(x) d\mu(x) + \int_A L(x, \nabla_\mu u) d\mu \right) \right] \\ &+ \frac{c}{q} \left(\int_A \sup_{|\xi| \leq r} G(x, \xi) d\mu + \int_A L(x, \nabla_\mu u_n) d\mu + \int_A L(x, \nabla_\mu u) d\mu \right) \end{aligned}$$

with $c = \max\{\beta(1 + \gamma + \gamma^2) + 1, \frac{\beta\gamma^2}{\alpha}\}$, where $\int_A \sup_{|\xi| \leq r} G(x, \xi) d\mu < \infty$ by (2.6). Thus, letting $n \rightarrow \infty$ and $q \rightarrow \infty$ and using (4.21), we get

$$\overline{\lim}_{q \rightarrow \infty} \overline{\lim}_{n \rightarrow \infty} \int_A L(x, \nabla_\mu (t w_n^{i_{n,q}})) d\mu \leq (1 + \Delta_L^a(t)) \overline{E}(u, A) + \Delta_L^a(t) \int_A a(x) d\mu(x). \quad (4.29)$$

On the other hand, taking (4.23) into account and using (4.20) we see that

$$\lim_{q \rightarrow \infty} \lim_{n \rightarrow \infty} \|tw_n^{i_n, q} - tu\|_{L_\mu^p(X; \mathbb{R}^m)} = 0.$$

By diagonalization, there exists an increasing mapping $n \mapsto q_n$ with $q_n \rightarrow \infty$ such that:

$$\begin{aligned} \varliminf_{n \rightarrow \infty} \int_A L(x, \nabla_\mu \hat{w}_n) d\mu &\leq \overline{\lim}_{n \rightarrow \infty} \int_A L(x, \nabla_\mu \hat{w}_n) d\mu \leq \overline{\lim}_{q \rightarrow \infty} \overline{\lim}_{n \rightarrow \infty} \int_A L(x, \nabla_\mu (tw_n^{i_n, q})) d\mu; \\ \lim_{n \rightarrow \infty} \|\hat{w}_n - u\|_{L_\mu^p(X; \mathbb{R}^m)} &= 0, \end{aligned}$$

where $\hat{w}_n := tw_n^{i_n, q_n}$ is such that $\hat{w}_n - tu \in W_{\mu, 0}^{1, p}(A; \mathbb{R}^m)$. Hence

$$\overline{E}_0(tu, A) \leq \overline{\lim}_{q \rightarrow \infty} \overline{\lim}_{n \rightarrow \infty} \int_A L(x, \nabla_\mu (tw_n^{i_n, q})) d\mu,$$

and (4.18) follows from (4.29).

From (4.18) we deduce that

$$\overline{\lim}_{t \rightarrow 1^-} \overline{E}_0(tu, A) \leq \overline{E}(u, A).$$

Moreover, from (4.17) we have

$$\overline{E}(u, A) \leq \varliminf_{t \rightarrow 1^-} \overline{E}(tu, A) \leq \varliminf_{t \rightarrow 1^-} \overline{E}_0(tu, A),$$

which gives (4.19). \square

Step 3: using the Vitali envelope. For each $u \in W_\mu^{1, p}(X; \mathbb{R}^m)$ we consider the set function $\check{m}_u : \mathcal{O}(X) \rightarrow [0, \infty]$ defined by

$$\check{m}_u(A) := \overline{\lim}_{t \rightarrow 1^-} m_{tu}(A), \quad (4.30)$$

where, for each $z \in W_\mu^{1, p}(X; \mathbb{R}^m)$, $m_z : \mathcal{O}(X) \rightarrow [0, \infty]$ is given by

$$m_z(A) := \inf \left\{ E(v, A) : v - z \in W_{\mu, 0}^{1, p}(A; \mathbb{R}^m) \right\}. \quad (4.31)$$

For each $\varepsilon > 0$ and each $A \in \mathcal{O}(X)$, denote the class of countable families $\{Q_i := Q_{\rho_i}(x_i)\}_{i \in I}$ of disjoint open balls of A with $x_i \in A$, $\rho_i = \text{diam}(Q_i) \in]0, \varepsilon[$ and $\mu(\partial Q_i) = 0$ such that $\mu(A \setminus \cup_{i \in I} Q_i) = 0$ by $\mathcal{V}_\varepsilon(A)$, consider $\check{m}_u^\varepsilon : \mathcal{O}(X) \rightarrow [0, \infty]$ given by

$$\check{m}_u^\varepsilon(A) := \inf \left\{ \sum_{i \in I} \check{m}_u(Q_i) : \{Q_i\}_{i \in I} \in \mathcal{V}_\varepsilon(A) \right\},$$

and define $\check{m}_u^* : \mathcal{O}(X) \rightarrow [0, \infty]$ by

$$\check{m}_u^*(A) := \sup_{\varepsilon > 0} \check{m}_u^\varepsilon(A) = \lim_{\varepsilon \rightarrow 0} \check{m}_u^\varepsilon(A).$$

The set function \check{m}_u^* is called the Vitali envelope of \check{m}_u , see §3.3 for more details. (Note that as X satisfies the Vitali covering theorem, see [Proposition 3.7\(c\)](#) and [Remark 3.8](#), we have $\mathcal{V}_\varepsilon(A) \neq \emptyset$ for all $A \in \mathcal{O}(X)$ and all $\varepsilon > 0$.)

Step 3 consists of proving the following lemma.

Lemma 4.7. *If (2.5), (2.6), (2.7), (2.12) and (2.13) hold then*

$$\overline{E}(u, A) = \check{m}_u^*(A) \tag{4.32}$$

for all $u \in \mathfrak{G}$ and all $A \in \mathcal{O}(X)$.

Proof of Lemma 4.7. Fix $u \in \mathfrak{G}$. Given any $A \in \mathcal{O}(X)$, it is easy to see that $m_{tu}(A) \leq \overline{E}_0(tu, A)$ for all $t \in]0, 1[$, hence

$$\check{m}_u(A) = \overline{\lim}_{t \rightarrow 1^-} m_{tu}(A) \leq \lim_{t \rightarrow 1^-} \overline{E}_0(tu, A) = \overline{E}(u, A)$$

by [Lemma 4.4](#), and consequently

$$\check{m}_u^*(A) \leq \overline{E}(u, A)$$

because in the proof of [Lemma 4.1](#) it is established that $\overline{E}(u, \cdot)$ can be uniquely extended to a finite positive Radon measure on X , see [Remark 4.2](#). Hence, to establish (4.32), it remains to prove that

$$\overline{E}(u, A) \leq \check{m}_u^*(A) \tag{4.33}$$

with $\check{m}_u^*(A) < \infty$. Fix any $\varepsilon > 0$. By definition of $\check{m}_u^\varepsilon(A)$ there exists $\{Q_i\}_{i \in I} \in \mathcal{V}_\varepsilon(A)$ such that

$$\sum_{i \in I} \check{m}_u(Q_i) \leq \check{m}_u^\varepsilon(A) + \frac{\varepsilon}{2}. \tag{4.34}$$

Fix any $t > 0$. For each $i \in I$, by definition of $m_{tu}(Q_i)$ there exists $v_t^i \in W_\mu^{1,p}(Q_i; \mathbb{R}^m)$ such that $v_t^i - tu \in W_{\mu,0}^{1,p}(Q_i; \mathbb{R}^m)$ and

$$E(v_t^i, Q_i) \leq m_{tu}(Q_i) + \frac{\varepsilon \mu(Q_i)}{2\mu(A)}. \tag{4.35}$$

Define $u_t^\varepsilon : X \rightarrow \mathbb{R}^m$ by

$$u_t^\varepsilon := \begin{cases} tu & \text{in } X \setminus A \\ v_t^i & \text{in } Q_i. \end{cases}$$

Then $u_t^\varepsilon - tu \in W_{\mu,0}^{1,p}(A; \mathbb{R}^m)$. Moreover, because of [Proposition 3.7\(a\)](#), $\nabla_\mu u_t^\varepsilon(x) = \nabla_\mu v_t^i(x)$ for μ -a.e. $x \in Q_i$. From [\(4.35\)](#) we see that

$$E(u_t^\varepsilon, A) \leq \sum_{i \in I} m_{tu}(Q_i) + \frac{\varepsilon}{2},$$

hence $\overline{\lim}_{t \rightarrow 1^-} E(u_t^\varepsilon, A) \leq \check{m}_u^\varepsilon(A) + \varepsilon$ by using [\(4.34\)](#), and consequently

$$\overline{\lim}_{\varepsilon \rightarrow 0} \overline{\lim}_{t \rightarrow 1^-} E(u_t^\varepsilon, A) \leq \check{m}_u^*(A). \quad (4.36)$$

On the other hand, we have

$$\begin{aligned} \|u_t^\varepsilon - u\|_{L_{\mu}^{\chi p}(X; \mathbb{R}^m)}^p &\leq 2^p \left[\|u_t^\varepsilon - tu\|_{L_{\mu}^{\chi p}(X; \mathbb{R}^m)}^p + \|tu - u\|_{L_{\mu}^{\chi p}(X; \mathbb{R}^m)}^p \right] \\ &= 2^p \left[\left(\int_A |u_t^\varepsilon - tu|^{\chi p} d\mu \right)^{\frac{1}{\chi}} + (1-t)^p \|u\|_{L_{\mu}^{\chi p}(X; \mathbb{R}^m)}^p \right] \\ &= 2^p \left[\left(\sum_{i \in I} \int_{Q_i} |v_t^i - tu|^{\chi p} d\mu \right)^{\frac{1}{\chi}} + (1-t)^p \|u\|_{L_{\mu}^{\chi p}(X; \mathbb{R}^m)}^p \right] \\ &\leq 2^p \left[\sum_{i \in I} \left(\int_{Q_i} |v_t^i - tu|^{\chi p} d\mu \right)^{\frac{1}{\chi}} + (1-t)^p \|u\|_{L_{\mu}^{\chi p}(X; \mathbb{R}^m)}^p \right] \end{aligned}$$

with $\chi \geq 1$ given by [\(3.3\)](#). As X enjoys a p -Sobolev inequality, see [Proposition 3.7\(b\)](#), and $\text{diam}(Q_i) \in]0, \varepsilon[$ for all $i \in I$, we have

$$\sum_{i \in I} \left(\int_{Q_i} |v_t^i - tu|^{\chi p} d\mu \right)^{\frac{1}{\chi}} \leq \varepsilon^p C_S^p \sum_{i \in I} \int_{Q_i} |\nabla_\mu v_t^i - t \nabla_\mu u|^p d\mu$$

with $C_S > 0$ given by [\(3.3\)](#), hence

$$\sum_{i \in I} \left(\int_{Q_i} |v_t^i - tu|^{\chi p} d\mu \right)^{\frac{1}{\chi}} \leq 2^p \varepsilon^p C_S^p \left(\sum_{i \in I} \int_{Q_i} |\nabla_\mu v_t^i|^p d\mu + t^p \int_A |\nabla_\mu u|^p d\mu \right),$$

and consequently

$$\begin{aligned} \|u_t^\varepsilon - u\|_{L_{\mu}^{\chi p}(X; \mathbb{R}^m)}^p &\leq 2^{2p} \varepsilon^p C_S^p \left(\sum_{i \in I} \int_{Q_i} |\nabla_\mu v_t^i|^p d\mu + t^p \int_A |\nabla_\mu u|^p d\mu \right) \\ &\quad + 2^p (1-t)^p \|u\|_{L_{\mu}^{\chi p}(X; \mathbb{R}^m)}^p. \end{aligned} \quad (4.37)$$

Taking (2.5), the left inequality in (2.12), (4.34) and (4.35) into account, from (4.37) we deduce that

$$\overline{\lim}_{t \rightarrow 1^-} \|u_t^\varepsilon - u\|_{L_{\mu}^{\chi p}(X; \mathbb{R}^m)}^p \leq 2^p C_S^p \varepsilon^p \left(\frac{1}{\alpha c} (\check{m}_u^\varepsilon(A) + \varepsilon) + \int_A |\nabla_\mu u|^p d\mu \right),$$

which gives

$$\lim_{\varepsilon \rightarrow 0} \overline{\lim}_{t \rightarrow 1^-} \|u_t^\varepsilon - u\|_{L_{\mu}^{\chi p}(X; \mathbb{R}^m)}^p = 0 \quad (4.38)$$

because $\lim_{\varepsilon \rightarrow 0} \check{m}_u^\varepsilon(A) = \check{m}_u^*(A) < \infty$. According to (4.36) and (4.38), by diagonalization there exists a mapping $\varepsilon \mapsto t_\varepsilon$, with $t_\varepsilon \rightarrow 1^-$ as $\varepsilon \rightarrow 0$, such that:

$$\lim_{\varepsilon \rightarrow 0} \|w_\varepsilon - u\|_{L_{\mu}^{\chi p}(X; \mathbb{R}^m)}^p = 0; \quad (4.39)$$

$$\underline{\lim}_{\varepsilon \rightarrow 0} E(w_\varepsilon, A) \leq \check{m}_u^*(A) \quad (4.40)$$

with $w_\varepsilon := u_{t_\varepsilon}^\varepsilon$. Since $\chi p \geq p$, $w_\varepsilon \rightarrow u$ in $L_\mu^p(X; \mathbb{R}^m)$ by (4.39), and (4.33) follows from (4.40) by noticing that $\overline{E}(u; A) \leq \underline{\lim}_{\varepsilon \rightarrow 0} E(w_\varepsilon, A)$. \square

Step 4: differentiation with respect to μ . This step consists of applying [Theorem 3.17](#) (with $\Theta = \check{m}_u$ where $u \in \mathfrak{S}$). More precisely, Step 4 consists of proving the following lemma.

Lemma 4.8. *If (2.6), (2.7) and the left inequality in (2.12) hold then*

$$\check{m}_u^*(A) = \int_A \lim_{\rho \rightarrow 0} \frac{\check{m}_u(Q_\rho(x))}{\mu(Q_\rho(x))} d\mu(x) \quad (4.41)$$

for all $u \in \mathfrak{S}$ and all $A \in \mathcal{O}(X)$. As a consequence, if furthermore (2.5), the right inequality in (2.12) and (2.13) hold then

$$\overline{E}(u, A) = \int_A \lim_{\rho \rightarrow 0} \overline{\lim}_{t \rightarrow 1^-} \frac{m_{tu}(Q_\rho(x))}{\mu(Q_\rho(x))} d\mu(x) \quad (4.42)$$

for all $u \in \mathfrak{S}$ and all $A \in \mathcal{O}(X)$.

Proof of Lemma 4.8. Fix $u \in \mathfrak{S}$. The integral representation of $\overline{E}(u, \cdot)$ in (4.42) follows from (4.41) by using [Lemma 4.7](#) and the definition of \check{m}_u in (4.30). So, we only need to establish (4.41). For this, it is sufficient to prove that \check{m}_u is subadditive and there exists a finite Radon measure ν on X which is absolutely continuous with respect to μ such that

$$\check{m}_u(A) \leq \nu(A) \quad (4.43)$$

for all $A \in \mathcal{O}(X)$, and then to apply [Theorem 3.17](#). For each $t \in]0, 1[$, from the definition of m_{tu} in [\(4.31\)](#), it is easy to see that for every $A, B, C \in \mathcal{O}(X)$ with $B, C \subset A$, $B \cap C = \emptyset$ and $\mu(A \setminus B \cup C) = 0$,

$$m_{tu}(A) \leq m_{tu}(B) + m_{tu}(C),$$

and so

$$\overline{\lim}_{t \rightarrow 1^-} m_{tu}(A) \leq \overline{\lim}_{t \rightarrow 1^-} m_{tu}(B) + \overline{\lim}_{t \rightarrow 1^-} m_{tu}(C), \text{ i.e., } \check{m}_u(A) \leq \check{m}_u(B) + \check{m}_u(C),$$

which shows the subadditivity of \check{m}_u . On the other hand, given any $t \in]0, 1[$, by using the right inequality in [\(2.12\)](#) we have

$$m_{tu}(A) \leq \beta\mu(A) + \beta \int_A G(x, t\nabla_\mu u(x)) d\mu(x).$$

But, from [\(2.7\)](#) we see that $G(x, t\nabla_\mu u(x)) \leq \gamma(1 + G(x, \nabla_\mu u(x)) + G(x, 0))$ for μ -a.a. $x \in X$, hence

$$m_{tu}(A) \leq \beta\mu(A) + \beta\gamma\mu(A) + \beta\gamma \int_A (G(x, \nabla_\mu u(x)) + G(x, 0)) d\mu(x).$$

Letting $t \rightarrow 1^-$, we conclude that

$$\check{m}_u(A) \leq c \left(\mu(A) + \int_A (G(x, \nabla_\mu u(x)) + G(x, 0)) d\mu(x) \right),$$

with $c := \beta(1 + \gamma)$. Thus [\(4.43\)](#) is satisfied with the Radon measure $\nu := c(1 + G(x, \nabla_\mu u(x)) + G(x, 0))d\mu$ which is necessarily finite since $u \in \mathfrak{S}$ and $G(\cdot, 0) \in L_\mu^1(X)$ because $G(\cdot, 0) \leq \sup_{|\xi| \leq r} G(\cdot, \xi)$ and $\sup_{|\xi| \leq r} G(\cdot, \xi) \in L_\mu^1(X)$ by [\(2.6\)](#). \square

Step 5: linking the relaxed integrand to the differentiation of the Vitali envelope. According to [\(4.42\)](#), the proof of [Theorem 2.7](#) will be completed (see Substep 5-2 and also Step 6) if we prove that for each $u \in \mathfrak{S}$ and μ -a.e. $x \in X$, we have:

$$\begin{aligned} \lim_{\rho \rightarrow 0} \overline{\lim}_{t \rightarrow 1^-} \frac{m_{tu}(Q_\rho(x))}{\mu(Q_\rho(x))} &\geq \lim_{t \rightarrow 1^-} \overline{\lim}_{\rho \rightarrow 0} \frac{m_{tu_x}(Q_\rho(x))}{\mu(Q_\rho(x))}, \\ \text{i.e., } \lim_{\rho \rightarrow 0} \frac{\check{m}_u(Q_\rho(x))}{\mu(Q_\rho(x))} &\geq \lim_{t \rightarrow 1^-} \overline{\lim}_{\rho \rightarrow 0} \frac{m_{tu_x}(Q_\rho(x))}{\mu(Q_\rho(x))}; \end{aligned} \quad (4.44)$$

$$\begin{aligned}
\lim_{\rho \rightarrow 0} \overline{\lim}_{t \rightarrow 1^-} \frac{\mathfrak{m}_{tu}(Q_\rho(x))}{\mu(Q_\rho(x))} &\leq \underline{\lim}_{t \rightarrow 1^-} \overline{\lim}_{\rho \rightarrow 0} \frac{\mathfrak{m}_{tu_x}(Q_\rho(x))}{\mu(Q_\rho(x))}, \\
\text{i.e., } \lim_{\rho \rightarrow 0} \frac{\check{\mathfrak{m}}_u(Q_\rho(x))}{\mu(Q_\rho(x))} &\leq \underline{\lim}_{t \rightarrow 1^-} \overline{\lim}_{\rho \rightarrow 0} \frac{\mathfrak{m}_{tu_x}(Q_\rho(x))}{\mu(Q_\rho(x))}, \tag{4.45}
\end{aligned}$$

where $u_x \in W_\mu^{1,p}(\Omega; \mathbb{R}^m)$ is given by [Proposition 3.7\(d\)](#) (and satisfies [\(3.4\)](#) and [\(3.5\)](#)).

Substep 5-1: proof of (4.44) and (4.45). We only give the proof of [\(4.44\)](#). As the proof of [\(4.45\)](#) uses the same method, details are left to the reader.

Fix $u \in \mathfrak{S}$. Fix any $\varepsilon > 0$, any $\tau \in]0, 1[$, any $\rho \in]0, \varepsilon[$, any $t \in]0, 1[$ and any $s \in]t, 1[$. By definition of $\mathfrak{m}_{su}(Q_{\tau\rho}(x))$, where there is no loss of generality in assuming that $\mu(\partial Q_{\tau\rho}(x)) = 0$, there exists $w : X \rightarrow \mathbb{R}^m$ such that $w - su \in W_{\mu,0}^{1,p}(Q_{\tau\rho}(x); \mathbb{R}^m)$ and

$$\int_{Q_{\tau\rho}(x)} L(y, \nabla_\mu w(y)) d\mu(y) \leq \mathfrak{m}_{su}(Q_{\tau\rho}(x)) + \varepsilon \mu(Q_{\tau\rho}(x)). \tag{4.46}$$

By [Proposition 3.7\(e\)](#) there is a Urysohn function $\varphi \in \text{Lip}(X)$ for the pair $(X \setminus Q_\rho(x), \overline{Q}_{\tau\rho}(x))$ such that

$$\|D_\mu \varphi\|_{L_\mu^\infty(X; \mathbb{R}^N)} \leq \frac{\theta}{\rho(1-\tau)} \tag{4.47}$$

for some $\theta > 0$ (which does not depend on ρ). Define $v \in W_\mu^{1,p}(Q_\rho(x); \mathbb{R}^m)$ by

$$v := \varphi \frac{t}{s} u + (1 - \varphi) \frac{t}{s} u_x.$$

Then $v - \frac{t}{s} u_x \in W_{\mu,0}^{1,p}(Q_\rho(x); \mathbb{R}^m)$. Using [Theorem 3.3\(d\)](#) and [\(3.1\)](#) we have

$$\begin{aligned}
\nabla_\mu(sv) &= \begin{cases} \nabla_\mu(tu) & \text{in } \overline{Q}_{\tau\rho}(x) \\ tD_\mu\varphi \otimes (u - u_x) + s(\varphi \frac{t}{s} \nabla_\mu u + (1 - \varphi) \frac{t}{s} \nabla_\mu u(x)) & \text{in } Q_\rho(x) \setminus \overline{Q}_{\tau\rho}(x) \end{cases} \\
&= \begin{cases} \nabla_\mu(tu) & \text{in } \overline{Q}_{\tau\rho}(x) \\ (1-t) \frac{t}{1-t} D_\mu\varphi \otimes (u - u_x) + t(\varphi \nabla_\mu u + (1 - \varphi) \nabla_\mu u(x)) & \text{in } Q_\rho(x) \setminus \overline{Q}_{\tau\rho}(x). \end{cases}
\end{aligned}$$

As $\frac{t}{s} w - tu \in W_{\mu,0}^{1,p}(Q_{\tau\rho}(x); \mathbb{R}^m)$ we have $sv + (\frac{t}{s} w - tu) - tu_x \in W_{\mu,0}^{1,p}(Q_\rho(x); \mathbb{R}^m)$. Noticing that $\mu(\partial Q_{\tau\rho}(x)) = 0$ and, because of [Proposition 3.7\(a\)](#), $\nabla_\mu(\frac{t}{s} w - tu)(y) = 0$ for μ -a.e. $y \in Q_\rho(x) \setminus \overline{Q}_{\tau\rho}(x)$, we see that

$$\begin{aligned}
\frac{\mathfrak{m}_{tu_x}(Q_\rho(x))}{\mu(Q_{\tau\rho}(x))} &\leq \frac{1}{\mu(Q_{\tau\rho}(x))} \int_{Q_\rho(x)} L\left(y, \nabla_\mu(sv) + \nabla_\mu\left(\frac{t}{s} w - tu\right)\right) d\mu \\
&= \frac{1}{\mu(Q_{\tau\rho}(x))} \int_{\overline{Q}_{\tau\rho}(x)} L\left(y, \nabla_\mu(tu) + \nabla_\mu\left(\frac{t}{s} w - tu\right)\right) d\mu
\end{aligned}$$

$$\begin{aligned}
& + \frac{1}{\mu(Q_{\tau\rho}(x))} \int_{Q_\rho(x) \setminus \overline{Q_{\tau\rho}(x)}} L(y, \nabla_\mu(sv)) d\mu \\
& = \frac{1}{\mu(Q_{\tau\rho}(x))} \int_{Q_{\tau\rho}(x)} L\left(y, \frac{t}{s} \nabla_\mu w\right) d\mu \\
& + \frac{1}{\mu(Q_{\tau\rho}(x))} \int_{Q_\rho(x) \setminus Q_{\tau\rho}(x)} L(y, \nabla_\mu(sv)) d\mu.
\end{aligned}$$

It follows that

$$\begin{aligned}
\frac{m_{tu_x}(Q_\rho(x))}{\mu(Q_{\tau\rho}(x))} & \leq \frac{1}{\mu(Q_{\tau\rho}(x))} \int_{Q_\rho(x)} L(y, \nabla_\mu w) d\mu \\
& + \Delta_L^a \left(\frac{t}{s}\right) \left(\frac{\mu(Q_\rho(x))}{\mu(Q_{\tau\rho}(x))} \int_{Q_\rho(x)} a d\mu + \frac{1}{\mu(Q_{\tau\rho}(x))} \int_{Q_\rho(x)} L(y, \nabla_\mu w) d\mu \right) \\
& + \frac{1}{\mu(Q_{\tau\rho}(x))} \int_{Q_\rho(x) \setminus Q_{\tau\rho}(x)} L(y, \nabla_\mu(sv)) d\mu,
\end{aligned}$$

where $a \in L_\mu^1(X;]0, \infty])$ is given by (2.13) (and $\overline{\lim}_{r \rightarrow 1^-} \Delta_L^a(r) \leq 0$ because L is ru-usc). Taking (4.46), (2.7) and the right inequality in (2.12) into account we deduce that

$$\begin{aligned}
\frac{m_{tu_x}(Q_\rho(x))}{\mu(Q_{\tau\rho}(x))} & \leq \left(1 + \Delta_L^a \left(\frac{t}{s}\right)\right) \left(\frac{m_{su}(Q_{\tau\rho}(x))}{\mu(Q_{\tau\rho}(x))} + \varepsilon\right) \\
& + \Delta_L^a \left(\frac{t}{s}\right) \frac{\mu(Q_\rho(x))}{\mu(Q_{\tau\rho}(x))} \int_{Q_\rho(x)} a d\mu \\
& + \frac{c}{\mu(Q_{\tau\rho}(x))} \int_{Q_\rho(x) \setminus Q_{\tau\rho}(x)} G\left(y, \frac{t}{1-t} D_\mu \varphi \otimes (u - u_x)\right) d\mu \\
& + \frac{c}{\mu(Q_{\tau\rho}(x))} \int_{Q_\rho(x) \setminus Q_{\tau\rho}(x)} (G(y, \nabla_\mu u) + G(y, \nabla_\mu u(x))) d\mu \\
& + c \left(\frac{\mu(Q_\rho(x))}{\mu(Q_{\tau\rho}(x))} - 1\right)
\end{aligned}$$

with $c := \beta + \beta\gamma + \beta\gamma^2$, where $\gamma > 0$ and $\beta > 0$ given by (2.7) and (2.12) respectively. Thus, noticing that $\mu(Q_\rho(x)) \geq \mu(Q_{\tau\rho}(x))$ and letting $s \rightarrow 1^-$, we obtain

$$\frac{m_{tu_x}(Q_\rho(x))}{\mu(Q_\rho(x))} \leq \left(1 + \overline{\lim}_{s \rightarrow 1^-} \Delta_L^a \left(\frac{t}{s}\right)\right) \left(\frac{\check{m}_u(Q_{\tau\rho}(x))}{\mu(Q_{\tau\rho}(x))} + \varepsilon\right)$$

$$\begin{aligned}
& + \overline{\lim}_{s \rightarrow 1^-} \Delta_L^a \left(\frac{t}{s} \right) \frac{\mu(Q_\rho(x))}{\mu(Q_{\tau\rho}(x))} \int_{Q_\rho(x)} a d\mu \\
& + \frac{c}{\mu(Q_{\tau\rho}(x))} \int_{Q_\rho(x) \setminus Q_{\tau\rho}(x)} G \left(y, \frac{t}{1-t} D_\mu \varphi \otimes (u - u_x) \right) d\mu \\
& + \frac{c}{\mu(Q_{\tau\rho}(x))} \int_{Q_\rho(x) \setminus Q_{\tau\rho}(x)} (G(y, \nabla_\mu u) + G(y, \nabla_\mu u(x))) d\mu \\
& + c \left(\frac{\mu(Q_\rho(x))}{\mu(Q_{\tau\rho}(x))} - 1 \right). \tag{4.48}
\end{aligned}$$

On the other hand, by (4.47) we have

$$\begin{aligned}
\left| \frac{t}{1-t} D_\mu \varphi(y) \otimes (u(y) - u_x(y)) \right| & \leq \frac{t}{1-t} \|D_\mu \varphi\|_{L_\mu^\infty(X)} \|u - u_x\|_{L_\mu^\infty(Q_\rho(x); \mathbb{R}^m)} \\
& \leq \frac{t\theta}{(1-t)(1-\tau)} \frac{1}{\rho} \|u - u_x\|_{L_\mu^\infty(Q_\rho(x); \mathbb{R}^m)}
\end{aligned}$$

for μ -a.a. $y \in Q_\rho(x) \setminus Q_{\tau\rho}(x)$. But, since $p > \kappa$, $\lim_{\rho \rightarrow 0} \frac{1}{\rho} \|u - u_x\|_{L_\mu^\infty(Q_\rho(x); \mathbb{R}^m)} = 0$ by (3.5), hence there exists $\rho_0 > 0$ (which depends on t and τ) such that

$$\left| \frac{t}{1-t} D_\mu \varphi(y) \otimes (u(y) - u_x(y)) \right| \leq r$$

for μ -a.a. $y \in Q_\rho(x) \setminus Q_{\tau\rho}(x)$ and all $\rho \in]0, \rho_0[$ with $r > 0$ given by (2.6). Hence

$$\int_{Q_\rho(x) \setminus Q_{\tau\rho}(x)} G \left(y, \frac{t}{1-t} D_\mu \varphi \otimes (u - u_x) \right) d\mu \leq \int_{Q_\rho(x) \setminus Q_{\tau\rho}(x)} \sup_{|\xi| \leq r} G(y, \xi) d\mu(y) \tag{4.49}$$

for all $\rho \in]0, \rho_0[$. Moreover, it easy to see that:

$$\begin{aligned}
\int_{Q_\rho(x) \setminus Q_{\tau\rho}(x)} \sup_{|\xi| \leq r} G(y, \xi) d\mu(y) & \leq \mu(Q_\rho(x)) \int_{Q_\rho(x)} \left| \sup_{|\xi| \leq r} G(y, \xi) - \sup_{|\xi| \leq r} G(x, \xi) \right| d\mu(y) \\
& + \mu(Q_\rho(x) \setminus Q_{\tau\rho}(x)) \sup_{|\xi| \leq r} G(x, \xi); \tag{4.50}
\end{aligned}$$

$$\begin{aligned}
\int_{Q_\rho(x) \setminus Q_{\tau\rho}(x)} G(y, \nabla_\mu u(y)) d\mu(y) & \leq \mu(Q_\rho(x)) \int_{Q_\rho(x)} |G(y, \nabla_\mu u(y)) - G(x, \nabla_\mu u(x))| d\mu(y) \\
& + \mu(Q_\rho(x) \setminus Q_{\tau\rho}(x)) G(x, \nabla_\mu u(x)); \tag{4.51}
\end{aligned}$$

$$\begin{aligned}
\int_{Q_\rho(x) \setminus Q_{\tau\rho}(x)} G(y, \nabla_\mu u(x)) d\mu(y) & \leq \mu(Q_\rho(x)) \int_{Q_\rho(x)} |G(y, \nabla_\mu u(x)) - G(x, \nabla_\mu u(x))| d\mu(y) \\
& + \mu(Q_\rho(x) \setminus Q_{\tau\rho}(x)) G(x, \nabla_\mu u(x)). \tag{4.52}
\end{aligned}$$

Combining (4.49) with (4.50), (4.51) and (4.52) we deduce that

$$\begin{aligned}
\frac{m_{tu_x}(Q_\rho(x))}{\mu(Q_\rho(x))} &\leq \left(1 + \overline{\lim}_{s \rightarrow 1^-} \Delta_L^a \left(\frac{t}{s}\right)\right) \left(\frac{\check{m}_u(Q_{\tau\rho}(x))}{\mu(Q_{\tau\rho}(x))} + \varepsilon\right) \\
&\quad + \overline{\lim}_{s \rightarrow 1^-} \Delta_L^a \left(\frac{t}{s}\right) \frac{\mu(Q_\rho(x))}{\mu(Q_{\tau\rho}(x))} \int_{Q_\rho(x)} a(y) d\mu(y) \\
&\quad + c \frac{\mu(Q_\rho(x))}{\mu(Q_{\tau\rho}(x))} \int_{Q_\rho(x)} \left| \sup_{|\xi| \leq r} G(y, \xi) - \sup_{|\xi| \leq r} G(x, \xi) \right| d\mu(y) \\
&\quad + c \frac{\mu(Q_\rho(x))}{\mu(Q_{\tau\rho}(x))} \int_{Q_\rho(x)} |G(y, \nabla_\mu u(y)) - G(x, \nabla_\mu u(x))| d\mu(y) \\
&\quad + c \frac{\mu(Q_\rho(x))}{\mu(Q_{\tau\rho}(x))} \int_{Q_\rho(x)} |G(y, \nabla_\mu u(x)) - G(x, \nabla_\mu u(x))| d\mu(y) \\
&\quad + c \left(\frac{\mu(Q_\rho(x))}{\mu(Q_{\tau\rho}(x))} - 1\right) \sup_{|\xi| \leq r} G(x, \xi) \\
&\quad + 2c \left(\frac{\mu(Q_\rho(x))}{\mu(Q_{\tau\rho}(x))} - 1\right) G(x, \nabla_\mu u(x)) \\
&\quad + c \left(\frac{\mu(Q_\rho(x))}{\mu(Q_{\tau\rho}(x))} - 1\right). \tag{4.53}
\end{aligned}$$

As $\sup_{|\xi| \leq r} G(\cdot, \xi) \in L_\mu^1(X)$ by (2.6), and μ is a doubling measure, we have

$$\lim_{\eta \rightarrow 0} \int_{Q_\eta(x)} \left| \sup_{|\xi| \leq r} G(y, \xi) - \sup_{|\xi| \leq r} G(x, \xi) \right| d\mu(y) = 0. \tag{4.54}$$

In the same way, as $u \in \mathfrak{S}$, i.e., $G(\cdot, \nabla_\mu u(\cdot)) \in L_\mu^1(X)$ (and μ is a doubling measure) we can assert that

$$\lim_{\eta \rightarrow 0} \int_{Q_\eta(x)} |G(y, \nabla_\mu u(y)) - G(x, \nabla_\mu u(x))| d\mu(y) = 0, \tag{4.55}$$

and by (2.11) we have

$$\lim_{\eta \rightarrow 0} \int_{Q_\eta(x)} |G(y, \nabla_\mu u(x)) - G(x, \nabla_\mu u(x))| d\mu(y) = 0. \tag{4.56}$$

Moreover, since $a \in L^1_\mu(X;]0, \infty])$, it is clear that

$$\lim_{\eta \rightarrow 0} \int_{Q_\eta(x)} a(y) d\mu(y) = a(x). \quad (4.57)$$

Letting $\rho \rightarrow 0$ in (4.53) and using (4.54), (4.55), (4.56) and (4.57) we see that

$$\begin{aligned} \overline{\lim}_{\rho \rightarrow 0} \frac{m_{tu_x}(Q_\rho(x))}{\mu(Q_\rho(x))} &\leq \left(1 + \overline{\lim}_{s \rightarrow 1^-} \Delta_L^a \left(\frac{t}{s}\right)\right) \left(\lim_{\rho \rightarrow 0} \frac{\check{m}_u(Q_\rho(x))}{\mu(Q_\rho(x))} + \varepsilon\right) \\ &\quad + \overline{\lim}_{s \rightarrow 1^-} \Delta_L^a \left(\frac{t}{s}\right) \overline{\lim}_{\rho \rightarrow 0} \frac{\mu(Q_\rho(x))}{\mu(Q_{\tau\rho}(x))} a(x) \\ &\quad + c \left(\frac{\mu(Q_\rho(x))}{\mu(Q_{\tau\rho}(x))} - 1\right) \sup_{|\xi| \leq r} G(x, \xi) \\ &\quad + 2c \left(\overline{\lim}_{\rho \rightarrow 0} \frac{\mu(Q_\rho(x))}{\mu(Q_{\tau\rho}(x))} - 1\right) G(x, \nabla_\mu u(x)) \\ &\quad + c \left(\overline{\lim}_{\rho \rightarrow 0} \frac{\mu(Q_\rho(x))}{\mu(Q_{\tau\rho}(x))} - 1\right). \end{aligned} \quad (4.58)$$

Letting $t \rightarrow 1^-$ and $\tau \rightarrow 1^-$ in (4.58) and using (3.6) we deduce that

$$\begin{aligned} \underline{\lim}_{t \rightarrow 1^-} \overline{\lim}_{\rho \rightarrow 0} \frac{m_{tu_x}(Q_\rho(x))}{\mu(Q_\rho(x))} &\leq \left(1 + \underline{\lim}_{t \rightarrow 1^-} \overline{\lim}_{s \rightarrow 1^-} \Delta_L^a \left(\frac{t}{s}\right)\right) \left(\lim_{\rho \rightarrow 0} \frac{\check{m}_u(Q_\rho(x))}{\mu(Q_\rho(x))} + \varepsilon\right) \\ &\quad + \underline{\lim}_{t \rightarrow 1^-} \overline{\lim}_{s \rightarrow 1^-} \Delta_L^a \left(\frac{t}{s}\right) a(x). \end{aligned} \quad (4.59)$$

But, by diagonalization there exists a mapping $s \mapsto t_s$ with $t_s \rightarrow 1^-$ as $s \rightarrow 1^-$ such that:

$$\begin{aligned} \lim_{s \rightarrow 1^-} \frac{t_s}{s} &= 1; \\ \underline{\lim}_{t \rightarrow 1^-} \overline{\lim}_{s \rightarrow 1^-} \Delta_L^a \left(\frac{t}{s}\right) &\leq \overline{\lim}_{s \rightarrow 1^-} \Delta_L^a \left(\frac{t_s}{s}\right). \end{aligned}$$

But $\overline{\lim}_{r \rightarrow 1^-} \Delta_L^a(r) \leq 0$ because L is ru-usc, hence

$$\underline{\lim}_{t \rightarrow 1^-} \overline{\lim}_{s \rightarrow 1^-} \Delta_L^a \left(\frac{t}{s}\right) \leq 0,$$

and so from (4.59) we conclude that

$$\underline{\lim}_{t \rightarrow 1^-} \overline{\lim}_{\rho \rightarrow 0} \frac{m_{tu_x}(Q_\rho(x))}{\mu(Q_\rho(x))} \leq \lim_{\rho \rightarrow 0} \frac{\check{m}_u(Q_\rho(x))}{\mu(Q_\rho(x))} + \varepsilon,$$

and (4.44) follows by letting $\varepsilon \rightarrow 0$.

Substep 5-2: establishing the relaxation formula. By (4.44) and (4.45) we have

$$\lim_{\rho \rightarrow 0} \frac{\check{m}_u(Q_\rho(x))}{\mu(Q_\rho(x))} = \lim_{t \rightarrow 1^-} \overline{\lim}_{\rho \rightarrow 0} \frac{m_{tu_x}(Q_\rho(x))}{\mu(Q_\rho(x))}.$$

On the other hand, it is easily seen that

$$\overline{\lim}_{\rho \rightarrow 0} \frac{m_{tu_x}(Q_\rho(x))}{\mu(Q_\rho(x))} = \mathcal{Q}_\mu L(x, \nabla_\mu u(x)),$$

where $\mathcal{Q}_\mu L$ is the μ -quasiconvexification of L defined in (2.14). Hence

$$\lim_{\rho \rightarrow 0} \frac{\check{m}_u(Q_\rho(x))}{\mu(Q_\rho(x))} = \mathcal{Q}_\mu L(x, \nabla_\mu u(x)).$$

Step 6: end of the proof. From (2.12) we see that

$$\alpha \overline{\mathcal{G}}(u) \leq \overline{E}(u, X) \leq \beta(1 + \overline{\mathcal{G}}(u))$$

for all $u \in W_\mu^{1,p}(X; \mathbb{R}^m)$, where $\overline{\mathcal{G}}$ is defined (2.9). Hence, $\overline{E}(u, X) = \infty$ if $u \in W_\mu^{1,p}(X; \mathbb{R}^m) \setminus \overline{\mathfrak{S}}$, where $\overline{\mathfrak{S}}$ is the effective domain of $\overline{\mathcal{G}}$, and (2.15) follows because it is assumed that $\mathfrak{S} = \overline{\mathfrak{S}}$, where \mathfrak{S} denotes the effective domain of \mathcal{G} defined in (2.8). \square

Conflict of interest statement

There is no conflict of interest.

References

- [1] Omar Anza Hafs, On the integral representation of relaxed functionals with convex bounded constraints, *ESAIM Control Optim. Calc. Var.* 16 (1) (2010) 37–57.
- [2] Omar Anza Hafs, Jean-Philippe Mandallena, Interchange of infimum and integral, *Calc. Var. Partial Differ. Equ.* 18 (4) (2003) 433–449.
- [3] Omar Anza Hafs, Jean-Philippe Mandallena, Relaxation of second order geometric integrals and non-local effects, *J. Nonlinear Convex Anal.* 5 (3) (2004) 295–306.
- [4] Omar Anza Hafs, Jean-Philippe Mandallena, Relaxation of variational problems in two-dimensional nonlinear elasticity, *Ann. Mat. Pura Appl.* (4) 186 (1) (2007) 187–198.
- [5] Omar Anza Hafs, Jean-Philippe Mandallena, Relaxation theorems in nonlinear elasticity, *Ann. Inst. Henri Poincaré, Anal. Non Linéaire* 25 (1) (2008) 135–148.
- [6] Omar Anza Hafs, Jean-Philippe Mandallena, Homogenization of nonconvex integrals with convex growth, *J. Math. Pures Appl.* (9) 96 (2) (2011) 167–189.
- [7] Omar Anza Hafs, Jean-Philippe Mandallena, Homogenization of unbounded singular integrals in $W^{1,\infty}$, *Ric. Mat.* 61 (2) (2012) 185–217.
- [8] Omar Anza Hafs, Jean-Philippe Mandallena, On the relaxation of unbounded multiple integrals, arXiv:1207.2652 [math.AP], 2012.
- [9] Omar Anza Hafs, Jean-Philippe Mandallena, Radial representation of lower semicontinuous envelope, *Boll. Unione Mat. Ital.* 7 (1) (2014) 1–18.
- [10] Omar Anza Hafs, Jean-Philippe Mandallena, On the relaxation of variational integrals in metric Sobolev spaces, *Adv. Calc. Var.* 8 (1) (2015) 69–91.

- [11] Omar Anza Hafsa, Jean-Philippe Mandallena, Hamdi Zorgati, Homogenization of unbounded integrals with quasiconvex growth, *Ann. Mat. Pura Appl.* (4) 194 (6) (2015) 1619–1648.
- [12] Guy Bouchitté, Michel Belleueud, Regularization of a set function—application to integral representation, *Ric. Mat.* 49 (suppl.) (2000) 79–93, contributions in honor of the memory of Ennio De Giorgi (in Italian).
- [13] Anders Björn, Jana Björn, *Nonlinear Potential Theory on Metric Spaces*, EMS Tracts in Mathematics, vol. 17, European Mathematical Society (EMS), Zürich, 2011.
- [14] Guy Bouchitte, Giuseppe Buttazzo, Pierre Seppecher, Energies with respect to a measure and applications to low-dimensional structures, *Calc. Var. Partial Differ. Equ.* 5 (1) (1997) 37–54.
- [15] Guy Bouchitté, Irene Fonseca, Luisa Mascarenhas, A global method for relaxation, *Arch. Ration. Mech. Anal.* 145 (1) (1998) 51–98.
- [16] Jana Björn, L^q -differentials for weighted Sobolev spaces, *Mich. Math. J.* 47 (1) (2000) 151–161.
- [17] Giuseppe Buttazzo, *Semicontinuity, Relaxation and Integral Representation in the Calculus of Variations*, Pitman Research Notes in Mathematics Series, vol. 207, Longman Scientific & Technical, Harlow, 1989.
- [18] Sergio Conti, Georg Dolzmann, On the theory of relaxation in nonlinear elasticity with constraints on the determinant, *Arch. Ration. Mech. Anal.* 217 (2) (2015) 413–437.
- [19] J. Cheeger, Differentiability of Lipschitz functions on metric measure spaces, *Geom. Funct. Anal.* 9 (3) (1999) 428–517.
- [20] Gregory A. Chechkin, Vasili V. Jikov, Dag Lukkassen, Andrey L. Piatnitski, On homogenization of networks and junctions, *Asymptot. Anal.* 30 (1) (2002) 61–80.
- [21] Tobias H. Colding, William P. Minicozzi II, Liouville theorems for harmonic sections and applications, *Commun. Pure Appl. Math.* 51 (2) (1998) 113–138.
- [22] G.A. Chechkin, A.L. Piatnitski, A.S. Shamaev, *Homogenization. Methods and Applications*, Translations of Mathematical Monographs, vol. 234, American Mathematical Society, Providence, RI, 2007, translated from the 2007 Russian original by Tamara Rozhkovskaya.
- [23] Mitia Duerinckx, Antoine Gloria, Stochastic homogenization of nonconvex unbounded integral functionals with convex growth, *Arch. Ration. Mech. Anal.* 221 (3) (2016) 1511–1584.
- [24] E. De Giorgi, G. Letta, Une notion générale de convergence faible pour des fonctions croissantes d'ensemble, *Ann. Sc. Norm. Super. Pisa, Cl. Sci.* (4) 4 (1) (1977) 61–99.
- [25] Herbert Federer, *Geometric Measure Theory*, Die Grundlehren der mathematischen Wissenschaften, vol. 153, Springer-Verlag New York Inc., New York, 1969.
- [26] B. Franchi, P. Hajłasz, P. Koskela, Definitions of Sobolev classes on metric spaces, *Ann. Inst. Fourier (Grenoble)* 49 (6) (1999) 1903–1924.
- [27] Fragalà Ilaria, Lower semicontinuity of multiple μ -quasiconvex integrals, *ESAIM Control Optim. Calc. Var.* 9 (2003) 105–124 (electronic).
- [28] Jasun Gong, Piotr Hajłasz, Differentiability of p -harmonic functions on metric measure spaces, *Potential Anal.* 38 (1) (2013) 79–93.
- [29] Vladimir Gol'dshtein, Marc Troyanov, Axiomatic theory of Sobolev spaces, *Expo. Math.* 19 (4) (2001) 289–336.
- [30] Piotr Hajłasz, Sobolev spaces on metric-measure spaces, in: *Heat Kernels and Analysis on Manifolds, Graphs, and Metric Spaces*, Paris, 2002, in: *Contemp. Math.*, vol. 338, Amer. Math. Soc., Providence, RI, 2003, pp. 173–218.
- [31] Juha Heinonen, Nonsmooth calculus, *Bull., New Ser., Am. Math. Soc.* 44 (2) (2007) 163–232.
- [32] Piotr Hajłasz, Pekka Koskela, Sobolev meets Poincaré, *C. R. Acad. Sci. Paris Sér. I Math.* 320 (10) (1995) 1211–1215.
- [33] Juha Heinonen, Pekka Koskela, Quasiconformal maps in metric spaces with controlled geometry, *Acta Math.* 181 (1) (1998) 1–61.
- [34] Piotr Hajłasz, Pekka Koskela, Sobolev Met Poincaré, *Mem. Amer. Math. Soc.*, vol. 145(688), 2000, x+101.
- [35] Heikki Hakkarainen, Juha Kinnunen, Panu Lahti, Pekka Lehtelä, Relaxation and integral representation for functionals of linear growth on metric measure spaces, *Anal. Geom. Metr. Spaces* 4 (2016) 288–313.
- [36] Juha Heinonen, Pekka Koskela, Nageswari Shanmugalingam, Jeremy T. Tyson, *Sobolev Spaces on Metric Measure Spaces*, New Mathematical Monographs, vol. 27, Cambridge University Press, Cambridge, 2015, an approach based on upper gradients.
- [37] Omar Anza Hafsa, Jean Philippe Mandallena, Γ -limits of functionals determined by their infima, *J. Convex Anal.* 23 (1) (2016) 103–137.

- [38] Stephen Keith, A differentiable structure for metric measure spaces, *Adv. Math.* 183 (2) (2004) 271–315.
- [39] Jean-Philippe Mandallena, On the relaxation of nonconvex superficial integral functionals, *J. Math. Pures Appl.* (9) 79 (10) (2000) 1011–1028.
- [40] Jean-Philippe Mandallena, Quasiconvexification of geometric integrals, *Ann. Mat. Pura Appl.* (4) 184 (4) (2005) 473–493.
- [41] Jean-Philippe Mandallena, Localization principle and relaxation, *Adv. Calc. Var.* 6 (2) (2013) 217–246.
- [42] Jerrold E. Marsden, Thomas J.R. Hughes, *Mathematical Foundations of Elasticity*, Dover Publications, Inc., New York, 1994, corrected reprint of the 1983 original.
- [43] Marcelina Mocanu, Variational integrals in metric measure spaces, *Stud. Cercet. Ştiinţ. - Univ. Bacău, Ser. Mat.* (15) (2005) 67–89.
- [44] Jean-Philippe Mandallena, Mikhail Sychev, New classes of integral functionals for which the integral representation of lower semicontinuous envelopes is valid, *Dokl. Math.* 94 (1) (2016) 430–433.
- [45] Jean-Philippe Mandallena, Mikhail Sychev, New relaxation theorems with applications to strong materials, *Proc. R. Soc. Edinb., Sect. A* (2017), in press.
- [46] Nageswari Shanmugalingam, Newtonian spaces: an extension of Sobolev spaces to metric measure spaces, *Rev. Mat. Iberoam.* 16 (2) (2000) 243–279.
- [47] Sychev Mikhail, Attainment and relaxation results in special classes of deformations, *Calc. Var. Partial Differ. Equ.* 19 (2) (2004) 183–210.
- [48] Sychev Mikhail, Semicontinuity and relaxation theorems for integrands satisfying the fast growth condition, *Sib. Mat. Zh.* 46 (3) (2005) 679–697.
- [49] Sychev Mikhail, First general lower semicontinuity and relaxation results for strong materials, *J. Convex Anal.* 17 (1) (2010) 183–202.
- [50] V.V. Zhikov, Averaging of problems in the theory of elasticity on singular structures, *Izv. Ross. Akad. Nauk, Ser. Mat.* 66 (2) (2002) 81–148.