## **Effective Properties of Phononic Crystals in Bragg Regime**

N. Nemati<sup>1</sup>, C. Perrot<sup>1</sup>, D. Duhamel<sup>2</sup>, D. Lafarge<sup>3</sup>, Y. E. Lee<sup>4</sup>, N. Fang<sup>4</sup>

We present how the full account of temporal dispersion and spacial dispersion leads to the precise description of effective-medium parameters of a phononic crystal made of periodic arrangement of rigid inclusions embedded in a viscothermal fluid in high-frequency regime where Bragg scattering phenomena appear. We discuss the interplay between micro-geometry, frequency, fluid motions, and dissipative processes, and its impact on the emergence of macroscopic temporal and spacial dispersion effects. In this respect, we compare the local approach based on a two-scale asymptotic homogenization method, and a general nonlocal homogenization scheme.

<sup>&</sup>lt;sup>1</sup> Laboratoire Modélisation et Simulation Multi Echelle, Université Paris-Est UMR 8208, 77454 Marne-la-Vallée, France

<sup>&</sup>lt;sup>2</sup> Laboratoire Navier, Ecole des Ponts, UMR 8205, 77455 Marne-la-Vallée, France

<sup>&</sup>lt;sup>3</sup> Laboratoire d'Acoustique de l'Université du Maine, UMR 6613, Le Mans, France

<sup>&</sup>lt;sup>4</sup> Department of Mechanical Engineering, Massachusetts Institute of Technology, Cambridge, MA 02139, USA