



**HAL**  
open science

## Data set for transcriptome analysis of *Escherichia coli* exposed to nickel

Manon Gault, Agnès Rodrigue

► **To cite this version:**

Manon Gault, Agnès Rodrigue. Data set for transcriptome analysis of *Escherichia coli* exposed to nickel. *Data in Brief*, 2016, 9, pp.314 - 317. 10.1016/j.dib.2016.08.069 . hal-01667550

**HAL Id: hal-01667550**

**<https://hal.science/hal-01667550v1>**

Submitted on 14 Mar 2018

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



ELSEVIER

Contents lists available at ScienceDirect

## Data in Brief

journal homepage: [www.elsevier.com/locate/dib](http://www.elsevier.com/locate/dib)

## Data Article

Data set for transcriptome analysis of *Escherichia coli* exposed to nickel

Manon Gault, Agnès Rodrigue\*

Microbiologie, Adaptation et Pathogénie, UMR5240, INSA Lyon, Université Lyon 1, CNRS, Université de Lyon,  
F-69621 Villeurbanne, France

## ARTICLE INFO

## Article history:

Received 29 June 2016

Received in revised form

24 August 2016

Accepted 30 August 2016

Available online 7 September 2016

## Keywords:

RNA-Seq

*Escherichia coli*

nickel

## ABSTRACT

Ni is recognized as an element that is toxic to humans, acting as an allergen and a carcinogenic agent, and it is also toxic to plants. The toxicity of Ni has been understudied in microorganisms. The data presented here were obtained by submitting the model bacterium *Escherichia coli* K-12 to nickel stress. To identify expressed genes, RNA-Seq was performed. Bacteria were exposed to 50  $\mu$ M NiCl<sub>2</sub> during 10 min. Exposure to Ni lead to the deregulation of 57% of the *E. coli* transcripts. Further analysis using DAVID identified most affected biological pathways. The list of differentially expressed genes and physiological consequences of Ni stress are described in “Ni exposure impacts the pool of free Fe and modifies DNA supercoiling via metal-induced oxidative stress in *Escherichia coli* K-12” (M. Gault, G. Effantin, A. Rodrigue, 2016) [1].

© 2016 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY license

(<http://creativecommons.org/licenses/by/4.0/>).

## Specifications Table

Subject area	Biology
More specific subject area	Bioinformatics and microbiology
Type of data	Tables, figure

DOI of original article: <http://dx.doi.org/10.1016/j.freeradbiomed.2016.06.030>

\* Corresponding author.

E-mail address: [agnes.rodrigue@insa-lyon.fr](mailto:agnes.rodrigue@insa-lyon.fr) (A. Rodrigue).

<http://dx.doi.org/10.1016/j.dib.2016.08.069>

2352-3409/© 2016 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

How data was acquired	<i>High-throughput RNA-sequencing</i>
Data format	<i>Filtered and analyzed with statistical tests</i>
Experimental factors	<i>The bacteria were grown in minimal medium until early log-phase where 50 <math>\mu</math>M NiCl<sub>2</sub> was added, after 10 min bacteria were harvested and frozen</i>
Experimental features	<i>Total RNA was extracted using the frozen acid-phenol method. ARNr were excluded. Directional libraries were sequenced on Illumina Hiseq2500 in single reads.</i>
Data source location	<i>Laboratory "Microbiologie, Adaptation et Pathogénie", UMR5240, INSA Lyon, France</i>
Data accessibility	<i>Data are with this article and deposited in NCBI's Gene Expression Omnibus (GEO), accessible through GEO Series accession number GEO: GSE76167 <a href="http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE76167">http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE76167</a></i>

---

### Value of the data

---

- Ni, as many transition metals, is essential as a trace element to living organisms whereas it becomes toxic when present in excess. At present, the description of Ni toxicity in bacteria is under-studied although this metal is a widespread element bacteria are in contact with.
  - The data shows differentially expressed genes under Ni stress that could be compared to differentially expressed genes in other metal-stress conditions or other stress conditions.
  - Analysis of the biological pathways impacted when cells are exposed to Ni will help to understand the molecular mechanisms of Ni- or metal-stress.
  - Identification of Ni-deregulated genes could lead to biotechnological applications such as the design of whole cell biosensors.
- 


## 1. Data

The RNA-Seq and gene expression datasets were deposited in NCBI's Gene expression Omnibus [2], accessible through GEO series accession number GEO: GSE76176. Fig. 1 shows the distribution of deregulated genes in *E. coli* upon exposure to 50  $\mu$ M Ni. 2545 genes were deregulated considering a Fold-Change (FC) of 1.5, representing 57 % of the 4440 annotated transcripts of *E. coli* K-12 strain W3110. Gene Ontology was applied to classify differentially expressed genes according to their biological function (see Fig. 1 in [1]). GO Terms that were enriched in the list of differentially expressed genes were identified using the DAVID tools (Database for Annotation, Visualization and Integrated Discovery) [3,4]. Pathways that were significantly affected were mapped using KEGG and are listed in Table 1.

## 2. Experimental design, materials and methods

### 2.1. Strains and growth conditions

*E. coli* K-12 cells were grown at 37 °C in minimal medium supplemented with glucose until  $O_{D_{600\text{ nm}}} = 0.3$  and then treated with 50  $\mu$ M NiCl<sub>2</sub> during 10 min. These conditions lead to maximal expression of the Ni-stress marker gene *rcnA* (see Fig. S1A and S1B in [1]).



**Fig. 1.** *E. coli* transcriptome response to Ni. The diagrams show the number of genes whose synthesis was increased or decreased after metal exposure. 57 % of the 4440 *E. coli* transcripts were differentially expressed using expression ratio of 1.5 ( $p < 0.05$ ). Among these 57 %, 16 % had expression ratios superior to 4.5. The functional analysis was performed on these 16% that represent 196 down-regulated and 148 up-regulated genes.

## 2.2. RNA extraction and RNA-Seq

Three samples of each condition were treated, as described in [1].

## 2.3. Data analysis

Strand-orientated RNA-Seq was performed on Illumina Hiseq2500. Basecalls were performed using HCS 2.0.5 and RTA 1.17.20. Reads were aligned to whole reference genome *Escherichia coli* K-12 W3110 NC\_007779 using CASAVA v 1.8.2 software (Illumina). Gene expression was determined using Cufflinks v. 2.0.2. software. Differentially expressed genes were identified as described in [1].

**Table 1**  
Pathways most affected upon Ni stress.

Term	%	<i>p</i> -value
<b>Down-regulated genes</b>		
Ribosome	9	$5.9 \times 10^{-19}$
Purine metabolism	8.5	$1.3 \times 10^{-14}$
Flagellar assembly	5.5	$3.2 \times 10^{-9}$
Sulfur metabolism	3.5	$1.4 \times 10^{-8}$
Pyrimidine metabolism	5	$6 \times 10^{-8}$
ABC transporters	7	$9.1 \times 10^{-6}$
Biosynthesis of siderophore groups non ribosomal peptides	2	$6.1 \times 10^{-5}$
<b>Up-regulated genes</b>		
Two component system	5.5	$3.4 \times 10^{-7}$
Phenylalanine metabolism	2.7	$4.3 \times 10^{-5}$

Pathway analysis has been performed from the list of differentially expressed genes using the online tool DAVID and as per information from KEGG. The threshold was settled to  $\geq 4$  genes being involved in a given pathway.

% : involved genes/total of up- or down-regulated genes.

*p*-value : modified Fisher exact *p*-value.

#### 2.4. Identification of the affected pathways from the differentially expressed genes

Online tool DAVID (<http://david.abcc.ncifcrf.gov/>) [3,4] was used to find out the affected pathways among the differentially expressed gene lists. The gene lists were uploaded with selecting the background as all the genes of *E. coli*. Functional Annotation Chart was visualized using the *p*-value threshold of 0.01 and a minimum number of genes of 4. The information regarding the affected pathways was obtained from Kyoto Encyclopedia of Genes and Genomes (KEGG) within the analysis in DAVID, using the mentioned thresholds.

#### Acknowledgments

This work was supported by a BQR (BIOVIME) INSA Lyon. MG is the recipient of a doctoral fellowship from the French Ministry of Higher Education and Research.

#### Transparency document. Supplementary material

Transparency data associated with this article can be found in the online version at <http://dx.doi.org/10.1016/j.dib.2016.08.069>.

#### References

- [1] M. Gault, G. Effantin, A. Rodrigue, Ni exposure impacts the pool of free Fe and modifies DNA supercoiling via metal-induced oxidative stress in *Escherichia coli* K-12, *Free Radic. Biol. Med.* 97 (2016) 351–361.
- [2] R. Edgar, M. Domrachev, A.E. Lash, Gene expression omnibus: NCBI gene expression and hybridization array data repository, *Nucleic Acids Res.* 30 (2002) 207–210.
- [3] D.W. Huang, B.T. Sherman, R.A. Lempicki, Systematic and integrative analysis of large gene lists using DAVID bioinformatics resources, *Nat. Protoc.* 4 (2008) 44–57.
- [4] D.W. Huang, B.T. Sherman, R.A. Lempicki, Bioinformatics enrichment tools: paths toward the comprehensive functional analysis of large gene lists, *Nucleic Acids Res.* 37 (2009) 1–13.