



HAL
open science

Recent climate hiatus revealed dual control by temperature and drought on the stem growth of Mediterranean *Quercus ilex*

Morine Lempereur, Jean-Marc Limousin, Frédéric Guibal, Jean-Marc Ourcival, Serge Rambal, Julien Ruffault, Florent Mouillot

► To cite this version:

Morine Lempereur, Jean-Marc Limousin, Frédéric Guibal, Jean-Marc Ourcival, Serge Rambal, et al.. Recent climate hiatus revealed dual control by temperature and drought on the stem growth of Mediterranean *Quercus ilex*. *Global Change Biology*, 2017, 23 (1), pp.42-55. 10.1111/gcb.13495 . hal-01681557

HAL Id: hal-01681557

<https://hal.science/hal-01681557>

Submitted on 14 May 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 RECENT CLIMATE HIATUS REVEALED DUAL CONTROL BY TEMPERATURE AND
2 DROUGHT ON THE STEM GROWTH OF MEDITERRANEAN *QUERCUS ILEX*

3
4
5 **Running head:** STEM GROWTH AND CLIMATE HIATUS

6
7 Morine Lempereur^{1, 2*}, Jean-Marc Limousin¹, Frédéric Guibal³, Jean-Marc Ourcival¹, Serge
8 Rambal^{1, 4}, Julien Ruffault^{1, 5, 6}, and Florent Mouillot⁷

9
10 ¹ Centre d'Ecologie Fonctionnelle et Evolutive CEFE, UMR 5175, CNRS - Université de Montpellier -
11 Université Paul-Valéry Montpellier – EPHE, 1919 Route de Mende, 34293 Montpellier Cedex 5, France

12 ² Agence de l'Environnement et de la Maîtrise de l'Energie, 20, avenue du Grésillé- BP 90406, 49004
13 Angers Cedex 01, France

14 ³ Institut Méditerranéen de Biodiversité et d'Ecologie marine et continentale (IMBE), UMR 7263
15 CNRS, Aix-Marseille Université – IRD – Avignon Université, Europôle de l'Arbois, BP 8013545, Aix-
16 en-Provence Cedex 4, France

17 ⁴ Universidade Federal de Lavras, Departamento de Biologia, CP 3037, CEP 37200-000, Lavras, MG,
18 Brazil

19 ⁵ Irstea, UR RECOVER, 3275 route Cézanne, CS 40061, 13182 Aix-en-Provence cedex 5, France

20 ⁶ CEREGE UMR 7330, CNRS – Aix-Marseille Université, Europôle de l'Arbois, BP 8013545, Aix-en-
21 Provence Cedex 4, France

22 ⁷ Centre d'Ecologie Fonctionnelle et Evolutive CEFE, UMR 5175, CNRS - Université de Montpellier -
23 Université Paul-Valéry Montpellier – EPHE - IRD, 1919 Route de Mende, 34293 Montpellier Cedex 5,
24 France

25
26 *Author for correspondence: Morine Lempereur

27 Tel: +33(0)467613292

28 Email: morine.lempereur@cefe.cnrs.fr

29
30 **Keywords:** Basal area increment, climate change, climate-growth response, climate hiatus,
31 drought, growth duration, growth phenology, *Quercus ilex*

32
33 **Type of Paper:** Primary Research Article

34

35 **Abstract**

36 A better understanding of stem growth phenology and its climate drivers would improve
37 projections of the impact of climate change on forest productivity. Under a Mediterranean
38 climate, tree growth is primarily limited by soil water availability during summer, but cold
39 temperatures in winter also prevent tree growth in evergreen forests. In the widespread
40 Mediterranean evergreen tree species *Quercus ilex*, the duration of stem growth has been shown
41 to predict annual stem increment, and to be limited by winter temperatures on the one hand, and
42 by the summer drought onset on the other. We tested how these climatic controls of *Q. ilex*
43 growth varied with recent climate change by correlating a 40-year tree ring record and a 30-
44 year annual diameter inventory against winter temperature, spring precipitation, and simulated
45 growth duration. Our results showed that growth duration was the best predictor of annual tree
46 growth. We predicted that recent climate changes have resulted in earlier growth onset (-10
47 days) due to winter warming and earlier growth cessation (-26 days) due to earlier drought
48 onset. These climatic trends partly offset one another, as we observed no significant trend of
49 change in tree growth between 1968 and 2008. A moving-window correlation analysis revealed
50 that in the past, *Q. ilex* growth was only correlated with water availability, but that since the
51 2000s, growth suddenly became correlated with winter temperature in addition to spring
52 drought. This change in the climate-growth correlations matches the start of the recent
53 atmospheric warming pause also known as the ‘climate hiatus’. The duration of growth of *Q.*
54 *ilex* is thus shortened because winter warming has stopped compensating for increasing drought
55 in the last decade. Decoupled trends in precipitation and temperature, a neglected aspect of
56 climate change, might reduce forest productivity through phenological constraints and have
57 more consequences than climate warming alone.

58

59 **LIST OF ABBREVIATIONS USE IN PAPER**

60 **FS:** Long-term field survey of diameter at breast height (DBH), measured from 1986 to 2013.
61 **RW:** Ring-width series, measured from 1942 to 2008.
62 **AD:** Automatic dendrometer series, measured from 2004 to 2013.
63 **Puéchabon station:** weather station located in the study site since 1984.
64 **SML station:** weather station located in St Martin-de-Londres, 12 km away from the study site,
65 data available from 1966 to 2013 (Meteo France).
66 **T_{JFM}:** Mean of daily temperature from January to March (°C).
67 **P_{AMJ}:** Sum of precipitation from April to June (mm).
68 **WSI:** Water stress integral, a time-cumulated drought severity index (MPa day)
69 **to:** Day of year when stem growth starts

70 **t₁**: Day of year when stem growth stops in early summer
71 **Δt_{0-t1}**: Duration of the period between t₀ and t₁ computed for each set of simulated phenological
72 thresholds.
73 **BAI**: Basal area increment of the stems, expressed in mm² year⁻¹
74 **PET**: Potential evapotranspiration (mm)
75

76 INTRODUCTION

77 Stem growth, recorded in annual tree rings, is a synthetic surrogate of carbon input in
78 standing biomass (Babst *et al.*, 2014) and an index for tree vitality and fitness (Bigler &
79 Bugmann 2003; Benito-Garzon *et al.*, 2013). In return, it is the record of the yearly climatic
80 conditions that have enabled stem growth over a period lasting from decades to centuries (Fritts
81 1976; Briffa *et al.*, 2002). Understanding the factors that control stem growth is therefore crucial
82 to assess the impact of past and future climate change on forests, but remains a major scientific
83 issue due to a series of currently unresolved uncertainties.

84 First, the climate sensitivity of tree growth is generally considered to depend on
85 photosynthesis and respiration fluxes because most process-based models assume that tree
86 growth is carbon limited (Davi *et al.*, 2006; Gaucherel *et al.*, 2008; Keenan *et al.*, 2011). Carbon
87 allocation to sapwood is therefore the key process driving the carbon sink, and is at the heart of
88 tree growth simulations in most terrestrial biosphere models (e.g. Sitch *et al.*, 2003;
89 Friedlingstein *et al.*, 2006; Fisher *et al.*, 2010). Several allocation schemes exist (e.g. fixed
90 allocation, the pipe model, and hierarchical allocation between plant organs, Schippers *et al.*,
91 2015), but there is accumulating experimental evidence that cambial activity (sink limitation)
92 is more sensitive to environmental stressors than carbon assimilation (source limitation; Körner
93 2003; Fatichi *et al.*, 2014; Guillemot *et al.*, 2015; Körner 2015; Delpierre *et al.*, 2016a). Taking
94 sink limitation into account in terrestrial biosphere models is thus likely to improve their ability
95 to simulate past and future forest productivity (Leuzinger *et al.*, 2013; Fatichi *et al.*, 2014), but
96 a better characterization of the phenology and climate sensitivity of wood formation is a
97 prerequisite (Rossi *et al.*, 2011; Rossi *et al.*, 2014; Lempereur *et al.*, 2015; Delpierre *et al.*,
98 2016a; Delpierre *et al.*, 2016b).

99 Second, global climate change may simultaneously exert opposing influences on forest
100 functioning, and these impacts are difficult to disentangle. For example, increased water use
101 efficiency due to atmospheric CO₂ enrichment is sometime reported to increase tree growth in
102 spite of concurrently increasing aridity (Martinez-Vilalta *et al.*, 2008; Koutouvas 2013), but the
103 opposite finding is far more common (Peñuelas *et al.*, 2011). Warming also exerts opposing
104 impacts on tree growth: on one hand it increases the length of the growing season (Keeling *et al.*
105 *et al.*, 1996; Dragoni *et al.*, 2011) while on the other hand it exacerbates atmospheric evaporative
106 demand and hence water limitation (Angert *et al.*, 2005; Zhao & Running, 2010). Warming
107 may therefore affect tree growth positively (e.g. Rossi *et al.*, 2014) or negatively (e.g. Brzostek
108 *et al.*, 2014) depending on the ecosystem under consideration.

109 Third, relationships between tree growth and climate may change under the influence of
110 climate change (e.g. Briffa *et al.*, 1998; Büntgen *et al.*, 2006; Carrer & Urbinati 2006; D'Arrigo
111 *et al.*, 2008). A classic example of this phenomenon is the so-called 'divergence problem' in
112 northern forests, which is a weakening of the positive temperature response of tree growth in
113 strongly temperature limited ecosystems since the middle of the 20th century (D'Arrigo *et al.*,
114 2008). The consequence of such temporal variations in tree growth sensitivity to climate is that
115 empirical dendrochronological models based on statistical links between tree growth and
116 climate cannot be extrapolated to long term historical and future climate conditions with
117 confidence (e.g. Gea-Izquierdo *et al.*, 2013; Subedi & Sharma 2013).

118 Fourth, climate change itself is a complex phenomenon that may result in changes in the
119 mean climate (IPCC 2014), in the frequency of extreme climatic events (Meehl & Tebaldi 2004)
120 and in seasonality (Giorgi *et al.*, 2011; Ruffault *et al.*, 2013), all of which may have contrasted
121 impacts on tree growth and function. In addition, climate change is not monotonous and the
122 recent slowing down of the warming trend, early identified by Easterling & Wehner (2009) and
123 later called the 'climate hiatus' (Trenberth & Fasullo, 2013), is currently a topic of increasing
124 scientific interest. But if most scientific studies have focused on explaining the causes and
125 processes behind this 'climate hiatus' or pause in atmospheric warming (Meehl *et al.*, 2011;
126 Balmaseda *et al.*, 2013; England *et al.*, 2014; Steinman *et al.*, 2015), few have attempted to
127 assess its ecological consequences, especially for processes that are highly sensitive to climate,
128 such as tree growth.

129 In water limited ecosystems, where forest productivity is primarily limited by drought
130 (Churkina & Running 1998; Zhao & Running 2010), the supply of water from rainfall appears
131 to be the key variable to explain tree growth (Babst *et al.*, 2013). In the case of the
132 Mediterranean species *Quercus ilex*, the conclusions of dendrochronological studies converge
133 more specifically to identify winter and spring rainfall amounts as the key drivers of annual
134 stem growth (Gutiérrez *et al.*, 2011). In contrast, *Q. ilex* stem growth is generally poorly
135 correlated with winter and spring temperatures. Daily measurements of basal area increment
136 over an eight year period (2004-2011) with automatic dendrometers in a *Q. ilex* forest,
137 demonstrated that the duration of spring growth was an accurate predictor of annual stem
138 growth (Lempereur *et al.*, 2015). After observing that the duration of spring growth was
139 determined by winter temperature for growth start and a lower threshold of predawn plant water
140 potential of -1.1 MPa for growth cessation, Lempereur *et al.* (2015) proposed a new hypothesis
141 wherein the annual growth of *Q. ilex* is under the dual control of winter temperature and spring
142 water limitation.

143 In this contextual framework, we used both tree ring records over a 40-year period and
144 an annual diameter inventory over a 30-year period to (i) test whether the stem growth
145 phenology approach proposed by Lempereur *et al.* (2015) can explain *Q. ilex* growth over a
146 long retrospective period with different growth estimation methods, (ii) assess the relative
147 importance of drought and temperature in limiting *Q. ilex* growth under a Mediterranean
148 climate, and (iii) evaluate the stationarity of these limitations in recent decades. In particular,
149 we aimed to identify whether any temporal shift in the climate controls of *Q. ilex* growth
150 (drought or temperature plus drought) are concomitant to, and the functional consequence of,
151 the recent pause in atmospheric warming.

152

153 **MATERIALS AND METHODS**

154 *Site description*


155 The study site is located 35 km north-west of Montpellier (southern France), on a flat
156 plateau, in the Puéchabon state forest (43°44'29"N, 3°35'45"E, 270 m a.s.l.). This forest has
157 been managed as a coppice for centuries and the last clearcut took place in 1942. Vegetation is
158 largely dominated by a dense overstory of the evergreen oak *Quercus ilex*. In 2014, the average
159 top canopy height was 5.5 m and the stem density was 4700 (± 700) stems ha⁻¹. Understory
160 evergreen species, *Buxus sempervirens*, *Phyllirea latifolia*, *Pistacia lentiscus* and *Juniperus*
161 *oxycedrus*, compose a sparse shrubby layer (height < 2 m) with less than 25% cover. The
162 climate is Mediterranean, with 80% of the rainfall occurring between September and April.
163 Average annual precipitation for the 1984-2013 period was 916 mm (550 to 1549 mm). Mean
164 annual temperature over the same period was 13.2 °C with a minimum in January (5.5 °C) and
165 a maximum in July (22.9 °C). The very shallow bedrock has a hard Jurassic limestone origin.
166 The volumetric fraction of stones and rocks averages 0.75 in the top 0-50 cm and 0.90 below.
167 The stone free fine fraction of the soil in the 0-50 cm layer is a homogeneous silty clay loam
168 (USDA texture triangle) comprising 38.8% clay, 35.2% silt and 26% sand.

169

170 *Monitoring annual stem growth*

171 Stem diameter at breast height (DBH) was measured annually from 1986 to 2013 on 319
172 trees distributed in eight circular plots (diameter 20 m) within an area of 2 ha. The 319 trees
173 used in this field survey (hereafter FS, Fig. 1) were selected so as to include all the diameter

174 size classes in 1986. DBH was measured every winter under dry conditions using a diameter
 175 tape at a height identified by a paint mark on the stem. In January 2008, the stem diameters
 176 were distributed as follows: 58% < 8 cm DBH, 24% between 8 and 10 cm and 18% >10 cm.
 177 Because the smaller size classes exhibited a very low growth and a high mortality rate, we
 178 restricted our analysis to the 125 trees with a DBH > 8 cm in 2008 to obtain a stronger growth
 179 signal (Table 1). From 2004 to 2013, stem growth was measured more frequently and with
 180 greater accuracy using automatic band dendrometers on a subset of trees in two neighboring
 181 plots (for more details, see Lempereur *et al.*, 2015). Automatic dendrometers (ELPA-98,
 182 University of Oulu, Finland; hereafter AD) were set up at a height of 1.3 m above the ground
 183 on 6 to 12 trees with DBH > 7 cm. The two datasets (FS and AD) were strongly correlated
 184 during their overlapping period (Table 1), thus demonstrating the relative accuracy of FS in
 185 estimating the annual growth of the largest trees.
 186



187
 188 **Fig. 1** Chronological timeline of the main data sources used in this study. Stem growth series: automatic
 189 dendrometers (AD), field survey (FS) of stem diameter at breast height (DBH) and ring width series (RW).
 190 Climatic data: Puechabon meteorological station and St Martin-de-Londres (SML) meteorological station.
 191

192 *Dendrochronological procedures*

193 In 2005 and 2008, 15 and 12 stems, respectively, were selected among the largest size
 194 classes of the field survey (FS) sample (10 cm < DBH < 16 cm) and cut down to measure ring-
 195 width chronology. Cross sections were collected at stump height, air dried, sanded and polished
 196 (40 to 400 grit). The longest growth radii in each cross section were selected and compared
 197 pairwise under a binocular magnifier. In old coppiced oaks, rings are narrow and sometimes
 198 not clearly visible after 40 years of cambial age; consequently only 12 of the sampled stems
 199 could be cross dated with confidence, seven stems in 2005 and five stems in 2008. The first
 200 ring was formed in 1942 after the clearcut in winter 1941-1942, so data in the ring width series

201 (hereafter RW) were obtained from 1942 to 2008 (Table 1, Fig. 1). Cross dating was facilitated
 202 by the presence of frost rings corresponding to the severe winters of 1963, 1985 and 1987.
 203 These rings, typical for years with severe freezing episodes, were abnormally wide because the
 204 cells of their initial area were crushed and dislocated, a typical constraint on tree ring
 205 interpretation in woody species under a Mediterranean climate (Cherubini *et al.*, 2003). This
 206 particular feature made them useful as markers for cross dating but prevented the reliable
 207 measurement of annual stem growth, so these years were excluded from subsequent analyses.
 208 Cross correlation coefficients on annual tree ring indices calculated among the 12 individual
 209 series were all higher than 0.6 (P-value<0.05). The ring width measured at stump height and
 210 along the longest growth radius was then rescaled to the tree DBH at the time of the cut. This
 211 required correction for the tapering between stump height (20 cm) and breast height (130 cm)
 212 as well as correction for the bark thickness (see Fig. S1 for details). The average tree ring width
 213 after rescaling, which was used to calculate the basal area increment (BAI), was 954 μm
 214 (CV = 48%) between 1942 and 2008, and 792 μm (CV = 38%) over the period 1968-2008 used
 215 in our study.

216

217 **Table 1** Main characteristics of the three growth series used in the study, automatic dendrometer (AD), field survey
 218 (FS) and ring width (RW), over their respective complete temporal series and their overlap periods. Coefficients
 219 of correlation (r) between FS and AD over the period 2004-2013 and between RW and FS over the period 1987-
 220 2008 are given (P-value < 0.05: *; P-value < 0.001: ***).

Characteristics of the growth series

	Method	Period	Mean BAI (mm ²)	DBH 2008 (cm)	Sample size
AD	Auto Dendro	2004-2013	185.8(±70,00)	11.1 (±2.45)	12
FS	Diameter tape	1987-2013	154.4 (±44,47)	10.5 (±1.85)	125
RW	Ring width	1968-2008	204.6 (±55,39)	12.7 (±1.75)	12

Correlations on overlap periods

	Overlap period	Growth series	Mean BAI (mm ²)	r
AD vs. FS	2004-2013	AD	185.8(±70.00)	0.90 (***)
		FS	142.3 (±55.37)	
FS vs. RW	1987-2008	FS	158.2 (±44.68)	0.55 (*)
		RW	196.2(±45.00)	

221

222 *Climate variables*

223 The Puéchabon meteorological station (hereafter Puéchabon) is located in a clearing
 224 200 m away from the study plot and has provided daily on-site climate data since 1984 (Fig. 1).
 225 Precipitation was measured with a tipping bucket rain gauge (ARG100; Environmental
 226 Measurements, Sunderland, UK) calibrated to 0.2 mm per tip and placed 1 m above the ground,
 227 air temperature was recorded with a MP100 sensor (Rotronic, Bassersdorf, Switzerland) at a

228 height of 2 m, and net radiation was measured with a pyranometer (SKS1110; Skye Instruments,
229 UK) at a height of 2 m above the ground.

230 To extend our analysis before 1984, we used climate variables (daily rainfall, daily
231 minimum and maximum temperature) from the St Martin-de-Londres meteorological station
232 (hereafter SML; 43°47'06''N; 3°43'48''E, altitude 194 m a.s.l., located about 12 km away from
233 the study site; source Meteo-France) over the period 1966-2013 (Fig. 1). The mean daily
234 temperature was calculated as the average of the minimum and maximum daily temperature.
235 The climate data from SML station were compared to on-site measurements in the overlap
236 period 1984-2013 (Table S1 and Fig. S2). The comparison showed close agreement for
237 temperature but a higher rainfall amount in SML than in Puéchabon. No significant biases
238 (slope not different from 1, and distance to origin not different from 0) were observed between
239 the two meteorological stations in the two crucial variables related to stem growth: the mean
240 temperature from January to March (T_{JFM}) and the sum of precipitation from April to June
241 (P_{AMJ} ; Fig. S2).

242 Daily solar radiation at the SML station was calculated from the processing chain
243 described in Kumar *et al.* (1997). In the first step, theoretical clear sky solar radiation (R_a) was
244 calculated from the daily timing of sunrise and sunset, the bi-hourly sun azimuthal angle and
245 the atmospheric transmittance according to longitude using R Cran Packages 'RAtmosphere'
246 and 'oce'. No topographical effects were taken into consideration. Actual solar radiation (R_s , in
247 $\text{MJ m}^{-2} \text{ day}^{-1}$) was then calculated from maximum and minimum daily temperature (T_{\max} and
248 T_{\min} , respectively) and the clear sky theoretical radiation following the Hargreaves equation:
249 $R_s = k_{Rs} \sqrt{(T_{\max} - T_{\min})} R_a$, where the adjustment coefficient k_{Rs} was set to 0.16 for interior
250 location (Allen *et al.*, 1998). Results of the Hargreaves equation were validated against solar
251 radiation measured in Puéchabon since 1998.

252

253 ***Modeling predawn leaf water potential and water stress integral***

254 Soil water storage integrated over the rooting depth, c.a. 4.5 m, was measured during the
255 vegetative periods of 1984-1986 and from July 1998 to August 2009 at approximately monthly
256 intervals, using a neutron moisture gauge (503DR Hydroprobe, CPN, Concord, CA, USA).
257 Discrete measurements were interpolated at a daily time step using the soil water balance model
258 described by Rambal (1993) and further used by Rambal *et al.* (2014). The model was driven
259 by daily values of incoming solar radiation, minimum and maximum temperatures, and rainfall
260 amount. Potential evapotranspiration was computed using the Priestley-Taylor equation

261 (Priestley & Taylor, 1972). The reduced major axis (RMA) regression between neutron
262 moisture gauge measurements and model simulations yielded an R^2 of 0.93, the slope was
263 0.94 ± 0.05 ($P < 0.0001$, $n = 91$) and the intercept not significantly different from 0. Soil water
264 storage and soil water potential were linked by a Campbell-type retention curve (Campbell,
265 1985) whose parameters are strongly dependent on soil texture (Saxton *et al.*, 1986; Rambal *et*
266 *al.*, 2003). Predawn leaf water potential used for model validation was measured about eight
267 times a year between April and October from 2003 to 2009 on a subsample of four trees among
268 those equipped with automatic dendrometers (see Limousin *et al.*, 2012). RMA regressions
269 between measured and simulated values of predawn leaf water potential yielded an R^2 of 0.84,
270 the slope was 0.93 ± 0.05 ($P < 0.0001$, $n = 54$), and the intercept was not significantly different
271 from 0. We used the simulations of predawn water potential rather than soil water content, as
272 the former is more closely linked with plant functioning (Rambal *et al.*, 2003). The daily
273 simulations of predawn water potential were performed with the climate data from the
274 Puéchabon and SML stations for the periods 1984-2012 and 1966-2012, respectively. The water
275 stress integral (WSI), defined by Myers (1988) as the seasonally or yearly sum of predawn
276 water potential, was used as a drought severity index to quantify annual or seasonal water stress.
277

278 ***Duration of spring basal area increment: calculations of dates t_0 and t_1***

279 The dates of stem growth phenology that bounded the spring growth period, the DOY
280 (day of the year) when stem growth starts (hereafter t_0) and the DOY when stem growth stops
281 in early summer (hereafter t_1 , Fig. S3), were estimated for each year. The relationships between
282 t_0 or t_1 and climate variables were calibrated by Lempereur *et al.* (2015) on observations of t_0
283 and t_1 obtained with automatic dendrometers from 2004 to 2011. Briefly, t_0 was defined as the
284 first day at which basal area exceeded the culmination of the previous year, and t_1 as the first
285 day when BAI became null or negative (see Lempereur *et al.*, 2015 for methodological details).
286 t_0 and t_1 were estimated using climate data from both Puéchabon and the nearby SML station
287 over their corresponding timeframes. t_0 was predicted by a nonlinear relationship with the mean
288 temperature from January to March (T_{JFM}). The relationships were fitted between t_0 and T_{JFM}
289 measured in Puéchabon ($t_0 = 849.2 * \exp(-0.6436 T_{JFM}) + 121$; $R^2 = 0.95$; RMSE = 2.6 days;
290 Lempereur *et al.*, 2015) and between t_0 and T_{JFM} measured in SML ($t_0 = 4300 * \exp(-0.943$
291 $T_{JFM}) + 124.4$; $R^2 = 0.95$; RMSE = 2.3 days; Fig. S4). t_1 was predicted by the DOY when the
292 plant water potential simulated using climate data from Puéchabon reached a threshold of -
293 1.1 MPa ($R^2 = 0.75$; RMSE = 7 days, Lempereur *et al.*, 2015), corresponding to a DOY when

294 the plant water potential simulated using climate data from SML reached a threshold of -
295 $1.1 \text{ MPa} + \varepsilon$ (with $\varepsilon = -0.1$; $R^2 = 0.73$; $\text{RMSE} = 13.5$ days) due to the slight differences in
296 temperature and precipitation between the two stations (see Fig. S5). The duration $\Delta t_{t_0-t_1}$
297 corresponds to the duration of the period between t_0 and t_1 and was computed for each pair of
298 simulated phenological thresholds.

299

300 ***Data processing and statistical analyses***

301 Annual stem growth (DBH and ring widths) was converted into annual basal area
302 increments (BAI, expressed in $\text{mm}^2 \text{ year}^{-1}$) for the two data series. Analyses were performed
303 with averaged BAI values of 125 and 5-12 individual trees for FS and RW, respectively. The
304 year 1992 presented two periods when predawn plant water potential fell below the threshold
305 of -1.1 MPa , in spring and autumn (Fig. S6). This unusual bi-modal drought prevented us from
306 determining the duration $\Delta t_{t_0-t_1}$ so 1992 was excluded from subsequent analyses. The growth
307 series BAI_{FS} included measurements that appeared to be outliers compared to BAI_{RW} or BAI_{AD}
308 for the years 2002 and 2010 (see Fig. S7). In 2002, BAI_{FS} was particularly low when BAI_{RW}
309 was close to the expected value, and in 2010, BAI_{FS} was clearly higher than the BAI_{AD} value.
310 Because BAI_{FS} is less accurate than the other two methods, and was calculated as the difference
311 between two subsequent measurements of DBH, these two values were considered to be
312 unreliable and excluded from the analyses performed on BAI_{FS} .


313 To investigate the links between tree growth and climate, we computed the correlations
314 between our two different growth datasets (FS and RW) and a set of relevant climate predictors.
315 For each correlation tested, we reported the Pearson's correlation coefficient and its
316 significance based on the standard bootstrap method with 1000 samples taken from the original
317 distribution of climate and tree ring data. Within each sample, the number of observations was
318 random and followed a geometric distribution. We used the 'treeclim' R package built upon the
319 DENDROCLIM2002 statistical tool dedicated to tree ring analysis (Biondi & Waikul, 2004;
320 Zang & Biondi, 2015). Climate predictors were derived from temperature and precipitation at
321 monthly, seasonal and annual time scales, and from our functional index of spring growth
322 duration ($\Delta t_{t_0-t_1}$). To test for the stationarity of the drought and temperature controls over the
323 studied period, we also performed the same bootstrap sampling Pearson correlation analysis on
324 a 10-year moving window along the whole series.

325 The temporal trends in stem growth and climate variables were characterized using both
326 trend tests and breakpoint analyses. Temporal trends in time series were estimated using the

327 Theil-Sen test (Sen 1968) from the ‘openair’ R cran package, applied on a 1000 block bootstrap
328 simulations to account for auto-correlated variables (Kunsch 1989), with the block length set to
329 $n/3$, n being the length of the time series. Temporal breakpoints in the time series were assessed
330 by computing the yearly F-statistic (sequential F-test) and the *supF* statistic was then used to
331 test for their significance.
332

333 RESULTS

334 *Sensitivity of stem growth to the dual temperature-drought drivers*




335
336 **Fig. 2** Time series of annual stem basal area increment (BAI) for the ring width series (RW; 1942-2008), the long-
337 term field survey (FS; 1986-2013) and the automatic dendrometer series (AD; 2004-2013). Error bars are standard
338 errors among the sampled trees (see Table 1 for sample size).
339

340 Annual stem growth (BAI_{RW}) from 1942 to 2008 exhibited two distinct phases, with a
341 significant breakpoint around 1967 ($F_{\text{stat}} = 81.96$; $P\text{-value} < 0.0001$; Fig. 2). The first phase
342 (1942-1967) lasted 25 years after the clearcut and showed a linear increase of $5.5 (\pm 0.62) \text{ mm}^2$
343 y^{-1} in annual BAI. This phase, which corresponds approximately to the cut frequency formerly
344 used in the traditional management of this *Q. ilex* coppice (Floret *et al.*, 1992), was
345 characterized by incomplete canopy cover and notable self-thinning. The second phase (1968-
346 2008) showed a stabilization of BAI_{RW} around a mean value of $203.8 (\pm 54.84) \text{ mm}^2 \text{y}^{-1}$. The
347 influence of climate on stem growth was consequently only considered from 1968 on, to avoid
348 the confounding effects caused by changes in competition, space-filling and self-thinning
349 during the first phase.

350 The two long-term growth series BAI_{RW} and BAI_{FS} were significantly correlated with each
351 other during their overlap period 1986-2008 ($r = 0.55$, $P\text{-value} < 0.05$, Table 1). Both series of
352 BAI exhibited large between year variations (coefficient of variation, $\text{CV} = 27\%$ and 29% for

353 BAI_{RW} and BAI_{FS}, respectively). The minimum value of BAI was recorded in 2006 for both
 354 series: BAI_{RW} = 114 mm² y⁻¹ (SE = 36.5 mm² y⁻¹) and BAI_{FS} = 58 mm² y⁻¹ (SE = 13.2 mm² y⁻¹)
 355 ¹). The maximum BAI was recorded in 1977 for RW (BAI_{RW} = 348 mm² y⁻¹, SE = 22.1 mm² y⁻¹)
 356 ¹) and in 2001 for FS (BAI_{FS} = 246 mm² y⁻¹; SE = 20.8 mm² y⁻¹, vs. BAI_{RW} = 213 mm² y⁻¹;
 357 SE = 24 mm² y⁻¹.2).
 358



359
 360 **Fig. 3** Relationship between yearly basal area increment (BAI) and the duration of the spring growth period ($\Delta t_{t_0-t_1}$). Ring width (RW, dark grey circle) is shown for the period 1984-2008 and field survey (FS, light grey square)
 361 for the period 1986-2013. The linear relationships between $\Delta t_{t_0-t_1}$ and BAI_{RW} (dark grey line; BAI_{RW} = 1.34* $\Delta t_{t_0-t_1}$ + 122.9; R² = 0.56; P-value < 0.001) or BAI_{FS} (light grey line; BAI_{FS} = 1.09* $\Delta t_{t_0-t_1}$ + 95.5; R² = 0.35; P-
 362 value < 0.01) are represented. Error bars are standard errors.
 363
 364
 365

366 The DOY of the start (t_0) and stop (t_1) of the spring stem growth, simulated over the period
 367 1984-2013 using climate data from the Puéchabon station, occurred on average in mid-May
 368 (DOY 133, SD = 9.9 days) and early July (DOY 184, SD = 23.5 days), respectively. The growth
 369 duration $\Delta t_{t_0-t_1}$ varied considerably among years (CV = 50%) with values ranging from 2 days
 370 in 1995 to 95 days in 2008 (Fig. 3). BAI was linearly correlated with $\Delta t_{t_0-t_1}$ for both stem growth
 371 series (R² = 0.56; P-value < 0.0001 and R² = 0.35; P-value < 0.01 for RW and FS respectively;
 372 Fig. 3). Moreover, $\Delta t_{t_0-t_1}$ was the best explanatory variable for the inter-annual variations in
 373 BAI_{RW} compared to other climate variables over the period 1984-2008 (Table 2; 3). We thus
 374 concluded that the dual control of annual stem growth by temperature and precipitation, stated
 375 by Lempereur *et al.* (2015) for the period 2004-2011, remained valid for a longer retrospective
 376 period and when growth is measured with less accuracy than with automatic dendrometers. The
 377 intercepts of the linear relationships between BAI and $\Delta t_{t_0-t_1}$, which represent the residual
 378 autumnal growth (Lempereur *et al.*, 2015), differed significantly between the two BAI series
 379 (F = 17.97; P-value < 0.001), with 122.9 (\pm 34.42) mm² y⁻¹ and 95.5 (\pm 39.09) mm² y⁻¹ for RW
 380 and FS, respectively (Fig. 3). By contrast, the slopes of these relationships did not differ

381 significantly between RW and FS ($F = 0.46$; $P\text{-value} > 0.05$), and the common slope equals
 382 $1.23 \text{ mm}^2 \text{ day}^{-1}$. The sensitivity of BAI_{RW} to $\Delta t_{t_0-t_1}$ was thus identical to the sensitivity of BAI_{FS} .
 383 Furthermore, neither the extension of the study period to 1968-2008 nor the switch in
 384 meteorological stations from Puéchabon to SML, removed the significant correlation between
 385 BAI_{RW} and $\Delta t_{t_0-t_1}$ ($R^2 = 0.31$; $P\text{-value} < 0.001$; Table 2).


386
 387 **Table 2** Mean Pearson's correlation coefficients (and significance derived from a 1000 classical bootstrap
 388 sampling) between chronologies of annual stem growth (BAI) and the main phenology explanatory variables. The
 389 correlations are given using climate variables measured in Puéchabon field surveys (FS) over the period 1986-
 390 2013 and ring width (RW) over the period 1984-2008, and climate data from the SML meteorological station over
 391 the period 1968-2008 for RW (see Fig. 1). The explanatory variables tested were the start and stop growth dates
 392 (t_0 and t_1 , respectively) and the duration of stem growth ($\Delta t_{t_0-t_1}$). The coefficient of correlation (r) and the level of
 393 statistical significance (* $P\text{-value} < 0.05$; ** $P\text{-value} < 0.01$; *** $P\text{-value} < 0.001$) are given. Significant correlations
 394 are in bold.

	FS	RW	RW
Period	1986-2013	1984-2008	1968-2008
t_0	-0.43**	-0.51***	0.18
t_1	0.56***	0.68***	0.66***
$\Delta t_{t_0-t_1}$	0.60***	0.73***	0.56***

395
 396 ***Stem growth response to climate trends from 1968 to 2013***
 397 Over the longer period 1968-2008, $\Delta t_{t_0-t_1}$ was not a better predictor of BAI than the sum
 398 of spring precipitation (P_{AMJ}) or t_1 alone ($r = 0.75$; $P\text{-value} < 0.001$ and $r = 0.66$; $P\text{-value} < 0.001$
 399 respectively; Table 2, 3), and we also observed a loss of correlation between BAI and t_0
 400 ($r = 0.18$; $P\text{-value} = 0.31$; Table 2).

401 P_{AMJ} was the main explanatory climate variable for t_1 (over the period 1968-2008:
 402 $r = 0.80$; $P\text{-value} < 0.001$) and it exhibited no temporal trend between 1968 and 2013 ($P\text{-}$
 403 $\text{value} = 0.31$; Fig. 4b). However, spring water limitation increased (WSI in spring: -
 404 $0.48 \text{ MPa day } y^{-1}$; $P\text{-value} < 0.001$; Fig. 4d, Table S3) as a result of increasing spring potential
 405 evapotranspiration (PET; $+0.028 \text{ mm day}^{-1} y^{-1}$; $P\text{-value} < 0.001$; Fig. 4c, Table S3). This trend
 406 in PET was mainly due to the significant warming trend in spring temperatures throughout the
 407 period 1968-2013 ($+0.07^\circ\text{C } y^{-1}$; $P\text{-value} < 0.001$; Fig. 4a, Table S3). As a result, drought onset
 408 t_1 exhibited a temporal trend toward earlier dates ($-0.57 \text{ day } y^{-1}$; $P\text{-value} < 0.01$; Fig. 5). The
 409 correlation between BAI_{RW} and P_{AMJ} remained significant, however, throughout most of the
 410 1968-2008 period (Fig. 6a).

411



412 **Fig. 4** Temporal trends in (a) spring (AMJ) temperature (grey circles) and winter (JFM) temperature (light grey
 413 inverted triangles), (b) mean annual precipitation (dark grey triangles) and AMJ precipitation (grey circles), (c)
 414 mean annual potential evapotranspiration (ETP; dark grey triangles) and AMJ ETP (grey circles) and (d) annual
 415 water stress integral (WSI, dark grey triangles) and AMJ three months WSI (grey circles). The solid lines represent
 416 significant temporal trends (P -value <0.05). The dashed lines represent the extended trend for temperature after
 417 1998, or before 1978 for PET. The effect of the warming pause on the JFM temperature (panel a) is shown by the
 418 hatched area between the points and the regression line, the vertical dotted lines indicate the start of the winter
 419 warming pause in 2002.

420
 421 In parallel, T_{JFM} increased significantly throughout the period ($+0.04 \text{ } ^\circ\text{C y}^{-1}$; P -
 422 value <0.05 ; Fig. 4a) which led to a significant trend of t_0 toward earlier dates (-0.21 day y^{-1} ; P -
 423 value <0.05 ; Fig. 5). However, a pause in the atmospheric winter warming occurred in the last
 424 decade. This warming pause is apparent in the discrepancy between actual and expected T_{JFM}
 425 obtained from the temporal trend built over the period 1978-1998. The year 1978 was chosen
 426 because it corresponds to the onset of a rapid warming phase in the Northern Hemisphere (Mann
 427 *et al.*, 1999). Similarly, the year 1998 was considered as the onset of the climate hiatus
 428 observations (IPCC 2014). We used the period 1978-1998 to build the reference trend line and
 429 extrapolate expected T_{JFM} values over the 1999-2013 period. We then tested for a breakpoint in
 430 the anomalies between observed and expected T_{JFM} over the whole period. Such a breakpoint
 431 was identified in 2002, indicating a significant slowing down in winter warming after this date


432 (Fstat = 21.52; P<0.001; Fig. 4a). This temporal pattern was also observed at seven other
 433 meteorological stations in the region surrounding our study site (Table S4 and Fig. S8), thus
 434 confirming the regional occurrence of the globally observed climate hiatus (Trenberth &
 435 Fasullo, 2013). The date of growth onset (t_0) was very sensitive to the warming pause in T_{JFM}
 436 and a breakpoint in the t_0 trend was observed in 2002 (Fstat = 21.52; P<0.001; Fig. 5). We also
 437 observed that the start of the winter warming pause coincided with a change in the climate
 438 controls on annual stem growth. Indeed, the correlation between BAI and T_{JFM} , which was only
 439 significant over one 10-year window between 1968 and 1994, became constantly significant
 440 from the time window 1995-2004 on (Fig. 6b).

441
 442 **Table 3** Mean Pearson's correlation coefficients and significance (derived from a 1000 classical bootstrap
 443 sampling) between annual stem growth (BAI) or spring growth duration ($\Delta t_{t_0 t_1}$) and monthly or seasonal
 444 precipitation (P, in mm) and temperature (T, in °C) for RW data series. The climate data from Puéchabon and
 445 Saint Martin-de-Londres meteorological stations were used for the periods 1984-2013 and 1966-2013,
 446 respectively. The coefficients of correlation (r) and the level of statistical significance (*P-value < 0.05; **P-
 447 value < 0.01; ***P-value < 0.001) are given. Significant correlations are in bold.


	BAI				$\Delta t_{t_0 t_1}$			
	1984-2008		1968-2008		1984-2013		1966-2013	
	T	P	T	P	T	P	T	P
November (t-1)	0.06	-0.08	-0.23*	-0.03	0.07	0.24	-0.31*	-0.03
December (t-1)	0.30**	0.08	-0.07	-0.04	0.25	0.04	-0.06	-0.03
January	0.50***	-0.06	0.02	0.15	0.51***	0.03	0.34***	0.14
February	0.37*	-0.27*	-0.02	0.06	0.20	-0.16	0.22	0.02
March	0.34**	-0.17	-0.10	0.21	0.41***	-0.10	0.11	0.19*
April	0.05	0.23	0.00	0.33*	-0.11	0.48***	-0.07	0.24*
May	0.12	0.47***	-0.23	0.57***	0.08	0.37	-0.25*	0.48**
June	-0.46**	0.21	-0.65***	0.44***	-0.01	0.00	-0.24*	0.22
July	-0.56***	0.32**	-0.54***	0.36**	-0.26	-0.13	-0.42***	0.01
August	-0.08	-0.04	-0.30**	0.07	0.03	0.24*	-0.37***	0.08
September	-0.39*	-0.39*	-0.42***	-0.25*	-0.18	-0.35	-0.34***	-0.10
October	-0.13	-0.13	-0.14	-0.04	-0.18	-0.34**	-0.22	-0.15
Jan. Feb. March	0.48***	-0.20	-0.05	0.18	0.47*	-0.11	0.32**	0.15
Apr. May June	-0.15	0.53***	-0.38***	0.75***	-0.02	0.49*	-0.23	0.55***
June Aug. Sept.	-0.42**	-0.34	-0.51*	-0.14	-0.18	-0.30	-0.45***	-0.05
from Nov(t-1) to Oct	0.04	-0.11	-0.37*	0.25*	0.15	0.01	-0.21*	0.17

448
 449 The temporal trends in t_0 and t_1 , both toward earlier dates (-10 and -26 days,
 450 respectively) offset each other to some extent until 2002 and resulted in a non-significant trend
 451 in $\Delta t_{t_0 t_1}$. After 2002, the prolonged trend in t_1 continued to decrease while t_0 stabilized, thereby
 452 leading to a sudden shortening of the growing season (Fig. 5). The timing of this phenological
 453 reduction was concomitant with the appearance of a significant relationship between T_{JFM} and


454 BAI (Fig. 6b). The decline in $\Delta t_{t_0-t_1}$ induced by the winter warming pause resulted in a decrease
 455 in BAI_{RW} after 2002. This decreasing trend over time become significant after the 10-year time
 456 window 1997-2006 (Fig. 7a,b). BAI_{FS} exhibited a similar decreasing trend in the same period,
 457 but of lower amplitude and not significant.
 458



459 **Fig. 5** Onset of phenological stem growth (t_0 , pale grey inverted triangles) and end of stem growth (t_1 , grey circles)
 460 from 1966 to 2013, expressed as day of the year (DOY). The effect of the warming pause on t_0 is shown by the
 461 hatched area between the points and the regression line over the 1978-1998 period.
 462
 463



464
 465 **Fig. 6** Temporal variations in the bootstrapped Pearson's correlation coefficients between (a) BAI_{RW} and P_{AMJ} ,
 466 and (b) BAI_{RW} and T_{JFM} over 10-year moving windows. The black curve represents the correlation coefficient and
 467 the grey curves the upper and lower confidence intervals at a significance level $\alpha = 0.05$. Asterisks identify the
 468 time windows with significant correlations.
 469
 470



471 **Fig. 7** Temporal trend over 10-year moving windows of (a) mean annual stem growth from 1968 to 2008 and from
 472 1987 to 2013 for ring width (RW, dark grey circles) and field surveys (FS, grey squares), respectively. Error bars
 473 are standard errors. (b) Slopes of the linear trends of BAI for the two series. The asterisks indicate time windows
 474 with a significant trend ($p < 0.05$). The dotted vertical lines in both panels indicate the first time window with a
 475 significant relationship between BAI_{RW} and T_{JFM} after the warming pause.
 476

477 DISCUSSION

478 *Growth duration as a predictor of annual stem growth*

479 The duration of spring stem growth ($\Delta t_{t_0-t_1}$) was the best explanatory variable of yearly
 480 basal area increment for both BAI_{FS} and BAI_{RW} after 1984. This confirms earlier results
 481 obtained with automatic dendrometers over a shorter period and a smaller sample of trees
 482 (Lempereur *et al.*, 2015), and therefore validates $\Delta t_{t_0-t_1}$ as a robust estimator of *Q. ilex* annual
 483 stem growth over long time scales and large samples. The two independent samplings we used
 484 here differed markedly in the number of trees measured, the measurement method and its
 485 precision. FS corresponds to a large number of trees (125 trees) representing a broader tree size
 486 distribution but measured with a less precise resolution (0.3 mm of DBH), while RW was
 487 measured on a small sample of large dominant trees (12 trees) but with a better resolution

488 (0.01 mm of radius). Average stem growth was lower for BAI_{FS} than for BAI_{RW} , but both
489 responded linearly and with the same sensitivity to $\Delta t_{t_0-t_1}$ (Fig. 3).

490 The spring stem growth duration $\Delta t_{t_0-t_1}$ is defined as the length of the period between the
491 onset of spring stem growth (t_0) and the date of growth cessation caused by summer drought
492 (t_1). Lempereur *et al.* (2015) observed that t_0 was well predicted by winter temperature, and that
493 t_1 occurred when the leaf predawn water potential dropped below a threshold of -1.1 MPa. A
494 similar approach was used by Rossi *et al.* (2011, 2014) who linked the stem growth of boreal
495 black spruce with the duration of xylem growth delimited by cold temperatures in spring and
496 autumn, and by Delpierre *et al.* (2016a) who linked the growth of temperate oak to the timing
497 of the summer growth cessation controlled by water limitation. By identifying the climatic
498 limits for cell division and elongation, the temporal delimitation of tree growth duration is an
499 innovative way of improving simulation of forest carbon sink response to climate change
500 (Fatichi *et al.*, 2014). Tree growth has generally been simulated using two types of models:
501 statistical dendroclimatological models, which are empirical and may lead to marked
502 uncertainties when extrapolated to future climate scenarios (e.g. Gea-Izquierdo *et al.*, 2013;
503 Babst *et al.*, 2014); and process based models that link tree growth to carbon assimilation (e.g.
504 Sitch *et al.*, 2003; Friedlingstein *et al.*, 2006; Gaucherel *et al.*, 2008; Keenan *et al.*, 2011),
505 ignoring experimental evidence showing a direct, and generally more limiting, climate control
506 of plant growth (Körner, 2003; Muller *et al.*, 2011). Projecting tree growth based on growth
507 phenology as in Rossi *et al.* (2011) and Lempereur *et al.* (2015) merges the biological realism
508 of the first approach, i.e. a direct link between climate and growth, with the mechanistic
509 understanding of tree physiology used in process based models, and be a significant step
510 forward in projecting the response of forest ecosystems to climate change (Delpierre *et al.*,
511 2016b). Showing that growth duration is an adequate predictor of tree growth for long time
512 series datasets (40 years in our study) and coarse temporal resolution (annual diameter
513 measurements) is thus an important preliminary.

514 Nevertheless, the spring growth duration approach remains limited by both its conditions of
515 applicability and the large proportion of the growth variability not explained by $\Delta t_{t_0-t_1}$. The
516 particular case of the year 1992, with its bi-modal drought in spring and autumn and its well-
517 watered summer (Fig. S6), suggests that $\Delta t_{t_0-t_1}$ can only be used as a descriptor of stem growth
518 phenology in years with typical Mediterranean seasonality and a drought period occurring in
519 summer. While the lowest threshold of predawn water potential below which stem growth is
520 prevented certainly remains valid for any drought seasonality (Hsiao & Xu, 2000; Muller *et al.*,
521 2011; Lempereur *et al.*, 2015), we hypothesize that the relationship between stem growth and

522 the duration of periods with adequate growth conditions (the rate of growth) varies in years
523 with peculiar phenology. Variations in the rate of growth may also account for the unexplained
524 variability in the BAI against Δt_{0-t1} relationships (44% and 65% for RW and FS, respectively;
525 Fig. 3). Finally, the relative extent of autumn growth compared to spring growth would deserve
526 more detailed investigation. Lempereur *et al.* (2015) observed that autumn growth explained
527 only approximately 30% of annual stem growth and was strongly determined by spring
528 conditions, but its magnitude over long time periods and under climate change conditions
529 remains to be described.

530

531 ***Dual limitation of growth by winter cold and spring-summer drought***

532 Summers in the Mediterranean region are typically characterized by the concomitance of
533 low rainfall, high temperature and high solar radiation, while winters are cold and humid.
534 Consequently, vegetation functioning is limited by water deficit in summer and by low
535 temperatures in winter (Mitrakos, 1980; Terradas & Savé, 1992). A cessation of growth during
536 cold and dry periods is frequently observed in Mediterranean trees which display a bi-phasic
537 growth pattern over the year (Cherubini *et al.*, 2002; Campelo *et al.*, 2007; Montserrat-Marti *et al.*,
538 *et al.*, 2009; Camarero *et al.*, 2009, Gutiérrez *et al.*, 2011).

539 The sensitivity of growth to water deficit is well established in the literature (Lockhart
540 1965; Hsiao & Xu, 2000; Muller *et al.*, 2011), as is the negative impact of drought on ring width
541 (Fritts, 1976). Previous dendroclimatological studies generally stated that spring to early-
542 summer precipitation was the main driver of annual stem growth of *Q. ilex* (Gutiérrez *et al.*,
543 2011). More precisely, it is the start of the dry season, defined by a threshold of soil water
544 deficit that was identified as the main determinant of variations in inter-annual growth (Maselli
545 *et al.*, 2014; Lempereur *et al.*, 2015). The predominant effect of the timing of drought induced
546 growth cessation on annual stem growth is not limited to Mediterranean ecosystems, as a similar
547 effect has been observed in mesic temperate forests (Brzostek *et al.*, 2014; Delpierre *et al.*,
548 2016a). In our study, we defined the start of the dry season as a critical threshold of predawn
549 plant water potential of -1.1 MPa. This threshold is biologically meaningful because water
550 potential affects the cell turgor pressure necessary for cell growth (Hsiao & Xu, 2000), and it
551 has the other advantage of comprehensively accounting for climate and local soil conditions in
552 a single metric (Ruffault *et al.*, 2013). It is, however, a more complex index to calculate than
553 climatic indices of drought as it requires substantial knowledge of soil and vegetation
554 characteristics, and accurate meteorological data at the daily time scale. Consequently, the

555 uncertainty on t_1 was higher than the uncertainty on t_0 , especially when meteorological data
556 from SML (located at a distance of 12 km from the study site) was used instead of on-site data,
557 but t_1 nonetheless remained a good predictor of annual stem growth for whichever period (Table
558 2).

559 Depending on the species and on the local bio-climate, a minimum temperature
560 threshold ranging from +4 °C to +7 °C is required for stem growth to occur (Körner, 2003;
561 Rossi *et al.*, 2007; Deslauriers *et al.*, 2008; Gruber *et al.*, 2010; Swidrak *et al.*, 2011; Lempereur
562 *et al.*, 2015). The onset of stem growth can be assessed from winter temperature (Delpierre *et al.*
563 *et al.*, 2016b), thus the mean winter temperature directly impacts the duration of cambium activity
564 and wood formation (Rossi *et al.*, 2011). However, there is no consensus in
565 dendrochronological studies of the Mediterranean *Q. ilex* that winter temperature is a good
566 predictor of annual BAI, and both negative (Zhang & Romane 1990; Paton *et al.*, 2009; Gea-
567 Izquierdo *et al.*, 2009) and positive correlations (Campelo *et al.*, 2009; Nijland *et al.*, 2011;
568 Gea-Izquierdo *et al.*, 2011) between annual stem growth and winter temperature have been
569 reported. In our study, a positive correlation between growth and winter temperature was
570 observed over the period 1984-2008 ($r = 0.48$, $P < 0.001$; Table 3), in accordance with
571 Lempereur *et al.* (2015), but the relationship was not significant for the period 1968-2008, either
572 with t_0 or with T_{JFM} .

573 Taken together, these results suggest that summer drought is the main limiting factor for
574 *Q. ilex* growth, which is to be expected under the dry Mediterranean climate. Nevertheless, the
575 contrasted growth responses to winter temperature depending on the study concerned, or on the
576 period of interest, led us to investigate the impact of recent climate change on *Q. ilex* growth in
577 more detail.

578

579 ***Can warmer winter temperature compensate for earlier summer drought under climate*** 580 ***change?***

581 From 1968 to 2008, the annual and spring amounts of precipitation were stable (Fig. 4b),
582 but water limitation increased (Fig. 4d) due to increasing potential evapotranspiration with
583 rising temperature in spring and summer (Fig. 4a and c), in accordance with regional
584 observations (Ruffault *et al.*, 2013). Consequently, we simulated an earlier occurrence of
585 drought onset (t_1 , -26 days on average) along with an earlier growth onset, although to a lesser
586 extent (t_0 , -10 days; Fig. 5), which, taken together resulted in a non-significant decrease in
587 growth duration. The positive effect of warming on ecosystem functioning and tree growth

588 through a longer growing season has been widely observed in temperature limited boreal and
589 temperate forests (Keeling *et al.*, 1996; Menzel *et al.*, 2006; D'Arrigo *et al.*, 2008; Dragoni *et*
590 *al.*, 2011). However, in water limited regions, like the Mediterranean, warming is generally
591 considered to be an aggravating factor for drought, mainly because of increased evaporative
592 demand (Angert *et al.*, 2005; Zhao & Running, 2010; Park *et al.*, 2012). Consequently, climate
593 warming in the Mediterranean usually reduces tree growth (Jump *et al.*, 2006; Sarris *et al.*,
594 2007; Peñuelas *et al.*, 2008; Piovesan *et al.*, 2008; Martin-Benito *et al.*, 2010). Our results thus
595 mitigate this widely accepted conclusion and illustrate the peculiar parallel controls driven by
596 temperature, such that the benefit of an earlier stem growth is cancelled out by earlier drought
597 onset mostly caused by increasing evaporative demand.

598 The effect of temperature on t_1 , mediated by PET, may also explain why the correlation
599 with growth was better for t_1 than for P_{AMJ} over the period 1984-2008, when temperatures
600 increased significantly, but not over the longer period 1968-2008. Actually, spring PET, like
601 temperature, did not increase significantly until the early 1980s (Fig. 4a, c), suggesting that, in
602 the past, inter-annual variability in precipitation may have been a stronger driver of drought
603 onset. Alternatively, the lack of on-site precipitation measurements before 1984 may mean that
604 t_1 estimates based on SML are less closely linked to on-site conditions than three-month
605 cumulated precipitation. The correlation in precipitation amounts between Puéchabon and SML
606 actually increases with longer temporal resolutions.

607 Winter warming may also have a positive impact on stem growth by delaying growth
608 cessation in autumn, thereby partly compensating for earlier drought onset. However, the
609 phenology of autumn growth cessation appears to be less variable than that of t_0 (Lempereur *et*
610 *al.*, 2015), possibly because it is concurrently driven by photoperiod. Moreover, warm winters
611 may even impact *Q. ilex* growth negatively if the species requires winter chilling, as is the case
612 of the deciduous *Quercus* species (Fu *et al.*, 2015).

613

614 ***A Mediterranean “divergence problem”***

615 When looking at the temporal variations in the correlations between stem growth and
616 climate variables, we observed an abrupt and significant increase in the sensitivity of stem
617 growth to temperature in the early 2000s, while at the same time, its response to precipitation
618 weakened (Fig. 6a, b). Temporal changes in the response of tree growth to climate have been
619 observed in a wide range of climates and tree species in recent decades (e.g. Briffa *et al.*, 1998;
620 Büntgen *et al.*, 2006; Carrer & Urbinati 2006; Jump *et al.*, 2007; D'Arrigo *et al.*, 2008). The

621 ‘divergence problem’ in northern forests has been defined as the tendency for tree growth at
622 previously temperature limited sites to undergo a weakening of their temperature response
623 concurrent with an increasing sensitivity to drought (D’Arrigo *et al.*, 2008). Our results suggest
624 a Mediterranean ‘divergence problem’ according to which tree growth in water limited
625 Mediterranean ecosystems undergo a weakening of their response to spring-summer
626 precipitation and an increasing sensitivity to winter temperatures. Similar reports of temporal
627 changes from water driven to increasingly temperature driven tree growth in water limited
628 ecosystems have already been reported for beech forests in northeast Spain (Jump *et al.*, 2007),
629 black pine forests in Spain (Martin-Benito *et al.*, 2010), and Scots pine, European larch and
630 black pine in Switzerland (Feichtinger *et al.*, 2014). These observations differ from ours by
631 reporting an increase of the overall negative effect of warming on tree growth. Nevertheless,
632 they collectively point to the increasing influence of temperature on the growth of previously
633 water limited trees.

634

635 ***Concurrent increase in drought and warming: a keystone aspect of climate change revealed***
636 ***during the recent warming hiatus***

637 A keystone result of our study is the sudden significance of the growth-temperature
638 relationship occurring from the 1995-2004 time window on (Fig. 6b). In parallel, we observed
639 lower winter temperatures than would have been expected on the basis of continuous climate
640 warming (Fig. 4a). This regional pattern is therefore comparable to the globally observed
641 “warming hiatus” (Easterling & Wehner 2009; Trenberth & Fasullo 2013). Our results suggest
642 that these recent cooler winter temperatures resulted in later growth onset and led to an
643 increased temperature constraint on the duration of stem growth. In the meantime, the constraint
644 exerted by the water deficit increased constantly from 1968 onward with neither breakpoints
645 nor changes in the trend line (Fig. 4d, 5). As a result, the stem growth of *Q. ilex* was significantly
646 correlated only with water deficit in the past, but the shorter growth period after the 2000s
647 revealed the dual control by winter temperatures and spring-summer water deficit. The pause
648 in climate warming in turn disrupted the precarious balance between increasing winter
649 temperatures and increasing spring-summer water deficit, which temporarily sustained stem
650 growth until the end of the 1990s (Fig. 7). This is, to our knowledge, the first example of a
651 ‘divergence problem’ in the tree growth-climate relationship triggered not by a continuous
652 climate change but instead by the warming pause. This recent warming pause, a still
653 controversial aspect of climate change (Lewandowsky *et al.*, 2015; Wehner & Easterling 2015),

654 is mainly caused by the variations in Atlantic and Pacific multidecadal oscillations (Steinman
655 *et al.*, 2015), and produced globally heterogeneous patterns of breakpoint in the recent warming
656 trend (Ying *et al.*, 2015), with an enhanced effect on winter temperature in Eurasia (Li *et al.*,
657 2015). It is however, likely to be reversed in the coming decades (Steinman *et al.*, 2015).
658 Whether a future re-acceleration of climate warming in the Mediterranean would compensate
659 for increasing water deficit by phenological stimulation of earlier tree growth again, or on the
660 contrary, further exacerbate drought stress remains an open and important question. Studying
661 trends in targeted and concurrent climate variables such as temperature and drought should fully
662 capture the complexity of climate change impacts (Mazdiyani & Aghakouchak 2015). Climate
663 projections for the Mediterranean region forecast an increase in potential evapotranspiration
664 and a decrease in summer precipitation by the end of the 21th century (Gao & Giorgi 2008;
665 Ruffault *et al.*, 2014), which are likely to move the onset of drought forward more strongly than
666 the onset of spring growth (Lempereur *et al.*, 2015). Together with the current growth limiting
667 winter temperatures, these future trends in drought features could lead to a sharp reduction in
668 forest productivity and an increase in tree mortality in Mediterranean *Q. ilex* forests.

669

670 **ACKNOWLEDGEMENTS**

671 M. Lempereur benefited from a doctoral research grant provided by the French Environment
672 and Energy Management Agency (ADEME). Long term meteorological data were provided by
673 Météo France. The Puéchabon experimental site belongs to the SOERE F-ORE-T, which is
674 supported annually by Ecofor, Allenvi and the French national research infrastructure ANAEE-
675 F. The authors would like to thank Alain Rocheteau, Karim Piquemal and David Degueldre for
676 their technical assistance, and the two anonymous reviewers for their detailed and helpful
677 reviews of the manuscript.

678 **REFERENCES**

- 679 Allen RG, Pereira LS, Raes D, Smith M (1998) Crop evapotranspiration-Guidelines for
680 computing crop water requirements-FAO Irrigation and drainage paper 56. FAO, Rome,
681 **300**.
- 682 Angert A, Biraud S, Bonfils C, Henning CC, Buermann W, Pinzon J, Tucker CJ, Fung I (2005)
683 Drier summers cancel out the CO₂ uptake enhancement induced by warmer springs.
684 Proceedings of the National Academy of Sciences of the United States of America, **102**,
685 10823-10827.

- 686 Babst F, Bouriaud O, Papale D *et al.* (2014) Above-ground woody carbon sequestration
687 measured from tree rings is coherent with net ecosystem productivity at five eddy-
688 covariance sites. *New Phytologist* **201**, 1289-1303.
- 689 Babst F, Poulter B, Trouet V *et al.* (2013) Site- and species-specific responses of forest growth
690 to climate across the European continent. *Global Ecology and Biogeography*, **22**, 706-
691 717.
- 692 Balmaseda MA, Trenberth KE, Källén E (2013) Distinctive climate signals in reanalysis of
693 global ocean heat content. *Geophysical Research Letters*, **40**, 1754-1759.
- 694 Benito-Garzón M, Ruiz-Benito P, Zavala MA (2013) Interspecific differences in tree growth
695 and mortality responses to environmental drivers determine potential species
696 distributional limits in Iberian forests. *Global Ecology and Biogeography*, **22**, 1141-
697 1151.
- 698 Bigler C, Bugmann H (2003) Growth-dependent tree mortality models based on tree rings.
699 *Canadian Journal of Forest Research*, **33**, 210-221.
- 700 Biondi F, Waikul K (2004) DENDROCLIM2002: A C++ program for statistical calibration of
701 climate signals in tree-ring chronologies. *Computers & Geosciences*, **30**, 303-311.
- 702 Briffa KR, Osborn TJ, Schweingruber FH, Jones PD, Shiyatov SG, Vaganov EA (2002) Tree-
703 ring width and density data around the northern hemisphere: Part 1, local and regional
704 climate signals. *The Holocene*, **12**, 737-757.
- 705 Briffa KR, Schweingruber FH, Jones PD, Osborn TJ, Shiyatov SG, Vaganov EA (1998)
706 Reduced sensitivity of recent tree-growth to temperature at high northern latitudes.
707 *Nature*, **391**, 678-682.
- 708 Brzostek ER, Dragoni D, Schmid HP, Rahman AF, Sims D, Wayson CA, Johnson DJ, Phillips
709 RP (2014) Chronic water stress reduces tree growth and the carbon sink of deciduous
710 hardwood forests. *Global Change Biology*, **20**, 2531-2539.
- 711 Büntgen U, Frank DC, Nievergelt D, Esper J (2006) Summer temperature variations in the
712 European Alps, a.d. 755–2004. *Journal of Climate*, **19**, 5606-5623.
- 713 Camarero JJ, Olano JM, Parras A (2009) Plastic bimodal xylogenesis in conifers from
714 continental Mediterranean climates. *New Phytologist*, **185**, 471-480.
- 715 Campbell GS (1985) *Soil physics with basic: Transport models for soil-plant systems*, Elsevier.
- 716 Campelo F, Nabais C, Freitas H, Gutiérrez E (2007) Climatic significance of tree-ring width
717 and intra-annual density fluctuations in *Pinus pinea* from a dry Mediterranean area in
718 Portugal. *Annals of Forest Science*, **64**, 229-238.
- 719 Campelo FC, Nabais CN, García-González IG-G, Cherubini PC, Gutiérrez EG, Freitas HF
720 (2009) Dendrochronology of *Quercus ilex* L. and its potential use for climate
721 reconstruction in the Mediterranean region. *Canadian Journal of Forest Research*, **39**,
722 2486-2493.
- 723 Carrer M, Urbinati C (2006) Long-term change in the sensitivity of tree-ring growth to climate
724 forcing in *Larix decidua*. *New Phytologist*, **170**, 861-872.
- 725 Cherubini P, Gartner BL, Tognetti R, Bräker OU, Schoch W, Innes JL (2003) Identification,
726 measurement and interpretation of the tree rings in woody species from mediterranean
727 climates. *Biological Reviews*, **78**, 119-148.
- 728 Cherubini P, Fontana G, Rigling D, Dobbertin M, Brang P, Innes JL (2002) Tree-life history
729 prior to death: two fungal root pathogens affect tree-ring growth differently. *Journal of*
730 *Ecology*, **90**, 839-850.
- 731 Churkina G, Running SW (1998) Contrasting climatic controls on the estimated productivity
732 of global terrestrial biomes. *Ecosystems*, **1**, 206-215.
- 733 D'Arrigo R, Wilson R, Liepert B, Cherubini P (2008) On the 'divergence problem' in northern
734 forests: A review of the tree-ring evidence and possible causes. *Global and Planetary*
735 *Change*, **60**, 289-305.

736 Davi H, Dufrêne E, Francois C *et al.* (2006) Sensitivity of water and carbon fluxes to climate
737 changes from 1960 to 2100 in European forest ecosystems. *Agricultural and Forest*
738 *Meteorology*, **141**, 35-56.

739 Delpierre N, Berveiller D, Granda E, Dufrêne E (2016a) Wood phenology, not carbon input,
740 controls the interannual variability of wood growth in a temperate oak forest. *New*
741 *Phytologist*, **210**, 459-470. DOI: 10.1111/nph.13771

742 Delpierre N, Vitasse Y, Chuine I, Guillemot J, Bazot S, Rutishauser T, Rathgeber CK (2016b)
743 Temperate and boreal forest tree phenology: from organ-scale processes to terrestrial
744 ecosystem models. *Annals of Forest Science*, **73**, 5-25.

745 Deslauriers A, Rossi S, Anfodillo T, Saracino A (2008) Cambial phenology, wood formation
746 and temperature thresholds in two contrasting years at high altitude in southern Italy.
747 *Tree Physiology*, **28**, 863-871.

748 Dragoni D, Schmid HP, Wayson CA, Potter H, Grimmond CSB, Randolph JC (2011) Evidence
749 of increased net ecosystem productivity associated with a longer vegetated season in a
750 deciduous forest in south-central Indiana, USA. *Global Change Biology*, **17**, 886-897.

751 Easterling DR, Wehner MF (2009) Is the climate warming or cooling? *Geophysical Research*
752 *Letters*, **36**.

753 England MH, McGregor S, Spence P *et al.* (2014) Recent intensification of wind-driven
754 circulation in the Pacific and the ongoing warming hiatus. *Nature Climate Change*, **4**,
755 222-227.

756 Fatichi S, Leuzinger S, Körner C (2014) Moving beyond photosynthesis: from carbon source
757 to sink-driven vegetation modeling. *New Phytologist*, **201**, 1086-1095.

758 Feichtinger LM, Eilmann B, Buchmann N, Rigling A (2014) Growth adjustments of conifers
759 to drought and to century-long irrigation. *Forest Ecology and Management*, **334**, 96-
760 105.

761 Fisher R, Mcdowell N, Purves D *et al.* (2010) Assessing uncertainties in a second-generation
762 dynamic vegetation model caused by ecological scale limitations. *New Phytologist*, **187**,
763 666-681.

764 Floret C, Galan MJ, Le Floc'h E, Romane F (1992) Dynamics of holm oak (*Quercus ilex* L.)
765 coppices after clearcutting in southern France. *Vegetatio*, **99-100**, 97-105.

766 Friedlingstein P, Cox P, Betts R *et al.* (2006) Climate–Carbon Cycle Feedback Analysis:
767 Results from the C4MIP Model Intercomparison. *Journal of Climate*, **19**, 3337-3353.

768 Fritts HC (1976) *Tree rings and climate*, London, Academic press.

769 Fu YH, Zhao H, Piao S (2015) Declining global warming effects on the phenology of spring
770 leaf unfolding. *Nature*, **526**, 104-107.

771 Gao X, Giorgi F (2008) Increased aridity in the Mediterranean region under greenhouse gas
772 forcing estimated from high resolution simulations with a regional climate model.
773 *Global and Planetary Change*, **62**, 195-209.

774 Gaucherel C, Guiot J, Misson L (2008) Changes of the potential distribution area of French
775 Mediterranean forests under global warming. *Biogeosciences*, **5**, 1493-1504.

776 Gea-Izquierdo G, Cherubini P, Cañellas I (2011) Tree-rings reflect the impact of climate change
777 on *Quercus ilex* L. along a temperature gradient in Spain over the last 100years. *Forest*
778 *Ecology and Management*, **262**, 1807-1816.

779 Gea-Izquierdo G, Fernandez-De-Una L, Canellas I (2013) Growth projections reveal local
780 vulnerability of Mediterranean oaks with rising temperatures. *Forest Ecology and*
781 *Management*, **305**, 282-293.

782 Gea-Izquierdo G, Martín-Benito D, Cherubini P, Isabel C (2009) Climate-growth variability in
783 *Quercus ilex* L. west Iberian open woodlands of different stand density. *Annals of Forest*
784 *Science*, **66**, 802.

- 785 Giorgi F, Im ES, Coppola E, Diffenbaugh NS, Gao XJ, Mariotti L, and Shi Y (2011) Higher
786 Hydroclimatic Intensity with Global Warming. *Journal of Climate*, **24**, 5309–5324.
- 787 Gruber A, Strobl S, Veit B, Oberhuber W (2010) Impact of drought on the temporal dynamics
788 of wood formation in *Pinus sylvestris*. *Tree Physiology*, **30**, 490-501.
- 789 Guillemot J, Martin-Stpaul N, Dufrêne E, François C, Soudani K, Ourcival J, Delpierre N
790 (2015) The dynamic of annual carbon allocation to wood in European forests is
791 consistent with a combined source-sink limitation of growth: implications for
792 modelling. *Biogeosciences*, **12**, 2213-2255.
- 793 Gutiérrez E, Campelo F, Camarero JJ, Ribas M, Muntán E, Nabais C, Freitas H (2011) Climate
794 controls act at different scales on the seasonal pattern of *Quercus ilex* L. stem radial
795 increments in NE Spain. *Trees*, **25**, 637-646.
- 796 Hsiao TC, Xu LK (2000) Sensitivity of growth of roots versus leaves to water stress:
797 biophysical analysis and relation to water transport. *Journal of Experimental Botany*,
798 **51**, 1595-1616.
- 799 IPCC (2014) *Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part A: Global and*
800 *Sectoral Aspects. Contribution of Working Group II to the Fifth Assessment Report of*
801 *the Intergovernmental Panel on Climate Change.* (eds Field CB, Barros VR, Dokken
802 DJ, Mach KJ, Mastrandrea MD, Bilir TE, Chatterjee M, Ebi KL, Estrada YO, Genova
803 RC, Girma B, Kissel ES, Levy AN, MacCracken S, Mastrandrea PR, White LL).
804 Cambridge, United Kingdom, and New York, NY, USA, Cambridge University Press.
- 805 Jump AS, Hunt JM, Peñuelas J (2006) Rapid climate change-related growth decline at the
806 southern range edge of *Fagus sylvatica*. *Global Change Biology*, **12**, 2163-2174.
- 807 Jump AS, Hunt JM, Peñuelas J (2007) Climate relationships of growth and establishment across
808 the altitudinal range of *Fagus sylvatica* in the Montseny Mountains, northeast Spain.
809 *Ecoscience*, **14**, 507-518.
- 810 Keeling CD, Chin J, Whorf T (1996) Increased activity of northern vegetation inferred from
811 atmospheric CO₂ measurements. *Nature*, **382**, 146-149.
- 812 Keenan T, Maria Serra J, Lloret F, Ninyerola M, Sabate S (2011) Predicting the future of forests
813 in the Mediterranean under climate change, with niche-and process-based models: CO₂
814 matters! *Global Change Biology*, **17**, 565-579.
- 815 Körner C (2003) Carbon limitation in trees. *Journal of Ecology*, **91**, 4-17.
- 816 Körner C (2015) Paradigm shift in plant growth control. *Current Opinion in Plant Biology*, **25**,
817 107-114.
- 818 Koutavas A (2013) CO₂ fertilization and enhanced drought resistance in greek firs from
819 Cephalonia Island, Greece. *Global Change Biology*, **19**, 529-539.
- 820 Kumar L, Skidmore AK, Knowles E (1997) Modelling topographic variation in solar radiation
821 in a GIS environment. *International Journal of Geographical Information Science*, **11**,
822 475-497.
- 823 Kunsch H R (1989) The jackknife and the bootstrap for general stationary observations. *Annals*
824 *of Statistics*, **17**, 1217-1241.
- 825 Lempereur M, Martin-Stpaul NK, Damesin C, Joffre R, Ourcival J-M, Rocheteau A, Rambal S
826 (2015) Growth duration is a better predictor of stem increment than carbon supply in a
827 mediterranean oak forest: implications for assessing forest productivity under climate
828 change. *New Phytologist*, **207**, 579-590.
- 829 Leuzinger S, Hättenschwiler S (2013) Beyond global change: lessons from 25 years of CO₂
830 research. *Oecologia*, **171**, 639-651.
- 831 Lewandowsky S, Risbey JS, Oreskes N (2015) On the definition and identifiability of the alleged
832 “hiatus” in global warming. *Scientific Reports* **5**, 16784.
- 833 Li C, Stevens B, Marotzke J (2015) Eurasian winter cooling in the warming hiatus of 1998-
834 2012. *Geophysical Research Letters*, **42**, 8131-8139

- 835 Limousin JM, Rambal S, Ourcival JM, Rodriguez-Calcerrada J, Perez-Ramos IM, Rodriguez-
836 Cortina R, Misson L, Joffre R (2012) Morphological and phenological shoot plasticity
837 in a Mediterranean evergreen oak facing long-term increased drought. *Oecologia*, **169**,
838 565-577.
- 839 Lockhart JA (1965) An analysis of irreversible plant cell elongation. *Journal of Theoretical*
840 *Biology*, **8**, 264-275.
- 841 Mann ME, Bradley RS, Hughes MK (1999) Northern hemisphere temperatures during the past
842 millennium: Inferences, uncertainties, and limitations. *Geophysical Research Letters*,
843 **26**, 759-762.
- 844 Martín-Benito D, Del Río M, Heinrich I, Helle G, Cañellas I (2010) Response of climate-growth
845 relationships and water use efficiency to thinning in a *Pinus nigra* afforestation. *Forest*
846 *Ecology and Management*, **259**, 967-975.
- 847 Martínez-Vilalta J, Lopez BC, Adell N, Badiella L, Ninyerola M (2008) Twentieth century
848 increase of scots pine radial growth in NE Spain shows strong climate interactions.
849 *Global Change Biology*, **14**, 2868-2881.
- 850 Maselli F, Cherubini P, Chiesi M, Gilabert MA, Lombardi F, Moreno A, Teobaldelli M,
851 Tognetti R (2014) Start of the dry season as a main determinant of inter-annual
852 Mediterranean forest production variations. *Agricultural and Forest Meteorology*, **194**,
853 197-206.
- 854 Mazdiyasnı O and Aghakouchak A (2015) Substantial increase in concurrent drought and
855 heatwaves in the united states. *Proceedings of the National Academy of Sciences of the*
856 *United States of America*, **112**, 11484-11489
- 857 Meehl GA, Arblaster JM, Fasullo JT, Hu A, Trenberth KE (2011) Model-based evidence of
858 deep-ocean heat uptake during surface-temperature hiatus periods. *Nature Clim.*
859 *Change*, **1**, 360-364.
- 860 Meehl GA, Tebaldi C (2004) More intense, more frequent, and longer lasting heat waves in the
861 21st century. *science*, **305**, 994-997.
- 862 Menzel A, Sparks TH, Estrella N *et al.* (2006) European phenological response to climate
863 change matches the warming pattern. *Global Change Biology*, **12**, 1969-1976.
- 864 Mitrakos K (1980) Plant life under Mediterranean climatic conditions. *Port. Acta Biol. Ser. A*,
865 **16**, 33-44.
- 866 Montserrat-Martı G, Camarero JJ, Palacio S, Perez-Rontome C, Milla R, Albuixech J, Maestro
867 M (2009) Summer-drought constrains the phenology and growth of two coexisting
868 Mediterranean oaks with contrasting leaf habit: implications for their persistence and
869 reproduction. *Trees-Structure and Function*, **23**, 787-799.
- 870 Muller B, Pantin F, Génard M, Turc O, Freixes S, Piques M, Gibon Y (2011) Water deficits
871 uncouple growth from photosynthesis, increase C content, and modify the relationships
872 between C and growth in sink organs. *Journal of Experimental Botany*, **62**, 1715-1729.
- 873 Myers BJ (1988) Water stress integral—a link between short-term stress and long-term growth.
874 *Tree Physiology*, **4**, 315-323.
- 875 Nijland W, Jansma E, Addink EA, Delmas MD, De Jong SM (2011) Relating ring width of
876 Mediterranean evergreen species to seasonal and annual variations of precipitation and
877 temperature. *Biogeosciences*, **8**, 1141-1152.
- 878 Park Williams A, Allen CD, Macalady AK *et al.* (2012) Temperature as a potent driver of
879 regional forest drought stress and tree mortality. *Nature Clim. Change*, **3**, 292-297.
- 880 Patón D, Garcia-Herrera R, Guenca J, Galavis M, Roig FA (2009) Influence of climate on radial
881 growth of holm oaks (*Quercus ilex* subsp. *ballota* Desf) from SW Spain
882 *Geochronometria*, **34**, 49-56.

- 883 Peñuelas J, Canadell JG, Ogaya R (2011) Increased water-use efficiency during the 20th
884 century did not translate into enhanced tree growth. *Global Ecology and Biogeography*,
885 **20**, 597-608.
- 886 Peñuelas J, Hunt JM, Ogaya R, Jump AS (2008) Twentieth century changes of tree-ring $\delta^{13}\text{C}$
887 at the southern range-edge of *Fagus sylvatica*: increasing water-use efficiency does not
888 avoid the growth decline induced by warming at low altitudes. *Global Change Biology*,
889 **14**, 1076-1088.
- 890 Piovesan G, Biondi F, Filippo AD, Alessandrini A, Maugeri M (2008) Drought-driven growth
891 reduction in old beech (*Fagus sylvatica* L.) forests of the central Apennines, Italy.
892 *Global Change Biology*, **14**, 1265-1281.
- 893 Priestley CHB, Taylor RJ (1972). On the Assessment of Surface Heat Flux and Evaporation
894 Using Large-Scale Parameters. *Monthly Weather Review*, **100**, 81-92.
- 895 Rambal S (1993) The differential role of mechanisms for drought resistance in a Mediterranean
896 evergreen shrub: a simulation approach. *Plant, cell & environment*, **16**, 35-44.
- 897 Rambal S, Lempereur M, Limousin JM, Martin-Stpaul NK, Ourcival JM, Rodríguez-
898 Calcerrada J (2014) How drought severity constrains gross primary production(GPP)
899 and its partitioning among carbon pools in a *Quercus ilex* coppice? *Biogeosciences*, **11**,
900 6855-6869.
- 901 Rambal S, Ourcival JM, Joffre R, Mouillot F, Nouvellon Y, Reichstein M, Rocheteau A (2003)
902 Drought controls over conductance and assimilation of a Mediterranean evergreen
903 ecosystem: scaling from leaf to canopy. *Global Change Biology*, **9**, 1813-1824.
- 904 Rossi S, Deslauriers A, Anfodillo T, Carraro V (2007). Evidence of threshold temperatures for
905 xylogenesis in conifers at high altitudes. *Oecologia*, **152**, 1-12.
- 906 Rossi S, Girard M-J, Morin H (2014) Lengthening of the duration of xylogenesis engenders
907 disproportionate increases in xylem production. *Global Change Biology*, **20**, 2261-
908 2271.
- 909 Rossi S, Morin H, Deslauriers A, Plourde PY (2011) Predicting xylem phenology in black
910 spruce under climate warming. *Global Change Biology*, **17**, 614-625.
- 911 Ruffault J, Martin StPaul N, Rambal S, Mouillot F. (2013): Differential regional responses in
912 drought length, intensity and timing to recent climate changes in a Mediterranean
913 forested ecosystem. *Climatic Change*, **117**: 103-107.
- 914 Ruffault J, Martin-StPaul N, Duffet C, Goge F, Mouillot F (2014) Projecting future drought in
915 Mediterranean forests: bias correction of climate models matters! *Theoretical and*
916 *Applied Climatology*, **117**, 113-122.
- 917 Sarris D, Christodoulakis D, Körner C (2007) Recent decline in precipitation and tree growth
918 in the eastern Mediterranean. *Global Change Biology*, **13**, 1187-1200.
- 919 Saxton K, Rawls WJ, Romberger J, Papendick R (1986) Estimating generalized soil-water
920 characteristics from texture. *Soil Science Society of America Journal*, **50**, 1031-1036.
- 921 Schippers P, Sterck F, Vlam M, Zuidema PA (2015) Tree growth variation in the tropical forest:
922 understanding effects of temperature, rainfall and CO₂. *Global Change Biology*, **21**,
923 2749-2761. DOI: 10.1111/gcb.12877.
- 924 Sen PK (1968) Estimates of the Regression Coefficient Based on Kendall's Tau. *Journal of the*
925 *American Statistical Association*, **63**, 1379-1389.
- 926 Sitch S, Smith B, Prentice IC *et al.* (2003) Evaluation of ecosystem dynamics, plant geography
927 and terrestrial carbon cycling in the LPJ dynamic global vegetation model. *Global*
928 *Change Biology*, **9**, 161-185.
- 929 Steinman BA, Mann ME, Miller SK (2015) Atlantic and Pacific multidecadal oscillations and
930 Northern hemisphere temperatures. *Science*, **347**, 988-991.
- 931 Subedi N, Sharma M (2013) Climate-diameter growth relationships of black spruce and jack
932 pine trees in boreal Ontario, Canada. *Global Change Biology*, **19**, 505-516.

933 Swidrak I, Gruber A, Kofler W, Oberhuber W (2011) Effects of environmental conditions on
934 onset of xylem growth in *Pinus sylvestris* under drought. *Tree Physiology*, **31**, 483-493.
935 Terradas J, Savé R (1992) The influence of summer and winter stress and water relationships
936 on the distribution of *Quercus ilex* L. *Vegetatio*, **99/100**, 137-145.
937 Trenberth KE, Fasullo JT (2013) An apparent hiatus in global warming? *Earth's Future*, **1**, 19-
938 32.
939 Wehner MF, Easterling DR (2015) The global warming hiatus's irrelevance. *Science*, **350**,
940 1482-1483.
941 Ying LX, Shen ZH, Piao SL (2015) The recent hiatus in global warming of the land surface:
942 Scale dependent breakpoint occurrences in space and time. *Geophysical Research*
943 *Letters*, **42**, 6471-6478.
944 Zang C, Biondi F (2015) treeclim: an R package for the numerical calibration of proxy-climate
945 relationships. *Ecography*, **38**, 431-436.
946 Zhang S, Romane F (1990) Variations de la croissance radiale de *Quercus ilex* L en fonction
947 du climat. *Annals of Forest Science*, **48**, 225-234.
948 Zhao M, Running SW (2010) Drought-induced reduction in global terrestrial net primary
949 production from 2000 through 2009. *science*, **329**, 940-943.
950

951 **SUPPORTING INFORMATION**

952 **Fig. S1** Relationships used to rescale the ring width series to the tree DBH at the time of the
953 cut: tree tapering (base diameter minus DBH) against DBH, and bark thickness against DBH.

954 **Table S1** Summary of climate data for Puéchabon and St Martin-de-Londres meteorological
955 stations.

956 **Fig. S2** Relationships between the meteorological data of Puéchabon and St Martin-de-Londres
957 for the overlap period 1984-2013.

958 **Fig. S3** Characteristic example of annual stem growth showing the main phenological events
959 for stem growth.

960 **Fig. S4** Relationship between the spring onset of growth (t_0) and the January to March
961 temperature (T_{JFM}) for the meteorological station of St Martin-de-Londres.

962 **Fig. S5** Relationship between observed and predicted day of the year of growth cessation due
963 to drought (t_1) when simulating plant water potential with the meteorological data from St
964 Martin-de-Londres.

965 **Fig. S6** Seasonal variations of simulated predawn leaf water in Puéchabon in 1992.

966 **Fig. S7** Relationship between yearly basal area increment (BAI) and the duration of the spring
967 basal growth period (Δt_{0-1}) showing the years 2002 and 2010 that are outliers for the series
968 BAI_{FS} .

969 **Table S3** Results of the Theil-Sen tests for the trends of temperature, precipitation,
970 evapotranspiration and drought index.

971 **Fig. S8** Temporal trends in winter temperature for seven meteorological stations in the region
972 Languedoc-Roussillon.
973 **Table S4** Breakpoints and changes in mean winter temperature for seven meteorological
974 stations in the region Languedoc-Roussillon.