



## **Poecillastrosides, Steroidal Saponins from the Mediterranean Deep-Sea Sponge *Poecillastra compressa* (Bowerbank, 1866)**

Kevin Calabro, Elaheh Lotfi Kalahroodi, Daniel Rodrigues, Caridad Diaz, Mercedes de La Cruz, Bastien Cautain, Remi Laville, Fernando Reyes, Thierry Pérez, Bassam Soussi, et al.

### **► To cite this version:**

Kevin Calabro, Elaheh Lotfi Kalahroodi, Daniel Rodrigues, Caridad Diaz, Mercedes de La Cruz, et al.. Poecillastrosides, Steroidal Saponins from the Mediterranean Deep-Sea Sponge *Poecillastra compressa* (Bowerbank, 1866). *Marine drugs*, 2017, 15 (7), pp.1-12/199. <10.3390/md15070199>. <hal-01681632>

**HAL Id: hal-01681632**

**<https://hal.science/hal-01681632v1>**

Submitted on 19 Apr 2018

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

## Article

# Poecillastrosides, Steroidal Saponins from the Mediterranean Deep-Sea Sponge *Poecillastra compressa* (Bowerbank, 1866)

Kevin Calabro <sup>1,2</sup>, Elaheh Lotfi Kalahroodi <sup>3</sup>, Daniel Rodrigues <sup>3,4</sup>, Caridad Díaz <sup>5</sup>, Mercedes de la Cruz <sup>5</sup>, Bastien Cautain <sup>5</sup>, Rémi Laville <sup>2</sup>, Fernando Reyes <sup>5</sup>, Thierry Pérez <sup>4</sup>, Bassam Soussi <sup>3,6,7</sup> and Olivier P. Thomas <sup>1,3,\*</sup>

<sup>1</sup> School of Chemistry, National University of Ireland Galway, University Road, H91 TK33 Galway, Ireland; KEVIN.CALABRO@nuigalway.ie

<sup>2</sup> Cosmo International Ingredients, 855 avenue du Docteur Maurice Donat, 06250 Mougins, France; remi.laville@airliquide.com

<sup>3</sup> Géoazur, Université Côte d'Azur, CNRS, OCA, IRD, 250 rue Albert Einstein, 06560 Valbonne, France; elaheh.lotfi-kalahroodi@univ-rennes1.fr (E.L.K.); daniel4rodrigues@gmail.com (D.R.); bassam.soussi@gu.se (B.S.)

<sup>4</sup> Institut Méditerranéen de Biodiversité et d'Ecologie marine et continentale, CNRS—Aix-Marseille University, IRD—University Avignon, Station Marine d'Endoume, rue de la batterie des lions, 13007 Marseille, France; thierry.perez@imbe.fr

<sup>5</sup> Fundación MEDINA, Centro de Excelencia en Investigación de Medicamentos Innovadores en Andalucía, Avda. del Conocimiento 34, Parque Tecnológico de Ciencias de la Salud, E-18016 Armilla, Granada, Spain; caridad.diaz@medinaandalucia.es (C.D.); mercedes.delacruz@medinaandalucia.es (M.d.l.C.); bastien.cautain@medinaandalucia.es (B.C.); fernando.reyes@medinaandalucia.es (F.R.)

<sup>6</sup> Department of Marine Sciences, University of Gothenburg, P.O. Box 460, SE40530 Gothenburg, Sweden

<sup>7</sup> Oman Centre for Marine Biotechnology, P.O. Box 236, PC 103 Muscat, Oman

\* Correspondence: olivier.thomas@nuigalway.ie; Tel.: +353-(0)91-493-563

Received: 17 May 2017; Accepted: 21 June 2017; Published: 26 June 2017

**Abstract:** The first chemical investigation of the Mediterranean deep-sea sponge *Poecillastra compressa* (Bowerbank, 1866) led to the identification of seven new steroidal saponins named poecillastrosides A–G (1–7). All saponins feature an oxidized methyl at C-18 into a primary alcohol or a carboxylic acid. While poecillastrosides A–D (1–4) all contain an *exo* double bond at C-24 of the side-chain and two osidic residues connected at O-2', poecillastrosides E–G (5–7) are characterized by a cyclopropane on the side-chain and a connection at O-3' between both sugar units. The chemical structures were elucidated through extensive spectroscopic analysis (High-Resolution Mass Spectrometry (HRESIMS), 1D and 2D NMR) and the absolute configurations of the sugar residues were assigned after acidic hydrolysis and cysteine derivatization followed by LC-HRMS analyses. Poecillastrosides D and E, bearing a carboxylic acid at C-18, were shown to exhibit antifungal activity against *Aspergillus fumigatus*.

**Keywords:** sponge; saponins; deep-sea; *Poecillastra compressa*

## 1. Introduction

In the marine environment, steroid and triterpenoid glycosides are widespread metabolites mainly produced by echinoderms [1–3], although saponins have also been isolated from other marine invertebrates such as octocorals or sponges [4,5]. To date, about 70 saponins have been reported from sponges [6] including sarasinosides from *Asteropus* spp. [7,8], *Melophlus* spp. [9,10], and *Lipastrotethya* sp. [11], ulososides from *Ulosa* sp. [12,13] and *Ectoplyasia ferox* [14], pandarosides

and acanthifoliosides from *Pandaros acanthifolium* [15–18], wondosterols from the association of two sponges [19], erylosides, sokodosides, nobiloside, and formosides from *Erylus* spp. [20–29], ptilosaponosides from *Ptilocaulis spiculifer* [30], mycalosides from *Mycale laxissima* [31,32], feroxosides from *Ectyoplasia ferox* [33], and silenosides from *Silene vulgaris* [34]. While some sponge saponins can be oxidized on the D ring or can contain unusual side chains, the aglycone of most of them belongs to the 30-norlanostane triterpenoid family, with steroidal saponins being rather rare for sponges. Some sponge saponins were subjected to different bioassays and they usually demonstrated interesting biological activities, mostly cytotoxicity against tumor cell lines [35–37].

In our continuous efforts to describe the chemical diversity of marine sponges from the Mediterranean, we undertook the first chemical study of the deep-sea Tetractinellid sponge *Poecillastra compressa* (Bowerbank, 1866). The genus *Poecillastra* is known to produce a broad range of secondary metabolites such as macrolactams [38,39], nitrosohydroxyalkylamines [40], sesquiterpenes, and steroids [41,42]. We report herein the isolation and structure elucidation of seven new steroidal glycosides named poecillastrosides A–G (1–7) from the deep-sea sponge *P. compressa* (Figure 1). Their structures were deduced from spectroscopic data including 1D- and 2D-NMR experiments as well as high-resolution mass spectra (HRESIMS) analyses. Three different aglycone moieties were identified, and oxidation at the C-18 position is a common feature among all isolated saponins. Poecillastroside A (1) contains an ergostane aglycone, whereas poecillastrosides B–D (2–4) contain a poriferastane, and poecillastrosides E–G (5–7) a cholestane with a cyclopropyl ring on the side-chain.




Figure 1. Structure of poecillastrosides A–G (1–7).

## 2. Results and Discussion

The freeze-dried sponge sample (43.1 g) was macerated and repeatedly extracted with a mixture of CH<sub>2</sub>Cl<sub>2</sub>/CH<sub>3</sub>OH (1:1) under sonication. The extract (7.9 g) was fractionated by Reversed Phase C18 Vacuum Liquid Chromatography with solvent mixtures of decreasing polarity. The methanolic fraction was then purified by successive RP-Phenylhexyl and C18 HPLC yielding pure compounds 1–7.

Compound 1 was isolated as a yellowish amorphous solid. Its molecular formula C<sub>40</sub>H<sub>68</sub>O<sub>13</sub> was determined by HRESIMS. The <sup>1</sup>H NMR spectrum of 1 suggested a steroidal saponin (Table 1). First, the characteristic anomeric signals at δ<sub>H</sub> 4.49 (d, *J* = 7.6 Hz, 1H, H-1'), 4.56 (d, *J* = 7.9 Hz, 1H, H-1''), and δ<sub>C</sub> 101.8 (C-1'), 105.2 (C-1'') evidenced the presence of two sugar residues. The <sup>1</sup>H NMR data of the steroid revealed one methyl singlet at δ<sub>H</sub> 0.88 (s, 3H, H<sub>3</sub>-19), three methyl doublets at δ<sub>H</sub> 1.02 (d, *J* = 6.8 Hz, 3H, H<sub>3</sub>-21) and 1.03 (d, *J* = 6.8 Hz, 6H, H<sub>3</sub>-26 and -27), ten methylene groups, an oxygenated methylene with the AB system at δ<sub>H</sub> 3.95 and 3.59, a 1,1-disubstituted olefin at δ<sub>H</sub> 4.70 and 4.71 (H<sub>2</sub>-24<sup>1</sup>), seven methine groups, two oxygenated methines at δ<sub>H</sub> 3.72 (m, 1H, H-3), 4.26 (td, *J* = 7.7, 3.7 Hz, 1H, H-16), and three

quaternary carbons at C-10, C-13 and C-24. When compared to usual steroids, this aglycone lacks one characteristic methyl signal for C-18. A hydroxylation was proposed at this position based on the presence of an AB system at  $\delta_{\text{H}}$  3.59 (d,  $J = 11.5$  Hz, 1H, H-18b) and 3.95 (d,  $J = 11.5$  Hz, 1H, H-18a) and further key H-12b, H-14, H-17/C-18, and H<sub>2</sub>-18/C-13, C-14, C-17 HMBC correlations. Another unusual feature for the steroid moiety was evidenced in the HSQC spectrum with signals of an oxygenated methine at  $\delta_{\text{H}}$  4.26 (td,  $J = 7.7, 3.7$  Hz, 1H, H-16) and  $\delta_{\text{C}}$  72.8 (CH, C-16). The location of this hydroxyl group at C-16 was confirmed after interpretation of key H-16/H-17 and H-16/H-15a COSY and TOCSY correlations. While most of the relative configurations were in accordance with a common steroid core, the relative configuration at C-16 was established after examination of the NOESY spectrum. Absence of clear nuclear Overhauser effect (nOe) between H-16 and H-14 but also H-18 together with some overlap between H-17 and H-22 did not allow a straightforward determination of the relative configuration at this position. However, H-16/H-15a and H-8/H-15b nOes suggested a  $\beta$  orientation for the hydroxyl group at C-16. As a confirmation of this orientation, the coupling constant values of H-16 were in perfect accordance with those observed for the same signal of a closely related analogue weinbergsterol B, isolated from the sponge *Petrosia weinbergi* [43]. NMR signals of the sugar residues were assigned by extensive COSY, TOCSY, and HSQC interpretation. HMBC experiment evidenced H-5'/C-1', H-1''/C-2', H-5''/C-1'' long-range correlations, thus revealing the pyranose nature of these two sugars and their connection at C-2'. Finally, the connectivity of the sugar with the aglycone at C-3 was confirmed through the key HMBC H-1'/C-3 correlation. Moving to the relative configuration of the residues, the large coupling constants between H-1'/H-2' and H-1''/H-2'' (7.9 and 7.6 Hz, respectively) were consistent with a  $\beta$  configuration for both anomeric centers. This interpretation was confirmed with the one-bond coupling constant  $^1J_{\text{CH}} \approx 160$  Hz for the two anomeric positions [44]. In addition, the coupling constant values of  $^3J_{\text{H3'-H4'}}$  3.2 Hz and  $^3J_{\text{H5'-H4'}}$  close to zero suggested an axial position for the hydroxyl at C-4 and, therefore, a  $\beta$ -galactopyranosyl residue attached at C-3 of the aglycone [45]. For the second sugar residue, all coupling constants were measured with values between 7 and 9 Hz which implies equatorial positions for all oxygen atoms and, therefore, a  $\beta$ -glucopyranosyl residue connected at C-2' of the first residue.

Assuming a usual absolute configuration for the aglycone, we turned towards the pyranose moieties. After hydrolysis of the acetal bonds, the resulting monosaccharides were derivatized with L-cysteine methyl ester and phenylisothiocyanate in pyridine [46]. By comparison with standards, a D absolute configuration was assigned for both glucose and galactose monosaccharides.

Compound **2** was isolated as a yellowish amorphous solid. The molecular formula of **2** was determined by HRESIMS as C<sub>41</sub>H<sub>70</sub>O<sub>13</sub>. The spectroscopic data were very similar to those of **1**, thereby suggesting that both compounds were close analogues. Examination of the <sup>1</sup>H NMR spectrum revealed the presence of an additional methyl group at  $\delta_{\text{H}}$  1.59 (d,  $J = 6.3$  Hz, 3H, H<sub>3</sub>-24<sup>2</sup>) placed on the double bond at C-24<sup>1</sup>, therefore, leading to a poriferastane skeleton. The relative configuration of **2** was found to be the same as that of poecillastroside A based on nOe correlations. A key H<sub>3</sub>-24<sup>2</sup>/H<sub>2</sub>-23 nOe led us to assign the configuration of the double bond as *E*.

Compound **3** was isolated as a pale yellowish amorphous solid with the same molecular formula C<sub>41</sub>H<sub>70</sub>O<sub>13</sub>. Both compounds **2** and **3** are, therefore, isomers. The <sup>1</sup>H NMR spectra were almost identical except for a deshielding of the signal corresponding to H-25, from  $\delta_{\text{H}}$  2.24 in **2** to  $\delta_{\text{H}}$  2.85 for **3**. We first supposed that a change in the configuration of the double had occurred. Due to the low amount of compound available, the corresponding carbons were not visible neither in the <sup>13</sup>C NMR spectrum nor in the HSQC, HMBC spectra. We, therefore, decided to enhance the sensitivity of the HSQC spectrum using the recently developed Pure Shift HSQC experiment [47]. Gratifyingly, we were then able to observe both HSQC spots corresponding to C-24<sup>1</sup> and C-25 (Figure S24). The shielding of the C-25 signal from  $\delta_{\text{C}}$  36.0 for **2** to  $\delta_{\text{C}}$  29.8 for **3** clearly confirmed a *Z* configuration for the double bond of **3**.


**Table 1.** NMR spectroscopic data for poecillastrosides A–D (1–4) in CD<sub>3</sub>OD (500 MHz for <sup>1</sup>H NMR data and 125 MHz for <sup>13</sup>C NMR data).

No.	1		2		3		4	
	$\delta_H$ , mult. (J in Hz)	$\delta_C$	$\delta_H$ , mult. (J in Hz)	$\delta_C$	$\delta_H$ , mult. (J in Hz)	$\delta_C$	$\delta_H$ , mult. (J in Hz)	$\delta_C$
1	1.70, m	38.1	1.69, m	38.1	1.69, m	38.1	1.69, m	38.2
	0.98, m		0.98, m		0.98, m		0.98, m	
2	1.90, m	30.5	1.90, m	30.5	1.90, m	30.5	1.92, m	30.5
	1.50, m		1.50, m		1.50, m		1.48, m	
3	3.72, m	80.2	3.72, m	80.2	3.72, m	80.2	3.72, m	80.3
	1.71, m		1.71, m		1.71, m		1.70, m	
4	1.34, m	35.5	1.34, m	35.6	1.34, m	35.5	1.32, m	35.6
	1.12, m		1.12, m		1.12, m		1.12, m	
5	1.34, m	46.2	1.34, m	46.2	1.34, m	46.2	1.32, m	46.1
	1.32, m		1.31, m		1.31, m		1.29, m	
6	1.73, m	33.3	1.74, m	33.3	1.75, m	33.3	1.74, m	33.1
	0.94, m		0.94, m		0.95, m		0.92, m	
8	1.67, m	36.1	1.67, m	36.1	1.67, m	36.1	1.38, m	38.5
	0.75, m		0.74, m		0.74, m		0.72, m	
9	56.2	56.2	56.2	56.2	56.2	56.2	55.9	55.9
	36.8		36.9		36.8		36.8	
10	1.51, m	22.8	1.52, m	22.8	1.52, m	22.8	1.63, m	24.4
	1.31, m		1.32, m		1.32, m		1.34, m	
11	2.01, m	38.9	2.01, m	38.8	2.01, m	38.8	2.64, m	38.2
	1.11, m		1.10, m		1.10, m		1.09, m	
12	48.1	48.1	48.1	48.1	48.1	48.1	55.8	55.8
	1.10, m		1.10, m		1.10, m		1.39, m	
13	2.17, m	38.5	2.16, m	38.6	2.16, m	38.6	1.81, m	58.4
	1.34, m		1.33, m		1.33, m		1.19, m	
14	4.26, td (7.7, 3.7)	72.8	4.26, td (7.9, 3.7)	72.8	4.26, td (7.9, 3.7)	72.8	1.80, m	24.4
	1.19, m		1.19, m		1.19, m		0.89, m	
15	3.95, d (11.6)	62.3	3.95, d (11.6)	62.3	3.95, d (11.6)	62.3	1.48, m	57.4
	3.59, d (11.6)		3.60, d (11.6)		3.60, d (11.6)			
16	0.88, s	12.8	0.88, s	12.8	0.88, s	12.9	0.76, s	12.8
	1.94, m		1.93, m		1.93, m		1.49, m	
17	1.02, d (6.8)	19.0	1.07, d (6.7)	19.1	1.02, d (6.7)	19.1	1.09, d (6.3)	19.1
	1.87, m		1.73, m		1.83, m		1.45, m	
18	1.21, m	35.5	1.18, m	35.5	1.18, m	36.8	1.14, m	36.0
	2.15, m		2.13, m		2.04, m		2.07, m	
19	1.98, m	32.4	1.94, m	26.8	1.83, m	29.1	1.90, m	29.9
20	158.0	158.0	148.2	148.2	146.9	146.9	147.9	147.9
21	4.71, br s 4.70, br s	106.7	5.19, q (6.7)	116.6	5.17, q (6.7)	117.7	5.18, q (6.7)	116.8
			1.59, d (6.3)		1.58, d (6.3)		1.56, d (6.7)	
22	2.29, h (6.5)	34.8	2.24, m	36.0	2.85, m	29.8	2.19, m	35.6
	1.03, d (6.8)		0.99, d (6.8)		0.99, d (6.8)		0.98, d (6.8)	
23	1.03, d (6.8)	22.3	0.99, d (6.8)	22.6	0.99, d (6.8)	21.4	0.98, d (6.8)	22.6
24	4.49, d (7.6)	101.8	4.49, d (7.6)	101.8	4.49, d (7.6)	101.8	4.48, d (7.5)	101.8
	3.70, m		3.69, t (10.2)		3.69, t (10.2)		3.70, t (10.2)	
25	3.65, dd (9.6, 3.3)	74.8	3.65, dd (9.6, 3.3)	74.8	3.65, dd (9.6, 3.3)	74.8	3.64, dd (9.5, 3.3)	74.8
	3.84, d (3.2)		3.84, d (3.2)		3.84, d (3.2)		3.84, d (3.1)	
26	3.50, t (6.1)	76.4	3.50, t (6.1)	76.4	3.50, t (6.1)	76.4	3.49, t (6.2)	76.4
	3.73, m		3.73, m		3.73, m		3.73, m	
27	3.71, m	62.7	3.71, m	62.7	3.71, m	62.7	3.71, m	62.7
28	4.56, d (7.9)	105.2	4.56, d (7.9)	105.2	4.56, d (7.9)	105.2	4.56, d (7.9)	105.2
	3.25, dd (9.1, 7.9)		3.25, dd (9.1, 7.9)		3.25, dd (9.1, 7.9)		3.25, dd (9.0, 7.8)	
29	3.37, t (8.8)	77.7	3.37, t (8.8)	77.7	3.37, t (8.8)	77.7	3.37, t (8.9)	77.7
	3.33, t (9.3)		3.33, t (9.3)		3.33, t (9.3)		3.33, t (9.4)	
30	3.29, m	78.4	3.29, m	78.4	3.29, m	78.4	3.28, m	78.4
	3.84, dd (11.2, 2.3)		3.84, dd (11.1, 2.3)		3.84, dd (11.1, 2.3)		3.84, dd (13.5, 2.8)	
31	3.71, m	62.4	3.71, m	62.4	3.71, m	62.4	3.71, m	62.4

Compound **4** was isolated as a pale yellowish amorphous solid with a molecular formula C<sub>41</sub>H<sub>68</sub>O<sub>13</sub>. The <sup>1</sup>H NMR spectrum of **4** was very similar to the one of **2** except for the absence of the signals corresponding to the AB system of H<sub>2</sub>-18 and a shielding observed for  $\delta_H$  2.64 (m, 1H, H-12a). The only explanation consistent with all these observations, including the molecular formula, was the replacement of the hydroxyl group at C-18 by a carboxylic acid. This interpretation was further supported by a key H-17/C-18 HMBC correlation. Based on the chemical shift of the signal H-25 the configuration of the double bond was found to be the same as in **2**.

Compound **5** was isolated as a white amorphous solid with a molecular formula of C<sub>43</sub>H<sub>66</sub>O<sub>15</sub>. Despite strong differences when compared with **1–4**, the NMR data of **5** evidenced that the molecule was a steroidal saponin (Table 2). The aglycone exhibited an unusual skeleton with the presence of a terminal methylated cyclopropyl ring on the lateral chain. This assumption was based on the shielded signals of H-25 and H-26 but also by COSY, HSQC, and HMBC data analyses with the key

H-27/C-24, H-27/C-26 HMBC correlations. Further analysis of  $^1\text{H}$  NMR data revealed the *E* geometry of the olefinic bond ( $J_{\text{H-22,-23}} = 15.2$  Hz). No clear nOe correlations were observed for assessing the relative configuration around the cyclopropane ring. Gratifyingly, comparison with literature data and synthetic analogues of sterols with an identical side-chain led us to propose a *trans* configuration for the substituents at C-24 and C-25 of this ring [48–51]. To confirm this configuration in our case, we decided to look further into the coupling constants of the signals corresponding to the cyclopropane protons. Only the signals of the methylene and their multiplicity were clearly identified in the  $^1\text{H}$  NMR spectrum (Figure 2). In the case of a *trans* configuration of the two substituents around the cyclopropane,  $\text{H}_a$  and  $\text{H}_b$  would have the same splitting pattern as they would have in the presence of a pseudo  $\text{C}_2$  axial symmetry perpendicular to the cyclopropane plane. The  $^3J$  coupling constants between protons in a *cis* configuration are known to be between 8 and 10 Hz while values below 7 Hz are always observed when placed in a *trans* configuration. The multiplicity for both signals is observed as a doublet or triplet with coupling constants around 8 and 4 Hz, respectively. This same splitting pattern for both signals is only consistent for a *trans* configuration. Indeed, for a *cis* configuration, one of the two *gem* protons  $\text{H}_b$  would exhibit two large  $^3J$  coupling constants of 8 Hz. We, therefore, confirm a *trans* configuration for the two substituents and estimate the *gem*  $^2J$  coupling constants between  $\text{H}_a$  and  $\text{H}_b$  to be around 4 Hz. The presence of a carboxyl group at C-18 was inferred first from the HRESIMS data and then from the deshielding of H-12a, exactly in the same manner as for compound 4. Another difference with 4 arose from the absence of the signal corresponding to the oxygenated methine at C-16. This feature was confirmed by COSY, HSQC, and HMBC correlations. Looking at the glycosidic part of the saponin, the relative configuration was similar to those of 1–4, therefore, confirming one galactose linked to the aglycone and one glucose linked to the galactose. HMBC showed long-range correlations between H-1''/C-3', H-2'/C<sub>Ac</sub> ( $\delta_{\text{C}}$  172.2), and H-6''/C<sub>Ac</sub> ( $\delta_{\text{C}}$  172.8), thereby indicating the presence of two acetyl groups at C-2' and C-6''. Unlike compounds 1–4, the glycosidic link between both sugar residues was placed at C-3' of the galactose. Deshielding of the signal of C-3' at  $\delta_{\text{C}}$  82.4 in the  $^{13}\text{C}$  NMR spectrum confirmed this new substitution pattern.



**Figure 2.** Assignment of the relative configuration of the disubstituted cyclopropane through  $^1\text{H}$  NMR coupling constants [52].

**Table 2.** NMR spectroscopic data for poecillastrosides E–G (5–7) in CD<sub>3</sub>OD (500 MHz for <sup>1</sup>H NMR data and 125 MHz for <sup>13</sup>C NMR data of 5; 600 MHz for <sup>1</sup>H data and 150 MHz for <sup>13</sup>C data of 6 and 7).

No.	5		6		7	
	$\delta_{\text{H}}, \text{mult. (J in Hz)}$	$\delta_{\text{C}}$	$\delta_{\text{H}}, \text{mult. (J in Hz)}$	$\delta_{\text{C}}$	$\delta_{\text{H}}, \text{mult. (J in Hz)}$	$\delta_{\text{C}}$
1	1.70, m 0.97, m	38.0	1.72, m 0.97, m	38.2	1.72, m 0.98, m	38.2
2	1.85, m 1.44, m	30.4	1.86, m 1.46, m	30.7	1.87, m 1.46, m	30.8
3	3.62, m 1.58, m	79.9	3.63, m 1.58, m	80.0	3.62, m 1.58, m	80.0
4	1.17, m 1.12, m	35.8	1.17, m 1.09, m	35.9	1.19, m 1.10, m	36.0
5	1.32, m 1.29, m	46.0	1.32, m 1.29, m	46.1	1.31, m 1.27, m	46.1
6	1.76, m 0.94, m	30.3	1.68, m 0.87, m	29.9	1.67, m 0.87, m	30.4
7	1.53, m 0.73, m	33.1	1.43, m 0.68, m	33.5	1.43, m 0.68, m	33.5
8		38.8		37.1		37.0
9		55.9		56.0		56.0
10		36.7		36.8		36.8
11	1.63, m 1.31, m	24.4	1.53, m 1.36, m	22.3	1.53, m 1.34, m	22.3
12	2.63, m 1.10, m	38.4	2.44, d (12.8) 0.94, m	35.9	2.44, dt (12.7, 3.4) 0.94, m	35.9
13		55.6		47.9		47.9
14	1.38, m	58.4	1.11, m	57.6	1.12, m	57.6
15	1.75, m 1.30, m	30.8	1.70, m 1.30, m	29.9	1.71, m 1.29, m	29.9
16	1.78, m 1.53, m	25.8	1.54, m 0.98, m	25.0	1.54, m 0.98, m	24.9
17	1.46, m	57.3	1.15, m	58.2	1.16, m	58.1
18		180.1	3.65, d (11.5) 3.45, d (11.6)	60.2	3.65, d (11.1) 3.45, d (11.7)	60.4
19	0.73, s	12.7	0.83, s	12.7	0.83, s	12.7
20	1.92, m	42.4	2.26, m	41.7	2.26, m	41.7
21	1.07, d (6.3)	21.2	1.07, d (5.9)	22.1	1.07, d (6.4)	22.1
22	5.21, dd (15.1, 8.5)	134.6	5.22, dd (14.8, 9.0)	136.0	5.22, dd (15.2, 8.9)	136.0
23	4.90, m	132.4	4.94, dd (14.8, 8.1)	131.6	4.94, dd (15.2, 8.3)	131.6
24	0.96, m	23.4	0.93, m	23.4	0.93, m	23.4
25	0.62, m	15.5	0.62, m	15.5	0.62, m	15.5
26	0.44, td (9.0, 4.5) 0.36, dt (9.0, 4.5)	15.2	0.45, m 0.36, m	15.2	0.45, m 0.35, m	15.2
27	1.03, d (5.9)	18.8	1.03, d (5.8)	18.9	1.03, d (5.9)	18.9
1'	4.56, d (8.0)	101.1	4.55, d (7.9)	101.2	4.56, d (8.0)	101.2
2'	5.11, dd (8.4, 8.1)	72.5	5.12, dd (9.0, 7.7)	72.6	5.11, dd (10.1, 8.0)	72.4
2'-Ac	2.06, s	21.2	2.06, s	21.2	2.06, s	21.2
3'	3.76, dd (10.2, 3.3)	172.2	3.80, dd (10.0, 2.8)	171.2	3.76, dd (10.1, 3.2)	172.2
4'	4.07, d (3.2)	82.4	4.11, d (3.1)	82.2	4.07, d (3.4)	82.4
5'	3.55, t (6.1)	70.2	3.56, t (6.2)	70.2	3.55, t (6.4)	70.2
6'	3.74, m 3.73, m	76.4	3.74, m 3.72, m	76.4	3.74, m 3.72, m	76.4
1''	4.39, d (7.6)	62.3		62.2		62.1
2''	3.21, t (8.3)	106.0	4.38, d (7.9)	106.0	4.38, d (8.0)	106.0
3''	3.32, t (10.1)	74.6	3.19, t (8.3)	74.8	3.21, t (8.3)	74.7
4''	3.28, t (9.6)	77.7	3.35, m	77.9	3.33, m	77.9
5''	3.46, m	71.6	3.28, m	71.3	3.29, m	71.5
6''	4.38, d (11.9) 4.20, dd (11.9, 6.1)	75.3	3.64, m 3.84, m 3.67, m	80.0 62.5	3.46, m 4.38, dd (11.9, 2.7) 4.20, dd (11.9, 6.2)	75.3 64.7
6''-Ac	2.06, s	20.8			2.06, s	20.8
		172.8				172.8

Compound **6** was isolated as a white amorphous solid with a molecular formula of C<sub>41</sub>H<sub>66</sub>O<sub>13</sub>. The spectroscopic data were very similar to those of **5**, thereby suggesting a close aglycone moiety. However, some changes were noticed by HSQC and HMBC analyses. Indeed, in the aglycone moiety, we observed the same AB system for H<sub>2</sub>-18 as that present in compounds **1**–**3**. The long-range H-17/C-18 HMBC correlation confirmed the presence of an oxygenated methylene at C-13. In the D- $\beta$ -glucose residue, the chemical shifts, and the COSY data were consistent with a terminal primary alcohol at C-6'', thereby implying the loss of the acetate at this position.

Compound **7** was isolated as a white amorphous solid with a molecular formula C<sub>43</sub>H<sub>68</sub>O<sub>14</sub>. The <sup>1</sup>H NMR spectrum evidenced the fact that **7** is a close analogue of **6**. The long-range H-6''/C<sub>Ac</sub>

( $\delta_C$  172.8) HMBC correlation revealed the presence of an acetate group linked at O-6'' as in compound 5. The relative configuration of 7 was the same as those of 5 and 6.

Poecillastrosides A–G were tested in a panel of antimicrobial and cytotoxicity assays, including antibacterial activity against Gram positive (methicillin resistant (MRSA) and methicillin sensitive (MSSA) *Staphylococcus aureus*), and Gram negative bacteria (*Escherichia coli*, *Klebsiella pneumoniae*, *Pseudomonas aeruginosa*, and *Acinetobacter baumannii*), antifungal activity against *Aspergillus fumigatus*, and cytotoxicity against the hepatic tumoral cell line hep\_G2. Poecillastrosides D (4) ( $MIC_{90}$  = 6  $\mu\text{g/mL}$ ) and E (5) ( $MIC_{90}$  = 24  $\mu\text{g/mL}$ ) were the only two molecules active in the assay against *A. fumigatus*, revealing a key role of the carboxylic acid functionality at C-18 in the antifungal activity of this structural class. On the other hand, cytotoxicity assays also revealed weak activity of some members of the family against the hep\_G2 human cell line, with  $IC_{50}$  values of 38, 28, and 89  $\mu\text{g/mL}$  for poecillastrosides B, C, and D (2–4), respectively. None of the compounds of this family displayed activity against the bacterial pathogens at the highest concentration tested (96  $\mu\text{g/mL}$  for compound 1–5, and 64  $\mu\text{g/mL}$  for compounds 6 and 7).

### 3. Material and Methods

#### 3.1. General Experimental Procedures

Optical rotations were recorded with a PerkinElmer 343 polarimeter equipped with a 10 cm microcell and a sodium lamp. UV measurements were obtained by extraction of the Diode Array Detector (DAD) signal of the Ultra-High Pressure Liquid Chromatography (UHPLC) Dionex Ultimate 3000 (Thermo Scientific, Waltham, MA, USA). NMR experiments were performed on a 500 MHz (Advance, Bruker, Billerica, MA, USA) or a 600 MHz (Agilent, Santa Clara, CA, USA) spectrometer. Chemical shifts ( $\delta$  in ppm) are referenced to the carbon ( $\delta_C$  49.0) and residual proton ( $\delta_H$  3.31) signals of  $\text{CD}_3\text{OD}$ . High-resolution mass spectra (HRESIMS) were obtained from a mass spectrometer Agilent 6540. HPLC separation and purification were carried out on a Jasco LC-2000 series equipped with a UV detector coupled with an Evaporative Light Scattering Detector, ELSD (Sedere, Alfortville, France).

#### 3.2. Biological Material

*Poecillastra compressa* (Bowerbank, 1866) was collected in the Mediterranean Sea, off the French coasts, on 15 October 2014 at 200 m depth using a Remotely Operated Vehicle (Super Achille, COMEX S.A., Marseille, France). The voucher specimen "CS2ACHP09\_ECH04" is kept at the Marine Station of Endoume (OSU Institut Pythéas, Marseille, France).

#### 3.3. Extraction and Isolation

The dry sponge sample (43.1 g) was ground with a mortar and extracted with a mixture of  $\text{CH}_3\text{OH}/\text{CH}_2\text{Cl}_2$  (1:1,  $v/v$ ) at room temperature, yielding 7.9 g (18% yield from dry-weight) of extract after solvent evaporation. The crude extract was fractionated by RP-C18 vacuum liquid chromatography (elution with a decreasing polarity gradient of  $\text{H}_2\text{O}/\text{CH}_3\text{OH}$  from 1:0 to 0:1, then  $\text{CH}_3\text{OH}/\text{CH}_2\text{Cl}_2$  from 1:0 to 0:1). The  $\text{CH}_3\text{OH}$  (422 mg) fraction was then subjected to RP-HPLC on a preparative phenylhexyl column, 250 mm  $\times$  19 mm, 5  $\mu\text{m}$  (Xselect, Waters, Milford, CT, USA), using a mobile phase of water (A) and acetonitrile (B). The method was developed on 30 min acquisition time: isocratic 60% B for 15 min, then linear gradient to 98% B in 1 min, held at 98% B for 10 min, back to 60% B in 1 min, and held at that percentage of B for 3 min. Selected fractions from this chromatography were then purified by RP-HPLC on a semi-preparative HTec C18 column, 250 mm  $\times$  10 mm, 5  $\mu\text{m}$  (Nucleodur, Macherey-Nagel, Düren, Germany), with the following methods for each subsequent purification: isocratic 47% B to afford pure 1 (4.3 mg,  $9.98 \times 10^{-3}\%$   $w/w$ ), isocratic 49% B to afford 2 (6.2 mg,  $1.44 \times 10^{-2}\%$   $w/w$ ) and 3 (1.4 mg,  $3.49 \times 10^{-3}\%$   $w/w$ ), isocratic 50% B to afford 4 (1.6 mg,

$3.71 \times 10^{-3}\%$  w/w), isocratic 51% B to afford **5** (0.9 mg,  $2.09 \times 10^{-3}\%$  w/w), and isocratic 53% B to afford **6** (0.7 mg,  $1.62 \times 10^{-3}\%$  w/w) and **7** (0.8 mg,  $1.86 \times 10^{-3}\%$  w/w).

Poecillastroside A (**1**): Yellow, amorphous solid;  $[\alpha]_D^{20} +12.8$  (c 0.1, CH<sub>3</sub>OH); UV (DAD)  $\lambda_{\max}$  195 nm; <sup>1</sup>H NMR and <sup>13</sup>C NMR data, see Table 1; HRESIMS (−)  $m/z$  755.4582 [M − H]<sup>−</sup> (calcd. for C<sub>40</sub>H<sub>67</sub>O<sub>13</sub>, 755.4587,  $\Delta$  − 0.7 ppm).

Poecillastroside B (**2**): Yellow, amorphous solid;  $[\alpha]_D^{20} +13.2$  (c 0.1, CH<sub>3</sub>OH); UV (DAD)  $\lambda_{\max}$  210 nm; <sup>1</sup>H NMR and <sup>13</sup>C NMR data, see Table 1; HRESIMS (−)  $m/z$  769.4743 [M − H]<sup>−</sup> (calcd. for C<sub>41</sub>H<sub>69</sub>O<sub>13</sub>, 769.4744,  $\Delta$  − 0.1 ppm).

Poecillastroside C (**3**): Yellow, amorphous solid;  $[\alpha]_D^{20} +13.0$  (c 0.1, CH<sub>3</sub>OH); UV (DAD)  $\lambda_{\max}$  212 nm; <sup>1</sup>H NMR and <sup>13</sup>C NMR data, see Table 1; HRESIMS (−)  $m/z$  769.4745 [M − H]<sup>−</sup> (calcd. for C<sub>41</sub>H<sub>69</sub>O<sub>13</sub>, 769.4744,  $\Delta$  + 0.1 ppm).

Poecillastroside D (**4**): Yellow, amorphous solid;  $[\alpha]_D^{20} +8.9$  (c 0.1, CH<sub>3</sub>OH); UV (DAD)  $\lambda_{\max}$  222 nm; <sup>1</sup>H NMR and <sup>13</sup>C NMR data, see Table 1; HRESIMS (+)  $m/z$  791.4567 [M + Na]<sup>+</sup> (calcd. for C<sub>41</sub>H<sub>68</sub>NaO<sub>13</sub>, 791.4563,  $\Delta$  + 0.5 ppm).

Poecillastroside E (**5**): White, amorphous solid;  $[\alpha]_D^{20} -6.2$  (c 0.1, CH<sub>3</sub>OH); UV (DAD)  $\lambda_{\max}$  220 nm; <sup>1</sup>H NMR and <sup>13</sup>C NMR data, see Table 2; HRESIMS (+)  $m/z$  845.4307 [M + Na]<sup>+</sup> (calcd. for C<sub>43</sub>H<sub>66</sub>NaO<sub>15</sub>, 845.4299,  $\Delta$  + 0.9 ppm).

Poecillastroside F (**6**): White, amorphous solid;  $[\alpha]_D^{20} -27.3$  (c 0.1, CH<sub>3</sub>OH); UV (DAD)  $\lambda_{\max}$  222 nm; <sup>1</sup>H NMR and <sup>13</sup>C NMR data, see Table 2; HRESIMS (+)  $m/z$  789.4405 [M + Na]<sup>+</sup> (calcd. for C<sub>41</sub>H<sub>66</sub>NaO<sub>13</sub>, 789.4401,  $\Delta$  + 0.5 ppm).

Poecillastroside G (**7**): White, amorphous solid;  $[\alpha]_D^{20} -14.1$  (c 0.1, CH<sub>3</sub>OH); UV (DAD)  $\lambda_{\max}$  225 nm; <sup>1</sup>H NMR and <sup>13</sup>C NMR data, see Table 2; HRESIMS (+)  $m/z$  831.4518 [M + Na]<sup>+</sup> (calcd. for C<sub>43</sub>H<sub>68</sub>NaO<sub>14</sub>, 831.4507,  $\Delta$  + 1.3 ppm).

### 3.4. Determination of the Absolute Configuration of the Pyranoses

Hydrolysis of glycosides and derivatization of the subsequent monosaccharides were performed individually following previously described methodologies [46]. The monosaccharide derivatives separation was carried out by UHPLC-HRMS on Acquity BEH (Ethylene Bridged Hybrid) C18 1.7  $\mu$ m, 2.1 mm  $\times$  100 mm (Waters). The column was heated at 40 °C. The eluent consisted of water with 0.1% formic acid (A) and acetonitrile/methanol/isopropanol (50:25:25, v/v/v) with 0.1% formic acid (B). The analysis was performed in isocratic mode at 13% B and at a flow rate of 360  $\mu$ L/min. The injection volume was set at 3  $\mu$ L. The identity of all monosaccharide derivatives was confirmed after extraction of the ion [M + H]<sup>+</sup> at  $m/z$  433.1098 (Figure S55).

### 3.5. Evaluation of the Biological Activities

Compounds **1–7** were tested for their ability to inhibit the growth of Gram positive bacteria (*S. aureus* ATCC29213 (MSSA), and *S. aureus* MB5393 (MRSA)) and Gram negative bacteria (*E. coli* ATCC25922, *K. pneumoniae* ATCC700603, *P. aeruginosa* PAO1, and *A. baumannii* CL5973), and fungi (*A. fumigatus* ATCC46645), following previously described methodologies [53,54]. Cytotoxic activity against the hepatic human tumoral cell line hep\_G2 was determined as previously reported [55].

## 4. Conclusions

Poecillastrosides A–G (**1–7**) share an unusual oxidized methyl at C-18, and they are the first saponins exhibiting this feature. The structures of poecillastrosides E–G (**5–7**) also incorporate a

terminal methylated cyclopropyl ring already known in some sponge steroids and already investigated for biosynthetic studies [56]. This cyclopropanation process could lead to the cholestane skeleton, then ergostane, and finally poriferastane, all of them being present in the metabolome of this sponge. Many sterols containing a cyclopropyl ring have been isolated to date [57], but to our best knowledge, this is the first time that saponins containing a 3-membered ring on the side-chain have been reported. Poecillastrosides D (4) and E (5), bearing a carboxylic acid at C-18, were found to be the most bioactive compounds in the antimicrobial bioassays with an interesting antifungal activity against *Aspergillus fumigatus*.

**Supplementary Materials:** The following are available online at [www.mdpi.com/1660-3397/15/7/199/s1](http://www.mdpi.com/1660-3397/15/7/199/s1): HRMS and  $^1\text{H}$ ,  $^{13}\text{C}$ , COSY, TOCSY, HSQC, HMBC, and NOESY NMR data for compounds 1–7 as well as procedures for absolute configuration of compound 3.

**Acknowledgments:** This work was partially funded by the Swiss company Ferring. The sampling was supported by the “Agence des Aires Marines Protégées (AAMP)”, a French establishment dedicated to the protection of the marine environment. The authors are grateful to the COMEX crew who operated the MINIBEX vessel and its ROV SUPER ACHILLE. We are grateful to G. Genta-Jouve for fruitful discussions about the relative configuration of the cyclopropane. The company Cosmo International Ingredients supported the work of K. Calabro. R. Doohan (NUI Galway) is acknowledged for her help in the record of NMR spectra and H. Solanki for his help in the HRMS acquisition.

**Author Contributions:** O.P.T. conceived and designed the experiments; E.L.K., D.R. and K.C. performed the experiments; C.D., M.d.I.C., B.C. and F.R. performed the bioassays; K.C., O.P.T., R.L. and B.S. analyzed the data; T.P. collected and identified the biomaterial; K.C., F.R. and O.P.T. wrote the paper.

**Conflicts of Interest:** The authors declare no conflict of interest.

## References

1. Stonik, V.A.; Kalinin, V.I.; Avilov, S.A. Toxins from sea cucumbers (holothuroids): Chemical structures, properties, taxonomic distribution, biosynthesis and evolution. *J. Nat. Toxins* **1999**, *8*, 235–248. [PubMed]
2. Makarieva, T.N.; Stonik, V.A.; Kapustina, I.I.; Boguslavsky, V.M.; Dmitrenko, A.S.; Kalinin, V.I.; Cordeiro, M.L.; Djerassi, C. Biosynthetic studies of marine lipids. 42. Biosynthesis of steroid and triterpenoid metabolites in the sea cucumber *Eupentacta fraudatrix*. *Steroids* **1993**, *58*, 508–517. [CrossRef]
3. Burnell, D.J.; Apsimon, J.W. Chapter 6—Echinoderm Saponins. In *Marine Natural Products*; Scheuer, P.J., Ed.; Academic Press: Waltham, MA, USA, 1983; pp. 287–389.
4. Qi, S.; Zhang, S.; Huang, J.; Xiao, Z.; Wu, J.; Li, Q. Complete  $^1\text{H}$  and  $^{13}\text{C}$  NMR assignments of four new steroidal glycosides from a gorgonian coral *Junceella juncea*. *Magn. Reson. Chem.* **2005**, *43*, 266–268. [CrossRef] [PubMed]
5. Wang, S.-K.; Dai, C.-F.; Duh, C.-Y. Cytotoxic Pregnane Steroids from the Formosan Soft Coral *Stereonephthya crystallina*. *J. Nat. Prod.* **2006**, *69*, 103–106. [CrossRef] [PubMed]
6. Ivanchina, N.V.; Kicha, A.A.; Stonik, V.A. Steroid glycosides from marine organisms. *Steroids* **2011**, *76*, 425–454. [CrossRef] [PubMed]
7. Kitagawa, I.; Kobayashi, M.; Okamoto, Y.; Yoshikawa, M.; Hamamoto, Y. Structures of Sarasinosides A1', B1', and C1'; New Norlanostane-Triterpenoid Oligoglycosides from the Palauan Marine Sponge *Asteropus sarasinosum*. *Chem. Pharm. Bull.* **1987**, *35*, 5036–5039. [CrossRef] [PubMed]
8. Espada, A.; Jiménez, C.; Rodríguez, J.; Crews, P.; Riguera, R. Sarasinosides D–G: Four new triterpenoid saponins from the sponge *Asteropus sarasinosum*. *Tetrahedron* **1992**, *48*, 8685–8696. [CrossRef]
9. Lee, H.-S.; Seo, Y.; Cho, K.W.; Rho, J.-R.; Shin, J.; Paul, V.J. New triterpenoid saponins from the sponge *Melophylus isis*. *J. Nat. Prod.* **2000**, *63*, 915–919. [CrossRef] [PubMed]
10. Dai, H.-F.; Edrada, R.A.; Ebel, R.; Nimtz, M.; Wray, V.; Proksch, P. Norlanostane triterpenoidal saponins from the marine sponge *Melophylus sarassinorum*. *J. Nat. Prod.* **2005**, *68*, 1231–1237. [CrossRef] [PubMed]
11. Lee, J.-H.; Jeon, J.-E.; Lee, Y.-J.; Lee, H.-S.; Sim, C.J.; Oh, K.-B.; Shin, J. Nortriterpene Glycosides of the Sarasinoside Class from the Sponge *Lipastrotethya* sp. *J. Nat. Prod.* **2012**, *75*, 1365–1372. [CrossRef] [PubMed]
12. Antonov, A.S.; Kalinovskii, A.I.; Stonik, V.A.; Evtushenko, E.V.; Elyakov, G.B. Structure of uloside A, a new triterpenoid glycoside from the *Ulosa* sp. sponge. *Russ. Chem. Bull.* **1994**, *43*, 1265–1269. [CrossRef]

13. Antonov, A.S.; Kalinovsky, A.I.; Stonik, V.A. Ulososide B, a new unusual norlanostane-triterpene glycoside and its genuine aglycone from the Madagascar sponge *Ulosa* sp. *Tetrahedron Lett.* **1998**, *39*, 3807–3808. [[CrossRef](#)]
14. Colorado, J.; Muñoz, D.; Marquez, D.; Marquez, M.; Lopez, J.; Thomas, O.P.; Martinez, A. Ulososides and Urabosides—Triterpenoid Saponins from the Caribbean Marine Sponge *Ectyoplasia ferox*. *Molecules* **2013**, *18*, 2598–2610. [[CrossRef](#)] [[PubMed](#)]
15. Cachet, N.; Regalado, E.L.; Genta-Jouve, G.; Mehiri, M.; Amade, P.; Thomas, O.P. Steroidal glycosides from the marine sponge *Pandaros acanthifolium*. *Steroids* **2009**, *74*, 746–750. [[CrossRef](#)] [[PubMed](#)]
16. Regalado, E.L.; Tasdemir, D.; Kaiser, M.; Cachet, N.; Amade, P.; Thomas, O.P. Antiprotozoal Steroidal Saponins from the Marine Sponge *Pandaros acanthifolium*. *J. Nat. Prod.* **2010**, *73*, 1404–1410. [[CrossRef](#)] [[PubMed](#)]
17. Regalado, E.L.; Jimenez-Romero, C.; Genta-Jouve, G.; Tasdemir, D.; Amade, P.; Nogueiras, C.; Thomas, O.P. Acanthifoliosides, minor steroidal saponins from the Caribbean sponge *Pandaros acanthifolium*. *Tetrahedron* **2011**, *67*, 1011–1018. [[CrossRef](#)]
18. Regalado, E.L.; Turk, T.; Tasdemir, D.; Gorjanc, M.; Kaiser, M.; Thomas, O.P.; Fernandez, R.; Amade, P. Cytotoxic and haemolytic steroidal glycosides from the Caribbean sponge *Pandaros acanthifolium*. *Steroids* **2011**, *76*, 1389–1396. [[CrossRef](#)] [[PubMed](#)]
19. Ryu, G.; Choi, B.W.; Lee, B.H.; Hwang, K.-H.; Lee, U.C.; Jeong, D.S.; Lee, N.H. Wondosterols A–C, three steroidal glycosides from a Korean marine two-sponge association. *Tetrahedron* **1999**, *55*, 13171–13178. [[CrossRef](#)]
20. D’Auria, M.V.; Paloma, L.G.; Minale, L.; Riccio, R.; Debitus, C. Structure characterization by two-dimensional NMR spectroscopy, of two marine triterpene oligoglycosides from a pacific sponge of the genus *Erylus*. *Tetrahedron* **1992**, *48*, 491–498. [[CrossRef](#)]
21. Gulavita, N.K.; Wright, A.E.; Kelly-Borges, M.; Longley, R.E.; Yarwood, D.; Sills, M.A. Eryloside E from an Atlantic sponge *Erylus goffrilleri*. *Tetrahedron Lett.* **1994**, *35*, 4299–4302. [[CrossRef](#)]
22. Stead, P.; Hiscox, S.; Robinson, P.S.; Pike, N.B.; Sidebottom, P.J.; Roberts, A.D.; Taylor, N.L.; Wright, A.E.; Pomponi, S.A.; Langley, D. Eryloside F, a novel penasterol disaccharide possessing potent thrombin receptor antagonist activity. *Bioorg. Med. Chem. Lett.* **2000**, *10*, 661–664. [[CrossRef](#)]
23. Shin, J.; Lee, H.-S.; Woo, L.; Rho, J.-R.; Seo, Y.; Cho, K.W.; Sim, C.J. New triterpenoid saponins from the sponge *Erylus nobilis*. *J. Nat. Prod.* **2001**, *64*, 767–771. [[CrossRef](#)] [[PubMed](#)]
24. Antonov, A.S.; Kalinovsky, A.I.; Stonik, V.A.; Afiyatullo, S.S.; Aminin, D.L.; Dmitrenok, P.S.; Mollo, E.; Cimino, G. Isolation and Structures of Erylosides from the Caribbean Sponge *Erylus formosus*. *J. Nat. Prod.* **2007**, *70*, 169–178. [[CrossRef](#)] [[PubMed](#)]
25. Jaspars, M.; Crews, P. A triterpene tetrasaccharide, formoside, from the Caribbean Choristida sponge *Erylus formosus*. *Tetrahedron Lett.* **1994**, *35*, 7501–7504. [[CrossRef](#)]
26. Takada, K.; Nakao, Y.; Matsunaga, S.; van Soest, R.W.M.; Fusetani, N. Nobiloside, a New Neuraminidase Inhibitory Triterpenoid Saponin from the Marine Sponge *Erylus nobilis*. *J. Nat. Prod.* **2002**, *65*, 411–413. [[CrossRef](#)] [[PubMed](#)]
27. Fouad, M.; Al-Trabeen, K.; Badran, M.; Wray, V.; Edrada, R.; Proksch, P.; Ebel, R. New steroidal saponins from the sponge *Erylus lendenfeldi*. *ARKIVOC* **2004**, *37*, 17–27.
28. Sandler, J.S.; Forsburg, S.L.; Faulkner, D.J. Bioactive steroidal glycosides from the marine sponge *Erylus lendenfeldi*. *Tetrahedron* **2005**, *61*, 1199–1206. [[CrossRef](#)]
29. Okada, Y.; Matsunaga, S.; van Soest, R.W.M.; Fusetani, N. Sokodosides, Steroid Glycosides with an Isopropyl Side Chain, from the Marine Sponge *Erylus placenta*. *J. Org. Chem.* **2006**, *71*, 4884–4888. [[CrossRef](#)] [[PubMed](#)]
30. Gabant, M.; Schmitz-Afonso, I.; Gallard, J.-F.; Menou, J.-L.; Laurent, D.; Debitus, C.; Al-Mourabit, A. Sulfated Steroids: Ptilosteroids A–C and Ptilosaponosides A and B from the Solomon Islands Marine Sponge *Ptilocaulis spiculifer*. *J. Nat. Prod.* **2009**, *72*, 760–763. [[CrossRef](#)] [[PubMed](#)]
31. Kalinovsky, A.I.; Antonov, A.S.; Afiyatullo, S.S.; Dmitrenok, P.S.; Evtuschenko, E.V.; Stonik, V.A. Mycaloside A, a new steroid oligoglycoside with an unprecedented structure from the Caribbean sponge *Mycale laxissima*. *Tetrahedron Lett.* **2002**, *43*, 523–525. [[CrossRef](#)]
32. Antonov, A.S.; Afiyatullo, S.S.; Kalinovsky, A.I.; Ponomarenko, L.P.; Dmitrenok, P.S.; Aminin, D.L.; Agafonova, I.G.; Stonik, V.A. Mycalosides B–I, Eight New Spermostatic Steroid Oligoglycosides from the Sponge *Mycale laxissima*. *J. Nat. Prod.* **2003**, *66*, 1082–1088. [[CrossRef](#)] [[PubMed](#)]

33. Campagnuolo, C.; Fattorusso, E.; Tagliatela-Scafati, O. Feroxosides A–B, two norlanostane tetraglycosides from the Caribbean sponge *Ectyoplasia ferox*. *Tetrahedron* **2001**, *57*, 4049–4055. [CrossRef]
34. Glensk, M.; Wray, V.; Nimtz, M.; Schöpke, T. Silenosides A–C, Triterpenoid Saponins from *Silene vulgaris*. *J. Nat. Prod.* **1999**, *62*, 717–721. [CrossRef] [PubMed]
35. Wang, W.; Hong, J.; Lee, C.-O.; Im, K.S.; Choi, J.S.; Jung, J.H. Cytotoxic Sterols and Saponins from the Starfish *Certanardoa semiregularis*. *J. Nat. Prod.* **2004**, *67*, 584–591. [CrossRef] [PubMed]
36. Wang, W.; Jang, H.; Hong, J.; Lee, C.-O.; Bae, S.-J.; Shin, S.; Jung, J.H. New cytotoxic sulfated saponins from the starfish *Certanardoa semiregularis*. *Arch. Pharm. Res.* **2005**, *28*, 285–289. [CrossRef] [PubMed]
37. Kicha, A.A.; Ivanchina, N.V.; Huong, T.T.T.; Kalinovsky, A.I.; Dmitrenok, P.S.; Fedorov, S.N.; Dyshlovoy, S.A.; Long, P.Q.; Stonik, V.A. Two new asterosaponins, archasterosides A and B, from the Vietnamese starfish *Archaster typicus* and their anticancer properties. *Bioorg. Med. Chem. Lett.* **2010**, *20*, 3826–3830. [CrossRef] [PubMed]
38. Rashid, M.A.; Gustafson, K.R.; Crouch, R.C.; Groweiss, A.; Pannell, L.K.; Van, Q.N.; Boyd, M.R. Application of High-Field NMR and Cryogenic Probe Technologies in the Structural Elucidation of Poecillastrin A, a New Antitumor Macrolide Lactam from the Sponge *Poecillastra* Species. *Org. Lett.* **2002**, *4*, 3293–3296. [CrossRef] [PubMed]
39. Takada, K.; Choi, B.W.; Rashid, M.A.; Gamble, W.R.; Cardellina, J.H.; Van, Q.N.; Lloyd, J.R.; McMahon, J.B.; Gustafson, K.R. Structural Assignment of Poecillastrins B and C, Macrolide Lactams from the Deep-Water Caribbean Sponge *Poecillastra* Species. *J. Nat. Prod.* **2007**, *70*, 428–431. [CrossRef] [PubMed]
40. Natori, T.; Kataoka, Y.; Kato, S.; Kawai, H.; Fusetani, N. Poecillanosine, a new free radical scavenger from the marine sponge *Poecillastra* spec. aff. *tenuilaminaris*. *Tetrahedron Lett.* **1997**, *38*, 8349–8350. [CrossRef]
41. Killday, K.B.; Longley, R.; McCarthy, P.J.; Pomponi, S.A.; Wright, A.E.; Neale, R.F.; Sills, M.A. Sesquiterpene-Derived Metabolites from the Deep Water Marine Sponge *Poecillastra sollasi*. *J. Nat. Prod.* **1993**, *56*, 500–507. [CrossRef] [PubMed]
42. Makarieva, T.N.; Stonik, V.A.; D'Yachuk, O.G.; Dmitrenok, A.S. Annasterol sulfate, a novel marine sulfated steroid, inhibitor of glucanase activity from the deep water sponge *Poecillastra laminaris*. *Tetrahedron Lett.* **1995**, *36*, 129–132. [CrossRef]
43. Sun, H.H.; Gross, S.S.; Gunasekera, M.; Koehn, F.E. Weinbersterol disulfates A and B, antiviral steroid sulfates from the sponge *Petrosia weinbergi*. *Tetrahedron* **1991**, *47*, 1185–1190. [CrossRef]
44. Tvaroska, I.; Taravel, F.R. Carbon-Proton Coupling Constants In The Conformational Analysis of Sugar Molecules. In *Advances in Carbohydrate Chemistry and Biochemistry*; Derek, H., Ed.; Academic Press: Waltham, MA, USA, 1995; Volume 51, pp. 15–61.
45. Stenutz, R. Coupling Constants of Pyranoses. Available online: <http://www.stenutz.eu/sop/a704.html> (accessed on 19 February 2017).
46. Wang, Y.-H.; Avula, B.; Fu, X.; Wang, M.; Khan, I.A. Simultaneous Determination of the Absolute Configuration of Twelve Monosaccharide Enantiomers from Natural Products in a Single Injection by a UPLC-UV/MS Method. *Planta Med.* **2012**, *78*, 834–837. [CrossRef] [PubMed]
47. Paudel, L.; Adams, R.W.; Király, P.; Aguilar, J.A.; Foroozandeh, M.; Cliff, M.J.; Nilsson, M.; Sándor, P.; Waltho, J.P.; Morris, G.A. Simultaneously Enhancing Spectral Resolution and Sensitivity in Heteronuclear Correlation NMR Spectroscopy. *Angew. Chem. Int. Ed.* **2013**, *52*, 11616–11619. [CrossRef] [PubMed]
48. Bonini, C.; Kinnel, R.B.; Li, M.; Scheuer, P.J.; Djerassi, C. Minor and trace sterols in marine invertebrates. 38. Isolation, structure elucidation, and partial synthesis of papakusterol, a new biosynthetically unusual marine sterol with a cyclopropyl-containing side chain. *Tetrahedron Lett.* **1983**, *24*, 277–280. [CrossRef]
49. Catalan, C.A.N.; Lakshmi, V.; Schmitz, F.J.; Djerassi, C. Minor and trace sterols in marine invertebrates. 39. 24 $\xi$ ,25 $\xi$ -24,26-Cyclocholest-5-en-3 $\beta$ -ol, a novel cyclopropyl sterol. *Steroids* **1982**, *40*, 455–463. [CrossRef]
50. Fujimoto, Y.; Kimura, M.; Terasawa, T.; Khalifa, F.A.M.; Ikekawa, N. Stereocontrolled synthesis and determination of the C-24 and C-25 stereochemistry of glaucasterol. *Tetrahedron Lett.* **1984**, *25*, 1805–1808. [CrossRef]
51. Kobayashi, M.; Mitsuhashi, H. Marine sterols. XII. Glaucasterol, a novel C27 sterol with a unique side chain, from the soft coral *Sarcophyton glaucum*. *Steroids* **1982**, *40*, 665–672. [CrossRef]
52. Wiberg, K.B.; Nist, B.J. The Nuclear Magnetic Resonance Spectra of Cyclopropane Derivatives. *J. Am. Chem. Soc.* **1963**, *85*, 2788–2790. [CrossRef]

53. Audoin, C.; Bonhomme, D.; Ivanisevic, J.; Cruz, M.; Cautain, B.; Monteiro, M.; Reyes, F.; Rios, L.; Perez, T.; Thomas, O.P. Balibalosides, an Original Family of Glucosylated Sesterterpenes Produced by the Mediterranean Sponge *Oscarella balibalo*. *Mar. Drugs* **2013**, *11*, 1477–1489. [[CrossRef](#)] [[PubMed](#)]
54. Braña, A.F.; Sarmiento-Vizcaíno, A.; Pérez-Victoria, I.; Otero, L.; Fernández, J.; Palacios, J.J.; Martín, J.; de la Cruz, M.; Díaz, C.; Vicente, F.; et al. Branimycins B and C, Antibiotics Produced by the Abyssal Actinobacterium *Pseudonocardia carboxydivorans* M-227. *J. Nat. Prod.* **2017**, *80*, 569–573. [[CrossRef](#)] [[PubMed](#)]
55. Cautain, B.; de Pedro, N.; Schulz, C.; Pascual, J.; Sousa, T.d.S.; Martin, J.; Pérez-Victoria, I.; Asensio, F.; González, I.; Bills, G.F.; et al. Identification of the Lipodepsipeptide MDN-0066, a Novel Inhibitor of VHL/HIF Pathway Produced by a New *Pseudomonas* Species. *PLoS ONE* **2015**, *10*, e0125221. [[CrossRef](#)] [[PubMed](#)]
56. Wessjohann, L.A.; Brandt, W.; Thiemann, T. Biosynthesis and Metabolism of Cyclopropane Rings in Natural Compounds. *Chem. Rev.* **2003**, *103*, 1625–1648. [[CrossRef](#)] [[PubMed](#)]
57. Gunasekera, S.P.; Cranick, S.; Pomponi, S.A. New Sterol Ester from a Deep Water Marine Sponge, *Xestospongia* sp. *J. Nat. Prod.* **1991**, *54*, 1119–1122. [[CrossRef](#)]



© 2017 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).