



**HAL**  
open science

## Draft Genome Sequence of the Iridescent Marine Bacterium *Cellulophaga lytica* CECT 8139

Maylis Chapelais-Baron, Isabelle Goubet, Éric Duchaud, Eric Rosenfeld

► **To cite this version:**

Maylis Chapelais-Baron, Isabelle Goubet, Éric Duchaud, Eric Rosenfeld. Draft Genome Sequence of the Iridescent Marine Bacterium *Cellulophaga lytica* CECT 8139. *Genome Announcements*, 2017, 5 (36), pp.e00811-17. 10.1128/genomeA.00811-17. hal-01681826

**HAL Id: hal-01681826**

**<https://hal.science/hal-01681826>**

Submitted on 11 Mar 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



# Draft Genome Sequence of the Iridescent Marine Bacterium *Cellulophaga lytica* CECT 8139

Maylis Chapelais-Baron,<sup>a</sup> Isabelle Goubet,<sup>a</sup> Eric Duchaud,<sup>b</sup> Eric Rosenfeld<sup>a</sup>

UMR 7266, CNRS, Littoral Environnement et Sociétés (LIENSs), Microbial Physiology Group, Université de La Rochelle, La Rochelle, France<sup>a</sup>; INRA, Virologie et Immunologie Moléculaires UR892, Université Paris-Saclay, Jouy-en-Josas, France<sup>b</sup>

**ABSTRACT** Some species of the genus *Cellulophaga* have been reported as having biotechnological interests and noteworthy physiological properties. We report here the draft genome sequence of *Cellulophaga lytica* CECT 8139, a bacterium that produces an intensely iridescent colony biofilm on agar surfaces.

The genus *Cellulophaga* is a member of the family *Flavobacteriaceae* in the phylum *Bacteroidetes*. *Cellulophaga* strains have been isolated from a variety of coastline biotopes, including the water column, sediments, seaweed or diatom surfaces, and marine invertebrates (1–3). Some reports have focused on potential biotechnological properties of *Cellulophaga* species, including antifouling and algicidal activities (4–6), inhibition of quorum sensing (7), or the production of novel enzymes involved in the bioconversion of algal polysaccharides (e.g., the breakdown of carrageenans or agar) (8, 9). Other studies described the development of genetic tools for understanding the gliding motility mechanisms or for identifying the components involved in the type IX secretion system (T9SS) (10, 11).

*Cellulophaga lytica* is a yellow/orange-pigmented aerobic agarolytic carrageenolytic rod and is motile by gliding (12). *C. lytica* CECT 8139 (CL8139) was isolated from a red sea anemone on the Charente Maritime Atlantic coast (France) (13). On agar, strain CL8139 produces a colony biofilm that displays an intensely “pointillistic” iridescence under epi-illumination. Under conditions that mimic its natural environment, the cell population shows a unique periodicity that induces this structural coloration (14–16). However, the cellular and molecular mechanisms of this self-organization, as well as its biological role, are still unknown.

The draft genome sequence of *C. lytica* CL8139 was obtained using HiSeq 2000 (Illumina) sequencing (GenoScreen, France), which generated 129,354,414 (2 × 100 bp) paired-end reads assembled *de novo* in 87 scaffolds (>1 kb) using Velvet. The total size for the combined scaffolds is 3,773,420 bp, with an average G+C content of 31.97%. Gene prediction and functional annotation were carried out using the MicroScope annotation platform (<https://www.genoscope.cns.fr/agc/microscope/home/index.php>) (17). The *C. lytica* CL8139 draft genome is predicted to contain 3,321 protein-coding genes, a coding density of 91.11%, and an average gene length of 1,031 bp. Strain CL8139 is the third *C. lytica* strain whose genome is sequenced (12, 18). Using average nucleotide identity (ANI) (19) as a proxy index for genomic relatedness, strain CL8139 displays 99.22% and 99.21% identities with the *C. lytica* type strain LIM21 and with strain HI1, respectively, suggesting a very close intraspecies phylogenetic proximity. The core genome of the species *C. lytica* is composed of 2,951 protein-coding genes (>80% identity over >80% of the size of the smallest protein), and strain CL8139 encompasses 232 strain-specific genes (7.0%) that essentially encode proteins of

Received 30 June 2017 Accepted 12 July 2017 Published 7 September 2017

**Citation** Chapelais-Baron M, Goubet I, Duchaud E, Rosenfeld E. 2017. Draft genome sequence of the iridescent marine bacterium *Cellulophaga lytica* CECT 8139. *Genome Announc* 5:e00811-17. <https://doi.org/10.1128/genomeA.00811-17>.

**Copyright** © 2017 Chapelais-Baron et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Eric Rosenfeld, [eric.rosenfeld@univ-lr.fr](mailto:eric.rosenfeld@univ-lr.fr).

E.D. and E.R. contributed equally to this article.

unknown function, Rhs and Vgr family proteins, and their remnants, transposases, or restriction-modification systems.

Genome analysis identified a clustered regularly interspaced short palindromic repeat (CRISPR)-encoding locus, a *dnd* gene cluster likely acquired by horizontal genetic transfer (20), genes encoding the gliding motility machinery and the T9SS, as well as proteins similar to the cold-adapted, thermotolerant, and denaturant-stable GH5 endoglucanase Celal\_2753 of *Cellulophaga algicola* IC166<sup>T</sup> (21) and to the kappa-carrageenase CgkA of *Zobellia galactanivorans* (22).

In contrast to other marine *Flavobacteria* species, strain CL8139 is devoid of (i) key quorum-sensing genes (i.e., *luxS* and *luxI*), (ii) gene encoding the acyl-L-homoserine lactonase (AHL) recently identified in *Tenacibaculum* species (23), and (iii) proteorhodopsin-encoding genes (24).

The genome of this environmental bacterium may help identify the genes required for the production of iridescent colony biofilms, improve knowledge on genome evolution, and facilitate future functional genomics studies.

**Accession number(s).** This whole-genome shotgun project has been deposited in ENA under the accession number [FZLT00000000](https://ENA.uniprot.org/record/FZLT00000000). The version reported in this work is the first version, FZLT01000000.

## ACKNOWLEDGMENTS

This work was financially supported by two CNRS grants, AIR75515 (“Bactéridescence” project) and 44162 (“Modiris in mobile” project), awarded to E.R. The Ph.D. thesis of M.C.-B. was supported by a grant from Région Poitou-Charentes (France). The LABGem (CEA/IG/Genoscope & CNRS UMR 8030) and the France Génomique National infrastructure (funded as part of Investissement d’Avenir program managed by Agence Nationale pour la Recherche, contract ANR-10-INBS-09) are acknowledged for support within the MicroScope annotation platform.

We are grateful to Jean-François Bernardet for critical reading of the manuscript.

## REFERENCES

- Lewin RA, Lounsbury DM. 1969. Isolation, cultivation and characterization of *Flexibacteria*. *J Gen Microbiol* 58:145–170. <https://doi.org/10.1099/00221287-58-2-145>.
- Bowman JP. 2000. Description of *Cellulophaga algicola* sp. nov., isolated from the surfaces of Antarctic algae, and reclassification of *Cytophaga uliginosa* (ZoBell and Upham 1944) Reichenbach 1989 as *Cellulophaga uliginosa* comb. nov. *Int J Syst Evol Microbiol* 50:1861–1868. <https://doi.org/10.1099/00207713-50-5-1861>.
- Kientz B, Agogue H, Lavergne C, Marié P, Rosenfeld E. 2013. Isolation and distribution of iridescent *Cellulophaga* and other iridescent marine bacteria from the Charente-Maritime coast, French Atlantic. *Syst Appl Microbiol* 36:244–251. <https://doi.org/10.1016/j.syapm.2013.02.004>.
- Sokolova A, Cilz N, Daniels J, Stafslie SJ, Brewer LH, Wendt DE, Bright FV, Detty MR. 2012. A comparison of the antifouling/foul-release characteristics of non-biocidal xerogel and commercial coatings toward micro- and macrofouling organisms. *Biofouling* 28:511–523. <https://doi.org/10.1080/08927014.2012.690197>.
- Majumdar P, Lee E, Patel N, Ward K, Stafslie SJ, Daniels J, Chisholm BJ, Boudjouk P, Callow ME, Callow JA, Thompson SEM. 2008. Combinatorial materials research applied to the development of new surface coatings IX: an investigation of novel antifouling/fouling-release coatings containing quaternary ammonium salt groups. *Biofouling* 24:185–200. <https://doi.org/10.1080/08927010801894660>.
- Skerratt JH, Bowman JP, Hallegraef G, James S, Nichols PD. 2002. Algalicidal bacteria associated with blooms of a toxic dinoflagellate in a temperate Australian estuary. *Mar Ecol Prog Ser* 244:1–15. <https://doi.org/10.3354/meps244001>.
- Lafleur JE, Costa SK, Bitzer AS, Silby MW. 2015. Draft genome sequence of *Cellulophaga* sp. E6, a marine algal epibiont that produces a quorum-sensing inhibitory compound active against *Pseudomonas aeruginosa*. *Genome Announc* 3(1):e01565-14. <https://doi.org/10.1128/genomeA.01565-14>.
- Shan D, Ying J, Li X, Gao Z, Wei G, Shao Z. 2014. Draft genome sequence of the carrageenan-degrading bacterium *Cellulophaga* sp. strain KL-A, isolated from decaying marine algae. *Genome Announc* 2(2):e00145-14. <https://doi.org/10.1128/genomeA.00145-14>.
- Lee YS, Choi YL. 2017. Complete genome sequence and analysis of three kinds of  $\beta$ -agarase of *Cellulophaga lytica* DAU203 isolated from marine sediment. *Mar Genomics*, in press. <https://doi.org/10.1016/j.margen.2017.05.005>.
- McBride MJ, Baker SA. 1996. Development of techniques to genetically manipulate members of the genera *Cytophaga*, *Flavobacterium*, *Flexibacter*, and *Sporocystophaga*. *Appl Environ Microbiol* 62:3017–3022.
- Zhu Y, McBride MJ. 2016. Comparative analysis of *Cellulophaga algicola* and *Flavobacterium johnsoniae* gliding motility. *J Bacteriol* 198:1743–1754. <https://doi.org/10.1128/JB.01020-15>.
- Pati A, Abt B, Teshima H, Nolan M, Lapidus A, Lucas S, Hammon N, Deshpande S, Cheng JF, Tapia R, Han C, Goodwin L, Pitluck S, Liolios K, Pagani I, Mavromatis K, Ovchinnikova G, Chen A, Palaniappan K, Land M, Hauser L, Jeffries CD, Detter JC, Brambilla EM, Kannan KP, Rohde M, Spring S, Göker M, Woyke T, Bristow J, Eisen JA, Markowitz V, Hugenholtz P, Kyrpides NC, Klenk HP, Ivanova N. 2011. Complete genome sequence of *Cellulophaga lytica* type strain (LIM-21). *Stand Genomic Sci* 4:221–232. <https://doi.org/10.4056/signs.1774329>.
- Kientz B, Vukusic P, Luke S, Rosenfeld E. 2012. Iridescence of a marine bacterium and classification of prokaryotic structural colors. *Appl Environ Microbiol* 78:2092–2099. <https://doi.org/10.1128/AEM.07339-11>.
- Kientz B, Ducret A, Luke S, Vukusic P, Mignot T, Rosenfeld E. 2012. Glitter-like iridescence within the *Bacteroidetes* especially *Cellulophaga* spp.: optical properties and correlation with gliding motility. *PLoS One* 7:e52900. <https://doi.org/10.1371/journal.pone.0052900>.
- Kientz B, Marié P, Rosenfeld E. 2012. Effect of abiotic factors on the unique glitter-like iridescence of *Cellulophaga lytica*. *FEMS Microbiol Lett* 333:101–108. <https://doi.org/10.1111/j.1574-6968.2012.02614.x>.
- Kientz B, Luke S, Vukusic P, Péteri R, Beaudry C, Renault T, Simon D, Mignot T, Rosenfeld E. 2016. A unique self-organization of bacterial sub-communities creates iridescence in *Cellulophaga lytica* colony biofilms. *Sci Rep* 6:19906. <https://doi.org/10.1038/srep19906>.

17. Vallenet D, Belda E, Calteau A, Cruveiller S, Engelen S, Lajus A, Le Fèvre F, Longin C, Mornico D, Roche D, Rouy Z, Salvignol G, Scarpelli C, Thil Smith AA, Weiman M, Médigue C. 2013. MicroScope—an integrated microbial resource for the curation and comparative analysis of genomic and metabolic data. *Nucleic Acids Res* 41:D636–D647. <https://doi.org/10.1093/nar/gks1194>.
18. Asahina AY, Hadfield MG. 2014. Complete genome sequence of *Cellulophaga lytica* HI1 using PacBio single-molecule real-time sequencing. *Genome Announc* 2(6):e01148-14. <https://doi.org/10.1128/genomeA.01148-14>.
19. Goris J, Konstantinidis KT, Klappenbach JA, Coenye T, Vandamme P, Tiedje JM. 2007. DNA-DNA hybridization values and their relationship to whole-genome sequence similarities. *Int J Syst Evol Microbiol* 57:81–91. <https://doi.org/10.1099/ijs.0.64483-0>.
20. Barbier P, Lunazzi A, Fujiwara-Nagata E, Avendaño-Herrera R, Bernardet JF, Touchon M, Ducaud E. 2013. From the *Flavobacterium* genus to the phylum *Bacteroidetes*: genomic analysis of *dnd* gene clusters. *FEMS Microbiol Lett* 348:26–35. <https://doi.org/10.1111/1574-6968.12239>.
21. Wang Y, Yu W, Han F. 2016. Expression and characterization of a cold-adapted, thermotolerant and denaturant-stable GH5 endoglucanase Celal\_2753 that withstands boiling from the psychrophilic bacterium *Cellulophaga algicola* IC166<sup>T</sup>. *Biotechnol Lett* 38:285–290. <https://doi.org/10.1007/s10529-015-1971-5>.
22. Barbeyron T, Gerard A, Potin P, Henrissat B, Kloareg B. 1998. The kappa-carrageenase of the marine bacterium *Cytophaga drobachiensis*. Structural and phylogenetic relationships within family-16 glycoside hydrolases. *Mol Biol Evol* 15:528–537. <https://doi.org/10.1093/oxfordjournals.molbev.a025952>.
23. Mayer C, Romero M, Muras A, Otero A. 2015. Aii20J, a wide-spectrum thermostable *N*-acylhomoserine lactonase from the marine bacterium *Tenacibaculum* sp. 20J, can quench AHL-mediated acid resistance in *Escherichia coli*. *Appl Microbiol Biotechnol* 99:9523–9539. <https://doi.org/10.1007/s00253-015-6741-8>.
24. Kwon YM, Kim SY, Jung KH, Kim SJ. 2016. Diversity and functional analysis of light-driven pumping rhodopsins in marine *Flavobacteria*. *Microbiol Open* 5:212–223. <https://doi.org/10.1002/mbo3.321>.