



HAL
open science

Resistance growth of branching random networks

Dayue Chen, Yueyun Hu, Shen Lin

► **To cite this version:**

Dayue Chen, Yueyun Hu, Shen Lin. Resistance growth of branching random networks. 2018. hal-01688502v1

HAL Id: hal-01688502

<https://hal.science/hal-01688502v1>

Preprint submitted on 19 Jan 2018 (v1), last revised 18 May 2018 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Resistance growth of branching random networks

Dayue Chen,^a Yueyun Hu^b and Shen Lin^c

December 28, 2017

Abstract

Consider a rooted infinite Galton–Watson tree with mean offspring number $m > 1$, and a collection of i.i.d. positive random variables ξ_e indexed by all the edges in the tree. We assign the resistance $m^d \xi_e$ to each edge e at distance d from the root. In this random electric network, we study the asymptotic behavior of the effective resistance and conductance between the root and the vertices at depth n . Our results generalize an existing work of Addario-Berry, Broutin and Lugosi on the binary tree to random branching networks.

Keywords. electric networks, Galton–Watson tree, random conductance.

AMS 2010 Classification Numbers. 60F05, 60J80.

1 Introduction

An electric network is an undirected locally finite connected graph $G = (V, E)$ with a countable set of vertices V and a set of edges E , endowed with nonnegative numbers $\{r(e), e \in E\}$, called resistances, that are associated to the edges of G . The reciprocal $c(e) = 1/r(e)$ is called the conductance of the edge e . It is well-known that the electrical properties of the network $(G, \{r(e)\})$ are closely related to the nearest-neighbor random walk on G , whose transition probabilities from a vertex are proportional to the conductances along the edges to be taken. See, for instance, the book of Lyons and Peres [11] for a detailed exposition of this connection.

To study random walks in certain random environments, it is natural to consider a random electric network by choosing the resistances independent and identically distributed. For example, the infinite cluster of bond percolation on \mathbb{Z}^d can be seen as a random electric network in which each open edge has unit resistance and each closed edge

^aDepartment of Probability and Statistics, School of Mathematical Sciences, Peking University, Beijing, China, *E-mail*: dayue@pku.edu.cn

^bLAGA, Université Paris XIII, Villetaneuse, France, *E-mail*: yueyun@math.univ-paris13.fr

^cLPSM, Sorbonne Université, Paris, France, *E-mail*: shen.lin.math@gmail.com

Cooperation between D.C. and Y.H. was supported by NSFC 11528101, S.L. is supported in part by the grant ANR-14-CE25-0014 (ANR GRAAL).

has infinite resistance. Grimmett, Kesten and Zhang [7] proved that when $d \geq 3$, the effective resistance of this network between a fixed point and infinity is a.s. finite, thus the simple random walk on this infinite percolation cluster is a.s. transient. In [3], Benjamini and Rossignol considered a different model of the cubic lattice \mathbb{Z}^d , where the resistance of each edge is an independent copy of a Bernoulli random variable. They showed that point-to-point effective resistance has submean variance in \mathbb{Z}^2 , whereas the mean and the variance are of the same order when $d \geq 3$. The case of a complete graph on n vertices has also been studied by Grimmett and Kesten [6]. For a particular class of resistance distribution on the edges (see Theorem 3 in [6]), as $n \rightarrow \infty$, the limit distribution of the random effective resistance between two specified vertices was identified as the sum of two i.i.d. random variables, each with the distribution of the effective resistance between the root and infinity in a Galton–Watson tree with a supercritical Poisson offspring distribution.

In this paper, we investigate the effective resistance and conductance in a supercritical Galton–Watson tree \mathbb{T} rooted at \emptyset . Let $\mathbf{p} = (p_k)_{k \geq 0}$ be the offspring distribution of \mathbb{T} , with finite mean $m > 1$. We assume $p_0 = 0$ to avoid the conditioning on survival. Formally, every vertex in \mathbb{T} can be represented as a finite word written with positive integers. The depth $|x|$ of a vertex x in \mathbb{T} is the number of edges on the unique non-self-intersecting path from the root \emptyset to x , which also equals the length of the word representing x . Let $\mathbb{T}_n := \{x \in \mathbb{T} : |x| = n\}$ denote the n -th level of \mathbb{T} . We write \overleftarrow{x} for the parent vertex of x if $x \neq \emptyset$. For each edge $e = \{\overleftarrow{x}, x\}$ of \mathbb{T} , we define its depth $d(e) := |x|$. Let ν be the number of children of the root, whose expected value is m . For $1 \leq i \leq \nu$, the edge $\{\emptyset, i\}$ between the root \emptyset and its child i has depth 1. If x and y are vertices of \mathbb{T} , we write $x \preceq y$ if x is on the non-self-intersecting path connecting \emptyset and y . In this case, we say that y is a descendant of x . We define $\mathbb{T}_n[x] := \{y \in \mathbb{T}_n : x \preceq y\}$ as the set of vertices at depth n that are descendants of x .

If the resistance of an edge at depth d equals λ^d with a deterministic $\lambda > 0$, Lyons [8] showed that the effective resistance between the root and infinity in \mathbb{T} is a.s. infinite if $\lambda > m$ and a.s. finite $\lambda < m$. The corresponding λ -biased random walk on \mathbb{T} is thus recurrent if $\lambda > m$, and transient if $\lambda < m$. For the critical value $\lambda = m$, we know by a subsequent work of Lyons [9] that the network still has an infinite effective resistance between the root and infinity. More precisely, the critical λ -biased random walk is null recurrent provided $\sum(k \log k)p_k < \infty$.

When the edges of \mathbb{T} have random resistances, we are mainly interested in the similar case of critical exponential weighting: to each edge e at depth $d(e)$, we assign the resistance

$$r(e) := m^{d(e)}\xi(e), \tag{1.1}$$

where, conditionally on \mathbb{T} , $\{\xi(e)\}$ are i.i.d. copies of a nonnegative random variable ξ . We will call $(\mathbb{T}, \{r(e)\})$ a branching random network of offspring distribution \mathbf{p} and electric resistance ξ . For convenience, we assume that $(\mathbb{T}, \{r(e)\})$ and ξ are independent and defined under the same probability measure \mathbb{P} .

Let R_n (resp. C_n) be the effective resistance (resp. effective conductance) between the root \emptyset and the vertices at depth n in $(\mathbb{T}, \{r(e)\})$. When \mathbb{T} is a deterministic binary tree,

Addario-Berry, Broutin and Lugosi [1] showed that as $n \rightarrow \infty$,

$$\mathbb{E}[R_n] = \mathbb{E}[\xi] n - \frac{\text{Var}[\xi]}{\mathbb{E}[\xi]} \log n + O(1) \quad \text{and} \quad \mathbb{E}[C_n] = \frac{1}{\mathbb{E}[\xi]} \frac{1}{n} + \frac{\text{Var}[\xi]}{\mathbb{E}[\xi]^3} \frac{\log n}{n^2} + O(n^{-2}),$$

provided ξ is bounded away from both zero and infinity. Their arguments are based on the concentration phenomenon of C_n and R_n when the underlying tree is regular. The Efron–Stein inequality is the main tool to deduce the following upper bounds on the variance

$$\text{Var}[R_n] = O(1) \quad \text{and} \quad \text{Var}[C_n] = O(n^{-4}).$$

A sub-Gaussian tail bound is also established for R_n , which gives

$$\mathbb{E}\left[|R_n - \mathbb{E}[R_n]|^k\right] = O(1) \quad \text{for all } k \geq 1.$$

As observed in the concluding remarks in [1], if the tree \mathbb{T} is random, C_n and R_n are no longer concentrated. For any nonnegative random variable X , we set $\{X\} := \frac{X}{\mathbb{E}[X]}$ whenever $0 < \mathbb{E}[X] < \infty$.

Theorem 1.1. *Assuming that $\mathbb{E}[\xi + \xi^{-1} + \nu^2] < \infty$, we have the almost convergence*

$$\{C_n\} \xrightarrow[n \rightarrow \infty]{} W, \tag{1.2}$$

where $W := \lim_{n \rightarrow \infty} m^{-n} \#\mathbb{T}_n$.

We write $W_n := m^{-n} \#\mathbb{T}_n$. When $\mathbb{E}[\nu^2] < \infty$, it is well-known that $(W_n)_{n \geq 1}$ is an L^2 -bounded martingale. The convergence $W_n \rightarrow W$ holds in the L^2 -sense, and the limit W is almost surely strictly positive, with

$$\mathbb{E}[W] = 1 \quad \text{and} \quad \mathbb{E}[W^2] = \frac{\sum k^2 p_k - m}{m(m-1)}.$$

Similarly, for each vertex $x \in \mathbb{T}$, the random variable

$$W^{(x)} := \lim_{n \rightarrow \infty} m^{|x|-n} \#\mathbb{T}_n[x]$$

has the same distribution as W . Using the tree notation $|x| = n$ to denote a vertex x at depth n , we have $W = m^{-n} \sum_{|x|=n} W^{(x)}$.

Theorem 1.1 answers some questions mentioned at the end of [1]. When the offspring number ν is not deterministic, it implies that the limit distribution of $\{C_n\}$ is absolutely continuous with respect to the Lebesgue measure, which is a “scaled analogue” of Question 4.1 in Lyons, Pemantle and Peres [10]. For the absolute continuity of W , see for instance Theorem 10.4 in Chapter 1 of [2].

For our next result, let us define

$$a_1 := m^{-2} \mathbb{E}[\nu(\nu-1)], \tag{1.3}$$

$$b_1 := \mathbb{E}[\xi],$$

$$c_1 := \frac{a_1 b_1}{1 - m^{-1}}. \tag{1.4}$$

Notice that by Theorems 22 and 23 in Dubuc [5], $\mathbb{E}[W^{-1}] < \infty$ if and only if $p_1 m < 1$.

Theorem 1.2. *Assuming that $\mathbb{E}[\xi^2 + \xi^{-1} + \nu^3] < \infty$, we have*

$$\lim_{n \rightarrow \infty} n \mathbb{E}[C_n] = \frac{1}{c_1}. \quad (1.5)$$

If additionally $p_1 m < 1$, then

$$\lim_{n \rightarrow \infty} \frac{\mathbb{E}[R_n]}{n} = c_1 \mathbb{E}\left[\frac{1}{W}\right].$$

If $p_1 m \geq 1$, by Fatou's lemma, we deduce from (1.2) and (1.5) that

$$\liminf_{n \rightarrow \infty} \frac{\mathbb{E}[R_n]}{n} = \infty.$$

See also the remark at the end of Section 3.

To state a more precise asymptotic expansion for $\mathbb{E}[C_n]$, we define

$$a_2 := m^{-3} \mathbb{E}\left[\nu(\nu-1)(\nu-2)\mathbf{1}_{\{\nu \geq 2\}}\right], \quad (1.6)$$

$$b_2 := \mathbb{E}[\xi^2],$$

$$c_2 := (1 - m^{-2})^{-1} \left(\frac{3a_1^2}{m-1} + a_2 \right), \quad (1.7)$$

$$c_3 := \frac{2a_1 c_1}{m-1} - \frac{2b_1 c_2}{m}, \quad (1.8)$$

$$c_4 := \frac{b_1}{1 - m^{-1}} \left(\frac{c_3}{c_1} + a_1 \right) - b_2 \frac{c_2}{c_1}. \quad (1.9)$$

If $\nu = m \geq 2$ is deterministic,

$$c_1 = b_1 = \mathbb{E}[\xi], \quad c_2 = 1, \quad c_3 = 0 \quad \text{and} \quad c_4 = b_1 - \frac{b_2}{b_1} = -\frac{\text{Var}[\xi]}{\mathbb{E}[\xi]}.$$

Theorem 1.3. *Assume that $\mathbb{E}[\xi^3 + \xi^{-1} + \nu^4] < \infty$. Then there exists a constant $c_0 \in \mathbb{R}$ such that, as $n \rightarrow \infty$,*

$$\mathbb{E}[C_n] = \frac{1}{c_1 n} - \frac{c_4 \log n}{c_1^2 n^2} - \frac{c_0}{c_1^2} \frac{1}{n^2} + O\left(\frac{(\log n)^2}{n^3}\right).$$

The constant c_0 appearing in the expansion above will be defined at the end of Section 4, but its explicit value is unknown to us.

To further describe the rate of convergence in (1.2), we write $\xi_x := \xi(\{\overleftarrow{x}, x\})$ for every vertex $x \neq \emptyset$. Remark that, conditioning on the first ℓ levels of the tree \mathbb{T} , the random variables $W^{(x)}, |x| = \ell$ are i.i.d. and independent of $\xi_x, |x| = \ell$. Notice that $W^{(x)}(1 - \frac{\xi_x}{c_1} W^{(x)})$ is of zero mean, because $c_1 = \mathbb{E}[\xi] \mathbb{E}[W^2]$. When $\mathbb{E}[\xi^2 + \nu^4] < \infty$, one can easily verify that

$$\sum_{\ell=1}^{\infty} \frac{1}{m^\ell} \sum_{|x|=\ell} W^{(x)} \left(1 - \frac{\xi_x}{c_1} W^{(x)} \right)$$

converges in L^2 .

Theorem 1.4. *Assuming that $\mathbb{E}[\xi^3 + \xi^{-1} + \nu^4] < \infty$, we have*

$$n(\{C_n\} - W) \xrightarrow[n \rightarrow \infty]{(P)} \sum_{\ell=1}^{\infty} \frac{1}{m^\ell} \sum_{|x|=\ell} W^{(x)} \left(1 - \frac{\xi_x}{c_1} W^{(x)}\right),$$

and, with the same constant c_0 in Theorem 1.3,

$$R_n - \left(\frac{c_1}{W} n + \frac{c_4}{W} \log n + \frac{1}{W} \left(c_0 - \frac{1}{W} \sum_{\ell=1}^{\infty} \frac{1}{m^\ell} \sum_{|x|=\ell} W^{(x)} \left(c_1 - \xi_x W^{(x)} \right) \right) \right) \xrightarrow[n \rightarrow \infty]{(P)} 0, \quad (1.10)$$

where $\xrightarrow{(P)}$ indicates convergence in probability.

The rest of the paper is organized as follows. In the next section, we recall Thomson's principle for the effective resistance, and we derive the recurrence relation for C_n . In Section 3, we collect some estimates on the moments of C_n . The convergence (1.5) and Theorem 1.3 will be shown in Section 4 by analyzing the recurrence equations on the moments of C_n . Similar arguments have already been used in the proof of Theorem 5 in [1]. By second moment calculations, we establish Theorems 1.1 and 1.4 in Section 5, and, by proving the uniform integrability of $(n^{-1}R_n)_{n \geq 1}$, we complete the proof of Theorem 1.2 in Section 6. Finally, in Section 7 we briefly discuss the case when we change the scaling by assigning to each edge e in \mathbb{T} the resistance $\lambda^{d(e)}\xi(e)$ with $\lambda > m$.

2 Preliminaries

Consider a general network $G = (V, E)$ with the resistances $\{r(e)\}$. For $x, y \in V$, we write $x \sim y$ to indicate that $\{x, y\}$ belongs to E . To each edge $e = \{x, y\}$, one may associate two directed edges $\vec{x\dot{y}}$ and $\vec{y\dot{x}}$. We shall denote by \vec{E} the set of all directed edges. A flow θ is a function on \vec{E} that is antisymmetric, meaning that $\theta(\vec{x\dot{y}}) = -\theta(\vec{y\dot{x}})$. The divergence of θ at a vertex x is defined by

$$\operatorname{div} \theta(x) := \sum_{y: y \sim x} \theta(\vec{x\dot{y}}).$$

Let A and Z be two disjoint non-empty subsets of V : A will represent the source of the network and Z the sink. The flow θ is from A to Z with strength $\|\theta\|$ if it satisfies Kirchhoff's node law that $\operatorname{div} \theta(x) = 0$ for all $x \notin A \cup Z$, and that

$$\|\theta\| = \sum_{a \in A} \sum_{y \sim a, y \notin A} \theta(\vec{a\dot{y}}) = \sum_{z \in Z} \sum_{y \sim z, y \notin Z} \theta(\vec{y\dot{z}}).$$

The effective resistance between A and Z can be defined as

$$R(A \leftrightarrow Z) := \inf_{\|\theta\|=1} \sum_{e \in E} r(e)\theta(e)^2, \quad (2.1)$$

where the infimum is taken over all flows θ from A to Z with unit strength. The infimum is always attained at what is called the unit current flow, which satisfies, in addition to

the node law, Kirchhoff's cycle law. This flow-based formulation of the effective resistance is also called Thomson's principle. The effective conductance $C(A \leftrightarrow Z)$ between A and Z is the reciprocal $R(A \leftrightarrow Z)^{-1}$.

Conditionally on the branching random network $(\mathbb{T}, \{r(e)\})$, let X be the associated random walk on the tree \mathbb{T} . Let $\omega(x, y), x \sim y$ denote the transition probabilities of X , and let $\pi(x), x \in \mathbb{T}$ denote the reversible measure. Writing the conductances $c(e) = 1/r(e)$, we have

$$\pi(x) = \sum_{y: y \sim x} c(\{x, y\}) \quad \text{and} \quad \omega(x, y) = \frac{c(\{x, y\})}{\pi(x)}.$$

We suppose that the random walk X starts from the vertex x at time 0 under the probability measure $P_{x, \omega}$. As a probabilistic interpretation, the effective conductance $C_n := C(\{\emptyset\} \leftrightarrow \mathbb{T}_n)$ between the root and the level set $\{x \in \mathbb{T} : |x| = n\}$ satisfies

$$C_n = \pi(\emptyset) P_{\emptyset, \omega}(\tau_n < T_\emptyset^+),$$

where $\tau_n := \inf\{k \geq 0 : |X_k| = n\}$ and $T_\emptyset^+ := \inf\{k \geq 1 : X_k = \emptyset\}$. We see immediately that $C_n \geq C_{n+1}$.

For $1 \leq i \leq \nu$, let $C_{n+1, i} := C(\{i\} \leftrightarrow \mathbb{T}_{n+1}[i])$ denote the effective conductance between the vertex i and $\mathbb{T}_{n+1}[i]$. We also set $\eta_i := \xi(\{\emptyset, i\})^{-1}$, $1 \leq i \leq \nu$, which are i.i.d., independent of ν . Observe that conditioning on ν , $(C_{n+1, i})_{1 \leq i \leq \nu}$ are i.i.d., independent of η_i , and distributed as $\frac{C_n}{m}$. Using the series and parallel law of electric networks, we obtain the recurrence relation that for $n \geq 1$,

$$C_{n+1} = \sum_{i=1}^{\nu} \left(\frac{m}{\eta_i} + \frac{1}{C_{n+1, i}} \right)^{-1} = \frac{1}{m} \sum_{i=1}^{\nu} \frac{\eta_i C_n^{(i)}}{\eta_i + C_n^{(i)}}, \quad (2.2)$$

where for $1 \leq i \leq \nu$, $C_n^{(i)} := m C_{n+1, i}$ are i.i.d. copies of C_n , independent of $(\eta_i)_{1 \leq i \leq \nu}$. It is clear that $C_1 = m^{-1} \sum_{i=1}^{\nu} \eta_i$. If we set $\xi_i := \xi(\{\emptyset, i\}) = \eta_i^{-1}$ for $1 \leq i \leq \nu$, the recurrence equation (2.2) can also be written as

$$C_{n+1} = \frac{1}{m} \sum_{i=1}^{\nu} \frac{C_n^{(i)}}{1 + \xi_i C_n^{(i)}}. \quad (2.3)$$

3 Bounds on the expected conductance

Let η denote the reciprocal ξ^{-1} .

Lemma 3.1. *If $\mathbb{E}[\eta] = \mathbb{E}[\xi^{-1}] < \infty$, then $\mathbb{E}[C_n] \leq \frac{\mathbb{E}[\eta]}{n}$ for all $n \geq 1$.*

Proof. First of all, $\mathbb{E}[C_1] = \mathbb{E}[\eta]$. From (2.2) we obtain for all $n \geq 1$ that

$$\mathbb{E}[C_{n+1}] = \mathbb{E} \left[\frac{\eta C_n}{\eta + C_n} \right].$$

By concavity of the function $x \mapsto \frac{xy}{x+y}$, $y > 0$ being fixed,

$$\mathbb{E} \left[\frac{\eta C_n}{\eta + C_n} \right] \leq \mathbb{E} \left[\frac{\eta \mathbb{E}[C_n]}{\eta + \mathbb{E}[C_n]} \right] \leq \frac{\mathbb{E}[\eta] \mathbb{E}[C_n]}{\mathbb{E}[\eta] + \mathbb{E}[C_n]},$$

It follows that $(\mathbb{E}[C_{n+1}])^{-1} \geq (\mathbb{E}[\eta])^{-1} + (\mathbb{E}[C_n])^{-1} \geq \dots \geq (n+1)(\mathbb{E}[\eta])^{-1}$. \square

Lemma 3.2. *Assume that $\mathbb{E}[\eta] = \mathbb{E}[\xi^{-1}] < \infty$. For $2 \leq k \leq 4$, if $\mathbb{E}[\nu^k] < \infty$, then*

$$\mathbb{E}[(C_n)^k] = O(n^{-k}) \quad \text{as } n \rightarrow \infty.$$

Proof. Starting from (2.2), we obtain

$$\mathbb{E}[(C_{n+1})^2] = \frac{1}{m^2} \mathbb{E}[\nu] \mathbb{E} \left[\left(\frac{\eta C_n}{\eta + C_n} \right)^2 \right] + \frac{\mathbb{E}(\nu(\nu - 1))}{m^2} (\mathbb{E}[C_{n+1}])^2,$$

by developing the square and using the independence after conditioning on ν . Together with Lemma 3.1, it follows that

$$\mathbb{E}[(C_{n+1})^2] \leq \frac{1}{m} \mathbb{E}[C_n^2] + \frac{\mathbb{E}[\nu(\nu - 1)]}{m^2} (\mathbb{E}[C_{n+1}])^2 \leq \frac{1}{m} \mathbb{E}[C_n^2] + \frac{\mathbb{E}[\nu(\nu - 1)]}{m^2} \frac{(\mathbb{E}[\eta])^2}{(n+1)^2}.$$

Since $m > 1$, we get $\mathbb{E}[C_n^2] = O(n^{-2})$ by induction. Furthermore, if $\mathbb{E}[\nu^3] < \infty$, by developing the third power and using the independence,

$$\begin{aligned} \mathbb{E}[(C_{n+1})^3] &= \mathbb{E} \left[\left(\frac{1}{m} \sum_{i=1}^{\nu} \frac{\eta_i C_n^{(i)}}{\eta_i + C_n^{(i)}} \right)^3 \right] \\ &\leq \frac{1}{m^2} \mathbb{E} \left[\left(\frac{\eta C_n}{\eta + C_n} \right)^3 \right] + \frac{3\mathbb{E}[\nu^2]}{m^3} \mathbb{E} \left[\left(\frac{\eta C_n}{\eta + C_n} \right)^2 \right] \mathbb{E} \left[\frac{\eta C_n}{\eta + C_n} \right] + \frac{\mathbb{E}[\nu^3]}{m^3} \left(\mathbb{E} \left[\frac{\eta C_n}{\eta + C_n} \right] \right)^3 \\ &\leq \frac{1}{m^2} \mathbb{E}[C_n^3] + \frac{3\mathbb{E}[\nu^2]}{m^3} \mathbb{E}[C_n^2] \mathbb{E}[C_n] + \frac{\mathbb{E}[\nu^3]}{m^3} (\mathbb{E}[C_n])^3. \end{aligned}$$

Thus, $\mathbb{E}[C_n^3] = O(n^{-3})$ follows from $\mathbb{E}[C_n] = O(n^{-1})$ and $\mathbb{E}[C_n^2] = O(n^{-2})$. The last bound $\mathbb{E}[C_n^4] = O(n^{-4})$ is similarly obtained by assuming that $\mathbb{E}[\nu^4] < \infty$. \square

Lemma 3.3. *If $\mathbb{E}[\xi] \in (0, \infty)$ and $\mathbb{E}[\nu^2] < \infty$, then there exists a constant $c > 0$ such that $\mathbb{E}[C_n] \geq \frac{c}{n}$ for all $n \geq 1$.*

In the following proof, we will use the uniform flow on \mathbb{T} to give an upper bound for $R_n = C_n^{-1}$. Similar arguments can be found in Lemma 2.2 of Pemantle and Peres [12].

Proof. We define on \mathbb{T} the uniform flow Θ_{unif} of unit strength (with the source $\{\emptyset\}$) by setting

$$\Theta_{\text{unif}}(\{\overleftarrow{x}, x\}) = m^{-|x|} \frac{W^{(x)}}{W} \quad \text{for every } x \in \mathbb{T} \setminus \{\emptyset\}.$$

According to Thomson's principle (2.1),

$$R_n \leq \sum_{k=1}^n \sum_{|x|=k} m^k \xi_x \Theta_{\text{unif}}(\{\overleftarrow{x}, x\})^2 = \sum_{k=1}^n \sum_{|x|=k} m^{-k} \xi_x \left(\frac{W^{(x)}}{W} \right)^2. \quad (3.1)$$

We write $A := \sup_{k \geq 1} m^{-k} \#\mathbb{T}_k$, which is square integrable by L^2 -maximal inequality of Doob. It follows that

$$\frac{R_n}{n} \leq \frac{1}{n} \sum_{k=1}^n \frac{A}{W^2} \left(\frac{1}{\#\mathbb{T}_k} \sum_{|x|=k} \xi_x (W^{(x)})^2 \right).$$

Using Proposition 2.3 in [12], a variant of the strong law of large numbers for exponentially growing blocks of identically distributed random variables being independent inside each block, we have

$$\frac{1}{\#\mathbb{T}_k} \sum_{|x|=k} \xi_x (W^{(x)})^2 \xrightarrow[k \rightarrow \infty]{\text{a.s.}} \mathbb{E}[\xi] \mathbb{E}[W^2].$$

Hence, almost surely

$$\limsup_{n \rightarrow \infty} \frac{R_n}{n} \leq A \mathbb{E}[\xi] \frac{\mathbb{E}[W^2]}{W^2},$$

which yields

$$\liminf_{n \rightarrow \infty} n C_n \geq (A \mathbb{E}[\xi])^{-1} \frac{W^2}{\mathbb{E}[W^2]},$$

Taking expectation and using Fatou's lemma, we obtain

$$\liminf_{n \rightarrow \infty} n \mathbb{E}[C_n] \geq \frac{\mathbb{E}[W^2 A^{-1}]}{\mathbb{E}[\xi] \mathbb{E}[W^2]} > 0.$$

The proof is thus completed. \square

Remark. The Nash-Williams inequality (see Section 2.5 in [11]) gives the lower bound

$$R_n \geq \sum_{k=1}^n \left(\sum_{d(e)=k} r(e)^{-1} \right)^{-1} = \sum_{k=1}^n \left(\sum_{|x|=k} m^{-k} (\xi_x)^{-1} \right)^{-1}.$$

Suppose that $\mathbb{E}[\xi^{-1}] < \infty$. Proposition 2.3 in [12] implies that

$$\frac{1}{\#\mathbb{T}_k} \sum_{|x|=k} (\xi_x)^{-1} \xrightarrow[k \rightarrow \infty]{\text{a.s.}} \mathbb{E}[\xi^{-1}].$$

With the almost sure convergence $m^{-k} \#\mathbb{T}_k \rightarrow W$, it follows that

$$\frac{1}{n} \sum_{k=1}^n \left(\sum_{|x|=k} m^{-k} (\xi_x)^{-1} \right)^{-1} \xrightarrow[n \rightarrow \infty]{\text{a.s.}} \frac{1}{W \mathbb{E}[\xi^{-1}]}.$$

By Fatou's lemma, we obtain

$$\liminf_{n \rightarrow \infty} \frac{\mathbb{E}[R_n]}{n} \geq \mathbb{E} \left[\liminf_{n \rightarrow \infty} \frac{R_n}{n} \right] \geq \frac{\mathbb{E}[W^{-1}]}{\mathbb{E}[\xi^{-1}]}.$$

The integrability of W^{-1} is therefore a necessary condition for having $\mathbb{E}[R_n] = O(n)$.

4 Asymptotic expansion of the expected conductance

Within this section, let the assumption $\mathbb{E}[\xi^2 + \xi^{-1} + \nu^3] < \infty$ be always in force. We first establish (1.5) in Theorem 1.2. Afterwards we will prove Theorem 1.3 under the stronger assumption that $\mathbb{E}[\xi^3 + \xi^{-1} + \nu^4] < \infty$.

For every integer $n \geq 1$, we write

$$x_n := \mathbb{E}[C_n], \quad y_n := \mathbb{E}[C_n^2], \quad z_n := \mathbb{E}[C_n^3].$$

By Lemma 3.2, we have $x_n = O(n^{-1})$, $y_n = O(n^{-2})$ and $z_n = O(n^{-3})$.

Observe from (2.3) that $\mathbb{E}[C_{n+1}] = \mathbb{E}\left[\frac{C_n}{1+\xi C_n}\right]$ with ξ and C_n being independent. Then developing the power of C_{n+1} , we arrive at

$$\mathbb{E}[C_{n+1}^2] = \frac{1}{m} \mathbb{E}\left[\left(\frac{C_n}{1+\xi C_n}\right)^2\right] + \frac{\mathbb{E}[\nu(\nu-1)]}{m^2} (\mathbb{E}[C_{n+1}])^2 = \frac{1}{m} \mathbb{E}\left[\left(\frac{C_n}{1+\xi C_n}\right)^2\right] + a_1 (\mathbb{E}[C_{n+1}])^2$$

and

$$\begin{aligned} \mathbb{E}[C_{n+1}^3] &= \frac{1}{m^2} \mathbb{E}\left[\left(\frac{C_n}{1+\xi C_n}\right)^3\right] + \frac{3\mathbb{E}[\nu(\nu-1)]}{m^3} \mathbb{E}\left[\left(\frac{C_n}{1+\xi C_n}\right)^2\right] \mathbb{E}\left[\frac{C_n}{1+\xi C_n}\right] \\ &\quad + m^{-3} \mathbb{E}\left[\sum_{1 \leq i, j, k \leq \nu} \mathbf{1}_{\{i \neq j \neq k\}}\right] \left(\mathbb{E}\left[\frac{C_n}{1+\xi C_n}\right]\right)^3 \\ &= \frac{1}{m^2} \mathbb{E}\left[\left(\frac{C_n}{1+\xi C_n}\right)^3\right] + \frac{3a_1}{m} \mathbb{E}\left[\left(\frac{C_n}{1+\xi C_n}\right)^2\right] \mathbb{E}\left[\frac{C_n}{1+\xi C_n}\right] + a_2 \left(\mathbb{E}\left[\frac{C_n}{1+\xi C_n}\right]\right)^3, \end{aligned}$$

with the constants a_1, a_2 defined as in (1.3) and (1.6).

Using the identity $\frac{1}{1+x} = 1 - x + \frac{x^2}{1+x}$, we obtain

$$\begin{aligned} \mathbb{E}\left[\frac{C_n}{1+\xi C_n}\right] &= \mathbb{E}[C_n] - \mathbb{E}[\xi] \mathbb{E}[C_n^2] + \mathbb{E}\left[\frac{\xi^2 C_n^3}{1+\xi C_n}\right] \\ &= \mathbb{E}[C_n] - \mathbb{E}[\xi] \mathbb{E}[C_n^2] + O(n^{-3}), \end{aligned}$$

because $\mathbb{E}[C_n^3] = O(n^{-3})$ and $\mathbb{E}[\xi^2] < \infty$. Similarly,

$$\mathbb{E}\left[\left(\frac{C_n}{1+\xi C_n}\right)^2\right] = \mathbb{E}[C_n^2] + O(n^{-3}).$$

Hence, we have

$$x_{n+1} = x_n - b_1 y_n + O(n^{-3}), \tag{4.1}$$

$$y_{n+1} = \frac{y_n}{m} + a_1 x_{n+1}^2 + O(n^{-3}) = \frac{y_n}{m} + a_1 x_n^2 + O(n^{-3}). \tag{4.2}$$

Remark that

$$x_{n+1} = \mathbb{E}\left[\frac{C_n}{1+\xi C_n}\right] \geq \mathbb{E}[C_n] - \mathbb{E}[\xi] \mathbb{E}[C_n^2] = x_n - b_1 y_n.$$

Since $x_n \geq \frac{c}{n}$ by Lemma 3.3 and $y_n = O(n^{-2})$, we get $\frac{x_n}{x_{n+1}} \leq 1 + \frac{C}{n}$ for some positive constant C independent of n . It follows that for any $i < n/2$,

$$1 \leq \frac{x_{n-i}}{x_n} \leq \prod_{j=n-i}^{n-1} \left(1 + \frac{C}{j}\right) \leq \exp\left(Ci/(n-i)\right) \leq 1 + C' \frac{i}{n} \tag{4.3}$$

with another constant $C' > 0$.

Still by Lemma 3.3, we can divide all terms in (4.1) by $x_n x_{n+1}$, which leads to

$$\frac{1}{x_{n+1}} - \frac{1}{x_n} = b_1 \frac{y_n}{x_n x_{n+1}} + O(n^{-1}). \quad (4.4)$$

By induction, (4.2) implies that

$$y_{n+1} = a_1 \sum_{i=0}^{n-1} m^{-i} x_{n-i}^2 + O(n^{-3}).$$

Using (4.3), we deduce that

$$\frac{y_n}{x_n x_{n+1}} = a_1 \sum_{i=0}^{\infty} m^{-i} + O(n^{-1}) = \frac{a_1}{1-m^{-1}} + O(n^{-1}).$$

It follows from (4.4) that

$$\frac{1}{x_{n+1}} - \frac{1}{x_n} = \frac{a_1 b_1}{1-m^{-1}} + O(n^{-1}) = c_1 + O(n^{-1}), \quad (4.5)$$

with the constant c_1 defined in (1.4). Consequently,

$$\frac{1}{x_n} = c_1 n + O(\log n), \quad (4.6)$$

which gives the convergence (1.5).

Assuming from now on that $\mathbb{E}[\xi^3 + \xi^{-1} + \nu^4] < \infty$, we proceed to find higher-order asymptotic expansions for x_n . Using the identity $\frac{1}{1+x} = 1 - x + x^2 - \frac{x^3}{1+x}$, we obtain

$$\begin{aligned} \mathbb{E} \left[\frac{C_n}{1 + \xi C_n} \right] &= \mathbb{E}[C_n] - \mathbb{E}[\xi] \mathbb{E}[C_n^2] + \mathbb{E}[\xi^2] \mathbb{E}[C_n^3] - \mathbb{E} \left[\frac{\xi^3 C_n^4}{1 + \xi C_n} \right] \\ &= \mathbb{E}[C_n] - \mathbb{E}[\xi] \mathbb{E}[C_n^2] + \mathbb{E}[\xi^2] \mathbb{E}[C_n^3] + O(n^{-4}), \end{aligned}$$

as $\mathbb{E}[\xi^3] < \infty$ and $\mathbb{E}[C_n^4] = O(n^{-4})$ by Lemma 3.2. We prove in the same manner that

$$\begin{aligned} \mathbb{E} \left[\left(\frac{C_n}{1 + \xi C_n} \right)^2 \right] &= \mathbb{E}[C_n^2] - 2 \mathbb{E}[\xi] \mathbb{E}[C_n^3] + O(n^{-4}), \\ \mathbb{E} \left[\left(\frac{C_n}{1 + \xi C_n} \right)^3 \right] &= \mathbb{E}[C_n^3] + O(n^{-4}). \end{aligned}$$

Hence, we deduce that

$$x_{n+1} = x_n - b_1 y_n + b_2 z_n + O(n^{-4}), \quad (4.7)$$

$$\begin{aligned} y_{n+1} &= \frac{y_n}{m} + a_1 x_{n+1}^2 - \frac{2b_1}{m} z_n + O(n^{-4}) \\ &= \frac{y_n}{m} + a_1 x_n^2 - \left(2a_1 b_1 x_n y_n + \frac{2b_1}{m} z_n \right) + O(n^{-4}), \end{aligned} \quad (4.8)$$

$$\begin{aligned} z_{n+1} &= \frac{z_n}{m^2} + \frac{3a_1}{m} x_{n+1} y_n + a_2 x_{n+1}^3 + O(n^{-4}) \\ &= \frac{z_n}{m^2} + \frac{3a_1}{m} x_n y_n + a_2 x_n^3 + O(n^{-4}). \end{aligned} \quad (4.9)$$

Dividing all terms in (4.9) by x_{n+1}^3 gives

$$\frac{z_{n+1}}{x_{n+1}^3} = \frac{x_n^3}{x_{n+1}^3} \left(\frac{1}{m^2} \frac{z_n}{x_n^3} + \frac{3a_1}{m} \frac{y_n}{x_n^2} + a_2 + O(n^{-1}) \right).$$

Recall that $\frac{x_n}{x_{n+1}} = 1 + O(n^{-1})$ by (4.3). Hence,

$$\frac{z_{n+1}}{x_{n+1}^3} = \frac{1}{m^2} \frac{z_n}{x_n^3} + \frac{3a_1}{m} \frac{y_n}{x_n^2} + a_2 + O(n^{-1}).$$

Since

$$\frac{y_n}{x_n^2} = \frac{a_1}{1 - m^{-1}} + O(n^{-1}), \quad (4.10)$$

we get by induction that

$$\frac{z_{n+1}}{x_{n+1}^3} = \sum_{i=0}^{n-1} m^{-2i} \left(\frac{3a_1}{m} \frac{y_{n-i}}{x_{n-i}^2} + a_2 \right) + O(n^{-1}) \xrightarrow{n \rightarrow \infty} \sum_{i=0}^{\infty} m^{-2i} \left(\frac{3a_1}{m} \frac{a_1}{1 - m^{-1}} + a_2 \right) = c_2, \quad (4.11)$$

with the constant c_2 defined in (1.7).

Dividing all terms in (4.8) by x_{n+1}^2 gives

$$\frac{y_{n+1}}{x_{n+1}^2} = \frac{x_n^2}{x_{n+1}^2} \left(\frac{y_n}{mx_n^2} + a_1 - 2a_1b_1 \frac{y_n}{x_n} - \frac{2b_1}{m} \frac{z_n}{x_n^2} \right) + O(n^{-2}). \quad (4.12)$$

For every $n \geq 1$, define

$$\begin{aligned} \varepsilon_n &:= \frac{1}{x_{n+1}} - \frac{1}{x_n} - c_1, \\ \delta_n &:= \frac{y_{n+1}}{x_{n+1}^2} - \frac{y_n}{mx_n^2} - a_1. \end{aligned}$$

It has been shown that $\varepsilon_n = O(n^{-1})$. Putting

$$\frac{x_n^2}{x_{n+1}^2} = \left(1 + (c_1 + \varepsilon_n)x_n \right)^2 = 1 + 2c_1x_n + O(n^{-2})$$

into (4.12), we see that

$$\delta_n = 2a_1c_1x_n + \left(\frac{2c_1}{m} - 2a_1b_1 \right) \frac{y_n}{x_n} - \frac{2b_1}{m} \frac{z_n}{x_n^2} + O(n^{-2}).$$

By (4.10) and (4.11), it follows that

$$\frac{\delta_n}{x_n} \xrightarrow{n \rightarrow \infty} 2a_1c_1 + \frac{a_1}{1 - m^{-1}} \left(\frac{2c_1}{m} - 2a_1b_1 \right) - \frac{2b_1c_2}{m} = c_3,$$

with the constant c_3 defined in (1.8). In view of (4.6), $\delta_n = \frac{c_3}{c_1} \frac{1}{n} + o(n^{-1})$. If we set

$$\Delta_{n+1} := \frac{y_{n+1}}{x_{n+1}^2} - \frac{a_1}{1 - m^{-1}},$$

then $\Delta_{n+1} = \frac{1}{m}\Delta_n + \delta_n$ by the definition of δ_n . Hence, $\Delta_{n+1} = m^{-n}\Delta_1 + \sum_{i=0}^{n-1} m^{-i}\delta_{n-i}$ by induction, and

$$n\Delta_n \xrightarrow{n \rightarrow \infty} \frac{c_3}{c_1(1-m^{-1})}. \quad (4.13)$$

Going back to (4.7), we obtain by the definition of ε_n that

$$\begin{aligned} c_1 + \varepsilon_n &= \frac{1}{x_{n+1}} \left(1 - \frac{x_{n+1}}{x_n}\right) \\ &= \frac{x_n}{x_{n+1}} \left(b_1 \frac{y_n}{x_n^2} - b_2 \frac{z_n}{x_n^2}\right) + O(n^{-2}) \\ &= (1 + (c_1 + \varepsilon_n)x_n) \left(b_1 \frac{y_n}{x_n^2} - b_2 \frac{z_n}{x_n^2}\right) + O(n^{-2}) \\ &= b_1 \frac{y_n}{x_n^2} + c_1 b_1 \frac{y_n}{x_n} - b_2 \frac{z_n}{x_n^2} + O(n^{-2}). \end{aligned}$$

As $c_1 = \frac{a_1 b_1}{1-m^{-1}}$, we deduce that

$$\varepsilon_n = b_1 \Delta_n + c_1 b_1 \frac{y_n}{x_n} - b_2 \frac{z_n}{x_n^2} + O(n^{-2}). \quad (4.14)$$

Together with (4.13), (4.10), (4.11) and (4.6), it implies that

$$n\varepsilon_n \xrightarrow{n \rightarrow \infty} b_1 \frac{c_3}{c_1(1-m^{-1})} + b_1 \frac{a_1}{1-m^{-1}} - b_2 \frac{c_2}{c_1} = c_4,$$

with the constant c_4 defined in (1.9). Therefore,

$$\frac{1}{x_n} - \frac{1}{x_1} = c_1(n-1) + \sum_{i=1}^{n-1} \varepsilon_i = c_1 n + c_4 \log n + o(\log n),$$

yielding that

$$x_n = \frac{1}{c_1 n} + O\left(\frac{\log n}{n^2}\right). \quad (4.15)$$

To get an error term $O(1)$ instead of $o(\log n)$ in the expansion of $\frac{1}{x_n}$, we first notice that (4.11) can be improved as

$$\frac{z_{n+1}}{x_{n+1}^3} = c_2 + O(n^{-1}). \quad (4.16)$$

Using it together with (4.10) and (4.15), we derive from

$$\delta_n = x_n \left(2a_1 c_1 + \left(\frac{2c_1}{m} - 2a_1 b_1\right) \frac{y_n}{x_n^2} - \frac{2b_1}{m} \frac{z_n}{x_n^3}\right) + O(n^{-2})$$

that $\delta_n = \frac{c_3}{c_1} \frac{1}{n} + O(n^{-2} \log n)$. It follows that

$$\Delta_{n+1} = m^{-n}\Delta_1 + \sum_{i=0}^{n-1} m^{-i}\delta_{n-i} = \frac{c_3}{c_1(1-m^{-1})} \frac{1}{n} + O(n^{-2} \log n).$$

Reformulating (4.14) as

$$\varepsilon_n = b_1 \Delta_n + x_n \left(c_1 b_1 \frac{y_n}{x_n^2} - b_2 \frac{z_n}{x_n^3} \right) + O(n^{-2}),$$

and using (4.15), (4.10) and (4.16), we get that

$$\varepsilon_n = \frac{c_4}{n} + O(n^{-2} \log n),$$

which implies the absolute convergence of $\sum_{i=1}^{\infty} (\varepsilon_i - \frac{c_4}{i})$. Hence,

$$\frac{1}{x_n} = \frac{1}{x_1} + c_1(n-1) + \sum_{i=1}^{n-1} \varepsilon_i = c_1 n + c_4 \log n + c_0 + o(1),$$

with the constant

$$c_0 := -c_1 + \frac{1}{x_1} + \sum_{i=1}^{\infty} \left(\varepsilon_i - \frac{c_4}{i} \right) = -c_1 + \mathbb{E}[\eta]^{-1} + \sum_{i=1}^{\infty} \left(\varepsilon_i - \frac{c_4}{i} \right).$$

Finally we have

$$\mathbb{E}[C_n] = x_n = \frac{1}{c_1 n} - \frac{c_4 \log n}{c_1^2 n^2} - \frac{c_0}{c_1^2 n^2} + O\left(\frac{(\log n)^2}{n^3}\right). \quad (4.17)$$

5 Almost sure convergence and rate of convergence

To prove Theorems 1.1 and 1.4, let us write

$$\begin{aligned} Y_n &:= \{C_n\} - W, \\ \Pi_n &:= C_n \left(\frac{1}{x_{n+1}} - \frac{1}{x_n} - \frac{1}{x_{n+1}} \frac{\xi C_n}{1 + \xi C_n} \right). \end{aligned}$$

For every vertex $x \in \mathbb{T}$ and $j \geq 1$, we also define

$$\begin{aligned} C_j^{(x)} &:= m^{|x|} C(\{x\} \leftrightarrow \mathbb{T}_{j+|x|}[x]), \\ Y_j^{(x)} &:= \{C_j^{(x)}\} - W^{(x)}, \\ \Pi_j^{(x)} &:= C_j^{(x)} \left(c_1 + \varepsilon_j - \frac{1}{x_{j+1}} \frac{\xi_x C_j^{(x)}}{1 + \xi_x C_j^{(x)}} \right). \end{aligned}$$

Using (2.3), we have

$$\{C_n\} = \frac{1}{x_n} \frac{1}{m} \sum_{i=1}^{\nu} \frac{C_{n-1}^{(i)}}{1 + \xi_i C_{n-1}^{(i)}} = \frac{1}{m} \sum_{i=1}^{\nu} \{C_{n-1}^{(i)}\} + \frac{1}{m} \sum_{i=1}^{\nu} \Pi_{n-1}^{(i)},$$

Using the simple equality $W = m^{-1} \sum_{i=1}^{\nu} W^{(i)}$, we deduce that

$$Y_n = \frac{1}{m} \sum_{i=1}^{\nu} Y_{n-1}^{(i)} + \frac{1}{m} \sum_{i=1}^{\nu} \Pi_{n-1}^{(i)}.$$

Since $W = m^{-k} \sum_{|x|=k} W^{(x)}$, by induction,

$$Y_n = \frac{1}{m^k} \sum_{|x|=k} Y_{n-k}^{(x)} + \sum_{\ell=1}^k \frac{1}{m^\ell} \sum_{|y|=\ell} \Pi_{n-\ell}^{(y)} \quad \text{for any } 1 \leq k < n.$$

Proof of Theorem 1.1. Assume that $\mathbb{E}[\xi + \xi^{-1} + \nu^2] < \infty$. Notice that our proof preceding (4.3) to establish $\frac{x_n}{x_{n+1}} = 1 + O(n^{-1})$ is still valid. Besides, $y_n = \mathbb{E}[C_n^2] = O(n^{-2})$ by Lemma 3.2, and $\frac{y_n}{x_{n+1}} = O(n^{-1})$ by Lemma 3.3. Hence, we derive from the inequality

$$\mathbb{E}[|\Pi_n|] \leq x_n \left(\frac{1}{x_{n+1}} - \frac{1}{x_n} \right) + \frac{1}{x_{n+1}} \mathbb{E} \left[\frac{\xi (C_n)^2}{1 + \xi C_n} \right] \leq \frac{x_n}{x_{n+1}} - 1 + \frac{y_n}{x_{n+1}} \mathbb{E}[\xi]$$

that $\mathbb{E}[|\Pi_n|] \leq \frac{C}{n}$ with some constant $C > 0$.

Conditioning on the first k levels of the tree \mathbb{T} , $(Y_{n-k}^{(x)}, |x| = k)$ are i.i.d. copies of Y_{n-k} . Using the fact that Y_n is of zero mean and uniformly bounded in L^2 , we can find a constant $C' > 0$ such that

$$\mathbb{E} \left[\left(\frac{1}{m^k} \sum_{|x|=k} Y_{n-k}^{(x)} \right)^2 \right] = m^{-k} \mathbb{E} \left[(Y_{n-k})^2 \right] \leq C' m^{-k}. \quad (5.1)$$

Meanwhile,

$$\mathbb{E} \left[\sum_{\ell=1}^k \frac{1}{m^\ell} \sum_{|y|=\ell} |\Pi_{n-\ell}^{(y)}| \right] \leq \sum_{\ell=1}^k \frac{C}{n-\ell} \leq \frac{Ck}{n-k}.$$

It follows that

$$\mathbb{E}[|Y_n|] \leq \sqrt{C' m^{-k}} + \frac{Ck}{n-k}.$$

By taking $k = C'' \log n$ for some constant C'' sufficiently large, we see that

$$\mathbb{E}[|Y_n|] = O\left(\frac{\log n}{n}\right).$$

Choose a subsequence $n_j = j^2$. Borel–Cantelli’s lemma gives that Y_{n_j} converges to 0 almost surely. The monotonicity of C_n shows that for any $n_j \leq n < n_{j+1}$,

$$\frac{x_{n_{j+1}}}{x_{n_j}} \cdot \{C_{n_{j+1}}\} \leq \{C_n\} \leq \frac{x_{n_j}}{x_{n_{j+1}}} \cdot \{C_{n_j}\}.$$

By (4.3), the almost sure convergence of Y_n readily follows. \square

Together with (4.6), Theorem 1.1 implies that

$$nC_n \xrightarrow[n \rightarrow \infty]{\text{a.s.}} \frac{W}{c_1}, \quad (5.2)$$

provided $\mathbb{E}[\xi^2 + \xi^{-1} + \nu^3] < \infty$.

Proof of Theorem 1.4. Assume now $\mathbb{E}[\xi^3 + \xi^{-1} + \nu^4] < \infty$. First, observe that taking the subsequence $k_n = \frac{4}{\log m} \log n$ in (5.1) yields

$$n \left(\frac{1}{m^{k_n}} \sum_{|x|=k_n} Y_{n-k_n}^{(x)} \right) \xrightarrow{n \rightarrow \infty} 0 \quad \text{in } L^2.$$

By Borel–Cantelli’s lemma, the preceding convergence also holds in the almost sure sense. We claim that

$$\sum_{\ell=1}^{k_n} \frac{1}{m^\ell} \sum_{|y|=\ell} n \Pi_{n-\ell}^{(y)} \xrightarrow[n \rightarrow \infty]{(P)} \sum_{\ell=1}^{\infty} \frac{1}{m^\ell} \sum_{|y|=\ell} W^{(y)} \left(1 - \frac{\xi_y}{c_1} W^{(y)} \right). \quad (5.3)$$

In fact, for each vertex y at fixed depth ℓ ,

$$n C_{n-\ell}^{(y)} \xrightarrow[n \rightarrow \infty]{\text{a.s.}} \frac{W^{(y)}}{c_1} \quad \text{and} \quad n \Pi_{n-\ell}^{(y)} \xrightarrow[n \rightarrow \infty]{\text{a.s.}} W^{(y)} \left(1 - \frac{\xi_y}{c_1} W^{(y)} \right).$$

So for any integer $K \geq 1$,

$$\sum_{\ell=1}^K \frac{1}{m^\ell} \sum_{|y|=\ell} n \Pi_{n-\ell}^{(y)} \xrightarrow[n \rightarrow \infty]{\text{a.s.}} \sum_{\ell=1}^K \frac{1}{m^\ell} \sum_{|y|=\ell} W^{(y)} \left(1 - \frac{\xi_y}{c_1} W^{(y)} \right).$$

Note that

$$\mathbb{E} \left[\left(\frac{1}{m^\ell} \sum_{|y|=\ell} n \Pi_{n-\ell}^{(y)} \right)^2 \right] \leq m^{-\ell} n^2 \mathbb{E}[\Pi_{n-\ell}^2] + \frac{\mathbb{E}[(\#\mathbb{T}_\ell)^2]}{m^{2\ell}} n^2 \left(\mathbb{E}[\Pi_{n-\ell}] \right)^2.$$

On the one hand,

$$\begin{aligned} \mathbb{E}[\Pi_n^2] &\leq 2 \left(\frac{1}{x_{n+1}} - \frac{1}{x_n} \right)^2 \mathbb{E}[C_n^2] + \frac{2}{(x_{n+1})^2} \mathbb{E} \left[\frac{\xi^2 C_n^4}{(1 + \xi C_n)^2} \right] \\ &\leq 2 \left(\frac{1}{x_{n+1}} - \frac{1}{x_n} \right)^2 \mathbb{E}[C_n^2] + \frac{2}{(x_{n+1})^2} \mathbb{E}[\xi^2] \mathbb{E}[C_n^4]. \end{aligned}$$

Using (4.5) and the facts that x_n is of order n^{-1} , $\mathbb{E}[C_n^2] = O(n^{-2})$ and $\mathbb{E}[C_n^4] = O(n^{-4})$, we deduce that $\mathbb{E}[\Pi_n^2] = O(n^{-2})$. On the other hand,

$$\begin{aligned} \mathbb{E}[\Pi_n] &= \frac{x_n}{x_{n+1}} - 1 - \frac{1}{x_{n+1}} \mathbb{E}[\xi C_n^2] + \frac{1}{x_{n+1}} \mathbb{E}[\xi^2 C_n^3] - \frac{1}{x_{n+1}} \mathbb{E} \left[\frac{\xi^3 C_n^4}{1 + \xi C_n} \right] \\ &= \frac{x_n}{x_{n+1}} - 1 - \frac{1}{x_{n+1}} b_1 y_n + \frac{1}{x_{n+1}} b_2 z_n + O(n^{-3}). \end{aligned}$$

It follows by (4.7) that $\mathbb{E}[\Pi_n] = O(n^{-3})$. In particular, $\mathbb{E}[\Pi_{n-\ell}] = O(n^{-3})$ for any $\ell = o(n)$. Besides, $m^{-2\ell} \mathbb{E}[(\#\mathbb{T}_\ell)^2]$ is uniformly bounded in ℓ . Hence, there exists some constant $\tilde{C} > 0$ so that

$$\mathbb{E} \left[\left(\frac{1}{m^\ell} \sum_{|y|=\ell} n \Pi_{n-\ell}^{(y)} \right)^2 \right] \leq \tilde{C} m^{-\ell} + \tilde{C} n^{-4} \quad \text{for all } \ell \leq k_n.$$

It follows that

$$\lim_{K \rightarrow \infty} \limsup_{n \rightarrow \infty} \left\| \sum_{\ell=K}^{k_n} \frac{1}{m^\ell} \sum_{|y|=\ell} n \Pi_{n-\ell}^{(y)} \right\|_{L^2} = 0,$$

which yields (5.3). Therefore,

$$nY_n = n \left(\frac{1}{m^{k_n}} \sum_{|x|=k_n} Y_{n-k_n}^{(x)} \right) + \sum_{\ell=1}^{k_n} \frac{1}{m^\ell} \sum_{|y|=\ell} n \Pi_{n-\ell}^{(y)} \xrightarrow[n \rightarrow \infty]{(P)} \sum_{\ell=1}^{\infty} \frac{1}{m^\ell} \sum_{|y|=\ell} W^{(y)} \left(1 - \frac{\xi_y}{c_1} W^{(y)} \right).$$

In view of (4.17), we have

$$n^2 C_n - \left(\frac{W}{c_1} n - \frac{c_4 W}{c_1^2} \log n - \frac{c_0 W}{c_1^2} + \frac{1}{c_1} \sum_{\ell=1}^{\infty} \frac{1}{m^\ell} \sum_{|y|=\ell} W^{(y)} \left(1 - \frac{\xi_y}{c_1} W^{(y)} \right) \right) \xrightarrow[n \rightarrow \infty]{(P)} 0,$$

and the convergence (1.10) follows immediately. \square

6 The expected resistance

When $\mathbb{E}[\xi^2 + \xi^{-1} + \nu^3] < \infty$, it follows from (5.2) that

$$\frac{R_n}{n} \xrightarrow[n \rightarrow \infty]{\text{a.s.}} \frac{c_1}{W}.$$

The following lemma yields the uniform integrability of $(\frac{R_n}{n}, n \geq 1)$, and completes the proof of Theorem 1.2.

Lemma 6.1. *Suppose that $p_1 m < 1$ and $\mathbb{E}[\xi^r + \nu^{2r}] < \infty$ for some $r > 1$. Then there exists some $s > 1$ such that*

$$\sup_{n \geq 1} \mathbb{E} \left[\left(\frac{R_n}{n} \right)^s \right] < \infty.$$

Proof. As $p_1 m < 1$, by Theorems 22 and 23 in Dubuc [5], there is some $\alpha > 1$ such that

$$\mathbb{E}[W^{-\alpha}] < \infty.$$

In fact, we may take any $\alpha \in (1, -\frac{\log p_1}{\log m})$, with the convention that $-\frac{\log p_1}{\log m} = +\infty$ if $p_1 = 0$. Moreover, $\mathbb{E}[\nu^{2r}] < \infty$ implies that $\mathbb{E}[W^{2r}] < \infty$, according to Bingham and Doney [4].

Recall that the martingale $W_k = m^{-k} \#\mathbb{T}_k$ converges in L^1 to W . Let

$$\mathcal{F}_k := \sigma\{\#\mathbb{T}_i, i \leq k\}, \quad k \geq 0$$

denote the natural filtration associated to $(W_k)_{k \geq 0}$. Since $W_k = \mathbb{E}[W | \mathcal{F}_k]$, it follows from Jensen's inequality that $(W_k)^{-\alpha} \leq \mathbb{E}[W^{-\alpha} | \mathcal{F}_k]$. Consequently,

$$\sup_{k \geq 1} \mathbb{E} \left[(W_k)^{-\alpha} \right] < \infty. \tag{6.1}$$

Fix an arbitrary $s \in (1, r \wedge \alpha)$. By convexity, we deduce from (3.1) that

$$\begin{aligned} \left(\frac{R_n}{n}\right)^s &\leq \frac{1}{n} \sum_{k=1}^n \left(\sum_{|x|=k} m^{-k} \xi_x \left(\frac{W^{(x)}}{W}\right)^2 \right)^s \\ &\leq \frac{1}{n} \sum_{k=1}^n (\#\mathbb{T}_k)^{s-1} \sum_{|x|=k} m^{-ks} (\xi_x)^s \left(\frac{W^{(x)}}{W}\right)^{2s}. \end{aligned}$$

Since $\mathbb{E}[\xi^s] < \infty$, the proof boils down to showing that

$$\sup_{k \geq 1} \mathbb{E} \left[(\#\mathbb{T}_k)^{s-1} \sum_{|x|=k} m^{-ks} \left(\frac{W^{(x)}}{W}\right)^{2s} \right] < \infty. \quad (6.2)$$

Recall that $W = \sum_{|x|=k} m^{-k} W^{(x)}$, and conditioning on \mathcal{F}_k , $(W^{(x)})_{|x|=k}$ are i.i.d. copies of W . Let $\phi(u) := -\log \mathbb{E}[e^{-uW}]$ for any $u \geq 0$. Using the elementary identity

$$a^{-2s} = \frac{1}{\Gamma(2s)} \int_0^\infty t^{2s-1} e^{-at} dt \quad \text{for any } a > 0,$$

we get that for any vertex x at depth k ,

$$\begin{aligned} \mathbb{E} \left[\left(\frac{W^{(x)}}{W}\right)^{2s} \middle| \mathcal{F}_k \right] &= \frac{1}{\Gamma(2s)} \int_0^\infty dt t^{2s-1} \mathbb{E} \left[(W^{(x)})^{2s} e^{-t \sum_{|y|=k} m^{-k} W^{(y)}} \middle| \mathcal{F}_k \right] \\ &= \frac{1}{\Gamma(2s)} \int_0^\infty dt t^{2s-1} e^{-(\#\mathbb{T}_k-1)\phi(tm^{-k})} \mathbb{E} \left[W^{2s} e^{-tm^{-k}W} \right] \\ &= \frac{1}{\Gamma(2s)} m^{2ks} \int_0^\infty du u^{2s-1} e^{-(\#\mathbb{T}_k-1)\phi(u)} \mathbb{E} \left[W^{2s} e^{-uW} \right]. \end{aligned}$$

It follows that

$$\begin{aligned} I_k &:= \mathbb{E} \left[(\#\mathbb{T}_k)^{s-1} \sum_{|x|=k} m^{-ks} \left(\frac{W^{(x)}}{W}\right)^{2s} \right] \\ &= \frac{1}{\Gamma(2s)} m^{ks} \int_0^\infty du u^{2s-1} \mathbb{E} \left[(\#\mathbb{T}_k)^s e^{-(\#\mathbb{T}_k-1)\phi(u)} \right] \mathbb{E} \left[W^{2s} e^{-uW} \right]. \end{aligned} \quad (6.3)$$

For any $a > 0$, we claim that there exists some positive constant $C = C(a, s) > 0$ such that for any $k \geq 1$,

$$m^{ks} \mathbb{E} \left[(\#\mathbb{T}_k)^s e^{-a(\#\mathbb{T}_k-1)} \right] \leq C. \quad (6.4)$$

Indeed, by discussing whether $\#\mathbb{T}_k \geq k^2$ or not, we have

$$m^{ks} \mathbb{E} \left[(\#\mathbb{T}_k)^s e^{-a\#\mathbb{T}_k} \right] \leq m^{ks} \sup_{y \geq k^2} y^s e^{-ay} + m^{ks} k^{2s} \mathbb{P}(\#\mathbb{T}_k < k^2).$$

The first term in the right-hand side is uniformly bounded, while

$$m^{ks} k^{2s} \mathbb{P}(\#\mathbb{T}_k < k^2) \leq m^{ks} k^{2s+2\alpha} \mathbb{E} \left[(\#\mathbb{T}_k)^{-\alpha} \right].$$

Note that $\mathbb{E}[(\#T_k)^{-\alpha}] = O(m^{-\alpha k})$ by (6.1). Since $s < \alpha$, we obtain (6.4).

Going back to the right-hand side of (6.3), we split the integral \int_0^∞ into two parts \int_0^1 and \int_1^∞ . For the part \int_1^∞ we apply (6.4) with $a = \phi(1)$, and for the part \int_0^1 we dominate $\mathbb{E}[W^{2s}e^{-uW}]$ by $\mathbb{E}[W^{2s}]$, to arrive at

$$I_k \leq \frac{C}{\Gamma(2s)} \int_1^\infty du u^{2s-1} \mathbb{E}[W^{2s}e^{-uW}] + C' m^{ks} \int_0^1 du u^{2s-1} \mathbb{E}[(\#T_k)^s e^{-\#T_k \phi(u)}],$$

with the constant

$$C' := \frac{e^{\phi(1)} \mathbb{E}[W^{2s}]}{\Gamma(2s)} < \infty.$$

Notice that by Fubini's theorem and a change of variables $v = uW$,

$$\int_1^\infty du u^{2s-1} \mathbb{E}[W^{2s}e^{-uW}] \leq \int_0^\infty du u^{2s-1} \mathbb{E}[W^{2s}e^{-uW}] = \Gamma(2s).$$

To treat the integral from 0 to 1, we remark that $\lim_{u \rightarrow 0} \frac{\phi(u)}{u} = \mathbb{E}[W] = 1$. Then there exists some positive constant c , such that $\phi(u) \geq \frac{u}{c}$ for all $0 \leq u \leq 1$. It follows that

$$\begin{aligned} I_k &\leq C + C' m^{ks} \int_0^1 du u^{2s-1} \mathbb{E}[(\#T_k)^s e^{-\frac{u}{c} \#T_k}] \\ &= C + C' \mathbb{E} \left[\int_0^{\#T_k} dv (W_k)^{-s} v^{2s-1} e^{-\frac{v}{c}} \right] \\ &\leq C + C' c^{2s} \Gamma(2s) \mathbb{E}[(W_k)^{-s}]. \end{aligned}$$

Using again (6.1) we get that $\sup_{k \geq 1} I_k < \infty$, yielding (6.2) and completing the proof. \square

7 General exponential weighting

Given the Galton–Watson tree \mathbb{T} and $\lambda > 0$, one can do the λ -exponential weighting of resistance by assigning the resistance $\lambda^{d(e)} \xi(e)$ to each edge e at depth $d(e)$. As before, conditionally on \mathbb{T} , $\{\xi(e)\}$ are i.i.d. positive random variables. In this random electric network, let $C_n(\lambda)$ denote the effective conductance between the root and the vertices at depth n . Instead of (2.3), the recurrence equation now reads as

$$C_{n+1}(\lambda) = \frac{1}{\lambda} \sum_{i=1}^{\nu} \frac{C_n^{(i)}(\lambda)}{1 + \xi_i C_n^{(i)}(\lambda)},$$

where for $1 \leq i \leq \nu$, $C_n^{(i)}(\lambda)$ are i.i.d. copies of $C_n(\lambda)$, independent of $(\xi_i)_{1 \leq i \leq \nu}$.

Theorem 7.1. *Fix $\lambda > m$. Assuming that $\mathbb{E}[\xi + \xi^{-1} + \nu^2] < \infty$, we have*

$$\{C_n(\lambda)\} \xrightarrow[n \rightarrow \infty]{\text{a.s.}} W.$$

If $\mathbb{E}[\xi^2 + \xi^{-1} + \nu^3] < \infty$, then, as $n \rightarrow \infty$, the limit of

$$\left(\frac{\lambda}{m}\right)^n \mathbb{E}[C_n(\lambda)]$$

exists and is strictly positive.

Basically the proof of Theorem 7.1 goes along the same lines as Theorem 1.1 and that of (1.5), except a few minor modifications. We leave the details to the reader.

References

- [1] L. ADDARIO-BERRY, N. BROUTIN AND G. LUGOSI. Effective resistance of random trees. *Ann. Appl. Probab.* **19** (2009), 1092–1107.
- [2] K. ATHREYA, P. NEY, *Branching Processes*. Die Grundlehren der mathematischen Wissenschaften, Band 196, Springer–Verlag, New York–Heidelberg, 1972. xi+287 pp.
- [3] I. BENJAMINI AND R. ROSSIGNOL. Submean variance bound for effective resistance on random electric networks. *Commun. Math. Phys.* **280** (2008), 445–462.
- [4] N. BINGHAM AND R. DONEY. Asymptotic properties of supercritical branching processes I: The Galton–Watson process. *Adv. Appl. Probab.* **6** (1974), 711–731.
- [5] S. DUBUC. Problèmes relatifs à l’itération de fonctions suggérés par les processus en cascade. *Ann. Inst. Fourier, Grenoble* **21** (1971), 171–251.
- [6] G. GRIMMETT AND H. KESTEN. Random electrical networks on complete graphs. *J. London Math. Soc. (2)* **30** (1984), 171–192.
- [7] G. GRIMMETT, H. KESTEN AND Y. ZHANG. Random walk on the infinite cluster of the percolation model. *Probab. Theory Relat. Fields* **96** (1993), 33–44.
- [8] R. LYONS. Random walks and percolation on trees. *Ann. Probab.* **18** (1990), 931–958.
- [9] R. LYONS. Random walks, capacity and percolation on trees. *Ann. Probab.* **20** (1992), 2043–2088.
- [10] R. LYONS, R. PEMANTLE AND Y. PERES. Unsolved problems concerning random walks on trees. *IMA Vol. Math. Appl.* **84** (1997), 223–237.
- [11] R. LYONS AND Y. PERES. *Probability on Trees and Networks*. Cambridge University Press, New York, 2016, xv+699 pp.
- [12] R. PEMANTLE AND Y. PERES. Galton–Watson trees with the same mean have the same polar sets. *Ann. Probab.* **23** (1995), 1102–1124.