

Table 1: Localization, physico-chemical and biological characteristics of the six derelict soils. Means \pm standard deviations (n=5, except F where n=4). Means with different letters are significantly different ($P < 0.05$) using one-way ANOVA or Kruskal-Wallis test. Localization in Lambert 93 projection in m; age is the time lapse from the last anthropogenic action (years); clay and WHC in %, CEC in $\text{cmol}^+ \cdot \text{kg}^{-1}$ dry soil; Olsen Phosphorous in $\text{mg} \cdot \text{kg}^{-1}$ dry soil; bacterial and fungal density in 16S and 18S rDNA gene copy numbers. g^{-1} dry soil, respectively; bacterial and fungal diversity are expressed as ribotype numbers; mycorrhizal colonization is the frequency of colonized roots (%); Collembola and macrofauna density are the numbers of individuals. m^{-2} ; Collembola, macrofauna and plant diversity are taxonomic richness; plant biomass is in kg of shoot dry weight. m^{-2} ; the relative density of GP and GN PAH-RHD α are the ratio between GP or GN PAH-dioxygenase genes and the 16S rDNA gene copy number. g^{-1} dry soil respectively in %; the proportion of the meso- and macro-decomposers is in %; AMC in $\text{ppm CO}_2 \cdot \text{mg}^{-1}$ dry soil. h^{-1} . Statistical test: 1 for one-way ANOVA test; 2 for Kruskal-Wallis test.

| | A | B | C | D | E | F | | |
|--------------------------------|--|---|-------------------------------|--------------------------------------|--------------------------------|--|------------------|---------|
| Site type and name | Experimental constructed soil (Biotechnosol) | Non-hazardous waste landfill (Retomfey) | Coking plant site (Homécourt) | Settling pond site (Moyeuvre-Petite) | Coking plant site (Micheville) | Experimental constructed and contaminated soil (Jeandelaincourt) | Statistical test | P-value |
| Localization | E:918202 N:6905686 | E:942352 N:6897630 | E:918487 N:6905891 | E:920222 N:6911528 | E:942262 N:6897475 | E:938477 N:6864617 | | |
| Age | 8 | 4 | 35 | 34 | 40 | 2 | | |
| Texture | Silt loam | Silty clay loam | Loamy sand | Sandy loamy | Loam | Silt loam | | |
| Clay | 11 | 40 | 6 | 4 | 27 | 15 | | |
| Water holding capacity (WHC) | 95±7 a | 59±8 b | 63±5 b | 58±4 b | 70±1 b | 64±1 b | 1 | 0.02 |
| Cation exchange capacity (CEC) | 22.0±2.9 a | 17.2±3.3 b | 17.2±1.8 b | 11.3±0.7 c | 15.5±1.4 b | 10.1±0.4 c | 2 | 0.01 |
| pH | 7.9±0.2 a | 7.9±0.1 a | 7.9±0.1 a | 8.4±0.2 b | 7.2±0.2 c | 8.3±0.4 b | 2 | <0.001 |
| Olsen phosphorous | 62±5 a | 25±17 c | 39±16 bc | 44±10 b | 36±2 bc | 44±10 bc | 2 | <0.001 |
| C:N | 30±6 bc | 28±6 c | 26±5 c | 65±20 a | 20±3 e | 37±16 b | 2 | <0.001 |
| Organic matter (%) | 39.7±5.9 a | 3.2±1.4 c | 16.7±6.3 b | 13.3±2.8 b | 4.8±3.0 c | 2.8±0.8 c | 2 | <0.001 |
| Ca _{exchangeable} | 292±44 d | 713±144 c | 903±405 c | 2031±240 a | 786±84 c | 1123±130 b | 2 | <0.001 |
| K _{exchangeable} | 706±105 a | 491±24 b | 155±36 e | 377±30 c | 131±50 e | 218±36 d | 2 | <0.001 |
| Mg _{exchangeable} | 84±14 cd | 152±38 b | 103±6 c | 312±47 a | 47±15 e | 72±34 d | 2 | <0.001 |
| N _o | 27-10 | 24-20 | 11-21 | 12-11 | 11-14 | 55-20 | 2 | <0.001 |

| | | | | | | | | | |
|---|---|---------------|---------------|---------------|---------------|---------------|---------------|---|--------|
| Pollutants mg kg ⁻¹ dry soil | Cd _{available} | 0.22±0.05 b | 0.03±0.01 e | 0.16±0.05 c | 0.15±0.01 c | 0.10±0.03 d | 9.07±1.5 a | 2 | <0.001 |
| | Pb _{available} | 13±3 b | 14±10 b | 46±14 a | 18±5 b | 14±6 b | 110±60 a | 2 | <0.001 |
| | Zn _{available} | 29±9 c | 8±6 d | 45±20 ab | 25±6 c | 7±2 d | 308±189 a | 2 | <0.001 |
| | Cd _{total} | 1.1±0.5 b | 0.1±0.2 b | 0.1±0.7 b | 1.4±0.1 b | 0.1±0.1 b | 23.6±6.9 a | 2 | <0.001 |
| | Pb _{total} | 150±53 c | 41±26 d | 346±106 ab | 339±50 b | 37±10 d | 460±73 a | 2 | <0.001 |
| | Zn _{total} | 345±37 c | 131±2 d | 1162±396 b | 1196±122 b | 116±23 d | 1813±302 a | 2 | <0.001 |
| | Σ16 PAH (US-EPA) | 170±49 a | 12±12 b | 179±136 a | 97±54 a | 142±90 a | 10±4 b | 2 | 0.03 |
| Microorganisms | Bacterial density ($\times 10^{10}$) | 5.4±1.2 a | 2.2±0.7 b | 4.8±0.7 a | 1.4±0.5 b | 2.6±1 b | 1.5±0.4 b | 1 | <0.001 |
| | Bacterial richness | 9.6±0.9 d | 13.2±1.8 ab | 15.2±0.8 a | 15.8±1.9 ab | 14.0±3.5 bc | 9.0±2.4 cd | 1 | <0.001 |
| | Fungal density ($\times 10^8$) | 10.9±4.5 a | 3.4±0.7 b | 4.9±0.6 b | 1.7±0.5 c | 5.1±1.2 b | 1.2±0.4 c | 2 | <0.001 |
| | Fungal richness | 3.6±0.9 b | 4±1.2 b | 7.4±1.1 a | 4.4±0.5 b | 3.6±0.5 b | 3.3±1.5 b | 1 | <0.001 |
| | Mycorrhizal colonization | 76±20 ab | 92±8 a | 86±12 ab | 82±9 ab | 96±4 a | 71±7 b | 1 | 0.04 |
| Fauna | Collembola density ($\times 10^3$) | 21±15 c | 17±8 c | 48±29 a | 91±41 a | 22±10 bc | 41±22 ab | 2 | 0.004 |
| | Collembola richness | 3.4±1.3 b | 4.2±0.8 ab | 6±1.4 ab | 7±0.8 a | 7±2.2 a | 5.8±1 a | 1 | 0.005 |
| | Macrofauna density | 611±219 a | 668±205 a | 314±138 b | 214±37 bc | 118±108 c | 205±15 bc | 1 | <0.001 |
| | Macrofauna richness | 10.2±2 a | 11.6±4 a | 6.8±3 b | 6.4±1 b | 3.0±2 c | 6.5±1 b | 2 | 0.001 |
| Flora | Plant biomass | 3.0±1.2 a | 3.2±1.2 a | 0.6±0.2 b | 0.7±0.4 b | 0.7±0.4 b | 0.6±0.2 b | 2 | <0.001 |
| | Plant richness | 2.2±0.8 b | 4.4±0.9 b | 10.2±1.6 a | 7.2±1.8 a | 13.0±3.7 a | 12.3±1 a | 2 | <0.001 |
| Functional parameters mU g ⁻¹ dry soil | FDA hydrolysis | 4.3±1.0 a | 2.1±0.6 b | 1.8±0.6 b | 1.0±0.6 b | 2.8±0.7 b | 2.7±0.9 ab | 1 | <0.001 |
| | Alkaline phosphatase activity | 1.7±0.4 a | 1.3±0.3 ab | 1.6±0.4 a | 0.9±0.4 b | 0.5±0.2 b | 1.0±0.2 ab | 1 | <0.001 |
| | Urease activity | 175±55 a | 75±32 b | 172±47 a | 78±6 b | 155±20 a | 71±21 b | 2 | 0.002 |
| | Average mineralization capacity (AMC) | 2654±501 a | 1714±521 b | 1880±289 b | 820±395 c | 879±247 cd | 380±96 d | 1 | <0.001 |
| | Relative density of GP PAH-RHD α genes | 1.2±0.3 a | 0.3±0.1 c | 0.7±0.1 bc | 1±0.4 ab | 0.9±0.2 ab | 0.3±0.1 c | 1 | <0.001 |
| | Relative density of GN PAH-RHD α genes | 0.010±0.003 a | 0.001±0.000 b | 0.001±0.000 b | 0.001±0.002 b | 0.077±0.037 a | 0.001±0.001 b | 2 | <0.001 |
| | Proportion of mesodecomposers | 85±10 a | 80±9 a | 84±7 a | 88±3 a | 92±4 a | 82±9 a | 1 | 0.170 |
| Functional parameters mU g ⁻¹ dry soil | Proportion of macrodecomposers | 42±11 a | 37±18 a | 27±12 ab | 7±7 b | 37±18 a | 25±1 ab | 2 | <0.001 |