



## Tectonics of the Deccan Large Igneous Province: an introduction

Soumyajit Mukherjee, Achyuta Ayan Misra, G r me Calv s, Michal Nem ok

### ► To cite this version:

Soumyajit Mukherjee, Achyuta Ayan Misra, G r me Calv s, Michal Nem ok. Tectonics of the Deccan Large Igneous Province: an introduction. Tectonics of the Deccan Large Igneous Province, 445 (1), pp.1 - 9, 2017, 10.1144/SP445.14 . hal-01697291

**HAL Id: hal-01697291**

**<https://hal.science/hal-01697291>**

Submitted on 31 Jan 2018

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destin e au d p t et   la diffusion de documents scientifiques de niveau recherche, publi s ou non,  manant des  tablissements d'enseignement et de recherche fran ais ou  trangers, des laboratoires publics ou priv s.

# Tectonics of the Deccan Large Igneous Province: an introduction

SOUMYAJIT MUKHERJEE<sup>1\*</sup>, ACHYUTA AYAN MISRA<sup>2</sup>, GÉRÔME CALVÈS<sup>3</sup> & MICHAL NEMČOK<sup>4,5</sup>

<sup>1</sup>*Department of Earth Sciences, Indian Institute of Technology Bombay, Mumbai 400 076, Maharashtra, India*

<sup>2</sup>*Exploration, Reliance Industries Ltd, Mumbai 400 701, Maharashtra, India*

<sup>3</sup>*Université Toulouse 3, Paul Sabatier, Géosciences Environnement Toulouse, 14 avenue Edouard Belin, 31400, Toulouse, France*

<sup>4</sup>*EGI at University of Utah, 423 Wakara Way, Suite 300, Salt Lake City, UT 84108, USA*

<sup>5</sup>*EGI Laboratory at SAV, Dúbravská cesta 9, 840 05 Bratislava, Slovakia*

\*Correspondence: soumyajitm@gmail.com



**Gold Open Access:** This article is published under the terms of the CC-BY 3.0 license.

Many questions have been raised about the tectonic controls and structural evolution of the large igneous provinces (LIPs) of the world; these provinces are of enormous interest to tectonicians (see recent review by Ernst 2014) and the Deccan Traps are no exception (e.g. Dessai & Bertrand 1995; Peshwa & Kale 1997; Collier *et al.* 2008; Corfield *et al.* 2010; Hooper *et al.* 2010; Calvès *et al.* 2011; Dixit *et al.* 2014; Misra 2015; Misra *et al.* 2014, 2015, 2016; Misra & Mukherjee 2015). The Deccan large igneous province (DLIP) (the Deccan Traps) is the product of the voluminous outpouring of mainly basaltic lavas at the Cretaceous–Tertiary (K–T) boundary and constitutes one of the largest known LIPs in the world (Bryan *et al.* 2010). The Deccan Volcanic Province (DVP) covers >500 000 km<sup>2</sup> (Watts & Cox 1989; Deshpande 1998; Chandrasekharam 2003 and references cited therein; see reviews by Vaidhyanadhan & Ramakrishnan 2008; Philpotts & Ague 2009). Pre-erosion estimates for the area covered by the lavas vary widely, but are typically 1–2.6 × 10<sup>6</sup> km<sup>2</sup> (Chandrasekharam 2003 from references cited therein; Vaidhyanadhan & Ramakrishnan 2008). The total thickness of the Deccan Traps exceeds 2000 m (Kaila *et al.* 1981; Deshpande 1998; Harinarayana *et al.* 2007). The Deccan Traps are one of the most important LIPs in studies of the origin and eruption of large volumes of basaltic magma since the Mesozoic.

The Deccan volcanism produced outpourings of lavas over an extended period of time from c. 68 to 60 Ma (reviews by Pande 2002; Chenet *et al.* 2007 and references cited therein; Hooper *et al.* 2010; Valdiya 2010, 2011; Baksi 2014), possibly in pulses (Chenet *et al.* 2007). Peak volcanism occurred at

c. 65 ± 0.5 Ma (Chenet *et al.* 2007; review by Valdiya 2010). Most of the research on the onshore part of the DLIP has focused on the lithology, geochemistry, isotopic ages and overall relationship of the volcanism with continental rifting. Organized studies on the tectonic structure of the underlying continental lithosphere, the structural development of the volcanic province and the post-eruption tectonic development of the province have been relatively undocumented, pending fresh studies (e.g. Misra *et al.* 2014). The western DLIP is an exposed passive continental margin and is a globally significant example of a volcanic margin, crucial in our understanding of the dynamics of such margins. The offshore areas of the passive margin have been investigated using single-channel and seismic refraction studies, although none of these studies has produced images deep enough to image the entire crust. New data from many sources have now been made available (Misra *et al.* 2015, 2016). This book presents further research in the onshore and offshore passive margin regions. The northern areas of the DLIP in Kutch (Gujarat, India) are highly active and the central parts are moderately active in terms of present day deformation manifested by recurring earthquakes, possibly related to the ongoing Himalayan orogeny (Bilham *et al.* 2001; Bodin & Horton 2004; Mukherjee *et al.* 2013, 2015). The Early Tertiary was characterized by tensile to strike-slip stresses due to rifting, which changed to compressive to strike-slip stresses from the Miocene Himalayan collision onwards (Bilham *et al.* 2001; Misra *et al.* 2014, 2015, 2016).

Comprehending the tectonics of the Deccan Traps, which outcrop mainly in the Indian states of Maharashtra, Gujarat, Madhya Pradesh, Andhra

Pradesh and Karnataka, is also crucial in deciphering how the Seychelles microcontinent rifted from India, with implications for offshore basin and continental margin evolution around the northern Indian Ocean.

This Special Publication contains 13 research papers on the tectonic aspects of the DLIP, commonly referred to as the Deccan Traps. The broad perspectives of these papers include field-based structural geology, geochemistry, analytical models, the relationship of geomorphology to structure and tectonics, and geophysics (palaeomagnetism, gravity and magnetic anomalies, and seismic imaging), with respect to both palaeo- and neotectonics. These papers develop the relatively neglected tectonic investigations of the Deccan LIP and demonstrate that it is important to understand the structure and deformation of the province as well as the petrology, which has been well studied over many years.

## Overview of the tectonics and structural geology of the DLIP


Madagascar, Seychelles and India constituted a single large continental block until Cenomanian–Turonian times. Madagascar rifted from India at about 90 Ma. In a subsequent rifting event at about 63.4 Ma, the Seychelles microcontinent and the Laxmi Ridge, a microcontinental block now located between India and Seychelles, rifted from India. This rifting event of the Laxmi and Seychelles microcontinents was almost coeval with Deccan flood volcanism (reviewed in Bhattacharya & Yatheesh 2015). Geophysical studies have been already conducted to ascertain the thicknesses of the trap/crust/lithosphere, the basement configuration and mantle anisotropies (reviewed in Misra 2015). The elastic thickness ( $T_e$ ) of the plate beneath the Deccan Traps is *c.* 8–10 km (Tiwari & Misra 1999). The anomalous gravity high in the Mumbai coastal area indicates either a shallow secondary plume that supplied lava for the Deccan volcanics (Negi *et al.* 1992) or a bolide impact at around the K–T boundary that might have triggered Deccan lava eruption (Negi *et al.* 1993). By contrast, Sethna (2003) considered acid volcanic rocks, and Bhattacharji *et al.* (2004) mafic bodies, to give rise to the anomalous gravity high. Flows in the Deccan Traps are mostly sub-horizontal (Watts & Cox 1989), although Devey & Lightfoot (1986) deciphered a regional ‘very open’ antiform around Panvel (Maharashtra, India), along which the Deccan lava layers steepen to constitute the Panvel Flexure.

Regional faults have been identified in the north and west Deccan Traps from geophysical and remote sensing studies (Fig. 1; Harinarayana *et al.*

2007; Chandrasekhar *et al.* 2011; Kumar *et al.* 2011). The Cambay rift is bound by NNW-trending faults affecting the Deccan Traps (Dixit *et al.* 2014). The east–west-trending Tapti Fault defines the north boundary of the main outcrop of the DLIP (Azeez *et al.* 2011). The west coast passive margin is defined by the NNW-trending West Coast Fault (Fig. 1; Chandrasekharam 1985; Biswas 1987). The location of this fault is debatable because compelling field evidence is difficult to obtain. The Western Ghat escarpment (Figs 2 & 3; Kale 2010) is defined by a sharp change in topography, with the eastern side uplifted by 800–1200 m. This escarpment is considered to be a rift-bounding fault of the west coast passive margin (Gunnell & Fleitout 1998). The NNW-trending Koyana Fault parallels this escarpment. The Kurduwardi Lineament is another important structural element (Peshwa & Kale 1997). The northernmost region of the Kurduwardi Lineament is manifested by a NW-trending > 20 km long series of fractures/faults in Thane district (Maharashtra State, India) (Peshwa & Kale 1997). Slickensides have developed over the fault planes. These faults have a *c.* 10–30 m throw and indicate both dextral and sinistral shear. Flexure, indicated by gently dipping flow layers (Fig. 3) related to the West Coast Fault, opened fractures that were subsequently filled with igneous material to form dykes (Peshwa & Kale 1997). Dessai & Bertrand (1995) and Hooper *et al.* (2010) reported north–south shear zones from the Deccan coastal areas, extending for *c.* 40 km from Murud to Mumbai (Fig. 4). Dessai & Bertrand (1995) reported steep fault planes with subvertical displacement and occasional slickensides. Hooper *et al.* (2010) established that the earliest (66–65 Ma) dykes have been sheared and that the later dykes were oriented either along the shear planes or intruded into them.

## Why study structures in the DLIP?

The structural and tectonic features of the Deccan region have significant economic and societal importance and the region includes the important city of Mumbai and its hinterland. Basin development and the locations of fractures and faults in the DLIP have far-reaching implications in hydrocarbon exploration (Pandey & Agarwal 2000). There are four petroliferous basins in India within the DLIP: the Barmer, Kutch, Cambay and Mumbai shelf basins. The Deccan syncline and Saurashtra basins have also been studied for the occurrence of hydrocarbons. The area is important for groundwater resources and the Deccan structures are important for groundwater exploration (Rai *et al.* 2011). The seismicity of the area is a concern and the region around Koyana in the Deccan Traps suffered



**Fig. 1.** Major tectonic elements of the Deccan large igneous province as deciphered from geophysical studies. Note the gravity anomaly highs associated with each of the rift zones (modified from Bhattacharji *et al.* 1996).

seismicity in 1967, possibly because of seepage-induced reactivation of the fault zone, which received global attention. Structural geological and tectonic studies can, in the long term, indicate the subsurface structures of the Deccan Traps as an important input into seismic studies, including the identification of seismic risk. Structural and tectonic studies will be better able to constrain possible locations for carbon sequestration in the Deccan Traps (Jayaraman 2007).

### Content of this edited book

Detailed geomorphologic and morphometric studies of river networks within the area underlain by the Deccan Traps are presented by **Kale *et al.* (2016)**. This study indicates that the Western Upland region of the Deccan Plateau consists of at least three blocks with dissimilar Quaternary uplift histories. The authors therefore argue that the classical thought that the Deccan Traps are structurally monotonous is incorrect. **Mitra *et al.* (2016)** document smectite and kaolinite variations associated

with faults bounding the Matanomadh Basin in Gujarat and use alteration patterns to interpret the presence of a palaeoslope created by Cenozoic rifting, through which water flowed and selectively weathered the basalt lavas. **Pathak *et al.* (2016)** studied 37 lava flows in the eastern DVP by palaeomagnetic methods. Their virtual geomagnetic pole positions indicate a brief eruption period at the Cretaceous–Palaeogene boundary in the eastern and western Deccan Traps. The western Deccan Traps are near the west coast of India and their eastern counterparts are at the easternmost extreme of the DLIP. The virtual geomagnetic pole position does not indicate a brief eruption period around the Cretaceous–Palaeogene boundary, but represents a record of the past geomagnetic field direction at the sampling site/location during the interval of time over which the characteristic remnant magnetization directions were acquired. **Nemčok & Rybár (2016)** describe the rift–drift transition processes and their controlling factors in the magma-rich Gop Rift–Laxmi Basin that predated the Deccan eruption event. Details of the documented break-up mechanism apply to magma-rich break-ups worldwide.





**Fig. 2.** View of sub-horizontal lava flows of the Deccan large igneous province looking towards the west from Lion Point in Lonavala, c. 100 km east of Mumbai. This location is on the top of the Western Ghat Escarpment; Photograph by Achyuta Ayan Misra.

**Misra *et al.* (2016)** studied carbonate seismic facies and subsidence on the ridges in the NW Arabian Sea to analyse how the Deccan mantle plume affected the tectonics of the offshore region of the DLIP. **Gupta *et al.* (2016)** detail seismicity based on several geophysical techniques in the Koyna region, where the periodic loading and unloading of water has been envisaged for seismicity induced by artificial water reservoirs since 1967. Their preliminary borehole investigations revealed faults and foliation planes and more detail is expected from this research group in the coming years regarding subsurface structural geology at Koyna. Analyses of P-wave receiver functions by **Mandal (2016)** enabled him to delineate the crust–mantle structure below the Kutch rift zone. He attributed this complex structure to be a key causative factor of the recent lower crustal seismicity in the region. **Rajaram *et al.* (2016)** performed detailed magnetic studies on part of the Deccan Traps and identified seven subsurface lineaments. They correlated these lineaments in the magnetic anomalies with already known brittle shear zones, faults and river

courses. Identifying such lineaments can pinpoint unrecognized fault or shear zones, or sediment-filled basins, underneath the Deccan basalt cover. **Maurya *et al.* (2016)** reviewed the neotectonic roles of active faults (Mukherjee 2013, 2014a, b, 2015, 2016) in the Kutch basin with respect to the post-Deccan Traps inversion of the Kachchh basin and the present day seismicity. With the help of ground-penetrating radar, they also documented a few subsurface faults. Such studies can correlate the lineaments in studies such as **Rajaram *et al.* (2016)**.

Several papers are concerned with the field and structural interpretation of basaltic dykes associated with the DLIP. The field structural studies of **Misra & Mukherjee (2016)** revealed differences in the structural deformation of different age groups of basaltic dykes in the province. An older c. 65.5 Ma group of dykes (Group I) are faulted/sheared and lacking in a uniform trend. Younger c. 65–63 Ma groups II and III dykes are syn- to post-deformation intrusions and intrude into the faults/shears. They point out that, in addition to the dyke trends, the



**Fig. 3.** Southern ridge of the *c.* 1000 m high Matheran Plateau showing sub-horizontal to gently dipping flows of the Deccan large igneous province. View looking towards the south; Photograph by Achyuta Ayan Misra.





**Fig. 4.** A typical wave-cut platform at Janjira-Murud c. 140 km south of Mumbai along the west coast of India. This view looking towards the south shows the hilly terrain around the area and the Janjira sea fort in the background. Such sub-horizontal exposures along rocky stretches provide a wealth of geological knowledge on the tectonics of the Deccan large igneous province (e.g. Dessai & Bertrand 1995; Hooper *et al.* 2010; Misra *et al.* 2014; **Misra & Mukherjee 2016**). Photograph by Achyuta Ayan Misra.

direction of the Seychelles–India separation can also be studied from the attitude of the brittle shear planes within the Deccan Traps. The analytical modelling of **Ju *et al.* (2016)** explains how propagating dykes deflect due to mechanical layering in the crust. They also postulate a four-stage evolution of the dykes from the Narmada rift zone. **Kaplay *et al.* (2016)** report dykes from the Kinwat region in the eastern part of the Deccan Traps. Based on the occurrence of faults and lineaments, the authors consider Kinwat to be a future location of intraplate seismicity. **Babar *et al.* (2016)** report brittle-sheared NE-trending dykes from the Aurangabad region that match the trend of the Narmada–Tapti lineament. They also report on field studies of the local directions of lava flows.

The collection of papers in this volume provides new insights into the structural features and the tectonic evolution of the DLIP. This provides an impetus for further field geological and geophysical studies. The main findings include: evidence that the DLIP is divisible into a number of structural blocks separated by faults, rather than being a single coherent block; interpretation of the rift-induced palaeotopography, which provides further constraints for the eruption period; details of how continental rifting occurs in magma-rich systems; the role of the Deccan/Réunion plumes in the tectonics of the DLIP; the emplacement of dykes with different time and structural relationships to faults and shear zones from borehole studies; the demarcation of the crust–mantle structure; the identification of subsurface tectonic boundaries; and basin inversion after Deccan basalt eruption. This volume provides an impetus for further field-based geological and geophysical studies of this important province.

Thanks to all the participating authors and reviewers. Comments by reviewers on the book proposal helped us to attain a balance in choosing papers for this volume. Support from Phil Leat, Angharad Hills, Tamzin Anderson and Rachael Kriefman (Geological Society of London) is gratefully acknowledged. Phil Leat is owed additional thanks for reviewing this introductory chapter.

## References

- AZEEZ, K.K.A., KUMAR, T.S., BASAVA, S., HARINARAYANA, T. & DAYAL, A.M. 2011. Hydrocarbon prospects across Narmada-Tapti rift in Deccan trap, central India: inferences from integrated interpretation of magnetotelluric and geochemical prospecting studies. *Marine and Petroleum Geology*, **28**, 1073–1082.
- BABAR, M.D., KAPLAY, R.D., MUKHERJEE, S. & KULKARNI, P.S. 2016. Evidence of deformation of dykes from the Central Deccan Volcanic Province, Aurangabad, Maharashtra, India. In: MUKHERJEE, S., MISRA, A.A., CALVÈS, G. & NEMČOK, M. (eds) *Tectonics of the Deccan Large Igneous Province*. Geological Society, London, Special Publications, **445**. First published online November 8, 2016, <https://doi.org/10.1144/SP445.13>
- BAKSI, A.K. 2014. The Deccan Trap – Cretaceous–Paleogene boundary connection; new  $^{40}\text{Ar}/^{39}\text{Ar}$  ages and critical assessment of existing argon data pertinent to this hypothesis. *Journal of Asian Earth Sciences*, **84**, 9–23.
- BODIN, P. & HORTON, S. 2004. Source parameters and tectonic implications of aftershocks of the Mw 7.6 Bhuj earthquake of 26 January 2001. *Bulletin of the Seismological Society of America*, **94**, 818–827.
- BHATTACHARJ, S., CHATTERJEE, N., WAMPLER, J.M., NAYAK, P.N. & DESHMUKH, S.S. 1996. Indian intraplate and continental margin rifting, lithospheric extension, and mantle upwelling in Deccan flood basalt volcanism near the K/T boundary: evidence from mafic dike swarms. *Journal of Geology*, **104**, 379–398.
- BHATTACHARJ, S., SHARMA, R. & CHATTERJEE, N. 2004. Two- and three dimensional gravity modelling along west continental margin and intraplate Narmada-Tapi rifts: its relevance to Deccan flood basalt volcanism. *Proceedings of the Indian Academy of Sciences (Earth and Planetary Sciences)*, **113**, 771–784.
- BHATTACHARYA, G.C. & YATHEESH, Y. 2015. Plate-tectonic evolution of the deep ocean basins adjoining the western continental margin of India – a proposed model for the early opening scenario. In: MUKHERJEE, S. (ed.) *Petroleum Geosciences: Indian Contexts*. Springer Geology Series, 1–61.
- BILHAM, R., GAUR, V.K. & MOLNAR, P. 2001. Himalayan seismic hazard. *Science*, **293**, 1442–1444.
- BISWAS, S.K. 1987. Regional tectonic framework, structure and evolution of the western marginal basins of India. *Tectonophysics*, **135**, 307–327.
- BRYAN, S.E., PEATE, I.U. *ET AL.* 2010. The largest volcanic eruptions on Earth. *Earth-Science Reviews*, **102**, 207–229.
- CALVÈS, G., SCHWAB, A.M. *ET AL.* 2011. Seismic volcanostigraphy of the western Indian rifted margin: the pre-Deccan igneous province. *Journal of Geophysical Research*, **116**, B01101.
- CHANDRASEKHAR, P., MARTHA, T.R., VENKATERSWARLU, N., SUBRAMANIAN, S.K. & KAMARAJU, M.V.V. 2011. Regional geological studies over parts of Deccan Syncline using remote sensing and geophysical data for understanding hydrocarbon prospects. *Current Science*, **100**, 95–99.
- CHANDRASEKHARAM, D. 1985. Structure and evolution of the western continental margin of India deduced from gravity, seismic, geomagnetic and geochronological studies. *Physics of the Earth and Planetary Interiors*, **41**, 186–198.
- CHANDRASEKHARAM, D. 2003. Deccan flood basalts. *Memor of the Geological Society of India*, **53**, 197–214.
- CHENET, A.L., QUIDELLEUR, Z., FLUTEAU, F., COURTILLOT, V. & BAJPAI, S. 2007.  $^{40}\text{K}$ – $^{40}\text{Ar}$  dating of the main Deccan large igneous province: further evidence of KTB age and short duration. *Earth and Planetary Science Letters*, **263**, 1–15.
- COLLIER, J.S., SANSOM, V., ISHIZUKA, O., TAYLOR, R.N., MINSHULL, T.A. & WHITMARSH, R.B. 2008. Age of Seychelles–India break-up. *Earth and Planetary Science Letters*, **272**, 264–277.

- CORFIELD, R.I., CARMICHAEL, S., BENNETT, J., AKHTER, S., FATIMI, M. & CRAIG, T. 2010. Variability in the crustal structure of the West Indian continental margin in the northern Arabian Sea. *Petroleum Geoscience*, **16**, 257–265.
- DESHPANDE, G.G. 1998. *Geology of Maharashtra*. Geological Survey of India, Bangalore, 129–162.
- DESSAI, A. & BERTRAND, H. 1995. The 'Panvel flexure' along the western Indian continental margin: an extensional fault structure related to Deccan magmatism. *Tectonophysics*, **241**, 165–178.
- DEVY, C.W. & LIGHTFOOT, P. 1986. Volcanological and tectonic control of stratigraphy and structure in the Western Deccan Traps. *Bulletin of Volcanology*, **48**, 195–207.
- DIXIT, M.M., KUMAR, S., CATCHINGS, R.D., SUMAN, K., SARKAR, D. & SEN, M.K. 2014. Seismicity, faulting, and structure of the Koyna-Warna seismic region, Western India from local earthquake tomography and hypocenter locations. *Journal of Geophysical Research: Solid Earth*, **119**, 6372–6398.
- ERNST, R.E. 2014. *Large Igneous Provinces*. Cambridge University Press, Cambridge.
- GUNNELL, Y. & FLEITOUT, L. 1998. Shoulder uplift of the Western Ghats passive margin, India: a denudational model. *Earth Surface Processes and Landforms*, **23**, 391–404.
- GUPTA, H.K., ARORA, K. ET AL. 2016. Investigations of continued reservoir triggered seismicity at Koyna, India. In: MUKHERJEE, S., MISRA, A.A., CALVÈS, G. & NEMČOK, M. (eds) *Tectonics of the Deccan Large Igneous Province*. Geological Society, London, Special Publications, **445**. First published online October 24, 2016, <https://doi.org/10.1144/SP445.11>
- HARINARAYANA, T., PATRO, B.P.K., VEERASWAMI, K., MANOI, C., NAGANJANEYULU, K., MURTHY, D.N. & VIRUPAKSHI, G. 2007. Regional geoelectric structure beneath Deccan Volcanic Province of the Indian subcontinent using magnetotellurics. *Tectonophysics*, **445**, 66–80.
- HOOVER, P., WIDDOWSON, M. & KELLEY, S. 2010. Tectonic setting and timing of the final Deccan flood basalt eruptions. *Geology*, **38**, 839–842.
- JAYARAMAN, K. 2007. India's carbon dioxide trap. *Nature Geoscience*, **445**, 350.
- JU, W., HOU, G. & HARI, K.R. 2016. Dyke emplacement in the Narmada rift zone and implications for the evolution of the Deccan Traps. In: MUKHERJEE, S., MISRA, A.A., CALVÈS, G. & NEMČOK, M. (eds) *Tectonics of the Deccan Large Igneous Province*. Geological Society, London, Special Publications, **445**. First published online May 30, 2016, <https://doi.org/10.1144/SP445.1>
- KAILA, K.L., MURTY, P.R.K., RAO, V.K. & KHAR-ETCHKO, G.E. 1981. Crustal structure from deep seismic soundings along the Koyna II (Kelsi-Loni) profile in the Deccan Trap area, India. *Tectonophysics*, **73**, 365–384.
- KALE, V.S. 2010. The Western Ghat: the Great Escarpment of India. In: MIGNON, P. (ed.) *Geomorphological Landscapes of the World*. Springer, Dordrecht, 257–264.
- KALE, V.S., DOLE, G., UPASANI, D. & PILLAI, S.P. 2016. Deccan Plateau uplift: insights from parts of Western Uplands, Maharashtra, India. In: MUKHERJEE, S., MISRA, A.A., CALVÈS, G. & NEMČOK, M. (eds) *Tectonics of the Deccan Large Igneous Province*. Geological Society, London, Special Publications, **445**. First published online July 20, 2016, <https://doi.org/10.1144/SP445.2>
- KAPLAY, R.D., BABAR, M.D., MUKHERJEE, S. & KUMAR, T.V. 2016. Morphotectonic expression of geological structures in eastern part of the South East Deccan Volcanic Province (around Nanded, Maharashtra, India). In: MUKHERJEE, S., MISRA, A.A., CALVÈS, G. & NEMČOK, M. (eds) *Tectonics of the Deccan Large Igneous Province*. Geological Society, London, Special Publications, **445**. First published online November 8, 2016, <https://doi.org/10.1144/SP445.12>
- KUMAR, D., THIAGARAJAN, S. & RAI, S.N. 2011. Deciphering geothermal resources in Deccan Trap region using electrical resistivity tomography technique. *Journal of the Geological Society of India*, **78**, 541–548.
- MANDAL, P. 2016. Influence of Deccan volcanism/synrift magmatism on the crust–mantle structure and its implications for the seismogenesis of earthquakes occurring in the Kachhh rift zone. In: MUKHERJEE, S., MISRA, A.A., CALVÈS, G. & NEMČOK, M. (eds) *Tectonics of the Deccan Large Igneous Province*. Geological Society, London, Special Publications, **445**. First published online July 11, 2016, <https://doi.org/10.1144/SP445.6>
- MAURYA, D.M., CHOWKSEY, V., PATIDAR, A.K. & CHAMYAL, L.S. 2016. A review and new data on neotectonic evolution of active faults in the Kachhh Basin, Western India: legacy of post-Deccan Trap tectonic inversion. In: MUKHERJEE, S., MISRA, A.A., CALVÈS, G. & NEMČOK, M. (eds) *Tectonics of the Deccan Large Igneous Province*. Geological Society, London, Special Publications, **445**. First published online September 19, 2016, <https://doi.org/10.1144/SP445.7>
- MISRA, A.A. 2015. *Sheared nature of the Indian western continental margin around Mumbai, India: onshore and offshore geoscientific studies*. PhD thesis, Indian Institute of Technology, Bombay.
- MISRA, A.A. & MUKHERJEE, S. 2015. *Tectonic Inheritance in Continental Rifts and Passive Margins*. Springer-Briefs in Earth Sciences. Springer International Publishing, Berlin.
- MISRA, A.A. & MUKHERJEE, S. 2016. Dyke–brittle shear relationships in the Western Deccan Strike-slip Zone around Mumbai (Maharashtra, India). In: MUKHERJEE, S., MISRA, A.A., CALVÈS, G. & NEMČOK, M. (eds) *Tectonics of the Deccan Large Igneous Province*. Geological Society, London, Special Publications, **445**. First published online June 22, 2016, <https://doi.org/10.1144/SP445.4>
- MISRA, A.A., BHATTACHARYA, G., MUKHERJEE, S. & BOSE, N. 2014. Near N–S paleo-extension in the western Deccan region in India: does it link strike-slip tectonics with India–Seychelles rifting? *International Journal of Earth Sciences*, **103**, 1645–1680.
- MISRA, A.A., SINHA, N. & MUKHERJEE, S. 2015. Repeat ridge jumps and microcontinent separation: insights from NE Arabian Sea. *Marine and Petroleum Geology*, **59**, 406–428.
- MISRA, A.A., BANERJEE, S., KUNDU, N. & MUKHERJEE, B. 2016. Subsidence around oceanic ridges along passive

- margins: NE Arabian Sea. In: MUKHERJEE, S., MISRA, A.A., CALVÈS, G. & NEMČOK, M. (eds) *Tectonics of the Deccan Large Igneous Province*. Geological Society, London, Special Publications, **445**. First published online October 12, 2016, <https://doi.org/10.1144/SP445.10>
- MITRA, S., MITRA, K., GUPTA, S., BHATTACHARYA, S., CHAUHAN, P. & JAIN, N. 2016. Alteration and submergence of basalts in Kachchh, Gujarat, India: implications for the role of the Deccan Traps in the India–Seychelles break-up. In: MUKHERJEE, S., MISRA, A.A., CALVÈS, G. & NEMČOK, M. (eds) *Tectonics of the Deccan Large Igneous Province*. Geological Society, London, Special Publications, **445**. First published online August 31, 2016, <https://doi.org/10.1144/SP445.9>
- MUKHERJEE, S. 2013. *Deformation Microstructures in Rocks*. Springer, Berlin.
- MUKHERJEE, S. 2014a. *Atlas of Shear Zone Structures in Meso-Scale*. Springer, Berlin.
- MUKHERJEE, S. 2014b. Review of flanking structures in meso- and micro-scales. *Geological Magazine*, **151**, 957–974.
- MUKHERJEE, S. 2015. *Atlas of Structural Geology*. Elsevier, Amsterdam.
- MUKHERJEE, S. 2016. A review on out-of-sequence deformation in the Himalaya. In: MUKHERJEE, S., CAROSI, R., VAN DER BEEK, P., MUKHERJEE, B.K. & ROBINSON, D. (eds) *Tectonics of the Himalaya*. Geological Society, London, Special Publications, **412**, 67–109.
- MUKHERJEE, S., MUKHERJEE, B. & THIEDE, R. 2013. Geosciences of the Himalaya-Karakoram-Tibet Orogen. *International Journal of Earth Sciences*, **102**, 1757–1758.
- MUKHERJEE, S., CAROSI, R., VAN DER BEEK, P.A., MUKHERJEE, B.K. & ROBINSON, D.M. (eds) 2015. *Tectonics of the Himalaya: an Introduction*. Geological Society, London, Special Publications, **412**, 1–3.
- NEGI, J.G., AGARWAL, P.K., SINGH, A.P. & PANDEY, O.P. 1992. Bombay gravity high and eruption of Deccan flood basalts (India) from a shallow secondary plume. *Tectonophysics*, **206**, 341–350.
- NEGI, J.G., AGARWAL, P.K., PANDEY, O.P. & SINGH, A.P. 1993. A possible K-T boundary bolide impact site offshore near Bombay and triggering of rapid Deccan volcanism. *Physics of the Earth and Planetary Interiors*, **76**, 189–197.
- NEMČOK, M. & RYBÁR, S. 2016. Rift–drift transition in a magma-rich system: the Gop Rift–Laxmi Basin case study, West India. In: MUKHERJEE, S., MISRA, A.A., CALVÈS, G. & NEMČOK, M. (eds) *Tectonics of the Deccan Large Igneous Province*. Geological Society, London, Special Publications, **445**. First published online July 20, 2016, <https://doi.org/10.1144/SP445.5>
- PANDE, K. 2002. Age and duration of the Deccan Traps, India: a review of radiometric and paleomagnetic constraints. *Proceedings of the Indian Academy of Science (Earth and Planetary Science)*, **111**, 115–123.
- PANDEY, O.P. & AGARWAL, P.K. 2000. Thermal regime, hydrocarbon maturation and geodynamic events along the western margin of India since the Late Cretaceous. *Journal of Geodynamics*, **30**, 439–459.
- PATHAK, V., PATIL, S.K. & SRIVASTAVA, J.P. 2016. Tectonomagmatic setting of lava packages in the Mandla lobe of the eastern Deccan volcanic province, India: palaeomagnetism and magnetostratigraphic evidence. In: MUKHERJEE, S., MISRA, A.A., CALVÈS, G., NEMČOK, M. (eds) *Tectonics of the Deccan Large Igneous Province*. Geological Society, London, Special Publications, **445**. First published online July 21, 2016, <https://doi.org/10.1144/SP445.3>
- PESHPWA, V.V. & KALE, V.S. 1997. Neotectonics of the Deccan Traps Province: focus on the Kurduwadi Lineament. *Journal of Geophysics*, **18**, 77–86.
- PHILPOTTS, A.R. & AGUE, J.J. 2009. *Principles of Igneous and Metamorphic Petrology*. Cambridge University Press, Cambridge
- RAI, S.N., THIAGARAJAN, S. & KUMARI, Y.R. 2011. Exploration for groundwater in the basaltic Deccan traps terrain in Katrol Taluk, Nagpur District, India. *Current Science*, **101**, 1198–1205.
- RAJARAM, M., ANAND, S.P., ERRAM, V.C. & SHINDE, B.N. 2016. Insight into the structures below the Deccan Trap-covered region of Maharashtra, India from geopotential data. In: MUKHERJEE, S., MISRA, A.A., CALVÈS, G. & NEMČOK, M. (eds) *Tectonics of the Deccan Large Igneous Province*. Geological Society, London, Special Publications, **445**. First published online September 16, 2016, <https://doi.org/10.1144/SP445.8>
- SETHNA, S.F. 2003. The occurrence of acid and intermediate rocks in the Deccan Volcanic Province with associated high positive gravity anomalies and their probable significance. *Journal of the Geological Society of India*, **61**, 220–222.
- TIWARI, V.M. & MISRA, D.C. 1999. Estimation of effective elastic thickness from gravity and topography data under the Deccan Volcanic Province, India. *Earth and Planetary Science Letters*, **171**, 289–299.
- VAIDHYANADHAN, R. & RAMAKRISHNAN, M. 2008. *Geology of India*. Vol. 2. Geological Society of India, Bangalore, 733–784.
- VALDIYA, K.S. 2010. *The Making of India: Geodynamic Evolution*. Macmillan, New Delhi.
- VALDIYA, K.S. 2011. Some burning questions remaining unanswered. *Journal of the Geological Society of India*, **78**, 299–320.
- WATTS, A.B. & COX, K.G. 1989. The Deccan Traps: an interpretation in terms of progressive lithospheric flexure in response to a migrating load. *Earth and Planetary Science Letters*, **93**, 85–97.