







# Source reduction of personal care products in the urban water cycle of Paris conurbation

Régis Moilleron & Adèle Bressy


















# Context





Darbre Ph. *et al.* (2004)

Potential link with breast cancer



**Envoyé Spécial (2005)** 

Whistle-blowers



**Consumers Industrials** 




Geara-Matta (2012)

Flow and sources of parabens...: comparison between Paris and Beirut




# Context





# Context





## Cosmet'eau



- Personal care products: a pretext to study risks associated with micropollutants and source control solutions
- Two issues
  - Environmental impact
    - Effluent monitoring, ecotoxicological impacts of the different types of cosmetic product
  - Health impact through:
    - Alert (scientific origin)
    - The responses (from industrials)
    - Perception of risk by consumers of cosmetic products (surveys) and urban managers
- Multidisciplinary approach to the strategies of actors likely to interact with micropollutants, in particular risk anticipation (health and environment)
- Possibility of prevention at source of the impact of micropollutants on the environment (and on health)
  - Acting on products consumed and consumption practices,
  - Preventive policy partly promoted by local and regional authorities

# Aims





To determine the impact of PCP use on the quality of urban effluents



To study the transfer from their release into greywater up to the receiving media




To assess the concentrations observed in the receiving environment

To evaluate the impact of the Paris conurbation

## What Molecules?



#### **Preservatives and biocides**





pharmaceuticals foodstuffs personal care products...



Aqua, glycerin, glucose, calendula officinalis extract, potassium sorbate, sodium benzoate, phenoxyethanol, methylparaben, butylparaben, ethylparaben, propylparaben, isobutylparaben



# Part I Housing scale: Greywater study




To determine the origin of domestic wastewater contamination for source reduction purposes












## Greywater contamination - results





1. Ubiquitous contamination for every greywater


2. Intra-greywater variability highlighted


3. MeP (> TCS) > PrP = EtP → formulation of cosmetics


## Greywater contamination - results




#### **GREYWATER (ng/L)**











# **Findings**


- Contamination of all greywater waters by parabens, TCS and TCC
  - No unique source... 
     ¬ questioning of information available to the consumers (INCI regulation)
- MeP, EtP & PrP predominate
  - Consistent with the uses associated with these preservatives... in personal care products
- Significant concentration variability: from < LOQ up to 2 mg/L (shower) for parabens and from 36 ng/L up to 20  $\mu$ g/L (shower) for TCS
  - Nature of products used
  - User habits need to acquire more data?
- Major contribution of shower and washing machine (loads)
  - **→** What are the product origins?

# Part II Urban watershed scale: Wastewater study

#### **Selected activities**




#### Main sewer trunks







#### Wastewater (ng/L)




1. Industrial effluents (more or less) contaminated...





#### Wastewater (ng/L)



1. Industrial effluents (more or less) contaminated...

2. Wastewater levels similar to greywater ones


#### Wastewater - results




Synoptic (ng/L)

Greywater

Hair Dresser







**Beauty Salon** 

Hospital


Catchment
















#### 1. Same profiles for parabens











- 1. Same profiles for parabens
- 2. When the scale increases, the variability decreases











- 1. Same profiles for parabens
- 2. When the scale increases, the variability decreases
- 3. Surprising behaviour of triclosan and triclocarban...









- 1. Same profiles for parabens
- 2. When the scale increases, the variability decreases
- 3. Surprising behaviour of triclosan and triclocarban...
- → Substantial flows reach WWTPs: what about the receiving environment?



# **Findings**

- Ubiquitous wastewater contamination by parabens, TCS and TCC
  - Domestic and industrial origins
  - → Relative contributions to identify levers for action?
- MeP, EtP & PrP predominate
  - Consistent with the uses associated with these preservatives... in personal care products
- Significant variability of concentrations
  - Within a given sector of activity but also between different sectors
  - → What about the composition of the products used?
- Smoothing of observed levels when scale increases
  - Concentrations close to those in greywater
- Transfer from emission sources to WWTP
  - → Fate at the different stages of WWTP? In fine impact on the environment?


# Part III Surface Water Contamination


Determine the levels of impregnation of the Seine River

## WWTP discharge



#### Discharge vs main trunks (ng/L)





#### Trunks (ng/L)

MeP: 2893

EtP: 1273

PrP: 1080

IsoBuP: 9

BuP: 62

TCS: 597

TCC: 39

#### **Significant decrease of concentrations**

Efficiency of treatment processes...

#### Discharge (ng/L)

MeP: 35

EtP: 19

PrP: 21

IsoBuP: < LQ


BuP: < LQ


TCS: 78

TCC:



#### CSOs (ng/L)





# CSO levels in between sewer trunks and discharge

#### DO (ng/l)

MeP: 1757

EtP: 476

PrP: 381

IsoBuP: < LQ

BuP: 32

TCS: 109

TCC: 24


#### **High temporal variability**

- Largely related to the intensity of the discharge...
- and therefore the volume of wastewater mixed to runoff.






#### **UPSTREAM (ng/L)**



#### **DOWNSTREAM (ng/L)**




1. Increase in concentrations from upstream to downstream of the Paris conurbation... for parabens






#### **UPSTREAM (ng/L)**




#### **DOWNSTREAM (ng/L)**




- 1. Increase in concentrations from upstream to downstream of the Paris conurbation... for parabens
- 2. Along with spatial variability > to be linked to contamination sources



#### **UPSTREAM (ng/L)**




#### **DOWNSTREAM (ng/L)**



- 1. Increase in concentrations from upstream to downstream of the Paris conurbation... for parabens
- 2. Along with spatial variability > to be linked to contamination sources
- 3. Less net increase for triclosan...



#### **Evolution of the contamination of the Seine River from upstream to downstream**





# **Findings**

- High WWTP removal efficiency...
- However WWTP discharges contaminate the receiving environment
  - → Need to refine treatments (by promoting tertiary treatment)
- CSOs: significant source of contamination
  - **→** Keep on limiting untreated discharges for major events

- Impact of the Paris conurbation on the Seine River quality
  - Increase of the concentrations from upstream to downstream

# Conclusion









Discharge into the environment



**Receiving** waters

**WWTP** 

Removal Efficiency **Impact of Paris Conurbation** 






Fate in sewer

network

**Ubiquitous Water Contamination in Urban Areas** 



Dynamics of emerging pollutants (parabens, triclosan and triclocarban) in the grey watersreceiving environment continuum

# Greywater

Significant concentration variability

# 30<sup>th</sup> November – 1<sup>st</sup> December 2017 INSA Lyon, France

# Colombian - French Conference on Urban Water Systems




























1. (very) similar profiles...











- 1. (very) similar profiles...
- 2. Same concentration levels...









- 1. (very) similar profiles...
- 2. Same concentration levels...
- 3. Except for TCS