



**HAL**  
open science

# Some inequalities involving the extended gamma function and the Kummer confluent hypergeometric $k$ -function

Kottakkaran Sooppy Nisar, Feng Qi, Gauhar Rahman, Shahid Mubeen, Muhammad Arshad

► **To cite this version:**

Kottakkaran Sooppy Nisar, Feng Qi, Gauhar Rahman, Shahid Mubeen, Muhammad Arshad. Some inequalities involving the extended gamma function and the Kummer confluent hypergeometric  $k$ -function. *Journal of Inequalities and Applications*, 2018, 2018 (Paper No. 135), 12 p. 10.1186/s13660-018-1717-8 . hal-01701647v2

**HAL Id: hal-01701647**

**<https://hal.science/hal-01701647v2>**

Submitted on 19 Jan 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## RESEARCH

# Some inequalities involving the extended gamma function and the Kummer confluent hypergeometric $k$ -function<sup>[1]</sup>

Kottakkaran Sooppy Nisar<sup>1</sup>, Feng Qi<sup>2,3\*</sup>, Gauhar Rahman<sup>4</sup>, Shahid Mubeen<sup>5</sup> and Muhammad Arshad<sup>4</sup>

\*Correspondence:

[qifeng618@gmail.com](mailto:qifeng618@gmail.com)

<sup>2</sup>Institute of Mathematics, Henan Polytechnic University, 454010, Jiaozuo, Henan, China

<sup>3</sup>School of Mathematical Sciences, Tianjin Polytechnic University, 300387, Tianjin, China

Full list of author information is available at the end of the article

## Abstract

In the paper, the authors present some inequalities involving the extended gamma function and the Kummer confluent hypergeometric  $k$ -function via some classical inequalities such as the Chebyshev inequality for synchronous (or asynchronous, respectively) mappings, give a new proof of the log-convexity of the extended gamma function by using the Hölder inequality, and introduce a Turán type mean inequality for the Kummer confluent  $k$ -hypergeometric function.

**Keywords:** extended gamma function; inequality; Kummer confluent hypergeometric  $k$ -function; logarithmic convexity

**AMS Subject Classification:** Primary 33B15; Secondary 33C15, 33B20, 33C05, 41A60

## 1 Introduction

The gamma function  $\Gamma$  can be defined [24, 28, 31, 32] by

$$\Gamma(z) = \int_0^{\infty} t^{z-1} e^{-t} dt, \quad \Re(z) > 0.$$

Alternatively, it can also be defined [19] by

$$\Gamma(z) = \lim_{n \rightarrow \infty} \frac{n! n^{z-1}}{(z)_n},$$

where  $(z)_n$  for  $z \neq 0$  is the Pochhammer symbol defined [27] as

$$(z)_n = \begin{cases} z(z+1)(z+2) \cdots (z+n-1), & n \geq 1; \\ 1, & n = 0. \end{cases}$$

The relation between  $(z)_n$  and  $\Gamma(z)$  is

$$(z)_n = \frac{\Gamma(z+n)}{\Gamma(z)}.$$

<sup>[1]</sup>Kottakkaran Sooppy Nisar, Feng Qi, Gauhar Rahman, Shahid Mubeen, and Muhammad Arshad, *Some inequalities involving the extended gamma function and the Kummer confluent hypergeometric  $k$ -function*, Journal of Inequalities and Applications 2018, Paper No. 135, 12 pages; available online at <https://doi.org/10.1186/s13660-018-1717-8>.

The beta function  $B(x, y)$  can be defined [17, 20, 21] by

$$B(x, y) = \int_0^1 t^{x-1}(1-t)^{y-1} dt, \quad \Re(x), \Re(y) > 0$$

and can be expressed by

$$B(x, y) = \frac{\Gamma(x)\Gamma(y)}{\Gamma(x+y)}, \quad \Re(x), \Re(y) > 0.$$

In 1995, Chaudhry and Zubair [3] introduced the extended gamma function

$$\Gamma_b(z) = \int_0^\infty t^{z-1}e^{-t-bt^{-1}} dt, \quad \Re(z) > 0, \quad b \geq 0. \tag{1.1}$$

If  $b = 0$ , then  $\Gamma_b$  becomes the classical gamma function  $\Gamma$ .

In 1997, Chaudhry et al [2] introduced the extended beta function

$$B_b(x, y) = \int_0^1 t^{x-1}(1-t)^{y-1}e^{-b/t(1-t)} dt, \quad \Re(b), \Re(x), \Re(y) > 0. \tag{1.2}$$

It is clear that  $B_0(x, y) = B(x, y)$ .

In 2009, Barnard et al [1] established three inequalities

$$\begin{aligned} [\phi(a, a+b, x)]^2 &> \phi(a+v, a+b, x)\phi(a-v, a+b, x), \\ [\phi(a, c, x)]^2 &> \phi(a+v, c, x)\phi(a-v, c, x), \end{aligned}$$

and

$$\begin{aligned} A(\phi(a+v, a+b, x), \phi(a-v, a+b, x)) &> \phi(a, a+b, x) \\ &> G(\phi(a+v, a+b, x), \phi(a-v, a+b, x)), \end{aligned}$$

where  $A(\alpha, \beta) = \frac{\alpha+\beta}{2}$  and  $G(\alpha, \beta) = \sqrt{\alpha\beta}$  are the arithmetic and geometric means and

$$\phi(a, b, x) = \sum_{n=0}^\infty \frac{(a)_n}{(b)_n} \frac{x^n}{n!}$$

is the Kummer confluent hypergeometric function [25, 28].

The Kummer confluent hypergeometric  $k$ -function is defined by

$$\phi_k(a, b, x) = \sum_{n=0}^\infty \frac{(a)_{n,k}}{(b)_{n,k}} \frac{x^n}{n!},$$

where

$$(a)_{n,k} = a(a+k)(a+2k)\cdots[a+(n-1)k]$$

for  $n \geq 1$  and  $k > 0$  with  $(a)_{0,k} = 1$  is the Pochhammer  $k$ -symbol, which can also be rewritten as

$$(a)_{n,k} = \frac{\Gamma_k(a + nk)}{\Gamma_k(a)}$$

and the gamma  $k$ -function  $\Gamma_k(a)$  is defined [5, 20, 21] by

$$\Gamma_k(a) = \int_0^\infty t^{a-1} e^{-t^k/k} dt$$

In 2012, Mubeen [14] introduced the  $k$ -analogue of Kummer’s transformation as

$$\phi_k(a, b, x) = e^x \phi_k(a, b - a, -x). \tag{1.3}$$

In Section 2, we prepare two lemmas. In Section 3, we discuss applications of some integral inequalities such as the Chebychev integral inequality. In Section 4, we prove the logarithmic convexity of the extended gamma function. In the last section, we introduce a mean inequality of Turán type for the Kummer confluent hypergeometric  $k$ -function.

## 2 Lemmas

In order to obtain our main results, we need the following lemmas.

**Lemma 2.1** (Chebychev integral inequality [6, 7, 11, 23]) *Let  $f, g, h : I \subseteq \mathbb{R} \rightarrow \mathbb{R}$  be mappings such that  $h(x) \geq 0$ ,  $h(x)f(x)g(x)$ ,  $h(x)f(x)$ , and  $h(x)g(x)$  are integrable on  $I$ . If  $f(x)$  and  $g(x)$  are synchronous (or asynchronous, respectively) on  $I$ , that is,*

$$[f(x) - f(y)][g(x) - g(y)] \gtrless 0$$

for all  $x, y \in I$ , then

$$\int_I h(x) dx \int_I h(x)f(x)g(x) dx \gtrless \int_I h(x)f(x) dx \int_I h(x)g(x) dx.$$

**Lemma 2.2** (Hölder’s inequality [29, 30]) *Let  $p$  and  $q$  be positive real numbers such that  $\frac{1}{p} + \frac{1}{q} = 1$  and  $f, g : [c, d] \rightarrow \mathbb{R}$  be integrable functions. Then*

$$\left| \int_c^d f(x)g(x) dx \right| \leq \left[ \int_c^d |f(x)|^p dx \right]^{1/p} \left[ \int_c^d |g(z)|^q dx \right]^{1/q}.$$

## 3 Inequalities involving the extended gamma function via the Chebychev integral inequality

In this section, we prove some inequalities involving the extended gamma function via the Chebychev integral inequality in Lemma 2.1.

**Theorem 3.1** *Let  $m, p$  and  $r$  be positive real numbers such that  $p > r > -m$ . If  $r(p - m - r) \geq 0$ , then*

$$\Gamma_b(m)\Gamma_b(p) \geq \Gamma_b(p - r)\Gamma_b(m + r). \tag{3.1}$$

*Proof* Let us define the mappings  $f, g, h : [0, \infty) \rightarrow [0, \infty)$  given by

$$f(t) = t^{p-r-m}, \quad g(t) = t^r, \quad \text{and} \quad h(t) = t^{m-1}e^{-t-bt^{-1}}.$$

If  $r(p - m - r) \geq 0$ , then we can claim that the mappings  $f$  and  $g$  are synchronous (asynchronous) on  $(0, \infty)$ . Thus, by applying the Chebychev inequality on  $I = (0, \infty)$  to the functions  $f, g$  and  $h$  defined above, we can write

$$\begin{aligned} \int_0^\infty t^{m-1}e^{-t-bt^{-1}} dt \int_0^\infty t^{p-r-m}t^r t^{m-1}e^{-t-bt^{-1}} dt \\ \geq \int_0^\infty t^{p-r-m}t^{m-1}e^{-t-bt^{-1}} dt \int_0^\infty t^r t^{m-1}e^{-t-bt^{-1}} dt. \end{aligned}$$

This implies that

$$\begin{aligned} \int_0^\infty t^{m-1}e^{-t-bt^{-1}} dt \int_0^\infty t^{p-1}e^{-t-bt^{-1}} dt \\ \geq \int_0^\infty t^{p-r-1}e^{-t-bt^{-1}} dt \int_0^\infty t^{m+r-1}e^{-t-bt^{-1}} dt. \end{aligned}$$

By (1.1), we acquire the required inequality (3.1). □

**Corollary 3.1** *If  $p > 0$  and  $q \in \mathbb{R}$  with  $|q| < p$ , then*

$$\Gamma_b(p) \leq [\Gamma_b(p - q)\Gamma_b(p + q)]^{1/2}.$$

*Proof* By setting  $m = p$  and  $r = q$  in Theorem 3.1, we obtain  $r(p - m - r) = -q^2 \leq 0$  and then the inequality (3.1) provides the desired Corollary 3.1. □

**Theorem 3.2** *If  $m, n > 0$  are similarly (oppositely) unitary, then*

$$\Gamma_b(m + n + b) \geq \frac{\Gamma_b(m + b + 1)\Gamma_b(n + b + 1)}{\Gamma_b(b + 2)}.$$

*Proof* Consider the mappings  $f, g, h : [0, \infty) \rightarrow [0, \infty)$  defined by

$$f(t) = t^{m-1}, \quad g(t) = t^{n-1}, \quad \text{and} \quad h(t) = t^{b+1}e^{-t-bt^{-1}}.$$

Now if the condition  $(m-1)(n-1) \geq 0$  holds, then the Chebychev integral inequality applied to the functions  $f, g$ , and  $h$  means

$$\int_0^\infty t^{b+1}e^{-t-bt^{-1}} dt \int_0^\infty t^{m-1}t^{n-1}t^{b+1}e^{-t-bt^{-1}} dt$$

$$\geq \int_0^\infty t^{m-1}t^{b+1}e^{-t-bt^{-1}} dt \int_0^\infty t^{n-1}t^{b+1}e^{-t-bt^{-1}} dt.$$

This implies that

$$\begin{aligned} \int_0^\infty t^{b+1}e^{-t-bt^{-1}} dt \int_0^\infty t^{m+n+b-1}e^{-t-bt^{-1}} dt \\ \geq \int_0^\infty t^{m+b}e^{-t-bt^{-1}} dt \int_0^\infty t^{n+b}e^{-t-bt^{-1}} dt. \end{aligned}$$

By the definition of the extended gamma function, we have

$$\Gamma_b(b+2)\Gamma_b(m+n+b) \geq \Gamma_b(m+b+1)\Gamma_b(n+b+1),$$

or

$$\Gamma_b(m+n+b) \geq \frac{\Gamma_b(m+b+1)\Gamma_b(n+b+1)}{\Gamma_b(b+2)}.$$

The required proof is complete. □

**Corollary 3.2** *If  $b = 0$ , then*

$$\Gamma(m+n) \geq mn\Gamma(m)\Gamma(n).$$

**Theorem 3.3** *If  $m$  and  $n$  are positive real numbers such that  $m$  and  $n$  are similarly (oppositely) unitary, then*

$$\Gamma_b(b+1)\Gamma_b(m+n+b+1) \geq \Gamma_b((m+b+1)\Gamma_b(n+b+1), \quad b \geq 0.$$

*Proof* Consider the mappings  $f, g, h : [0, \infty) \rightarrow [0, \infty)$  defined by

$$f(t) = t^m, \quad g(t) = t^n, \quad \text{and} \quad h(t) = t^b e^{-t-bt^{-1}}.$$

If the conditions of Theorem 3.1 hold, then the mappings  $f$  and  $g$  are synchronous (asynchronous) on  $[0, \infty)$ . Thus, by applying the Chebychev integral inequality in Lemma 2.1 to the functions  $f, g$  and  $h$  defined above, we have

$$\begin{aligned} \int_0^\infty t^b e^{-t-bt^{-1}} dt \int_0^\infty t^m t^n t^b e^{-t-bt^{-1}} dt \\ \geq \int_0^\infty t^m t^b e^{-t-bt^{-1}} dt \int_0^\infty t^n t^b e^{-t-bt^{-1}} dt. \end{aligned}$$

This implies that

$$\begin{aligned} \int_0^\infty t^b e^{-t-bt^{-1}} dt \int_0^\infty t^{m+n+b} e^{-t-bt^{-1}} dt \\ \geq \int_0^\infty t^{m+b} e^{-t-bt^{-1}} dt \int_0^\infty t^{n+b} e^{-t-bt^{-1}} dt. \end{aligned}$$

Thus by the definition of the extended gamma function, we have

$$\Gamma_b(b + 1)\Gamma_b(m + n + b + 1) \geq \Gamma_b(m + b + 1)\Gamma_b(n + b + 1).$$

The required proof is complete. □

**Corollary 3.3** *If  $b = 0$ , then*

$$\Gamma(m + n) \geq \frac{mn\Gamma(m)\Gamma(n)}{m + n}.$$

### 4 Log-convexity of the extended gamma function

It is well known that, if  $f > 0$  and  $\ln f$  is convex, then  $f$  is said to be a logarithmically convex function. Every logarithmically convex must be convex. See [15] and [18, Remark 1.9]. In this section, we verify the log-convexity of the extended gamma function.

**Theorem 4.1** *The extended gamma function  $\Gamma_b : (0, \infty) \rightarrow \mathbb{R}$  is logarithmically convex.*

*Proof* Let  $p$  and  $q$  be positive numbers such that  $\frac{1}{p} + \frac{1}{q} = 1$ . Since

$$\Gamma_b\left(\frac{x}{p} + \frac{y}{q}\right) \leq [\Gamma_b(x)]^{1/p}[\Gamma_b(y)]^{1/q},$$

see [4], letting  $\lambda = \frac{1}{p}$  and  $(1 - \lambda) = \frac{1}{q}$  leads to

$$\Gamma_b[\lambda x + (1 - \lambda)y] \leq [\Gamma_b(x)]^\lambda[\Gamma_b(y)]^{(1-\lambda)}.$$

As a result, the function  $\Gamma_b$  is logarithmically convex. □

### 5 A mean inequality of the Turán type for the Kummer confluent hypergeometric $k$ -function

In this section, we present a mean inequality involving the Kummer confluent hypergeometric  $k$ -function. For this purpose, we consider the relation

$$\phi_k(a + k, b, x) - \phi_k(a, b, x) = \frac{kx}{b}\phi_k(a + k, b + k, x), \quad k > 0. \tag{5.1}$$

**Theorem 5.1** *For  $a, b, k > 0$  and  $v \in \mathbb{N}$  with  $a, b \geq v - k$ , the inequality*

$$[\phi_k(a, a + b, x)]^2 > \phi_k(a + v, a + b, x)\phi_k(a - v, a + b, x) \tag{5.2}$$

*is valid for all nonzero  $x \in \mathbb{R}$ .*

*First proof* Assume that  $x > 0$ . For  $c \neq 0, -1, -2, \dots$ , define

$$f_{v,k}(x) = [\phi_k(a, c, x)]^2 - \phi_k(a + v, c, x)\phi_k(a - v, a + b, x)$$

and

$$f_{v+k,k}(x) = \phi_k(a, c, x)^2 - \phi_k(a + v + k, c, x)\phi_k(a - v - k, a + b, x).$$

From (5.1), it follows that

$$\begin{aligned} f_{v+k,k}(x) - f_{v,k}(x) &= \phi_k(a + v, c, x)\phi_k(a - v, c, x) \\ &\quad - \phi_k(a + v + k, c, x)\phi_k(a - v - k, c, x) \\ &= \phi_k(a - v, c, x)[\phi_k(a + v, c, x) - \phi_k(a + v + k, c, x)] \\ &\quad + \phi_k(a + v + k, c, x)[\phi_k(a - v, c, x) - \phi_k(a - v - k, c, x)] \\ &= \phi_k(a - v, c, x)\left(\frac{-kx}{c}\right)\phi_k(a + v + k, c + k, x) \\ &\quad + \phi_k(a + v + k, c, x)\left(\frac{kx}{c}\right)\phi_k(a - v, c + k, x) \\ &= \frac{kx}{c}g_{v,k}(x), \end{aligned}$$

where

$$\begin{aligned} g_{v,k}(x) &= \phi_k(a + v + k, c, x)\phi_k(a - v, c + k, x) \\ &\quad - \phi_k(a - v, c, x)\phi_k(a + v + k, c + k, x). \end{aligned}$$

Accordingly, by the Cauchy product, we have

$$\begin{aligned} g_{v,k}(x) &= \sum_{s=0}^{\infty} \sum_{r=0}^s \frac{(a + v + k)_{s,k}(a - v)_{s-r,k}}{r!(s - r)!} \\ &\quad \times \left[ \frac{1}{(c)_{s,k}(c + k)_{s-r,k}} - \frac{1}{(c)_{s-r,k}(c + k)_{s,k}} \right] x^s \\ &= \sum_{s=0}^{\infty} \sum_{r=0}^s \frac{(a + v + k)_{s,k}(a - v)_{s-r,k}}{r!(s - r)!} \left[ \frac{(c + mk) - (c + nk - mk)}{c(c + k)_{s,k}(c + k)_{s-r,k}} \right] x^s \\ &= \frac{k}{c} \sum_{s=0}^{\infty} \sum_{r=0}^s T_{s,r,k}(2r - s)x^s, \end{aligned}$$

where

$$T_{s,r,k} = \frac{(a + v + k)_{s,k}(a - v)_{s-r,k}}{r!(s - r)!(c + k)_{s,k}(c + k)_{s-r,k}}.$$

If  $s$  is even, then

$$\begin{aligned} \sum_{r=0}^s T_{s,r,k}(2r - s) &= \sum_{r=0}^{s/2-1} T_{s,r,k}(2r - s) + \sum_{r=s/2+1}^s T_{s,r,k}(2r - s) \\ &= \sum_{r=0}^{s/2-1} T_{s,r,k}(2r - s) + \sum_{r=0}^{s/2-1} T_{s,s-r,k}(2(s - r) - s) \end{aligned}$$



$$= \sum_{r=0}^{\lceil (s-1)/2 \rceil} (T_{s,s-r,k} - T_{s,r,k})(s - 2r),$$

where  $\lceil x \rceil$  denotes the ceiling function whose value is the greatest integer not more than  $x$ . Similarly, if  $s$  is odd,

$$\sum_{r=0}^s T_{s,r,k}(2r - s) = \sum_{r=0}^{\lceil (s-1)/2 \rceil} (T_{s,s-r,k} - T_{s,r,k})(s - 2r).$$

Accordingly,

$$\begin{aligned} f_{v+k,k}(x) - f_{v,k}(x) &= \frac{kx}{c} g_{v,k}(x) \\ &= \frac{k^2 x}{c^2} \sum_{s=1}^{\infty} \sum_{r=0}^{\lceil (s-1)/2 \rceil} (T_{s,s-r,k} - T_{s,r,k})(s - 2r)x^s. \end{aligned} \tag{5.3}$$

Carefully simplifying gives

$$\begin{aligned} T_{s,s-r,k} - T_{s,r,k} &= \frac{(a + v + k)_{s,k}(a - v)_{s,k} - (a + v + 1)_{s,k}(a - v)_{s,k}}{r!(s - r)!(c + k)_{s-r,k}(c + k)_{s,k}} \\ &= \frac{(a + v + k)_{s,k}(a - v)_{s,k}}{r!(s - r)!(c + k)_{s-r,k}(c + k)_{s,k}} \left[ \frac{(a + v + k)_{s-r,k}}{(a + v + k)_{s,k}} - \frac{(a - v)_{s-r,k}}{(a - v)_{s,k}} \right] \\ &= \frac{(a + v + k)_{s,k}(a - v)_{s,k}}{r!(s - r)!(c + k)_{s-r,k}(c + k)_{s,k}} [h_k(a + v + k) - h_k(a - v)], \end{aligned} \tag{5.4}$$

where  $h_k(x) = \frac{(x)_{s-r,k}}{(x)_{s,k}}$ . For  $x > 0$  and  $s - r > r$ , that is,  $\lceil \frac{s-1}{2} \rceil \geq r$ , the logarithmic derivatives of  $h_k$  is

$$\frac{h'_k(x)}{h_k(x)} = \psi_k(x + (s - r)k) - \psi_k(x + nk) > 0,$$

where  $\psi_k = \frac{\Gamma'_k}{\Gamma_k}$  is the digamma  $k$ -function (see [5, 10, 15]). Hence, the function  $h_k$  is increasing under the condition stated. This fact together with the aid of (5.3) and (5.4) yields

$$\begin{aligned} f_{v+k,k}(x) - f_{v,k}(x) &= \frac{kx}{c} g_{v,k}(x) \\ &= \frac{k^2 x}{c^2} \sum_{s=1}^{\infty} \sum_{r=0}^{\lceil (s-1)/2 \rceil} (T_{s,s-r,k} - T_{s,r,k})(s - 2r)x^s > 0, \end{aligned} \tag{5.5}$$

where  $a \geq v \geq 0$ ,  $x > 0$ ,  $c + k > 0$ , and  $c \neq 0$ . Consequently, from (5.5), it follows that

$$f_{v+k,k}(x) = [f_{v+k,k}(x) - f_{v,k}(x)] + [f_{v,k}(x) - f_{v-k,k}(x)] + \dots + [f_{1,k}(x) - f_{0,k}(x)]$$

is positive for  $a \geq v \geq v - k \geq v - 2k \geq \dots \geq 0$  and  $f_{0,k}(x) = 0$ . Now replacing  $v$  by  $v - k$  shows that

$$f_{v,k}(x) > 0, \quad x > 0, v \in \mathbb{N}, a \geq v - k. \tag{5.6}$$

Therefore, the function  $f_{v,k}$  is absolutely monotonic on  $(0, \infty)$ , that is,  $f_{v,k}^{(\ell)}(x) > 0$  for  $\ell = 0, 1, 2, \dots$ . This proves Theorem 5.1 for the case  $x > 0$ .

Now suppose that  $x < 0$ ,  $a, b > 0$ , and  $v \in \mathbb{N}$  with  $a, b \geq v - k$ . Since  $\phi_k(a, c, x)$  is symmetric in  $a$  and  $b$ , by interchanging  $a$  and  $b$  in Theorem 5.1, we obtain

$$\phi_k(b, a + b, -x)^2 - \phi_k(b + v, a + b, -x)\phi_k(b - v, a + b, -x) > 0.$$

By using Kummer's transformation (1.3), we have

$$e^{-2x}\phi_k(a, a + b, x)^2 - e^{-2x}\phi_k(a - v, a + b, x)\phi_k(a + v, a + b, x) > 0.$$

Thus, Theorem 5.1 also holds for  $x < 0$ . □

*Second proof* Since

$$\begin{aligned} (a)_{n,k} &= a(a + k)(a + 2k) \cdots (a + (n - 1)k) \\ &= k^n \frac{a}{k} \left(\frac{a}{k} + 1\right) \left(\frac{a}{k} + 2\right) \cdots \left(\frac{a}{k} + (n - 1)\right) \\ &= k^n \left(\frac{a}{k}\right)_n, \end{aligned}$$

it follows that

$$\phi_k(a, b; x) = \sum_{n=0}^{\infty} \frac{(a)_{n,k}}{(b)_{n,k}} \frac{x^n}{n!} = \sum_{n=0}^{\infty} \frac{k^n (a/k)_n}{k^n (b/k)_n} \frac{x^n}{n!} = \phi\left(\frac{a}{k}, \frac{b}{k}, x\right).$$

Replacing  $a$  and  $b$  by  $\frac{a}{k}$  and  $\frac{b}{k}$ , respectively, gives Theorem 5.1. □

**Corollary 5.1** *If  $a > 0$  and  $c + k > 0$  with  $c \neq 0$ , then the inequality*

$$[\phi_k(a, c, x)]^2 \geq \phi_k(a - v, c, x)\phi_k(a + v, c, x)$$

*holds for any  $v \in \mathbb{N}$  with  $a \geq v - k$ .*

*Proof* This follows directly from the proof of Theorem 5.1 and the fact that the equation (5.6) holds under the conditions  $c + k > 0$  and  $c \neq 0$ . □

**Corollary 5.2** *If  $v \in \mathbb{N}$  and  $a, b \geq v$ , then*

$$\begin{aligned} A(\phi_k(a + v, a + b, x), \phi_k(a - v, a + b, x)) &> \phi_k(a, a + b, x) \\ &> G(\phi_k(a + v, a + b, x), \phi_k(a - v, a + b, x)) \end{aligned} \tag{5.7}$$

for all nonzero  $x \in \mathbb{R}$ , where  $A$  and  $G$  are, respectively, the arithmetic and geometric means.

*Proof* First assume  $x \geq 0$  and  $a, b \geq v$  for  $v \in \mathbb{N}$ . Then the left hand side inequality in (5.7) is a direct consequence of the facts that

$$A((a + v)_{s,k}, (a - v)_{s,k}) = (a)_{s,k}$$

for  $s = 0, 1$  and

$$A((a + v)_{s,k}, (a - v)_{s,k}) > (a)_{s,k}$$

for  $s \geq 2$ . Hence, by induction, we have

$$\begin{aligned} A(\phi_k(a + v, a + b, x), \phi_k(a - v, a + b, x)) &= \sum_{s=0}^{\infty} \frac{A((a + v)_{s,k}, (a - v)_{s,k})x^s}{(a + b)_{s,k}s!} \\ &> \sum_{s=0}^{\infty} \frac{(a)_{s,k}x^s}{(a + b)_{s,k}s!} = \phi_k(a, a + b, x). \end{aligned}$$

For  $x \geq 0$ , the right hand side inequality in (5.7) follows from taking square root of (5.2). The proof of Corollary 5.2 for  $x \geq 0$  is thus complete.

Now assume  $x < 0$  with  $a, b \geq v$ . Interchanging  $a$  and  $b$  in (5.7) one arrives at

$$\begin{aligned} A(\phi_k(b + v, a + b, -x), \phi_k(b - v, a + b, -x)) &> \phi_k(b, a + b, -x) \\ &> G(\phi_k(b + v, a + b, -x), \phi_k(b - v, a + b, -x)). \end{aligned}$$

Making use of the  $k$ -analogue of Kummer's transformation and the homogeneity of  $A$  and  $G$  acquires

$$\begin{aligned} e^{-x}A(\phi_k(a - v, a + b, x), \phi_k(a + v, a + b, x)) &> e^{-x}\phi_k(a, a + b, x) \\ &> e^{-x}G(\phi_k(a - v, a + b, x), \phi_k(a + v, a + b, x)). \end{aligned}$$

Consequently, Theorem 5.7 also follows for  $x < 0$ . □

*Remark 5.1* In Section 5, we have established a Turán type and mean inequality for  $k$ -analogue of the Kummer confluent hypergeometric function. If we let  $k \rightarrow 1$ , then we can conclude to the corresponding inequalities of the confluent hypergeometric function.

*Remark 5.2* In [22], some inequalities of the Turán type for confluent hypergeometric functions of the second kind were also discovered.

*Remark 5.3* By the way, we note that the papers [8, 9, 12, 13, 26, 32, 33] belong to the same series in which inequalities and complete monotonicity for functions involving the gamma function  $\Gamma(x)$  and the logarithmic function  $\ln(1 + x)$  were discussed.

*Remark 5.4* This paper is a slightly revised version of the preprint [16].

## 6 Conclusions

In this paper, we present some inequalities involving the extended gamma function  $\Gamma_b(z)$  via some classical inequalities such as the Chebychev inequality for synchronous (or asynchronous, respectively) mappings, give a new proof of the log-convexity of the extended gamma function  $\Gamma_b(z)$  by using the Hölder inequality, and introduce a Turán type mean inequality for the Kummer confluent  $k$ -hypergeometric function  $\phi(z)$ .

### Acknowledgement

The authors appreciate the anonymous referees for their careful corrections to and valuable comments on the original version of this paper.

### Competing interests

The authors declare that they have no competing interests.

### Authors' contributions

All authors contributed equally to the manuscript and read and approved the final manuscript.

### Author details

<sup>1</sup>Department of Mathematics, College of Arts and Science at Wadi Aldawaser, Prince Sattam Bin Abdulaziz University, 11991, Riyadh Region, Kingdom of Saudi Arabia. <sup>2</sup>Institute of Mathematics, Henan Polytechnic University, 454010, Jiaozuo, Henan, China. <sup>3</sup>School of Mathematical Sciences, Tianjin Polytechnic University, 300387, Tianjin, China. <sup>4</sup>Department of Mathematics, International Islamic University, Islamabad, Pakistan. <sup>5</sup>Department of Mathematics, University of Sargodha, Sargodha, Pakistan.

### References

1. R. W. Barnard, M. B. Gordy, and K. C. Richards, *A note on Turán type and mean inequalities for the Kummer function*, J. Math. Anal. Appl. **349** (2009), no. 1, 259–263; available online at <http://dx.doi.org/10.1016/j.jmaa.2008.08.024>.
2. M. A. Chaudhry, A. Qadir, M. Rafique, and S. M. Zubair, *Extension of Euler's beta function*, J. Comput. Appl. Math. **78** (1997), no. 1, 19–32; available online at [http://dx.doi.org/10.1016/S0377-0427\(96\)00102-1](http://dx.doi.org/10.1016/S0377-0427(96)00102-1).
3. M. A. Chaudhry and S. M. Zubair, *On the decomposition of generalized incomplete gamma functions with applications of Fourier transforms*, J. Comput. Appl. Math. **59** (1995), no. 3, 253–284; available online at [http://dx.doi.org/10.1016/0377-0427\(94\)00026-W](http://dx.doi.org/10.1016/0377-0427(94)00026-W).
4. M. A. Chaudhry and S. M. Zubair, *On a Class of Incomplete Gamma Functions with Applications*, Chapman & Hall/CRC, Boca Raton, FL, 2002.
5. R. Díaz and E. Pariguan, *On hypergeometric functions and Pochhammer  $k$ -symbol*, Divulg. Mat. **15** (2007), no. 2, 179–192.
6. S. S. Dragomir, R. P. Agarwal, and N. S. Barnett, *Inequalities for beta and gamma functions via some classical and new integral inequalities*, J. Inequal. Appl. **5** (2000), no. 2, 103–165; available online at <http://dx.doi.org/10.1155/S1025583400000084>.
7. S. S. Dragomir and S. Wang, *Applications of Ostrowski's inequality to the estimation of error bounds for some special means and for some numerical quadrature rules*, Appl. Math. Lett. **11** (1998), no. 1, 105–109; available online at [http://dx.doi.org/10.1016/S0893-9659\(97\)00142-0](http://dx.doi.org/10.1016/S0893-9659(97)00142-0).
8. B.-N. Guo and F. Qi, *Monotonicity of functions connected with the gamma function and the volume of the unit ball*, Integral Transforms Spec. Funct. **23** (2012), no. 9, 701–708; available online at <http://dx.doi.org/10.1080/10652469.2011.627511>.
9. P. Ivády, *A note on a gamma function inequality*, J. Math. Inequal. **3** (2009), no. 2, 227–236; available online at <http://dx.doi.org/10.7153/jmi-03-23>.
10. C. G. Kokologiannaki, *Properties and inequalities of generalized  $k$ -gamma, beta and zeta functions*, Int. J. Contemp. Math. Sci. **5** (2010), no. 13–16, 653–660.
11. P. Kumar, S. P. Singh, and S. S. Dragomir, *Some inequalities involving beta and gamma functions*, Nonlinear Anal. Forum **6** (2001), no. 1, 143–150.
12. P. A. Kupán, G. Márton, and R. Szász, *A result regarding monotonicity of the Gamma function*, Acta Univ. Sapientiae Math. **9** (2017), no. 2, 291–302; available online at <https://doi.org/10.1515/ausm-2017-0022>.
13. P. A. Kupán and R. Szász, *Monotonicity theorems and inequalities for the gamma function*, Math. Inequal. Appl. **17** (2014), no. 1, 149–159; available online at <http://dx.doi.org/10.7153/mia-17-11>.
14. S. Mubeen,  *$k$ -analogue of Kummer's first formula*, J. Inequal. Spec. Funct. **3** (2012), no. 3, 41–44.
15. K. Nantomah, and E. Prempeh, *The  $k$ -analogues of some inequalities for the digamma function*, Math. Aeterna, **4** (2014), no. 3–4, 269–273.
16. K. S. Nisar, F. Qi, G. Rahman, S. Mubeen, and M. Arshad, *Some inequalities involving the extended gamma and beta functions*, HAL preprint (2018), available online at <https://hal.archives-ouvertes.fr/hal-01701647>.
17. F. W. J. Olver, D. W. Lozier, R. F. Boisvert, and C. W. Clark (eds.), *NIST Handbook of Mathematical Functions*, Cambridge University Press, New York, 2010; available online at <http://dlmf.nist.gov/>.
18. F. Qi, *Bounds for the ratio of two gamma functions*, J. Inequal. Appl. **2010** (2010), Article ID 493058, 84 pages; available online at <http://dx.doi.org/10.1155/2010/493058>.

19. F. Qi, *Limit formulas for ratios between derivatives of the gamma and digamma functions at their singularities*, *Filomat* **27** (2013), no. 4, 601–604; available online at <http://dx.doi.org/10.2298/FIL1304601Q>.
20. F. Qi, *Parametric integrals, the Catalan numbers, and the beta function*, *Elem. Math.* **72** (2017), no. 3, 103–110; available online at <http://dx.doi.org/10.4171/EM/332>.
21. F. Qi, A. Akkurt, and H. Yildirim, *Catalan numbers, k-gamma and k-beta functions, and parametric integrals*, *J. Comput. Anal. Appl.* **25** (2018), no. 6, 1036–1042.
22. F. Qi, R. Bhukya, and V. Akavaram, *Some inequalities of the Turán type for confluent hypergeometric functions of the second kind*, *Stud. Univ. Babeş-Bolyai Math.* **64** (2019), no. 1, 63–70; available online at <https://doi.org/10.24193/subbmth.2019.1.06>.
23. F. Qi, L.-H. Cui, and S.-L. Xu, *Some inequalities constructed by Tchebysheff's integral inequality*, *Math. Inequal. Appl.* **2** (1999), no. 4, 517–528; available online at <http://dx.doi.org/10.7153/mia-02-42>.
24. F. Qi and B.-N. Guo, *Integral representations and complete monotonicity of remainders of the Binet and Stirling formulas for the gamma function*, *Rev. R. Acad. Cienc. Exactas Fis. Nat. Ser. A Mat. RACSAM* **111** (2017), no. 2, 425–434; available online at <http://dx.doi.org/10.1007/s13398-016-0302-6>.
25. F. Qi and B.-N. Guo, *On the sum of the Lah numbers and zeros of the Kummer confluent hypergeometric function*, *Acta Univ. Sapientiae Math.* **10** (2018), no. 1, 125–133; available online at <https://doi.org/10.2478/ausm-2018-0011>.
26. F. Qi and W.-H. Li, *Integral representations and properties of some functions involving the logarithmic function*, *Filomat* **30** (2016), no. 7, 1659–1674; available online at <https://doi.org/10.2298/FIL1607659Q>.
27. F. Qi, X.-T. Shi, and F.-F. Liu, *Several identities involving the falling and rising factorials and the Cauchy, Lah, and Stirling numbers*, *Acta Univ. Sapientiae Math.* **8** (2016), no. 2, 282–297; available online at <http://dx.doi.org/10.1515/ausm-2016-0019>.
28. E. D. Rainville, *Special Functions*, The Macmillan Co., New York, 1960.
29. W. Rudin, *Real and complex analysis*, 3rd edn, McGraw-Hill Book Co., New York, 1987.
30. J.-F. Tian and M.-H. Ha, *Properties of generalized sharp Hölder's inequalities*, *J. Math. Inequal.* **11** (2017), no. 2, 511–525; available online at <https://doi.org/10.7153/jmi-11-42>.
31. Z.-H. Yang and J.-F. Tian, *Monotonicity and inequalities for the gamma function*, *J. Inequal. Appl.* **2017**, 2017:317, 15 pages; available online at <https://doi.org/10.1186/s13660-017-1591-9>.
32. Z.-H. Yang and J.-F. Tian, *Monotonicity and sharp inequalities related to gamma function*, *J. Math. Inequal.* **12** (2018), no. 1, 1–22; available online at <http://dx.doi.org/10.7153/jmi-2018-12-01>.
33. J.-L. Zhao, B.-N. Guo, and F. Qi, *A refinement of a double inequality for the gamma function*, *Publ. Math. Debrecen* **80** (2012), no. 3–4, 333–342; available online at <http://dx.doi.org/10.5486/PMD.2012.5010>.