



HAL
open science

Control of fluctuations and heavy tails for heat variation in the two-time measurement framework

Tristan Benoist, Annalisa Panati, Renaud Raquépas

► **To cite this version:**

Tristan Benoist, Annalisa Panati, Renaud Raquépas. Control of fluctuations and heavy tails for heat variation in the two-time measurement framework. 2018. hal-01703374v1

HAL Id: hal-01703374

<https://hal.science/hal-01703374v1>

Preprint submitted on 7 Feb 2018 (v1), last revised 29 Oct 2018 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Control of fluctuations and heavy tails for heat variation in the two-time measurement framework

Tristan Benoist¹, Annalisa Panati², Renaud Raquépas^{3,4}

February 6, 2018

1. Université de Toulouse UPS
CNRS, Institut de mathématiques de Toulouse
F-31 062 Toulouse Cedex 9, France

2. Aix Marseille Univ, Université de Toulon
CNRS, CPT
Marseille, France

3. McGill University
Department of Mathematics and Statistics
1005–805 rue Sherbrooke Ouest
Montréal (Québec) H3A 0B9

4. Univ. Grenoble Alpes
CNRS, Institut Fourier
F-38 000 Grenoble, France

Abstract

We study heat fluctuations in the two-time measurement framework. For bounded perturbations, we give sufficient ultraviolet regularity conditions on the perturbation for the moments of the heat variation to be uniformly bounded in time, and for the Fourier transform of the heat variation distribution to be analytic and uniformly bounded in time in a complex neighborhood of 0.

On a set of canonical examples, with bounded and unbounded perturbations, we show that our ultraviolet conditions are essentially necessary. If the form factor of the perturbation does not meet our assumptions, the heat variation distribution exhibits heavy tails. The tails can be as heavy as preventing the existence of a fourth moment of the heat variation.

1 Introduction

The goal of this paper is to study statistical fluctuations of heat in locally perturbed quantum statistical systems. We particularly aim at highlighting the difference in the control of heat fluctuations between classical and quantum systems.

In classical statistical mechanics, it is generally admitted that heat fluctuations are controlled by the strength of the perturbation. Consider a closed system whose dynamics is governed by a time independent perturbed Hamiltonian $H_\lambda = H_0 + \lambda V$ and let $(\Xi_\lambda^t)_{t \in \mathbf{R}}$ be the flow it generates. Then, by conservation of energy (or the first law of thermodynamics), the total heat variation $\Delta Q := H_0 \circ \Xi_\lambda^t - H_0$ is equal to the work $\lambda V - \lambda V \circ \Xi_\lambda^t$. If the perturbation V is bounded, the heat variation ΔQ is therefore surely bounded uniformly in time. Hence, all the moments of ΔQ are uniformly bounded in time. Moreover, the Fourier transform of the law \mathbb{P}_t of ΔQ is entire analytic and uniformly bounded in time at any point of the complex plane. Boundedness of the perturbation V therefore trivially implies that the distribution \mathbb{P}_t of heat variation is light-tailed.¹

¹We say that the distribution of a random variable is heavy-tailed if its Fourier transform does not extend analytically to any open neighborhood of the real axis in the complex plane; and light-tailed if it extends analytically to the whole complex plane. For a probability measure P on \mathbf{R} , we use the terms characteristic function and Fourier transform interchangeably to mean the function that assigns to $\alpha \in \mathbf{R}$ the integral $\int_{\mathbf{R}} e^{i\alpha x} dP(x)$, also known as the Fourier–Stieltjes transform of P , always using this convention without any factor of 2π .

If V is unbounded, the fluctuations of ΔQ are expected to be controlled by the strength parameter λ . We illustrate that expectation with two examples in Section 2. In the first example, \mathbb{P}_t is neither heavy nor light-tailed, whereas in the second one, \mathbb{P}_t remains light-tailed even if V is no longer bounded.

Certainly motivated by this classical behavior, it is sometimes expected that λ controls the fluctuations of heat in quantum systems whether V is bounded or not. Our goal in this paper is to show that, for quantum systems, the value of λ is not the most determinant parameter for the control of the tails of heat variation when it is defined through a two-time measurement protocol. Actually, the ultraviolet (UV) regularity of V can play a crucial role in the control of the tails of ΔQ , even when V is bounded.

The definition of thermodynamic quantities as random variables is still debated in quantum statistical mechanics. Naive quantization from the classical definition might suggest defining \mathbb{P}_t as the spectral measure of the operator $\lambda V - \lambda e^{itH_\lambda} V e^{-itH_\lambda}$ with respect to the initial state, but this definition has no clear physical meaning. Moreover, the distribution of entropy production defined in the same fashion does not verify the transient fluctuation relation when the dynamic is time-reversal invariant [JOPP11, §4.6.10]². An alternative definition is provided by the two-time measurement protocol framework. It has become one of the prominent approaches to quantum thermodynamics and has been extensively studied in the literature; see for example [EHM09, CHT11, JOPP11]. It is by introducing this definition that Kurchan [Kur00] and Tasaki [Tas00] provided the first extensions of fluctuation relations to quantum systems. In the literature, this framework is also sometimes named *full statistics* or *full counting statistics*.

In this framework, the variation of heat is defined according to the following Gedankenexperiment. A first measurement of H_0 , the unperturbed Hamiltonian, is performed at initial time, yielding a result $E \in \text{spec } H_0$; the system then evolves according to H_λ , the perturbed Hamiltonian, for a time t ; and H_0 is once again measured, yielding a result $E' \in \text{spec } H_0$. The resulting heat variation is defined as the difference $\Delta Q := E' - E$ between the two measurement results.

For confined systems where the unperturbed Hamiltonian H_0 and the density matrix ρ representing the initial state are commuting matrices, the characteristic function of the measure \mathbb{P}_t assigning probabilities to the differences ΔQ in the above thought experiment is

$$\mathcal{E}_t(\alpha) := \text{tr}(e^{itH_\lambda} e^{i\alpha H_0} e^{-itH_\lambda} e^{-i\alpha H_0} \rho).$$

A rewriting of this expression in terms of algebraic objects that survive the thermodynamic limit serves as the basis for the construction of the generalization of this measure to infinitely extended systems; see Section 3.1.

As mentioned before, the probability measure \mathbb{P}_t is not equal to the spectral measure of $\lambda V - \lambda e^{itH_\lambda} V e^{-itH_\lambda}$ with respect to the initial state. Hence, even if V is bounded, \mathbb{P}_t is not guaranteed to have compact support in the thermodynamic limit. Note also that we no longer have the almost sure first law equality $\Delta Q = -\Delta W$ where the work ΔW would traditionally be defined by a measurement of H_0 and then of H_λ .³ In particular, with the two-time measurement definitions of thermodynamic quantities, there exists no joint law of work and heat variation, only the respective marginals.

Our main results provide sufficient conditions, in the thermodynamic limit, for the control of the tails of \mathbb{P}_t . We furthermore test the necessity of these conditions in different usual models of quantum statistical mechanics.

A sufficient condition for the exponential control of the tails of \mathbb{P}_t has been introduced in our partly co-authored work [BJP⁺15]. There, we considered quantum dynamical systems arising as the limit of a sequence of finite dimensional systems. In the present article, we work directly in the

²Section 4.6.10. *Full counting statistics and quantum fluxes: a comparison* of the published version corresponds to Section 5.10 of the version available on the *arXiv*.

³This choice allows for the recovery of Jarzynsky's equality and Crook's fluctuation relation when V depends on time.

thermodynamic limit, via the operator algebraic formulation of quantum statistical mechanics. In this setting, the condition given in [BJP⁺15] can be reformulated as follows.

Let δ be the generator of the unperturbed dynamics over the C^* -algebra \mathcal{O} of observables of the system (in finite dimension, $\delta(\cdot) = i[H_0, \cdot]$). Assume that the initial state ω over \mathcal{O} is invariant under the unperturbed dynamics, i.e. $\omega \circ e^{t\delta} = \omega$ for all $t \in \mathbf{R}$. Let V be a self-adjoint element of \mathcal{O} defining the perturbation. The result of [BJP⁺15] can be summarized by the implication

$$V \in \text{Dom } e^{i\frac{1}{2}\gamma\delta} \cap \text{Dom } e^{-i\frac{1}{2}\gamma\delta} \implies \sup_{t \in \mathbf{R}} \mathbb{E}_t(e^{\gamma|\Delta Q|}) < \infty \quad (1)$$

with \mathbb{E}_t the expectation with respect to \mathbb{P}_t and $\gamma \in \mathbf{R}_+$. We provide a proof of this implication in Section 3. In terms of control of the tails, Markov's inequality implies that if the left-hand side of (1) holds for some $\gamma > 0$, then there exists $C > 0$ such that

$$\sup_{t \in \mathbf{R}} \mathbb{P}_t(|\Delta Q| > E) \leq C e^{-\gamma E}$$

for any $E > 0$.

It follows trivially from (1) that if the left-hand side holds, any moment $\mathbb{E}_t(|\Delta Q|^m)$ is uniformly bounded in time. In Theorem 3.5, we show that such a uniform bound for the moments holds under weaker conditions:

$$V \in \text{Dom } \delta^n \implies \sup_{t \in \mathbf{R}} \mathbb{E}_t(\Delta Q^{2n+2}) < \infty \quad (2)$$

for any $n \in \mathbf{N}$. In terms of control of the tails, Markov's inequality implies that if $V \in \text{Dom } \delta^n$ for some $n \in \mathbf{N}$, then there exists $C > 0$ such that

$$\sup_{t \in \mathbf{R}} \mathbb{P}_t(|\Delta Q| > E) \leq C E^{-n}$$

for any $E > 0$.

Remark that both the left-hand side of (1) and (2) do not depend on the initial state. To obtain these implications, we merely assume the initial state is invariant under the unperturbed dynamics. In particular, the implications hold even if the system is not initially at equilibrium. Furthermore, in each statement, the left-hand side does not depend on λ ; only the value of the supremum in the corresponding right-hand side depends on it.

The implications of (1) on the large deviations of \mathbb{P}_t were discussed in [BJP⁺15]. In [BPP], two of us and Y. Pautrat discuss the implications of an assumption akin to the left-hand side of (1) on the large deviations of heat currents in autonomous out of equilibrium open systems. In particular, we prove a translation symmetry of the cumulant generating function, first proposed in [AGMT09]. We relate it to the large deviations of the conservation of heat currents and derive Onsager's reciprocity relations under a time-reversal invariance assumption.

Both implications (1) and (2) establish sufficient conditions for the control of the tails of \mathbb{P}_t . Such conditions are not necessary in classical systems. In the third part of the paper, starting in Section 4, we establish an essential necessity of these conditions for different usual models of quantum statistical mechanics, with V bounded or unbounded. Our first model corresponds to a fermionic impurity interacting with a quasi-free Fermi gas at equilibrium. The second model is the bosonic counterpart of the first one and is similar to the interaction of an harmonic oscillator with a quasi-free Bose gas. The third model is a van Hove Hamiltonian which corresponds to a quasi-free Bose gas at equilibrium interacting with prescribed classical charges. The classical counterparts of these two last models of bosons are discussed in Sections 2.1 and 2.2 respectively.

In the first model, a quasi-free Fermi gas, V is bounded and, in Theorem 4.4, we show the

equivalence

$$\begin{aligned} V \in \text{Dom } \delta^n &\iff \exists t_1 < t_2 : \int_{t_1}^{t_2} \mathbb{E}_t(\Delta Q^{2n+2}) dt < \infty \\ &\iff \sup_{t \in \mathbf{R}} \mathbb{E}_t(\Delta Q^{2n+2}) < \infty. \end{aligned} \quad (3)$$

In this model, the assumption $V \in \text{Dom } \delta^n$ is equivalent to $\int_{\mathbf{R}_+} e^{2n} |f(e)|^2 de < \infty$ with $e \mapsto |f(e)|^2$ the form factor of the perturbation V , where $e \in \mathbf{R}_+$ is the energy. Similarly, the assumption $V \in \text{Dom } e^{i\frac{1}{2}\gamma\delta} \cap \text{Dom } e^{-i\frac{1}{2}\gamma\delta}$ is equivalent to $\int_{\mathbf{R}_+} e^{\gamma e} |f(e)|^2 de < \infty$. Hence the left-hand side of (1) and (2) indeed correspond to UV regularity conditions. It is with similar models in mind that we choose to refer to the left-hand sides of (1) and (2) as UV regularity conditions. Note however that for quantum spin systems, these conditions are typically the consequence of some locality assumptions for V and H_0 ; see [BR97, §6.2] and [BPP, §4.1].

In the second and third models, V is unbounded and $e \mapsto |f(e)|^2$ is again its form factor. For both these models, we prove the equivalence

$$\begin{aligned} \int_{\mathbf{R}_+} e^{2n} |f(e)|^2 de < \infty &\iff \exists t_1 < t_2 : \int_{t_1}^{t_2} \mathbb{E}_t(\Delta Q^{2n+2}) dt < \infty \\ &\iff \sup_{t \in \mathbf{R}} \mathbb{E}_t(\Delta Q^{2n+2}) < \infty. \end{aligned} \quad (4)$$

For the first bosonic model, we prove an implication similar to (1), namely

$$\int_{\mathbf{R}_+} e^{\gamma e} |f(e)|^2 de < \infty \implies \exists \gamma' \in (0, \gamma) : \sup_{t \in \mathbf{R}} \mathbb{E}_t(e^{\gamma' |\Delta Q|}) < \infty. \quad (5)$$

For the second bosonic model, we prove the equivalence

$$\begin{aligned} \int_{\mathbf{R}_+} e^{\gamma e} |f(e)|^2 de < \infty &\iff \exists t_1 < t_2 : \int_{t_1}^{t_2} \mathbb{E}_t(e^{\gamma |\Delta Q|}) dt < \infty \\ &\iff \sup_{t \in \mathbf{R}} \mathbb{E}_t(e^{\gamma |\Delta Q|}) < \infty. \end{aligned} \quad (6)$$

For this last model, we moreover prove that \mathbb{P}_t is the law of an inhomogeneous Poisson process for every t , while its classical counterpart is the law of a Gaussian random variable.

The existence of these equivalences highlights the contrast between classical statistical mechanics and the quantum two-time measurement framework. For a quantum system, if one aims at controlling the tails of the heat variation, before discussing the value of λ , one has to assume that the interaction has sufficient UV regularity. Typically, if one uses a cutoff $f(e) = 0$ for e large enough, completely preventing the contribution of energy scales at which the validity of the model is no longer guaranteed, then the left-hand side of (1) holds for any $\gamma > 0$. This discussion is not necessary for classical systems.

Heuristically speaking, the physical picture can be understood in terms of the Fermi golden rule. According to this rule, the transition rates between different energy levels E and E' is given roughly by $T(E', E) := |\langle E', VE \rangle|^2$ with $\{|E\rangle\}_E$ the energy eigenstates of H_0 . The underlying physical intuition for the left-hand side of (1) [resp. (2)] is roughly a condition controlling $T(E', E)e^{\gamma|E'-E|}$ [resp. $T(E', E)|E'-E|^{2n}$] for large values of $|E'-E|$. Hence, these assumptions control the decay of the transition rates as $|E'-E|$ goes to ∞ .

Note that, while important for the control of the tails at finite λ , the UV regularisation assumptions become irrelevant in the limit $t \rightarrow \infty$ and then $\lambda \rightarrow 0$, if one assumes return to equilibrium. Indeed, adapting [JPPP15], it can be shown that in this limit \mathbb{P}_t converges weakly to a Dirac measure in 0.

As pointed out in the conclusion of [BFJP16], though \mathbb{P}_t is generally thought not directly accessible as it involves the projective measurement of non-local quantities, proposals have emerged allowing for a sampling of \mathbb{P}_t using an interaction with an auxiliary qubit [DCH⁺13, MDCP13, CBK⁺13, RCP14]. The proposal of [DCH⁺13] involves only a local interaction between the qubit and the system and therefore appears more appropriate for thermodynamic systems. Hence, \mathbb{P}_t may be experimentally accessible indirectly and a phenomenon akin to heavy tails may be observed.

Structure. The paper is organised as follows. In Section 2, we discuss shortly classical systems with emphasis on two models with V unbounded whose quantum counterpart will be studied in Section 4. In Section 3, we introduce the C^* -algebraic formalism of quantum statistical mechanics and prove (1) and (2). In Section 4, we introduce the three models for which we test the necessity of the UV regularity conditions. The proof of the implication (5) and the equivalences (3), (4) and (6) for these models are postponed to Section 5. Appendix A contains technical results on the self-adjointness of some operators involved in the definition of \mathbb{P}_t for unbounded V . Finally, in Appendix B, we show that the measure \mathbb{P}_t defined in our models through the algebraic formalism indeed emerges as the thermodynamic limit of measures describing two-time measurement protocols for some finite dimensional systems.

Acknowledgements. We would like to thank Vojkan Jakšić for informative discussions and suggestions about this project, and Claude-Alain Pillet and Yan Pautrat for useful comments about our work. The research of T.B. has been supported by ANR-11-LABX-0040-CIMI within the program ANR-11-IDEX-0002-02 and ANR project StoQ (ANR-14-CE25-0003-01). The research of A.P. was partially supported by ANR project SQFT (ANR-12-JS01-0008-01) and ANR grant NONSTOPS (ANR-17-CE40-0006-01, ANR17-CE40-0006-02, ANR-17-CE40-0006-03). The research of R.R. was partially supported by NSERC and FRQNT. R.R. would like to thank the Laboratoire de physique théorique at Université Paul Sabatier, where part of this research was conducted, for its support and hospitality.

2 A short classical detour

To give some perspective on our results we first make a short detour through classical statistical mechanics. We refer the reader to [Thi12, §3] for a detailed definition of classical Hamiltonian dynamical systems and to [CM76] for an overview of the possible problems arising when considering an infinite dimensional phase space.

Let (K, ϖ) be a connected smooth symplectic manifold. The manifold K is endowed with the meaning of a classical system phase space. Since we are concerned with a thermodynamic setting, the manifold K is not assumed to be finite dimensional.

The unperturbed Hamiltonian of the system is a continuously differentiable function $H_0 : K \rightarrow \mathbf{R}$. The perturbation is a second continuously differentiable function $V : K \rightarrow \mathbf{R}$, giving rise to the perturbed Hamiltonian $H := H_0 + V$.

Under minimal technical assumptions on H , it defines a continuous Hamiltonian flow $(\Xi^t)_{t \in \mathbf{R}}$ defined on K , and the evolution of any sufficiently regular function g satisfies

$$\frac{d}{dt}(g \circ \Xi^t) = \{g, H\} \circ \Xi^t,$$

where $\{\cdot, \cdot\}$ is the Poisson bracket induced by ϖ ; see the general conservation theorem in [CM76].

We define the heat variation between times 0 and t as the difference between the unperturbed energy at times t and 0: $\Delta Q := H_0 \circ \Xi^t - H_0$. We give some arguments for this choice in Remark 2.4.

Since $H \circ \Xi^t = H$ and $H = H_0 + V$, equivalently,

$$\Delta Q := V - V \circ \Xi^t. \quad (7)$$

Note that under minimal technical assumptions⁴ on V ,

$$\Delta Q = - \int_0^t \frac{d}{ds} V \circ \Xi^s ds = - \int_0^t \{V, H_0\} \circ \Xi^s ds.$$

This equality is the expression of the first law of thermodynamics for a closed system, the right-hand side being the work.

This definition through the work and the first law is useful when one is concerned with thermodynamic systems. Indeed, in some standard examples, the phase space needs to be enlarged to $\bar{K} \supset K$, a measurable space in which K is dense (see for example [JP98, SL77]). Then, the functions H_0 and V , and the flow Ξ^t are typically extended by continuity to \bar{K} . Though H_0 may be infinite at some points in \bar{K} , we assume that the perturbation V remains finite everywhere on \bar{K} . The function ΔQ on K is then extended by continuity to a function on \bar{K} using the extensions of V and Ξ^t .

With these definitions in mind, we introduce the statistical aspect of the model: the initial state of the system is a probability measure μ on the phase space \bar{K} and the heat variation ΔQ becomes a random variable. We are interested in characterizing the law \mathbb{P}_t of ΔQ with respect to μ .

Remark 2.1. We denote \mathbb{E}_t the expectation with respect to \mathbb{P}_t . The random variable ΔQ is denoted without index t to highlight that we do not study a stochastic process but only a family $(\mathbb{P}_t)_{t \in \mathbf{R}}$ of measures on \mathbf{R} .

Remark 2.2. Whenever the initial definition $\Delta Q = H_0 \circ \Xi^t - H_0$ makes sense, \mathbb{P}_t has characteristic function

$$\mathbb{E}_t(e^{i\alpha\Delta Q}) = \mu(e^{i\alpha(H_0 \circ \Xi^t - H_0)}) = \mu(e^{i\alpha H_0} \circ \Xi^t e^{-i\alpha H_0}). \quad (8)$$

This formula parallels an expression we will encounter for quantum systems (see Remark 3.2).

The following proposition is a trivial consequence of the definition of ΔQ .

Proposition 2.3. *If $C := \sup_{x \in \bar{K}} |V(x)| < \infty$, then for all $t \in \mathbf{R}$, $|\Delta Q| < 2C$ surely.*

As soon as the perturbation is bounded, the heat fluctuations are surely uniformly controlled in time. We will see in Section 4.1 that this implication is false in the context of quantum two-time measurements.

Remark 2.4. The separation between work and heat involves a certain amount of arbitrariness. More precisely, the choice of H_0 and V such that $H = H_0 + V$ is partly arbitrary. There is no reason another couple of functions would not be preferred. For example, if $W : K \rightarrow \mathbf{R}$ is a continuously differentiable function that remains finite on \bar{K} , then $H_0 + W$ and $V - W$ could also be a suitable couple of functions. The choice of H_0 in the definition of heat must therefore be motivated by other arguments that are a priori model dependent.

Nevertheless, since H is not time dependent, a quite interesting generic argument is given by the entropy balance equation. Assume, if it exists, that μ is the Gibbs measure at inverse temperature $\beta > 0$ for the Hamiltonian H_0 . Then the choice of the splitting $H = H_0 + V$ for the definition of ΔQ is motivated by the μ -almost sure validity of the entropy balance equation

$$\beta\Delta Q = -\log \frac{d\mu_{-t}}{d\mu}, \quad (9)$$

⁴See the conservation theorems in [CM76].

where $\mu_s := \mu \circ \Xi^{-s}$ is the time evolution of the measure μ induced by the flow $(\Xi^s)_{s \in \mathbf{R}}$. Averaging, we recover the mean entropy balance equation

$$\beta \mathbb{E}_t(\Delta Q) = - \int \log \frac{d\mu_{-t}}{d\mu} d\mu =: S(\mu|\mu_{-t}) \quad (10)$$

where $S(\mu|\nu)$ is the relative entropy of μ with respect to ν . A similar argument holds for composite systems with different parts at different inverse temperatures β_i . Then, the choice of H_0 is motivated by the μ -almost sure entropy balance equation $-\log \frac{d\mu_{-t}}{d\mu} = \sum_i \beta_i \Delta Q_{t,i}$ and the definition of the total heat $\Delta Q_{t,\text{tot}} = \sum_i \Delta Q_{t,i}$.

If another choice of splitting of H is made, Equation (9) does not hold exactly and some boundary terms may have non-trivial asymptotic contributions; see [JPS16].

To the authors' knowledge, if V is unbounded, the existence of the moments or the extension to an open set of \mathbf{C} of the Fourier transform of \mathbb{P}_t can only be discussed on a model-by-model basis. In the next two subsections, we discuss two models with unbounded V that can be studied using only properties of Gaussian random variables.

2.1 Harmonic systems

Let Y be a real separable Hilbert space. Let B be a normal operator on Y with domain $\text{Dom } B$ and trivial kernel. Let X be the completion of $\text{Dom } B$ with respect to the norm $\|B \cdot\|_Y$. We also denote the extension of B to X by the letter B . We interpret the vector space $K = Y \oplus X$ as the phase space of a (finite, countable or uncountable) number of harmonic oscillators. We endow K with the Hilbert space structure given by the inner product

$$\langle \pi \oplus \phi, \pi' \oplus \phi' \rangle := \langle \pi, \pi' \rangle_Y + \langle B\phi, B\phi' \rangle_Y.$$

Let ϖ be the symplectic bilinear form on K defined by

$$\varpi(x, y) = \langle \mathcal{L}_0^{-1} x, y \rangle,$$

where

$$\mathcal{L}_0 := \begin{pmatrix} 0 & -B^* B \\ \mathbf{1} & 0 \end{pmatrix}.$$

The operator \mathcal{L}_0 is skew-adjoint with domain

$$\text{Dom}(\mathcal{L}_0) = \{(\pi, \phi) \in K : \pi \in \text{Dom } B, B\phi \in \text{Dom } B^*\}.$$

The space (K, ϖ) is our symplectic (Hilbert) manifold.

Let the unperturbed Hamiltonian be

$$\begin{aligned} H_0 : K &\rightarrow \mathbf{R}_+ \\ x &\mapsto \frac{1}{2} \|x\|^2. \end{aligned}$$

Let v be a trace class self-adjoint operator on K such that $\|\mathcal{L}_0 v\| < \infty$. The perturbation V is then defined as

$$\begin{aligned} V : K &\rightarrow \mathbf{R} \\ x &\mapsto \frac{1}{2} \langle x, vx \rangle, \end{aligned}$$

and $H(x) = \frac{1}{2} \langle x, (\mathbf{1} + v)x \rangle$.

Let $\mathcal{L} = \mathcal{L}_0(\mathbf{1} + v)$. Since \mathcal{L}_0 is skew-adjoint and $\|\mathcal{L}_0 v\| < \infty$, perturbation theory implies that $(e^{t\mathcal{L}})_{t \in \mathbf{R}}$ is a semigroup of bounded operators on K . It is easy to check that the flow $(\Xi^t)_{t \in \mathbf{R}}$

associated to H is then given by $\Xi^t(x) = e^{t\mathcal{L}}x$ for any $x \in K$. Moreover, using the inequality $\|A^*TA\|_{\text{tr}} \leq \|A\|^2\|T\|_{\text{tr}}$ for $A \in \mathcal{B}(K)$ and T trace class, $v_t := e^{t\mathcal{L}^*}ve^{t\mathcal{L}}$ remains trace class for any $t \in \mathbf{R}$. Here, $\|\cdot\|_{\text{tr}}$ denotes the trace norm.

We turn to the construction of Gaussian states. First, K needs to be enlarged. Since it is separable by assumption, there exists an orthonormal basis $(u_i)_{i \in \mathbf{N}}$ of K . Any such countable orthonormal basis (ONB) induces an isomorphism of K into $\ell^2(\mathbf{N}; \mathbf{R})$: $K \ni x \mapsto (\langle u_i, x \rangle)_{i \in \mathbf{N}}$. Let $(l_i)_{i \in \mathbf{N}}$ be a sequence of strictly positive numbers such that $\sum_{i \in \mathbf{N}} l_i = 1$. Let $\langle \cdot, \cdot \rangle_l$ be the inner product on K defined by

$$\langle x, y \rangle_l := \sum_{i \in \mathbf{N}} l_i \langle u_i, x \rangle \langle u_i, y \rangle$$

and let \bar{K} be the completion of K with respect to the norm induced by $\langle \cdot, \cdot \rangle_l$.

Consider a positive definite bounded linear operator D on K . As pointed out for example in [JPRB11], by Kolmogorov's extension Theorem, there is a unique measure μ_D on \bar{K} whose restriction to $\text{linspan}\{u_i\}_{i \in I}$, with $I \subset \mathbf{N}$ a finite set, is the centered Gaussian measure with inverse covariance matrix $[\langle u_i, Du_j \rangle]_{i,j \in I}$. The fact that μ_D is indeed concentrated on \bar{K} is guaranteed by the estimate $\sum_{n \in I} l_n \langle u_n, Du_n \rangle \leq \|D\|$. By uniqueness of the measure, the inclusion $\text{supp } \mu_D \subseteq \bar{K}$ does not depend on the choice of $(u_i)_{i \in \mathbf{N}}$ and $(l_i)_{i \in \mathbf{N}}$.

The measure μ_D then has characteristic function

$$\begin{aligned} \xi_D : \bar{K}^* &\rightarrow \mathbb{R}_+ \\ y &\mapsto e^{-\frac{1}{2}\langle y, Dy \rangle}, \end{aligned}$$

where $\bar{K}^* \subset K$ is the dual of \bar{K} .

Conversely, the measure μ_D can equivalently be constructed directly using the above characteristic function and the Bochner–Minlos Theorem, see for example [JP98, SL77].

Since the choice of ONB $(u_i)_{i \in \mathbf{N}}$ and sequence $(l_i)_{i \in \mathbf{N}}$ is arbitrary, we can choose, for any self-adjoint trace class operator a with trivial kernel, $(u_i)_{i \in \mathbf{N}}$ an ONB diagonalizing $|a|$, and $(l_i)_{i \in \mathbf{N}}$ its eigenvalues divided by $\text{tr } |a|$. We then have $\sup_{x \in \bar{K}} |\langle x, ax \rangle| / \|x\|_{\bar{K}} < \text{tr } |a|$. Hence $x \mapsto \langle x, ax \rangle$ extends by continuity to a finite function on $\text{supp } \mu_D \subset \bar{K}$. This procedure is easily adapted for a any trace class operator.

Since v is trace class, V can be extended by continuity to $\text{supp } \mu_D \ni x \mapsto \langle x, vx \rangle \in \mathbf{R}$. Similarly $\text{supp } \mu_D \ni x \mapsto \langle x, v_t x \rangle \in \mathbf{R}$ is extended by continuity from K to $\text{supp } \mu_D$. On the contrary, the extension of the unperturbed Hamiltonian can take infinite values on $\text{supp } \mu_D$.

Definition 2.5. The probability distribution \mathbb{P}_t describing the *heat variation* is the law of the random variable ΔQ defined in Equation (7) with respect to μ_D . Namely, \mathbb{P}_t is the law of

$$x \mapsto \langle x, vx \rangle - \langle e^{t\mathcal{L}}x, v e^{t\mathcal{L}}x \rangle,$$

with respect to μ_D .

Remark 2.6. Because v and $e^{t\mathcal{L}^*}ve^{t\mathcal{L}}$ are trace class, this random variable is μ_D -integrable.

In this quadratic model, the Fourier transform of \mathbb{P}_t extends analytically to an open neighborhood of \mathbf{R} in \mathbf{C} . Hence, all the moments of ΔQ exist. A finer study of the following proposition, including a proof of a large deviation principle with linear rate function can be found in [BJP17].

Proposition 2.7. *Given \mathbb{P}_t of Definition 2.5, for all $t \in \mathbf{R}$, there exists $\gamma_t > 0$ such that*

$$\mathbb{E}_t[e^{\gamma_t |\Delta Q|}] < \infty. \quad (11)$$

Moreover, if $-1 \notin \text{sp } v$, then there exists $\gamma > 0$ such that,

$$\sup_{t \in \mathbf{R}_+} \mathbb{E}_t[e^{\gamma |\Delta Q|}] < \infty. \quad (12)$$

Proof. Since $\mathbb{E}_t[e^{\gamma|\Delta Q|}] \leq \mathbb{E}_t[e^{\gamma\Delta Q}] + \mathbb{E}_t[e^{-\gamma\Delta Q}]$, the first bound (11) follows directly from properties of Gaussian measures and the fact that $v - e^{t\mathcal{L}^*} v e^{t\mathcal{L}}$ is trace class. The uniform bound (12) follows from the fact that $-1 \notin \text{sp } v$ implies $v - e^{t\mathcal{L}^*} v e^{t\mathcal{L}}$ is uniformly bounded with respect to t . Indeed, in that case, $\mathbf{1} + v$ is invertible with bounded inverse so that $e^{t\mathcal{L}^*}(\mathbf{1} + v)e^{t\mathcal{L}} = \mathbf{1} + v$ implies $\|e^{t\mathcal{L}}\|^2 \leq \|\mathbf{1} + v\| \|(\mathbf{1} + v)^{-1}\|$. \square

Remark 2.8. Note that the size of the interval of $\gamma \in \mathbf{R}_+$ for which (12) holds is typically increasing in $\|v\|$. This model then illustrates the expectation stated in the Introduction that the heat fluctuations are controlled by λ when V is unbounded. Further discussion of the relationship between λ and γ for such harmonic models can be found in [BJP17].

Corollary 2.9. *Given \mathbb{P}_t of Definition 2.5,*

$$\mathbb{E}_t[|\Delta Q|^k] < \infty$$

for all $k \in \mathbf{N}$ and any $t \in \mathbf{R}_+$. Moreover, if $-1 \notin \text{sp } v$, then

$$\sup_{t \in \mathbf{R}} \mathbb{E}_t[|\Delta Q|^k] < \infty.$$

for all $k \in \mathbf{N}$,

Proof. The corollary is an immediate consequence of previous proposition and the inequality $x^k \leq \gamma^{-k} k! e^{\gamma x}$ for x and γ positive. \square

We will see in Section 4.2 that, for the quantum harmonic oscillator in the context of quantum two-time measurement, further assumptions on V are needed to prove analogues of Proposition 2.7 and Corollary 2.9.

Example 2.10. We give an example of this abstract setting that can be compared with the quantum harmonic oscillator of Section 4.2. We consider a single harmonic oscillator interacting with a Gaussian 1-dimensional thermal bath at inverse temperature $\beta > 0$. The oscillator has Hilbert space $K_{\text{osc}} = \mathbf{R} \oplus \mathbf{R}$ with inner product

$$\langle (p, q), (p', q') \rangle_{\text{osc}} = pp' + qq'.$$

Let $\dot{H}^1(\mathbf{R})$ be the completion of $H^1(\mathbf{R})$ with respect to the norm $\|\nabla \cdot\|$. Then the bath Hilbert space is the vector space $K_{\text{bath}} := L^2(\mathbf{R}) \oplus \dot{H}^1(\mathbf{R})$ equipped with the inner product

$$\langle (\pi, \phi), (\pi', \phi') \rangle_{\text{bath}} = \langle \pi, \pi' \rangle_{L^2} + \langle \nabla \phi, \nabla \phi' \rangle_{L^2}.$$

The Hilbert space for the compound system is then $K = K_{\text{osc}} \oplus K_{\text{bath}}$ with unperturbed Hamiltonian

$$\begin{aligned} H_0 : K_{\text{osc}} \oplus K_{\text{bath}} &\rightarrow \mathbf{R} \\ (p, q, \pi, \phi) &\mapsto \frac{1}{2}(\|(p, q)\|_{\text{osc}}^2 + \|(\pi, \phi)\|_{\text{bath}}^2). \end{aligned}$$

The operator defining the symplectic form is

$$\mathcal{L}_0 = \begin{pmatrix} 0 & -1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & \Delta \\ 0 & 0 & 1 & 0 \end{pmatrix},$$

where Δ denotes the Laplacian on \mathbf{R} .

We take as an initial state the measure μ_D for the bounded operator $D = \beta^{-1}\mathbf{1}$ on K . With $\psi_o = (0, 1, 0, 0)$ and $\psi_f = (0, 0, 0, f)$ for some non-zero $f \in H^2(\mathbf{R})$, we consider

$$v = \psi_o \langle \psi_f, \cdot \rangle + \psi_f \langle \psi_o, \cdot \rangle.$$

The perturbed Hamiltonian is then

$$H : K_{\text{osc}} \oplus K_{\text{bath}} \rightarrow \mathbf{R}$$

$$(p, q, \pi, \phi) \mapsto \frac{1}{2}(\|(p, q)\|_{\text{osc}}^2 + \|(\pi, \phi)\|_{\text{bath}}^2) + q \int_{\mathbf{R}} \nabla f(x) \nabla \phi(x) dx$$

and we have

$$\mathcal{L} = \begin{pmatrix} 0 & -1 & 0 & -\langle f, \cdot \rangle_{H^1} \\ 1 & 0 & 0 & 0 \\ 0 & \Delta f & 0 & \Delta \\ 0 & 0 & 1 & 0 \end{pmatrix}.$$

The assumption $-1 \notin \text{sp } v$ is satisfied if and only if $\|\nabla f\|_{L^2} \neq 1$.

2.2 Linear perturbations

Consider the setting of the previous subsection but let $V : K \mapsto \mathbf{R}$ instead be a real linear form defined by

$$V(x) := \langle f, x \rangle$$

with $f \in K$. The perturbed Hamiltonian is then

$$H : K \rightarrow \mathbf{R}$$

$$x \mapsto \frac{1}{2}\|x\|^2 + \langle f, x \rangle.$$

Remark 2.11. With an appropriate choice of gauge, these definitions correspond to the Hamiltonian description of the electromagnetic field x in presence of a charge current encoded in f . The hypothesis $f \in K$ corresponds to a mild UV regularity condition that prevents the apparition of infinite electrostatic energy. It is necessary to properly define the model. Physically, this condition is justified by taking into account that charges are not point like but have some ‘‘volume’’ [Spo04, §2.3].

The technical condition $f \in K$ here is analogous to the assumption $f \in \text{Dom } \hat{e}$ in the quantum counter part to this model discussed in Section 4.3, and is significantly weaker than the essentially necessary UV conditions that are used to control the tails of \mathbb{P}_t there.

It is again easy to show that the flow defined by the perturbed Hamiltonian H is given by

$$\Xi^t(x) = e^{t\mathcal{L}_0}x + (e^{t\mathcal{L}_0} - 1)f$$

for any $t \in \mathbf{R}$ and $x \in K$.

Definition 2.12. The probability distribution \mathbb{P}_t describing the *heat variation* is the law of the random variable ΔQ defined in Equation (7) with respect to μ_D . Namely, \mathbb{P}_t is the law of the random variable

$$x \mapsto \langle f, (1 - e^{t\mathcal{L}_0})(f + x) \rangle$$

with respect to the measure μ_D .

The following propositions are straightforward consequences of the properties of Gaussian random variables and show that the fluctuations of heat are in some sense trivial in this model.

Proposition 2.13. For any $t \in \mathbf{R}$, \mathbb{P}_t is a Gaussian probability measure with mean

$$\mathbb{E}_t \Delta Q = \langle f, (1 - e^{t\mathcal{L}_0})f \rangle$$

and variance

$$\mathbb{E}_t \Delta Q^2 - (\mathbb{E}_t \Delta Q)^2 = \|D^{\frac{1}{2}}(1 - e^{-t\mathcal{L}_0})f\|^2.$$

Proof. By definition μ_D is such that any projection of x on a finite dimensional subspace of K is a centered Gaussian random vector with same dimension as the subspace and covariance given by the projection of D on this subspace. Hence, considering the projection of x on the one dimensional linear subspace spanned by $(1 - e^{t\mathcal{L}_0})f$, for any $\alpha \in \mathbf{R}$,

$$\mathbb{E}_t(e^{i\alpha\Delta Q}) = e^{i\alpha\langle f, (1 - e^{t\mathcal{L}_0})f \rangle - \frac{1}{2}\alpha^2\|D^{\frac{1}{2}}(1 - e^{-t\mathcal{L}_0})f\|^2}. \quad \square$$

Remark 2.14. If $D\mathcal{L}_0 = \mathcal{L}_0D$, the variance is $2\langle D^{\frac{1}{2}}f, (1 - \cosh(t\mathcal{L}_0))D^{\frac{1}{2}}f \rangle$.

Corollary 2.15. For any $\gamma > 0$,

$$\sup_{t \in \mathbf{R}} \mathbb{E}_t(e^{\gamma|\Delta Q|}) < \infty.$$

Proof. The probability measure \mathbb{P}_t being Gaussian, the proposition follows from the fact that the Fourier transform of \mathbb{P}_t is entire analytic, and from the inequalities $|\langle f, (1 - e^{t\mathcal{L}_0})f \rangle| \leq 2\|f\|^2$, $\|D^{\frac{1}{2}}(1 - e^{-t\mathcal{L}_0})f\| \leq 2\|D^{\frac{1}{2}}\|\|f\|$ and $e^{|x|} \leq e^x + e^{-x}$. \square

Remark 2.16. Even if the Fourier transform of \mathbb{P}_t is analytic on \mathbf{C} , the variance is quadratic in $\|f\|$. Hence this model illustrates again the control of the heat fluctuations by λ as written in the Introduction.

In Section 4.3, we will see that both Proposition 2.13 and Corollary 2.15 do not hold for the similar quantized model in the two-time measurement framework. In the quantum case, \mathbb{P}_t is the probability measure of an inhomogeneous Poisson process, not of a Gaussian random variable. The existence of a uniform bound similar to the one obtained in Corollary 2.15, depends then on the properties of the Poisson process' intensity.

3 Control of the tails for bounded perturbations

3.1 Setup

Let us first briefly introduce the operator algebraic definition of quantum dynamical systems that we adopt. We refer the reader to [BR87, §2–3] for a thorough exposition of this mathematical formalism; see also [Pil06] for a shorter introduction.

A C^* -dynamical system is a triplet $(\mathcal{O}, \tau, \omega)$, where \mathcal{O} is a unital C^* -algebra, $(\tau^s)_{s \in \mathbf{R}}$ is a strongly continuous one-parameter group of $*$ -automorphisms of \mathcal{O} , and ω is a state, i.e. a positive linear functional on \mathcal{O} satisfying $\omega(\mathbf{1}) = 1$. We furthermore assume ω to be faithful and τ -invariant.⁵ Namely, we require that $\omega(A^*A) = 0$ implies $A = 0$, and that

$$\omega = \omega \circ \tau^s$$

for all $s \in \mathbf{R}$. We denote the generator of $(\tau^s)_{s \in \mathbf{R}}$ by δ . From the representation theory of C^* -algebras, there exists a Hilbert space \mathcal{H} , a $*$ -isomorphism $\pi : \mathcal{O} \rightarrow \mathcal{B}(\mathcal{H})$, and a unit vector $\Omega \in \mathcal{H}$ such that

$$\omega(A) = \langle \Omega, \pi(A)\Omega \rangle \quad (13)$$

⁵We restrict ourselves to a faithful ω for simplicity. Up to some technical details, this assumption can be lifted.

for all $A \in \mathcal{O}$ and such that the vector Ω is cyclic for $\pi(\mathcal{O})$. This triple $(\mathcal{H}, \pi, \Omega)$ is called a *Gelfand–Naimark–Segal (GNS) representation* of \mathcal{O} associated to the state ω and is unique up to unitary equivalence; see for example [BR87, §2.3.3]. On this Hilbert space \mathcal{H} , there exists a unique self-adjoint operator L satisfying

$$L\Omega = 0 \tag{14}$$

and

$$\pi(\tau^s(A)) = e^{isL}\pi(A)e^{-isL} \tag{15}$$

for all $A \in \mathcal{O}$ and $s \in \mathbf{R}$. The self-adjoint operator L is referred to as the ω -Liouvillean of $(\tau^s)_{s \in \mathbf{R}}$, or simply the *Liouvillean*, and is typically not bounded, nor semi-bounded, for thermodynamic systems.

Finally, to a self-adjoint element V of \mathcal{O} we associate a *perturbed dynamics* $(\tau_V^s)_{s \in \mathbf{R}}$ generated by $\delta_V := \delta + i[V, \cdot]$. Both δ and δ_V are derivations on \mathcal{O} . We refer to V as a (bounded) *perturbation* and associate to it a measure that is the central object of this paper. Note that, in the GNS representation, the perturbed dynamics is implemented by $L + \pi(V)$ in the sense that

$$\pi(\tau_V^s(A)) = e^{is(L+\pi(V))}\pi(A)e^{-is(L+\pi(V))}$$

for all $A \in \mathcal{O}$ and all $s \in \mathbf{R}$.

Definition 3.1. The probability distribution \mathbb{P}_t describing the *heat variation* between times 0 and t associated to $(\mathcal{O}, \tau, \omega)$ and the self-adjoint perturbation $V \in \mathcal{O}$ is the spectral measure of the operator

$$L + \pi(V) - \pi(\tau_V^t(V)) \tag{16}$$

with respect to the vector Ω . Equivalently, it is the unique probability measure with characteristic function

$$\mathcal{E}_t(\alpha) = \langle \Omega, e^{i\alpha(L+\pi(V)-\pi(\tau_V^t(V)))}\Omega \rangle. \tag{17}$$

For a non-negative Borel function g , we will use $\mathbb{E}_t(g(\Delta Q))$ to denote the integral $\int_{\mathbf{R}} g(\Delta Q) d\mathbb{P}_t(\Delta Q)$ with value in $[0, \infty]$.

Remark 3.2. If \mathcal{O} is finite dimensional, \mathbb{P}_t is the probability distribution of the heat variation as defined by the two-time measurement protocol outlined in the introduction.

To see this, let $\mathcal{O} = M_{n \times n}(\mathbf{C})$, with an unperturbed dynamics defined in the Heisenberg picture by $\tau^t(A) = e^{itH_0} A e^{-itH_0}$ for all $A \in \mathcal{O}$ and where the Hamiltonian H_0 is a self-adjoint element of $M_{n \times n}(\mathbf{C})$. Let ω be $\omega(A) := \text{tr}(A\hat{\omega})$ with $\hat{\omega}$ a full-rank density matrix that commutes with H_0 . The state ω is then faithful and τ -invariant.

As presented in [JOPP11, §4.3.11]⁶, an associated GNS representation for this system is known as the *standard* GNS representation: it has Hilbert space $\mathcal{H} = M_{n \times n}(\mathbf{C})$ equipped with the inner-product $\langle A, B \rangle = \text{tr}(A^*B)$ and the representation π is given by left matrix multiplication: $(\pi(A))(B) = AB$. The vector representative Ω of ω is the $n \times n$ matrix $\hat{\omega}^{1/2}$. Then, the Liouvillean is $L = [H_0, \cdot]$ (i.e. the difference between left and right matrix multiplication) and Definition 3.1, for a self-adjoint perturbation $V \in M_{n \times n}(\mathbf{C})$, reads

$$\begin{aligned} \mathcal{E}_t(\alpha) &= \langle \Omega, e^{i\alpha(\pi(H_0)+\pi(V)-\pi(\tau_V^t(V)))} e^{-i\alpha\pi(H_0)} \Omega \rangle \\ &= \text{tr}(\hat{\omega}^{1/2} e^{it(H_0+V)} e^{i\alpha(H_0+V-V)} e^{-it(H_0+V)} e^{-i\alpha H_0} \hat{\omega}^{1/2}) \\ &= \text{tr}(e^{i\alpha e^{it(H_0+V)} H_0 e^{-it(H_0+V)}} e^{-i\alpha H_0} \hat{\omega}). \end{aligned}$$

In the first equality we used $[H_0, \hat{\omega}] = 0$ to change $\hat{\omega}^{1/2} e^{-i\alpha H_0}$ to $e^{-i\alpha H_0} \hat{\omega}^{1/2}$. The last expression of the characteristic function is the quantum equivalent of (8) for classical systems.

⁶Section 4.3.11. The *standard representations* of \mathcal{O} of the published version corresponds to Section 2.11 of the version available on the *arXiv*.

Using the spectral decomposition $H_0 =: \sum_j \epsilon_j P_j$ where $\{P_j\}_j$ is a resolution of the identity, we obtain that

$$\begin{aligned} \mathcal{E}_t(\alpha) &= \sum_{i,j} \text{tr}(e^{i\alpha\epsilon_j} P_j e^{-it(H_0+V)} e^{-i\alpha\epsilon_i} P_i \hat{\omega} e^{it(H_0+V)}) \\ &= \sum_{\Delta Q \in \text{sp } H_0 - \text{sp } H_0} e^{i\alpha\Delta Q} \sum_{\substack{i,j \\ \epsilon_j - \epsilon_i = \Delta Q}} \text{tr}(P_j e^{-it(H_0+V)} P_i \hat{\omega} e^{it(H_0+V)}) \end{aligned}$$

is the characteristic function of the probability measure

$$\mathbb{P}_t(\{\Delta Q\}) = \sum_{\substack{i,j \\ \epsilon_j - \epsilon_i = \Delta Q}} \text{tr}(P_j e^{-it(H_0+V)} P_i \hat{\omega} P_i e^{it(H_0+V)} P_j).$$

Hence, \mathbb{P}_t is indeed the probability measure of the heat variation defined as the result of a two-time measurement protocol of the Hamiltonian H_0 [Kur00, Tas00, JOPP11, EHM09, CHT11].

Relevant models of infinitely extended quantum systems are obtained as the thermodynamic limit of a sequence of such confined systems. As conveyed e.g. in [JOPP11], [JPPP15] and [BJP⁺15], mild assumptions ensure that the corresponding sequence of probability measures converges weakly to the measure \mathbb{P}_t of Definition 3.1 on the limiting infinitely extended system. In Appendix B, we provide proofs of this weak convergence for the models studied in Section 4.

Remark 3.3. As we pointed out in Remark 2.4 for classical systems, the choice of operator whose spectral measure defines \mathbb{P}_t is partly arbitrary. We could have chosen the spectral measure of $L_W + \pi(V - W) - \pi(\tau_V^t(V - W))$ with $W \in \mathcal{O}$ self-adjoint and L_W the Liouvillean of τ_W as long as ω is τ_W invariant. As in the classical case, the choice of $L + \pi(V) - \pi(\tau_V^t(V))$ can be motivated by the entropy balance equation when ω is a (τ, β) -KMS state.

Indeed, in the framework of Tomita–Takesaki theory, if ω is a (τ, β) -KMS state, the Liouvillean is $L = -\beta^{-1} \log \Delta_\omega$, where Δ_ω denotes the modular operator for the state ω . With $\Delta_{\omega_{-t}|\omega}$ denoting the relative modular operator (non-commutative analogue of the Radon–Nikodym derivative) between the states $\omega_{-t} := \omega \circ \tau_V^{-t}$ and ω , one then has the identity [JPPP15]

$$\beta(L + \pi(V) - \pi(\tau_V^t(V))) = -\log \Delta_{\omega_{-t}|\omega}. \quad (18)$$

Hence, defining the distribution \mathbb{P}_t of heat variation as the spectral measure for the operator $L + \pi(V) - \pi(\tau_V^t(V))$ with respect to Ω amounts to lifting the classical entropy balance equation (9) to the level of the operators defining the distributions of heat on one side and entropy production on the other. Indeed, upon taking expectation with respect Ω , the above identity reduces to

$$\beta \mathbb{E}_t(\Delta Q) = -\langle \Omega, \log \Delta_{\omega_{-t}|\omega} \Omega \rangle, \quad (19)$$

where the right-hand side is identified as the relative entropy between ω and ω_{-t} , exactly as in (10).

Remark 3.4. Definition 3.1 extends naturally to the case of W^* -dynamical system $(\mathfrak{M}, \tau, \omega)$, where \mathfrak{M} is a concrete W^* -algebra, $(\tau^s)_{s \in \mathbf{R}}$ is a σ -weakly continuous one-parameter group of $*$ -automorphisms of \mathfrak{M} and ω is a τ -invariant state on \mathfrak{M} , together with a bounded self-adjoint perturbation $V \in \mathfrak{M}$. Theorems 3.5 and 3.6 then also generalise to this case.

When considering V merely affiliated to the W^* -algebra \mathfrak{M} , one may in some cases mimic the construction Definition 3.1 and obtain results analogous to the ones for C^* -algebras. For example, if the dynamics τ is implemented by the self-adjoint operator L , the condition $L + V$ being essentially self-adjoint on $\text{Dom } L \cap \text{Dom } V$ ensures that the construction of the perturbed group τ_V is indeed a W^* -dynamics (see [DJP03]). We do not discuss the general theory of heat variation for such unbounded perturbations, but treat some examples in Sections 4.2 and 4.3.

3.2 Results

Our two main general results for C^* -dynamical systems with bounded perturbations describe the control of the tails of \mathbb{P}_t . The first one gives a sufficient condition for the existence and uniform boundedness in time of even moments; the second, for the existence and uniform boundedness of an analytic extension of the Fourier transform to a neighborhood of 0 in \mathbf{C} .

Theorem 3.5. *Let $(\mathcal{O}, \tau, \omega)$ be a C^* -dynamical system, $V \in \mathcal{O}$ a self-adjoint perturbation, and \mathbb{P}_t the probability measure of Definition 3.1. If $V \in \text{Dom } \delta^n$, then*

$$\sup_{t \in \mathbf{R}} \mathbb{E}_t[\Delta Q^{2n+2}] < \infty.$$

Theorem 3.6. *Let $(\mathcal{O}, \tau, \omega)$ be a C^* -dynamical system, $V \in \mathcal{O}$ a self-adjoint perturbation and \mathbb{P}_t the probability measure of Definition 3.1. For $\gamma > 0$, if $V \in \text{Dom } e^{i\frac{\gamma}{2}\delta} \cap \text{Dom } e^{-i\frac{\gamma}{2}\delta}$, then*

$$\sup_{t \in \mathbf{R}} \mathbb{E}_t[e^{\gamma|\Delta Q|}] < \infty.$$

Remark 3.7. A similar result was already presented in [BJP⁺15]. It was proved via the thermodynamic limit. Below, we provide a proof in the present framework.

Remark 3.8. The assumptions $V \in \text{Dom } \delta^n(V)$ and $V \in \text{Dom } e^{i\frac{\gamma}{2}\delta} \cap \text{Dom } e^{-i\frac{\gamma}{2}\delta}$ can be reformulated in terms of regularity of the map $t \mapsto \tau^t(V)$. We have $V \in \text{Dom } \delta^n(V)$ if and only if $t \mapsto \tau^t(V)$ is n times norm differentiable and $V \in \text{Dom } e^{i\frac{\gamma}{2}\delta} \cap \text{Dom } e^{-i\frac{\gamma}{2}\delta}$ if and only if $t \mapsto \tau^t(V)$ admits a bounded analytic extension to the strip $\{z \in \mathbf{C} : |\text{Im } z| < \frac{\gamma}{2}\}$.

Remark 3.9. In Section 4.1 we discuss a model of quasi-free fermions where the regularity conditions of Theorems 3.5 and 3.6 correspond explicitly to UV regularity conditions. In this model, we prove that the assumptions of Theorem 3.5 are essentially necessary for the existence of moments at any finite time. This is in sharp contrast with the results for classical systems. Indeed, as exposed in Section 2, the heat variation is surely uniformly bounded as soon as the perturbation is bounded (see Proposition 2.3). Hence no UV regularity assumption on the perturbation is needed for the existence of moments or the analytic extension to a complex neighborhood of 0 of the Fourier transform of \mathbb{P}_t .

3.3 Proofs

We start with some useful properties of derivations.

Lemma 3.10. *Let δ be a derivation on an algebra \mathcal{O} and $A, B \in \text{Dom}(\delta^n)$, $n \in \mathbf{N}$. Then*

$$\delta^n(AB) = \sum_{k=0}^n \binom{n}{k} \delta^{n-k}(A) \delta^k(B).$$

Proof. It follows easily by induction from the property of derivations $\delta(AB) = \delta(A)B + A\delta(B)$. \square

Lemma 3.11. *Let δ be a derivation on a C^* -algebra \mathcal{O} and let $V_t \in \text{Dom}(\delta^n)$ and $\sup_t \|\delta^n(V_t)\| < \infty$. Then $V_t^p \in \text{Dom}(\delta^n)$ for all $p \in \mathbf{N}$ and*

$$\sup_t \|\delta^n(V_t^p)\| < \infty.$$

Proof. It follows easily from Lemma 3.10 by induction on p . \square

Recall that, given $V \in \mathcal{O}$, the one-parameter group $(\tau_V^t)_{t \in \mathbf{R}}$ is the perturbed dynamics whose generator is given by $\delta_V = \delta + i[\cdot, V]$.

Lemma 3.12. *If $V \in \text{Dom } \delta^n$, then $V - \tau_V^t(V) \in \text{Dom } \delta^n$ and*

$$\sup_{t \in \mathbf{R}} \|\delta^n(V - \tau_V^t(V))\| < \infty.$$

Proof. Using a Dyson expansion of $V - \tau_V^t(V)$, it is easy to show that $V - \tau_V^t(V) \in \text{Dom}(\delta^n)$. It remains to show that the norm of $\delta^n(V - \tau_V^t(V))$ is uniformly bounded in t .

Since $\|\delta^n(V)\| < \infty$ is independent of t , it suffices to show uniform boundedness of $\|\delta^n(\tau_V^t(V))\|$. We proceed by induction on n , noting that because δ is a strongly continuous group generator, $V \in \text{Dom } \delta^n$ implies $V \in \text{Dom } \delta^k$ for $k = 1, \dots, n-1, n$. For convenience let $c_n := \sup_{t \in \mathbf{R}} \|\delta^n(\tau_V^t(V))\|$.

For $n = 1$, by definition of τ_V^t from $\delta_V = \delta + i[V, \cdot]$,

$$\begin{aligned} \delta^1(\tau_V^t(V)) &= \delta_V^1(\tau_V^t(V)) - i[V, \tau_V^t(V)] \\ &= \tau_V^t(\delta_V(V)) - i[V, \tau_V^t(V)] \\ &= \tau_V^t(\delta(V)) - i[V, \tau_V^t(V)] \end{aligned}$$

and thus

$$\|\delta^1(\tau_V^t(V))\| \leq \|\tau_V^t(\delta(V))\| + \|[V, \tau_V^t(V)]\| \leq \|\delta(V)\| + 2\|V\|^2.$$

Hence $c_1 \leq \|\delta(V)\| + 2\|V\|^2$.

Now suppose that the statement holds for k and that $V \in \text{Dom } \delta^{k+1}$. Then,

$$\begin{aligned} \delta^{k+1}(\tau_V^t(V)) &= \delta_V(\delta^k(\tau_V^t(V))) - i[V, \delta^k(\tau_V^t(V))] \\ &= \delta_V^{k+1}(\tau_V^t(V)) - i \sum_{l=0}^k \delta_V^l[V, \delta^{k-l}(\tau_V^t(V))] \\ &= \tau_V^t(\delta_V^{k+1}(V)) - i \sum_{l=0}^k \delta_V^l[V, \delta^{k-l}(\tau_V^t(V))] \end{aligned}$$

and thus, using Lemma 3.10,

$$\begin{aligned} \|\delta^{k+1}(\tau_V^t(V))\| &\leq \|\tau_V^t(\delta_V^{k+1}(V))\| + \sum_{l=0}^k \|\delta_V^l[V, \delta^{k-l}(\tau_V^t(V))]\| \\ &\leq \|\delta_V^{k+1}(V)\| + \sum_{l=0}^k \sum_{j=0}^l \binom{l}{j} 2\|\delta_V^j(V)\| \|\delta_V^{l-j} \delta^{k-l}(\tau_V^t(V))\|. \end{aligned}$$

By repeated application of the product rule of Lemma 3.10, this may in turn be bounded uniformly in t by a combination of powers of c_1, \dots, c_k , $\|V\|$, $\|\delta(V)\|$, \dots , $\|\delta^k(V)\|$ and $\|\delta^{k+1}(V)\|$, which are all finite by induction hypothesis. \square

In the following, L and W are operators on a Hilbert space, with L possibly unbounded. We denote by $\text{ad}_L(W)$ the commutator $[L, W]$. We set $\text{ad}_L^0 W = W$. The following formula can be easily proven by induction.

Lemma 3.13. *Let L and W be two linear operators on a Hilbert space. Assume that $\text{ad}_L^k(W)$ extends to a bounded operator for all k with $0 \leq k \leq n$. Then,*

$$L^n W = \sum_{k=0}^n \binom{n}{k} \text{ad}_L^k(W) L^{n-k}.$$

With these lemmas, we are now ready to prove Theorems 3.5 and 3.6.

Proof of Theorem 3.5. Set $X_t := \pi(V - \tau_V^t(V))$. By definition of \mathbb{P}_t as a spectral measure, and since $L\Omega = 0$, we have

$$\int_{\mathbf{R}} |\Delta Q|^{2n+2} d\mathbb{P}_t(\Delta Q) = \|(L + X_t)^{n+1}\Omega\|^2 = \|(L + X_t)^n X_t \Omega\|^2$$

Now,

$$(L + X_t)^n X_t = \sum_{A_i=L, X_t} A_1 A_2 \cdots A_n X_t.$$

Each term in the sum can be written as

$$L^{\alpha_1} X_t^{\beta_1} \cdots L^{\alpha_m} X_t^{\beta_m} X_t$$

with $\sum_i \alpha_i + \sum_i \beta_i = n$. Using repeatedly Lemma 3.13 and $L\Omega = 0$,

$$(L + X_t)^n X_t \Omega = P_t X_t \Omega$$

with P_t a non-commutative polynomial of variables $\text{ad}_L^k X_t^p$ with $k + p \leq n$.

By Lemmas 3.11 and 3.12, $\|\delta^k((V - \tau_V^t(V))^p)\|$ is uniformly bounded for all p and all $k \leq n$. Since $\text{ad}_L^k(X_t^p) = \pi(\delta^k((V - \tau_V^t(V))^p))$, it implies $\text{ad}_L^k(X_t^p)$ is uniformly bounded. Hence $\sup_t \|P_t\| < \infty$, and $(L + X_t)^{n+1}\Omega = P_t \Omega$ proves the theorem. \square

Proof of Theorem 3.6. By hypothesis, for each $N \in \mathbf{N}$, the function of $s \in \mathbf{R}$ defined by the truncated Araki–Dyson series

$$E_V^{(N)}(s) := \mathbf{1} + \sum_{n=1}^N i^n \int_0^s \cdots \int_0^{s_{n-1}} \pi(\tau^{s_n}(V)) \cdots \pi(\tau^{s_1}(V)) ds_n \cdots ds_1 \quad (20)$$

has an analytic extension to the complex open neighborhood $S_0 := \{z \in \mathbf{C} : |\text{Im } z| < \frac{1}{2}\gamma\}$. Moreover,

$$\sup_{N \in \mathbf{N}} \|E_V^{(N)}(z)\| \leq e^{|z|v_0}$$

with $v_0 := \sup_{z \in S_0} \|\tau^z(V)\|$. Hence, by the Vitali–Porter convergence theorem and equivalence of the weak and strong analyticity, (20) converges on S_0 to a norm-analytic $\pi(\mathcal{O})$ -valued function $S_0 \ni z \mapsto E_V(z)$ satisfying

$$\|E_V(z)\| \leq e^{|z|v_0}.$$

On the other hand, the truncated Araki–Dyson series (20) converges to $e^{is(L+\pi(V))}e^{-isL}$ for all $s \in \mathbf{R}$. Then we have

$$\begin{aligned} e^{-is(L+\pi(V)-\pi(\tau_V^t(V)))}\Omega &= e^{-is(L+\pi(V)-e^{it(L+\pi(V))}\pi(V)e^{-it(L+\pi(V))})}\Omega \\ &= e^{it(L+\pi(V))}e^{-isL}e^{is(L+\pi(V))}e^{-it(L+\pi(V))}e^{-is(L+\pi(V))}\Omega \\ &= e^{it(L+\pi(V))}E_V(-s)^*e^{-it(L+\pi(V))}E_V(-s)\Omega, \end{aligned}$$

with the operators on the right-hand side admitting analytic extensions as functions of s to the set S_0 . Therefore,

$$\begin{aligned} \mathbb{E}_t[e^{\gamma|\Delta Q|}] &\leq \mathbb{E}_t[e^{-\gamma\Delta Q}] + \mathbb{E}_t[e^{\gamma\Delta Q}] \\ &= \|E_V(-\frac{i}{2}\gamma)^*e^{-it(L+\pi(V))}E_V(-\frac{i}{2}\gamma)\Omega\|^2 \\ &\quad + \|E_V(\frac{i}{2}\gamma)^*e^{-it(L+\pi(V))}E_V(\frac{i}{2}\gamma)\Omega\|^2 \\ &\leq \|\tau_V^t(E_V(-\frac{i}{2}\gamma)^*)\|^2\|E_V(-\frac{i}{2}\gamma)\|^2\|\Omega\|^2 \\ &\quad + \|\tau_V^t(E_V(\frac{i}{2}\gamma)^*)\|^2\|E_V(\frac{i}{2}\gamma)\|^2\|\Omega\|^2 \\ &\leq 2e^{2\gamma v_0}. \end{aligned} \quad \square$$

4 Heavy tails in some quasi-free gas models

In the present section, we study some models with V bounded and unbounded where the assumptions of Theorems 3.5 and 3.6 are related to explicit UV regularity conditions. We moreover prove some converse implications in each model. The results presented in this section are proved in Section 5.

In Section 4.1, in a model of quasi-free fermions, we show that the $2n + 2^{\text{nd}}$ moment of \mathbb{P}_t exists essentially if and only if $V \in \text{Dom } \delta^n$.

In Section 4.2, we study a model that can be seen as the bosonic analogue of the previous one or as a quantization of the classical model of Section 2.1. We show that, even if V is unbounded in this model, results similar to Theorems 3.5 and 3.6 hold, with a converse to Theorem 3.5.

In Section 4.3, we study a model that can be seen as a quantization of the classical model of Section 2.2. First we show that \mathbb{P}_t is the law of an inhomogeneous Poisson process, in sharp contrast with its classical counterpart for which \mathbb{P}_t is the law of a Gaussian random variable. Building on that result, we again show that, even if V is unbounded in this model, analogues of Theorems 3.5 and 3.6 hold. Moreover, we prove that the implications in both of these theorems have converses.

Before we give the details on these three models, we introduce common notations. Given a Hilbert space \mathfrak{h} , we denote by $\Gamma^-(\mathfrak{h})$ [resp. $\Gamma^+(\mathfrak{h})$] the anti-symmetric [resp. symmetric] Fock space associated to \mathfrak{h} ; by a and a^* , the usual annihilation and creation operators there; and by $d\Gamma(b)$ the second quantization of the one-particle operator b on \mathfrak{h} . Since it is clear from the context, we use the same letter for the fermionic and the bosonic ones. The expression a^\sharp stands either for the creation a^* or for the annihilation a . We denote by φ the corresponding field operators $\varphi(\psi) := \frac{1}{\sqrt{2}}(a(\psi) + a^*(\psi))$.

4.1 Impurity in a quasi-free Fermi gas

We consider a fermionic impurity interacting with a quasi-free Fermi gas. The corresponding unital C^* -algebra \mathcal{O} is generated by $\{a(\phi) : \phi \in \mathfrak{h}\}$ with the one-particle Hilbert space $\mathfrak{h} = \mathbf{C} \oplus L^2(\mathbf{R}_+, de)$. The unperturbed dynamic is given by the extension of

$$\tau^s(a^\sharp(\phi)) = a^\sharp(e^{ish_0}\phi) \quad (21)$$

for $\phi \in \mathfrak{h}$, where $h_0 := \epsilon_o \oplus \hat{e}$ on \mathfrak{h} , to a group of $*$ -automorphisms of \mathcal{O} . Here and in what follows, $(\hat{e}\phi)(e) = e\phi(e)$ and $\epsilon_o > 0$.

The initial state ω is taken to be the quasi-free state on \mathcal{O} generated by the Fermi–Dirac density $T = (1 + e^{\beta h_0})^{-1}$ for some inverse temperature $\beta > 0$. Then, ω is a (τ, β) -KMS state. We refer the reader to [BR97, §5] or [RS75, §X.7] for more details.

The impurity–gas interaction is given by the bounded perturbation

$$V = a^*(\psi_f)a(\psi_o) + a^*(\psi_o)a(\psi_f),$$

where $\psi_f = 0 \oplus f \in \mathbf{C} \oplus L^2(\mathbf{R}_+, de)$ and $\psi_o = 1 \oplus 0 \in \mathbf{C} \oplus L^2(\mathbf{R}_+, de)$. Note that $V = d\Gamma(v)$ for the one-particle rank two operator

$$v = \psi_o \langle \psi_f, \cdot \rangle + \psi_f \langle \psi_o, \cdot \rangle.$$

Remark 4.1. This algebra and dynamics can be represented on the Hilbert space

$$\Gamma^-(\mathbf{C}) \otimes \Gamma^-(L^2(\mathbf{R}_+, de)) \cong \mathbf{C}^2 \otimes \Gamma^-(L^2(\mathbf{R}_+, de))$$

with unperturbed Hamiltonian

$$d\Gamma(\epsilon_o) \otimes \mathbf{1} + \mathbf{1} \otimes d\Gamma(\hat{e}),$$

and using the identification $\Gamma^-(\mathbf{C}) \otimes \Gamma^-(L^2(\mathbf{R}_+, de)) \cong \Gamma^-(\mathfrak{h})$ through the fermionic exponential map.

Through this map, the operator $d\Gamma(\epsilon_0) \otimes \mathbf{1} + \mathbf{1} \otimes d\Gamma(\hat{e})$ on $\Gamma^-(\mathbf{C}) \otimes \Gamma^-(L^2(\mathbf{R}_+, de))$ is indeed unitarily equivalent to $d\Gamma(h_0)$ on $d\Gamma(\mathfrak{h})$ and the operator

$$a^*(1) \otimes a(f) + a(1) \otimes a^*(f)$$

on $\Gamma^-(\mathbf{C}) \otimes \Gamma^-(L^2(\mathbf{R}_+, de))$ is unitarily equivalent to the operator $V = d\Gamma(v)$ on $\Gamma^-(\mathfrak{h})$.

Remark 4.2. The space $L^2(\mathbf{R}_+, de)$ with the unperturbed one-particle Hamiltonian \hat{e} is chosen for simplicity. This choice captures the essential features of our problem. Our proofs can be adapted to more general separable Hilbert spaces as long as the unperturbed one-particle Hamiltonian is lower bounded and one works in the representation where it is a multiplication operator.

Remark 4.3. If there exists $\gamma > 0$ such that $f \in \text{Dom } e^{\frac{1}{2}\gamma\hat{e}}$, the norm equality $\|a(\phi)\| = \|\phi\|$ implies that Theorem 3.6 holds for said γ . If one defines V using a sharp UV cutoff, namely if there exists $\Lambda > 0$ such that $f(e) = 0$ for all $e > \Lambda$, then $f \in \text{Dom } e^{\frac{1}{2}\gamma}$ for any $\gamma \in \mathbf{R}_+$ and Theorem 3.6 holds for arbitrarily large γ . Next theorem shows that UV regularization conditions are not only sufficient but necessary for the existence of moments of \mathbb{P}_t .

Theorem 4.4. *Let \mathbb{P}_t be the probability measure of Definition 3.1 for $(\mathcal{O}, \tau, \omega)$ and V defined in this section. Then, for any $n \in \mathbf{N}$, the following are equivalent*

- (i) $\sup_{t \in \mathbf{R}} \mathbb{E}_t[\Delta Q^{2n+2}] < \infty$;
- (ii) there exists $t_1 < t_2$ such that $\int_{t_1}^{t_2} \mathbb{E}_t[\Delta Q^{2n+2}] dt < \infty$;
- (iii) $f \in \text{Dom } \hat{e}^n$.

Remark 4.5. In this setup, since $\delta(a^\sharp(\phi)) = a^\sharp(ih_0\phi)$, and $\|a(\phi)\| = \|\phi\|$, $f \in \text{Dom } \hat{e}^n$ implies $V \in \text{Dom } \delta^n$. Hence the condition $V \in \text{Dom } \delta^n$ is equivalent to an UV regularity condition on the form factor f .

Remark 4.6. If there exists $n \in \mathbf{N}$ such that $f \notin \text{Dom } \hat{e}^n$, then the $2n+2^{\text{nd}}$ moment of \mathbb{P}_t essentially never exists. Hence, the Fourier transform of \mathbb{P}_t does not extend analytically to a neighborhood of the real axis and \mathbb{P}_t is, by definition, heavy-tailed.

4.2 Quantum Open Harmonic Oscillator

We consider a quantum harmonic oscillator coupled to a gas of quasi-free scalar bosons. Our model is similar to the one studied in [Ara81, Dav73]. It is the bosonic analogue of the model of the previous section and the quantum analogue of the classical model of Section 2.1.

Consider the Hilbert space

$$\Gamma^+(\mathbf{C}) \otimes \Gamma^+(L^2(\mathbf{R}_+, de)) \cong L^2(\mathbf{R}) \otimes \Gamma^+(L^2(\mathbf{R}_+, de))$$

with unperturbed Hamiltonian

$$d\Gamma(\epsilon_0) \otimes \mathbf{1} + \mathbf{1} \otimes d\Gamma(\hat{e}).$$

The oscillator-gas interaction is given by the unbounded perturbation

$$a^*(1) \otimes a(f) + a(1) \otimes a^*(f)$$

with $f \in \text{Dom } \hat{e}^{-1/2} \cap \text{Dom } \hat{e}$.⁷

⁷The assumption $f \in \text{Dom } \hat{e}^{-1/2} \cap \text{Dom } \hat{e}$ is a technical IR/UV regularity condition needed to have a properly defined model.

Again by the usual identification $\Gamma^+(\mathbf{C}) \otimes \Gamma^+(L^2(\mathbf{R}_+, de)) \cong \Gamma^+(\mathbf{C} \oplus L^2(\mathbf{R}_+, de))$, we obtain unitarily equivalents $d\Gamma(h_0)$ and $V := a^*(\psi_f)a(\psi_o) + a^*(\psi_o)a(\psi_f)$ on $\Gamma^+(\mathfrak{h})$ where $\psi_f = 0 \oplus f \in \mathbf{C} \oplus L^2(\mathbf{R}_+, de)$ and $\psi_o = 1 \oplus 0 \in \mathbf{C} \oplus L^2(\mathbf{R}_+, de)$.

Let $\rho = (e^{\beta h_0} - 1)^{-1}$ be the Bose–Einstein density for some inverse temperature $\beta > 0$. Let us introduce a representation of the *canonical commutation relations* (CCR) algebra over $\text{Dom } \rho^{1/2}$ known as the (glued left) Araki–Woods representation, which in its earliest form goes back to [AW63]. This representation is determined by

$$\begin{aligned} W^{\text{AW}} : \text{Dom } \rho^{1/2} &\rightarrow \mathcal{B}(\Gamma^+(\mathfrak{h} \oplus \mathfrak{h})) \\ \phi &\mapsto W(\sqrt{1 + \rho} \phi \oplus \sqrt{\rho} \bar{\phi}), \end{aligned} \quad (22)$$

where W denotes the Weyl operators on the symmetric Fock space $\Gamma^+(\mathfrak{h} \oplus \mathfrak{h})$ and $\bar{\phi}$ is the complex conjugation of ϕ . These operators satisfy the Weyl form of the CCR:

$$W^{\text{AW}}(\phi)W^{\text{AW}}(\psi) = e^{-\frac{1}{2} \text{Im}(\phi, \psi)} W^{\text{AW}}(\phi + \psi) \quad (23)$$

for all ϕ and ψ in $\text{Dom } \rho^{1/2}$. Also, we have the evaluation identity

$$\langle \Omega, W^{\text{AW}}(\phi)\Omega \rangle = e^{-\frac{1}{4} \langle \phi, \phi \rangle - \frac{1}{2} \langle \phi, \rho \phi \rangle} \quad (24)$$

for all ϕ in $\text{Dom } \rho^{1/2}$.

Let $\mathfrak{M}^{\text{AW}} \subseteq \mathcal{B}(\Gamma^+(\mathfrak{h} \oplus \mathfrak{h}))$ be the von Neumann algebra generated by

$$\{W^{\text{AW}}(\phi) : \phi \in \text{Dom } \rho^{1/2}\}.$$

The maps defined by

$$\tau^s(W^{\text{AW}}(\phi)) = W^{\text{AW}}(e^{is h_0} \phi) \quad (25)$$

for all $\phi \in \text{Dom } \rho^{1/2}$ extend to a W^* -dynamics $(\tau^s)_{s \in \mathbf{R}}$ on \mathfrak{M}^{AW} i.e. a σ -weakly continuous one-parameter group of $*$ -automorphisms of \mathfrak{M}^{AW} . We consider the state $\omega : A \mapsto \langle \Omega, A\Omega \rangle$ on \mathfrak{M}^{AW} where Ω is the vacuum in $\Gamma^+(\mathfrak{h} \oplus \mathfrak{h})$. Then, ω is a (τ, β) -KMS state on \mathfrak{M}^{AW} and the corresponding Liouvillean is $L = d\Gamma(h_0 \oplus -h_0)$. For proofs of these facts, we refer the reader to [DG13, §17] and [Der06, §9].

For each $\phi \in \text{Dom } \rho^{1/2}$, the map $\mathbf{R} \ni s \mapsto W^{\text{AW}}(s\phi)$ is strongly continuous and we will denote its generator by $\varphi^{\text{AW}}(\phi)$. It is explicitly given on the Fock space $\Gamma^+(\mathfrak{h} \oplus \mathfrak{h})$ by

$$\varphi^{\text{AW}}(\phi) = \varphi(\sqrt{1 + \rho} \phi \oplus \sqrt{\rho} \bar{\phi}).$$

The affiliated creation and annihilation operators, $a^{\text{AW}*}(\phi)$ and $a^{\text{AW}}(\phi)$, are related to $\varphi^{\text{AW}}(\phi)$ by the formula $\varphi^{\text{AW}}(\phi) = \frac{1}{\sqrt{2}}(a^{\text{AW}*}(\phi) + a^{\text{AW}}(\phi))$ and $[a^{\text{AW}}(\phi), a^{\text{AW}*}(\psi)] = \langle \phi, \psi \rangle$. They are explicitly given by the formulae

$$\begin{aligned} a^{\text{AW}*}(\phi) &= a^*(\sqrt{1 + \rho} \phi \oplus 0) + a(0 \oplus \sqrt{\rho} \bar{\phi}), \\ a^{\text{AW}}(\phi) &= a(\sqrt{1 + \rho} \phi \oplus 0) + a^*(0 \oplus \sqrt{\rho} \bar{\phi}). \end{aligned}$$

The perturbation V is represented by an operator V^{AW} affiliated to \mathfrak{M}^{AW} given by

$$V^{\text{AW}} = a^{\text{AW}*}(\psi_o)a^{\text{AW}}(\psi_f) + a^{\text{AW}*}(\psi_f)a^{\text{AW}}(\psi_o),$$

where $\psi_f = 0 \oplus f \in \text{Dom } \rho^{1/2} \subset \mathfrak{h}$ since $f \in \text{Dom } \hat{e}^{-1/2}$ and $\psi_o = 1 \oplus 0 \in \mathfrak{h}$.

Since $f \in \text{Dom } \hat{e}^{-1/2} \cap \text{Dom } \hat{e}$, the operator

$$L + V^{\text{AW}} \quad (26)$$

is essentially self-adjoint on $\text{Dom } L \cap \text{Dom } V^{\text{AW}}$ by Lemma A.1 and the family of maps $(\tau_V^s)_{s \in \mathbf{R}}$ defined by

$$\tau_V^s(W^{\text{AW}}(\phi)) = e^{it(L+V^{\text{AW}})}W^{\text{AW}}(\phi)e^{-it(L+V^{\text{AW}})}$$

for $\phi \in \text{Dom } \rho^{\frac{1}{2}}$ extends to a W^* -dynamics on \mathfrak{M}^{AW} [DJP03]. Again with the one-particle operator $v = \psi_o \langle \psi_f, \cdot \rangle + \psi_f \langle \psi_o, \cdot \rangle$, we have

$$\tau_V^s(W^{\text{AW}}(\phi)) = W^{\text{AW}}(e^{is(h_o+v)}\phi)$$

for all $\phi \in \text{Dom } \rho^{1/2}$.

The operator

$$L + V^{\text{AW}} - e^{it(L+V^{\text{AW}})}V^{\text{AW}}e^{-it(L+V^{\text{AW}})}$$

is well-defined and self-adjoint. Its self-adjointness follows from Lemma A.1 and the observation

$$L + V^{\text{AW}} - e^{it(L+V^{\text{AW}})}V^{\text{AW}}e^{-it(L+V^{\text{AW}})} = e^{it(L+V^{\text{AW}})}Le^{-it(L+V^{\text{AW}})}.$$

Hence the following definition is sound.

Definition 4.7. The distribution \mathbb{P}_t of heat variation is the spectral measure of

$$L + V^{\text{AW}} - e^{it(L+V^{\text{AW}})}V^{\text{AW}}e^{-it(L+V^{\text{AW}})}$$

with respect to Ω .

Theorem 4.8. Assume $f \in \text{Dom } \hat{e}^{-1/2} \cap \text{Dom } \hat{e}$ and $\|\hat{e}^{-1/2}f\| \neq \epsilon_o^{1/2}$. Let \mathbb{P}_t be the probability measure of Definition 4.7. Then, for any $n \in \mathbf{N}$ the following are equivalent

- (i) $\sup_{t \in \mathbf{R}} \mathbb{E}_t[\Delta Q^{2n+2}] < \infty$;
- (ii) there exist $t_1 < t_2$ such that $\int_{t_1}^{t_2} \mathbb{E}_t[\Delta Q^{2n+2}] dt < \infty$;
- (iii) $f \in \text{Dom } \hat{e}^n$.

Remark 4.9. As one can see in the proof, without the assumption $\|\hat{e}^{-1/2}f\| \neq \epsilon_o^{1/2}$, the second statement still implies the third one, and the third statement implies the second one for all time intervals (t_1, t_2) .

Proposition 4.10. Let \mathbb{P}_t be the probability measure of Definition 4.7. If there exists $\gamma > 0$ such that $f \in \text{Dom } e^{\frac{1}{2}\gamma\hat{e}}$, then there exists $\gamma' \in (0, \gamma)$ such that

$$\sup_{t \in \mathbf{R}} \mathbb{E}_t[e^{\gamma'|\Delta Q|}] < \infty. \quad (27)$$

4.3 van Hove Hamiltonian

We consider a quasi-free gas of scalar bosons interacting with a prescribed classical external scalar field. This model is the quantum counterpart to the classical model of Section 2.2. It typically models the interaction of an electromagnetic field with some prescribed classical charges.

The one-particle Hilbert space is $\mathfrak{h} = L^2(\mathbf{R}_+, d\epsilon)$ and the one-particle Hamiltonian is $h_o = \hat{e}$. The unperturbed Hamiltonian is $d\Gamma(\hat{e})$ on the Fock space $\Gamma(L^2(\mathbf{R}_+, d\epsilon))$. The prescribed classical external field induces a perturbation $V := \varphi(f)$. The perturbed Hamiltonian is

$$d\Gamma(\hat{e}) + \varphi(f)$$

with $f \in \text{Dom } \hat{e} \cap \hat{e}^{-1/2}$.

Hamiltonians of this type are often called *van Hove Hamiltonians* [Der03]. We cast this into the W^* -algebraic framework again through the Araki–Woods representation.

Let ρ be the Bose–Einstein density at inverse temperature $\beta > 0$, $\rho := (e^{\beta h_0} - 1)^{-1}$. In the corresponding Araki–Woods representation (22), the dynamics τ is again given by (25) and we consider the quasi-free state ω associated to ρ . Recall that ω is a (τ, β) -KMS state. The CCR (23) and the evaluation identity (24) still hold.

Since $f \in \text{Dom } \hat{e} \cap \text{Dom } \hat{e}^{-1/2}$, the van Hove perturbation

$$L + \varphi^{AW}(f), \quad (28)$$

of the Liouvillean is essentially self-adjoint on $\text{Dom } L \cap \text{Dom } \varphi^{AW}(f)$ by Lemma A.2. The maps

$$\tau_V^s(W^{AW}(\phi)) = e^{is(L+\varphi^{AW}(f))}W^{AW}(\phi)e^{-is(L+\varphi^{AW}(f))}$$

on $\{W^{AW}(\phi) : \phi \in \text{Dom } \rho^{1/2}\}$ extend to a W^* -dynamics on \mathfrak{M}^{AW} .

Then, as in the previous subsection, we can consider the operator

$$L + \varphi^{AW}(f) - e^{it(L+\varphi^{AW}(f))}\varphi^{AW}(f)e^{-it(L+\varphi^{AW}(f))}, \quad (29)$$

and Lemma A.2 ensures its essential self-adjointness and soundness of the following definition of the distribution of heat variation.

Definition 4.11. The distribution \mathbb{P}_t of heat variation is the spectral measure of

$$L + \varphi^{AW}(f) - e^{it(L+\varphi^{AW}(f))}\varphi^{AW}(f)e^{-it(L+\varphi^{AW}(f))}$$

with respect to Ω .

Remark 4.12. This system has the particularity that it admits a closed form expression for the characteristic function \mathcal{E}_t of \mathbb{P}_t , which allows an in-depth study of the relation between the tails of \mathbb{P}_t and the regularity of the perturbation. As in the previous models, this regularity is expressed in terms of the map $\mathbf{R} \ni s \mapsto e^{is\hat{e}}f \in L^2(\mathbf{R}_+, de)$, which is n times norm differentiable if and only if $f \in \text{Dom } \hat{e}^n$.

Theorem 4.13. Assume $f \in \text{Dom } \hat{e} \cap \text{Dom } \hat{e}^{-1/2}$. Then, the probability measure \mathbb{P}_t of Definition 4.11 is the law of an inhomogeneous Poisson process on \mathbf{R} with intensity

$$d\nu_t(e) := \frac{(1 - \cos(et))}{e^2} \frac{|f(|e|)|^2}{|1 - e^{-\beta e}|} de.$$

Remark 4.14. This distribution of the heat variation is in sharp contrast with the Gaussian heat variation of its classical analogue we discussed in Section 2.2.

Theorem 4.15. Assume $f \in \text{Dom } \hat{e} \cap \text{Dom } \hat{e}^{-1/2}$. Let \mathbb{P}_t be the probability measure of Definition 4.11. Then, for any $n \in \mathbf{N}$ the following are equivalent

- (i) $\sup_{t \in \mathbf{R}} \mathbb{E}_t[\Delta Q^{2n+2}] < \infty$;
- (ii) there exist $t_1 < t_2$ such that $\int_{t_1}^{t_2} \mathbb{E}_t[\Delta Q^{2n+2}] dt < \infty$;
- (iii) $f \in \text{Dom } \hat{e}^n$.

Theorem 4.16. Assume $f \in \text{Dom}(\hat{e}) \cap \text{Dom}(\hat{e}^{-1/2})$. Let \mathbb{P}_t be the probability measure of Definition 4.11. Then, for $\gamma > 0$, the following are equivalent

- (i) $\sup_{t \in \mathbf{R}} \mathbb{E}_t[e^{\gamma|\Delta Q|}] < \infty$;
- (ii) there exist $t_1 < t_2$ such that $\int_{t_1}^{t_2} \mathbb{E}_t[e^{\gamma|\Delta Q|}] dt < \infty$;
- (iii) $f \in \text{Dom } e^{\frac{1}{2}\gamma\hat{e}}$.

5 Proofs for quasi-free gas models

5.1 One-particle space preliminaries

Some of the results of Section 4 require technical lemmas on one-particle operators and vectors. Throughout this section, $h_0 = \epsilon_0 \oplus \hat{e}$ on $\mathfrak{h} = \mathbf{C} \oplus L^2(\mathbf{R}_+, de)$, and the one-particle perturbation v is of the form $\psi_0 \langle \psi_f, \cdot \rangle + \psi_f \langle \psi_0, \cdot \rangle$ for $\psi_f = 0 \oplus f \in \mathbf{C} \oplus L^2(\mathbf{R}_+, de)$ and $\psi_0 = 1 \oplus 0 \in \mathbf{C} \oplus L^2(\mathbf{R}_+, de)$.

First note that $\text{Dom } h_0 = \text{Dom } h$ and, if $f \in \text{Dom } \hat{e}^{n-1}$, $\text{Dom } h^n = \text{Dom } h_0^n$. Moreover, we have the following first two lemmas.

The first lemma is a somehow surprising, but quite straight forward, property of the comparison between perturbed and unperturbed one-particle unitary dynamical groups.

Lemma 5.1. *Let $n \in \mathbf{N}^*$ and assume $f \in \text{Dom } \hat{e}^{n-1}$. Then $h_0^n(e^{ith_0} - e^{ith})$ is bounded and there exists an affine function $\mathbf{R}_+ \ni t \mapsto K_n(t)$ such that*

$$\|h_0^n(e^{ith_0} - e^{ith})\| \leq K_n(|t|). \quad (30)$$

Proof. We prove the lemma for $t > 0$. The proof for $t < 0$ is similar.

Let $\chi \in \text{Dom } h_0^n$ and $\phi \in \text{Dom } h_0$. Using Duhamel's formula,

$$\begin{aligned} \langle h_0^n \chi, (e^{ith_0} - e^{ith})\phi \rangle &= i \int_0^t \langle h_0^n \chi, e^{i(t-s)h_0} v e^{ish} \phi \rangle ds, \\ &= - \int_0^t \langle \chi, (\partial_s e^{i(t-s)h_0}) h_0^{n-1} v e^{ish} \phi \rangle ds, \end{aligned}$$

Since $\phi \in \text{Dom } h$, the functions $s \mapsto \langle \psi_0, e^{ish} \phi \rangle$ and $s \mapsto \langle \psi_f, e^{ish} \phi \rangle$ are in $C^1(\mathbf{R})$. Similarly, the functions $s \mapsto \langle \chi, e^{i(t-s)h_0} h_0^{n-1} \psi_0 \rangle$ and $s \mapsto \langle \chi, e^{i(t-s)h_0} h_0^{n-1} \psi_f \rangle$ are in $C^1(\mathbf{R})$ using $\chi \in \text{Dom}(h_0^n)$ for the second function. Integrating by parts,

$$\begin{aligned} &i \int_0^t \langle \chi, (\partial_s e^{i(t-s)h_0}) h_0^{n-1} v e^{ish} \phi \rangle ds \\ &= i [\langle \chi, e^{i(t-s)h_0} h_0^{n-1} v e^{ish} \phi \rangle]_0^t + \int_0^t \langle \chi, e^{i(t-s)h_0} h_0^{n-1} v h e^{ish} \phi \rangle ds. \end{aligned}$$

Since ψ_0 and ψ_f are in $\text{Dom}(h_0^{n-1})$,

$$\|h_0^{n-1} v\| < \infty \quad \text{and} \quad \|h_0^{n-1} v h\| < \infty.$$

The inequality,

$$|\langle h_0^n \chi, (e^{ith_0} - e^{ith})\phi \rangle| \leq \|\chi\| \|\phi\| (2\|h_0^{n-1} v\| + t\|h_0^{n-1} v h\|),$$

and Riesz's Lemma yield the lemma with the affine bound

$$K_n(t) = 2\|h_0^{n-1} v\| + t\|h_0^{n-1} v h\|. \quad \square$$

The second Lemma concerns the stability of the domain of powers of h_0 with respect to the perturbed dynamical unitary group.

Lemma 5.2. *Let $n \in \mathbf{N}^*$ and suppose $f \in \text{Dom } \hat{e}^{n-1}$. If $\phi \in \text{Dom } h_0^n$, then for all $\alpha \in [0, n]$,*

$$\sup_t \|h_0^\alpha e^{ith} \phi\| < \infty. \quad (31)$$

Proof. By a concavity argument, it suffices to show (31) for $\alpha = n$. Since $f \in \text{Dom } \hat{e}^{n-1}$ implies $\text{Dom } h^n = \text{Dom } h_0^n$, the commutation $[e^{ith}, h^n] = 0$ yields directly that $e^{ith}\phi \in \text{Dom } h_0^n$.

Now, by the triangle inequality

$$\|h_0^n e^{ith}\phi\| \leq \|(h_0^n - h^n)e^{ith}\phi\| + \|h^n\phi\|. \quad (32)$$

The operator $h_0^n - h^n$ is a polynomial in operator variables $h_0^k v^p h_0^m$ with $k + p + m \leq n$, $p \geq 1$. Since $f \in \text{Dom } \hat{e}^{n-1}$, all the operator variables $h_0^k v^p h_0^m$ are bounded. Hence, $h_0^n - h^n$ is bounded and $\sup_t \|h_0^n e^{ith}\phi\| < \infty$ follows from $\phi \in \text{Dom } h^n$. \square

The last lemma of this subsection allows us to ensure that the existence of moments of \mathbb{P}_t is not the fortuitous consequence of some cycles in the evolution, but truly the consequence of $f \in \text{Dom } \hat{e}^n$. We will use it in the proof that (ii) implies (iii) in Theorems 4.4 and 4.8. For that purpose we show that a positive function of e that will appear later in our proofs is actually lower bounded away from 0.

We denote $\bar{h} := h_0 + \bar{v}$ where $\bar{v} := \psi_o \langle \bar{\psi}_f, \cdot \rangle + \bar{\psi}_f \langle \psi_o, \cdot \rangle$.

Lemma 5.3. *Let $A = \sqrt{T}$ or $A = \sqrt{\rho}$. Assume furthermore that $\psi \in \text{Dom } A \setminus \{0\}$ is not an eigenvector of h . Then, for any non-trivial interval (t_1, t_2) of \mathbf{R} ,*

$$\inf_{e \in \mathbf{R}_+} \int_{t_1}^{t_2} \|A(1 - e^{-it(\bar{h}-e)}\psi)\|^2 dt > 0.$$

Proof. Let

$$F(e) := \left\| \int_{t_1}^{t_2} A(1 - e^{-it(\bar{h}-e)}\psi) dt \right\|.$$

From triangle inequality on integrals and Jensen's inequality, it is sufficient to lower bound this function on \mathbf{R}_+ to prove the lemma.

To deal with the unboundedness of ρ , we introduce

$$\rho_{\text{cut}}(e) := \begin{cases} (e^{\beta e} - 1)^{-1} & \text{if } e \geq 1 \\ (e^\beta - 1)^{-1} & \text{if } e < 1. \end{cases}$$

Let $\rho_1 := (e^{\beta e_0} - 1)^{-1} \oplus \rho_{\text{cut}}(\hat{e})$. Since $e \mapsto (e^{\beta e} - 1)^{-1}$ is decreasing, it is a bounded operator and $\rho \geq \rho_1$. Hence lower bounding $e \mapsto F(e)$ with $A = \sqrt{\rho_1}$ implies the existence of the same lower bound for $e \mapsto F(e)$ with $A = \sqrt{\rho}$. Thus we now prove a lower bound away from 0 on $e \mapsto F(e)$ with A assumed to be a bounded positive operator with trivial kernel.

Let $y : \mathbf{R} \rightarrow \mathbf{C}$ be the function defined by

$$y(x) = \int_{t_1}^{t_2} 1 - e^{-itx} dt.$$

The function y is continuous and bounded by $2(t_2 - t_1)$ on \mathbf{R} , and $y(x) = 0$ if and only if $x = 0$. Fubini's Theorem implies,

$$F(e) = \|Ay(\bar{h} - e)\psi\|.$$

Since $\ker A = \{0\}$, it follows from $y(x) = 0 \Leftrightarrow x = 0$ that $F(e) = 0 \Leftrightarrow \bar{h}\psi = e\psi$ which contradicts our assumption that ψ is not an eigenvector of h . Hence, by continuity of F , on any compact $E \subset \mathbf{R}_+$,

$$\inf_{e \in E} F(e) > 0.$$

Hence $\inf_{e \in \mathbf{R}_+} F(e) = 0$ only if $\liminf_{e \rightarrow \infty} F(e) = 0$.

Since $\mathbf{R} \ni x \mapsto y(x - e)$ converges point wise to $t_2 - t_1$ when $e \rightarrow \infty$ and y is bounded, by Lebesgue's dominated convergence Theorem,

$$\lim_{e \rightarrow \infty} y(\bar{h} - e)\psi = (t_2 - t_1)\psi$$

in norm. Hence,

$$\liminf_{e \rightarrow \infty} F(e) = \lim_{e \rightarrow \infty} \|Ay(\bar{h} - e)\psi\| = (t_2 - t_1)\|A\psi\|.$$

The triviality of the kernel of A yields the result. \square

5.2 Proofs for the impurity in a quasi-free Fermi gas

The GNS representation we will work in is known as the (glued left) Araki–Wyss representation. This representation of the CAR was introduced in its earliest form in [AW64]. The Hilbert space for this representation is the anti-symmetric Fock space $\mathcal{H} = \Gamma^-(\mathfrak{h} \oplus \mathfrak{h})$ and the representation is determined by

$$\pi(a^*(\phi)) = a^*(\sqrt{1 - T}\phi \oplus 0) + a(0 \oplus \sqrt{T}\bar{\phi}), \quad (33)$$

$$\pi(a(\phi)) = a(\sqrt{1 - T}\phi \oplus 0) + a^*(0 \oplus \sqrt{T}\bar{\phi}), \quad (34)$$

for all $\phi \in \mathfrak{h}$. Finally the vector representative Ω of ω is the vacuum in $\Gamma^-(\mathfrak{h} \oplus \mathfrak{h})$ and the (unperturbed) Liouvillean is $L = d\Gamma(h_0 \oplus -h_0)$. For proofs of these facts, we refer the reader to [DG13, §17] and [Der06, §10]. For $\phi \in \text{Dom } h_0$, the operators L and $\pi(a^\sharp(\phi))$ satisfy the commutation relation

$$i[L, \pi(a^\sharp(\phi))]\Psi = \pi(\delta(a^\sharp(\phi)))\Psi = \pi(a^\sharp(ih_0\phi))\Psi, \quad (35)$$

for any $\Psi \in \text{Dom } L$, as seen from differentiating the relation

$$e^{isL}\pi(a^\sharp(\phi))e^{-isL}\Psi = \pi(a^\sharp(e^{ish_0}\phi))\Psi$$

obtained from (15) and (21). It follows directly from a repeated application of (35) that

$$V \in \text{Dom}(\delta^n) \iff f \in \text{Dom}(\hat{e}^n). \quad (36)$$

Proof of Theorem 4.4. The fact that the first statement implies the second one is trivial. The fact that the third one implies the first one is immediate from Theorem 3.5 and the equivalence (36).

To prove that the the second statement implies the third one, we proceed by induction on n . The result holds for $n = 0$. Assume it holds for some $n - 1 \in \mathbf{N}$ and suppose that there exists a non-trivial interval (t_1, t_2) on which $t \mapsto \mathbb{E}_t(\Delta Q^{2n+2})$ is integrable. Then, $t \mapsto \mathbb{E}_t(\Delta Q^{2n})$ is also integrable by Jensen's inequality and $f \in \text{Dom } \hat{e}^{n-1}$ by the induction hypothesis. The equivalence (36) yields $V \in \text{Dom } \delta^{n-1}$. Let $X_t := V - \tau_V^t(V)$. Then Lemma 3.12 implies $X_t \in \text{Dom } \delta^{n-1}$ and $\sup_{t \in \mathbf{R}} \|\delta^{n-1}(X_t)\| < \infty$.

Claim A $\pi(X_t)\Omega \in \text{Dom } L^n$ for a.e.⁸ $t \in (t_1, t_2)$ and

$$\int_{t_1}^{t_2} \|L^n \pi(X_t)\Omega\|^2 dt < \infty.$$

First, we prove $\Omega \in \text{Dom}(L + \pi(X_t))^{n+1}$ for a.e. $t \in (t_1, t_2)$. Let $E \mapsto P(E)$ be the projection valued spectral measure of $L + \pi(X_t)$. Let $\Phi \in \mathcal{H}$ be such that there exists $\Lambda > 0$ such that $P([- \Lambda, \Lambda])\Phi = \Phi$. Then by the Cauchy–Schwarz inequality,

$$|\langle \Phi, (L + \pi(X_t))^{n+1}\Omega \rangle|^2 \leq \|\Phi\|^2 \|P([- \Lambda, \Lambda])(L + \pi(X_t))^{n+1}\Omega\|^2.$$

⁸Here and in what follows, we use “a.e.” for almost every, with respect to the Lebesgue measure.

The norm $\|P([- \Lambda, \Lambda])(L + \pi(X_t))^{n+1}\Omega\|$ is non-decreasing as a function of Λ . Then (ii) implies that for a.e. $t \in (t_1, t_2)$, $\lim_{\Lambda \rightarrow \infty} \|P([- \Lambda, \Lambda])(L + \pi(X_t))^{n+1}\Omega\|^2 = \mathbb{E}_t(\Delta Q^{2n+2})$. Hence for a.e. $t \in (t_1, t_2)$,

$$|\langle \Phi, (L + \pi(X_t))^{n+1}\Omega \rangle|^2 \leq \|\Phi\|^2 \mathbb{E}_t(\Delta Q^{2n+2})$$

and Riesz's Lemma implies $\Omega \in \text{Dom}(L + \pi(X_t))^{n+1}$.

Now we prove **Claim A**. By the induction hypothesis, Lemma 5.2 and the equivalence (36), commuting repeatedly L with $\pi(X_t)$, it follows that $\pi(X_t) \text{Dom } L^{n-1} \subset \text{Dom } L^{n-1}$. Hence any non-commutative polynomial in L and $\pi(X_t)$ with at most $n-1$ occurrences of L in each monomial preserves $\text{Dom } L^{n-1}$. Working in a representation of \mathcal{H} in which L is a multiplication operator, we can develop $(L + \pi(X_t))^{n+1}$. Using $L\Omega = 0$ and $i[L, \pi(\delta^p(X_t))] = \pi(\delta^{p+1}(X_t))$ for $p = 0, \dots, n-2$ we obtain,

$$(L + \pi(X_t))^{n+1}\Omega = L^n \pi(X_t)\Omega + Q_t \Omega \quad (37)$$

with Q_t a non-commutative polynomial in $\{\pi(\delta^p(X_t))\}_{p=0, \dots, n-1}$. Since $\Omega \in \text{Dom}(L + \pi(X_t))^{n+1}$ and $Q_t \Omega \in \mathcal{H}$, we obtain $\pi(X_t)\Omega \in \text{Dom } L^n$. Moreover, Lemma 5.1 implies,

$$\sup_{t \in \mathbf{R}} \|Q_t \Omega\| < \infty.$$

Hence, **Claim A** follows from (37), the triangle inequality, assumption (ii) and Jensen's inequality.

Claim B $f \in \text{Dom } \hat{e}^n$

Using the explicit expressions in Fock space, the non-trivial part of $L^n \pi(X_t)\Omega$ lives in the two-particle subspace of $\Gamma^-(\mathfrak{h} \oplus \mathfrak{h})$ only, namely $P_2 L^n \pi(X_t)\Omega = L^n \pi(X_t)\Omega$ with P_2 the projector onto the two-particle subspace of $\Gamma^-(\mathfrak{h} \oplus \mathfrak{h})$. The vector there is

$$(\sqrt{\mathbf{1} - T} h_0^n \psi_o \oplus 0) \wedge (0 \oplus \sqrt{T} \bar{\psi}_f) - (\sqrt{\mathbf{1} - T} h_0^n e^{ith} \psi_o \oplus 0) \wedge (0 \oplus \sqrt{T} e^{-it\bar{h}} \bar{\psi}_f) \quad (38)$$

$$+ (\sqrt{\mathbf{1} - T} \psi_o \oplus 0) \wedge (0 \oplus \sqrt{T} h_0^n \bar{\psi}_f) + (\sqrt{\mathbf{1} - T} \psi_f \oplus 0) \wedge (0 \oplus \sqrt{T} h_0^n \psi_o) \quad (39)$$

$$- (\sqrt{\mathbf{1} - T} e^{ith} \psi_f \oplus 0) \wedge (0 \oplus \sqrt{T} h_0^n e^{-it\bar{h}} \psi_o) - (\sqrt{\mathbf{1} - T} e^{ith} \psi_o \oplus 0) \wedge (0 \oplus \sqrt{T} h_0^n e^{-it\bar{h}} \bar{\psi}_f) \quad (40)$$

$$+ (\sqrt{\mathbf{1} - T} h_0^n \psi_f \oplus 0) \wedge (0 \oplus \sqrt{T} \psi_o) - (\sqrt{\mathbf{1} - T} h_0^n e^{ith} \psi_f \oplus 0) \wedge (0 \oplus \sqrt{T} e^{-it\bar{h}} \psi_o). \quad (41)$$

Since $\psi_s \in \text{Dom } h_0^n$, Lemma 5.2 implies the norm of (38) in $(\mathfrak{h} \oplus \mathfrak{h}) \wedge (\mathfrak{h} \oplus \mathfrak{h})$ is uniformly bounded in $t \in \mathbf{R}$. Since $\sqrt{T} h_0^n$ is bounded, the norm of (39)+(40) is also uniformly bounded in $t \in \mathbf{R}$. Now, since, by **Claim A**, $L^n \pi(X_t)\Omega \in \mathcal{H}$ for a.e. $t \in (t_1, t_2)$, (41) has finite norm for a.e. $t \in (t_1, t_2)$. Recall that Lemma 5.1 implies $\|h_0^n(e^{ith_0} - e^{ith})\| \leq K_n(|t|)$ with $\mathbf{R}_+ \ni t \mapsto K_n(t)$ an affine function. Because of this and the fact that $\sqrt{\mathbf{1} - T}$ is bounded, there exists a family $(\phi_t)_{t \in \mathbf{R}}$ in $(\mathfrak{h} \oplus \mathfrak{h}) \wedge (\mathfrak{h} \oplus \mathfrak{h})$ with square integrable norm on (t_1, t_2) and such that (41) is equal to

$$\begin{aligned} & (\sqrt{\mathbf{1} - T} h_0^n \psi_f \oplus 0) \wedge (0 \oplus \sqrt{T} \psi_o) \\ & - (\sqrt{\mathbf{1} - T} h_0^n e^{ith_0} \psi_f \oplus 0) \wedge (0 \oplus \sqrt{T} e^{-it\bar{h}} \psi_o) + \phi_t. \end{aligned} \quad (42)$$

Since the norms of (38), (39) and (40) are uniformly bounded in time, and since $\|\phi_t\|^2$ is integrable on (t_1, t_2) , using the triangle inequality, **Claim A** and Jensen's inequality, it follows that $t \mapsto \|(42) - \phi_t\|^2$ is integrable on (t_1, t_2) :

$$\begin{aligned} & \int_{t_1}^{t_2} \|(\sqrt{\mathbf{1} - T} h_0^n \psi_f \oplus 0) \wedge (0 \oplus \sqrt{T} \psi_o) \\ & - (\sqrt{\mathbf{1} - T} h_0^n e^{ith_0} \psi_f \oplus 0) \wedge (0 \oplus \sqrt{T} e^{-it\bar{h}} \psi_o)\|^2 dt < \infty. \end{aligned}$$

Calculating explicitly the norm squared,

$$\int_{t_1}^{t_2} \int_{\mathbf{R}_+} \left| \frac{1}{e^{-\beta e} + 1} \right| e^{2n} |f(e)|^2 \|\sqrt{T}(\mathbf{1} - e^{-it(\bar{h}-e)})\psi_o\|^2 de dt < \infty. \quad (43)$$

By Fubini's theorem,

$$\int_{\mathbf{R}_+} \left| \frac{1}{e^{-\beta e} + 1} \right| e^{2n} |f(e)|^2 \int_{t_1}^{t_2} \|\sqrt{T}(\mathbf{1} - e^{-it(\bar{h}-e)})\psi_o\|^2 dt de < \infty. \quad (44)$$

Lemma 5.3 then implies

$$\int_{\mathbf{R}_+} e^{2n} |f(e)|^2 de < \infty. \quad \square$$

5.3 Proofs for the open harmonic oscillator

We begin with a few remarks that are immediate from the definitions. One computes directly from the definitions that for any $\phi \in \text{Dom } h_0 \cap \text{Dom } h_0^{-1/2}$,

$$i[L, a^{\text{AW}}(\phi)]\Psi = a^{\text{AW}}(ih_0\phi)\Psi \quad (45)$$

for all $\Psi \in \mathcal{H}_{\text{fin}} \cap \text{Dom } L$ with \mathcal{H}_{fin} the subspace of $\Gamma^+(\mathfrak{h} \oplus \mathfrak{h})$ spanned by the vectors with a fixed number of particles:

$$\mathcal{H}_{\text{fin}} := \text{linspan}\{a^*(\phi_n) \cdots a^*(\phi_1)\Omega : n \in \mathbf{N}, \phi_j \in \mathfrak{h} \oplus \mathfrak{h}, j = 1, \dots, n\}.$$

In order to prove Theorem 4.8, we will need the following lemma to control the eventual buildup of infrared irregularities.

Lemma 5.4. *Assume $\|\hat{e}^{-1/2}f\| \neq \epsilon_o^{1/2}$ and $\phi \in \text{Dom } h_0^{-1/2}$, then*

$$\sup_{t \in \mathbf{R}} \|h_0^{-1/2} e^{ith} \phi\| < \infty.$$

Proof. Since $\ker h_0 = \{0\}$, $h_0^{\pm 1/2} h_0^{\mp 1/2} = \mathbf{1}$ and

$$h_0^{-1/2} h h_0^{-1/2} = \mathbf{1} + (\epsilon_o^{-1/2} \psi_o) \langle h_0^{-1/2} \psi_f, \cdot \rangle + (h_0^{-1/2} \psi_f) \langle \epsilon_o^{-1/2} \psi_o, \cdot \rangle.$$

Since $\psi_f \in \text{Dom } h_0^{-1/2}$, $h_0^{-1/2} h h_0^{-1/2}$ is bounded and so is $h_0^{-1/2} h^{1/2}$. The operator $h_0^{-1/2} h h_0^{-1/2}$ has spectrum

$$\text{sp}(h_0^{-1/2} h h_0^{-1/2}) = \{1, 1 \pm \epsilon_o^{-1/2} \|\hat{e}^{-1/2}f\|\}. \quad (46)$$

Suppose now that there exists $\psi \in \ker h \setminus \{0\}$. Then $\psi \in \text{Dom } h = \text{Dom } h_0$, so $\psi \in \text{Dom } h_0^{1/2}$. Then $\psi' = h_0^{1/2} \psi$ is an eigenvector of $h_0^{-1/2} h h_0^{-1/2}$ of eigenvalue 0. Since, $\|\hat{e}^{-1/2}f\| \neq \epsilon_o^{1/2}$ implies 0 is not an eigenvalue of $h_0^{-1/2} h h_0^{-1/2}$, the hypothesis $\|\hat{e}^{-1/2}f\| \neq \epsilon_o^{1/2}$ implies $\ker h = \{0\}$ and $h^{\pm 1} h^{\mp 1} = \mathbf{1}$.

Hence, $\|\hat{e}^{-1/2}f\| \neq \epsilon_o^{1/2}$ implies that $(h_0^{-1/2} h h_0^{-1/2})^{-1} = h_0^{1/2} h^{-1} h_0^{1/2}$ is bounded, and so is $h_0^{1/2} h^{-1/2}$.

The inequality

$$|\langle \psi, h^{-1/2} \phi \rangle| \leq \|h_0^{1/2} h^{-1/2}\| \|\psi\| \|h_0^{-1/2} \phi\|.$$

for all $\psi \in \text{Dom } h_0^{-1/2}$ shows that $\phi \in \text{Dom } h^{-1/2}$. Therefore,

$$\|h_0^{-1/2} e^{ith} \phi\| = \|h_0^{-1/2} h^{1/2} e^{ith} h^{-1/2} \phi\| \leq \|h_0^{-1/2} h^{1/2}\| \|h^{-1/2} \phi\|$$

which is independent of t . □

Remark 5.5. As shown in the proof, assumption $\|\hat{e}^{-1/2}f\| \neq \epsilon_o^{1/2}$ implies $\ker h = \{0\}$ which might be a more recognizable assumption to the reader. Assuming $f \in \text{Dom}(\hat{e}^{-1})$, the implication is an equivalence. If $f \notin \text{Dom}(\hat{e}^{-1})$ and $\|\hat{e}^{-1/2}f\| = \epsilon_o^{1/2}$, 0 may be in the singular continuous spectrum of h (see [JKP06]).

Proof of Theorem 4.8. The fact that the first statement implies the second one is trivial.

(ii) \implies (iii): As for the fermionic case, we proceed by induction on n . The implication is true for $n = 0$. Assume it holds for some $n - 1 \in \mathbf{N}$ and suppose that there exists a non-trivial interval (t_1, t_2) on which $t \mapsto \mathbb{E}_t(\Delta Q^{2n+2})$ is integrable. Then, $t \mapsto \mathbb{E}_t(\Delta Q^{2n})$ is integrable by Jensen's inequality and $f \in \text{Dom} \hat{e}^{n-1}$ by induction hypothesis. Consider the set

$$\mathcal{G}_{n-1} := \{(\rho + 1)^{1/2} h_0^p e^{ish} \psi_f \oplus 0, (\rho + 1)^{1/2} h_0^p e^{ish} \psi_o \oplus 0, \\ 0 \oplus \rho^{1/2} (-h_0)^p e^{-is\bar{h}} \overline{\psi_f}, 0 \oplus \rho^{1/2} (-h_0)^p e^{-is\bar{h}} \psi_o\}_{s \in \mathbf{R}; p=0, \dots, n-1}$$

of vectors in $\mathfrak{h} \oplus \mathfrak{h}$. Lemmas 5.2 and 5.4 ensure that

$$m := \sup_{\phi \in \mathcal{G}_{n-1}} \|\phi\| < \infty \quad (47)$$

We set $X_t := V^{\text{AW}} - e^{it(L+V^{\text{AW}})} V^{\text{AW}} e^{-it(L+V^{\text{AW}})}$.

Claim A $X_t \Omega \in \text{Dom} L^n \cap \mathcal{H}_{\text{fin}}$ for a.e. $t \in (t_1, t_2)$ and

$$\int_{t_1}^{t_2} \|L^n X_t \Omega\|^2 dt < \infty.$$

Since both L and $L + X_t$ are self-adjoint, the proof uses the same arguments as the ones used in the proof of **Claim A** in the proof of Theorem 4.4. The only difference lays in the proof that $\sup_{t \in \mathbf{R}} \|Q_t \Omega\| < \infty$. We no longer have that $\pi(\delta^p(X_t))$ is uniformly bounded in norm for $p = 0, \dots, n - 1$. Instead, using

$$\|a^\sharp(\phi_k) \dots a^\sharp(\phi_1) \Omega\| \leq \sqrt{(k+1)!} \prod_{i=1}^k \|\phi_i\| \quad (48)$$

for $(\phi_i)_{i=1}^k \subset \mathfrak{h} \oplus \mathfrak{h}$, the uniform bound (47) leads to

$$\sup_{t \in \mathbf{R}} \|Q_t \Omega\| < \infty. \quad (49)$$

We leave the complete adaptation of the proof to the interested reader.

Claim B $f \in \text{Dom} \hat{e}^n$.

Using the explicit expressions in Fock space $L^n X_t \Omega$ lives in the two-particle subspace of $\Gamma^-(\mathfrak{h} \oplus \mathfrak{h})$ only. As in the proof of Theorem 4.4, using (47), (48) and Lemma 5.1 we deduce from **Claim A** that,

$$\int_{t_1}^{t_2} \|((\rho + 1)^{1/2} h_0^n \psi_f \oplus 0) \otimes_s (0 \oplus \rho^{1/2} \psi_o) \\ - ((\rho + 1)^{1/2} h_0^n e^{ith_0} \psi_f \oplus 0) \otimes_s (0 \oplus \rho^{1/2} e^{-it\bar{h}} \psi_o)\|^2 dt < \infty.$$

But because the two terms arising from the symmetrized tensor product are orthogonal, we have

$$\int_{t_1}^{t_2} \|(\rho + 1)^{1/2} h_0^n \psi_f \otimes \rho^{1/2} \psi_o - (\rho + 1)^{1/2} h_0^n e^{ith_0} \psi_f \otimes \rho^{1/2} e^{-it\bar{h}} \psi_o\|^2 dt < \infty.$$

Therefore, using the spectral representation of h_0 ,

$$\int_{t_1}^{t_2} \int_0^\infty (\rho(e) + 1) |f(e)|^2 e^{2n} \|\rho^{1/2} (\mathbf{1} - e^{-it(\bar{h}-e)}) \psi_0\|^2 de dt < \infty.$$

By Fubini's theorem, $\rho(e) + 1 \geq 1$ and Lemma 5.3, we conclude

$$\int_0^\infty e^{2n} |f(e)|^2 de < \infty \quad (50)$$

and **Claim B** holds for n . By induction, (ii) implies (iii).

(iii) \implies (i): Note that Lemmas 5.2 and 5.4 ensures that if the third statement holds, then

$$m := \sup_{\phi \in \mathcal{G}_n} \|\phi\| < \infty \quad (51)$$

Developing the power of $L + X_t$, using $L\Omega = 0$ and $i[L, a^{AW^\#}(\phi)] = a^{AW^\#}(ih_0\phi)$ for any $\phi \in \text{Dom } h_0$ we obtain $(L + X_t)^{n+1}\Omega = Q_t\Omega$ with Q_t a non-commutative polynomial in $a^\#(\phi)$, $\phi \in \mathcal{G}_n$. It follows then from, (48) and (51) that

$$\sup_{t \in \mathbf{R}} \|Q_t\Omega\| < \infty. \quad \square$$

Remark 5.6. If we restrict \mathcal{G}_n to $t < T$, then following the same proof, $\sup_{t \in [0, T]} \mathbb{E}_t(\Delta Q^{2n+2}) < \infty$. In that case the assumption $\|\hat{e}^{-1/2}f\| \neq \epsilon_0^{1/2}$ is not necessary. Indeed, one can use $f \in \text{Dom } \hat{e}^{-1/2}$ and $\phi \in \text{Dom } h_0^{-1/2}$ implies $t \mapsto \|h_0^{-1/2}e^{ith}\phi\|$ is bounded by a continuous function instead of Lemma 5.4. This implication is easily proved using Riesz's Lemma and Duhamel's formula.

We turn to the proof of Proposition 4.10. We first establish two lemmas.

Lemma 5.7. *If there exists $\gamma > 0$ such that $f \in \text{Dom } e^{\frac{1}{2}\gamma\hat{e}}$, then for any $\psi \in \text{Dom } e^{\frac{1}{2}\gamma h_0}$, and any $\gamma' \in [0, \frac{1}{2}\gamma)$,*

$$\sup_{t \in \mathbf{R}} (\|\rho^{1/2} e^{\gamma' h_0} e^{ith}\psi\| + \|(1 + \rho)^{1/2} e^{\gamma' h_0} e^{ith}\psi\|) < \infty.$$

Proof. First, let us show that $e^{\pm\frac{1}{2}\gamma h_0} e^{\mp\frac{1}{2}\gamma h}$ is bounded. Let $\phi \in \text{Dom } e^{\pm\frac{1}{2}\gamma h_0}$ and $\phi' \in \text{Dom } e^{\mp\frac{1}{2}\gamma h}$. Then, by repeated differentiation,

$$\begin{aligned} & \langle \phi, e^{\pm\frac{1}{2}\gamma h_0} e^{\mp\frac{1}{2}\gamma h} \phi' \rangle \\ &= \sum_n (\mp 1)^n \int_{0 < \alpha_1 < \dots < \alpha_n < \frac{1}{2}\gamma} \langle \phi, e^{\pm\alpha_1 h_0} v e^{\mp\alpha_1 h_0} \dots e^{\pm\alpha_n h_0} v e^{\mp\alpha_n h_0} \phi' \rangle \\ & \hspace{15em} d\alpha_1 \dots d\alpha_n. \end{aligned}$$

The series on the right-hand side converges uniformly in ϕ and ϕ' since assumption $\psi_f \in \text{Dom } e^{\frac{1}{2}\gamma h_0}$ implies

$$R(\gamma) := \sup_{\alpha \in [-\frac{1}{2}\gamma, \frac{1}{2}\gamma]} \|e^{\alpha h_0} v e^{-\alpha h_0}\| < \infty.$$

Then, the Cauchy–Schwarz inequality implies

$$|\langle \phi, e^{\pm\frac{1}{2}\gamma h_0} e^{\mp\frac{1}{2}\gamma h} \phi' \rangle| \leq \|\phi\| \|\phi'\| e^{\frac{1}{2}\gamma R(\gamma)}.$$

Thus $e^{\pm\frac{1}{2}\gamma h_0} e^{\mp\frac{1}{2}\gamma h}$ is bounded by $e^{\frac{1}{2}\gamma R(\gamma)}$.

Inserting the identity $e^{-\frac{1}{2}\gamma h} e^{\frac{1}{2}\gamma h}$ we obtain,

$$\|e^{\frac{1}{2}\gamma h_0} e^{ith}\psi\| = \|e^{\frac{1}{2}\gamma h_0} e^{-\frac{1}{2}\gamma h} e^{ith} e^{\frac{1}{2}\gamma h} \psi\| \leq e^{\frac{1}{2}\gamma R(\gamma)} \|e^{\frac{1}{2}\gamma h} \psi\|.$$

Inserting the identity $e^{-\frac{1}{2}\gamma h_0} e^{\frac{1}{2}\gamma h_0}$, we obtain,

$$\|e^{\frac{1}{2}\gamma h_0} e^{ith} \psi\| \leq e^{\frac{1}{2}\gamma R(\gamma)} \|e^{\frac{1}{2}\gamma h} e^{-\frac{1}{2}\gamma h_0} e^{\frac{1}{2}\gamma h_0} \psi\| \leq e^{\gamma R(\gamma)} \|e^{\frac{1}{2}\gamma h_0} \psi\|$$

and we conclude

$$\sup_{t \in \mathbf{R}} \|e^{\frac{1}{2}\gamma h_0} e^{ith} \psi\| < \infty. \quad (52)$$

Since $e e^{\frac{1}{2}\gamma' e} \leq \frac{2}{\epsilon} e^{\frac{1}{2}(\gamma' + \epsilon)e}$ for any $\gamma > 0$, $\epsilon > 0$ and $e \geq 0$, for any $x \in [0, \frac{1}{2}[$,

$$h_0 e^{x\gamma h_0} \leq \frac{2}{(\frac{1}{2} - x)\gamma} e^{\frac{1}{2}\gamma h_0}.$$

Then, since $\sqrt{\rho} h_0$ is bounded,

$$\|\sqrt{\rho} e^{x\frac{1}{2}\gamma h_0} \psi\| \leq \|\sqrt{\rho} h_0\| \frac{2}{(\frac{1}{2} - x)\gamma} \|e^{\frac{1}{2}\gamma h_0} e^{ith} \psi\|.$$

It then follows from (52) that

$$\sup_{t \in \mathbf{R}} \|\rho^{1/2} e^{x\gamma h_0} e^{ith} \psi\| < \infty.$$

Similarly

$$\sup_{t \in \mathbf{R}} \|(\mathbf{1} + \rho)^{1/2} e^{x\gamma h_0} e^{ith} \psi\| < \infty. \quad \square$$

Lemma 5.8. *If $f \in \text{Dom } \hat{e} \cap \text{Dom } \hat{e}^{-1/2}$, then there exists $s_0 > 0$ such that*

$$\sup_{t \in \mathbf{R}} \left\| e^{is(L+X_t)} \Omega - \Omega + \sum_{n=1}^N i^n \int_0^s \cdots \int_0^{s_{n-1}} \tau^{s_n}(X_t) \cdots \tau^{s_1}(X_t) \Omega \, ds_n \cdots ds_1 \right\|$$

converges to 0 as $N \rightarrow \infty$, for all $s \in (-s_0, s_0)$.

Proof. Since $L\Omega = 0$, $e^{is(L+X_t)} \Omega = e^{is(L+X_t)} e^{-isL} \Omega$ and for $\Psi \in \mathcal{H}_{\text{fin}}$,

$$\partial_s e^{is(L+X_t)} e^{-isL} \Psi = i e^{is(L+X_t)} e^{-isL} \tau^s(X_t) \Psi.$$

Since $f \in \text{Dom } \hat{e}^{-1/2}$ implies $\tau^{s_n}(X_t) \cdots \tau^{s_1}(X_t) \Omega \in \mathcal{H}_{\text{fin}}$, a repeated application of the fundamental theorem of calculus yields that for any $N \in \mathbf{N}$,

$$\begin{aligned} & e^{is(L+X_t)} e^{-isL} \Omega \\ &= 1 + \sum_{n=1}^N i^n \int_0^s \cdots \int_0^{s_{n-1}} \tau^{s_n}(X_t) \cdots \tau^{s_1}(X_t) \Omega \, ds_n \cdots ds_1 + R_N \end{aligned} \quad (53)$$

with R_N being

$$i^{N+1} \int_0^s \cdots \int_0^{s_N} e^{is_{N+1}(L+X_t)} e^{-is_{N+1}L} \tau^{s_{N+1}}(X_t) \cdots \tau^{s_1}(X_t) \Omega \, ds_{N+1} \cdots ds_1.$$

Then,

$$\|R_N\| \leq \frac{|s|^{N+1}}{(N+1)!} \sup_{s_1, \dots, s_{N+1} \in [0, s]} \|\tau^{s_{N+1}}(X_t) \cdots \tau^{s_1}(X_t) \Omega\|.$$

Let

$$\mathcal{G}_0 := \{(\rho + 1)^{1/2} e^{ish_0} e^{ith} \psi_f \oplus 0, (\rho + 1)^{1/2} e^{ish_0} e^{ith} \psi_o \oplus 0, \\ 0 \oplus \rho^{1/2} e^{ish_0} e^{-it\bar{h}} \overline{\psi_f}, 0 \oplus \rho^{1/2} e^{ish_0} e^{-it\bar{h}} \overline{\psi_o}\}_{s \in \mathbf{R}, t \in \mathbf{R}}.$$

Lemma 5.4 implies

$$m := \sup_{\phi \in \mathcal{G}_0} \|\phi\| < \infty. \quad (54)$$

The operators $\tau^s(X_t)$ are non-commutative polynomials of degree 2 in $a^\sharp(\phi)$, $\phi \in \mathcal{G}_0$. They each consists of 16 monomials. It thus follows from (48) that

$$\|R_N\| \leq \frac{\sqrt{(2N+1)!}}{(N+1)!} (16m^2|s|)^{N+1}$$

Since $\limsup_{N \rightarrow \infty} 2^{-N} \frac{\sqrt{(2N+1)!}}{(N+1)!} < \infty$, $\lim_{N \rightarrow \infty} \|R_N\| = 0$ for $|s| < \frac{1}{32m^2}$. \square

Proof of Proposition 4.10. Let $F_{t,N} : \mathbf{R} \rightarrow \mathbb{C}$ be defined by the truncated series

$$F_{t,N}(\alpha) := 1 + \sum_{n=1}^N i^n \int_0^\alpha \cdots \int_0^{\alpha_{n-1}} \langle \Omega, \tau^{\alpha_n}(X_t) \cdots \tau^{\alpha_1}(X_t) \Omega \rangle d\alpha_n \cdots d\alpha_1.$$

Let $\gamma' \in (0, \gamma)$ and

$$\mathcal{G}_{\gamma'} := \{(\rho + 1)^{1/2} e^{\alpha h_0} e^{ish} \psi_f \oplus 0, (\rho + 1)^{1/2} e^{\alpha h_0} e^{ish} \psi_o \oplus 0, \\ 0 \oplus \rho^{1/2} e^{\alpha h_0} e^{-is\bar{h}} \overline{\psi_f}, 0 \oplus \rho^{1/2} e^{\alpha h_0} e^{-is\bar{h}} \overline{\psi_o}\}_{s \in \mathbf{R}, |\operatorname{Im} \alpha| < \frac{1}{2}\gamma'}.$$

Lemma 5.7 implies,

$$m := \sup_{\phi \in \mathcal{G}_{\gamma'}} \|\phi\| < \infty. \quad (55)$$

Note that the operators $\tau^{\alpha_i}(X_t)$ are non-commutative polynomials of degree 2 in $a^\sharp(\phi)$, $\phi \in \mathcal{G}_{\gamma'}$. Each one of this polynomial consists of 16 monomials.

Using Wick's Theorem, each integrand in the truncated series $F_{t,N}$ admits an analytic extension to $\{(\alpha_1, \dots, \alpha_n) \in \mathbf{C}^n \mid |\operatorname{Im} \alpha_j| < \frac{1}{2}\gamma'\}$. Hence each $F_{t,N}$ admits an analytic extension to $\{\alpha \in \mathbf{C} : |\operatorname{Im} \alpha| < \frac{1}{2}\gamma'\}$.

Lemma 5.8 yields convergence of the sequence $(F_{t,N})_{N \in \mathbf{N}}$ on a subset of $\{\alpha \in \mathbf{C} : |\operatorname{Im} \alpha| < \frac{1}{2}\gamma'\}$ that has an accumulation point and the limit there coincides with the characteristic function of \mathbb{P}_t .

From (55),

$$\left\| \int_0^\alpha \cdots \int_0^{\alpha_{n-1}} \langle \Omega, \tau^{\alpha_n}(X_t) \cdots \tau^{\alpha_1}(X_t) \Omega \rangle d\alpha_n \cdots d\alpha_1 \right\| \\ \leq \frac{|\alpha|^n}{n!} \|\tau^{\alpha_n}(X_t) \cdots \tau^{\alpha_1}(X_t) \Omega\| \\ \leq \frac{\sqrt{(2n+1)!}}{n!} (16m^2|\alpha|)^n.$$

Since $\limsup_{n \rightarrow \infty} 2^{-n} \frac{\sqrt{(2n+1)!}}{n!} < \infty$, the sequence $(F_{t,N})_{N \in \mathbf{N}}$ is uniformly bounded on $\mathcal{D} := \{\alpha \in \mathbf{C} : |\alpha| < \frac{1}{32m^2} \text{ and } |\operatorname{Im} \alpha| < \frac{1}{2}\gamma'\}$. We conclude by the Vitali–Porter convergence theorem that $(F_{t,N})_{N \in \mathbf{N}}$ converges on \mathcal{D} to an analytic function. Therefore, for $\gamma' > 0$ small enough,

$$\sup_{t \in \mathbf{R}} \mathbb{E}_t[e^{\gamma'|\Delta Q|}] \leq \sup_{t \in \mathbf{R}} (\langle \Omega, e^{\gamma'(L+X_t)} \Omega \rangle + \langle \Omega, e^{-\gamma'(L+X_t)} \Omega \rangle)$$

is finite. \square

5.4 Proofs for the van Hove Hamiltonian

Proof of Theorem 4.13. Using $L\Omega = 0$, the characteristic function of \mathbb{P}_t can be rewritten as

$$\begin{aligned}\mathcal{E}_t(\alpha) &= \langle e^{-it(L+\varphi^{\text{AW}}(f))}\Omega, e^{i\alpha L}e^{-it(L+\varphi^{\text{AW}}(f))}e^{-i\alpha L}\Omega \rangle \\ &= \langle e^{-it(L+\varphi^{\text{AW}}(f))}\Omega, e^{-it(L+\varphi^{\text{AW}}(e^{i\alpha h_0}f))}\Omega \rangle.\end{aligned}\tag{56}$$

Using the Lie–Trotter–Kato formula, (23) and $L\Omega = 0$, for any t and α in \mathbf{R} ,

$$\begin{aligned}e^{-it(L+\varphi^{\text{AW}}(e^{i\alpha h_0}f))}\Omega &= \lim_{n \rightarrow \infty} (e^{-i\frac{t}{n}L}e^{-i\frac{t}{n}\varphi^{\text{AW}}(e^{i\alpha h_0}f)})^n \Omega \\ &= e^{i\theta} \lim_{n \rightarrow \infty} e^{-i\varphi^{\text{AW}}(\sum_{k=1}^n \frac{t}{n} e^{i\alpha h_0} e^{-i\frac{kt}{n}h_0} f)} \Omega,\end{aligned}\tag{57}$$

where $e^{i\theta}$ is a phase that is independent of α and thus cancels in (56).

Let $\mathcal{C} := \text{linspan}\{a^*(\phi_n) \cdots a^*(\phi_1)\Omega : n \in \mathbf{N}, \phi_i \in \text{Dom}(h_0^{-1/2} \oplus h_0^{-1/2})\}$. This is a core for all of the field operators $\varphi^{\text{AW}}(\sum_{k=1}^n \frac{t}{n} e^{i\alpha h_0} e^{-i\frac{kt}{n}h_0} f)$ and $\varphi^{\text{AW}}(ie^{i\alpha h_0} h_0^{-1}(e^{-ith_0} - 1)f)$, where $x \mapsto x^{-1}(e^{-itx} - 1)$ is extended to $x = 0$ by continuity. From the convergence of the Riemann sums to a Riemann integral,

$$\lim_{n \rightarrow \infty} \varphi^{\text{AW}}\left(\sum_{k=1}^n \frac{t}{n} e^{i\alpha h_0} e^{-i\frac{kt}{n}h_0} f\right) \Psi = \varphi^{\text{AW}}(ie^{i\alpha h_0} h_0^{-1}(e^{-ith_0} - 1)f) \Psi.$$

for every $\Psi \in \mathcal{C}$. It then follows from Proposition VIII.25 of [RS72] that

$$\text{s-r-lim}_{n \rightarrow \infty} \varphi^{\text{AW}}\left(\sum_{k=1}^n \frac{t}{n} e^{i\alpha h_0} e^{-i\frac{kt}{n}h_0} f\right) = \varphi^{\text{AW}}(ie^{i\alpha h_0} h_0^{-1}(e^{-ith_0} - 1)f).$$

Hence, from (57),

$$e^{-it(L+\varphi^{\text{AW}}(e^{i\alpha h_0}f))}\Omega = e^{i\theta} e^{-i\varphi^{\text{AW}}(ie^{i\alpha h_0} h_0^{-1}(e^{-ith_0} - 1)f)} \Omega.$$

Setting $f_t := -ih_0^{-1}(e^{-ith_0} - 1)f$, from this equality and (56), relations (23) and (24), imply

$$\begin{aligned}\mathcal{E}_t(\alpha) &= \langle W^{\text{AW}}(f_t)\Omega, W^{\text{AW}}(e^{i\alpha h_0} f_t)\Omega \rangle \\ &= \langle \Omega, W^{\text{AW}}((e^{i\alpha h_0} - 1)f_t)\Omega \rangle e^{\frac{i}{2} \text{Im}(\langle f_t, (e^{i\alpha h_0} - 1)f_t \rangle)} \\ &= e^{\frac{i}{2} \langle (1+\rho)^{1/2} f_t, (e^{i\alpha h_0} - 1)(1+\rho)^{1/2} f_t \rangle + \frac{i}{2} \langle \rho^{1/2} f_t, (e^{-i\alpha h_0} - 1)\rho^{1/2} f_t \rangle}.\end{aligned}$$

Writing explicitly the inner products as integrals with respect to $d\epsilon$, using $1 + \rho = (1 - e^{-\beta h_0})^{-1}$ and a change of variable $e \rightarrow -e$ in the second inner product,

$$\mathcal{E}_t(\alpha) = \exp\left(\int_{\mathbf{R}} (e^{i\alpha e} - 1) d\nu_t(e)\right).$$

Lévy–Khintchine’s canonical representation of infinitely divisible distributions yields the proposition. \square

Proof of Theorem 4.15. The implication (i) \implies (ii) is obvious.

Since Proposition 4.13 implies \mathbb{P}_t is the measure of an inhomogeneous Poisson process with intensity $d\nu_t(e)$, its $2n + 2^{\text{nd}}$ cumulant is given by $\int_{\mathbf{R}} e^{2n+2} d\nu_t(e)$. Then (iii) implies this cumulant is uniformly bounded and the equivalence between the existence of a uniform bound on the $2n + 2^{\text{nd}}$ cumulant and $2n + 2^{\text{nd}}$ moment yields (i).

For the implication (ii) \implies (iii), we first show that (ii) implies the $2n + 2^{\text{nd}}$ cumulant of \mathbb{P}_t is integrable over (t_1, t_2) . The $2n + 2^{\text{nd}}$ cumulant of \mathbb{P}_t is a polynomial in $\{\mathbb{E}_t(\Delta Q^p)\}_{p=0}^n$ where in each monomial the product of the moments $\prod_{i=1}^k \mathbb{E}_t(\Delta Q_i^{p_i})$ is such that $\sum_{i=1}^k p_i = 2n + 2$. Using Jensen's inequality on each moment in the product,

$$\prod_{i=1}^k \mathbb{E}_t(\Delta Q_i^{p_i}) \leq \mathbb{E}_t(\Delta Q^{2n+2})^{\frac{1}{2n+2} \sum_{i=1}^k p_i} = \mathbb{E}_t(\Delta Q^{2n+2}).$$

Hence (ii) implies the $2n + 2^{\text{nd}}$ cumulant of \mathbb{P}_t is integrable over (t_1, t_2) .

Since the $2n + 2^{\text{nd}}$ cumulant of \mathbb{P}_t is given by $\int_{\mathbf{R}} e^{2n+2} d\nu_t(e)$,

$$\int_{t_1}^{t_2} \int_{\mathbf{R}} e^{2n+2} d\nu_t(e) dt < \infty.$$

By Fubini's Theorem,

$$\int_{\mathbf{R}} \left(\int_{t_1}^{t_2} (1 - \cos(et)) dt \right) e^{2n} |\rho(e)| |f(|e|)|^2 de < \infty.$$

Since $e \mapsto \int_{t_1}^{t_2} (1 - \cos(et)) dt = (t_2 - t_1) \left(1 - \frac{\sin(et_2) - \sin(et_1)}{e(t_2 - t_1)}\right)$ is lower bounded away from 0 on $\mathbf{R} \setminus [-1, 1]$,

$$\int_1^{\infty} e^{2n} \rho(e) |f(e)|^2 de < \infty.$$

Since $\rho(e) \geq 1$ for $e \geq 1$,

$$\int_1^{\infty} e^{2n} |f(e)|^2 de < \infty.$$

Finally $f \in L^2(\mathbf{R}_+)$ implies $\int_0^1 e^{2n} |f(e)|^2 de \leq \int_0^1 |f(e)|^2 de$. Hence

$$\int_{\mathbf{R}_+} e^{2n} |f(e)|^2 de < \infty. \quad \square$$

Theorem 4.16. The implication (i) \implies (ii) is obvious.

From Proposition 4.13, the characteristic function of \mathbb{P}_t is,

$$\mathcal{E}_t(\alpha) = \exp \left(\int_{\mathbf{R}} (e^{ie\alpha} - 1) d\nu_t(e) \right).$$

Assumption $f \in \text{Dom } e^{\frac{1}{2}\gamma\hat{e}} \cap \text{Dom } \hat{e}^{-1/2}$ implies $f \in \text{Dom } \hat{e}^{-1/2} e^{\frac{1}{2}\gamma\hat{e}}$. Hence, since the functions $e \mapsto |1 - \cos(et)|/e^2$ and $e \mapsto |e|/|1 - e^{-\beta e}|$ are both bounded in $e \in \mathbf{R}$, (iii) and the expression of $d\nu_t$ imply $\sup_{t \in \mathbf{R}} \int_{\mathbf{R}} e^{\gamma|e|} d\nu_t(e) < \infty$. The inequality $\int_{\mathbf{R}} e^{\gamma|e|} d\nu_t(e) \geq \int_{\mathbf{R}} e^{\pm\gamma e} d\nu_t(e)$ and the continuity of $x \mapsto e^x$ imply

$$\sup_{t \in \mathbf{R}} \mathcal{E}_t(\pm i\gamma) < \infty.$$

The inequality $e^{|x|} \leq e^x + e^{-x}$ for $x \in \mathbf{R}$ yields (iii) \implies (i).

Using the $e^{|x|} \geq \frac{1}{2}(e^x + e^{-x})$ and Jensen's inequality (ii) implies, $t \mapsto \log \mathcal{E}_t(\pm i\gamma)$ are both integrable over (t_1, t_2) . Repeating the same steps as in the proof of Theorem 4.15,

$$\int_0^{\infty} e^{\gamma e} |f(e)|^2 de < \infty,$$

and we conclude that (ii) implies (iii). \square

Remark 5.9. From these last two proofs, one can understand why, in the second statements of Theorems 4.4, 4.8, 4.15 and 4.16, the integrability over an interval (t_1, t_2) cannot be replaced by finiteness for a $t \in \mathbf{R}$.

Let

$$f_n : e \mapsto \begin{cases} \frac{1}{|e|^{n+1}}(|e| - e - i|e|^{-1})^{-1} & \text{if } e \geq 1 \\ ie & \text{if } 0 \leq e < 1. \end{cases}$$

For $n \in \mathbf{N} \setminus \{0\}$, $f_n \in \text{Dom } \hat{e}^{-1/2} \cap \hat{e}$. Integrating between two consecutive integers $N > 1$,

$$\int_{N-1}^N e^{2n} |f_n(e)|^2 de \geq \left(\frac{(N-1)}{N} \right)^{2n} \frac{\arctan N}{N}.$$

Since $\lim_{N \rightarrow \infty} \arctan N = \frac{\pi}{2}$, $\lim_{N \rightarrow \infty} \frac{N-1}{N} = 1$ and $\sum_{N=2}^{\infty} \frac{1}{N} = \infty$, $f_n \notin \text{Dom } \hat{e}^n$ but

$$\int_{N-1}^N e^{2n} (1 - \cos(e2\pi)) |f_n(e)|^2 de \leq \frac{1}{N^2} \int_0^1 \frac{1 - \cos(x2\pi)}{x^2 + N^{-2}} dx \leq \frac{2\pi^2}{N^2},$$

where the last inequality follows from $\frac{1 - \cos(2x)}{x^2} \leq 2$. Then from $\sum_{N=2}^{\infty} \frac{1}{N^2} < \infty$ and Theorem 4.13, $\mathbb{E}_{2\pi}(\Delta Q^{2n+2})$ is finite even if $f \notin \text{Dom } \hat{e}^n$.

We can similarly construct a $f_\gamma \in \text{Dom } \hat{e}^{-1/2} \cap \hat{e}$ such that $\mathbb{E}_{2\pi}(e^{\gamma|\Delta Q|})$ is finite but $f_\gamma \notin \text{Dom } e^{\frac{1}{2}\gamma\hat{e}}$.

A Self-adjointness

The following two technical lemmas ensure mathematical soundness of the construction of the perturbed dynamics $(\tau_V^t)_{t \in \mathbf{R}}$ and of the measure \mathbb{P}_t in our models where the perturbation is unbounded [DJP03]. More detailed proofs are provided in [Raq17].

Lemma A.1. *Let $L + V$ the operator defined in (26) for the quantum open harmonic oscillator of Section 4.2. Assume $f \in \text{Dom}(\hat{e}) \cap \text{Dom}(\hat{e}^{-1/2})$. Then $L + V$ is essentially self-adjoint on $\text{Dom } L \cap \text{Dom } V$.*

Proof. We apply the Nelson commutator theorem in the form of [RS75, §X.37]. Set $\hat{N} := d\Gamma(h_0 \oplus h_0 + \mathbf{1}) + \mathbf{1}$. Then it is easy to show $D = \text{Dom}(L) \cap \text{Dom}(V)$ is a core for \hat{N} . The estimates

$$\|(L + V)\Psi\| \leq c\|\hat{N}\Psi\|$$

for some $c > 0$ and all $\Psi \in D$, and

$$|((L + V)\Psi, \hat{N}\Psi) - (\hat{N}\Psi, (L + V)\Psi)| \leq d\|\hat{N}^{\frac{1}{2}}\Psi\|^2,$$

for some $d > 0$ and all $\Psi \in D$ follow from the well-known estimate

$$\left\| \prod_{i=1}^n a^\#(g_i) (d\Gamma(\mathbf{1}) + \mathbf{1})^{-\frac{n}{2}} \right\| \leq c_n \prod_{i=1}^n \|g_i\|.$$

for $g_i \in \mathfrak{h}$, provided $\|\rho^{\frac{1}{2}}\psi_f\|$, $\|(1 + \rho)^{\frac{1}{2}}\psi_f\|$, $\|h_0\rho^{\frac{1}{2}}\psi_f\|$, $\|h_0(1 + \rho)^{\frac{1}{2}}\psi_f\|$ are finite. The hypothesis $f \in \text{Dom}(\hat{e}) \cap \text{Dom}(\hat{e}^{-1/2})$ precisely guarantees that these norms are finite. \square

Lemma A.2. *Let $L + \phi^{\text{AW}}(f)$ the operator defined in (28) in the van Hove model of Section 4.3. Assume $f \in \text{Dom}(\hat{e}) \cap \text{Dom}(\hat{e}^{-1/2})$. Then $L + \phi^{\text{AW}}(f)$ is essentially self-adjoint on $\text{Dom } L \cap \text{Dom } \phi^{\text{AW}}(f)$.*

Proof. The result follows applying Nelson's commutator theorem in a similar way as in Lemma A.1. \square

B A note on the thermodynamic limit

In this appendix, we show that for the models of Section 4, \mathbb{P}_t is the weak limit of a family of probability measures associated to a two-time measurement protocol on finite dimensional systems as described in Section 3.1. We provide the proof for bosonic models; the fermionic case only requires a straightforward adaptation of the arguments.

Consider a positive semi-definite densely defined self-adjoint operator h_0 on a separable complex Hilbert space \mathfrak{h} . The extended system we wish to obtain in the thermodynamic limit is a quasi-free Bose gas with one-particle unperturbed Hamiltonian h_0 , at equilibrium at inverse temperature $\beta > 0$, together with a perturbation of the form

$$V = \frac{1}{\sqrt{2}}(a^*(g) + a(g))$$

with $g \in \text{Dom } h_0^{-1/2}$ or of the form

$$V = a^*(g_1)a(g_2) + a^*(g_2)a(g_1)$$

with $g_1, g_2 \in \text{Dom } h_0^{-1/2}$. These operators are naturally affiliated to the Weyl algebra over $\text{Dom } h_0^{1/2}$, denoted $\mathfrak{W}(\text{Dom } h_0^{-1/2})$.

We work in a representation of this gas on the Araki-Woods von Neumann algebra $\mathfrak{M}^{\text{AW}} \subset \mathcal{B}(\Gamma^+(\mathfrak{h} \oplus \mathfrak{h}))$ generated by the Weyl operators

$$W^{\text{AW}}(\phi) := W(\sqrt{\mathbf{1} + \rho(h_0)}\phi \oplus \sqrt{\rho(h_0)}\bar{\phi})$$

for $\phi \in \text{Dom } h_0^{-1/2}$ and with initial state $\langle \Omega, \cdot \Omega \rangle$ where $\Omega \in \Gamma^+(\mathfrak{h} \oplus \mathfrak{h})$ is the vacuum vector. The Liouvillean for the unperturbed dynamics in this representation is

$$L := d\Gamma(h_0 \oplus -h_0).$$

In this representation, the creation operators affiliated to $\mathfrak{W}(\text{Dom } h_0^{-1/2})$ take the form

$$a^{\text{AW}*}(\phi) = a^*(\sqrt{\mathbf{1} + \rho(h_0)}\phi \oplus 0) + a(0 \oplus \sqrt{\rho(h_0)}\bar{\phi}).$$

for any $\phi \in \text{Dom } h_0^{-1/2}$. In this manner, V corresponds to an operator V^{AW} affiliated to \mathfrak{M}^{AW} .

Let $(\mathfrak{p}_D)_{D \in \mathbf{N}}$ be a sequence of orthogonal projections on \mathfrak{h} with D -dimensional range and satisfying the following properties:

- (i) the sequence $(\mathfrak{p}_D)_{D \in \mathbf{N}}$ converges strongly to the identity i.e.

$$\lim_{D \rightarrow \infty} \|\mathfrak{p}_D \phi - \phi\| = 0$$

for all $\phi \in \mathfrak{h}$;

- (ii) for all $D \in \mathbf{N}$, $\overline{\mathfrak{p}_D \phi} = \mathfrak{p}_D \bar{\phi}$ for all $\phi \in \mathfrak{h}$;

- (iii) for all $D \in \mathbf{N}$, $\mathfrak{p}_D h_0 \mathfrak{p}_D$ is positive definite;

- (iv) there exists a core $C \subseteq \mathfrak{h}$ for h_0 such that

$$\lim_{D \rightarrow \infty} \mathfrak{p}_D h_0 \mathfrak{p}_D \phi = h_0 \phi$$

for all $\phi \in C$.

With the help of this family of projectors, we define, for each $D \in \mathbf{N}$, an unperturbed Hamiltonian $H_0^{(D)} := d\Gamma(\mathfrak{p}_D h_0 \mathfrak{p}_D)$. We also set $H_0^{(D,\delta)} := d\Gamma(p_\delta(\mathfrak{p}_D h_0 \mathfrak{p}_D))$ for the purpose of defining an initial state below, understood in the sense of functional calculus for the real function

$$p_\delta : e \mapsto \max\{\delta, e\}.$$

The parameter $\delta > 0$ serves as an IR regularization. Both $H_0^{(D,\delta)}$ and $H_0^{(D)}$ are self-adjoint operators on $\Gamma^+(\mathfrak{p}_D \mathfrak{h})$ and, since $\mathfrak{p}_D h_0 \mathfrak{p}_D > 0$, the operator $e^{-\beta H_0^{(D,\delta)}}$ is trace class.

Let $V^{(D)}$ be the same polynomial as V , but where g [resp. g_1 and g_2] is replaced by $\mathfrak{p}_D g$ [resp. $\mathfrak{p}_D g_1$ and $\mathfrak{p}_D g_2$].

We make the following definition, in accordance with an extension of Remark 3.2 from matrices to trace class operators.

Definition B.1. Let $\mathbb{P}_t^{(D,\delta)}$ be the probability measure of the heat variation as defined by the two-time measurement of the unperturbed Hamiltonian $H_0^{(D)}$ with respect to the perturbation $V^{(D)}$ and the initial state

$$\begin{aligned} \omega^{(D,\delta)} : \mathfrak{W}(\mathfrak{p}_D \mathfrak{h}) &\rightarrow \mathbf{C} \\ A &\mapsto \frac{\text{tr}(A e^{-\beta H_0^{(D,\delta)}})}{\text{tr}(e^{-\beta H_0^{(D,\delta)}})}. \end{aligned}$$

Namely, it is the probability measure with characteristic function

$$\mathcal{E}_t^{(D,\delta)}(\alpha) = \omega^{(D,\delta)}(e^{i\alpha e^{it(H_0^{(D)} + V^{(D)})} H_0^{(D)} e^{-it(H_0^{(D)} + V^{(D)})}} e^{-i\alpha H_0^{(D)}}).$$

The algebra $\mathfrak{W}(\mathfrak{p}_D \mathfrak{h})$ together with the state $\omega^{(D,\delta)}$ admit an Araki–Woods representation on $\Gamma^+(\mathfrak{h} \oplus \mathfrak{h})$: the vacuum $\Omega \in \Gamma^+(\mathfrak{h} \oplus \mathfrak{h})$ is a vector representative of $\omega^{(D,\delta)}$ and the $*$ -isomorphism

$$\pi^{\text{AW}(D,\delta)} : \mathfrak{W}(\mathfrak{p}_D \mathfrak{h}) \rightarrow \mathfrak{M}^{\text{AW}(D,\delta)}$$

is the extension by linearity of

$$\pi^{\text{AW}(D,\delta)}(W(\phi)) := W^{\text{AW}(D,\delta)}(\phi)$$

for $\phi \in \mathfrak{p}_D \mathfrak{h}$, where

$$W^{\text{AW}(D,\delta)}(\phi) := W(\sqrt{\mathbf{1} + \rho \circ p_\delta(\mathfrak{p}_D h_0 \mathfrak{p}_D)} \phi \oplus \sqrt{\rho \circ p_\delta(\mathfrak{p}_D h_0 \mathfrak{p}_D)} \bar{\phi})$$

and $\rho : \mathbf{R}_+ \ni e \mapsto (e^{\beta e} - 1)^{-1}$. For each D and δ , $\mathfrak{M}^{\text{AW}(D,\delta)}$ is the von Neumann algebra generated by $\{W^{\text{AW}(D,\delta)}(\phi) : \phi \in \mathfrak{p}_D \mathfrak{h}\}$. It is a subalgebra of the algebra $\mathfrak{W}(\mathfrak{h} \oplus \mathfrak{h})$ of operators on $\Gamma^+(\mathfrak{h} \oplus \mathfrak{h})$. The Liouvillean for the dynamics implemented by $H_0^{(D)}$ is

$$L^{(D)} := d\Gamma(\mathfrak{p}_D h_0 \mathfrak{p}_D \oplus -\mathfrak{p}_D h_0 \mathfrak{p}_D).$$

From its definition as the generator of the unitary groups $t \mapsto W(tg)$, in this representation, each creation operator $a^*(g)$ on $\Gamma^+(\mathfrak{p}_D \mathfrak{h})$ is mapped to

$$a^*(\sqrt{\mathbf{1} + \rho \circ p_\delta(\mathfrak{p}_D h_0 \mathfrak{p}_D)} g \oplus 0) + a(0 \oplus \sqrt{\rho \circ p_\delta(\mathfrak{p}_D h_0 \mathfrak{p}_D)} \bar{g})$$

on $\Gamma^+(\mathfrak{h} \oplus \mathfrak{h})$. In this manner, the perturbation $V^{(D)}$ on $\Gamma^+(\mathfrak{p}_D \mathfrak{h})$ is mapped to an unbounded operator $V^{\text{AW}(D,\delta)}$ affiliated to $\mathfrak{M}^{\text{AW}(D,\delta)}$.

The following lemma allows us to prove a sufficiently strong notion of convergence of $V^{\text{AW}(D,\delta)}$ towards V^{AW} .

Lemma B.2. *Let \mathfrak{g} be a separable Hilbert space and Ω the vacuum state of $\Gamma^+(\mathfrak{g})$. Let $\Phi \in \text{linspan}\{a^*(g_k) \cdots a^*(g_1)\Omega : k \in \mathbf{N}, g_1, \dots, g_k \in \mathfrak{g}\}$. Then, the function*

$$\mathfrak{g}^n \ni (x_1, \dots, x_n) \mapsto a^\sharp(x_n) \cdots a^\sharp(x_1)\Phi \in \Gamma^+(\mathfrak{g})$$

where each \sharp stands for $*$ or nothing, is norm continuous for \mathfrak{g}^n equipped with the product of the norm topology.

Proof. We show the continuity by induction. Let $x_1, \dots, x_n; y_1, \dots, y_n \in \mathfrak{g}$ and for every $k \in \mathbf{N}$,

$$\mathcal{C}_k := \text{linspan}\{a^*(g_l) \cdots a^*(g_1)\Omega : l \in \{0, \dots, k\}, g_1, \dots, g_l \in \mathfrak{g}\}.$$

Since $\Phi \in \cup_{k \in \mathbf{N}} \mathcal{C}_k$, there exists $n_\Phi \in \mathbf{N}$ such that $\Phi \in \mathcal{C}_{n_\Phi}$.

For $n = 1$, the inequality (48) yields there exists $K > 0$ such that

$$\|(a^\sharp(y_1) - a^\sharp(x_1))\Phi\| \leq \sqrt{n_\Phi + 1} \|y_1 - x_1\| K.$$

Hence,

$$\mathfrak{g} \ni x_1 \mapsto a^\sharp(x_1)\Phi$$

is norm continuous.

Assume now that

$$\mathfrak{g}^{n-1} \ni (x_1, \dots, x_{n-1}) \mapsto a^\sharp(x_{n-1}) \cdots a^\sharp(x_1)\Phi \in \Gamma^+(\mathfrak{g})$$

has the desired continuity property and fix $(y_1, \dots, y_{n-1}, y_n)$. Adding and subtracting,

$$(a^\sharp(y_n) \cdots a^\sharp(y_1) - a^\sharp(x_n) \cdots a^\sharp(x_1))\Phi = (a^\sharp(y_n) - a^\sharp(x_n))\Phi' + a^\sharp(x_n)\Phi'' \quad (58)$$

with Φ' an element of $\mathcal{C}_{n_\Phi+n-1}$ and

$$\Phi'' = (a^\sharp(y_{n-1}) \cdots a^\sharp(y_1) - a^\sharp(x_{n-1}) \cdots a^\sharp(x_1))\Phi.$$

Using (48), there exists $K' > 0$ independent of (x_1, \dots, x_n) such that

$$\|(a^\sharp(y_n) - a^\sharp(x_n))\Phi'\| \leq \sqrt{n_\Phi + n} \|y_n - x_n\| K'.$$

Since $d\Gamma(\mathbf{1}) + \mathbf{1}$ is bounded by $n_\Phi + n + 1$ on $\mathcal{C}_{n_\Phi+n}$, we have

$$\mathbf{1} \leq (n_\Phi + n + 1)(d\Gamma(\mathbf{1}) + \mathbf{1})^{-1}$$

on $\mathcal{C}_{n_\Phi+n}$. It follows from $\|(d\Gamma(\mathbf{1}) + \mathbf{1})^{-1/2} a^\sharp(x_n)\| \leq \|x_n\|$ that

$$\|a^\sharp(x_n)\Phi''\| \leq \sqrt{n_\Phi + n + 1} \|x_n\| \|\Phi''\|.$$

By induction hypothesis, $\lim_{\sum_{k=1}^{n-1} \|x_k - y_k\| \rightarrow 0} \|\Phi''\| = 0$.

We deduce that the left-hand side of (58) converges to 0 in norm whenever (x_1, \dots, x_n) converges in norm to (y_1, \dots, y_n) . By induction, the lemma holds for any $n \in \mathbf{N}$. \square

Proposition B.3. *Let $\mathcal{C} := \text{linspan}\{a^*(g_n) \cdots a^*(g_1)\Omega : n \in \mathbf{N}, g_1, \dots, g_n \in C\}$ with C a core for h_0 verifying (iv). The space \mathcal{C} is a common core for $L^{(D)}$, L , $V^{\text{AW}(D, \delta)}$ and V . Moreover, for all $\Phi \in \mathcal{C}$,*

$$\lim_{D \rightarrow \infty} L^{(D)}\Phi = L\Phi$$

and

$$\lim_{\delta \downarrow 0} \lim_{D \rightarrow \infty} V^{\text{AW}(D, \delta)}\Phi = V^{\text{AW}}\Phi.$$

Proof. The fact that \mathcal{C} is a common core is direct from the definitions of the operators and the fact that \mathcal{C} is a core for h_0 .

The first convergence follows from a direct computation in Fock space and (iv).

For the second limit, let $g \in \text{Dom } h_0^{-1/2}$. By Proposition VIII.25 of [RS72] assumption (iv) implies the sequence $\mathfrak{p}_D h_0 \mathfrak{p}_D$ converges in the strong resolvent sense to h_0 . Then, for any $\delta > 0$, since $e \mapsto \rho \circ p_\delta(e)$ is continuous and bounded,

$$\lim_{D \rightarrow \infty} \sqrt{\rho \circ p_\delta(\mathfrak{p}_D h_0 \mathfrak{p}_D)} \mathfrak{p}_D \bar{g} = \sqrt{\rho \circ p_\delta(h_0)} \bar{g}.$$

By Lebesgue's monotone convergence Theorem,

$$\lim_{\delta \downarrow 0} \sqrt{\rho \circ p_\delta(h_0)} \bar{g} = \sqrt{\rho(h_0)} \bar{g} = \sqrt{\rho} \bar{g}.$$

The same arguments lead to

$$\lim_{\delta \downarrow 0} \lim_{D \rightarrow \infty} \sqrt{1 + \rho \circ p_\delta(\mathfrak{p}_D h_0 \mathfrak{p}_D)} \mathfrak{p}_D g = \sqrt{1 + \rho(h_0)} g.$$

The second convergence therefore follows from Lemma B.2 with $n = 1$ or $n = 2$. \square

For the proof of the self-adjointness of the different operators, we refer the reader to Appendix A.

Proposition B.4. *With $\mathbb{P}_t^{(D,\delta)}$ as in Definition B.1 and \mathbb{P}_t the spectral measure of*

$$L + V^{\text{AW}} - e^{it(L+V^{\text{AW}})} V^{\text{AW}} e^{-it(L+V^{\text{AW}})}$$

with respect to Ω ,

$$\text{w-lim}_{\delta \downarrow 0} \text{w-lim}_{D \rightarrow \infty} \mathbb{P}_t^{(D,\delta)} = \mathbb{P}_t.$$

Proof. By extension of Remark 3.2 to the trace-class operator $e^{-\beta H_0^{(D,\delta)}}$, the characteristic function of $\mathbb{P}_t^{(D,\delta)}$ can be expressed in the standard GNS representation, which is unitarily equivalent to the Araki–Woods representation. Hence,

$$\mathcal{E}_t^{(D,\delta)}(\alpha) = \langle e^{-it(L^{(D)} + V^{\text{AW}(D,\delta)})} \Omega, e^{i\alpha L^{(D)}} e^{-it(L^{(D)} + V^{\text{AW}(D,\delta)})} \Omega \rangle$$

By Proposition B.3 above and Proposition VIII.25 of [RS72], $L^{(D)} + V^{\text{AW}(D,\delta)}$ converges in the strong resolvent sense to $L + V^{\text{AW}}$. Hence,

$$\lim_{\delta \downarrow 0} \lim_{D \rightarrow \infty} e^{-it(L^{(D)} + V^{\text{AW}(D,\delta)})} \Omega = e^{-it(L + V^{\text{AW}})} \Omega.$$

Because $e^{i\alpha L^{(D)}}$ and $e^{-it(L^{(D)} + V^{\text{AW}(D,\delta)})}$ are unitary for $\alpha \in \mathbf{R}$ and $t \in \mathbf{R}$, the strong resolvent convergence of $L^{(D)}$ and $L^{(D)} + V^{\text{AW}(D,\delta)}$ also imply through an $\epsilon/2$ -argument that

$$\lim_{\delta \downarrow 0} \lim_{D \rightarrow \infty} e^{i\alpha L^{(D)}} e^{-it(L^{(D)} + V^{\text{AW}(D,\delta)})} \Omega = e^{i\alpha L} e^{-it(L + V^{\text{AW}})} \Omega.$$

By continuity of the inner product, we have

$$\lim_{\delta \downarrow 0} \lim_{D \rightarrow \infty} \mathcal{E}_t^{(D,\delta)}(\alpha) = \mathcal{E}_t(\alpha)$$

for all $\alpha \in \mathbf{R}$ and the result follows from Lévy's continuity theorem. \square

References

- [AGMT09] David Andrieux, Pierre Gaspard, Takaaki Monnai, and Shuichi Tasaki, *The fluctuation theorem for currents in open quantum systems*, New Journal of Physics **11** (2009), no. 4, 043014.
- [Ara81] Asao Arai, *On a model of a harmonic oscillator coupled to a quantized, massless, scalar field. I*, Journal of Mathematical Physics **22** (1981), no. 11, 2539–2548.
- [AW63] Huzihiro Araki and E.J. Woods, *Representations of the canonical commutation relations describing a nonrelativistic infinite free Bose gas*, Journal of Mathematical Physics **4** (1963), no. 5, 637–662.
- [AW64] Huzihiro Araki and Walter Wyss, *Representations of canonical anticommutation relations*, Helvetica Physica Acta **37** (1964), no. 2, 136.
- [BFJP16] Tristan Benoist, Martin Fraas, Vojkan Jakšić, and Claude-Alain Pillet, *Full statistics of erasure processes: Isothermal adiabatic theory and a statistical landauer principle*, Revue Roumaine Math Pures Appl. (2016), no. 62, 259–286.
- [BJP⁺15] Tristan Benoist, Vojkan Jakšić, Annalisa Panati, Yan Pautrat, and Claude-Alain Pillet, *Full statistics of energy conservation in two-time measurement protocols*, Phys. Rev. E **92** (2015), no. 3, 032115.
- [BJP17] Tristan Benoist, Vojkan Jakšić, and Claude-Alain Pillet, *Energy statistics in open harmonic networks*, Journal of Statistical Physics **168** (2017), no. 5, 1016–1030.
- [BPP] Tristan Benoist, Annalisa Panati, and Yan Pautrat, *Heat conservation and fluctuation relations for open quantum systems*, in preparation.
- [BR87] Ola Bratteli and Derek William Robinson, *Operator Algebras and Quantum Statistical Mechanics 1*, Springer, Berlin Heidelberg, 1987.
- [BR97] ———, *Operator Algebras and Quantum Statistical Mechanics 2*, Springer, Berlin Heidelberg, 1997.
- [CBK⁺13] Michele Campisi, Ralf Blattmann, Sigmund Kohler, David Zueco, and Peter Hänggi, *Employing circuit QED to measure non-equilibrium work fluctuations*, New Journal of Physics **15** (2013), no. 10, 105028.
- [CHT11] Michele Campisi, Peter Hänggi, and Peter Talkner, *Colloquium: Quantum fluctuation relations: Foundations and applications*, Reviews of Modern Physics **83** (2011), no. 3, 771.
- [CM76] Paul R. Chernoff and Jerrold E. Marsden, *Some basic properties of infinite dimensional Hamiltonian systems*, Colloques internationaux CNRS, Géométrie symplectique et physique mathématique, no. 237, 1976, pp. 313–330.
- [Dav73] EB Davies, *The harmonic oscillator in a heat bath*, Communications in Mathematical Physics **33** (1973), no. 3, 171–186.
- [DCH⁺13] Ross Dorner, Stephen Clark, Libby Heaney, Rosario Fazio, John Goold, and Vlatko Vedral, *Extracting quantum work statistics and fluctuation theorems by single-qubit interferometry*, Phys. Rev. Lett. **110** (2013), no. 23, 230601.
- [Der03] Jan Dereziński, *Van Hove Hamiltonians — exactly solvable models of the infrared and ultraviolet problem*, Annales Henri Poincaré, vol. 4, Springer, 2003, pp. 713–738.
- [Der06] ———, *Introduction to representations of the canonical commutation and anticommutation relations*, Large Coulomb Systems (Jan Dereziński and Heinz Siedentop, eds.), Lecture Notes in Physics, vol. 695, Springer Berlin Heidelberg, 2006, pp. 63–143.
- [DG13] Jan Dereziński and Christian Gérard, *Mathematics of quantization and quantum fields*, Cambridge Monographs on Mathematical Physics, Cambridge University Press, 2013.
- [DJP03] Jan Dereziński, Vojkan Jakšić, and Claude-Alain Pillet, *Perturbation theory of W^* -dynamics, Liouvilleans and KMS-states*, Reviews in Mathematical Physics **15** (2003), no. 05, 447–489.
- [EHM09] Massimiliano Esposito, Upendra Harbola, and Shaul Mukamel, *Nonequilibrium fluctuations, fluctuation theorems, and counting statistics in quantum systems*, Reviews of modern physics **81** (2009), no. 4, 1665.

- [JKP06] Vojkan Jakšić, Eugene Kritchevski, and Claude-Alain Pillet, *Mathematical theory of the Wigner–Weisskopf atom*, Large coulomb systems (2006), 145–215.
- [JOPP11] Vojkan Jakšić, Yoshihiko Ogata, Yan Pautrat, and Claude-Alain Pillet, *Entropic fluctuations in quantum statistical mechanics an introduction*, Quantum Theory from Small to Large Scales (Jürg Frohlich, Manfred Salmhofer, Vieri Mastropietro, Wojciech De Roeck, and Leticia F Cugliandolo, eds.), Lecture Notes of the Les Houches Summer School, vol. 95, Oxford University Press, 2011, pp. 213–410.
- [JP98] Vojkan Jakšić and Claude-Alain Pillet, *Ergodic properties of classical dissipative systems I*, Acta mathematica **181** (1998), no. 2, 245–282.
- [JPPP15] Vojkan Jakšić, Jane Panangaden, Annalisa Panati, and Claude-Alain Pillet, *Energy conservation, counting statistics, and return to equilibrium*, Lett. Math. Phys. **105** (2015), no. 7, 917–938.
- [JPRB11] Vojkan Jakšić, Claude-Alain Pillet, and Luc Rey-Bellet, *Entropic fluctuations in statistical mechanics: I. Classical dynamical systems*, Nonlinearity **24** (2011), no. 3, 699.
- [JPS16] Vojkan Jakšić, Claude-Alain Pillet, and Armen Shirikyan, *Entropic fluctuations in Gaussian dynamical systems*, Reports on Mathematical Physics **77** (2016), no. 3, 335–376.
- [Kur00] Jorge Kurchan, *A quantum fluctuation theorem*, arXiv preprint cond-mat/0007360 (2000).
- [MDCP13] Laura Mazzola, Gabriele De Chiara, and Mauro Paternostro, *Measuring the characteristic function of the work distribution*, Phys. Rev. Lett. **110** (2013), no. 23, 230602.
- [Pil06] Claude-Alain Pillet, *Quantum dynamical systems*, Open Quantum Systems I (Stéphane Attal, Alain Joye, and Claude-Alain Pillet, eds.), Lecture Notes in Mathematics, vol. 1880, Springer, 2006, pp. 107–182.
- [Raq17] Renaud Raquépas, *Heat full statistics and regularity of perturbations in quantum statistical mechanics*, Master’s thesis, McGill University, Montréal, 2017.
- [RCP14] Augusto J. Roncaglia, Federico Cerisola, and Juan Pablo Paz, *Work measurement as a generalized quantum measurement*, Phys. Rev. Lett. **113** (2014), no. 25, 250601.
- [RS72] Michael Reed and Barry Simon, *Methods of Modern Mathematical Physics*, vol. I: *Functional Analysis*, Academic Press, New York, 1972.
- [RS75] ———, *Methods of Modern Mathematical Physics*, vol. II: *Fourier Analysis, Self-Adjointness*, Academic Press, New York, 1975.
- [SL77] Herbert Spohn and Joel L. Lebowitz, *Stationary non-equilibrium states of infinite harmonic systems*, Communications in Mathematical Physics **54** (1977), no. 2, 97–120.
- [Spo04] Herbert Spohn, *Dynamics of charged particles and their radiation field*, Cambridge University Press, 2004.
- [Tas00] Hal Tasaki, *Jarzynski relations for quantum systems and some applications*, arXiv preprint cond-mat/0009244 (2000).
- [Thi12] Walter Thirring, *A course in mathematical physics 1 and 2: Classical dynamical systems and classical field theory*, Springer Science & Business Media, 2012.