



HAL
open science

Allegories: decidability and graph homomorphisms

Damien Pous, Valeria Vignudelli

► **To cite this version:**

Damien Pous, Valeria Vignudelli. Allegories: decidability and graph homomorphisms. 2018. hal-01703906v1

HAL Id: hal-01703906

<https://hal.science/hal-01703906v1>

Preprint submitted on 8 Feb 2018 (v1), last revised 27 Apr 2018 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Allegories: decidability and graph homomorphisms

Damien Pous Valeria Vignudelli
Univ Lyon, CNRS, ENS de Lyon, UCB Lyon 1, LIP

Abstract

Allegories were introduced by Freyd and Scedrov; they form a fragment of Tarski’s calculus of relations. We show that their equational theory is decidable by characterising it in terms of a specific class of graph homomorphisms.

We actually do so for an extension of allegories which we prove to be conservative, namely allegories with top. This generalisation makes it possible to exploit the correspondence between terms and K_4 -free graphs, for which isomorphism was known to be finitely axiomatisable.

1 Introduction


In the nineties, Freyd and Scedrov proposed the notion of *allegory* [12], an axiomatisation of categories with some additional structure present in the category of relations (with sets as objects and binary relations as morphisms). Allegories arise in regular categories [18, Chapter A3]; they were also applied to circuit design [4]. We show in this paper that their equational theory is decidable.

Forgetting the categorical structure, allegories form a finite and purely equational axiomatisation of the *positive calculus of relations* [3]. Their syntax is the following:

$$u, v ::= a \mid u \cdot v \mid u \cap v \mid u^\circ \mid 1$$

Letter a ranges over a set of variables. The first three operations intuitively denote relational composition (\cdot), intersection (\cap) and converse (\circ). The constant 1 corresponds to the identity relation.

We can associate to each term u a labelled, directed graph $g(u)$ with two designated vertices for input and output. In this construction, a is a directed edge labelled with a , \cdot is series composition of graphs, obtained by merging the output of the first graph and the input of the second one, \cap is parallel composition, merging the inputs and the outputs of the two graphs respectively, \circ exchanges input and output, and 1 is the graph with no edge and a single vertex. For instance, the graphs of the terms $a \cdot (b \cap c^\circ) \cap d$ and $1 \cap a \cdot b$ are the following ones:



A key result about allegories is that an inequation $u \geq v$ is universally valid for binary relations (i.e., it holds for all instantiation of its variables with binary relations) if and only if there is a graph homomorphism from $g(u)$ to $g(v)$. For instance, the inequation $a \cdot b \cap a \cdot c^\circ \geq a \cdot (b \cap c^\circ) \cap d$ can be proved by exhibiting the following homomorphism:





Figure 1. Valid and invalid homomorphisms. (The edges of the graph in the middle should be labelled with different letters and oriented arbitrarily; the orientation and labelling of the edges of the outer graphs are then determined by the two homomorphisms.)

This characterisation was sketched by Freyd and Scedrov [12, page 208] and later proved by Andr eka and Bredikhin [2, Theorem 1]. It actually appeared earlier under a different and more general form, to prove that the database problem of *conjunctive queries containment* is decidable [5, Lemma 13].

This characterisation however only applies to *representable* allegories, those allegories that are isomorphic to an algebra of concrete binary relations. Indeed, the equational theory of allegories is incomplete with respect to these models. Freyd and Scedrov give a counter-example [12, p. 210]: there are homomorphisms that correspond to inequations that are not derivable from the axioms of allegories. Incompleteness also follows from a general negative result by Andr eka and Mikul as [3]: any finite first-order axiomatisation must be incomplete whenever the considered fragment contains at least the operations of composition, intersection and converse.

When looking at counter-examples to completeness, one can see that the problems always arise from homomorphisms that equate more than two vertices at a time. In fact, Freyd and Scedrov suggest that “the equations chosen as the definition of allegory happen to be precisely those that account for all containments obtainable by identifying the vertices two at a time” [12, p. 210].

We obtain decidability by proving this claim, which happens to be more difficult than expected. An attempt at exhibiting a proof was proposed in Gutierrez’s doctoral dissertation [16]. However, his proof builds on a lemma which happens to be false and cannot be fixed [1, 13]. Gutierrez also claimed decidability earlier, in a short abstract [15] where he proposes an alternative characterisation, but proofs are not available and some of his assertions seem highly non-trivial to prove. (See Appendix E). To the best of our knowledge, the problem is thus currently considered as open.

The key difficulty is that some graphs cannot be represented by a term. Thus, even though every homomorphism can be decomposed into a sequence of homomorphisms equating at most two vertices at a time, there is no guarantee that the intermediate graphs appearing in such a decomposition are the graphs of some terms. Consider for instance the graphs in Figure 1. There are homomorphisms from the outer graphs to the inner one, obtained by merging vertices depicted with the same symbol. While those three graphs are graphs of terms, we shall see that only the first one can be decomposed into a sequence where all intermediate graphs are graphs of terms, and thus corresponds to an inequality provable from allegories axioms.

We proceed in two steps.

1. First we solve the problem for *allegories with top*, that is, allegories extended with a neutral element for intersection, whose graph is the disconnect graph with no edges and two vertices, input and output. Doing so gives us more flexibility: there are more graphs that can be represented by a term (for instance, the disconnected ones), and there is a clear characterisation of the class of graphs of terms: they are precisely the graphs of treewidth at most two, or equivalently, the graphs excluding K_4 as a minor. This move also makes it possible to exploit a recent axiomatisation of isomorphism on such graphs [7]: we show that the corresponding axioms are derivable in allegories with top (Proposition 18), and we can then reason modulo isomorphism. This latter possibility is crucial in most of our proofs.
2. Then we prove that allegories with top are a conservative extension of allegories: every equation over the signature of allegories that holds in all allegories with top actually holds in all allegories. We do so using model-theoretic means, by showing how to embed any given allegory into an allegory with top (Proposition 43). We solve in passing a problem that was left open in [7]: we give a finite axiomatisation of isomorphism for connected K_4 -free graphs.

Outline and contributions We first recall the correspondence between terms and K_4 -free graphs [7] (Section 2) and setup tools allowing us to extract terms from graphs (Section 3). Then we define allegories with top and we prove important laws that are required in the sequel (Section 4). Section 5 is devoted to our main contribution: there we characterise the inequational theory of allegories in terms of sequences of appropriate homomorphisms (Theorem 15). This characterisation leads to decidability (Section 6) and to a notion of normal form (Section 7). We finally proceed with the conservativity results (Section 8), which make it possible to lift our characterisation and decidability proof to pure allegories, and to provide a finite axiomatisation of isomorphism for connected K_4 -free graphs.

The conservativity results as well as all the equational proofs needed in the paper have been formally verified using the Coq proof assistant. The development can be downloaded and browsed online [20].

2 Terms and graphs

We let $a, b \dots$ range over the *letters* of a fixed alphabet A . We consider labelled directed graphs with two designated vertices. We just call them graphs in the sequel. Note that we allow multiple edges between two vertices, as well as self-loops.

Definition 1. A *graph* is a tuple $G = \langle V, E, s, t, l, \iota, o \rangle$, where V is a finite set of *vertices*, E is a finite set of *edges*, $s, t : E \rightarrow V$ are maps indicating the *source* and *target* of each edge, $l : E \rightarrow A$ is a map indicating the *label* of each edge, and $\iota, o \in V$ are the designated vertices, respectively called *input* and *output*.

Definition 2. An *homomorphism* from $G = \langle V, E, s, t, l, \iota, o \rangle$ to $G' = \langle V', E', s', t', l', \iota', o' \rangle$ is a pair $h = \langle f, g \rangle$ of functions $f : V \rightarrow V'$ and $g : E \rightarrow E'$ that respect the various components: $s' \circ g = f \circ s$, $t' \circ g = f \circ t$, $l = l' \circ g$, $\iota' = f(\iota)$, and $o' = f(o)$.

A *surjective* (resp. *injective*) homomorphism is an homomorphism whose two components are surjective (resp. injective) functions. A (*graph*) *isomorphism* is a surjective and injective homomorphism




Figure 2. Graph operations.

whose two components are bijective functions. We write $G \simeq G'$ when there exists an isomorphism between graphs G and G' .

We consider the following signatures for terms and algebras:

$$\Sigma = \{ \cdot, \cap, \circ, \tau, \text{dom} \} \quad \Sigma_{\tau} = \Sigma \cup \{ \tau_0 \} \quad \Sigma_{\text{dom}} = \Sigma \cup \{ \text{dom}_1 \}$$

We usually omit the \cdot symbol and we assign priorities so that the term $(a \cdot (b^\circ)) \cap c$ can be written just as $ab^\circ \cap c$.

Graphs form algebras for those signatures by considering the operations depicted in Figure 2, where input and outputs are represented by unlabelled ingoing and outgoing arrows. The operations composition (\cdot) and intersection (\cap) respectively correspond to series and parallel composition, converse ($^\circ$) just exchanges input and output, and *domain* ($\text{dom}(_)$) relocates the output to the input.


By interpreting a letter $a \in A$ as the graph \underline{a} from Figure 2, one can thus associate a graph $\mathbf{g}(u)$ to every term over the considered signatures and with variables in A .

Observe that intersecting a graph with 1 amounts to merging its input and its output. As a consequence, the domain operation is derivable in the signature Σ_{τ} thanks to the isomorphism below. Intuitively, relocating the output to the input can be implemented by first disconnecting the output (by multiplication with τ on the right), and then merging it with the input (by intersection with 1).

$$\text{dom}(G) \simeq 1 \cap G\tau$$

Accordingly, we will use the following shorthand when working with Σ_{τ} -terms: $\text{dom}(u) \triangleq 1 \cap u\tau$.

There are graphs which are not the graph of any term. For instance, this is the case for the following graphs, whatever the orientation and labelling of their edges.



We now recall some standard graph theory notions, to state the characterisation of the graphs of Σ_{τ} -terms from [7].

A *simple graph* is an unlabelled undirected graph with at most one edge between two vertices and without self-loops. We use standard notation and terminology from graph theory [9]. In particular, we denote by kj a potential edge between two vertices k and j ; a kj -path is a (possibly trivial) path whose ends are k and j ; $G + kj$ is the simple graph obtained from G by adding the edge kj if k and j were not already adjacent.

Definition 3. A *minor* of a simple graph G is a simple graph obtained from G by performing a sequence of the following operations: delete an edge or a vertex, contract an edge. A simple graph is *H-free* if it does not have H as a minor.

Robertson and Seymour's graph minor theorem [21], states that (simple) graphs are well-quasi-ordered by the minor relation. As a




Figure 3. Different terms denoting the same connected graph.

consequence, the classes of graphs of bounded treewidth [9], which are closed under taking minors, can be characterised by finite sets of excluded minors. Two simple and standard instances are the following ones: the graphs of treewidth at most one (the forests) are precisely those excluding the cycle with three vertices (C_3); those of treewidth at most two are those excluding the complete graph with four vertices (K_4) [10].



Definition 4. The *skeleton* of a graph G is the simple graph S obtained from G by forgetting input, output, labelling, edge directions, edge multiplicities, and self-loops. The *strong skeleton* of G is $S + \iota$ if $\iota \neq \circ$, and S otherwise.

As an example, K_4 is the strong skeleton of all instances of the graphs in (1).

Proposition 5 ([7, Corollary 26]). *Let G be a graph. The following are equivalent.*

1. There exists a Σ_\top -term u such that $G \simeq g(u)$.
2. The strong skeleton of G has treewidth at most two.
3. The strong skeleton of G is K_4 -free.

In the sequel, we write TW_2 for the set of graphs satisfying these conditions.

The results from [7] also entail that Proposition 5 can be adapted to connected graphs just by restricting to Σ_{dom} -terms. (Σ -terms alone are not enough, consider for instance the graph $\text{dom}(\underline{a})$)

3 Parsing graphs

Many different terms can denote the same graph. First because of associativity, commutativity, and neutral elements. But also, and more importantly, because of graphs whose input and output are equal. Consider for instance the graph in Figure 3, which is the graph of the five terms given on the left. Note that the terms in the second column do not exist in the syntax of pure allegories (the signature Σ): we need either the domain operation or its encoding through the constant \top . The ability to write such terms in the syntax of allegories with \top is crucial in Section 5.

We now prove a few results that allow us to extract terms from a given graph in TW_2 . We first focus on connected graphs.

Definition 6 (Primes, tests, petals, eyes). A graph G is *prime* if it is connected and for all graphs G_1, G_2 , $G \simeq G_1 \cdot G_2$ entails $G_1 \simeq 1$ or $G_2 \simeq 1$. A graph is a *test* if its input is equal to its output. A *petal* is a prime test. An *eye* is a prime with distinct input and output.

The graphs of 1 , a , a° , $ab \cap c$, $1 \cap a$, and $1 \cap ab$ and $\text{dom}(a)$ are all prime. The graphs of ab , $a(b \cap c)$, $1 \cap a \cap bc$ and $\text{dom}((1 \cap a)(b \cap c^\circ))$ are not, the latter two being the graph of $(1 \cap a)(1 \cap bc)$. A prime is either a petal or an eye. Petals can be characterised as follows.

Lemma 7. *A test G is a petal if either*

1. $G \simeq 1$, or
2. $G \simeq 1 \cap \underline{a}$ for some letter a , or
3. G has no self-loop on its input, is connected, and remains connected when removing the input.

As expected, every connected graph can be decomposed as a series composition of primes. This can typically be depicted as follows, where eyes are green and petals are yellow. The four depicted vertices are called *checkpoints*: they must be visited by any (undirected) path from the input to the output. *Proper checkpoints* are those different from input and output.



This decomposition is not unique however: there can be superfluous occurrences of 1 , and the order in which contiguous petals appear does not matter. If the starting graph belongs to TW_2 , then so do its prime components; this allows one to proceed recursively.


The following proposition makes it possible to decompose non-trivial eyes. This is a consequence of [7, Proposition 21(i)].

Proposition 8. *Let $G \in \text{TW}_2$ be an eye. Either G consists of a single edge, or there are two connected graphs $G_1, G_2 \in \text{TW}_2$ such that $G \simeq G_1 \cap G_2$.*

We write $G[k;j]$ for the graph G with input and output respectively set to k and j . As illustrated in Figure 3, there can be several ways of extracting a term from a test. We shall mostly use the following observation, to resort to the case where input and output differ:

Observation 9. *Let G be a test and let k be a vertex of G . We have $G \simeq \text{dom}(G[\iota; k])$.*

To extract a term using this observation, one must however make sure that $G[\iota; k]$ belongs to TW_2 . When G is already known to be in TW_2 , one can use for k any neighbour of the input: $G[\iota; k]$ is necessarily in TW_2 in such a case, since its strong skeleton is the same as that of G . This is how the two terms in the second column of Figure 3 are extracted. Other options are often possible, consider for instance the following graph:



Choosing the neighbour of the input yields $\text{dom}(\text{adom}(b))$, while choosing the other vertex yields $\text{dom}(ab)$. Conversely, some options are forbidden: the topmost vertex in the graph of Figure 3 cannot be chosen, the strong skeleton of the resulting graph being K_4 .

Disconnected graphs can be parsed as follows:

Proposition 10. *Let $G \in \text{TW}_2$ be a disconnected graph.*

1. If G has a connected component H which contains neither the input nor the output, then the graph G' obtained by removing H from G belongs to TW_2 and for every vertex k in H , we have $H[k; k] \in \text{TW}_2$ and $G \simeq G' \cap \top H[k; k] \top$.
2. Otherwise, G has exactly two connected components G' and G'' respectively containing the input and the output, we have $G'[\iota; \iota], G''[o; o] \in \text{TW}_2$ and $G \simeq G'[\iota; \iota] \top G''[o; o]$.

Proof. It suffices to show that the computed graphs are in TW_2 . This follows from the observation that their strong skeleton is always a subgraph of the strong skeleton of G , and the fact that the class of K_4 -free graphs is closed under taking subgraphs. \square

$$\begin{aligned}
 u \cap u &= u & (A0) \\
 u \cap (v \cap w) &= (u \cap v) \cap w & (A1) \\
 u \cap v &= v \cap u & (A2) \\
 u \cap \top &= u & (A3) \\
 u \cdot (v \cdot w) &= (u \cdot v) \cdot w & (A4) \\
 u \cdot 1 &= u & (A5) \\
 u^\circ &= u & (A6) \\
 (u \cap v)^\circ &= u^\circ \cap v^\circ & (A7) \\
 (u \cdot v)^\circ &= v^\circ \cdot u^\circ & (A8) \\
 uv \cap uw &\geq u(v \cap w) & (AD) \\
 (v \cap wu^\circ)u &\geq vu \cap w & (AM)
 \end{aligned}$$

Figure 4. Axioms of allegories with top (All_\top).

Finally note that the occurrences of letters in a term are in one to one correspondence with the edges of its graph. As a consequence, if $C[\]$ is a term context so that $C[a]$ is a term with a designated occurrence of the letter a , then given a term u , the graph of $C[u]$ is obtained from the graph of $C[a]$ by replacing the edge corresponding to the selected occurrence of a with the graph of u .

4 Allegories

Definition 11. An *allegory with top* is a Σ_\top -algebra satisfying the axioms in Figure 4, where an inequation of the form $u \geq v$ is a shorthand for the equation $u \cap v = v$. Given Σ_\top -terms with variables in A , we write $\vdash_{All_\top} u = v$ when this equation is derivable from the axioms in Figure 4 (equivalently, when it holds in all allegories with top, for all interpretation of the variables). Similarly for inequations.

An *allegory* is a Σ -algebra satisfying those axioms but (A3). We define $\vdash_{All} u = v$ accordingly.

Axioms (A0-A8) capture the most natural properties of the operators: intersection is idempotent, associative, commutative, and has \top as a neutral element (so that the derived relation \geq is a partial order with \top as maximum element and intersection as meet); composition is associative and has neutral element 1; converse is an involution that reverses compositions. They entail $1^\circ = 1$, $\top^\circ = \top$, monotonicity of intersection and converse, and a notion of duality: every statement which holds universally also holds when reversing all compositions. We use such laws freely in the sequel.

Axiom (AD), *semi-distributivity*, is equivalent to monotonicity of composition on the right, and thus also on the left by duality, so that all operations are monotone in the end.

Axiom (AM) is called *modular identity*. This is the only unusual axiom. It generalises the notion of modularity for lattices [8]. Its dual is the following law:

$$u(v \cap u^\circ w) \geq uv \cap w \quad (AM')$$

A symmetrical consequence of modular identity is the following inequation, known as *Dedekind law*:

$$(v \cap wu^\circ)(u \cap v^\circ w) \geq vu \cap w \quad (DD)$$




Figure 5. Surjective homomorphisms for Axioms (AD) and (AM).

The axioms of allegories are sound with respect to binary relations. As a consequence of the characterisation of representable allegories mentioned in the Introduction, we have:

Proposition 12. *If $\vdash_{All_\top} u \geq v$ then there exists an homomorphism from $g(u)$ to $g(v)$.*

Axioms (A1-A8) actually all correspond to graph isomorphisms. Idempotency (A0) corresponds to an injective homomorphism from right to left, and to a surjective homomorphism from left to right; Axioms (AD) and (AM) correspond to the surjective homomorphisms depicted in Figure 5. Note that those three families of surjective homomorphisms may equate arbitrarily many vertices: the graph of u might have many vertices in addition to input and output. (This issue will be addressed in Section 5.4.)

As a consequence of Proposition 12, we have $\vdash_{All_\top} 1 \geq u$ if and only if $g(u)$ is a test. We thus call *tests* the terms satisfying this condition. Equivalently, a test is a term that is provably equal to some term of the shape $1 \cap v$. Following notations from Kleene algebra with tests (KAT) [19], we let α, β range over tests.

Recall that $\text{dom}(u) \triangleq 1 \cap u\top$ is a derived operation in the syntax of allegories with top. Such an operation is not definable in the syntax of pure allegories, but we can define a similar operation by setting $\text{dom}'(u) \triangleq 1 \cap uu^\circ$. Both domain operations are tests by definition, but the graphs of $\text{dom}(u)$ and $\text{dom}'(u)$ are not isomorphic; they are depicted below.



These operations are however interchangeable in allegories with top: we will prove the equation below after Proposition 13.

$$\vdash_{All_\top} \text{dom}(u) = \text{dom}'(u) \quad (2)$$

Proposition 13. *The following laws are derivable in allegories.*

$$\text{dom}'(u \cap v) = 1 \cap uv^\circ \quad (3)$$

$$\alpha(v \cap w) = \alpha v \cap w \quad (4)$$

$$\alpha\beta = \alpha \cap \beta = \beta\alpha \quad (5)$$

$$\alpha = \alpha^\circ = \alpha\alpha = \text{dom}'(\alpha) \quad (6)$$

$$\text{dom}'(uv) = \text{dom}'(u\text{dom}'(v)) \quad (7)$$

Proof. We prove (3) by double inclusion. $1 \cap uv^\circ \geq \text{dom}'(u \cap v)$ follows by monotonicity from $u \geq u \cap v$ and $v^\circ \geq u^\circ \cap v^\circ$. For the other direction, we use Dedekind law (DD) with $w = 1$:

$$1 \cap (u \cap v)(u^\circ \cap v^\circ) \geq 1 \cap (uv^\circ \cap 1) = 1 \cap uv^\circ$$

For Equation (4), we have

$$\begin{aligned}
 \alpha(v \cap w) &\geq \alpha(v \cap \alpha^\circ w) && (1 \geq \alpha^\circ) \\
 &\geq \alpha v \cap w && (\text{by (AM)}) \\
 &\geq \alpha v \cap \alpha w && (1 \geq \alpha) \\
 &\geq \alpha(v \cap w) && (\text{by (AD)})
 \end{aligned}$$

It suffices to prove the first equation in (5), which follows from idempotency and (4)

$$\begin{aligned}
 \alpha\beta &= \alpha(1 \cap \beta) && (\text{by (A0)}) \\
 &= \alpha 1 \cap \beta = \alpha \cap \beta && (\text{by (4)})
 \end{aligned}$$

For the first equation in (6), we have

$$\begin{aligned}
 (1 \cap u)^\circ &= 1 \cap 1u^\circ = \text{dom}'(1 \cap u) && (\text{by (3)}) \\
 &= \text{dom}'(u \cap 1) = 1 \cap u1^\circ = 1 \cap u && (\text{by (3)})
 \end{aligned}$$

The other equations in (6) follow using (5). We get (7) as follows.

$$\begin{aligned}
 \text{dom}'(u\text{dom}'(v)) &\geq \text{dom}'(u\text{dom}'(u^\circ uv \cap v)) \\
 &= \text{dom}'(u(1 \cap u^\circ uvv^\circ)) && (\text{by (3)}) \\
 &\geq \text{dom}'(u \cap uvv^\circ) && (\text{by (AM)}) \\
 &= 1 \cap uvv^\circ u^\circ = \text{dom}'(uv) && (\text{by (3)}) \\
 &\geq 1 \cap u(1 \cap vv^\circ)u^\circ \\
 &= 1 \cap u(1 \cap vv^\circ)(1 \cap vv^\circ)^\circ u^\circ && (\text{by (6)}) \\
 &= \text{dom}'(u\text{dom}'(v)) && \square
 \end{aligned}$$

The above laws also holds in allegories with top. Equation (2) follows by instantiating v with \top in (3).

5 Graph theoretical characterisation

We define a class of homomorphisms on TW_2 graphs, that we will prove to characterise inequations in All_\top . Such homomorphisms, denoted by \rightarrow , are those that can be decomposed as a sequence of homomorphisms whose source and target are both in TW_2 and equate at most two vertices. This is the case for the first homomorphism in Figure 1 (one must merge the black circles first, see Appendix A), but not for the second one: merging any two vertices with the same shape in the graph on the right yields a graph containing K_4 as a minor.)

Definition 14. Define the following relations on graphs:

- $G \succ H$ if $G, H \in \text{TW}_2$ and there is a surjective homomorphism $h : G \rightarrow H$ such that h collapses at most two vertices;
- $G \hookrightarrow H$ if there is an injective homomorphism $h : G \rightarrow H$;
- $G \rightarrow H$ if $G \succ \cup \hookrightarrow^* H$;
- $G \rightleftharpoons H$ when $G \rightarrow H$ and $H \rightarrow G$.

The class TW_2 is closed under subgraphs, so $G \hookrightarrow H$ and $H \in \text{TW}_2$ implies $G \in \text{TW}_2$. Since we require G and H to be K_4 -free when $G \succ H$, we also have that $G \rightarrow H$ and $H \in \text{TW}_2$ implies $G \in \text{TW}_2$. (Note that when $h : G \rightarrow H$ is a surjective homomorphism equating exactly two vertices, then neither $G \in \text{TW}_2$ implies $H \in \text{TW}_2$ nor the converse. See Appendix B.)

Relation \rightarrow is a preorder and \rightleftharpoons is an equivalence relation. The remainder of this section is devoted to showing that the relation \rightleftharpoons is sound and complete w.r.t. provability in All_\top :

Theorem 15. *We have $\vdash_{\text{All}_\top} u = v$ if and only if $g(u) \rightleftharpoons g(v)$.*

$$1 \cap 1 = 1 \quad (\text{A9})$$

$$\text{dom}(u \cap v) = 1 \cap u \cdot v^\circ \quad (\text{A10})$$

$$u \cdot \top = \text{dom}(u) \cdot \top \quad (\text{A11})$$

$$(1 \cap u) \cdot v = (1 \cap u) \cdot \top \cap v \quad (\text{A12})$$

Figure 6. Axioms for $2p$ -algebras (with (A1-A8)).

We start with the right-to-left implication, for which it suffices to show that $g(u) \hookrightarrow g(v)$ entails $\vdash_{\text{All}_\top} u \geq v$ (Section 5.2) and $g(u) \succ g(v)$ entails $\vdash_{\text{All}_\top} u \geq v$ (Section 5.3). For both implications, a crucial preliminary step consists in dealing with isomorphisms (Section 5.1). Then we proceed with the left-to-right implication (Section 5.4).

5.1 Isomorphisms

Definition 16 ([7, Section 3]). A $2p$ -algebra is a Σ_\top -algebra satisfying the axioms (A1)-(A8) from Figure 4 and the axioms (A9)-(A12) in Figure 6. We write $\vdash_{2p} u = v$ when two terms u and v are congruent modulo those axioms, or equivalently, when the equation holds in all $2p$ -algebras.

Note that idempotency (A0) is not included in the axioms of $2p$ -algebra. TW_2 is the free $2p$ -algebra; in particular, we have

Theorem 17 ([7, Corollary 34]). *We have $\vdash_{2p} u = v$ iff $g(u) \simeq g(v)$.*

We now observe that every allegory with top is a $2p$ -algebra.

Proposition 18. *If $\vdash_{2p} u = v$ then $\vdash_{\text{All}_\top} u = v$.*

Proof. It suffices to prove axioms (A9)-(A12) from Figure 6. Axiom (A9) is a trivial instance of idempotency. Axiom (A10) follows from (3) and (2), and Axiom (A12) from (4). For Axiom (A11) we have $u\top \geq 1 \cap u\top$ and

$$\begin{aligned}
 u\top &= 1\top \cap u\top \leq (1 \cap u\top\top^\circ)\top && (\text{by (AM)}) \\
 &= \text{dom}(u)\top && \square
 \end{aligned}$$

Corollary 19. *If $g(u) \simeq g(v)$ then $\vdash_{\text{All}_\top} u = v$.*

This result is fundamental for the following proofs: it allows us to reason up to isomorphism, and to freely choose the way we want to read a given graph. Recall for instance the five terms denoting the same graph in Figure 3; thanks to the above corollary, we know that those five terms are provably equal in allegories with top, so that we can freely replace one by the other.

5.2 Injective homomorphisms

Write $G \hookrightarrow_v H$ if there exists an injective homomorphism which is bijective on edges and adds exactly one vertex, and $G \hookrightarrow_e H$ if there exists an injective homomorphism which is bijective on vertices and adds exactly one edge. We have

$$\hookrightarrow = \hookrightarrow_v^* \hookrightarrow_e^*$$

It thus suffices to show that \hookrightarrow_v and \hookrightarrow_e yield proofs in allegories with top. We now step on the previous result about isomorphisms to do so.

Proposition 20. *If $g(u) \hookrightarrow_v g(v)$ then $\vdash_{\text{All}_\top} u \geq v$.*

Proof. Observe that $g(u) \hookrightarrow_v g(v)$ entails $g(u \cap \top \top) \simeq g(v)$, and thus $\vdash_{All_\top} u \cap \top \top = v$ by Corollary 19. We finally get

$$All_\top \vdash u = u \cap \top \geq u \cap \top \top = v . \quad \square$$

Proposition 21. *If $g(u) \hookrightarrow_e g(v)$ then $\vdash_{All_\top} u \geq v$.*

Proof. Suppose the added edge is labelled with a , and write $v = C[a]$ by selecting the corresponding occurrence of a in v . We have $g(u) \simeq g(C[\top])$, and thus $\vdash_{All_\top} u = C[\top]$ by Corollary 19. We conclude by monotonicity of all operations: $All_\top \vdash u = C[\top] \geq C[a] = v$. \square

Note that in addition to Corollary 19, we are making a crucial use of the presence of \top in the syntax in the above two proofs. We could get rid of it when working with connected graphs, but this requires convoluted arguments (for instance, we can no longer handle vertices and edges separately and in an arbitrary order).

5.3 Surjective homomorphisms

Like above for injective homomorphisms, for two graphs $G, H \in TW_2$ write $G \succ_v H$ if there exists a surjective homomorphism which is bijective on edges and equates exactly two vertices, and $G \succ_e H$ if there exists a surjective homomorphism which is bijective on vertices and equates exactly two edges. We have

$$\succ = \succ_v \overset{=}{\succ_e} \overset{*}{\succ_e}$$

We now show that \succ_v and \succ_e yield proofs in allegories with top. This is easy for the latter, but the former relation requires a much deeper analysis.

Proposition 22. *If $g(u) \succ_e g(v)$ then $\vdash_{All_\top} u \geq v$.*

Proof. Let v' be the term obtained from v by replacing the occurrence of a which is used twice in $g(v)$ by $a \cap a$. We have $\vdash_{All_\top} v = v'$ by idempotency. Now observe that $g(u) \simeq g(v')$, so that $\vdash_{All_\top} u = v'$ by Corollary 19. \square

Proposition 23. *If $g(u) \succ_v g(v)$ then $\vdash_{All_\top} u \geq v$.*

Proof. Let h be the surjective homomorphism from $g(u)$ to $g(v)$ collapsing exactly two vertices. We prove the statement by induction on $|g(u)|$, where $|G|$ is the lexicographic product of:

- the number of edges and vertices of G ,
- 1 if G is a test, and 0 otherwise.

We proceed by cases on the structure of $g(u)$.


1. $g(u)$ is a disconnected graph. Then by Proposition 10 we have two cases:
 - a. $g(u)$ has a connected component which contains neither the input nor the output. Hence, there are w and α such that $g(u) \simeq g(w \cap \top \alpha \top)$. Let k, k' be the two collapsed vertices. If k, k' are either both in $g(w)$ or both in $g(\alpha)$, we derive the result by the inductive hypothesis. Suppose $k \in g(w)$ and $k' \in g(\alpha)$. By isomorphism, we can assume that k' is the input of $g(\alpha)$. Then $g(u) \simeq g(C[1 \cap \top \alpha \top])$, where 1 corresponds to vertex k , and $g(v) \simeq g(C[1 \cap \alpha])$. The result follows from $\vdash_{All_\top} \top \alpha \top \geq \alpha$.
 - b. $g(u)$ has exactly two connected components respectively containing the input and the output, and there are α and β such that $g(u) \simeq g(\alpha \top \beta)$. Analogously to the previous case, we can derive the result by the inductive hypothesis if the collapsed vertices k, k' are either both in $g(\alpha)$ or

both in $g(\beta)$. Suppose $k \in g(\alpha)$ and $k' \in g(\beta)$. By hypothesis, collapsing k and k' gives us a graph $g(v)$ in TW_2 . Since vertex k is connected to the input of $g(\alpha)$ and k' is connected to the output of $g(\beta)$, we derive that graphs $g(\alpha)[l, k]$ and $g(\beta)[k', o]$ are in TW_2 . Hence, there exist w, x s.t. $g(\alpha)[l, k] \simeq g(w)$, $g(\beta)[k', o] \simeq g(x)$. We have $g(u) \simeq g(w \top x)$ and $g(v) \simeq g(wx)$, and the result follows by $\vdash_{All_\top} w \top x \geq wx$.

2. $g(u)$ is a connected test. Since $g(u)$ has at least two vertices, there is some vertex k adjacent to the input, but different from it. Then there is some term w such that $g(u) \simeq g(\text{dom}(w))$ and $g(u)[l; k] \simeq g(w)$. The existence of a homomorphism from $g(u)$ to $g(v)$ implies that $g(v)$ is a test as well. Moreover, by the definition of homomorphism, $h(k)$ is either adjacent to the input in $g(v)$ or the input itself (in case the vertices identified by the homomorphism are exactly k and the input of $g(v)$). In both cases, there is a term x such that $g(v) \simeq g(\text{dom}(x))$ and $g(v)[l; h(k)] \simeq g(x)$. The function h is still a surjective homomorphism from $g(w)$ to $g(x)$ collapsing two vertices, since the only difference between $g(\text{dom}(w))$ and $g(w)$ is that the output has been relocated to k , and analogously the only difference between $g(\text{dom}(x))$ and $g(x)$ is that the output has been relocated to $h(k)$. The graph $g(w)$ has the same number of vertices and edges as $g(u)$, but has input different from output, so we can apply the inductive hypothesis to $g(w)$ and derive $\vdash_{All_\top} w \geq x$. Hence, $\vdash_{All_\top} \text{dom}(w) \geq \text{dom}(x)$.
3. $g(u)$ is a connected graph with input different from output, and is not prime, i.e., there are u_1, u_2 both not equivalent to 1 such that $g(u) = g(u_1) \cdot g(u_2)$. Let k, k' be the vertices merged by the homomorphism. If either k, k' are both in $g(u_1)$ or they are both in $g(u_2)$ (possibly including the case when one of k, k' is the checkpoint between $g(u_1)$ and $g(u_2)$ in $g(u)$), we can apply the inductive hypothesis. Otherwise, k is in $g(u_1)$ and k' is in $g(u_2)$, and neither of them is the checkpoint between $g(u_1)$ and $g(u_2)$. As discussed in Section 3, $g(u)$ can be decomposed as a sequence of prime components $g(u_1) \cdot \dots \cdot g(u_n)$, where each component is either a petal or an eye. Without loss of generality, we can assume that k is in the first prime component, and that k' is in the last prime component (modulo the presence of petals equivalent to 1). If it were not the case, the inductive hypothesis could be applied as above. We consider different cases depending on whether k, k' are in a petal or in an eye.
 - a. Both k and k' are in petals. Since k and k' can be merged, and since k' is connected to the output, the output of the first petal can be relocated to k . Analogously, the input of the last petal can be relocated to k' . Then there are w, x, z s.t. $g(u) \simeq g(w \top x \cap z)$, as represented below, with k the output of $g(w)$ and k' the input of $g(x)$, and $g(v) \simeq g(wx \cap z)$. We conclude by $\vdash_{All_\top} \top \geq 1$.



- b. Both k and k' are in an eye. Then (the strong skeleton of $g(u)$ has the following graph as a minor (the lower edge being the one added by the definition of strong skeleton).



We thus get K_4 by collapsing k and k' , which contradicts the fact that $g(v)$ is the graph of a term.


- c. k is in a petal and k' is in an eye. We consider two cases:
- i. Suppose the petal and the eye are contiguous, i.e., that there are no prime components between them (modulo the presence of 1). Then the petal is not equivalent to 1, otherwise k and k' would be in the same component. We have $g(u) \simeq g(\alpha w \cap x)$, with $k \in g(\alpha)$ and $g(w \cap x)$ the eye containing k' , with k' strictly inside $g(w)$ and $g(x)$ containing at least one edge (this decomposition of an eye always exists by Proposition 8).



Then we have $g(v) \simeq g(z \cap x)$ with $g(\alpha w) \succ_v g(z)$ and we conclude by the inductive hypothesis.


- ii. Suppose the petal and the eye are not contiguous. There are only petals (of which at least one not equivalent to 1) between them, we can move them by isomorphism before the petal containing k , and apply the inductive hypothesis. Hence, suppose there is at least one eye between the first and last prime component of $g(u)$. We consider different cases depending on where k and k' are respectively located in the petal and in the eye.

- A. If k does not coincide with the input, then $g(u)$ has as a minor



and by collapsing k and k' we would have that $g(v)$ has K_4 as a minor, which is a contradiction.


- B. k is the input and the eye containing k' has two parallel components w, z , for $k' \in g(w)$ and $g(w)$ with at least one proper checkpoint. We have three cases:



- k' is a proper checkpoint of $g(w)$. Then $g(u) \simeq g(x(w_1 w_2 \cap z))$ with k the input of x and with k' the output of w_1 , for $g(w) \simeq g(w_1 w_2)$. By collapsing k, k' we obtain $g(v) \simeq g((x \cap w_1^\circ)z \cap w_2)$ and we derive in All_\top :

$$\begin{aligned} & x(w_1 w_2 \cap z) \\ & \geq (x \cap w_1^\circ)((x \cap w_1^\circ)^\circ w_2 \cap z) \\ & \geq (x \cap w_1^\circ)z \cap w_2 \quad (\text{by (AM')}) \end{aligned}$$

- k' is strictly in a petal of $g(w)$ with input a proper checkpoint of $g(w)$. Then $g(u)$ has as a minor




and by collapsing k and k' we would have that $g(v)$ has K_4 as a minor.

- k' is strictly in an eye of $g(w)$ having as output or input a proper checkpoint of $g(w)$. Then $g(u)$ has as a minor one of the following graphs:



and, as in the previous case, by collapsing k and k' we obtain K_4 .

4. $g(u)$ is an eye not reduced to an edge. Then $g(u) \simeq g(u_1) \cap g(u_2)$ with both $g(u_1)$ and $g(u_2)$ containing at least one edge. If k, k' are in the same parallel component $g(u_i)$, then we can apply the inductive hypothesis. Otherwise, suppose that they are respectively in $g(u_1)$ and $g(u_2)$. We can assume that both $g(u_1)$ and $g(u_2)$ have at least one proper checkpoint, since otherwise there would be a way to decompose $g(u)$ into parallel components such that k, k' are in the same one. We have three cases:



- k, k' are respectively proper checkpoints of $g(u_1)$ and $g(u_2)$. Then there are w, w', x, x' s.t. $g(u) \simeq g(w w' \cap x x')$ and $g(v) \simeq g((w \cap x)(w' \cap x'))$, and we use Axiom (AD).
- k is strictly in the petal of a proper checkpoint of $g(u)$ (or symmetrically for k'). Then $g(u)$ has as a minor



and by collapsing k and k' we obtain K_4 .

- k is strictly in an eye of $g(u_1)$ having as output or input a proper checkpoint of $g(u_1)$ (or symmetrically for k'). Then $g(u)$ has as a minor one of the following graphs



and, as above, by collapsing k and k' we obtain K_4 .

5. It remains to consider the case when $g(u)$ is an eye reduced to an edge. Then $g(u) \simeq g(a)$, $g(v) \simeq g(a \cap 1)$ and the result follows by $\vdash_{All_\top} u \geq u \cap 1$. \square

Combining Corollary 19 and Propositions 20, 21, 22, and 23, we obtain that the relation \rightarrow is sound for allegories with top:

Theorem 24. *If $g(u) \rightarrow g(v)$ then $\vdash_{All_\top} u \geq v$.*

5.4 Completeness

To prove the converse of Theorem 24, we introduce an intermediate inequational presentation of allegories with top. This idea was already sketched in [12]; we give some details here for the sake of completeness.

The inequational theory All_\top^\geq is generated by the axioms (A1)-(A8) in Figure 4, where $u = v$ is a derived operator defined as $u \leq v$ and $v \leq u$, and the axioms in Figure 7. The point of this system is that its axioms all correspond to simple graph homomorphisms, that can easily be seen to belong to the relation \rightarrow :

Lemma 25. *For all axioms $u \geq v$ of All_\top^\geq , we have $g(u) \rightarrow g(v)$.*

Proof. Axioms (A1)-(A8) from Figure 4 as well as idempotency for 1 correspond to graph isomorphisms. Inequality axioms correspond to injective homomorphisms, unless v is a test and u is not. In such a case, e.g., for left inequality, we have $g(u) \succ_v g(u) \cap 1 \leftrightarrow g(u \cap v)$.

$$\begin{array}{ll}
 u \geq u \cap v & \text{(left inequality)} \\
 u \geq v \cap u & \text{(right inequality)} \\
 1 \cap 1 \geq 1 & \text{(1 idempotency)} \\
 a \cap a \geq a & \text{(atomic idempotency)} \\
 u \cdot v \cap x \cdot w \geq (u \cap x) \cdot (v \cap w) & \text{(separated semi-distributivity)} \\
 (u \cap w \cdot x^\circ) \cdot v \geq u \cdot (v \cap x) \cap w & \text{(separated modularity)}
 \end{array}$$

Figure 7. Inequational presentation of allegories with top (All_{\top}^{\geq}).

Atomic idempotency corresponds to the relation \succ_e . Finally, separated semi-distributivity and separated modularity correspond either to \succ_v or to \simeq . For instance, for separated semi-distributivity we have an isomorphism if both u and x are tests or if both v and w are tests, and we have \succ_v otherwise. \square

Note that unrestricted idempotency, semi-distributivity and modular identity (Axioms (A0), (AD) and (AM) in Figure 4) could not be handled in such a way since they correspond to homomorphisms potentially equating many vertices and edges. That they are nevertheless captured by the relation \rightarrow is obtained only a posteriori.

By further showing that \rightarrow is ‘closed under contexts’, we deduce that the system All_{\top}^{\geq} is sound for \rightarrow .

Lemma 26. *For all term-contexts C , we have*

1. *if $g(u) \succ g(v)$ then $g(C[u]) \succ g(C[v])$;*
2. *if $g(u) \hookrightarrow g(v)$ then $g(C[u]) \hookrightarrow g(C[v])$.*

Proof. By induction on C . Note that in case $C = [\cdot] \cap w$ for the first item, if the merged vertices are the input and output of u and if w has input equal output then we have $g(C[u]) \simeq g(C[v])$, which is a special case of \succ . \square

Theorem 27. *If $\vdash_{All_{\top}^{\geq}} u \geq v$ then $g(u) \rightarrow g(v)$.*

It remains to show that the inequational presentation All_{\top}^{\geq} is complete for allegories with top. We start by proving that (unrestricted) idempotency is derivable in All_{\top}^{\geq} :

Lemma 28. *For all terms u , we have $\vdash_{All_{\top}^{\geq}} u \cap u = u$.*

Proof. We prove $\vdash_{All_{\top}^{\geq}} u \cap u \geq u$ by induction on u . (The converse inequality trivially holds by the inequality axioms). The base cases are given by the axioms of All_{\top}^{\geq} . For series composition we have $\vdash_{All_{\top}^{\geq}} uv \cap uv \geq (u \cap u)(v \cap v)$ by separated semi-distributivity, and then we conclude by the inductive hypothesis. \square

Theorem 29. *If $\vdash_{All_{\top}} u \geq v$ then $\vdash_{All_{\top}^{\geq}} u \geq v$*

Proof. It suffices to derive modularity and semi-distributivity from their separated versions; this follows from Lemma 28. \square

Combining Theorems 24, 27, and 29 we finally obtain our main result (Theorem 15).

6 Decidability

We now show that the relation \rightarrow is decidable, and that there is a notion of normal form for allegories with top. The key observation is that surjective homomorphisms can always be applied first.

Lemma 30. *We have the following inclusion: $\hookrightarrow \succ \subseteq \rightarrow \hookrightarrow$.*

Proof. Let $i : G \hookrightarrow G'$ and $h : G' \succ H$. The graph $hi(G)$ is K_4 -free, being a subgraph of H , and the function $g : G \rightarrow hi(G)$, defined as hi , is a surjective homomorphism collapsing at most two vertices. Indeed, function g is trivially surjective, and either the vertices collapsed by h are both in $i(G)$, in which case g collapses them, or at least one of the collapsed vertices is not in $i(G)$, which implies that g is an isomorphism. Therefore, we have $G \succ hi(G)$. Since $hi(G)$ injects in H , we conclude that $G \succ \hookrightarrow H$. \square

As a consequence, we obtain the following characterisation, which gives decidability:

Proposition 31. *We have $G \rightarrow H$ iff $G \succ^* \hookrightarrow H$.*

Corollary 32. *The relation \rightarrow is decidable.*

Proof. It suffices to decide whether $G \succ^* \hookrightarrow H$. We have that the sets $\{G' \mid G \succ^* G'\}$ and $\{H' \mid H' \hookrightarrow H\}$ are finite and computable (up to isomorphism); it suffices to test whether they intersect. \square

One can actually get a non-deterministic polynomial algorithm: guess a sequence G_0, \dots, G_n of graphs obtained from $G = G_0$ by merging two vertices at a time, check that these graphs belong to TW_2 , compute the graph H' obtained from G_n by merging all parallel edges with the same label (so that $G_n \succ_e^* H'$) and check that $H' \hookrightarrow H$. The latter test can be done in polynomial time once G_n and thus H' are known to have bounded treewidth [6, 11, 14].

Corollary 33. *The equational theory of All_{\top} is in NP.*

7 Normal forms

When studying homomorphism equivalence on graphs, one often uses the notion of *core*, those graphs where every endomorphism is an isomorphism. Every graph has a core, which is a minimal graph in its equivalence class modulo homomorphism equivalence [17]. One defines a similar notion here for allegories, using our restricted form of homomorphism equivalence (\rightleftharpoons).

The *normal form* of a graph G , written $\text{nf}(G)$ is a graph which is minimal with respect to the number of vertices and edges in its equivalence class modulo \rightleftharpoons . The normal form of a graph is unique modulo isomorphism:

Proposition 34. *We have $G \rightleftharpoons H$ iff $\text{nf}(G) \simeq \text{nf}(H)$.*

Define the following (computable) relation:

$$G \rightsquigarrow H \triangleq G \succ^* H \hookrightarrow G$$

Proposition 35. *The relation \rightsquigarrow is a confluent and terminating preorder and for all graphs G we have $G \rightsquigarrow \text{nf}(G)$.*

Proof. That \rightsquigarrow is a preorder follows from \succ^* and \hookrightarrow being preorders. Termination follows from the observation that the size of a graph decreases along \succ (one must of course forbid trivial steps). Confluence is proved by induction on the sum of the sizes of the considered graphs, using Proposition 31 (see Appendix C). \square

Gutierrez proves a similar result in the context of finite categories with an epi-mono factorisation system [16, Chapter 4.2]. One cannot reuse his result directly: the injective (resp. surjective) homomorphisms we use here are not exactly the mono (resp. epi) morphisms of the natural category associated to the relation \rightarrow .

To illustrate this rewriting system, consider the four graphs depicted below. We highlight pairs of vertices that can potentially

$$1 \cap 1 = 1 \quad (\text{A9})$$


$$\text{dom}(u \cap v) = 1 \cap u \cdot v^\circ \quad (\text{A10})$$

$$\text{dom}(u \cdot v) = \text{dom}(u \cdot \text{dom}(v)) \quad (\text{A13})$$


$$\text{dom}(u) \cdot (v \cap w) = \text{dom}(u) \cdot v \cap w \quad (\text{A14})$$

Figure 8. Axioms for $2p\text{dom}$ -algebras (with (A1,A2,A4-A8)).

be merged by representing them with the same symbol, using the same convention as in Figure 1 for labels and orientation of edges. G_4 is the normal form of the four graphs, and there are three ways of reaching it from G_1 : directly, or by going through G_2 or G_3 .



In this example, every attempt to collapse two vertices results in a graph in TW_2 . This is not always the case, as shown by the following example.



H_3 is the normal form of those three graphs. The unique homomorphism h from H_1 to H_3 can be factorised through the graph $H_2 \in \text{TW}_2$ obtained by first merging the vertices depicted with blue squares. Since $H_3 \hookrightarrow H_1$, we deduce $H_1 \rightsquigarrow H_3$. We also have $H_2 \rightsquigarrow H_3$, but not $H_1 \rightsquigarrow H_2$, since H_2 does not embed in H_1 . If instead we try to collapse first the vertices depicted with black circles in H_1 , we obtain a graph that does not belong to TW_2 .

8 Conservativity arguments

We now show that the results presented in the previous sections extend to pure allegories (without top). We do so by proving that All_\top is a conservative extension of All , i.e., that for all terms u, v in the syntax of allegories (i.e., for all \top -free terms), $\vdash_{All_\top} u = v$ if and only if $\vdash_{All} u = v$. Consequently, the equational theory of allegories is decidable and Theorem 15 also holds for All .

We prove conservativity by showing that every allegory embeds into an allegory with top (Proposition 43). It happens that we can factorise this construction to show in passing how to handle isomorphism of connected K_4 -free graphs. This is what we do first.

Proofs in this section are mostly equational and often involve many cases. We give some sketches to give intuitions; more details can be found in Appendix D; full proofs formalised in Coq can be browsed online [20].

8.1 Isomorphism of connected graphs in TW_2

As explained in [7], connected K_4 -free graphs correspond to terms over the signature Σ_{dom} , where \top is no longer present, and $\text{dom}(_)$ becomes a primitive operation. It was however left open whether isomorphism of such graphs could be finitely axiomatised over this syntax. We answer this question by the affirmative.

Definition 36. A $2p\text{dom}$ -algebra is a Σ_{dom} -algebra satisfying the axioms (A1-A8) from Figure 4 except (A3), and the axioms in Figure 8. We write $\vdash_{2p\text{dom}} u = v$ when two terms u and v are congruent modulo those axioms, or equivalently, when the equation holds in all $2p\text{dom}$ -algebras.

Graphs form a $2p\text{dom}$ -algebra: the axioms are sound. To prove that they are complete for graph isomorphism, we rely on Theorem 17 and we prove that every $2p\text{dom}$ algebra can be embedded in a $2p$ algebra. Combined with the other results from [7], this also yields that connected K_4 -free graphs form the free $2p\text{dom}$ -algebra.

We fix a Σ_{dom} -algebra $(X, \cdot, \cap, _^\circ, \text{dom}(_), 1)$ in the remainder of this section. We write T for the set of tests in X . We construct the following Σ_\top -algebra:

Definition 37. Let \bar{X} be the set $X \uplus T^2$. For $u \in X$, we write \bar{u} for u seen as an element of \bar{X} . For $\alpha, \beta \in T$, we write $\alpha \diamond \beta$ for the pair (α, β) seen as an element in \bar{X} .

We turn \bar{X} into a Σ_\top -algebra by setting

$$\begin{aligned} \bar{u} \cdot \bar{v} &\triangleq \overline{u \cdot v} & \bar{u} \cap \bar{v} &\triangleq \overline{u \cap v} \\ (\alpha \diamond \beta) \cdot (\gamma \diamond \delta) &\triangleq \alpha \diamond \delta & (\alpha \diamond \beta) \cap (\gamma \diamond \delta) &\triangleq \alpha \gamma \diamond \beta \delta \\ (\alpha \diamond \beta) \cdot \bar{v} &\triangleq \alpha \diamond \text{dom}(v^\circ \beta) & (\alpha \diamond \beta) \cap \bar{v} &\triangleq \overline{\alpha v \beta} \\ \bar{u} \cdot (\gamma \diamond \delta) &\triangleq \text{dom}(u \gamma) \diamond \delta & \bar{u} \cap (\gamma \diamond \delta) &\triangleq \overline{\gamma u \delta} \\ (\alpha \diamond \beta)^\circ &\triangleq \beta \diamond \alpha & 1 &\triangleq \bar{1} \\ \bar{u}^\circ &\triangleq \overline{u^\circ} & \top &\triangleq 1 \diamond 1 \end{aligned}$$

When X is the algebra of connected graphs, \bar{X} intuitively represent graphs where all vertices are connected either to the input or to the output: an element \bar{u} denotes a connected graph, while an element $\alpha \diamond \beta$ denotes the disconnected graph $\alpha \top \beta$.

When composing two ‘disconnected elements’ $\alpha \diamond \beta$ and $\gamma \diamond \delta$ in series, we throw away a component that should intuitively be created and which is not connected to the input or to the output: $\beta \gamma$. This means that \bar{X} cannot be the free $2p$ -algebra: whatever the starting $2p\text{dom}$ -algebra X , \bar{X} always satisfies the law $\top u \top = \top$.

Note however that the function mapping an element $u \in X$ to \bar{u} is an injective Σ -homomorphism from X to \bar{X} .

We have the following characterisation of tests:

Lemma 38. In $2p\text{dom}$, an element u is a test iff $\text{dom}(u) = u$.

Proposition 39. If X is a $2p\text{dom}$ -algebra then \bar{X} is a $2p$ -algebra.

Proof. We must show that \bar{X} satisfies all the $2p$ axioms (Figure 6). Consider the associativity of product (A4). Each of the three variables occurring in this axiom can be either in X or in T^2 . The proof is trivial when all elements are in X . We show two relevant cases.

$$\begin{aligned} ((\alpha \diamond \beta) \cdot \bar{u}) \cdot (\gamma \diamond \delta) &= (\alpha \diamond \text{dom}(u^\circ \beta)) \cdot (\gamma \diamond \delta) \\ &= \alpha \diamond \delta \end{aligned}$$

$$(\alpha \diamond \beta) \cdot (\bar{u} \cdot (\gamma \diamond \delta)) = (\alpha \diamond \beta) \cdot (\text{dom}(u \gamma) \diamond \delta)$$

$$\begin{aligned} ((\alpha \diamond \beta) \cdot \bar{u}) \cdot \bar{v} &= (\alpha \diamond \text{dom}(u^\circ \beta)) \cdot \bar{v} \\ &= \alpha \diamond \text{dom}(v^\circ \text{dom}(u^\circ \beta)) \\ &= \alpha \diamond \text{dom}(v^\circ u^\circ \beta) && \text{(by (A13))} \end{aligned}$$

$$(\alpha \diamond \beta) \cdot (\bar{u} \cdot \bar{v}) = (\alpha \diamond \beta) \cdot \overline{u \cdot v} \quad \square$$

Corollary 40. For all Σ_{dom} -terms u, v we have

$$\vdash_{2\text{pdom}} u = v \quad \text{iff} \quad \vdash_{2\text{p}} u = v \quad \text{iff} \quad g(u) \simeq g(v) .$$

8.2 Pure allegories

We now prove that All_{\top} is a conservative extension of All , using the same construction as in Definition 37.

A difficulty here is that we need a domain operation in order to use this construction, and this operation is not in the syntax of allegories, nor derivable due to the absence of \top . Fortunately, we can use the alternative operation $\text{dom}'(\cdot)$ (defined by $\text{dom}'(u) \triangleq 1 \cap uu^\circ$). Indeed, these terms are equivalent in allegories with \top (Equation (2) from Section 4).

The following lemma makes it possible to reuse Proposition 39 in the proof of Proposition 43 below.

Lemma 41. If $\langle X, \cdot, \cap, _^\circ, 1 \rangle$ is an allegory, then $\langle X, \cdot, \cap, _^\circ, \text{dom}'(\cdot), 1 \rangle$ is a 2pdom-algebra.

Proof. It suffices to show that the axioms in Figure 8 are derivable in All . (A9) is an instance of idempotency; (A10) and (A13) have been proved in Proposition 13; (A14) follows from law (4) in Proposition 13 since $\text{dom}'(u)$ is by definition a test in All . \square

Lemma 42. Let X be an allegory; the partial order derived on \bar{X} is characterised in terms of the partial order on X as follows.

$$\begin{aligned} \bar{u} \geq \bar{v} & \quad \text{iff} \quad u \geq v \\ \alpha \diamond \beta \geq \gamma \diamond \delta & \quad \text{iff} \quad \alpha \geq \gamma \text{ and } \beta \geq \delta \\ \alpha \diamond \beta \geq \bar{v} & \quad \text{iff} \quad \alpha v \beta \geq v \\ \bar{u} \geq \gamma \diamond \delta & \quad \text{never holds} \end{aligned}$$

Proposition 43. If X is an allegory then \bar{X} is an allegory with top.

Proof. By Lemma 41 and Proposition 39, it suffices to prove idempotency, semi-distributivity, and modularity. We show two interesting cases for modularity.

- $((\alpha \diamond \beta) \cap \bar{w} \bar{v}^\circ) \bar{v} \geq (\alpha \diamond \beta) \bar{v} \cap \bar{w}$: we have

$$\begin{aligned} ((\alpha \diamond \beta) \cap \bar{w} \bar{v}^\circ) \bar{v} &= \overline{\alpha w \beta v} \\ (\alpha \diamond \beta) \bar{v} \cap \bar{w} &= \overline{\alpha w \text{dom}'(v^\circ \beta)} \end{aligned}$$

and then

$$\begin{aligned} \alpha w v^\circ \beta v &\geq \alpha w (1 \cap v^\circ \beta v) = \alpha w (1 \cap v^\circ \beta^\circ \beta v) & \text{(by (6))} \\ &= \alpha w (\text{dom}'(v^\circ \beta)) \end{aligned}$$

- $(\bar{u} \cap \bar{w}(\alpha \diamond \beta)^\circ)(\alpha \diamond \beta) \geq \bar{u}(\alpha \diamond \beta) \cap \bar{w}$: we have

$$\begin{aligned} (\bar{u} \cap \bar{w}(\alpha \diamond \beta)^\circ)(\alpha \diamond \beta) &= \text{dom}'(\text{dom}'(w\beta^\circ)u\alpha^\circ\alpha) \diamond \beta \\ \bar{u}(\alpha \diamond \beta) \cap \bar{w} &= \overline{\text{dom}'(u\alpha)w\beta} \end{aligned}$$

We use Lemma 42 and prove

$$\begin{aligned} &\text{dom}'(\text{dom}'(w\beta^\circ)u\alpha^\circ\alpha)\text{dom}'(u\alpha)w\beta\beta \\ &= \text{dom}'(u\alpha)\text{dom}'(w\beta)w\beta & \text{(by (7), (5), (6))} \\ &\geq \text{dom}'(u\alpha)w\beta \end{aligned}$$

where the last step follows by $\vdash_{All} \text{dom}'(u)u \geq u$:

$$\text{dom}'(u)u = (1 \cap uu^\circ)u \geq 1u \cap u = u \quad \text{(by (AM))}$$

\square

Corollary 44. For all Σ -terms u, v we have $\vdash_{All} u = v$ if and only if $\vdash_{All_{\top}} u = v$.

Decidability of allegories follows, as well as the expected graph-theoretical characterisation:

Corollary 45. We have $\vdash_{All} u = v$ if and only if $g(u) \simeq g(v)$.

9 Future work

We proved decidability (in NP) of the equational theories of allegories and allegories with top, and we designed a graph rewriting system making it possible to compute normal forms. The precise complexity of these equational theories remains open.

There is a simple polynomial algorithm to find whether there is an (arbitrary) homomorphism from $g(u)$ to $g(v)$: fix $G = g(v)$ and consider the following formula over subterms w of u and pairs j, k of vertices in G :

$$\phi(w, j, k) \triangleq \text{there is an homomorphism from } g(w) \text{ to } G[j, k]$$


The truth table of this formula can be computed recursively for all arguments, and this can be done in polynomial time by dynamic programming. It is not clear whether this algorithm could be adapted to compute the restricted relation \rightarrow efficiently.

References


- [1] B. Aameri and M. Winter. A first-order calculus for allegories. In H. de Swart, editor, *RAMiCS*, pages 74–91, Berlin, Heidelberg, 2011. Springer.
- [2] H. Andr eka and D. A. Bredikhin. The equational theory of union-free algebras of relations. *Algebra Universalis*, 33(4):516–532, 1995.
- [3] H. Andr eka and S. Mikul as. Axiomatizability of positive algebras of binary relations. *Algebra Universalis*, 66(1):7–34, 2011.
- [4] C. Brown and G. Hutton. Categories, allegories, and circuit design. In *Proc. LICS*. IEEE, 1994.
- [5] A. K. Chandra and P. M. Merlin. Optimal implementation of conjunctive queries in relational data bases. In *Proc. STOC*, pages 77–90. ACM, 1977.
- [6] C. Chekuri and A. Rajaraman. Conjunctive query containment revisited. *Theoretical Computer Science*, 239(2):211–229, 2000.
- [7] E. Cosme-Ll opez and D. Pous. K4-free graphs as a free algebra. In *Proc. MFCS*, volume 83 of *LIPICs*, pages 76:1–76:14. Schloss Dagstuhl, 2017.
- [8] B. Davey and H. Priestley. *Introduction to Lattices and Order*. Cambridge University Press, 2nd (2002) edition, 1990.
- [9] R. Diestel. *Graph Theory*. Graduate Texts in Mathematics. Springer, 2005.
- [10] R. Duffin. Topology of series-parallel networks. *Journal of Mathematical Analysis and Applications*, 10(2):303–318, 1965.
- [11] E. C. Freuder. Complexity of k-tree structured constraint satisfaction problems. In *Proc. NCAI*, pages 4–9. AAAI Press / The MIT Press, 1990.
- [12] P. Freyd and A. Scedrov. *Categories, Allegories*. North Holland. Elsevier, 1990.
- [13] J. Glanfield. Towards automated derivation in the theory of allegories. MSc. Thesis, Brock University, 2008.
- [14] M. Grohe. The complexity of homomorphism and constraint satisfaction problems seen from the other side. *J. ACM*, 54(1):1:1–1:24, Mar. 2007.
- [15] C. Gutierrez. Decidability of the equational theory of allegories. In *Proc. RelMiCS*, 1998.
- [16] C. Gutierrez. *The Arithmetic and Geometry of Allegories: Normal Forms and Complexity of a Fragment of the Theory of Relations*. PhD thesis, Middletown, CT, USA, 1999. AAI9934518.
- [17] P. Hell and J. Ne et il. The core of a graph. *Discrete Mathematics*, 109(1):117–126, 1992.
- [18] P. T. Johnstone. *Sketches of an elephant: a Topos theory compendium*. Oxford logic guides. Oxford Univ. Press, New York, NY, 2002.
- [19] D. Kozen. Kleene algebra with tests. *Transactions on Programming Languages and Systems*, 19(3):427–443, May 1997.
- [20] D. Pous and V. Vignudelli. Web appendix to this paper, including Coq proofs of all equational proofs and conservativity results, 2018. <http://perso.ens-lyon.fr/damien.pous/allegories/>.
- [21] N. Robertson and P. Seymour. Graph minors. XX. Wagner’s conjecture. *Journal of Combinatorial Theory, Series B*, 92(2):325 – 357, 2004.

A Example in Figure 1


The homomorphism from the first to the second graph in Figure 1 corresponds to an inequality provable in *All*, i.e., it is of the form \rightarrow . We show that indeed the homomorphism can be decomposed as a sequence of homomorphisms on graphs in TW_2 merging two vertices at a time. Among the vertices having the same shape and colour in the first graph




the only two vertices that we can collapse are the ones depicted with a black circle. Otherwise, we would obtain a graph whose strong skeleton has K_4 as a minor. If we do so, we obtain




Then we can merge in sequence the following pairs of vertices: the two green squares on the left, the two blue squares on the left, the two green squares on the right, the two blue squares on the right. At each step, the reached graph is a subgraph of the previous one, and thus is in TW_2 (it actually does not matter the order in which we merge these pairs). After applying these four homomorphisms, we have



We can now collapse the two remaining green squares and obtain a graph in TW_2 (symmetrically, we could have first collapsed the blue squares)




Finally, we collapse the green squares and obtain the target graph




This concludes the proof that $G_1 \rightarrow G_2$.

By contrast, the third graph in Figure 1




has an homomorphism to G_2 , but the graphs are not related by \rightarrow . Since $\rightarrow = \succ^* \hookrightarrow$ (Proposition 31), in order to prove $G_3 \not\rightarrow G_2$ it suffices to note that whenever we collapse two nodes with the same shape in G_3 we obtain a graph not in TW_2 ; the following one for instance.



We conclude that $G_3 \not\rightarrow G_2$: there is no injection from G_3 to G_2 .


B Leaving/entering TW_2 by merging vertices

Let G be the graph depicted below, and let G' be the graph obtained by collapsing the nodes connected by the dotted line.



The function $g : G \rightarrow G'$ that assigns the vertices connected by the dotted line to the merged node and that is injective on the other nodes is a surjective homomorphism equating only two vertices. Graph G in TW_2 , but G' is not.

Let H be the following graph



and let H' be the graph obtained by collapsing the nodes connected by the dotted line. As in the previous example, there is a surjective homomorphism $h : H \rightarrow H'$ equating only two vertices. In this case, graph H is not in TW_2 , but H' is.

C Confluence of the relation \sim

We prove confluence of relation \sim (Proposition 35). The proof relies on the following lemma.

Lemma 46. *Let G_1, G_2 be graphs in TW_2 . If $G_1 \rightleftharpoons G_2$ then there is an H such that $G_1 \rightsquigarrow H$ and $G_2 \rightsquigarrow H$.*

Proof. We prove the statement by induction on $sz(G_1) + sz(G_2)$, where $sz(G)$ is the sum of the number of nodes and edges in G . By the definition of $G_1 \rightleftharpoons G_2$ we have that there exist G'_1, G'_2 such that

$$G_1 \succ^* G'_1 \hookrightarrow G_2 \succ^* G'_2 \hookrightarrow G_1$$

By $G'_1 \hookrightarrow G_2 \succ^* G'_2$ we derive, by Proposition 31, that there is a G_3 such that $G'_1 \succ^* G_3 \hookrightarrow G'_2$. Hence, $G_1 \rightsquigarrow G_3$, which also implies $sz(G_1) \geq sz(G_3)$. If $sz(G_1) > sz(G_3)$, by $G_3 \rightleftharpoons G_2$ we can apply the inductive hypothesis and derive that there is an H such that $G_3 \rightsquigarrow H$ and $G_2 \rightsquigarrow H$. Since \rightsquigarrow is transitive, we have $G_1 \rightsquigarrow H$.

Otherwise, suppose $sz(G_1) = sz(G_3)$. Then $G_1 \simeq G'_1 \simeq G_3$, which implies $G_1 \hookrightarrow G_2$. Hence, $G_2 \rightsquigarrow G'_2$ and either $sz(G_2) > sz(G'_2)$, in which case we conclude by the inductive hypothesis and by the transitivity of \rightsquigarrow , or $sz(G_2) = sz(G'_2)$, in which case $G_1 \simeq G_2$. \square

We deduce confluence of \rightsquigarrow as follows: if $G \rightsquigarrow G_1$ and $G \rightsquigarrow G_2$ then $G_1 \rightleftharpoons G_2$, and by Lemma 46 we have that there is an H such that $G_1 \rightsquigarrow H$ and $G_2 \rightsquigarrow H$.

D Additional proofs for Section 8

Proof of Lemma 38. If $u = 1 \cap u$ then we have

$$\text{dom}(u) = \text{dom}(u \cap 1) = 1 \cap u1^\circ = 1 \cap u = u \quad (\text{by (A13)})$$

Conversely, we have

$$\begin{aligned} \text{dom}(u) &= \text{dom}(u)(1 \cap 1) \\ &= \text{dom}(u)1 \cap 1 \quad (\text{by (A14)}) \\ &= 1 \cap \text{dom}(u) \quad \square \end{aligned}$$

Proof of Lemma 42. The first item follows from the definition. We have

$$\begin{aligned}
 \alpha \diamond \beta \geq \gamma \diamond \delta & \text{ iff } \alpha \diamond \beta \cap \gamma \diamond \delta = \gamma \diamond \delta \\
 & \text{ iff } \alpha \gamma \diamond \beta \delta = \gamma \diamond \delta \\
 & \text{ iff } \alpha \gamma = \gamma \text{ and } \beta \delta = \delta \\
 & \text{ iff } \alpha \cap \gamma = \gamma \text{ and } \beta \cap \delta = \delta \quad (\text{by (5)}) \\
 & \text{ iff } \alpha \geq \gamma \text{ and } \beta \geq \delta \\
 \\
 \alpha \diamond \beta \geq \bar{v} & \text{ iff } \alpha \diamond \beta \cap \bar{v} = \bar{v} \\
 & \text{ iff } \overline{\alpha v \beta} \cap \bar{v} = \bar{v} \\
 & \text{ iff } \alpha v \beta \cap v = v \\
 & \text{ iff } \alpha v \beta \geq v
 \end{aligned}$$

Finally, $\bar{u} \geq \gamma \diamond \delta$ holds iff $\overline{\gamma u \delta} = \gamma \diamond \delta$, which never holds. \square

E Problems with Gutierrez' approach in [15]

We list the difficulties we encountered when trying to follow the strategy proposed by Gutierrez [15].

1. His Theorem 1 claims without proof that a certain axiomatisation over the signature Σ is sound and complete for graph isomorphism. We do not know how to obtain this result. We needed a full paper [7] to prove a similar claim for the signature Σ_{\top} (Theorem 17), and the conservativity technique we used in Section 8.1 to get one for the signature Σ_{dom} . It is not clear whether this technique could scale to prove Gutierrez' Theorem 1 using the result in [7]: our construction requires a domain operator.
2. It is not clear how the conditions of Definition 6 should be applied to graphs that are tests.
3. His Theorem 2 (soundness and completeness of a graph-theoretical notion w.r.t allegories) has a five-line proof.
 - For the 'only if' direction, it is not clear that his notion of *walk* in a graph is closed under contexts (*i.e.*, whether a counter-part to Lemma 26 holds).
 - The 'if' part relies on his Proposition 1, which is announced to hold by induction. We do not see how to deal with the case of composition in this induction: the induction hypothesis does not seem to be strong enough.
 - His Lemma 2 is used to define a notion of normal forms, which should then lead to decidability. It is however not clear that those normal forms are computable. While the set mentioned in Lemma 2 is indeed finite (it consists of subgraphs of a given graph), one has to consider infinitely many sequences (the arrows of his category $G(X)$) in order to decide membership in this set.