



HAL
open science

Positive regulation of raphe serotonin neurons by serotonin 2B receptors

Arnauld Belmer, Emily Quentin, Silvina L. Diaz, Bruno Guiard, Sebastian P Fernandez, Stéphane Doly, Sophie M. Banas, Pothitos Pitychoutis, Imane Moutkine, Aude Muzerelle, et al.

► To cite this version:

Arnauld Belmer, Emily Quentin, Silvina L. Diaz, Bruno Guiard, Sebastian P Fernandez, et al.. Positive regulation of raphe serotonin neurons by serotonin 2B receptors. *Neuropsychopharmacology*, 2018, 43 (7), pp.1623-1633. 10.1038/s41386-018-0013-0 . hal-01703916

HAL Id: hal-01703916

<https://hal.science/hal-01703916v1>

Submitted on 8 Feb 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Title. Positive regulation of raphe serotonin neurons by serotonin 2B receptors.

Short title. 5-HT_{2B} receptors in serotonergic neurons

Author List. Arnauld Belmer^{1,2,3,4*}, Emily Quentin^{1,2,3*}, Silvina L. Diaz^{1,2,3,5}, Bruno P. Guiard^{6,7,8}, Sebastian P. Fernandez^{1,2,3,9}, Stéphane Doly^{1,2,3,10}, Sophie M. Banas^{1,2,3}, Pothitos M. Pitychoutis^{1,2,3,11}, Imane Moutkine^{1,2,3}, Aude Muzerelle^{1,2,3}, Anna Tchenio^{1,2,3,12}, Anne Roumier^{1,2,3}, Manuel Mameli^{1,2,3,12}, and Luc Maroteaux^{1,2,3}

Author Affiliations. ¹INSERM UMR-S 839, F-75005, Paris, France; ²Sorbonne Universités, UPMC Univ Paris 6, F-75005, Paris; ³Institut du Fer à Moulin, F-75005, Paris; ⁴Translational Research Institute, Queensland University of Technology, Brisbane Qld 4059, Australia; ⁵Instituto de Biología Celular y Neurociencia, UBA-CONICET - Paraguay 2155, 3° piso, C1121ABG Buenos Aires, Argentina, Fac. de Cs. Exactas, Químicas y Naturales, Universidad de Morón, Argentina; ⁶Research Center on Animal Cognition, Center for Integrative Biology, F-31062 Toulouse France; ⁷Université Paul Sabatier, F-31062 Toulouse, France; ⁸UMR5169 CNRS, F-31062 Toulouse, France; ⁹IPMC – CNRS UMR7275 660 Route des Lucioles Sophia-Antipolis 06560 Valbonne France; ¹⁰Université Clermont Auvergne, INSERM, NEURO-DOL, F-63000 Clermont-Ferrand, France; ¹¹Department of Biology & Center for Tissue Regeneration and Engineering at Dayton (TREND), University of Dayton, 300 College Park, Dayton, 45469-2320, Ohio, USA; ¹²Dept. Fundamental Neurosciences (DNF) The University of Lausanne, Rue du Bugnon 9 1005 Lausanne Switzerland

*Equal contributors

Corresponding author: Luc Maroteaux, Institut du Fer à Moulin UMR-S839 INSERM/UPMC 17 rue du Fer à Moulin 75005 Paris Email: luc.maroteaux@upmc.fr; Tel: (33) 01 45 87 61 23 Fax: (33) 01 45 87 61 32

Number of words in abstract: 253; Number of words in introduction: 761; in results: 2086; in discussion 1670; Number of tables: 0; Number of figures: 5; Number of supplementary material 1.

Keywords: Serotonin, electrophysiology, viral overexpression, conditional knockout, MDMA, SSRI.

Abstract.

Serotonin is a neurotransmitter involved in many psychiatric diseases. In humans, a lack of 5-HT_{2B} receptors is associated with serotonin-dependent phenotypes, including impulsivity and suicidality. A lack of 5-HT_{2B} receptors in mice eliminates the effects of molecules that directly target serotonergic neurons including amphetamine-derivative serotonin releasers, and selective serotonin reuptake inhibitor antidepressants. In this work, we tested the hypothesis that 5-HT_{2B} receptors directly and positively regulate raphe serotonin neuron activity. By *ex-vivo* electrophysiological recordings, we report that stimulation by the 5-HT_{2B}-receptor agonist, BW723C86, increased the firing frequency of serotonin Pet1-positive neurons. Viral overexpression of 5-HT_{2B} receptors in these neurons increased their excitability. Furthermore, *in-vivo* 5-HT_{2B}-receptor stimulation by BW723C86 counteracted 5-HT_{1A} autoreceptor-dependent reduction in firing rate and hypothermic response in wildtype mice. By a conditional genetic ablation that eliminates 5-HT_{2B}-receptor expression specifically and exclusively from Pet1-positive serotonin neurons (*Htr2b*^{5-HTKO} mice), we demonstrated that behavioral and sensitizing effects of MDMA, as well as acute behavioral and chronic neurogenic effects of the antidepressant fluoxetine, require 5-HT_{2B}-receptor expression in serotonergic neurons. In *Htr2b*^{5-HTKO} mice, dorsal raphe serotonin neurons displayed a lower firing frequency compared to control *Htr2b*^{lox/lox} mice as assessed by *in-vivo* extracellular recordings and a stronger hypothermic effect of 5-HT_{1A}-autoreceptor stimulation was observed. The increase in head twitch response to DOI further confirmed the lower serotonergic tone resulting from the absence of 5-HT_{2B} receptors in serotonin neurons. Together, these observations indicate that the 5-HT_{2B} receptor acts as a direct positive modulator of serotonin Pet1-positive neurons in an opposite way as the known 5-HT_{1A} negative autoreceptor.

Introduction

Serotonin (5-Hydroxytryptamine, 5-HT) is involved in many psychiatric diseases including depression, addiction, impulsivity or psychosis. The 5-HT neurons that innervate forebrain originate predominantly from the rostral cell group of neurons in the dorsal raphe nucleus (DRN) (Commons, 2016; Okaty *et al*, 2015). These neurons express the serotonergic markers tryptophan hydroxylase (TPH2), and 5-HT transporter (SERT), and also the negative autoreceptors, 5-HT_{1A} and 5-HT_{1B} receptors, whose expression is restricted to somatodendritic compartments of 5-HT neurons, and to axonal terminals, respectively (Riad *et al*, 2000). The 5-HT_{1A} autoreceptor activation elicits an outward current carried through G protein-coupled inwardly-rectifying potassium channels (GIRK) of the Kir3 family leading to membrane hyperpolarization and inhibition of 5-HT neuron firing (Aghajanian and Lakoski, 1984). The presence of synaptic vesicles in dendrites of 5-HT neurons led to the suggestion that autoinhibition is mediated via dendritic release of 5-HT, for review see (Andrade *et al*, 2015). However, activity of 5-HT DRN neurons can also be positively modulated by 5-HT_{2A/2B/2C} receptors triggering directly or indirectly inward currents (Boothman *et al*, 2003; Craven *et al*, 2001; Kirby *et al*, 2003; Liu *et al*, 2000; Quéree *et al*, 2009). Upon electrical stimulation of leech 5-HT neurons, transmembrane Ca²⁺ entry through L-type channels first evokes an early dendritic exocytosis; subsequently, the released 5-HT activates dendritic 5-HT₂ autoreceptors coupled to Gq and phospholipase C, resulting in a positive feedforward loop that maintains sustained exocytosis (Leon-Pinzon *et al*, 2014). It has thus been proposed that DRN neurons can display responses ranging from inhibition to excitation depending on a balance of functional 5-HT_{1A} and 5-HT₂ receptors (Marinelli *et al*, 2004). However, a direct action of 5-HT₂ receptor subtypes at 5-HT neurons has not yet been clearly established.

In humans, a loss-of-function polymorphism of 5-HT_{2B} receptors is associated with 5-

HT-dependent phenotypes, including impulsivity and suicidality (Bevilacqua *et al*, 2010). Pharmacological experiments indicated that the 5-HT_{2B} receptor preferential agonist BW723C86 (Banas *et al*, 2011) mimicked serotonin specific reuptake inhibitor (SSRI) action in the forced swimming test (FST), which was abolished by injection of RS127445, a highly selective 5-HT_{2B} receptor antagonist, or in mice totally knocked-out (KO) for 5-HT_{2B} receptor gene (*Htr2b*^{-/-}) (Diaz and Maroteaux, 2011; Doly *et al*, 2008). Long-term behavioral and neurogenic SSRI effects were also abolished after either genetic ablation of 5-HT_{2B} receptors or chronic exposure to RS127445 (Diaz *et al*, 2012). Genetic (KO) or pharmacologic (antagonist) manipulation of 5-HT_{2B} receptors interferes similarly with effects of other molecules that target 5-HT neurons including amphetamine-derivatives, 5-HT releasers MDMA and dexfenfluramine (Banas *et al*, 2011; Doly *et al*, 2009; Doly *et al*, 2008). For example, the locomotor response to the psychostimulant MDMA was abolished in *Htr2b*^{-/-} or RS127445-treated wildtype mice (Doly *et al*, 2008). Furthermore, local infusion of BW723C86 in DRN by microdialysis increased extracellular 5-HT that was blocked by RS127445, supporting a functional role of this receptor within the raphe and the selectivity of BW723C86 for this effect (Doly *et al*, 2008).

Together, these studies suggested that 5-HT_{2B} receptors could be implicated in 5-HT-dependant behavior by acting directly and positively onto the 5-HT neurons. Nevertheless, the precise localization and way of action of 5-HT_{2B} receptors are still poorly investigated. Main difficulties include a lack of specific antibody as well as low level of expression of 5-HT_{2B} receptors in mouse (Loric *et al*, 1992) or human brains (Choi *et al*, 1994; Kursar *et al*, 1994). Nevertheless, expression of 5-HT_{2B} receptor mRNA was confirmed in several brain nuclei including DRN (Bevilacqua *et al*, 2010; Bonaventure *et al*, 2002). Besides, previous study using single cell RT-PCR (Diaz *et al*, 2012) permitted to establish a 5-HT_{2B} receptor expression not only in raphe but more specifically in 5-HT neurons. Nevertheless, precise role of 5-HT_{2B} receptors in 5-HT neurons remains elusive since pharmacologic manipulations or full gene knockout mice do not distinguish direct effects on 5-HT neuron from circuit effects.

In this work, we tested the hypothesis of a direct positive regulation of 5-HT neurons activity by 5-HT_{2B} receptors. First, we established that stimulation of 5-HT_{2B}-receptors can increase the firing frequency of 5-HT neurons, and counteract 5-HT_{1A} autoreceptor-dependent reduction in firing rate and hypothermic response in wildtype mice. Then, a conditional genetic ablation that eliminates 5-HT_{2B}-receptor expression specifically and exclusively from 5-HT neurons (*Htr2b*^{5-HTKO} mice), demonstrated a lack of effects of drugs, including SSRIs and MDMA, targeting 5-HT neurons, which present a lower firing rate and a stronger 8-OHDPAT-induced hypothermic response revealing its importance to maintain serotonergic tone. These observations established the direct modulation of 5-HT neurons by 5-HT_{2B} receptors acting in an opposite way as to 5-HT_{1A} negative autoreceptors.

Materials and Methods (see extended methods in supplementary materials)

Generation and use of mice -Knockout mice, *129S2.Cg-Htr2b^{tm1Lum}/Htr2b^{tm1Lum}* (*Htr2b*^{-/-}) were generated on the 129S2/SvPas strain. Floxed mice, *Htr2b^{tm2Lum}/Htr2b^{tm2Lum}*, (*Htr2b*^{lox/lox}) were generated on a mixed B6;129S2 background and backcrossed >10 times onto the 129S2 strain. *Htr2b^{lox/lox}* mice were inactivated for *Htr2b* in 5-HT neurons by crossing with *129S2.Cg-Tg(Fev-cre)^{IEsd}/0* (e*Pet1*-Cre BAC transgenic mice or *Pet1-Cre⁺/0*) (Scott *et al*, 2005) generating *129S2.Cg-Pet1-Cre⁺/0; Htr2b^{lox}/Htr2b^{lox}* conditional knockout mice (*Htr2b*^{5-HTKO}) and littermate controls (*Htr2b^{lox/lox}*). Then, *129S2.Cg-Tg(ROSA)26Sor^{tm1(CAG-EGFP)Fsh}* (Rosa26;CAG-loxP-STOP-loxP-EGFP or RCE) were crossed with *Pet1-Cre⁺/0* generating (*Pet1-GFP*) that express GFP in *Pet1* positive 5-HT neurons only after Cre

recombination. All experiments involving mice were approved by the local ethical committee (N° 1170.02).

Ex-vivo electrophysiology-Cell attached -Electrophysiological recordings and molecular characterization of 5-HT neurons was conducted in acute brain slices from male *Pet1-GFP* mice aged 3–4 weeks old according to published procedures (Fernandez *et al*, 2016; Vandermaelen and Aghajanian, 1983).

Viral constructs and stereotaxic injection -To express HA-tagged 5-HT_{2B} receptor specifically in 5-HT neurons, we use a Double floxed Inverse Orientation (DIO) adeno-associated virus (AAV) construct that allows Cre-mediated expression of the transgene (pAAV-EF1A-DIO-WPRE-pA vector; Addgene). The viruses packaged into AAV2.9 serotype with titers of 10¹²–10¹³ viral particles / ml were obtained (UNC Vector Core, Chapel Hill USA). AAV2.9 DIO-TdTomato was used as a control. AAVs were injected in the B7 raphe nuclei of *Pet1-GFP* following described procedure (Muzerelle *et al*, 2016).

Ex-vivo electrophysiology-Current clamp -Mice of either sex aged ~9 weeks were deeply anesthetized with ketamine and xylazine (150 mg/kg and 10 mg/kg, respectively; i.p.; Sigma-Aldrich, France) and immediately sacrificed. Coronal brain slices containing the raphe nucleus were prepared as previously described (Maroteaux and Mamei, 2012).

8-OHDPAT-induced hypothermia -Body temperature was then measured intrarectally in mice every 10 min after the agonist injection during 40 additional minutes (Bill *et al*, 1991).

In-vivo electrophysiology of DRN 5-HT neurons -The extracellular recordings of 5-HT neurons in DRN were performed in anesthetized mice (chloral hydrate, 400 mg/kg; i.p.) according to (Rainer *et al*, 2012). The DRN 5-HT neurons were identified according to the criteria of Aghajanian and Vandermaelen (Aghajanian and Vandermaelen, 1982b).

Locomotor response to novelty and MDMA-induced hyperlocomotion and sensitization -Locomotor activity was measured as previously described (Doly *et al*, 2008), in a circular corridor with four infrared beams placed at every 90° (Imetronic, Passac, France). MDMA-induced locomotor sensitization was performed in a two-injection protocol as previously described (Doly *et al*, 2009).

Prepulse Inhibition (PPI) of acoustic startle- Sensorimotor gating was assessed as previously described (Pitychoutis *et al*, 2015).

Forced Swimming Test -Mice FST was conducted as described (Diaz *et al*, 2011).

Fluoxetine treatments, Proliferation assay -Fluoxetine dose selection and FST were performed as described (Diaz *et al*, 2011). Neurogenesis study was performed as described (Diaz *et al*, 2012).

Synaptosome study -Crude synaptosomes were prepared as previously described (Gray and Whittaker, 1962). Synaptosomes 5-HT uptake and citalopram competition experiments were performed as described (Doly *et al*, 2008).

DOI-induced head-twitch -DOI-induced head twitch experiment was performed in clear plastic cages lined with bedding. Head-twitches (lateral movements of the head from side to side) were induced by \pm DOI (1, 3, 5 mg/kg, ip), and after 10 min, were scored for 10 minutes, by an experimenter unaware of mice genotype.

Statistical analysis- To determine differences between experimental groups, responses were analyzed by Two-tailed unpaired Student's t test or Mann-Whitney test when distribution was not normal. Also, a one- or two-way analysis of variance (ANOVA) repeated measure (RM) when appropriate, with Bonferroni's test was used for *post-hoc* multiple comparisons. In all cases, $P < 0.05$ was considered statistically significant.

Results

Ex-vivo electrophysiological responses to 5-HT_{2B}-receptor stimulation or overexpression

To examine the ability of 5-HT_{2B} receptors to regulate 5-HT neuron activity, we first performed cell-attached electrophysiological recordings of identified DRN 5-HT neurons in coronal slices of wildtype mice expressing GFP in Pet1-positive neurons (*Pet1-GFP* mice). *In-vivo*, DRN neurons fire with a regular pattern varying from 0.5 to 5 Hz. 5-HT neurons in *ex-vivo* coronal brain slices are electrically quiescent, but an excitatory noradrenergic tone facilitates firing (Haddjeri *et al*, 2004). In order to reproduce noradrenergic input, subsaturating concentrations of phenylephrine (Phe, 100-300 nM) were added to the bath (Vandermaelen *et al*, 1983). Low concentration (100 nM) of Phe was unable to initiate regular firing nor was the application of a preferential 5-HT_{2B} receptor agonist BW723C86 (1 μ M). Higher concentration of Phe (300 nM) was able to initiate and maintain a regular firing. Subsequent bath application of BW723C86 (1 μ M) induced a significant increase in firing frequency (1.73 \pm 0.23-fold) compared to Phe alone (**Fig. 1A**) (one way ANOVA repeated measures-RM-, effect of treatment $F_{1,6, 11.0} = 98$; $P < 0.0001$, n=8 cells). Bonferroni's *post-hoc* analysis showed a significant increase in firing frequency between each group (**Fig. 1A**). These data suggested that 5-HT_{2B}-receptor stimulation could increase Pet1-positive raphe 5-HT neuron firing activity.

To further establish that 5-HT_{2B} receptors can affect 5-HT neuron activity, we overexpressed a HA-tagged 5-HT_{2B} receptor specifically in Pet1-positive neurons, using a Double floxed Inverse Orientation (DIO) AAV construct that allows Cre-mediated expression of the tagged receptor. Viruses packaged into AAV2.9 serotype, AAV-DIO-5-HT_{2B}-HA or control AAV-DIO-TdTomato, were unilaterally injected into B7 raphe nucleus of *Pet1-GFP* mice (**Fig. S1A**). We confirmed the proper injection site by colocalization of either TdTomato expression or HA immunofluorescence with *Pet1-GFP* positive neurons (**Fig. S1B**). Coronal raphe-containing brain slices were used to record *Pet1-GFP* positive neuron excitability performed in current clamp mode. Recordings at various current steps showed that the number of action potentials obtained in function of the injected current was significantly increased in Pet1-positive neurons overexpressing 5-HT_{2B} receptors (two-way ANOVA RM, effect of overexpression, $F_{1,39} = 4.94$ $P = 0.032$, n=20 and 21 cells from 3 and 4 mice) (**Fig. 1B**). Input resistance value from recorded neurons was also significantly increased in 5-HT_{2B}-HA overexpressing neurons (unpaired t-test $t_{39} = 2.05$, $P = 0.047$) (**Fig. 1C**). *Ex-vivo* recordings indicated that the 5-HT_{2B}-receptor overexpression increased Pet1-positive neuron excitability.

In-vivo responses to 5-HT_{1A} and 5-HT_{2B} receptor agonists

Since our initial *ex-vivo* results indicated that 5-HT_{2B} receptors could increase 5-HT neuron activity, we hypothesized that it could work in an opposite way as to 5-HT_{1A} negative autoreceptor. We thus tested the effect of BW723C86 on 5-HT_{1A} receptor agonist 8-OHDPAT-induced inhibition of neuronal firing frequency *in-vivo*. Interestingly, 8-OHDPAT was significantly less potent in suppressing putative 5-HT neuron firing activity after BW723C86 injection as shown by two-way ANOVA RM (main effect of BW723C86 treatment $F_{1,9} = 7.34$; $P = 0.024$, n=6-5 mice). Bonferroni's *post-hoc* analysis revealed a reduced effect of 8-OHDPAT at 50 and 100 μ g/kg on putative 5-HT neuron firing activity (**Fig. 2A**). 8-OHDPAT ED₅₀ was shifted about 3.3-fold from 45 to 148 μ g/kg; BW723C86 alone did not modify firing (not illustrated).

Although the mechanism of 5-HT_{1A} receptor agonist-induced hypothermia is incompletely understood (Jacobsen *et al*, 2012), the 8-OHDPAT-induced hypothermia in mice is known to be mediated by 5-HT_{1A} autoreceptor (Richardson-Jones *et al*, 2010). We thus used the 8-OHDPAT-induced hypothermia as another *in-vivo* readout of the functional

status of 5-HT_{1A} autoreceptor and tested putative effects of stimulating 5-HT_{2B} receptors. In agreement with the above-mentioned effect on firing, a pretreatment of wildtype mice with BW723C86 (5 mg/kg sc) was able to significantly reduce the ability of 8-OHDPAT (0.3 mg/kg sc), to induce hypothermia in wildtype mice (**Fig. 2B**), as shown by two-way ANOVA RM analysis (interaction between the time and treatment, $F_{9, 54} = 10.34$; $P < 0.0001$, $n=4$ mice per group). Bonferroni's *post-hoc* analysis showed a significant reduction by BW723C86 at 30 and 40 min post-injection of the 8-OHDPAT-induced hypothermia (**Fig. 2B**). These data indicated that 5-HT_{1A} autoreceptor inhibitory activities can be attenuated by concomitant activation of 5-HT_{2B} receptors.

Absence of response to ecstasy, MDMA, in *Htr2b*^{5-HTKO} mice

To establish putative direct actions of 5-HT_{2B} receptors on 5-HT neurons, we developed mice with conditional ablation of this receptor gene specifically in Pet1-positive neurons. We inserted recombination sites (loxP) flanking the second exon of *Htr2b* gene (*Htr2b*^{lox/lox}) and crossed these mice with mice expressing the Cre recombinase under *Pet1* gene promoter (BAC-transgenic). *Pet1* gene expression in brain is restricted to most of differentiated 5-HT neurons and their postmitotic precursors (Scott *et al*, 2005). We thus generated *Pet1-Cre*^{+/0}; *Htr2b*^{lox/lox} mice (*Htr2b*^{5-HTKO}) (**Fig. S2A**). Restricted recombination in raphe 5-HT neurons was revealed by analysis of raphe DNA and by colocalization of a Cre-dependent GFP reporter with 5-HT staining (**Fig. S2B-C**) (Scott *et al*, 2005). We quantified the efficiency of recombination and found that 83.6±6.1% of the Pet1-Cre-dependent GFP was colocalized with 5-HT staining, 11.6±3.9% was only 5-HT positive and less than 5% (4.8±2.9%) was only GFP positive ($n=12$ from 3 different mice), supporting an efficient and specific recombination in 5-HT neurons (**Fig. S2C**).

MDMA, the active compound of ecstasy, is a substrate of SERT leading to massive 5-HT release from synaptic vesicle stores, which is partially calcium dependent (Crespi *et al*, 1997). The *Htr2b*^{5-HTKO} mice injected with MDMA (20 mg/kg) did not display increased locomotor responses in contrast to control *Htr2b*^{lox/lox} littermate mice (**Fig. 3A-B**). Two-way ANOVA RM analysis showed a main effect of genotype ($F_{1, 30} = 4.22$; $P = 0.049$, $n=16$ mice per group). Bonferroni's *post-hoc* analysis showed a significant difference between genotypes from $t=25$ to $t=45$ min post-injection (**Fig. 3A**). The analysis of the total locomotor activity over the first 60 min after MDMA injection confirmed the significant effect of genotypes (two-way ANOVA RM, $F_{1, 60} = 4.61$; $P = 0.036$, $n=16$ mice per group). Bonferroni's *post-hoc* analysis showed a significant increase in locomotion in control *Htr2b*^{lox/lox} mice, but not in *Htr2b*^{5-HTKO} mice (**Fig. 3B**). We further assessed the contribution in Pet1-positive neurons of 5-HT_{2B} receptors to MDMA-induced locomotor sensitization using a two-injection protocol (Valjent *et al*, 2010). A significant increase in locomotor activity was observed in *Htr2b*^{lox/lox} control mice as shown by two-way ANOVA RM analysis (main effect of genotype, $F_{1, 14} = 5.12$; $P = 0.04$). Bonferroni's *post-hoc* analysis showed a significant increase in locomotion in control *Htr2b*^{lox/lox} mice at the 2nd injection of MDMA (20 mg/kg) (1st injection 639 ± 222 vs. 2nd injection 1291 ± 295 1/4 of turns, $n=8$), but not in *Htr2b*^{5-HTKO} (1st injection 254 ± 132 vs. 2nd injection 353 ± 174 , $n=8$) (**Fig. 3C**). These results indicated a lack of behavioral and sensitizing effects of MDMA in *Htr2b*^{5-HTKO} mice as found in full KO, *Htr2b*^{-/-} mice, confirming the need for this receptor in Pet1-positive 5-HT neurons.

Absence of response to the SSRI Fluoxetine in *Htr2b*^{5-HTKO} mice

SSRIs are known to block 5-HT reuptake by SERT, leading to extracellular 5-HT accumulation following vesicular 5-HT release. A classical outcome for acute response to SSRIs is the reduced immobility time observed in forced swimming test (FST). We tested the effect of acute fluoxetine injection (3 mg/kg, ip), the optimal dose determined for 129S2

strain in FST (Diaz *et al*, 2011). Fluoxetine injection did not affect immobility time in *Htr2b*^{5-HTKO} mice (**Fig. 4A**) as shown by two-way ANOVA analysis (time and genotype interactions, $F_{1, 24} = 10.67$; $P < 0.003$, $n=7-7$). However, a significant reduction of immobility time was observed in control *Htr2b*^{lox/lox} littermates as Bonferroni's *post-hoc* analysis showed the only significant difference between vehicle and fluoxetine-treated *Htr2b*^{lox/lox} control mice (**Fig. 4A**).

Long-term effects of SSRI are known to be associated with hippocampus subgranular zone (SGZ) neuronal proliferation. We performed a daily ip injection for 4 weeks of fluoxetine (3 mg/kg), BrdU injections the 27th day, and quantified BrdU incorporation in SGZ neurons of the hippocampus at the 28th day. Chronic injection of fluoxetine did not produce changes in BrdU incorporation in *Htr2b*^{5-HTKO} mice (**Fig. 4B**) (two-way ANOVA analysis showed a trend for treatment and genotype interactions, $F_{1, 26} = 3.88$; $P = 0.059$, $n=7-8$). An increase was detected in controls *Htr2b*^{lox/lox} littermates as Bonferroni's *post-hoc* analysis showed the only significant difference between vehicle and fluoxetine-treated *Htr2b*^{lox/lox} controls (**Fig. 4B**). These results indicated that the lack of both acute behavioral and chronic neurogenic effects of SSRIs found in *Htr2b*^{-/-} mice was due to 5-HT_{2B}-receptor elimination from Pet1-positive neurons.

We recently reported that *Htr2b*^{-/-} mice displayed novelty-induced hyperlocomotion and a global deficit in sensorimotor gating (Pitychoutis *et al*, 2015). However, locomotor activity in a new environment was not different between *Htr2b*^{5-HTKO} and *Htr2b*^{lox/lox} littermate control mice over the first 60-minute (**Fig. S3A**). Similarly, prepulse inhibition (PPI) of startle reflex and startle amplitude were not different between *Htr2b*^{5-HTKO} and *Htr2b*^{lox/lox} littermate control mice (**Fig. S3B**). Although present in *Htr2b*^{-/-} mice, these deficits were not found in mice lacking the 5-HT_{2B} receptor selectively in 5-HT neurons, supporting the specificity of these conditional mice.

Since SERT targeting drug (MDMA and fluoxetine) action is affected by the lack of 5-HT_{2B} receptors in Pet1-positive neurons, we next determined possible alterations of SERT expression or function in brain synaptosome preparations from *Htr2b*^{5-HTKO} mice. Heterologous competition binding experiments showed no difference in the density of citalopram binding sites ($B_{max} 305 \pm 18$ vs. 389 ± 26 fmol/mg of protein) or affinity ($K_i 0.4 \pm 0.1$ vs. 0.17 ± 0.06 nM) between *Htr2b*^{5-HTKO} mice and their *Htr2b*^{lox/lox} littermate control mice (**Fig. 4C**). Additionally, 5-HT uptake experiments on brain synaptosomes showed no difference in 5-HT transport maximum velocity (V_{max} ; 5.57 ± 0.32 vs. 6.49 ± 0.38 fmol/sample/min) or apparent affinity ($K_m 40.7 \pm 2.6$ vs. 27.7 ± 2.5 nM) between *Htr2b*^{5-HTKO} mice and their *Htr2b*^{lox/lox} littermate controls (**Fig. 4C**). Together, these results demonstrated that selective ablation of 5-HT_{2B} receptors in raphe Pet1-positive neurons eliminates MDMA and fluoxetine actions but does not affect SERT expression and activity.

***Htr2b*^{5-HTKO} mice display a hyposerotonergic phenotype**

To determine whether the 5-HT_{2B} receptor had an overall effect on 5-HT neurons, we measured firing rates of putative 5-HT DRN neurons *in-vivo* in *Htr2b*^{5-HTKO} mice. Neurons were included in this analysis based on characteristics and averaged traces of their action potentials (Vandermaelen *et al*, 1983). *Htr2b*^{5-HTKO} mice display a higher percentage of putative 5-HT neurons discharging with a low firing mode (16.1% vs. 9.3% <1 Hz) and a lower percentage of neurons with a high firing mode (5.9% vs. 13.9% >4 Hz) relative to *Htr2b*^{lox/lox} mice. This observation was reflected by a significant shift in cumulative distribution of neurons with lower firing rate in *Htr2b*^{5-HTKO} mice compared to control *Htr2b*^{lox/lox} mice ($n = 108$ and 118 neurons, respectively; Kolmogorov-Smirnov test; $P=0.0008$) (**Fig. 5A**). These results revealed that ablation of 5-HT_{2B} receptors in raphe Pet1-positive neurons is sufficient to modify 5-HT neuron firing rate.

Head-twitch response is a rhythmic paroxysmal rotational head movement that occurs in mice and rats treated by a variety of serotonergic hallucinogens, including LSD and DOI (Halberstadt, 2015). This behavior is specifically linked to 5-HT_{2A} receptor activation, since selective 5-HT_{2A} receptor antagonists block head-twitch response induced by DOI and other hallucinogens, and it is absent in *Htr2a*^{-/-} mice (Halberstadt, 2015). Here, we tested ±DOI at 1, 3, and 5-mg/kg ip. We found that a dose of 5 mg/kg induced a larger increase in head-twitch in *Htr2b*^{5-HTKO} mice, as compared to *Htr2b*^{lox/lox} littermate control (+109%, $t_{13} = 2.68$, $n = 3-5$, $P = 0.016$, multiple t-test) (**Fig. 5B**). Similarly, a greater head-twitch response was observed in *Htr2b*^{-/-} compared with control mice at the same dose (**Fig. S4A**). Nevertheless, no significant change in 5-HT_{2A} or 5-HT_{1A} receptor expression or in 5-HT content and turnover was found in PFC from *Htr2b*^{-/-} compared to control mice (**Fig. S4B-D**). Since a drastic increase in number of DOI-induced head-twitch was observed in the *Tph2-R439H* knock-in mouse (Jacobsen *et al*, 2012), a mouse model with 60-80% reduction in TPH2 activity and thus with low serotonergic tone, the increased behavioral response to DOI confirmed a lower serotonergic tone in *Htr2b*^{5-HTKO} mice.

Finally, we used the 8-OHDPAT-induced hypothermia as an *in-vivo* readout of the functional status of 5-HT_{1A} autoreceptor in *Htr2b*^{5-HTKO} mice. The hypothermic response to 8-OHDPAT (0.1 mg/kg sc) was significantly increased in *Htr2b*^{5-HTKO} mice compared to control *Htr2b*^{lox/lox} mice as shown by two-way ANOVA RM analysis (interaction between time and treatment, $F_{4,36} = 5.30$; $P = 0.0018$, $n = 3-8$ mice). Bonferroni's *post-hoc* analysis showed a significant stronger hypothermic effect at 20 min post-injection of the 8-OHDPAT (**Fig. 5C**). These data supported that 5-HT_{1A} autoreceptor inhibitory activity is increased upon inactivation of 5-HT_{2B} receptors in Pet1-positive neurons.

Discussion

Our results show that 5-HT_{2B} receptors can positively modulate 5-HT neuron activity in brain slices of wildtype mice, and counteract 5-HT_{1A} negative autoreceptor actions. Furthermore, mice lacking 5-HT_{2B} receptors exclusively in Pet1-positive 5-HT neurons, *Htr2b*^{5-HTKO} mice, display an excess of inhibitory control exerted by 5-HT_{1A} autoreceptors supporting an opposite control of 5-HT_{2B} receptors on 5-HT neurons.

5-HT_{2B} receptors positively regulate 5-HT neurons- There is a growing consensus that 5-HT neurons are non-homogeneous as supported by anatomical, biochemical and electrophysiological studies (Andrade and Haj-Dahmane, 2013; Calizo *et al*, 2011; Fernandez *et al*, 2016; Marinelli *et al*, 2004). Sub-populations of 5-HT neurons, either within the DRN or between various raphe nuclei, are interconnected, and form complex circuits (Altieri *et al*, 2013; Bang *et al*, 2012; Gaspar and Lillesaar, 2012). The activity of 5-HT neurons can be modulated by both 5-HT_{1A} and 5-HT_{2A/2B/2C} receptors (Craven *et al*, 2001; Liu *et al*, 2000). Identified 5-HT neurons are known to respond to 5-HT_{1A} receptor agonists by a 5-HT-induced outward current (Beck *et al*, 2004; Craven *et al*, 2001; Kirby *et al*, 2003; Liu *et al*, 2000). A significant proportion of TPH2-positive neurons (about 50%) also respond to 5-HT₂-receptor activation by an inward current (Marinelli *et al*, 2004). Tonic spiking of 5-HT neurons establishes synaptic 5-HT levels. We show here, by cell-attached recordings of identified wildtype raphe Pet1-positive neurons, that stimulation of 5-HT_{2B} receptors by BW723C86 can increase their firing frequency. The effect on Pet1-positive neuron firing frequency of the 5-HT_{2B} receptor agonist can only be detected in the presence of phenylephrine (300 nM), suggesting that although the two receptors share common Gq signal transduction, they are likely acting at different effectors that remains to be identified. Independent electrophysiological current-clamp recordings showed that overexpression of 5-HT_{2B} receptors in Pet1-positive 5-HT neurons was sufficient to increase their excitability.

These results indicate that 5-HT_{2B} receptors can positively control the firing of 5-HT neurons. This was confirmed *in-vivo* by extracellular recordings in *Htr2b*^{5-HTKO} mice of putative 5-HT neurons that showed a significant shift to low firing rate. Together, the present results revealed a need for 5-HT_{2B} receptors in 5-HT neurons to positively regulate their activity.

Serotonergic tone results from an opposite control exerted by 5-HT_{1A} and 5-HT_{2B} receptors- The lack of effects of MDMA and fluoxetine in the absence of 5-HT_{2B} receptors in Pet1-positive 5-HT neurons, previously observed in full knockout mice *Htr2b*^{-/-} associated to reduced extracellular 5-HT accumulation as assessed by microdialysis (Diaz *et al*, 2012; Doly *et al*, 2008), raised the possibility of an interaction of 5-HT_{2B} receptors with SERT. However, the absence of modification in SERT uptake and expression lowers this possibility. The unique control of dendritic 5-HT release has important implications for DRN physiology and actions of SERT-targeting drugs, SSRIs and MDMA. Packaging by the vesicular monoamine transporter (VMAT2) is essential for 5-HT transmission; glutamate receptor activation in dorsal raphe brain slice can evoke somatodendritic release by vesicle exocytosis (Colgan *et al*, 2009). SSRI antidepressants, which have a more limited effect on 5-HT release from dendrites than from the soma and terminals, markedly increase extracellular 5-HT in DRN that involves both somatic and dendritic release (Colgan *et al*, 2009). The 5-HT released within DRN induces feedback inhibition of 5-HT neuron firing activity by stimulation of somatodendritic 5-HT_{1A} negative autoreceptors, which results from local release rather than extended diffusion of 5-HT throughout the extracellular space (Courtney and Ford, 2016). We found that the hypothermic response to 8-OHDPAT, known to be mediated by 5-HT_{1A} auto- but not hetero-receptors in mice (Richardson-Jones *et al*, 2010), is attenuated by pretreatment with the 5-HT_{2B} receptor agonist BW723C86. However, we never observed an effect of the 5-HT_{2B}-receptor agonist in the absence of 5-HT_{1A}-receptor agonist. These observations support either a dominant effect of 5-HT_{1A} over 5-HT_{2B} receptors or combination of 5-HT_{2B} auto- and hetero-receptors having opposite actions. Richardson-Jones *et al.*, (Richardson-Jones *et al*, 2010) generated a mouse strain differing in 5-HT_{1A} autoreceptor expression by approximately 30-40% below the wildtype level (1A-Low). These 1A-Low mice showed reduced 8-OHDPAT-induced hypothermia and their neurons exhibit a shift toward higher firing rates. To the opposite, in the absence of 5-HT_{2B} receptors in Pet1-neurons (in *Htr2b*^{5-HTKO} mice), we found a significant increase in hypothermic response to 8-OHDPAT and a significant shift toward lower frequency firing neurons. These findings support that the lack of 5-HT_{2B} receptor in Pet1-positive 5-HT neurons is associated with a higher 5-HT_{1A}-autoreceptor reactivity and thus a lower activity of 5-HT neurons. The lower serotonergic tone observed in the absence of 5-HT_{2B} receptors in Pet1-positive neurons would thus result from the opposite control exerted by 5-HT_{1A} and 5-HT_{2B} receptors on DRN neurons. This may explain the lack of actions of SERT-targeting drugs, SSRIs and MDMA, although the detailed mechanism remains to be identified. An interaction between 5-HT_{1A} and 5-HT_{2B} receptors directly or via trans regulation could be involved since we previously reported such cross-talks between 5-HT_{1B} and 5-HT_{2B} receptors (Janoshazi *et al*, 2007).

5-HT_{2B} receptors contribute to SSRI therapeutic effects- The excess of inhibitory control exerted by 5-HT_{1A} receptors in the absence of 5-HT_{2B} receptors in Pet1-positive 5-HT neurons may also explain the lack of response to chronic SERT blockers (fluoxetine) in *Htr2b*^{5-HTKO} mice. Chronic SSRI antidepressant responses are at least partially ascribed to desensitization of somatodendritic 5-HT_{1A} receptors (Richardson-Jones *et al*, 2010). Recent works using chemogenetic approaches (i.e., Designer Receptors Exclusively Activated by Designer Drugs-DREADDs) showed that CNO activation of SERT- or Pet1-positive neurons expressing the Gq-coupled M3Gq DREADD induced an increase in 5-HT neuron firing rate and a reduction in immobility in FST (Teissier *et al*, 2015; Urban *et al*, 2016). We showed previously that activation of 5-HT_{2B} Gq-coupled receptors with BW723C86 mimicked both

acute and chronic behavioral and neurogenic effects of SSRI antidepressants and led to extracellular 5-HT accumulation, which were eliminated in *Htr2b*^{-/-} mice or by RS127445 (Diaz *et al.*, 2012; Doly *et al.*, 2008). We found here that knocking-out the 5-HT_{2B} receptors exclusively from Pet1-positive neurons (*Htr2b*^{5-HTKO}) mice is sufficient to eliminate behavioral effects in FST and neurogenic effects of fluoxetine and that 5-HT_{2B}-receptor overexpression increases Pet1-positive neuron excitability. It appears thus that 5-HT_{2B} receptors contribute to SSRI therapeutic effects by their positive Gq-dependent signaling on adult raphe 5-HT neurons, which may be revealed upon somatodendritic 5-HT_{1A}-receptor desensitization.

The 5-HT neuron firing relies on a balance of functional 5-HT_{1A} and 5-HT_{2B} receptors- The reason why positive 5-HT_{2B} receptors acting in an opposite manner to negative 5-HT_{1A} autoreceptors has not been previously identified could have several explanations. Recently, the role for 5-HT_{1A} receptor-mediated autoinhibition of the DRN in homeostatic control of firing rate has been questioned. As discussed by Andrade *et al.*, (Andrade *et al.*, 2015), 5-HT_{1A} autoinhibition may participate in regulating glutamate signaling to 5-HT neurons (Colgan *et al.*, 2012) or in mediating inputs from distal serotonergic cell groups (Bang *et al.*, 2012). De Kock *et al.*, (de Kock *et al.*, 2006) first showed that following calcium influx through NMDA receptors, 5-HT could be released from DRN neuron dendrites in the absence of postsynaptic firing. Colgan *et al.*, (Colgan *et al.*, 2012) reported that 5-HT release from dendrites is secondary to calcium influx through L-type calcium channels that open in response to the local dendritic depolarization elicited by synaptically released glutamate. A contribution of dendritic 5-HT release to 5-HT_{1A}-autoreceptor activation would thus result from excitatory glutamatergic inputs to DRN via locally triggered calcium influx rather than by neuronal firing. Independently, it has been reported that upon electrical stimulation of leech 5-HT neurons, transmembrane Ca²⁺ entry through L-type channels can first evoke an early dendritic exocytosis; subsequently, the released 5-HT activates 5-HT₂ autoreceptors coupled to Gq and phospholipase C, resulting in a positive feedforward loop that maintains sustained exocytosis (Leon-Pinzon *et al.*, 2014). In frog motoneurons, a potentiation of NMDA-induced depolarization has been shown to depend on the activation of 5-HT_{2B} receptors causing an influx of extracellular Ca²⁺ through L-type Ca²⁺ channels and a reduction of the open-channel block of NMDA receptors (Holohean and Hackman, 2004). Since serotonergic DRN neurons can respond to 5-HT with responses ranging from inhibition to excitation with the net effect of 5-HT relying on a balance of functional 5-HT_{1A} and 5-HT_{2A/2B/2C} receptors (Marinelli *et al.*, 2004), combined expression levels of these receptors in various 5-HT subpopulations may set-up DRN firing levels. Since DRN receives serotonergic inputs from caudal raphe nuclei (Bang *et al.*, 2012), 5-HT released in DRN may also originate from extrinsic serotonergic afferents. Our results show that 5-HT_{2B} receptors can positively modulate 5-HT neuron activity, and counteract 5-HT_{1A} negative autoreceptor actions. Furthermore, our recent unpublished data revealed a somatodendritic expression of 5-HT_{2B} receptors as 5-HT_{1A} receptors. Whether 5-HT_{2B} receptors act at different 5-HT neurons subtypes, from different raphe nuclei, or by direct interaction with 5-HT_{1A} receptors, remain to be determined.

Concluding remarks- A remaining question is why both positive and negative autoreceptors are needed to regulate 5-HT neuron activity. In locus coeruleus (LC), the major noradrenergic nucleus of the brain, resting membrane potential and pattern of spontaneous firing neurons in newborns are affected by both α 1- and α 2-adrenergic receptors activation (Igata *et al.*, 2014). Application of α 2-adrenergic receptor agonists activates GIRK channels, resulting in membrane hyperpolarization and inhibition of spontaneous firing of action potentials (Aghajanian and VanderMaelen, 1982a). By contrast, stimulating α 1-adrenergic receptors, known to couple to a Gq protein, induces membrane depolarization and accelerates

spontaneous firing rates in neonatal rat LC neurons (Williams and Marshall, 1987), whose alterations may trigger pathological conditions (Igata *et al*, 2014). Mixtures of positive and negative feedback has been modeled and appears to be necessary to create oscillatory signal outputs (Brandman and Meyer, 2008), which are important for neuronal network functions such as DRN or LC. Our findings established that Gq-coupled 5-HT_{2B} receptors expressed by Pet1-positive 5-HT neurons act in an opposite manner as to 5-HT_{1A} autoreceptors. The 5-HT_{2B} receptor can thus be considered as a positive modulator of serotonergic tone that acts at 5-HT neuron excitability. This positive modulation has to be taken into account in the studies of the regulatory mechanisms of 5-HT neurons including those of antidepressants.

Funding and Disclosure

This work has been supported by funds from the *Centre National de la Recherche Scientifique*, the *Institut National de la Santé et de la Recherche Médicale*, the *Université Pierre et Marie Curie*, and by grants from the *Fondation pour la Recherche Médicale* "Equipe FRM DEQ2014039529", the French Ministry of Research (Agence Nationale pour la Recherche ANR-12-BSV1-0015 and ANR-17-CE16-0008 and the Investissements d'Avenir programme ANR-11-IDEX-0004-02). LM's team is part of the École des Neurosciences de Paris Ile-de-France network and of the Bio-Psy Labex and as such this work was supported by French state funds managed by the ANR within the Investissements d'Avenir programme under reference ANR-11-IDEX-0004-02. A Roumier has been supported by grants from the *Université Pierre et Marie Curie (Emergence-UPMC program)* and the Bio-Psy Labex. S. Doly has been supported by a fellowships of the Lefoulon-DeLalande foundation, SL Diaz from the Region Ile de France DIM STEM and from the ANPCyT (PICT 2013-3225), CONICET (PIP-11220130100157CO), and University of Moron (PID 2015), and E. Quentin by a PhD fellowship from the Region Ile de France DIM Cerveau et Pensée. All other authors have nothing to disclose.

Acknowledgment.

We thank the Mouse Clinical Institute (Strasbourg) for *Htr2b* floxed mice production, Evan Deneris for providing *Pet1-Cre* BAC transgenic mice, Mythili Savariradjane and the Imaging facility of the IFM, and Natacha Roblot and the IFM animal facility.

Author Contribution.

A. B., E. Q., S. L. D., S. P. F., S. D., S. M. B., P. M. P., I. M., A. M., and A. T., conducted the experiments, A. R., conducted and designed the experiments, B. P. G., conducted, designed the experiments, and wrote the paper, M. M. supervised, wrote the paper and provided funding, L. M. supervised, wrote the paper and provided funding.

References

- Aghajanian GK, Lakoski JM (1984). Hyperpolarization of serotonergic neurons by serotonin and LSD: studies in brain slices showing increased K⁺ conductance. *Brain Research* **305**(1): 181-185.
- Aghajanian GK, VanderMaelen CP (1982a). alpha 2-adrenoceptor-mediated hyperpolarization of locus coeruleus neurons: intracellular studies in vivo. *Science (New York, NY)* **215**(4538): 1394-1396.
- Aghajanian GK, Vandermaelen CP (1982b). Intracellular recording in vivo from serotonergic neurons in the rat dorsal raphe nucleus: methodological considerations. *J Histochem Cytochem* **30**(8): 813-814.
- Altieri SC, Garcia-Garcia AL, Leonardo ED, Andrews AM (2013). Rethinking 5-HT_{1A} receptors: emerging modes of inhibitory feedback of relevance to emotion-related behavior. *ACS Chem Neurosci* **4**(1): 72-83.

- Andrade R, Haj-Dahmane S (2013). Serotonin neuron diversity in the dorsal raphe. *ACS Chem Neurosci* **4**(1): 22-25.
- Andrade R, Huereca D, Lyons JG, Andrade EM, McGregor KM (2015). 5-HT1A Receptor-Mediated Autoinhibition and the Control of Serotonergic Cell Firing. *ACS Chem Neurosci* **6**(7): 1110-1115.
- Banas SM, Doly S, Boutourlinsky K, Diaz SL, Belmer A, Callebert J, *et al* (2011). Deconstructing antiobesity compound action: requirement of serotonin 5-HT2B receptors for dexfenfluramine anorectic effects. *Neuropsychopharmacology* **36**(2): 423-433.
- Bang SJ, Jensen P, Dymecki SM, Commons KG (2012). Projections and interconnections of genetically defined serotonin neurons in mice. *European J Neurosci* **35**(1): 85-96.
- Beck SG, Pan Y-Z, Akanwa AC, Kirby LG (2004). Median and dorsal raphe neurons are not electrophysiologically identical. *J Neurophysiology* **91**(2): 994-1005.
- Bevilacqua L, Doly S, Kaprio J, Yuan Q, Tikkanen R, Paunio T, *et al* (2010). A population-specific HTR2B stop codon predisposes to severe impulsivity. *Nature* **468**(8): 1061-1066.
- Bill DJ, Knight M, Forster EA, Fletcher A (1991). Direct evidence for an important species difference in the mechanism of 8-OH-DPAT-induced hypothermia. *Br J Pharmacol* **103**(4): 1857-1864.
- Bonaventure P, Guo H, Tian B, Liu X, Bittner A, Roland B, *et al* (2002). Nuclei and subnuclei gene expression profiling in mammalian brain. *Brain Res* **943**(1): 38-47.
- Boothman LJ, Allers KA, Rasmussen K, Sharp T (2003). Evidence that central 5-HT2A and 5-HT(2B/C) receptors regulate 5-HT cell firing in the dorsal raphe nucleus of the anaesthetised rat. *Br J Pharmacol* **139**(5): 998-1004.
- Brandman O, Meyer T (2008). Feedback loops shape cellular signals in space and time. *Science (New York, NY)* **322**(5900): 390-395.
- Calizo LH, Akanwa A, Ma X, Pan Y-Z, Lemos JC, Craig C, *et al* (2011). Raphe serotonin neurons are not homogenous: Electrophysiological, morphological and neurochemical evidence. *Neuropharmacology* **61**(3): 524-543.
- Choi D-S, Birraux G, Launay J-M, Maroteaux L (1994). The human serotonin 5-HT2B receptor: pharmacological link between 5-HT2 and 5-HT1D receptors. *FEBS L* **352**: 393-399.
- Colgan LA, Cavolo SL, Commons KG, Levitan ES (2012). Action potential-independent and pharmacologically unique vesicular serotonin release from dendrites. *J Neurosci* **32**(45): 15737-15746.
- Colgan LA, Putzier I, Levitan ES (2009). Activity-dependent vesicular monoamine transporter-mediated depletion of the nucleus supports somatic release by serotonin neurons. *J Neurosci* **29**(50): 15878-15887.
- Commons KG (2016). Ascending serotonin neuron diversity under two umbrellas. *Brain Structure and Function* **221**(7): 3347-3360.
- Courtney NA, Ford CP (2016). Mechanisms of 5-HT1A receptor-mediated transmission in dorsal raphe serotonin neurons. *J Physiol* **594**(4): 953-965.
- Craven RM, Grahame-Smith DG, Newberry NR (2001). 5-HT1A and 5-HT2 receptors differentially regulate the excitability of 5-HT-containing neurones of the guinea pig dorsal raphe nucleus in vitro. *Brain Research* **899**(1-2): 159-168.
- Crespi D, Mennini T, Gobbi M (1997). Carrier-dependent and Ca(2+)-dependent 5-HT and dopamine release induced by (+)-amphetamine, 3,4-methylenedioxymethamphetamine, p-chloroamphetamine and (+)-fenfluramine. *Br J Pharmacol* **121**(8): 1735-1743.

de Kock CPJ, Cornelisse LN, Burnashev N, Lodder JC, Timmerman AJ, Couey JJ, *et al* (2006). NMDA receptors trigger neurosecretion of 5-HT within dorsal raphe nucleus of the rat in the absence of action potential firing. *J Physiol* **577**(Pt 3): 891-905.

Diaz SL, Doly S, Narboux-Nême N, Fernandez S, Mazot P, Banas S, *et al* (2012). 5-HT_{2B} receptors are required for serotonin-selective antidepressant actions. *Mol Psychiatry* **17**: 154-163.

Diaz SL, Maroteaux L (2011). Implication of 5-HT_{2B} receptors in the serotonin syndrome. *Neuropharmacology* **61**: 495-502.

Doly S, Bertran-Gonzalez J, Callebert J, Bruneau A, Banas SM, Belmer A, *et al* (2009). Role of serotonin via 5-HT_{2B} receptors in the reinforcing effects of MDMA in mice *PLoS ONE* **4**(11): e7952.

Doly S, Valjent E, Setola V, Callebert J, Herve D, Launay JM, *et al* (2008). Serotonin 5-HT_{2B} receptors are required for 3,4-methylenedioxymethamphetamine-induced hyperlocomotion and 5-HT release in vivo and in vitro. *J Neurosci* **28**(11): 2933-2940.

Fernandez SP, Cauli B, Cabezas C, Muzerelle A, Poncer J-C, Gaspar P (2016). Multiscale single-cell analysis reveals unique phenotypes of raphe 5-HT neurons projecting to the forebrain. *Brain Structure and Function* **221**(8): 4007-4025.

Gaspar P, Lillesaar C (2012). Probing the diversity of serotonin neurons. *Philos Trans R Soc Lond, B, Biol Sci* **367**(1601): 2382-2394.

Gray EG, Whittaker VP (1962). The isolation of nerve endings from brain: an electron-microscopic study of cell fragments derived by homogenization and centrifugation. *J Anat* **96**: 79-88.

Haddjeri N, Lavoie N, Blier P (2004). Electrophysiological evidence for the tonic activation of 5-HT_{1A} autoreceptors in the rat dorsal raphe nucleus. *Neuropsychopharmacology* **29**(10): 1800-1806.

Halberstadt AL (2015). Recent advances in the neuropsychopharmacology of serotonergic hallucinogens. *Behavioural brain research* **277**: 99-120.

Holohean AM, Hackman JC (2004). Mechanisms intrinsic to 5-HT_{2B} receptor-induced potentiation of NMDA receptor responses in frog motoneurons. *Br J Pharmacol* **143**(3): 351-360.

Igata S, Hayashi T, Itoh M, Akasu T, Takano M, Ishimatsu M (2014). Persistent α 1-adrenergic receptor function in the nucleus locus coeruleus causes hyperexcitability in AD/HD model rats. *Journal of Neurophysiology* **111**(4): 777-786.

Jacobsen JPR, Siesser WB, Sachs BD, Peterson S, Cools MJ, Setola V, *et al* (2012). Deficient serotonin neurotransmission and depression-like serotonin biomarker alterations in tryptophan hydroxylase 2 (Tph2) loss-of-function mice. *Mol Psychiatry* **17**(7): 694-704.

Janoshazi A, Deraet M, Callebert J, Setola V, Guenther S, Saubamea B, *et al* (2007). Modified Receptor Internalization upon Co-expression of 5-HT_{1B} Receptor and 5-HT_{2B} Receptors. *Mol Pharmacol* **71**(6): 1463-1474.

Kirby LG, Pernar L, Valentino RJ, Beck SG (2003). Distinguishing characteristics of serotonin and non-serotonin-containing cells in the dorsal raphe nucleus: electrophysiological and immunohistochemical studies. *Neuroscience* **116**(3): 669-683.

Kursar JD, Nelson DL, Wainscott D, Baez M (1994). Molecular cloning, functional expression, and mRNA tissue distribution of the human 5-hydroxytryptamine_{2B} receptor. *Mol Pharmacol* **46**: 227-234.

Leon-Pinzon C, Cercós MG, Noguez P, Trueta C, De-Miguel FF (2014). Exocytosis of serotonin from the neuronal soma is sustained by a serotonin and calcium-dependent feedback loop. *Frontiers in Cellular Neuroscience* **8**: 169.

- Liu R, Jolas T, Aghajanian G (2000). Serotonin 5-HT₂ receptors activate local GABA inhibitory inputs to serotonergic neurons of the dorsal raphe nucleus. *Brain Research* **873**(1): 34-45.
- Loric S, Launay J-M, Colas J-F, Maroteaux L (1992). New mouse 5-HT₂-like receptor: Expression in brain, heart, and intestine. *FEBS L* **312**: 203-207.
- Marinelli S, Schnell SA, Hack SP, Christie MJ, Wessendorf MW, Vaughan CW (2004). Serotonergic and nonserotonergic dorsal raphe neurons are pharmacologically and electrophysiologically heterogeneous. *J Neurophysiol* **92**(6): 3532-3537.
- Maroteaux M, Mameli M (2012). Cocaine evokes projection-specific synaptic plasticity of lateral habenula neurons. *J Neurosci* **32**(36): 12641-12646.
- Muzerelle A, Scotto-Lomassese S, Bernard JF, Soiza-Reilly M, Gaspar P (2016). Conditional anterograde tracing reveals distinct targeting of individual serotonin cell groups (B5-B9) to the forebrain and brainstem. *Brain Structure and Function* **221**(1): 535-561.
- Okaty BW, Freret ME, Rood BD, Brust RD, Hennessy ML, Debairos D, *et al* (2015). Multi-Scale Molecular Deconstruction of the Serotonin Neuron System. *Neuron* **88**(4): 774-791.
- Pitychoutis P, Belmer A, Moutkine I, Adrien J, Maroteaux L (2015). Mice lacking the serotonin Htr2B receptor gene present an antipsychotic-sensitive schizophrenic-like phenotype. *Neuropsychopharmacology* **40**(12): 2764-2773.
- Quérée P, Peters S, Sharp T (2009). Further pharmacological characterization of 5-HT_{2C} receptor agonist-induced inhibition of 5-HT neuronal activity in the dorsal raphe nucleus in vivo. *Br J Pharmacol* **158**(6): 1477-1485.
- Rainer Q, Nguyen HT, Quesseveur G, Gardier AM, David DJ, Guiard BP (2012). Functional status of somatodendritic serotonin 1A autoreceptor after long-term treatment with fluoxetine in a mouse model of anxiety/depression based on repeated corticosterone administration. *Molecular Pharmacology* **81**(2): 106-112.
- Riad M, Garcia S, Watkins KC, Jodoin N, Doucet E, Langlois X, *et al* (2000). Somatodendritic localization of 5-HT_{1A} and preterminal axonal localization of 5-HT_{1B} serotonin receptors in adult rat brain. *The Journal of Comparative Neurology* **417**(2): 181-194.
- Richardson-Jones JW, Craige CP, Guiard BP, Stephen A, Metzger KL, Kung HF, *et al* (2010). 5-HT_{1A} autoreceptor levels determine vulnerability to stress and response to antidepressants. *Neuron* **65**(1): 40-52.
- Scott MM, Wylie CJ, Lerch JK, Murphy R, Lobur K, Herlitze S, *et al* (2005). A genetic approach to access serotonin neurons for in vivo and in vitro studies. *Proc Natl Acad Sci U S A* **102**(45): 16472-16477.
- Teissier A, Chemiakine A, Inbar B, Bagchi S, Ray RS, Palmiter RD, *et al* (2015). Activity of Raphé Serotonergic Neurons Controls Emotional Behaviors. *Cell reports* **13**(9): 1965-1976.
- Urban DJ, Zhu H, Marcinkiewicz CA, Michaelides M, Oshibuchi H, Rhea D, *et al* (2016). Elucidation of The Behavioral Program and Neuronal Network Encoded by Dorsal Raphe Serotonergic Neurons. *Neuropsychopharmacology* **41**(5): 1404-1415.
- Valjent E, Bertran-Gonzalez J, Aubier B, Greengard P, Hervé D, Girault J-A (2010). Mechanisms of locomotor sensitization to drugs of abuse in a two-injection protocol. *Neuropsychopharmacology* **35**(2): 401-415.
- Vandermaelen CP, Aghajanian GK (1983). Electrophysiological and pharmacological characterization of serotonergic dorsal raphe neurons recorded extracellularly and intracellularly in rat brain slices. *Brain Res* **289**(1-2): 109-119.
- Williams JT, Marshall KC (1987). Membrane properties and adrenergic responses in locus coeruleus neurons of young rats. *J Neurosci* **7**(11): 3687-3694.

Figure Legends.

Figure 1**A**

Mode: cell-attached






**B****C**

Figure 1 - Ex-vivo electrophysiological recordings of wildtype *Pet1-GFP* mice- A) Recordings of 5-HT neuron in *ex-vivo* wildtype slices. Cell-attached recordings made in identified *Pet1-GFP* neuron, did not detect firing in the presence of 100 nM phenylephrine (Phe) or of 1 μ M of the 5-HT_{2B}-receptor agonist BW723C86 (top line- 5 cells). In the presence of higher concentration of phenylephrine (300 nM), *Pet1*-positive neuron firing was observed, and BW723C86 (1 μ M) was able to increase this firing rate (8 cells). Representative traces (top) and quantification (bottom left) reveal a significant BW723C86-induced increase in firing frequency (one-way repeated measures-RM-ANOVA; Bar graph and scatter plots and means \pm SEM; Bonferroni post-test, * $P < 0.05$). **B)** Current clamp recordings of *ex-vivo* raphe slices from AAVs injected *Pet1-GFP* mice, overexpressing 5-HT_{2B}-HA receptor in *Pet1*-positive 5-HT neuron. (Left) Sample traces of action potential for a +200 pA current step in two experimental groups. (Right) Quantification of the number of action potentials obtained in function of injected currents showed a significant increase in action potential number in *Pet1*-positive 5-HT neurons of mice overexpressing 5-HT_{2B}-HA receptor, compared to controls (n=20 cells from 3 control mice and 21 cells from four 5-HT_{2B}-HA mice; two-way ANOVA RM, Bonferroni post-test, * $P < 0.05$; Data are means \pm SEM). **C)** Input resistance values. Bar graph and scatter plot for input resistance values showed that input resistance was also increased in mice overexpressing HA-5-HT_{2B} receptor in *Pet1*-positive 5-HT neuron (unpaired t-test, * $P < 0.05$; Data are means \pm SEM).

Figure 2




Figure 2 - *In-vivo* response of wildtype mice to combined 5-HT_{2B} and 5-HT_{1A} receptor agonists - A) *In-vivo* extracellular electrophysiological recordings of putative raphe neurons in anesthetized wildtype mice. (Left) BW723C86 (5 mg/kg sc) injected 20 min before test counteracted the 5-HT_{1A} agonist 8-OHDPAT (0.05 mg/kg; sc) inhibitory cumulative effects to putative 5-HT neuron firing (n=6-5 mice per group; Two-way ANOVA RM followed by Bonferroni's multiple comparisons test *P<0.05; Data are means \pm SEM). (Right) Examples of typical recordings of putative DRN 5-HT neurons obtained in each experimental group. Each arrow represents an injection of 8-OHDPAT (0.05 mg/kg; sc). The injection of the 5-HT_{1A} receptor antagonist WAY100635 (0.3 mg/kg sc) completely reversed the inhibitory effect of 8-OHDPAT. **B) *In-vivo* hypothothermic effects. The 5-HT_{2B}-receptor agonist BW723C86 (5 mg/kg sc) injected 20 min before test was able to counteract 5-HT_{1A} agonist 8-OHDPAT (0.3 mg/kg sc) hypothothermic effects on wildtype mice, (Scattered plot, n=4-4 mice two-way ANOVA RM followed by Bonferroni's multiple comparisons test, *P<0.05).**

Figure 3




Figure 3 - Behavioral response to the 5-HT releaser MDMA in *Htr2b^{5-HTKO}* mice - A) MDMA-induced locomotion. Mice were injected with MDMA (20 mg/kg ip) (arrow) after 30 minutes habituation. A lack of MDMA-induced locomotion was observed in *Htr2b^{5-HTKO}* mice, while control *Htr2b^{lox/lox}* mice showed a clear increase in locomotion. Data between -30 to +60 min were analyzed using two-way ANOVA RM (means \pm SEM, n=16 mice per group) and a Bonferroni post-test was applied on each graph, *P<0.05. **B)** Cumulative MDMA-induced locomotion. Cumulative locomotion during the first hour following MDMA injection showed a significant difference between the two genotypes. Data were analyzed using two-way ANOVA (n=16-16 mice, scattered plot, means \pm SEM). A Bonferroni post-test was also applied on each graph (*P<0.05 *Htr2b^{5-HTKO}* vs. *Htr2b^{lox/lox}*; #P<0.05 MDMA vs. Veh). **C)** Locomotor sensitization by two MDMA injection protocol. The stimulant effect of a challenge dose of MDMA (20 mg/kg ip) 7 days after the first (2nd) was significantly enhanced compared to the first injection in control *Htr2b^{lox/lox}* mice, while it had no effect in *Htr2b^{5-HTKO}* mice. Data were analyzed using two-way ANOVA RM (n=8-8 mice, scattered plot, means \pm SEM). Bonferroni post-test was also applied on each graph (*P<0.05 *Htr2b^{5-HTKO}* vs. *Htr2b^{lox/lox}*; # P<0.05 1st vs. 2nd injection).

Figure 4




Figure 4 - Antidepressant action in *Htr2b^{5-HTKO}* mice - A) Forced swimming test (FST). The time spent immobile in the FST was significantly reduced in control *Htr2b^{lox/lox}*, but not in *Htr2b^{5-HTKO}* mice 30 min after SSRI antidepressant fluoxetine (Flx 3 mg/kg ip) injection (Two-way ANOVA, Bonferroni post-tests * $P < 0.05$; $n = 7-7$ mice, scattered plots with mean \pm SEM). **B)** Neurogenesis in subgranular zone (SGZ) of the hippocampus. Fluoxetine (3 mg/kg/day ip), daily injected for 4 weeks, induced a significant increase in BrdU incorporation in neuron of the SGZ of the hippocampus of control *Htr2b^{lox/lox}* mice, but had no effect in *Htr2b^{5-HTKO}* mice (Two-way ANOVA, Bonferroni post-test; * $P < 0.05$; $n = 7-8$ mice, scattered plots with mean \pm SEM). **C)** SERT expression and function. (Left) SERT expression in conditional *Htr2b^{5-HTKO}* and *Htr2b^{lox/lox}* control mice was evaluated using heterologous competition binding assays of [³H]citalopram on synaptosomes membranes prepared from whole brain. No differences in the affinity (Ki) or expression (Bmax) between *Htr2b^{5-HTKO}* and *Htr2b^{lox/lox}* genotypes were observed. (Right) Saturation isotherms for [³H]5-HT uptake of this synaptosomal preparation were similar in conditional *Htr2b^{5-HTKO}* and *Htr2b^{lox/lox}* control mice. Nonlinear regression analysis did not reveal differences in the Km or Vmax. Shown are representative curves of at least 2 independent experiments performed in duplicates. Individual values are presented.

Figure 5




Figure 5 - Hyposerotonergic phenotype of *Htr2b*^{5-HTKO} mice - **A)** *In-vivo* extracellular recordings of putative DRN 5-HT neurons in anesthetized mice. (Left) The firing frequency of individual putative DRN 5-HT neuron was shifted from high to low firing rates in *Htr2b*^{5-HTKO} mice. (Right) A shift in cumulative distribution was observed with significantly lower firing rate in *Htr2b*^{5-HTKO} mice than in control *Htr2b*^{lox/lox} mice (n=108 and 118 neurons, respectively; Kolmogorov-Smirnov test; *P<0.05). **B)** Head twitch dose-response to DOI. Control *Htr2b*^{lox/lox} and *Htr2b*^{5-HTKO} mice were ip injected with ±DOI (1, 3, and 5 mg/kg ip). Ten min later, the head twitch response was scored for 10 min. DOI-induced head twitch response was significantly increased in conditional *Htr2b*^{5-HTKO} (*P<0.05; multiple t-test) compared to littermate control mice at 5 mg/kg of DOI (Data are presented as scattered plot, n=3-5 mice per group, and means ± SEM). **C)** *In-vivo* hypothermic effects of 8-OHDPAT. The hypothermic response to 8-OHDPAT (0.1 mg/kg sc) was significantly stronger in *Htr2b*^{5-HTKO} mice compared to control *Htr2b*^{lox/lox} mice, (n=3-8 mice, Scattered plot, two-way ANOVA RM followed by Bonferroni's multiple comparisons test, *P<0.05).