



**HAL**  
open science

# Non-wellfounded proof theory for (Kleene+action)(algebras+lattices)

Anupam Das, Damien Pous

► **To cite this version:**

Anupam Das, Damien Pous. Non-wellfounded proof theory for (Kleene+action)(algebras+lattices). 2018. hal-01703942v2

**HAL Id: hal-01703942**

**<https://hal.science/hal-01703942v2>**

Preprint submitted on 27 Apr 2018 (v2), last revised 29 Jun 2018 (v3)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Non-wellfounded proof theory for (Kleene+action)(algebras+lattices)

Anupam Das

University of Copenhagen

Damien Pous

CNRS, Plume team, LIP, ENS Lyon, France

---

## Abstract

We prove cut-elimination for a sequent-style proof system which is sound and complete for the equational theory of Kleene algebra, and where proofs are (potentially) non-wellfounded infinite trees. We extend these results to systems with meets and residuals, capturing ‘star-continuous’ action lattices in a similar way. We recover the equational theory of all action lattices by restricting to regular proofs (with cut)—those proofs that are unfoldings of finite graphs.

**2012 ACM Subject Classification** Proof theory; Algebra

**Keywords and phrases** Kleene algebra; proof theory; sequent system; non-wellfounded proofs


## 1 Introduction

The axioms of *Kleene algebras* are sound and complete for the theory of regular expressions under language equivalence [21, 26, 4]. As a consequence, the equational theory of Kleene algebras is decidable (in fact PSPACE-complete). Models of these axioms of particular interest include formal languages and binary relations. For binary relations, the Kleene star is interpreted as reflexive transitive closure, whence the axioms of Kleene algebra make it possible to reason abstractly about program correctness [23, 24, 3, 18, 1]. The aforementioned decidability result moreover makes it possible to automate interactive proofs [5, 25, 29].

There are however important extensions of Kleene algebras which are not yet fully understood. These include *action algebras* [30], where two ‘residual’ operations are added, *Kleene lattices*, where a ‘meet’ operation is added, and *action lattices* [22], where all three operations are added. Pratt introduced residuals in order to *internalise* the induction rules of the Kleene star, as we explain later; they allow us to express properties of relations such as well-foundedness in a purely algebraic way [11]. Kozen added the meet operation to action algebra to obtain a structure closed under taking matrices. In the context of program verification, meets are useful since they allow us to express conjunctions of local specifications.

Unfortunately, the decidability of the three corresponding equational theories is still open, and there is no known notion of ‘free model’ for them that is analogous to the rational languages for Kleene algebra. In this paper, we explore a proof-theoretic approach to such questions: we provide sequent calculi that capture these theories which we show admit a form of cut-elimination. Although this does not (yet) give us decidability, it does improve our understanding of these theories:

- we obtain a computational interpretation of proofs of inequalities in our systems as program transformers, which could prove useful to describe free models;
- we recover two conservativity results: action lattices are conservative over (star-continuous) Kleene lattices and action algebra, thanks to the sub-formula property; (these results are also implied by [28]).



■ **Figure 1** Context and contributions for Kleene algebra and action lattices.

- we obtain structural properties, e.g., as Whitman did when he proved cut-elimination for the theory of lattices, which we aim to exploit in consequent research.

We first focus on pure Kleene algebra, which is easier to handle and enables a simpler presentation. Being a well-established theory, we are able to relate our results to existing ones in the literature, identifying which issues become relevant when moving to extensions.

### Kleene algebra

In our sequent system, called LKA, proofs are finitely branching, but possibly infinitely deep (i.e. not wellfounded). To prevent fallacious reasoning, we give a simple validity criterion for proofs with cut, and prove that the corresponding system admits cut-elimination. The difficulty in the presence of infinitely deep proofs consists in proving that cut-elimination is productive; we do so by using the natural interpretation of regular expressions as data types for parse-trees [14], and by giving an interpretation of proofs as parse-tree transformers. Such an idea already appears in [17] but in a simpler setting, for a finitary natural deduction system rather than for a non-wellfounded sequent calculus.

The results we prove about LKA are summarised in Fig. 1(left). In addition to cut-elimination (Thm. 13), we prove that the system is sound for all *star-continuous* Kleene algebras (Thm. 10), and conversely, that it is complete w.r.t. the language theoretic interpretation of regular expressions (Thm. 11). We actually refine this latter result by showing that every proof from Kleene algebra axioms can be translated into a *regular* proof with cut (Thm. 12), i.e., a proof with cut which is the unfolding of a finite graph. Note, however, that regularity is not preserved by cut-elimination: the class of cut-free regular proofs in LKA is incomplete w.r.t. Kleene algebra.

### Action algebras, Kleene lattices, and action lattices

Despite its finite quasi-equational presentation, the equational theory of Kleene algebra is not finitely based: Redko proved that any finite set of equational axioms must be incomplete [31]. However, by adding two binary operations to the signature, Pratt showed how to obtain a finitely based extension which is conservative over the equational theory of Kleene algebras [30]. These two operations, called *left residual* ( $\backslash$ ) and *right residual* ( $/$ ), are ‘ad-joint’ to sequential composition and, as we mentioned, such structures are called *action algebras*. Kozen then proposed *action lattices* [22], where the signature is extended further to include a binary meet operation ( $\cap$ ). We call *Kleene lattices* the structures consisting of Kleene algebra extended just with meets.

While both action algebras and action lattices are finitely based and conservatively extend Kleene algebra, they bring some difficulties. By definition, their equational theories are at most  $\Sigma_1^0$ , so that they must differ from their star-continuous variants which are  $\Pi_1^0$ -complete [7, 28]. (Buszkowski proved the lower bound and Palka proved the upper bound.) In contrast, by Kozen’s completeness result we have that Kleene algebra and star-continuous Kleene algebra give rise to the same equational theory, which is PSPACE-complete. This matter remains open for Kleene lattices since Buszkowski’s lower bound does not apply.

Residuals and meets naturally correspond to linear implication and additive conjunction [19, 28], from (non-commutative intuitionistic) linear logic [16]. They are also essential connectives in the Lambek-calculus and related substructural logics [27]. We extend LKA accordingly into a system LAL and obtain the results summarised in Fig. 1(right): LAL is complete for star continuous action lattices (Thm. 33); it still admits cut-elimination (Thm. 35); thus it is also sound w.r.t. star continuous action lattices (Thms. 30). Furthermore we are able to show that its regular fragment with cut is in fact sound and complete for all action lattices (Thm. 31); this somewhat surprising result gives us a nontrivial yet finite proof theoretic representation of the theory of action lattices. The proof reasons inductively on the cycle structure of such regular proofs, and we crucially exploit the availability of both residuals and meets: for action algebra and Kleene lattices, it remains open whether the corresponding regular fragments with cut are sound.

Thms 30, 32, and 35 are proved by extending the proofs of Thms. 10, 12, and 13 to deal with the additional connectives. Amongst those, cut-elimination is the most delicate extension, relying on higher types to interpret residuals, and proving that LAL proofs still yield terminating programs in such a setting. Thm. 11 cannot be extended directly, due to the lack of a free model analogous to the regular languages for Kleene algebra when adding residuals or meet. This is why we instead rely on cut-elimination for completeness.

As explained above, while all notions are equivalent in the case of Kleene algebra (Fig. 1(left)), complexity arguments make it possible to separate the lower and upper parts of Fig. 1(right), except for Kleene lattices. Whether the upper part is decidable remains open, but it is interesting to note that we managed to characterise action lattices in such a way that the non-regular/regular distinction at the proof-level corresponds precisely to the difference between the star continuous and general cases, respectively. One potentially fruitful direction towards the decidability of action lattices is to characterise the image of regular proofs under cut-elimination. We aim to explore this possibility in future work.

## Related work

We briefly discussed the cut-free variant of the system LKA in [10] (with a simpler validity criterion), observing that its regular fragment is incomplete (due to the absence of cut). Our main contribution was a variant of it based on ‘hypersequents’, HKA, whose regular fragment is sound and complete without cut, and admits a PSPACE proof search procedure.

Palka proposed a sequent system for star continuous action lattices, for which she proved cut-elimination [28]. Its non-star rules are precisely those of LKA, but the system is well-founded and relies on an ‘omega-rule’ for Kleene star with infinitely many premisses, in the traditional school of infinitary proof theory, cf. [32]. Such an approach does not admit a notion of finite proof analogous to our regular proofs, corresponding to the upper parts of Fig. 1. Wurm also proposed a (finite, and thus wellfounded) sequent system for Kleene algebra [33]. Unfortunately his cut-admissibility theorem does not hold—see [10].

The normalisation theory of *linear logic* with (least and greatest) fixed point operators has been studied in [13] and, more comprehensively, in [12]. While the latter is a rather

general framework, its exposition still differs significantly from the current work for various reasons. One immediate difference is that their setting is commutative while ours is non-commutative, and so those results are not directly applicable. A more important difference is that they do not have any atoms in their language, reasoning only on closed formulae. This is rather significant from the point of view of normalisation, since the convergence of cut-elimination becomes more complicated in presence of atoms. The argument we give in Sect. 4 uses different ideas that are closely related to the language-based models of our algebras and the natural interpretation of language inclusions as programs [17]. A *game semantics* approach to cut-elimination for non-wellfounded proofs is given in [8], though in that work only finitely many cuts in a proof are considered and so it does not seem sufficient to handle the star rules in this work.

## 2 Preliminaries on Kleene algebra and extensions

Let  $A$  be a finite *alphabet*. *Regular expressions* [20] are generated as follows:

$$e, f ::= e \cdot e \mid e + e \mid e^* \mid 1 \mid 0 \mid a \in A$$

Sometimes we may simply write  $ef$  instead of  $e \cdot f$ . Each expression  $e$  generates a rational language  $\mathcal{L}(e) \subseteq A^*$ , defined in the usual way.

A *Kleene algebra* is a tuple  $(K, 0, 1, +, \cdot, *)$  where  $(K, 0, 1, +, \cdot)$  is an idempotent semiring and where the following properties hold, where  $x \leq y$  is a shorthand for  $x + y = y$ .

$$1 + xx^* \leq x^* \quad \text{if } xy \leq y \text{ then } x^*y \leq y \quad \text{if } yx \leq y \text{ then } yx^* \leq y \quad (1)$$

There are several equivalent variants of this definition [9]. Intuitively we have that  $x^*y$  (resp.,  $yx^*$ ) is the least fixpoint of the function  $z \mapsto y + xz$  (resp.,  $z \mapsto y + zx$ ). We write  $\text{KA} \models e \leq f$  if the inequality  $e \leq f$  holds universally in all Kleene algebras—or, equivalently, if it is derivable from the axioms of Kleene algebra. Kozen [21] and KroB [26] showed that this axiomatisation is complete for language inclusions, corresponding to the right-to-left implication in the following characterisation (the other direction is routine).

► **Theorem 1** ([21, 26]).  $\text{KA} \models e \leq f$  if and only if  $\mathcal{L}(e) \leq \mathcal{L}(f)$ .

A Kleene algebra is *star-continuous* if for all elements  $x, y, z$ ,  $xy^*z$  is the least upper bound of the sequence  $(xy^iz)_{i \in \mathbb{N}}$ , where  $y^0 = 1$  and  $y^{i+1} = yy^i$ . In presence of the other laws, star-continuity is equivalent to the following condition:

$$\forall xyz, (\forall i \in \mathbb{N}, xy^iz \leq t) \Rightarrow xy^*z \leq t.$$

We write  $\text{KA}^* \models e = f$  when the equality  $e = f$  holds in all star-continuous Kleene algebras. Formal languages form a star-continuous Kleene algebra, and so by completeness of Kleene algebra w.r.t. rational languages, we have  $\text{KA}^* \models e = f$  iff  $\text{KA} \models e = f$ ; this is the triangle on the left in Fig. 1.

An *action lattice* is a Kleene algebra with three additional binary operations, left and right *residuals* ( $\backslash, /$ ), and *meet* ( $\cap$ ) defined by the following equivalences:

$$\forall xyz, y \leq x \backslash z \Leftrightarrow xy \leq z \Leftrightarrow x \leq z / y \quad \forall xyz, z \leq x \cap y \Leftrightarrow z \leq x \wedge z \leq y$$

An *action algebra* is a Kleene algebra with residuals, a *Kleene lattice* is a Kleene algebra with meets. We extend regular expressions accordingly, writing  $\text{AL} \models e \leq f$  when the inequation  $e \leq f$  holds in all action lattices, and  $\text{AL}^* \models e \leq f$  when it holds in all star continuous action lattices. Note that while rational languages are closed under residuals and intersection, thus forming an action lattice, they are not the ‘free’ one: Thm. 1 fails. The equational theories generated by all action lattices and by the star-continuous ones actually differ, cf. [7, 28].

$$\begin{array}{c}
\text{cut} \frac{\Delta \rightarrow e \quad \Gamma, e, \Sigma \rightarrow f}{\Gamma, \Delta, \Sigma \rightarrow f} \quad \text{id} \frac{}{e \rightarrow e} \quad 0\text{-l} \frac{}{\Gamma, 0, \Delta \rightarrow e} \quad 1\text{-l} \frac{\Gamma, \Delta \rightarrow e}{\Gamma, 1, \Delta \rightarrow e} \quad 1\text{-r} \frac{}{\rightarrow 1} \\
\text{-l} \frac{\Gamma, e, f, \Delta \rightarrow g}{\Gamma, e \cdot f, \Delta \rightarrow g} \quad \text{+l} \frac{\Gamma, e, \Delta \rightarrow g \quad \Gamma, f, \Delta \rightarrow g}{\Gamma, e + f, \Delta \rightarrow g} \quad \text{*l} \frac{\Gamma, \Delta \rightarrow f \quad \Gamma, e, e^*, \Delta \rightarrow f}{\Gamma, e^*, \Delta \rightarrow f} \\
\text{-r} \frac{\Gamma \rightarrow e \quad \Delta \rightarrow f}{\Gamma, \Delta \rightarrow e \cdot f} \quad \text{+r}_i \frac{\Gamma \rightarrow e_i}{\Gamma \rightarrow e_1 + e_2} \quad i \in \{1, 2\} \quad \text{*r}_1 \frac{}{\rightarrow e^*} \quad \text{*r}_2 \frac{\Gamma \rightarrow e \quad \Delta \rightarrow e^*}{\Gamma, \Delta \rightarrow e^*}
\end{array}$$

■ **Figure 2** The rules of LKA.

### 3 The sequent system LKA

A *sequent* is an expression  $\Gamma \rightarrow e$ , where  $\Gamma$  is a list of regular expressions and  $e$  is a regular expression. For such a sequent we refer to  $\Gamma$  as the *antecedent* and  $e$  as the *succedent*, or simply the ‘left’ and ‘right’ hand sides, respectively. We say that a sequent  $e_1, \dots, e_n \rightarrow e$  is *valid* if  $\text{KA}^* \models e_1 \cdots e_n \leq e$ . i.e., the comma is interpreted as sequential composition, and the sequent arrow as containment. We may refer to expressions as ‘formulae’ when it is more natural from a proof theoretic perspective, e.g. ‘subformula’ or ‘principal formula’.

The rules of LKA are given in Fig. 2. We call  $\text{LKA}^-$  the subset of LKA where the *cut* rule is omitted (which corresponds to the system called LKA in [10]). Leaving the  $*$ -rules aside, these rules are those of the non-commutative variant of intuitionistic linear logic [16], restricted to the following connectives: multiplicative conjunction ( $\cdot$ ), additive disjunction ( $+$ ) and additive falsity ( $0$ ) (for which there is no right rule). The rules for Kleene star can be understood as those arising from the characterisation of  $e^*$  as a fixed point:  $e^* = \mu x.(1 + ex)$ . In contrast, Palka [28] follows the alternative interpretation of Kleene star as an infinite sum,  $e^* = \sum_i e^i$ , whence her left rule for Kleene star with infinitely many premisses, and the infinitely many right rules she uses for this operation.

As previously mentioned, we consider infinitely deep proofs, so it is necessary to impose a validity criterion to ensure that derivations remain sound.

► **Definition 2.** A (binary, possibly infinite) *tree* is a prefix-closed subset of  $\{0, 1\}^*$ ; elements of  $\{0, 1\}^*$  are called *nodes*. A *preproof* is a labelling  $\pi$  of a tree by sequents such that, for every node  $v$  with children  $v_1, \dots, v_n$  ( $n = 0, 1, 2$ ), the expression  $\frac{\pi(v_1) \cdots \pi(v_n)}{\pi(v)}$  is an instance of an LKA rule. Given a node  $v$  in a preproof  $\pi$ , we write  $\pi_v$  for the sub-preproof rooted at  $v$ , defined by  $\pi_v(w) = \pi(vw)$ . A preproof is *regular* if it has finitely many distinct subtrees, i.e. it can be expressed as the infinite unfolding of a finite graph. A preproof is *cut-free* if it does not use the *cut*-rule.

We will use standard proof theoretic terminology about *principal formulas* and *ancestry* in proofs, e.g. from [6] (see App. A for further details). The notion of validity below is similar to [12], adapted to our setting.

► **Definition 3.** A *thread* is a maximal path through the graph of (immediate) ancestry in a preproof. By definition it must start at a conclusion formula or at a cut formula and it only goes upwards. A thread is *valid* if it is principal for a  $*$ -l step infinitely often. A preproof is *valid* if every infinite branch eventually has a valid thread. A *proof* is a valid preproof. We write  $\text{LKA} \vdash^\infty \Gamma \rightarrow e$  if the sequent  $\Gamma \rightarrow e$  admits a proof,  $\text{LKA} \vdash^\omega \Gamma \rightarrow e$  if it admits a regular proof, and  $\text{LKA}^- \vdash^\infty \Gamma \rightarrow e$  if it admits a cut-free proof.

Notice that every valid thread eventually follows a unique (star) formula, by the subformula property. Let us consider some examples of (pre)proofs. In all cases, we will use the symbol  $\bullet$  to indicate circularities, colours to mark some of the threads, and double lines to denote finite derivations.

► **Example 4.** Here is a regular and cut-free proof of  $(b + c)^* \rightarrow (c + b)^*$ :

$$\frac{\frac{\frac{\frac{\vdots}{(b+c)^* \rightarrow (c+b)^*} \bullet}{b+c \rightarrow c+b} \quad \frac{\vdots}{(b+c)^* \rightarrow (c+b)^*} \bullet}{\frac{\frac{\vdots}{(b+c)^* \rightarrow (c+b)^*} \bullet}{b+c, (b+c)^* \rightarrow (c+b)^*} \bullet} \quad \frac{\vdots}{(b+c)^* \rightarrow (c+b)^*} \bullet}{\frac{\vdots}{(b+c)^* \rightarrow (c+b)^*} \bullet} \bullet$$

► **Example 5 (Atomicity of identity).** As in many common sequent systems, initial identity steps can be reduced to atomic form, although for this we crucially rely on access to non-wellfounded (yet regular) proofs. As usual, we proceed by induction on the size of an identity step, whence the crucial case is for the Kleene star,

$$\frac{\frac{\frac{\frac{\text{IH}}{e \rightarrow e} \quad \frac{\vdots}{e^* \rightarrow e^*} \bullet}{e, e^* \rightarrow e^*} \bullet}{\frac{\vdots}{e^* \rightarrow e^*} \bullet} \bullet}{\frac{\vdots}{e^* \rightarrow e^*} \bullet} \bullet$$

where the derivation marked *IH* is obtained by the inductive hypothesis.

Note that while  $\text{LKA}^-$  satisfies the subformula property, the size and number of sequents occurring in a cut-free proof is not a priori bounded, due to the  $*-l$  rule:

► **Example 6 (A non-regular proof).** The only cut-free proof of the sequent  $a, a^* \rightarrow a^*a$  is the one on the left below:

$$\frac{\frac{\frac{\frac{\vdots}{a, a \rightarrow a^*a} \quad \frac{\vdots}{a, a, a, a^* \rightarrow a^*a}}{\frac{\vdots}{a, a, a^* \rightarrow a^*a}} \bullet}{\frac{\vdots}{a, a^* \rightarrow a^*a}} \bullet}{\frac{\vdots}{a, a^* \rightarrow a^*a}} \bullet \quad \frac{\frac{\frac{\frac{\vdots}{a, a^* \rightarrow a^*a} \bullet \quad \frac{\vdots}{a, a^*a \rightarrow a^*a}}{\frac{\vdots}{a, a, a^* \rightarrow a^*a}} \bullet}{\frac{\vdots}{a, a^* \rightarrow a^*a}} \bullet}{\frac{\vdots}{a, a^* \rightarrow a^*a}} \bullet$$

This proof contains all sequents of the form  $a, \dots, a, a^* \rightarrow a^*a$ , whence non-regularity. A regular proof with cuts is given on the right; see [10] for more details on the lack of regularity in  $\text{LKA}^-$  and how to recover regularity in a cut-free setting, using ‘hypersequents’.

► **Example 7 (Two invalid preproofs).** The following preproofs are not valid; they derive invalid sequents.

$$\frac{\frac{\frac{\vdots}{a \rightarrow 1^*} \bullet}{\frac{\vdots}{a \rightarrow 1^*} \bullet} \bullet}{\frac{\vdots}{a \rightarrow 1^*} \bullet} \bullet \quad \frac{\frac{\frac{\frac{\frac{\text{id}}{a \rightarrow a} \quad \frac{\text{id}}{a^* \rightarrow a^*}}{\frac{\vdots}{a, a^* \rightarrow a^*} \bullet} \quad \frac{\vdots}{a^* \rightarrow b^*} \bullet}{\frac{\vdots}{a, a^* \rightarrow b^*} \bullet} \bullet}{\frac{\vdots}{a^* \rightarrow b^*} \bullet} \bullet$$

The left-hand preproof is cut-free and infinite; since it does not contain any  $*-l$ -rule, it cannot be valid. On the right-hand side, the principal formula of the  $*-l$ -rule is the cut formula of the *cut*-rule so that the only infinite thread is the one along the occurrences of  $b^*$ , and this formula is never principal for a  $*-l$  step.

The notion of validity we use here actually generalises the notion of *fairness* we used in [10], where we were working only with cut-free preproofs:

► **Proposition 8.** *A cut-free preproof is valid if and only if it is fair for  $*-l$ , i.e. every infinite branch contains infinitely many occurrences of  $*-l$ .*

**Proof.** The left-right implication is immediate. Conversely, every infinite path in a fair cut-free preproof has infinitely many  $*-l$  steps, and their principal formulae are finitely many by the subformula property. One can thus extract a valid thread. ◀

An alternative criterion for cut-free preproofs is obtained as follows:

► **Proposition 9.** *A cut-free preproof is valid if and only if it has no infinite branch with a tail of only  $*-r_2$ -steps.*

**Proof.** Define the ‘weight’ of a sequent to be the multiset of its formulae, ordered by the subformula relation. This measure strictly decreases when reading  $LKA^-$  rules bottom-up, except for the right premisses of rules  $*-l$  and  $*-r_2$ ; for the latter, it either remains unchanged (when  $\Gamma$  is empty) or it strictly decreases. Thus every infinite branch of a cut-free preproof either contains infinitely many  $*-l$  steps, or eventually contains only  $*-r_2$  steps. ◀

Observe that the proof on the left in Ex. 7 does not satisfy this condition.

The cut-free system  $LKA^-$  is sound and complete for Kleene algebras. Thanks to the completeness theorem for Kleene algebras, Thm. 1, it suffices to prove soundness with respect to star-continuous Kleene algebra.

► **Theorem 10 (Soundness).** *If  $LKA^- \vdash^\infty e_1, \dots, e_n \rightarrow e$ , then  $KA^* \models e_1 \cdots e_n \leq e$ .*

**Proof sketch.** Finite derivations in  $LKA$  are easily proven sound; the difficulty consists in handling infinite derivations. The proof follows the same strategy as the one given in [10] for the hypersequent system  $HKA$ . We proceed by induction on the multiset of the formulae in the conclusion, relying on star-continuity and the following lemma for  $*-l$ -steps:

$$\text{If } LKA^- \vdash^\infty \Gamma, e^*, \Delta \rightarrow f \text{ then, for all } n \in \mathbb{N}, LKA^- \vdash^\infty \Gamma, e^n, \Delta \rightarrow f.^1$$

See App. B for a detailed proof. ◀

For completeness of  $LKA^-$ , we can get a direct proof by starting from the free model of rational languages (Fig. 1). This strategy is no longer possible for Kleene lattices, action algebras and action lattices, for which we will need to go through cut-elimination.

► **Theorem 11 (Completeness).** *If  $\mathcal{L}(e_1 \cdots e_n) \subseteq \mathcal{L}(e)$  then  $LKA^- \vdash^\infty e_1, \dots, e_n \rightarrow e$ .*

**Proof sketch.** Again this can be proved like in [10] for  $HKA$ : all left rules of  $LKA^-$  are invertible so that they can be applied greedily; doing so, one obtains an infinite tree whose leaves are sequents of the shape  $a_1, \dots, a_k \rightarrow e$ , with  $k \geq 0$ , where  $a_1 \dots a_k$  is a word in  $\mathcal{L}(e_1 \cdots e_n)$  and thus in  $\mathcal{L}(e)$  by assumption. Those leaves can be replaced by finite derivations using only right rules. See App. C for a detailed proof. ◀

The previous proof builds infinite and non-regular derivations whenever the language of the starting antecedent is infinite. For instance, it would yield the proof given on the left in Ex. 6. By using a different technique, we show in the following theorem, that we can get regular proofs if we allow the cut-rule.

► **Theorem 12 (Regular completeness).** *If  $KA \models e \leq f$  then  $LKA \vdash^\omega e \rightarrow f$ .*

<sup>1</sup> Where  $e^0 = 1$  and  $e^{n+1} = e \cdot e^n$ .



**Proof.** We prove the statement for equalities. Consider the relation  $\equiv$  defined by  $e \equiv f$  if  $\text{LKA} \vdash^\omega e \rightarrow f$  and  $\text{LKA} \vdash^\omega f \rightarrow e$ . This relation is an equivalence on regular expressions thanks to the cut rule, and it is easily shown to be preserved by all contexts (i.e. it is a congruence). Also remark that we have  $e + f \equiv f$  iff  $\text{LKA} \vdash^\omega e \rightarrow f$ , thanks to the cut-rule and the rules about sum. It then suffices to show that regular expressions quotiented by  $\equiv$  form a Kleene algebra. The (in)equational axioms defining KA can be proved by finite derivations. The only difficulty is in dealing with the two implications from the definition of Kleene algebra (1). We implement them as follows:

$$\begin{array}{c}
 \vdots \\
 \text{*} \text{-} l \frac{}{e^*, f \rightarrow f} \bullet \\
 \text{id} \frac{}{f \rightarrow f} \\
 \text{cut} \frac{}{e, e^*, f \rightarrow f} \\
 \text{*} \text{-} l \frac{}{e^*, f \rightarrow f} \bullet
 \end{array}
 \quad
 \begin{array}{c}
 \text{IH} \\
 \text{cut} \frac{}{f, e \rightarrow f} \\
 \text{id} \frac{}{f \rightarrow f} \\
 \text{cut} \frac{}{f, e, e^* \rightarrow f} \\
 \text{*} \text{-} l \frac{}{f, e^* \rightarrow f} \bullet
 \end{array}$$

where the derivations marked *IH* are obtained from the inductive hypothesis. The preproofs we construct in this way are valid and regular, by inspection. In particular, the only infinite branch not in *IH* in the above derivations has a valid thread on  $e^*$ , coloured in green.  $\blacktriangleleft$

Note the asymmetry when we interpret the two implications: the premisses of the cut rule are swapped when we move from one to the other. This asymmetry comes from the fact that we have a single left rule for Kleene star, which unfolds the star from the left.

#### 4 Cut-elimination for LKA

This section is devoted to proving the following cut-elimination theorem.

► **Theorem 13.** *If  $\text{LKA} \vdash^\infty \Gamma \rightarrow e$  then  $\text{LKA}^- \vdash^\infty \Gamma \rightarrow e$ .*

Combined with Thm. 10, it establishes the soundness of our criterion for proofs with cuts. This serves as a ‘warm-up’ for the analogous result for the extended system (Sect. 6), which is obtained using the same template.

We show that proofs can be considered as certain *transducers*, transforming *parse-trees* of input words of languages computed by terms. We design them so that a given computation only explores a finite prefix of the proof, which we call the *head*. We then prove that cut-reductions, restricted to the head of a proof, preserve these computations, always terminate, and eventually produce some non-cut rules. We can then repeatedly apply this procedure to remove all cuts from an infinite proof, in a productive way.

#### 4.1 Programs from proofs

We first define programs and their reduction semantics, based on which we prove cut-elimination, in Sect. 4.2. We fix in this section a (valid) LKA proof  $\pi$  and we let  $v$  range over its nodes, which we recall are elements of  $\{0, 1\}^*$ , cf. Dfn. 2.

► **Definition 14 (Programs).** *Programs* are defined by the following syntax, where  $x$  ranges over a countable set of *variables*, and  $i$  ranges over  $\{1, 2\}$ .

$$M, N ::= x \mid \star \mid \langle M, N \rangle \mid \text{in}_i M \mid [] \mid M :: N \mid v[\vec{M}]$$

Intuitively, programs compute parse-trees for words belonging to the language of an expression. Given a node  $v$  of  $\pi$  such that  $\pi(v) = \Gamma \rightarrow e$ , the last entry corresponds to the

$$\begin{array}{c}
\frac{}{x : e \vdash x : e} \quad \frac{}{\vdash \star : 1} \quad \frac{E \vdash M : e \quad F \vdash N : f}{E, F \vdash \langle M, N \rangle : ef} \quad \frac{E \vdash M : e_i}{E \vdash \text{in}_i M : e_1 + e_2} \quad \frac{}{\vdash [] : e^*} \\
\\
\frac{E \vdash M : e \quad E' \vdash N : e^*}{E, E' \vdash M :: N : e^*} \quad \frac{\forall i, E_i \vdash M_i : e_i \quad \pi(v) = e_1, \dots, e_n \rightarrow f}{v, E_1, \dots, E_n \vdash v[\vec{M}] : f}
\end{array}$$

■ **Figure 3** Typing rules for programs.

application of the subproof  $\pi_v$ , rooted at  $v$ , to a list  $\vec{M}$  of programs for the antecedent ( $\Gamma$ ); it should eventually return a parse-tree for the succedent ( $e$ ). This is formalised using the following notion of types.

► **Definition 15** (Typing environment). A *typing environment*, written  $E$ , is a list of pairs of variables and expressions, written  $x : e$ , together with a finite antichain of nodes: for any two nodes  $v$  and  $w$  in the antichain,  $v$  is not a prefix of  $w$ . We write  $E, F$  for the concatenation of two typing environments, which is defined only when this antichain condition on nodes is preserved.

Intuitively, typing environments keep track of which variables and proof nodes are used in a program, to impose linearity constraints; those constraints become crucial when we add residuals and meets, in Sect. 5.

► **Definition 16** (Types). A program  $M$  has type  $e$  in an environment  $E$ , written  $E \vdash M : e$  if this judgement can be derived from the rules in Fig. 3.

► **Example 17.** With the proof from Ex. 4, letting  $\varepsilon$  denote the root node, we have:

$$\begin{array}{l}
\varepsilon, x : b, y : c \vdash \varepsilon[\text{in}_0 x :: \text{in}_1 y :: []] : (c + b)^* \\
\varepsilon, z : b + c, z' : b + c, q : (b + c)^* \vdash \varepsilon[z :: z' :: q] : (c + b)^*
\end{array}$$

► **Observation 18.** Let  $x_1, \dots, x_n$  be variables. We have  $a_1 \dots a_n \in \mathcal{L}(e)$  iff there exists a program  $M$  such that  $x_1 : a_1, \dots, x_n : a_n \vdash M : e$ . This (unused) observation has no counterpart when considering extensions of Kleene algebra, where there is no longer an appropriate notion of ‘language’ for expressions that constitutes a free model.

► **Definition 19** (Reduction). *Reduction*, written  $\rightsquigarrow$ , is the closure under all contexts of the following rules defined by case analysis on the last step of the subproof  $\pi_v$  rooted at  $v$ . These rules are written concisely for lack of space; in each case,  $v0$  and  $v1$  are the nodes of the premisses, when they exist. We moreover assume that the sizes of the vectors match those that arise from the various rules. See App. D for an extensive definition.

$$\begin{array}{ll}
id : v[M] \rightsquigarrow M & cut : v[\vec{M}, \vec{N}, \vec{P}] \rightsquigarrow v1[\vec{M}, v0[\vec{N}], \vec{P}] \\
1-l : v[\vec{M}, \star, \vec{N}] \rightsquigarrow v0[\vec{M}, \vec{N}] & 1-r : v[] \rightsquigarrow \star \\
-l : v[\vec{M}, \langle M, N \rangle, \vec{N}] \rightsquigarrow v0[\vec{M}, M, N, \vec{N}] & -r : v[\vec{M}, \vec{N}] \rightsquigarrow \langle v0[\vec{M}], v1[\vec{N}] \rangle \\
+l : v[\vec{M}, \text{in}_i M, \vec{N}] \rightsquigarrow vi[\vec{M}, M, \vec{N}] & +-r_i : v[\vec{M}] \rightsquigarrow \text{in}_i(v0[\vec{M}]) \\
*-l : v[\vec{M}, [], \vec{N}] \rightsquigarrow v0[\vec{M}, \vec{N}] \text{ and} & *-r_1 : v[] \rightsquigarrow [] \\
v[\vec{M}, M :: N, \vec{N}] \rightsquigarrow v1[\vec{M}, M, N, \vec{N}] & *-r_2 : v[\vec{M}, \vec{N}] \rightsquigarrow v0[\vec{M}] :: v1[\vec{N}]
\end{array}$$

When useful, we write  $\rightsquigarrow_{cut}$  indicate a reduction according to the *cut* rule above.

► **Example 20.** Continuing with the proof from Ex. 4 we have the following complete reductions. The second program still contains calls to proofs in the end because the inputs were under-specified.

$$\begin{array}{ll}
\varepsilon[\text{in}_0x :: \text{in}_1y :: []] & \varepsilon[z :: z' :: q] \\
\rightsquigarrow 1[\text{in}_0x, \text{in}_1y :: []] & \rightsquigarrow 1[z, z' :: q] \\
\rightsquigarrow 10[\text{in}_0x] :: 11[\text{in}_1y :: []] & \rightsquigarrow 10[z] :: 11[z' :: q] \\
\rightsquigarrow 100[x] :: 11[\text{in}_1y :: []] & \rightsquigarrow 10[z] :: 111[z', q] \\
\rightsquigarrow \text{in}_1x :: 11[\text{in}_1y :: []] & \rightsquigarrow 10[z] :: 1110[z'] :: 1111[q] \\
\rightsquigarrow \text{in}_1x :: 111[\text{in}_1y, []] & \\
\rightsquigarrow \text{in}_1x :: 1110[\text{in}_1y] :: 1111[[]] & \\
\rightsquigarrow^4 \text{in}_1x :: \text{in}_0y :: [] & 
\end{array}$$

As one might expect, we have subject reduction. We need the following notion of *extension* to state it properly.

► **Definition 21 (Extension).** Given two typing environments  $E, E'$ , we say that  $E'$  *extends*  $E$  if  $E$  and  $E'$  coincide after removing all nodes, and if all nodes in  $E'$  are either already in  $E$ , or are immediate successors of some nodes in  $E$ .

► **Proposition 22 (Subject reduction).** *If  $E \vdash M : e$  and  $M \rightsquigarrow M'$ , then  $E' \vdash M' : e$  for some environment  $E'$  extending  $E$ .*

For instance, along the reductions on the left in Ex. 20, the antichain part of the typing environment evolves as follows:  $\{\varepsilon\}, \{1\}, \{10, 11\}, \{100, 11\}, \{11\}, \{111\}, \{1110, 1111\}, \emptyset$ .

Our objective now is to prove that well-typed programs terminate. For the sake of simplicity, we work in the sequel with the ‘leftmost innermost’ strategy: a redex  $v[\bar{M}]$  is fired only when the programs in  $\bar{M}$  are irreducible and there are no other redexes to its left.

► **Definition 23 (Runs).** The *run* of a program  $M$  is the sequence of nodes corresponding to the redexes fired during the (potentially infinite) innermost leftmost reduction of  $M$ .

► **Lemma 24.** *If  $E \vdash M : e$  then every node  $w$  appears at most once in the run of  $M$ ; in this case we have that  $w = uv$  for some nodes  $u, v$  with  $u$  in  $E$  and, for every prefix  $v'$  of  $v$ ,  $uv'$  appears in the run of  $M$  before  $w$ .*

**Proof.** These are immediate consequences of Prop. 22. ◀

In particular, the run of a well-typed program has finitely many connected components. We finally obtain that well typed programs terminate, thanks to the validity criterion.

► **Proposition 25.** *If  $E \vdash M : e$ , then the run of  $M$  is finite.*

**Proof.** Suppose the run of  $M$  is infinite. Then by Lem. 24 and König’s Lemma one can extract an infinite branch of  $\pi$  which is contained in the run. By validity, this branch must eventually have a thread along a star formula  $f^*$  which is infinitely often principal. By analysis of the reduction rules, and thanks to the innermost strategy, we may find an infinite sequence of programs of type  $f^*$  whose sizes are strictly decreasing, which is impossible. ◀

## 4.2 Cut reduction

Our cut-elimination argument is driven by a standard set of cut reduction rules, which we do not have space to present in the main text. These include key and commutative cases, as

usual, and are fully presented in App. F. To produce an infinite cut-free proof, we must show that we may produce proofs with arbitrarily large cut-free prefixes in a continuous manner. The main difficulty is to show that such a procedure is *productive*, i.e., eventually produces non-cut steps. To this end, we use the previous notion of ‘run’ to drive cut-reductions.

► **Definition 26 (Head).** Let  $\pi$  be a proof of  $\Gamma \rightarrow e$ . The *head* of  $\pi$ , written  $hd(\pi)$ , is the run of the program  $\varepsilon[\vec{x}]$  in  $\pi$ , where  $\vec{x}$  is a list of variables of the same length as  $\Gamma$ .

Note that the above program is well-typed in the appropriate environment. The head is a sequence of nodes, but we shall sometimes see it as the underlying sequence of programs. Also note that the nodes of a cut step appearing in the head correspond to program reductions where the redex is a cut ( $\rightsquigarrow_{cut}$ ).

► **Definition 27 (Weight).** The *weight* of a proof  $\pi$ , written  $w(\pi)$ , is the multiset of cut nodes in its head, ordered by their distance to the end of the head.

► **Lemma 28.** Let  $\pi'$  be obtained from a proof  $\pi$  by a cut-reduction. We have that:

- (i)  $\pi'$  is a valid proof;
- (ii)  $|hd(\pi')| \leq |hd(\pi)|$ , where  $|s|$  is the length of a sequence  $s$ ;
- (iii) if the reduced cut was the last  $\rightsquigarrow_{cut}$  step in  $hd(\pi)$ , then  $w(\pi') < w(\pi)$ .

**Proof sketch.** By case analysis; key cases strictly decrease the length of the head while it is only conserved by commutative cases. We list and discuss all cases in App. F; one of the two \*-key cases is the following one:

$$\begin{array}{c} \Delta \rightarrow e \quad \Sigma \rightarrow e^* \\ \text{cut} \frac{}{\Gamma, \Delta, \Sigma \rightarrow e^*} \end{array} \quad \begin{array}{c} \Gamma, \Pi \rightarrow f \quad \Gamma, e, e^*, \Pi \rightarrow f \\ \text{*-l} \frac{}{\Gamma, e^*, \Pi \rightarrow f} \end{array} \quad \mapsto \quad \begin{array}{c} \Delta \rightarrow e \\ \text{cut} \frac{}{\Gamma, \Delta, \Sigma, \Pi \rightarrow f} \end{array} \quad \begin{array}{c} \Sigma \rightarrow e^* \quad \Gamma, e, e^*, \Pi \rightarrow f \\ \text{cut} \frac{}{\Gamma, e, \Sigma, \Pi \rightarrow f} \end{array}$$

If the reduced cut (with conclusion at  $v$ ) occurs in the head of  $\pi$  then the heads of the two proofs only differ by the following subsequences, inside some evaluation context:

$$\begin{array}{l} \rightsquigarrow_{cut_{e^*}} \quad v[\vec{M}, \vec{N}, \vec{O}, \vec{P}] \\ \rightsquigarrow \quad v1[\vec{M}, v0[\vec{N}, \vec{O}], \vec{P}] \\ \rightsquigarrow^n \quad v1[\vec{M}, v00[\vec{N} :: v01[\vec{O}], \vec{P}]] \\ \rightsquigarrow^o \quad v1[\vec{M}, N' :: O', \vec{P}] \\ \rightsquigarrow \quad v11[\vec{M}, N', O', \vec{P}] \end{array} \quad \rightsquigarrow_{cut_e} \quad v[\vec{M}, \vec{N}, \vec{O}, \vec{P}] \\ \rightsquigarrow^n \quad v1[\vec{M}, v0[\vec{N}], \vec{O}, \vec{P}] \\ \rightsquigarrow^n \quad v1[\vec{M}, N', \vec{O}, \vec{P}] \\ \rightsquigarrow_{cut_{e^*}} \quad v11[\vec{M}, N', v10[\vec{O}], \vec{P}] \\ \rightsquigarrow^o \quad v11[\vec{M}, N', O', \vec{P}]$$

(Note that the programs  $\vec{M}, \vec{N}, \vec{O}, \vec{P}$  are irreducible due to the innermost strategy, and that  $v00$  in the starting proof and  $v0$  in the resulting one both point to the same subproof:  $\pi_{v00} = \pi'_{v0}$ .) The new head is shorter by one step, and the initial cut on  $e^*$  is replaced by two cuts which are closer to the end of the head.

Commutative cases do not always shorten the head, but either they move the cut closer to its end, or the head no longer visits it. For instance, when the left premiss of the reduced cut ends with a  $-l$  step, the rule is

$$\begin{array}{c} \Delta, e, f, \Sigma \rightarrow g \\ \text{-l} \frac{}{\Delta, ef, \Sigma \rightarrow g} \\ \text{cut} \frac{}{\Gamma, \Delta, ef, \Sigma, \Pi \rightarrow h} \end{array} \quad \mapsto \quad \begin{array}{c} \Delta, e, f, \Sigma \rightarrow g \quad \Gamma, g, \Pi \rightarrow h \\ \text{cut} \frac{}{\Gamma, \Delta, e, f, \Sigma, \Pi \rightarrow h} \\ \text{-l} \frac{}{\Gamma, \Delta, ef, \Sigma, \Pi \rightarrow h} \end{array}$$

If the head of  $\pi$  goes through the step  $v[\vec{M}, \vec{N}, O, \vec{P}, \vec{Q}] \rightsquigarrow_{cut_{ef}} v1[\vec{M}, v0[\vec{N}, O, \vec{P}], \vec{Q}]$ , then there are two cases to consider:

- either  $O = \langle O_1, O_2 \rangle$  and the sequence continues with  $v1[\vec{M}, v00[\vec{N}, O_1, O_2, \vec{P}], \vec{Q}]$ ; then in  $\pi'$  we get  $v[\vec{M}, \vec{N}, O, \vec{P}, \vec{Q}] \rightsquigarrow v0[\vec{M}, \vec{N}, O_1, O_2, \vec{P}, \vec{Q}] \rightsquigarrow_{cut_{ef}} v01[\vec{M}, v00[\vec{N}, O_1, O_2, \vec{P}], \vec{Q}]$ ; the length is preserved and the cut has been pushed towards the end;
- or not, and the head of  $\pi'$  stops earlier, without visiting the cut on  $ef$  anymore, thus decreasing the weight.

For (iii), the assumption that the cut-reduction took place on the last cut node of the head is used in some of the cases to ensure that the weight of the other cut nodes of the head does not increase (e.g., in some of the right  $-r$  and  $*-r_2$  cases). ◀

► **Proposition 29** (Productive cut-reduction). *For a proof  $\pi$ , there exists a proof  $\pi'$  obtained from  $\pi$  by a sequence of cut-reductions, which does not start with a cut.*

**Proof.** By induction on the weight, reduce the last cut visited by the head until the head no longer contains any cut. The resulting proof cannot start with a cut, by definition. ◀

We can finally prove cut-elimination.

**Proof sketch of Thm. 13.** Focus on a lowest cut, at node  $v$ , and apply Prop. 29 to the corresponding subproof ( $\pi_v$ ). By iterating this process, we obtain in the limit a cut-free preproof  $\pi'$  with the same conclusion as the starting one. Moreover, thanks to Lem. 28(i), all heads of subproofs of  $\pi'$  are finite, so that  $\pi'$  is valid by Prop. 9: an infinite branch of  $*-r_2$  steps would give rise to a subproof with an infinite head. ◀

## 5 Action algebras, Kleene lattices, and action lattices

We now consider extensions of Kleene algebra by residuals and meets, as axiomatised in [30] and [22]. We first extend the system LKA with the following rules, which are standard from substructural logic [27, 15]. We write LAL for the corresponding system.

$$\begin{array}{ccc}
 \backslash-l \frac{\Delta \rightarrow e \quad \Gamma, f, \Sigma \rightarrow g}{\Gamma, \Delta, e \backslash f, \Sigma \rightarrow g} & /-l \frac{\Delta \rightarrow e \quad \Gamma, f, \Sigma \rightarrow g}{\Gamma, f/e, \Delta, \Sigma \rightarrow g} & \cap-l_i \frac{\Gamma, e_i, \Delta \rightarrow f}{\Gamma, e_1 \cap e_2, \Delta \rightarrow f} \quad i \in \{1, 2\} \\
 \\
 \backslash-r \frac{e, \Gamma \rightarrow f}{\Gamma \rightarrow e \backslash f} & /-r \frac{\Gamma, e \rightarrow f}{\Gamma \rightarrow f/e} & \cap-r \frac{\Gamma \rightarrow e \quad \Gamma \rightarrow f}{\Gamma \rightarrow e \cap f}
 \end{array}$$

We define judgements as previously. Except for Thm. 31, the results below also hold for action algebras and Kleene lattices using the appropriate fragment of LAL. We prove soundness w.r.t. star-continuous models exactly like for Kleene algebra (Thm. 10, App. B).

► **Theorem 30** (Soundness). *If  $LAL^- \vdash^\infty e_1, \dots, e_n \rightarrow e$ , then  $AL^* \models e_1 \cdots e_n \leq e$ .*

As announced in the introduction, regular proofs are sound for all (non-necessarily star-continuous) action lattices. We prove it using proof-theoretical arguments to translate every regular proof into an inductive proof from the axioms of action lattices.

► **Theorem 31** (Regular soundness). *If  $LAL \stackrel{\omega}{\vdash} e_1, \dots, e_n \rightarrow f$  then  $AL \models e_1 \cdots e_n \leq f$ .*

**Proof.** We prove the statement for all regular proofs in *\*-normal form*, where every backpointer points to a ‘validating’  $*-l$ -step: every infinite branch of the starting proof has a valid thread; since the proof is regular, this thread must be infinitely often principal for  $*-l$ -step of some sequent of the branch; cut the infinite branch by using a backpointer the second time this sequent appears in the branch.

We proceed by induction on the number of simple cycles in such a proof  $\pi$ . The interesting case is when  $\pi$  ends with a  $*-l$  step that is the target of a backpointer. Colour in red all ancestors of its principal formula that are the same expression,  $e^*$ . Let  $\{\Gamma_i, e^*, \Delta_i \rightarrow f_i\}_{i \in I}$  be the set of all sequents in  $\pi$  with  $e^*$  principal (thus for a  $*-l$  step) and let  $(\pi_i^l : \Gamma_i, \Delta_i \rightarrow f_i)_{i \in I}$  and  $(\pi_i^r : \Gamma_i, e, e^*, \Delta_i \rightarrow f_i)_{i \in I}$  be the set of subproofs rooted at their left and right premisses, respectively.

Define expressions  $g_i = \prod \Gamma_i$ ,  $d_i = \prod \Delta_i$ ,  $h_i = (g_i \setminus f_i) / d_i$ , and  $h = \bigcap_{i \in I} h_i$ . For  $i \in I$ , construct proofs  $\pi_i^{r'}$  from  $\pi_i^r$  by replacing each  $e^*$  by  $h$ , modifying critical steps as follows:

$$\frac{\frac{\Gamma_j, \Delta_j \rightarrow f_j \quad \Gamma_j, e, e^*, \Delta_j \rightarrow f_j}{\Gamma_j, e^*, \Delta_j \rightarrow f_j} \quad \text{*-l}}{\Gamma_j, \Delta_j \rightarrow f_j} \quad \mapsto \quad \left. \begin{array}{c} \frac{\frac{\frac{\frac{\Gamma_j \rightarrow g_j \quad f_j \rightarrow f_j}{\Gamma_j, g_j \setminus f_j \rightarrow f_j} \quad \text{-l}}{\Delta_j \rightarrow d_j} \quad \text{-r}}{\Gamma_j, h_j, \Delta_j \rightarrow f_j} \quad \text{-l}}{\Gamma_j, h, \Delta_j \rightarrow f_j} \quad \text{-l} \end{array} \right\} \rho_j$$

Note that the proofs  $\pi_i^l$  and  $\pi_i^{r'}$  have fewer simple cycles than  $\pi$ , so that by induction hypothesis we have that  $g_i d_i \leq f_i$  and  $g_i e h d_i \leq f_i$  hold universally in action lattices, for all  $i \in I$ . From those we deduce  $1 \leq h$  and  $eh \leq h$  using the laws about residuals and conjunction. Thus we have  $e^* \leq h$  by star induction (1). Finally note that following the above proof  $\rho_j$  we have in action lattices that  $g_j h d_j \leq f_j$  and thus  $g_j e^* d_j \leq f_j$ . We conclude by choosing  $j$  such that  $(\Gamma_j, \Delta_j, f_j) = (\Gamma, \Delta, f)$ .  $\blacktriangleleft$

Note that we crucially rely on the presence of both residuals and meet to compute invariants for Kleene stars in the above proof, so that it does not immediately carry over to action algebras and Kleene lattices.

Conversely, the regular fragment of LAL (with cut) is complete for action lattices.

► **Theorem 32** (Regular completeness). *If  $\text{AL} \models e \leq f$  then  $\text{LAL} \stackrel{\omega}{\vdash} e \rightarrow f$ .*

**Proof.** The axioms defining meet and residual immediately translate to finite derivations in LAL, one can thus extend the proof of Thm. 12.  $\blacktriangleleft$

Note that the regular fragment cannot be complete for star continuous models: a regular proof is a finite verifiable object and the equational theory of star-continuous action lattices is  $\Pi_1^0$ -complete [7]. The full, non-regular system is however complete for star-continuous models:

► **Theorem 33** (Star-continuous completeness). *If  $\text{AL}^* \models e \leq f$  then  $\text{LAL} \stackrel{\infty}{\vdash} e \rightarrow f$ .*

**Proof.** As for Thms. 12 and 32, consider the relation  $\equiv'$  defined by  $e \equiv' f$  if  $\text{LAL} \stackrel{\infty}{\vdash} e \rightarrow f$  and  $\text{LAL} \stackrel{\infty}{\vdash} f \rightarrow e$ . Expressions quotiented by this slightly larger relation also form an action lattice, which we prove star-continuous using the natural simulation of an  $\omega$ -rule for Kleene star: combine proofs  $(\pi_i)_{i \in \mathbb{N}}$  of the sequents  $(\Gamma, e^i, \Delta \rightarrow f)_{i \in \mathbb{N}}$  as follows:

$$\frac{\frac{\frac{\frac{\Gamma, e, \Delta \rightarrow f}{\Gamma, e, \Delta \rightarrow f} \quad \text{-l}}{\Gamma, e, e, \Delta \rightarrow f} \quad \text{-l}}{\Gamma, e, e, \Delta \rightarrow f} \quad \text{-l}}{\Gamma, e^*, \Delta \rightarrow f} \quad \text{-l}}{\Gamma, e, e, \Delta \rightarrow f \quad \Gamma, e, e, \Delta \rightarrow f \quad \Gamma, e, e, \Delta \rightarrow f \quad \dots} \quad \text{-l}$$

The remaining property to establish is cut-elimination: combined with Thm. 30 it gives soundness of proofs with cut w.r.t. star-continuous models, and combined with Thm. 33 it gives completeness of  $\text{LAL}^-$  w.r.t. these models.

## 6 Cut-elimination in LAL

The main alteration to the proof for LKA is that we need a more sophisticated notion of programs. We associate linear functions to residuals, and additive pairs to meets: a program for  $e \cap f$  waits to see whether the environment wants a value for  $e$  or a value for  $f$ —but not both, and reacts accordingly. We thus extend the syntax of programs (Dfn. 14) to include  $\lambda$ -abstractions, which will be used for residuals, and a new kind of pairs for meets.

$$M, N ::= x \mid \star \mid \langle M, N \rangle \mid \text{in}_i M \mid [] \mid M :: N \mid \pi[\vec{M}] \mid \lambda x.M \mid \langle\langle M, N \rangle\rangle$$

The type system (Fig. 3) is extended by the following rules, where in the final rule,  $E_1$  and  $E_2$  are extensions of  $E$ .

$$\frac{x : e, E \vdash M : f}{E \vdash \lambda x.M : e \setminus f} \quad \frac{E, x : e \vdash M : f}{E \vdash \lambda x.M : f/e} \quad \frac{E_1 \vdash M : e \quad E_2 \vdash N : f}{E \vdash \langle\langle M, N \rangle\rangle : e \cap f}$$

► **Lemma 34** (Substitution lemma). *If  $E \vdash N : e$  and  $F, x : e, F' \vdash M : f$  with  $F, E, F'$  defined, then  $F, E, F' \vdash M\{N/x\} : f$ , where  $M\{N/x\}$  is  $M$  with  $x$  substituted by  $N$ .<sup>2</sup>*

The following reductions are added, using the same conventions as in Dfn. 19:

$$\begin{aligned} \cap\text{-}l_i : v[\vec{M}, \langle\langle N_1, N_2 \rangle\rangle, \vec{P}] &\rightsquigarrow v0[\vec{M}, N_i, \vec{P}] & \cap\text{-}r : v[\vec{M}] &\rightsquigarrow \langle\langle v0[\vec{M}], v1[\vec{M}] \rangle\rangle \\ \setminus\text{-}l : v[\vec{M}, \vec{N}, \lambda x.F, \vec{P}] &\rightsquigarrow v1[\vec{M}, F\{v0[\vec{N}]/x\}, \vec{P}] & \setminus\text{-}r : v[\vec{M}] &\rightsquigarrow \lambda x.v0[x, \vec{M}] \\ /\text{-}l : v[\vec{M}, \lambda x.F, \vec{N}, \vec{P}] &\rightsquigarrow v1[\vec{M}, F\{v0[\vec{N}]/x\}, \vec{P}] & /\text{-}r : v[\vec{M}] &\rightsquigarrow \lambda x.v0[\vec{M}, x] \end{aligned}$$

One has to be careful about what we deem to be evaluation contexts: lambda abstractions and additive pairs are *not* considered evaluation contexts. This is crucial to obtain subject-reduction: otherwise some redexes duplicated by the  $\cap\text{-}r$  rule can be active at the same time, thus breaking the property of Lem. 24 used in our termination proof that a given node appears at most once in the run of a program.

Despite this subtlety, Prop. 22 (subject reduction) and Prop. 25 (termination) are proved for this extended system exactly as in the Kleene algebra case—see App. E. It thus remains to show that the new cut reductions do not increase the length of heads, and strictly decrease the weight (Lem. 28). The key cases are easy: they strictly decrease the length and replace the cut by smaller ones. Amongst the commutative cases, some care is required when a right introduction rule appears on the right of the cut. For instance, for meet:

$$\text{cut} \frac{\Delta \rightarrow f \quad \cap\text{-}r \frac{\Gamma, f, \Sigma \rightarrow e_1 \quad \Gamma, f, \Sigma \rightarrow e_2}{\Gamma, f, \Sigma \rightarrow e_1 \cap e_2}}{\Gamma, \Delta, \Sigma \rightarrow e_1 \cap e_2} \mapsto \text{cut} \frac{\Delta \rightarrow f \quad \Gamma, f, \Sigma \rightarrow e_1 \quad \text{cut} \frac{\Delta \rightarrow f \quad \Gamma, f, \Sigma \rightarrow e_2}{\Gamma, \Delta, \Sigma \rightarrow e_1}}{\cap\text{-}r \frac{\Gamma, \Delta, \Sigma \rightarrow e_1}{\Gamma, \Delta, \Sigma \rightarrow e_1 \cap e_2}}$$

If the head of  $\pi$  contains the sequence,

$$v[\vec{M}, \vec{N}, \vec{O}] \rightsquigarrow v1[\vec{M}, v0[\vec{N}], \vec{O}] \rightsquigarrow^n v1[\vec{M}, N', \vec{O}] \rightsquigarrow \langle\langle v10[\vec{M}, N', \vec{O}], v11[\vec{M}, N', \vec{O}] \rangle\rangle$$

where  $v$  is the reduced cut-node, then in the head of  $\pi'$  we just get:

$$v[\vec{M}, \vec{N}, \vec{O}] \rightsquigarrow \langle\langle v0[\vec{M}, \vec{N}, \vec{O}], v1[\vec{M}, \vec{N}, \vec{O}] \rangle\rangle$$

<sup>2</sup> More precisely, the occurrences of  $x$  selected by the typing derivation of  $M$ .

Here we see the need for  $\langle\langle -, - \rangle\rangle$  not being an evaluation context: the computations involving  $\vec{N}$  would otherwise be duplicated, thus potentially increasing the length of the run. If the head of  $\pi$  never touches the produced additive pair, then the head of  $\pi'$  is strictly shorter, and the cut on  $e_1 \cap e_2$  is not visited anymore. Otherwise, this pair can only be destroyed by a  $\cap\text{-}l_i$  rule:  $\langle\langle v10[\vec{M}, N', \vec{O}], v11[\vec{M}, N', \vec{O}] \rangle\rangle \rightsquigarrow v1i[\vec{M}, N', \vec{O}]$ , and the head of  $\pi'$  can ‘catch up’ by doing:

$$\langle\langle v0[\vec{M}, \vec{N}, \vec{O}], v1[\vec{M}, \vec{N}, \vec{O}] \rangle\rangle \rightsquigarrow vi[\vec{M}, \vec{N}, \vec{O}] \rightsquigarrow vi1[\vec{M}, vi0[\vec{N}], \vec{O}] \rightsquigarrow^n vi1[\vec{M}, N', \vec{O}]$$

The size of the head has not changed, but the cut is closer to the end. The analogous case for residuals is similar since the creation of a  $\lambda$ -abstraction temporarily blocks reductions; other cases can be found in App. F. Finally, by the same argument as for Thm. 13 we obtain:

► **Theorem 35** (Cut elimination). *If  $\text{LAL} \vdash^\infty \Gamma \rightarrow e$  then  $\text{LAL}^- \vdash^\infty \Gamma \rightarrow e$ .*

One useful application of this cut-elimination result is the following alternative proof of the upper bound result of Palka for star-continuous action lattices:

► **Corollary 36** (Palka [28]).  *$\text{AL}^*$  is in  $\Pi_1^0$ .*

**Proof.** We say that a sequent  $\Gamma \rightarrow e$  has a  $d$ -derivation, for  $d \in \mathbb{N}$ , if there is a  $\text{LAL}^-$  derivation ending in  $\Gamma \rightarrow e$  for which each branch has length  $d$ , or otherwise terminates at a correct initial sequent in length  $< d$ . To avoid validity issues, we assume that the left premiss of every  $*\text{-}r_2$  step has nonempty antecedent, so that all preproofs become valid without sacrificing provability (cf. Prop. 9). We define a  $\Pi_1^0$  predicate  $\text{Prov}(\Gamma \rightarrow e)$  as  $\forall d \in \mathbb{N}$ . “there is a  $d$ -derivation of  $\Gamma \rightarrow e$ ”. Notice that this is indeed  $\Pi_1^0$  since the size of a  $d$ -derivation is exponentially bounded. Furthermore, if  $\text{Prov}(\Gamma \rightarrow e)$  then, by the infinite pigeonhole principle, we may recover an infinite proof of  $\Gamma \rightarrow e$ , by inductively choosing premisses resulting in larger derivations that nonetheless prefix infinitely many  $d$ -derivations. ◀

## 7 Conclusions

We presented a simple sequent system LKA that admits non-wellfounded proofs and showed it to be sound and complete for Kleene algebra, KA, by consideration of the free model of rational languages. We showed that its regular fragment is already complete, in the presence of cut, by a direct simulation of KA. We also gave a cut-elimination result for LKA, obtaining an alternative proof of completeness of its cut-free fragment.

We were able to generalise these arguments to an extended system LAL of Kleene algebras with residuals and meets, resulting in a sound and complete cut-free system for the equational theory of star-continuous action lattices,  $\text{AL}^*$ . Thanks to the subformula property in cut-free proofs, this also gives us proof-theoretical characterisations for star-continuous action algebras and Kleene lattices. This yields alternative proofs of several results of Palka [28], namely conservativity of  $\text{AL}^*$  over its fragments, as well as their membership in  $\Pi_1^0$ .

Finally, we characterised the theory of all action lattices by just the regular proofs of LAL. Whether the equational theory of action lattices is decidable remains open. It would be interesting to see if techniques such as interpolants for our system LAL could yield decidability.

It would be natural to consider systems which are commutative and/or contain arbitrary fixed points, bringing the subject matter closer to that of [12]. We would however not be able to arrive at a similar subformula property once fixed point formulae are allowed to contain meets and residuals, since this property is essentially thanks to the presence of only ‘positive’ connectives in KA, from the point of view of focusing [2].



## References

- 1 C. J. Anderson, N. Foster, A. Guha, J.-B. Jeannin, D. Kozen, C. Schlesinger, and D. Walker. [NetKAT: semantic foundations for networks](#). In *Proc. POPL*, pages 113–126. ACM, 2014.
- 2 J.-M. Andreoli. Logic programming with focusing proofs in linear logic. *J. Log. and Comput.*, 2(3):297–347, 1992.
- 3 A. Angus and D. Kozen. [Kleene algebra with tests and program schematology](#). Technical Report TR2001-1844, CS Dpt., Cornell University, July 2001.
- 4 M. Boffa. [Une condition impliquant toutes les identités rationnelles](#). *Informatique Théorique et Applications*, 29(6):515–518, 1995.
- 5 T. Braibant and D. Pous. [An efficient Coq tactic for deciding Kleene algebras](#). In *Proc. 1st ITP*, volume 6172 of *LNCS*, pages 163–178. Springer, 2010.
- 6 S. R. Buss. An introduction to proof theory. *Handbook of proof theory*, 137:1–78, 1998.
- 7 W. Buszkowski. [On action logic: Equational theories of action algebras](#). *J. Log. Comput.*, 17(1):199–217, 2007.
- 8 P. Clairambault. [Least and greatest fixpoints in game semantics](#). In *Proc. FoSSaCS*, pages 16–31, 2009.
- 9 J. H. Conway. *Regular algebra and finite machines*. Chapman and Hall, 1971.
- 10 A. Das and D. Pous. [A cut-free cyclic proof system for Kleene algebra](#). In *Proc. TABLEAUX*, volume 10501 of *LNCS*, pages 261–277. Springer, 2017.
- 11 H. Doornbos, R. Backhouse, and J. van der Woude. [A calculational approach to mathematical induction](#). *TCS*, 179(1-2):103–135, 1997.
- 12 A. Doumane, D. Baelde, and A. Saurin. [Infinitary proof theory: the multiplicative additive case](#). In *Proc. CSL*, volume 62 of *LIPICs*, pages 42:1–42:17, Sept. 2016.
- 13 J. Fortier and L. Santocanale. [Cuts for circular proofs: semantics and cut-elimination](#). In *Proc. CSL*, volume 23 of *LIPICs*, pages 248–262, 2013.
- 14 A. Frisch and L. Cardelli. [Greedy regular expression matching](#). In *Proc. ICALP*, volume 3142 of *LNCS*, pages 618–629. Springer, 2004.
- 15 N. Galatos, P. Jipsen, T. Kowalski, and H. Ono. *Residuated Lattices: An Algebraic Glimpse at Substructural Logics*. Elsevier, 2007.
- 16 J.-Y. Girard. Linear logic. *TCS*, 50:1–102, 1987.
- 17 F. Henglein and L. Nielsen. [Regular expression containment: coinductive axiomatization and computational interpretation](#). In *Proc. POPL, 2011*, pages 385–398. ACM, 2011.
- 18 C. A. R. Hoare, B. Möller, G. Struth, and I. Wehrman. [Concurrent Kleene Algebra](#). In *Proc. CONCUR*, volume 5710 of *LNCS*, pages 399–414. Springer, 2009.
- 19 P. Jipsen. [From semirings to residuated Kleene lattices](#). *Studia Logica*, 76(2):291–303, 2004.
- 20 S. C. Kleene. [Representation of events in nerve nets and finite automata](#). In *Automata Studies*, pages 3–41. Princeton University Press, 1956.
- 21 D. Kozen. [A completeness theorem for Kleene algebras and the algebra of regular events](#). In *Proc. LICS*, pages 214–225. IEEE, 1991.
- 22 D. Kozen. [On action algebras](#). In J. van Eijck and A. Visser, editors, *Logic and Information Flow*, pages 78–88. MIT Press, 1994.
- 23 D. Kozen. [On Hoare logic and Kleene algebra with tests](#). *ACM Trans. Comput. Log.*, 1(1):60–76, 2000.
- 24 D. Kozen and M.-C. Patron. [Certification of compiler optimizations using Kleene algebra with tests](#). In *Proc. CL2000*, volume 1861 of *LNAI*, pages 568–582. Springer, 2000.
- 25 A. Krauss and T. Nipkow. [Proof pearl: Regular expression equivalence and relation algebra](#). *JAR*, 49(1):95–106, 2012.
- 26 D. Krob. [Complete systems of B-rational identities](#). *TCS*, 89(2):207–343, 1991.
- 27 J. Lambek. [The mathematics of sentence structure](#). *The American Mathematical Monthly*, 65:154–170, 1958.

- 28 E. Palka. An infinitary sequent system for the equational theory of  $*$ -continuous action lattices. *Fundam. Inform.*, pages 295–309, 2007.
- 29 D. Pous. Kleene Algebra with Tests and Coq tools for while programs. In *Proc. ITP*, volume 7998 of *LNCS*, pages 180–196. Springer, 2013.
- 30 V. Pratt. Action logic and pure induction. In *Proc. JELIA*, volume 478 of *LNCS*, pages 97–120. Springer, 1990.
- 31 V. N. Redko. On defining relations for the algebra of regular events. *Ukrainskii Matematicheskii Zhurnal*, 16:120–126, 1964.
- 32 K. Schütte. *Proof Theory*. Grundlehren der mathematischen Wissenschaften 225. Springer Berlin Heidelberg, 1977. Translation of *Beweistheorie*, 1968.
- 33 C. Wurm. Kleene algebras, regular languages and substructural logics. In *Proc. GandALF*, EPTCS, pages 46–59, 2014.

## A Proof ancestry

We recall some basic notions of proof ancestry, e.g. from [6], adapted to our setting. Let  $\rho$  be a rule instance of LAL, as written in Fig. 2 or in the beginning of Sect. 5.

If  $\rho$  is a left rule then its *principal formula* is the indicated expression in the antecedent of its lower sequent. If  $\rho$  is a right rule then its principal formula is just the succedent formula of its lower sequent. *id* and *cut* have no principal formula. Any other formulae in the lower sequent of  $\rho$  are called *side formulas*.

If  $\rho$  is a non-residual right rule then its *auxiliary formulae* are just the succedents of its upper sequent(s), and if it is a non-residual left rule then its auxiliary formulae are just the indicated expressions in the antecedent(s) of its upper sequent(s). If  $\rho$  is  $\setminus\text{-}r$  or  $/\text{-}r$  then the auxiliary formulae are the indicated occurrences of  $e$  and  $f$  in its upper sequent, and if  $\rho$  is  $\setminus\text{-}l$  or  $/\text{-}l$  then the auxiliary formulae are the indicated occurrences of  $e$  and  $f$  in its upper sequents. *id* has no auxiliary formulae, while *cut* has two auxiliary formulae: the two occurrences of  $e$  in its upper sequents; in this case we also call these formula occurrences the *cut formulae* of  $\rho$ .

An expression occurrence  $e$  in an upper sequent of  $\rho$  is an *immediate ancestor* of an expression occurrence  $f$  in its lower sequent if either  $f$  is principal and  $e$  is auxiliary, or if  $f$  is a side formula and  $e$  corresponds to the same occurrence of that side formula in an upper sequent, defined in the natural way. In a proof, the *ancestor* relation between two formula occurrences is defined as the reflexive transitive closure of the immediate ancestor relation; i.e.  $e$  is an ancestor of  $f$  if there is a path from  $f$  upwards, always choosing immediate ancestors, that ends in  $e$ . We also say that  $f$  is a (*immediate*) *descendant* of  $e$  if  $e$  is an (immediate, resp.) ancestor of  $f$ .

## B Soundness of LAL without cut w.r.t. star continuous models

We give here a detailed proof for Thms. 10 and 30: soundness of cut-free proofs w.r.t. star continuous models. The proofs are independent from the considered fragment (Kleene algebra, action algebra, action lattices) thanks to the sub-formula property.

► **Observation 37.** Every rule of LAL is sound (for Kleene algebra, action algebra, or action lattices depending on the connectives it mentions): if its premisses are valid then so is its conclusion.

We first prove the lemma mentioned in the proof sketch we gave for Thm. 10.

► **Lemma 38.** *If  $\text{LAL}^- \vdash^\infty \Gamma, e^*, \Delta \rightarrow f$  then, for each  $n \in \mathbb{N}$ ,  $\text{LAL}^- \vdash^\infty \Gamma, e^n, \Delta \rightarrow f$ .*

**Proof.** We define appropriate preproofs from by induction on  $n$ . Replace every direct ancestor of  $e^*$  by  $e^n$ , adjusting origins as follows,

$$*_l \frac{\Gamma, \Delta \rightarrow f \quad \Gamma, e, e^*, \Delta \rightarrow f}{\Gamma, e^*, \Delta \rightarrow f} \quad \mapsto \quad *_l \frac{\Gamma, \Delta \rightarrow f}{\Gamma, 1, \Delta \rightarrow f} \quad \text{or} \quad *_l \frac{\Gamma, e, e^{n-1}, \Delta \rightarrow f}{\Gamma, e^n, \Delta \rightarrow f}$$

when  $n = 0$  or  $n > 0$ , respectively. In the latter case we appeal to the inductive hypothesis.

Notice that, on branches where  $e^*$  is never principal, this is simply a substitution of  $e^n$  for  $e^*$  everywhere along the branch. The preproof resulting from this entire construction is fair since every infinite branch will share a tail with the proof we began with. ◀

We can now prove the soundness results w.r.t. star continuous models:

**Proof of Thms. 10 and 30.** Let  $\pi$  be an  $\text{LAL}^-$  proof of  $\Sigma \rightarrow f$ . We proceed by induction on the multiset of formulae in its conclusion, performing case analysis on the last rule. For all but two cases, we just use Obs. 37 and the induction hypotheses. The first remaining case is  $*_r_2$ , where we must appeal to a sub-induction since the measure does not always strictly decrease in the right premiss (Prop. 9). The last case is  $*_l$ , where  $\Sigma = \Gamma, e^*, \Delta$ . By Lem. 38,  $\pi$  can be transformed into proofs  $\pi_n$  of  $\Gamma, f^n, \Delta \rightarrow f$  for each  $n \in \mathbb{N}$ . Each  $\pi_n$  derives a sequent whose weight is strictly smaller than that of  $\Sigma \rightarrow f$ , which is thus valid by the inductive hypothesis. Finally, this means that  $\Gamma, f^*, \Delta \rightarrow g$  is valid, by star-continuity. ◀

## C Completeness of LKA without cut for Kleene algebra

► **Observation 39.** All left rules of  $\text{LKA}^-$  are (*weakly*) *invertible*: the validity of their conclusion implies the validity of all their premisses.

► **Lemma 40.** *If  $a_1 \dots a_n$  is a word in  $\mathcal{L}(e)$  for some expression  $e$ , then there is a finite proof of the sequent  $a_1, \dots, a_n \rightarrow e$  using only right logical rules.*

**Proof.** By a straightforward induction on  $e$ . ◀

**Proof of Thm. 11.** Suppose  $\mathcal{L}(e_1 \cdots e_n) \subseteq \mathcal{L}(e)$ . We construct an  $\text{LKA}^-$  proof of  $e_1, \dots, e_n \rightarrow e$  bottom-up, by first apply left-rules forever. By invertibility every sequent encountered will be valid, although this will not necessarily terminate due to  $*_l$ . Notice, however that we obtain fairness, since any infinite branch of only left rules must contain  $*_l$  infinitely often.

The branches which terminate to a normal form have leaves of the form  $a_1, \dots, a_n \rightarrow e$ . Since such a leaf must be valid, we have  $a_1 \dots a_n \in \mathcal{L}((e))$  and we can replace this leaf by a finite proof using Lem. 40. ◀

## D Summary of the reduction rules

- If  $\pi_v$  is  $id \frac{}{e \rightarrow e}$   
then  $v[M] \rightsquigarrow M$ .
- If  $\pi_v$  is  $1-r \frac{}{\rightarrow 1}$   
then  $v[] \rightsquigarrow \star$ .
- If  $\pi_v$  is  $-r \frac{\Gamma \rightarrow e \quad \Delta \rightarrow f}{\Gamma, \Delta \rightarrow ef}$   
then  $v[\vec{M}, \vec{N}] \rightsquigarrow \langle v0[\vec{M}], v1[\vec{N}] \rangle$ .
- If  $\pi_v$  is  $+r_i \frac{\Gamma \rightarrow e_i}{\Gamma \rightarrow e_1 + e_2}$   
then  $v[\vec{M}] \rightsquigarrow in_i(v0[\vec{M}])$ .
- If  $\pi_v$  is  $*-r_1 \frac{}{\rightarrow f^*}$   
then  $v[] \rightsquigarrow []$ .
- If  $\pi_v$  is  $*-r_2 \frac{\Gamma \rightarrow e \quad \Delta \rightarrow e^*}{\Gamma, \Delta \rightarrow e^*}$   
then  $v[\vec{M}, \vec{N}] \rightsquigarrow v0[\vec{M}] :: v1[\vec{N}]$ .
- If  $\pi_v$  is  $\cap-r \frac{\Gamma \rightarrow f_1 \quad \Gamma \rightarrow f_2}{\Gamma \rightarrow f_1 \cap f_2}$   
then  $v[\vec{M}] \rightsquigarrow \langle v0[\vec{M}], v1[\vec{M}] \rangle$ .
- If  $\pi_v$  is  $\setminus-r \frac{e, \Gamma \rightarrow f}{\Gamma \rightarrow e \setminus f}$   
then  $v[\vec{M}] \rightsquigarrow \lambda x. v0[x, \vec{M}]$ .
- If  $\pi_v$  is  $/-r \frac{\Gamma, e \rightarrow f}{\Gamma \rightarrow f/e}$   
then  $v[\vec{M}] \rightsquigarrow \lambda x. v0[\vec{M}, x]$ .
- If  $\pi_v$  is  $cut \frac{\Delta \rightarrow e \quad \Gamma, e, \Sigma \rightarrow f}{\Gamma, \Delta, \Sigma \rightarrow f}$   
then  $v[\vec{M}, \vec{N}, \vec{P}] \rightsquigarrow v1[\vec{M}, v0[\vec{N}], \vec{P}]$ .
- If  $\pi_v$  is  $1-l \frac{\Gamma, \Delta \rightarrow f}{\Gamma, 1, \Delta \rightarrow f}$   
then  $v[\vec{M}, \star, \vec{N}] \rightsquigarrow v0[\vec{M}, \vec{N}]$ .
- If  $\pi_v$  is  $-l \frac{\Gamma, e, f, \Delta \rightarrow f}{\Gamma, ef, \Delta \rightarrow f}$   
then  $v[\vec{M}, \langle M, N \rangle, \vec{N}] \rightsquigarrow v0[\vec{M}, M, N, \vec{N}]$ .
- If  $\pi_v$  is  $+l \frac{\Gamma, e_1, \Delta \rightarrow f \quad \Gamma, e_2, \Delta \rightarrow f}{\Gamma, e_1 + e_2, \Delta \rightarrow f}$   
then  $v[\vec{M}, in_i M, \vec{N}] \rightsquigarrow vi[\vec{M}, M, \vec{N}]$ .
- If  $\pi_v$  is  $*-l \frac{\Gamma, \Delta \rightarrow f \quad \Gamma, e, e^*, \Delta \rightarrow f}{\Gamma, e^*, \Delta \rightarrow f}$   
then  $v[\vec{M}, [], \vec{N}] \rightsquigarrow v0[\vec{M}, [], \vec{N}]$   
and  $v[\vec{M}, M :: N, \vec{N}] \rightsquigarrow v1[\vec{M}, M, N, \vec{N}]$ .
- If  $\pi_v$  is  $\cap-l_i \frac{\Gamma, e_i, \Delta \rightarrow f}{\Gamma, e_1 \cap e_2, \Delta \rightarrow f}$   
then  $v[\vec{M}, \langle M_1, M_2 \rangle, \vec{N}] \rightsquigarrow v0[\vec{M}, M_i, \vec{N}]$ .
- If  $\pi_v$  is  $\setminus-l \frac{\Delta \rightarrow e \quad \Gamma, f, \Sigma \rightarrow g}{\Gamma, \Delta, e \setminus f, \Sigma \rightarrow g}$   
then  $v[\vec{M}, \vec{N}, \lambda x. F, \vec{P}] \rightsquigarrow v1[\vec{M}, F\{v0[\vec{N}]/x\}, \vec{P}]$ .
- If  $\pi_v$  is  $/-l \frac{\Delta \rightarrow e \quad \Gamma, f, \Sigma \rightarrow g}{\Gamma, f/e, \Delta, \Sigma \rightarrow g}$   
then  $v[\vec{M}, \lambda x. F, \vec{N}, \vec{P}] \rightsquigarrow v1[\vec{M}, F\{v0[\vec{N}]/x\}, \vec{P}]$ .

## E Subject reduction proof

Here we present some further proof details about the subject reduction, which is the key step to obtain that programs terminate (Prop. 25).

We use the following notation:

► **Notation 41.** We write  $\vec{E} \vdash \vec{M} : \Gamma$  when  $\Gamma$  has the same length as  $\vec{E}$  and  $\vec{M}$  and for all  $i$ , we have  $E_i \vdash M_i : \Gamma_i$ .

**Proof of Lem. 34 (substitution lemma).** Given the typing derivation for  $F, x : e, F' \vdash M : f$ , replace every ancestor of  $x : e$  on the left by  $E$  and every ancestor of  $x$  in  $M$  by  $N$ . The

only critical case is the initial step, which can simply be replaced by a typing derivation for  $E \vdash N : e$ .  $\blacktriangleleft$

**Proof of Prop. 22 (subject reduction).** We argue by induction on the structure of a typing derivation for  $E \vdash M : e$ . If the reduction takes place under a context, we use the induction hypothesis and the substitution lemma. Otherwise, it suffices to conduct a ‘last step analysis’, extracting the immediate subderivations to form a new typing derivation for  $E \vdash N : e$ . Some interesting cases are the following:

- $cut : v[\vec{M}, \vec{N}, \vec{P}] \rightsquigarrow v1[\vec{M}, v0[\vec{N}], \vec{P}]$ . A typing derivation for the LHS has the form on the left below, so we recover a typing derivation for the RHS on the right below.

$$\frac{\vec{E} \vdash \vec{M} : \Gamma \quad \vec{F} \vdash \vec{N} : \Delta \quad \vec{G} \vdash \vec{P} : \Sigma}{v, \vec{E}, \vec{F}, \vec{G} \vdash v[\vec{M}, \vec{N}, \vec{P}] : f} \quad \frac{\vec{F} \vdash \vec{N} : \Delta}{\vec{E} \vdash \vec{M} : \Gamma \quad v0, \vec{F} \vdash v0[\vec{N}] : e \quad \vec{G} \vdash \vec{P} : \Sigma} \frac{}{v1, \vec{E}, v0, \vec{F}, \vec{G} \vdash v1[\vec{M}, \vec{N}, \vec{P}] : f}$$

Note that  $v1, \vec{E}, v0, \vec{F}, \vec{G}$  is well-defined since  $v, \vec{E}, \vec{F}, \vec{G}$  is, and that it forms an extension.

- $\cap\text{-}r : v[\vec{M}] \rightsquigarrow \langle\langle v0[\vec{M}], v1[\vec{M}] \rangle\rangle$ . A typing derivation for the LHS has the form on the left below, so we recover a typing derivation for the RHS on the right below:

$$\frac{\vec{E} \vdash \vec{M} : \Gamma}{v, \vec{E} \vdash v[\vec{M}] : e \cap f} \quad \frac{\vec{E} \vdash \vec{M} : \Gamma}{v0, \vec{E} \vdash v0[\vec{M}] : f} \quad \frac{\vec{E} \vdash \vec{M} : \Gamma}{v1, \vec{E} \vdash v1[\vec{M}] : f} \frac{}{v, \vec{E} \vdash \langle\langle v0[\vec{M}], v1[\vec{M}] \rangle\rangle : e \cap f}$$

$v0, \vec{E}$  and  $v1, \vec{E}$  are indeed extensions of  $v, \vec{E}$ .

- $\cap\text{-}l_i : v[\vec{M}, \langle\langle N_1, N_2 \rangle\rangle, \vec{P}] \rightsquigarrow v0[\vec{M}, N_i, \vec{P}]$ . A typing derivation for the LHS has the following form,

$$\frac{\vec{E} \vdash \vec{M} : \Gamma \quad \frac{F_1 \vdash N_1 : e_1 \quad F_2 \vdash N_2 : e_2}{F \vdash \langle\langle N_1, N_2 \rangle\rangle : e_1 \cap e_2} \quad \vec{G} \vdash \vec{P} : \Sigma}{v, \vec{E}, F, \vec{G} \vdash v[\vec{M}, \langle\langle N_1, N_2 \rangle\rangle, \vec{P}] : g}$$

where  $F_1$  and  $F_2$  are extensions of  $F$ . We deduce

$$\frac{\vec{E} \vdash \vec{M} : \Gamma \quad F_i \vdash N_i : e_i \quad \vec{G} \vdash \vec{P} : \Sigma}{v0, \vec{E}, F_i, \vec{G} \vdash v0[\vec{M}, N_i, \vec{P}] : g}$$

where  $v0, \vec{E}, F_i, \vec{G}$  is indeed an extension of  $v, \vec{E}, F, \vec{G}$ .

- $\backslash\text{-}l : v[\vec{M}, \vec{N}, \lambda x.P, \vec{Q}] \rightsquigarrow v1[\vec{M}, P\{v0[\vec{N}]/x\}, \vec{Q}]$ . A typing derivation for the LHS has the following form,

$$\frac{\vec{E} \vdash \vec{M} : \Gamma \quad \vec{F} \vdash \vec{N} : \Delta \quad \frac{x : e, \vec{G} \vdash P : f}{\vec{G} \vdash \lambda x.P : e \backslash f} \quad \vec{H} \vdash \vec{Q} : \Sigma}{v, \vec{E}, \vec{F}, \vec{G}, \vec{H} \vdash v[\vec{M}, \vec{N}, \lambda x.P, \vec{Q}] : g}$$

so we recover the following typing derivation for the RHS:

$$\frac{\vec{E} \vdash \vec{M} : \Gamma \quad \frac{\vec{F} \vdash \vec{N} : \Delta}{v0, \vec{F} \vdash v0[\vec{N}] : e} \quad \frac{x : e, \vec{G} \vdash P : f}{v0, \vec{F}, \vec{G} \vdash P\{v0[\vec{N}]/x\} : f} \quad \vec{H} \vdash \vec{Q} : \Sigma}{v1, \vec{E}, v0, \vec{F}, \vec{G}, \vec{H} \vdash v1[\vec{M}, P\{v0[\vec{N}]/x\}, \vec{Q}] : g}$$

where we mark by  $\text{subst}$  an application of the substitution lemma. Again,  $v1, \vec{E}, v0, \vec{F}, \vec{G}, \vec{H}$  is well-defined since  $v, \vec{E}, \vec{F}, \vec{G}, \vec{H}$  is, and that it forms an extension.

- $\backslash\text{-}r$  :  $v[\vec{M}] \rightsquigarrow \lambda x.v0[x, \vec{M}]$ . A typing derivation for the LHS has the form on the left below, so we recover a typing derivation for the RHS on the right below.

$$\frac{\vec{E} \vdash \vec{M} : \Gamma}{v, \vec{E} \vdash v[\vec{M}] : e \backslash f} \quad \frac{\overline{x : e \vdash x : e} \quad \vec{E} \vdash \vec{M} : \Gamma}{v0, x : e, \vec{E} \vdash v0[x, \vec{M}] : f} \quad \frac{v0, \vec{E} \vdash \lambda x.v0[x, \vec{M}] : e \backslash f}{v0, \vec{E} \vdash \lambda x.v0[x, \vec{M}] : e \backslash f}$$

(Note that  $v0, x : e, \vec{E}$  and  $x : e, v0, \vec{E}$  are the same typing environment.) ◀

## F Cut elimination rules

We list all cut-reduction rules. In each case, we sketch the analysis required by Lem. 28, by showing the relevant parts of the head before the cut-reduction ( $\pi$ ) and after ( $\pi'$ ); in those reductions, we assume the cut-reduction took place at a node  $v$  belonging to the head of  $\pi$  (otherwise the lemma holds trivially).

### F.1 Key cases for cut reduction

We first give the key cases for cut elimination, when the two cut formulae are principal for a logical or initial step.

- Identity:

$$\frac{\frac{id}{e \rightarrow e} \quad \frac{\vdots}{\Gamma, e, \Delta \rightarrow f}}{cut \quad \Gamma, e, \Delta \rightarrow f} \quad \mapsto \quad \frac{\vdots}{\Gamma, e, \Delta \rightarrow f}$$

$$\begin{array}{l} v[\vec{M}, N, \vec{O}] \\ \rightsquigarrow v1[\vec{M}, v0[N], \vec{O}] \\ \rightsquigarrow v1[\vec{M}, N, \vec{O}] \end{array} \quad v[\vec{M}, N, \vec{O}]$$

We have  $\pi_{v1} = \pi'_v$ , the head is reduced by two steps, and the cut is removed.

- Product:

$$\frac{\frac{\frac{\vdots}{\Delta \rightarrow e} \quad \frac{\vdots}{\Sigma \rightarrow f}}{cut \quad \Delta, \Sigma \rightarrow ef} \quad \frac{\frac{\vdots}{\Gamma, e, f, \Pi \rightarrow g}}{\Gamma, ef, \Pi \rightarrow g}}{\Gamma, \Delta, \Sigma, \Pi \rightarrow g} \quad \mapsto \quad \frac{\frac{\vdots}{\Delta \rightarrow e} \quad \frac{\frac{\frac{\vdots}{\Sigma \rightarrow f} \quad \frac{\vdots}{\Gamma, e, f, \Pi \rightarrow g}}{cut \quad \Gamma, e, \Sigma, \Pi \rightarrow g}}{cut \quad \Gamma, \Delta, \Sigma, \Pi \rightarrow g}}$$

$$\begin{array}{l} v[\vec{M}, \vec{N}, \vec{O}, \vec{P}] \\ \rightsquigarrow_{cut_{ef}} v1[\vec{M}, v0[\vec{N}, \vec{O}], \vec{P}] \\ \rightsquigarrow v1[\vec{M}, \langle v00[\vec{N}], v01[\vec{O}] \rangle, \vec{P}] \\ \rightsquigarrow^n v1[\vec{M}, \langle N', v01[\vec{O}] \rangle, \vec{P}] \\ \rightsquigarrow^o v1[\vec{M}, \langle N', O' \rangle, \vec{P}] \\ \rightsquigarrow v11[\vec{M}, N', O', \vec{P}] \end{array} \quad \begin{array}{l} v[\vec{M}, \vec{N}, \vec{O}, \vec{P}] \\ \rightsquigarrow_{cut_e} v1[\vec{M}, v0[\vec{N}], \vec{O}, \vec{P}] \\ \rightsquigarrow^n v1[\vec{M}, N', \vec{O}, \vec{P}] \\ \rightsquigarrow_{cut_f} v11[\vec{M}, N', v10[\vec{O}], \vec{P}] \\ \rightsquigarrow^o v11[\vec{M}, N', O', \vec{P}] \end{array}$$

The head is reduced by one step; the cut is replaced by two cuts on occurring closer to the end of the head.

- One:

$$\frac{\frac{\vdots}{\Gamma, \Delta \rightarrow f} \quad \frac{\vdots}{\Gamma, 1, \Delta \rightarrow f}}{\Gamma, \Delta \rightarrow f} \mapsto \frac{\vdots}{\Gamma, \Delta \rightarrow f}$$

$$\begin{array}{l} v[\vec{M}, \vec{N}] \\ \rightsquigarrow_{cut_1} v1[\vec{M}, v0[], \vec{N}] \\ \rightsquigarrow v1[\vec{M}, *, \vec{N}] \\ \rightsquigarrow v10[\vec{M}, \vec{N}] \end{array} \quad v[\vec{M}, \vec{N}]$$

We have  $\pi_{v10} = \pi'_v$ , the head is reduced by three steps, the cut disappears.

- Sum:

$$\frac{\frac{\vdots}{\Delta \rightarrow e_i} \quad \frac{\vdots}{\Gamma, e_0, \Sigma \rightarrow f} \quad \frac{\vdots}{\Gamma, e_1, \Sigma \rightarrow f}}{\Gamma, \Delta, \Sigma \rightarrow f} \mapsto \frac{\frac{\vdots}{\Delta \rightarrow e_i} \quad \frac{\vdots}{\Gamma, e_i, \Sigma \rightarrow f}}{\Gamma, \Delta, \Sigma \rightarrow f}$$

$$\begin{array}{l} v[\vec{M}, \vec{N}, \vec{O}] \\ \rightsquigarrow_{cut_{e_0+e_1}} v1[\vec{M}, v0[\vec{N}], \vec{O}] \\ \rightsquigarrow v1[\vec{M}, in_i v00[\vec{N}], \vec{O}] \\ \rightsquigarrow^n v1[\vec{M}, in_i N', \vec{O}] \\ \rightsquigarrow v1i[\vec{M}, N', \vec{O}] \end{array} \quad \begin{array}{l} v[\vec{M}, \vec{N}, \vec{O}] \\ \rightsquigarrow_{cut_{e_i}} v1[\vec{M}, v0[\vec{N}], \vec{O}] \\ \rightsquigarrow^n v1[\vec{M}, N', \vec{O}] \end{array}$$

We have  $\pi_{v1i} = \pi'_{v1}$ , the head is reduced by two steps, the cut is replaced by a cut occurring closer to the end.

- Zero has no key case since it does not have any right introduction rule.
- Kleene star 1 (which is similar to one):

$$\frac{\frac{\vdots}{\Gamma, \Delta \rightarrow f} \quad \frac{\vdots}{\Gamma, e, e^*, \Delta \rightarrow f}}{\Gamma, \Delta \rightarrow f} \mapsto \frac{\vdots}{\Gamma, \Delta \rightarrow f}$$

$$\begin{array}{l} v[\vec{M}, \vec{N}] \\ \rightsquigarrow_{cut_{e^*}} v1[\vec{M}, v0[], \vec{N}] \\ \rightsquigarrow v1[\vec{M}, [], \vec{N}] \\ \rightsquigarrow v10[\vec{M}, \vec{N}] \end{array} \quad v[\vec{M}, \vec{N}]$$

We have  $\pi_{v10} = \pi'_v$ , the head is reduced by three steps, the cut disappears.

- Kleene star 2 (which is similar to product)

$$\frac{\frac{\vdots}{\Delta \rightarrow e} \quad \frac{\vdots}{\Sigma \rightarrow e^*} \quad \frac{\vdots}{\Gamma, \Pi \rightarrow f} \quad \frac{\vdots}{\Gamma, e, e^*, \Pi \rightarrow f}}{\Gamma, \Delta, \Sigma, \Pi \rightarrow f} \mapsto \frac{\frac{\vdots}{\Delta \rightarrow e} \quad \frac{\vdots}{\Sigma \rightarrow e^*} \quad \frac{\vdots}{\Gamma, e, e^*, \Pi \rightarrow f}}{\Gamma, \Delta, \Sigma, \Pi \rightarrow f}$$

$$\begin{array}{ccc}
v[\vec{M}, \vec{N}, \vec{O}, \vec{P}] & & v[\vec{M}, \vec{N}, \vec{O}, \vec{P}] \\
\rightsquigarrow_{cut_{e^*}} v1[\vec{M}, v0[\vec{N}, \vec{O}], \vec{P}] & \rightsquigarrow_{cut_e} & v1[\vec{M}, v0[\vec{N}], \vec{O}, \vec{P}] \\
\rightsquigarrow v1[\vec{M}, v00[\vec{N}] :: v01[\vec{O}], \vec{P}] & \rightsquigarrow^n & v1[\vec{M}, N', \vec{O}, \vec{P}] \\
\rightsquigarrow^n v1[\vec{M}, N' :: v01[\vec{O}], \vec{P}] & \rightsquigarrow_{cut_{e^*}} & v11[\vec{M}, N', v10[\vec{O}], \vec{P}] \\
\rightsquigarrow^o v1[\vec{M}, N' :: O', \vec{P}] & \rightsquigarrow^o & v11[\vec{M}, N', O', \vec{P}] \\
\rightsquigarrow v11[\vec{M}, N', O', \vec{P}] & & 
\end{array}$$

The head is reduced by one step; the cut is replaced by two cuts occurring closer to the end of the head.

■ Meet:

$$\frac{\frac{\frac{\vdots}{\Delta \rightarrow e_0} \quad \frac{\vdots}{\Delta \rightarrow e_1}}{\Delta \rightarrow e_0 \cap e_1} \quad \frac{\frac{\vdots}{\Gamma, e_i, \Sigma \rightarrow f}}{\Gamma, e_0 \cap e_1, \Sigma \rightarrow f}}{cut \quad \Gamma, \Delta, \Sigma \rightarrow f} \mapsto \frac{\frac{\vdots}{\Delta \rightarrow e_i} \quad \frac{\vdots}{\Gamma, e_i, \Sigma \rightarrow f}}{cut \quad \Gamma, \Delta, \Sigma \rightarrow f}$$

$$\begin{array}{ccc}
v[\vec{M}, \vec{N}, \vec{O}] & & v[\vec{M}, \vec{N}, \vec{O}] \\
\rightsquigarrow_{cut_{e_0 \cap e_1}} v1[\vec{M}, v0[\vec{N}], \vec{O}] & & v1[\vec{M}, v0[\vec{N}], \vec{O}] \\
\rightsquigarrow v1[\vec{M}, \langle v00[\vec{N}], v01[\vec{N}] \rangle, \vec{O}] & \rightsquigarrow_{cut_{e_i}} & v1[\vec{M}, v0[\vec{N}], \vec{O}] \\
\rightsquigarrow v10[\vec{M}, v0i[\vec{N}], \vec{O}] & & 
\end{array}$$

The head is reduced by two steps; the cut is replaced by a cut occurring closer to the end.

■ Left residual:

$$\frac{\frac{\frac{\vdots}{e, \Sigma \rightarrow f} \quad \frac{\vdots}{\Delta \rightarrow e} \quad \frac{\vdots}{\Gamma, f, \Pi \rightarrow g}}{\Sigma \rightarrow e \setminus f} \quad \frac{\frac{\vdots}{\Gamma, \Delta, e \setminus f, \Pi \rightarrow g}}{\Gamma, \Delta, \Sigma, \Pi \rightarrow g}}{cut \quad \Gamma, \Delta, \Sigma, \Pi \rightarrow g} \mapsto \frac{\frac{\vdots}{\Delta \rightarrow e} \quad \frac{\frac{\vdots}{e, \Sigma \rightarrow f} \quad \frac{\vdots}{\Gamma, f, \Pi \rightarrow g}}{cut \quad \Gamma, e, \Sigma, \Pi \rightarrow g}}{cut \quad \Gamma, \Delta, \Sigma, \Pi \rightarrow g}$$

$$\begin{array}{ccc}
v[\vec{M}, \vec{N}, \vec{O}, \vec{P}] & & v[\vec{M}, \vec{N}, \vec{O}, \vec{P}] \\
\rightsquigarrow_{cut_{e \setminus f}} v1[\vec{M}, \vec{N}, v0[\vec{O}], \vec{P}] & \rightsquigarrow_{cut_e} & v1[\vec{M}, v0[\vec{N}], \vec{O}, \vec{P}] \\
\rightsquigarrow v1[\vec{M}, \vec{N}, \lambda x. v00[x, \vec{O}], \vec{P}] & \rightsquigarrow^n & v1[\vec{M}, N', \vec{O}, \vec{P}] \\
\rightsquigarrow v11[\vec{M}, v00[v10[\vec{N}], \vec{O}], \vec{P}] & \rightsquigarrow_{cut_f} & v11[\vec{M}, v10[N', \vec{O}], \vec{P}] \\
\rightsquigarrow^n v11[\vec{M}, v00[N', \vec{O}], \vec{P}] & & 
\end{array}$$

The head is reduced by one; the cut is replaced by two cuts occurring closer to the end. When working on the last cut visited by the head of  $\pi$ , the  $n$  reductions leading to  $N'$  do not visit any cut so that the weight decreases even though those reductions are moved one step away from the end of the head.

■ Right residual: symmetrical to the left residual

## F.2 Commutative cases

We now give the commutative cases for cut-elimination, when at least one of the cut formulae is not principal for a logical step. We give the relevant reductions of the head as before; when the logical step is a left-rule we only display the case where the program provided in the corresponding principal position enables the corresponding (program) reduction rule: otherwise the head of the reduced proof is blocked early, and the Lem. 28 is trivially satisfied.





The length of the head is conserved and the cut is moved  $m + 1$  steps towards the end. Note that the  $o$  steps leading to  $O'$  and the  $m$  steps leading to  $M'$  are swapped by this cut-reduction. This is why it is important for the third point of Lem. 28 that we consider the last cut of the head: under this assumption, the  $m$  steps leading to  $M'$  do not visit any cut (whose distance to the end of the head would increase).

■ Left 1- $l$ :

$$\frac{\frac{\frac{\vdots}{\Delta, \Sigma \rightarrow g}}{\Delta, 1, \Sigma \rightarrow g} \quad \frac{\vdots}{\Gamma, g, \Pi \rightarrow h}}{\Gamma, \Delta, 1, \Sigma, \Pi \rightarrow h} \text{cut}^{1-l} \mapsto \frac{\frac{\frac{\vdots}{\Delta, \Sigma \rightarrow g} \quad \frac{\vdots}{\Gamma, g, \Pi \rightarrow h}}{\Gamma, \Delta, \Sigma, \Pi \rightarrow h}}{\Gamma, \Delta, 1, \Sigma, \Pi \rightarrow h} \text{cut}^{1-l}$$

$$\begin{array}{ccc} v[\vec{M}, \vec{N}, \star, \vec{P}, \vec{Q}] & & v[\vec{M}, \vec{N}, \star, \vec{P}, \vec{Q}] \\ \rightsquigarrow_{\text{cut}} v1[\vec{M}, v0[\vec{N}, \star, \vec{P}], \vec{Q}] & \rightsquigarrow & v[\vec{M}, \vec{N}, \vec{P}, \vec{Q}] \\ \rightsquigarrow v1[\vec{M}, v00[\vec{N}, \vec{P}], \vec{Q}] & \rightsquigarrow_{\text{cut}} & v01[\vec{M}, v00[\vec{N}, \vec{P}], \vec{Q}] \end{array}$$

The length of the head is conserved; the cut is moved one step towards the end.

■ Right 1- $l$ :

$$\frac{\frac{\frac{\vdots}{\Sigma \rightarrow g} \quad \frac{\frac{\vdots}{\Gamma, \Delta, g, \Pi \rightarrow h}}{\Gamma, 1, \Delta, g, \Pi \rightarrow h}}{\Gamma, 1, \Delta, \Sigma, \Pi \rightarrow h} \text{cut}^{1-l} \mapsto \frac{\frac{\frac{\vdots}{\Sigma \rightarrow g} \quad \frac{\vdots}{\Gamma, \Delta, g, \Pi \rightarrow h}}{\Gamma, \Delta, \Sigma, \Pi \rightarrow h}}{\Gamma, 1, \Delta, \Sigma, \Pi \rightarrow h} \text{cut}^{1-l}$$

$$\begin{array}{ccc} v[\vec{M}, \star, \vec{O}, \vec{P}, \vec{Q}] & & v[\vec{M}, \star, \vec{O}, \vec{P}, \vec{Q}] \\ \rightsquigarrow_{\text{cut}} v1[\vec{M}, \star, \vec{O}, v0[\vec{P}], \vec{Q}] & \rightsquigarrow & v1[\vec{M}, \vec{O}, \vec{P}, \vec{Q}] \\ \rightsquigarrow^P v1[\vec{M}, \star, \vec{O}, P', \vec{Q}] & \rightsquigarrow_{\text{cut}} & v10[\vec{M}, \vec{O}, v10[\vec{P}], \vec{Q}] \\ \rightsquigarrow v10[\vec{M}, \vec{O}, P', \vec{Q}] & \rightsquigarrow^P & v10[\vec{M}, \vec{O}, P', \vec{Q}] \end{array}$$

The length of the head is conserved; the cut is moved one step towards the end. (Analogous case if the cut formula appears on the other side of 1.)

■ Rule 1- $r$  cannot appear on the right of a cut.

■ Left +- $l$ :

$$\frac{\frac{\frac{\vdots}{\Delta, e_0, \Sigma \rightarrow g} \quad \frac{\vdots}{\Delta, e_1, \Sigma \rightarrow g}}{\Delta, e_0 + e_1, \Sigma \rightarrow g} \quad \frac{\vdots}{\Gamma, g, \Pi \rightarrow h}}{\Gamma, \Delta, e_0 + e_1, \Sigma, \Pi \rightarrow h} \text{cut}^{+-l} \mapsto \frac{\frac{\frac{\vdots}{\Delta, e_0, \Sigma \rightarrow g} \quad \frac{\vdots}{\Gamma, g, \Pi \rightarrow h}}{\Gamma, \Delta, e_0, \Sigma, \Pi \rightarrow h} \quad \frac{\frac{\vdots}{\Delta, e_1, \Sigma \rightarrow g} \quad \frac{\vdots}{\Gamma, g, \Pi \rightarrow h}}{\Gamma, \Delta, e_1, \Sigma, \Pi \rightarrow h} \text{cut}^{+-l}}{\Gamma, \Delta, e_0 + e_1, \Sigma, \Pi \rightarrow h}$$

$$\begin{array}{ccc} v[\vec{M}, \vec{N}, \text{in}_i O, \vec{P}, \vec{Q}] & & v[\vec{M}, \vec{N}, \text{in}_i O, \vec{P}, \vec{Q}] \\ \rightsquigarrow_{\text{cut}} v1[\vec{M}, v0[\vec{N}, \text{in}_i O, \vec{P}], \vec{Q}] & \rightsquigarrow & vi[\vec{M}, \vec{N}, O, \vec{P}, \vec{Q}] \\ \rightsquigarrow v1[\vec{M}, v0i[\vec{N}, O, \vec{P}], \vec{Q}] & \rightsquigarrow_{\text{cut}} & vi1[\vec{M}, vi0[\vec{N}, O, \vec{P}], \vec{Q}] \end{array}$$

The length of the head is conserved; the cut is moved one step towards the end.

■ Right +- $l$ :

$$\frac{\frac{\frac{\vdots}{\Sigma \rightarrow g} \quad \frac{\frac{\vdots}{\Gamma, e_0, \Delta, g, \Pi \rightarrow h} \quad \frac{\vdots}{\Gamma, e_1, \Delta, g, \Pi \rightarrow h}}{\Gamma, e_0 + e_1, \Delta, g, \Pi \rightarrow h}}{\Gamma, e_0 + e_1, \Delta, \Sigma, \Pi \rightarrow h} \text{cut}^{+-l} \mapsto \frac{\frac{\frac{\vdots}{\Sigma \rightarrow g} \quad \frac{\vdots}{\Gamma, e_0, \Delta, g, \Pi \rightarrow h}}{\Gamma, e_0, \Sigma, \Pi \rightarrow h} \quad \frac{\frac{\vdots}{\Sigma \rightarrow g} \quad \frac{\vdots}{\Gamma, e_1, \Delta, g, \Pi \rightarrow h}}{\Gamma, e_1, \Sigma, \Pi \rightarrow h} \text{cut}^{+-l}}{\Gamma, e_0 + e_1, \Delta, \Sigma, \Pi \rightarrow h}$$

$$\begin{array}{ccc}
v[\vec{M}, \text{in}_i N, \vec{O}, \vec{P}, \vec{Q}] & & v[\vec{M}, \text{in}_i N, \vec{O}, \vec{P}, \vec{Q}] \\
\rightsquigarrow_{cut} v1[\vec{M}, \text{in}_i N, \vec{O}, v0[\vec{P}], \vec{Q}] & \rightsquigarrow & vi[\vec{M}, N, \vec{O}, \vec{P}, \vec{Q}] \\
\rightsquigarrow^P v1[\vec{M}, \text{in}_i N, \vec{O}, P', \vec{Q}] & \rightsquigarrow_{cut} & vi1[\vec{M}, N, \vec{O}, v10[\vec{P}], \vec{Q}] \\
\rightsquigarrow v1i[\vec{M}, N, \vec{O}, P', \vec{Q}] & \rightsquigarrow^P & vi1[\vec{M}, N, \vec{O}, P', \vec{Q}]
\end{array}$$

The length of the head is conserved; the cut is moved one step towards the end.  
(Analogous case if the cut formula appears on the other side of the sum.)

■ Right  $+r_i$ :

$$\text{cut} \frac{\frac{\vdots}{\Delta \rightarrow g} \quad \frac{\frac{\vdots}{\Gamma, g, \Sigma \rightarrow e_i}}{\Gamma, g, \Sigma \rightarrow e_0 + e_1}}{\Gamma, \Delta, \Sigma, e_0 + e_1} \quad +r_i \quad \mapsto \quad \text{cut} \frac{\frac{\vdots}{\Delta \rightarrow g} \quad \frac{\vdots}{\Gamma, g, \Sigma \rightarrow e_i}}{\Gamma, \Delta, \Sigma \rightarrow e_i} \quad +r_i \frac{\vdots}{\Gamma, \Delta, \Sigma \rightarrow e_0 + e_1}$$

$$\begin{array}{ccc}
v[\vec{M}, \vec{N}, \vec{O}] & & v[\vec{M}, \vec{N}, \vec{O}] \\
\rightsquigarrow_{cut} v1[\vec{M}, v0[\vec{N}], \vec{O}] & \rightsquigarrow & \text{in}_i v1[\vec{M}, \vec{N}, \vec{O}] \\
\rightsquigarrow^n v1[\vec{M}, N', \vec{O}] & \rightsquigarrow_{cut} & \text{in}_i v11[\vec{M}, v10[\vec{N}], \vec{O}] \\
\rightsquigarrow \text{in}_i v11[\vec{M}, N', \vec{O}] & \rightsquigarrow^n & \text{in}_i v11[\vec{M}, N', \vec{O}]
\end{array}$$

The length of the head is conserved; the cut is moved one step towards the end.

■ Left and right  $0-l$  are easy: they just produce a new  $0-l$  axiom; the length of the head is reduced (to one), the cut disappears.

■ Left  $*-l$ :

$$*-l \frac{\frac{\vdots}{\Delta, \Sigma \rightarrow f} \quad \frac{\frac{\vdots}{\Delta, e, e^*, \Sigma \rightarrow f}}{\Gamma, \Delta, e^*, \Sigma, \Pi \rightarrow g} \quad \frac{\vdots}{\Gamma, f, \Pi \rightarrow g}}{\Gamma, \Delta, e^*, \Sigma, \Pi \rightarrow g} \quad \text{cut} \quad \mapsto \quad \text{cut} \frac{\frac{\vdots}{\Delta, \Sigma \rightarrow f} \quad \frac{\vdots}{\Gamma, f, \Pi \rightarrow g}}{\Gamma, \Delta, \Sigma, \Pi \rightarrow g} \quad \text{cut} \frac{\frac{\vdots}{\Delta, e, e^*, \Sigma \rightarrow f} \quad \frac{\vdots}{\Gamma, f, \Pi \rightarrow g}}{\Gamma, \Delta, e, e^*, \Sigma, \Pi \rightarrow g} \quad *-l \frac{\vdots}{\Gamma, \Delta, e^*, \Sigma, \Pi \rightarrow g}$$

There are two cases depending on the value for  $e^*$ :

■  $\square$  (similar to left  $1-l$ ):

$$\begin{array}{ccc}
v[\vec{M}, \vec{N}, \square, \vec{P}, \vec{Q}] & & v[\vec{M}, \vec{N}, \square, \vec{P}, \vec{Q}] \\
\rightsquigarrow_{cut} v1[\vec{M}, v0[\vec{N}, \square, \vec{P}], \vec{Q}] & \rightsquigarrow & v0[\vec{M}, \vec{N}, \vec{P}, \vec{Q}] \\
\rightsquigarrow v1[\vec{M}, v00[\vec{N}, \vec{P}], \vec{Q}] & \rightsquigarrow_{cut} & v01[\vec{M}, v00[\vec{N}, \vec{P}], \vec{Q}]
\end{array}$$

■  $O :: U$  (similar to left  $-l$ ):

$$\begin{array}{ccc}
v[\vec{M}, \vec{N}, O :: U, \vec{P}, \vec{Q}] & & v[\vec{M}, \vec{N}, O :: U, \vec{P}, \vec{Q}] \\
\rightsquigarrow_{cut} v1[\vec{M}, v0[\vec{N}, O :: U, \vec{P}], \vec{Q}] & \rightsquigarrow & v1[\vec{M}, \vec{N}, O, U\vec{P}, \vec{Q}] \\
\rightsquigarrow v1[\vec{M}, v01[\vec{N}, O, U\vec{P}], \vec{Q}] & \rightsquigarrow_{cut} & v11[\vec{M}, v10[\vec{N}, O, U, \vec{P}], \vec{Q}]
\end{array}$$

In both cases the length of the head is conserved and the cut is moved one step towards the end.

■ Right  $*-l$ :

$$\text{cut} \frac{\frac{\vdots}{\Delta \rightarrow f} \quad \frac{\frac{\vdots}{\Gamma, f, \Sigma, \Pi \rightarrow g} \quad \frac{\vdots}{\Gamma, f, \Sigma, e, e^*, \Pi \rightarrow g}}{\Gamma, f, \Sigma, e^*, \Pi \rightarrow g}}{\Gamma, \Delta, \Sigma, e^*, \Pi \rightarrow g} \quad \mapsto \quad \text{cut} \frac{\frac{\vdots}{\Delta \rightarrow f} \quad \frac{\vdots}{\Gamma, f, \Sigma, \Pi \rightarrow g}}{\Gamma, \Delta, \Sigma, \Pi \rightarrow g} \quad \text{cut} \frac{\frac{\vdots}{\Delta \rightarrow f} \quad \frac{\vdots}{\Gamma, f, \Sigma, \Pi, e, e^* \rightarrow g}}{\Gamma, \Delta, \Sigma, e, e^*, \Pi \rightarrow g} \quad *-l \frac{\vdots}{\Gamma, \Delta, \Sigma, e^*, \Pi \rightarrow g}$$

Like previously, there are two cases depending on the value for  $e^*$ :

- $\square$  (similar to right 1- $l$ ):

$$\begin{array}{ccc}
v[\vec{M}, \vec{N}, \vec{P}, \square, \vec{Q}] & & v[\vec{M}, \vec{N}, \vec{P}, \square, \vec{Q}] \\
\rightsquigarrow_{cut} v1[\vec{M}, v0[\vec{N}], \vec{P}, \square, \vec{Q}] & \rightsquigarrow & v0[\vec{M}, \vec{N}, \vec{P}, \vec{Q}] \\
\rightsquigarrow^n v1[\vec{M}, N', \vec{P}, \square, \vec{Q}] & \rightsquigarrow_{cut} & v01[\vec{M}, v00[\vec{N}], \vec{P}, \vec{Q}] \\
\rightsquigarrow v10[\vec{M}, N', \vec{P}, \vec{Q}] & \rightsquigarrow^n & v01[\vec{M}, N', \vec{P}, \vec{Q}]
\end{array}$$

- $O :: U$  (similar to right  $\cdot$ - $l$ ):

$$\begin{array}{ccc}
v[\vec{M}, \vec{N}, \vec{P}, O :: U, \vec{Q}] & & v[\vec{M}, \vec{N}, \vec{P}, O :: U, \vec{Q}] \\
\rightsquigarrow_{cut} v1[\vec{M}, v0[\vec{N}], \vec{P}, O :: U, \vec{Q}] & \rightsquigarrow & v1[\vec{M}, \vec{N}, \vec{P}, O, U, \vec{Q}] \\
\rightsquigarrow^n v1[\vec{M}, N', \vec{P}, O :: U, \vec{Q}] & \rightsquigarrow_{cut} & v11[\vec{M}, v10[\vec{N}], \vec{P}, O, U, \vec{Q}] \\
\rightsquigarrow v11[\vec{M}, N', \vec{P}, O, U, \vec{Q}] & \rightsquigarrow^n & v11[\vec{M}, N', \vec{P}, O, U, \vec{Q}]
\end{array}$$

In both cases the length of the head is conserved and the cut is moved one step towards the end.

- Rule  $\ast$ - $r_1$  cannot appear on the right of a cut.
- Right  $\ast$ - $r_2$  (similar to right  $\cdot$ - $r$ ):

$$\begin{array}{ccc}
\begin{array}{c} \vdots \\ \vdots \\ \vdots \\ \vdots \end{array} & & \begin{array}{c} \vdots \\ \vdots \\ \vdots \\ \vdots \end{array} \\
\frac{\frac{\frac{\Delta \rightarrow e}{\vdots}}{\Gamma, e, \Sigma \rightarrow f} \quad \frac{\frac{\Pi \rightarrow f^*}{\vdots}}{\Gamma, e, \Sigma, \Pi \rightarrow f^*}}{\Gamma, \Delta, \Sigma, \Pi \rightarrow f^*} \text{cut} & \mapsto & \frac{\frac{\frac{\Delta \rightarrow e}{\vdots}}{\Gamma, \Delta, \Sigma \rightarrow f} \quad \frac{\frac{\Pi \rightarrow f^*}{\vdots}}{\Gamma, \Delta, \Sigma, \Pi \rightarrow f^*}}{\Gamma, \Delta, \Sigma, \Pi \rightarrow f^*} \text{cut} \\
\ast\text{-}r_2 & & \ast\text{-}r_2
\end{array}$$

$$\begin{array}{ccc}
v[\vec{M}, \vec{N}, \vec{O}, \vec{P}] & & v[\vec{M}, \vec{N}, \vec{O}, \vec{P}] \\
\rightsquigarrow_{cut} v1[\vec{M}, v0[\vec{N}], \vec{O}, \vec{P}] & \rightsquigarrow & v0[\vec{M}, \vec{N}, \vec{O}] :: v1[\vec{P}] \\
\rightsquigarrow^n v1[\vec{M}, N', \vec{O}, \vec{P}] & \rightsquigarrow_{cut} & v01[\vec{M}, v00[\vec{N}], \vec{O}] :: v1[\vec{P}] \\
\rightsquigarrow v10[\vec{M}, N', \vec{O}] :: v11[\vec{P}] & \rightsquigarrow^n & v01[\vec{M}, N', \vec{O}] :: v1[\vec{P}]
\end{array}$$

The length of the head is conserved and the cut is moved one step towards the end.

- Other right  $\ast$ - $r_2$  (similar to other right  $\cdot$ - $r$ ):

$$\begin{array}{ccc}
\begin{array}{c} \vdots \\ \vdots \\ \vdots \\ \vdots \end{array} & & \begin{array}{c} \vdots \\ \vdots \\ \vdots \\ \vdots \end{array} \\
\frac{\frac{\frac{\Sigma \rightarrow e}{\vdots}}{\Gamma \rightarrow f} \quad \frac{\frac{\Delta, e, \Pi \rightarrow f^*}{\vdots}}{\Gamma, \Delta, e, \Pi \rightarrow f^*}}{\Gamma, \Delta, \Sigma, \Pi \rightarrow f^*} \text{cut} & \mapsto & \frac{\frac{\frac{\Sigma \rightarrow e}{\vdots}}{\Gamma \rightarrow f} \quad \frac{\frac{\Delta, e, \Pi \rightarrow f^*}{\vdots}}{\Delta, \Sigma, \Pi \rightarrow f^*}}{\Gamma, \Delta, \Sigma, \Pi \rightarrow f^*} \text{cut} \\
\ast\text{-}r_2 & & \ast\text{-}r_2
\end{array}$$

$$\begin{array}{ccc}
v[\vec{M}, \vec{N}, \vec{O}, \vec{P}] & & v[\vec{M}, \vec{N}, \vec{O}, \vec{P}] \\
\rightsquigarrow_{cut} v1[\vec{M}, \vec{N}, v0[\vec{O}], \vec{P}] & \rightsquigarrow & v0[\vec{M}] :: v1[\vec{N}, \vec{O}, \vec{P}] \\
\rightsquigarrow^o v1[\vec{M}, N', O', \vec{P}] & \rightsquigarrow^m & M' :: v1[\vec{N}, \vec{O}, \vec{P}] \\
\rightsquigarrow v10[\vec{M}] :: v11[N', O', \vec{P}] & \rightsquigarrow_{cut} & M' :: v11[\vec{N}, v10[\vec{O}], \vec{P}] \\
\rightsquigarrow^m M' :: v11[N', O', \vec{P}] & \rightsquigarrow^o & M' :: v11[\vec{N}, O', \vec{P}]
\end{array}$$

The length of the head is conserved and the cut is moved  $m + 1$  steps towards the end. Like in the other right  $\cdot$ - $r$  case, the  $o$  steps leading to  $O'$  and the  $m$  steps leading to  $M'$  are swapped by this cut-reduction, which is fine when working on the last cut of the head.

■ Left  $\cap\text{-}l_i$ :

$$\begin{array}{c}
\vdots \\
\frac{\Delta, e_i, \Sigma \rightarrow f}{\cap\text{-}l_i} \quad \frac{\vdots}{\Gamma, f, \Pi \rightarrow g} \\
\frac{\Delta, e_0 \cap e_1, \Sigma \rightarrow f \quad \Gamma, f, \Pi \rightarrow g}{\text{cut}} \\
\Gamma, \Delta, e_0 \cap e_1, \Sigma, \Pi \rightarrow g
\end{array}
\mapsto
\begin{array}{c}
\vdots \quad \vdots \\
\frac{\Delta, e_i, \Sigma \rightarrow f \quad \Gamma, f, \Pi \rightarrow g}{\text{cut}} \\
\frac{\Gamma, \Delta, e_i, \Sigma, \Pi \rightarrow g}{\cap\text{-}l_i} \\
\Gamma, \Delta, e_0 \cap e_1, \Sigma, \Pi \rightarrow g
\end{array}$$

$$\begin{array}{ccc}
v[\vec{M}, \vec{N}, \langle\langle O_0, O_1 \rangle\rangle, \vec{P}, \vec{Q}] & & v[\vec{M}, \vec{N}, \langle\langle O_0, O_1 \rangle\rangle, \vec{P}, \vec{Q}] \\
\rightsquigarrow_{\text{cut}} v1[\vec{M}, v0[\vec{N}, \langle\langle O_0, O_1 \rangle\rangle, \vec{P}], \vec{Q}] & & \rightsquigarrow v0[\vec{M}, \vec{N}, O_i, \vec{P}, \vec{Q}] \\
\rightsquigarrow v10[\vec{M}, v0[\vec{N}, O_i, \vec{P}], \vec{Q}] & & \rightsquigarrow^o v0[\vec{M}, \vec{N}, O'_i, \vec{P}, \vec{Q}] \\
\rightsquigarrow^o v10[\vec{M}, v0[\vec{N}, O'_i, \vec{P}], \vec{Q}] & & \rightsquigarrow_{\text{cut}} v01[\vec{M}, v00[\vec{N}, O'_i, \vec{P}], \vec{Q}]
\end{array}$$

Note that while the programs  $\vec{M}, \vec{N}, \vec{P}, \vec{Q}$  are assumed to be irreducible thanks to the innermost reduction strategy,  $O_0$  and  $O_1$  are not, whence the reduction from  $O_i$  to  $O'_i$ . The length of the head is conserved and the cut is moved  $o + 1$  steps towards the end. If this cut was the last one visited in the head of  $\pi$  then no cuts are visited during the reduction of  $O_i$ , so that the overall weight indeed decreases.

■ Right  $\cap\text{-}l_i$ :

$$\begin{array}{c}
\vdots \\
\frac{\vdots}{\Delta \rightarrow f} \quad \frac{\Gamma, f, \Sigma, e_i, \Pi \rightarrow g}{\cap\text{-}l_i} \\
\frac{\Delta \rightarrow f \quad \Gamma, f, \Sigma, e_0 \cap e_1, \Pi \rightarrow g}{\text{cut}} \\
\Gamma, \Delta, \Sigma, e_0 \cap e_1, \Pi \rightarrow g
\end{array}
\mapsto
\begin{array}{c}
\vdots \quad \vdots \\
\frac{\Delta \rightarrow f \quad \Gamma, f, \Sigma, e_i, \Pi \rightarrow g}{\text{cut}} \\
\frac{\Gamma, \Delta, \Sigma, e_i, \Pi \rightarrow g}{\cap\text{-}l_i} \\
\Gamma, \Delta, \Sigma, e_0 \cap e_1, \Pi \rightarrow g
\end{array}$$

$$\begin{array}{ccc}
v[\vec{M}, \vec{N}, \vec{O}, \langle\langle P_0, P_1 \rangle\rangle, \vec{Q}] & & v[\vec{M}, \vec{N}, \vec{O}, \langle\langle P_0, P_1 \rangle\rangle, \vec{Q}] \\
\rightsquigarrow_{\text{cut}} v1[\vec{M}, v0[\vec{N}, \vec{O}, \langle\langle P_0, P_1 \rangle\rangle, \vec{Q}]] & & \rightsquigarrow v1[\vec{M}, \vec{N}, \vec{O}, P_i, \vec{Q}] \\
\rightsquigarrow^n v1[\vec{M}, N', \vec{O}, \langle\langle P_0, P_1 \rangle\rangle, \vec{Q}] & & \rightsquigarrow^p v1[\vec{M}, \vec{N}, \vec{O}, P'_i, \vec{Q}] \\
\rightsquigarrow v11[\vec{M}, N', \vec{O}, P_i, \vec{Q}] & & \rightsquigarrow_{\text{cut}} v11[\vec{M}, v10[\vec{N}, \vec{O}, P'_i, \vec{Q}]] \\
\rightsquigarrow^p v11[\vec{M}, N', \vec{O}, P'_i, \vec{Q}] & & \rightsquigarrow^n v11[\vec{M}, N', \vec{O}, P'_i, \vec{Q}]
\end{array}$$

The length of the head is preserved and the cut is moved  $p + 1$  steps towards the end. Reductions yielding  $P'_i$  and  $N'$  are swapped so that it is again crucial to operate on the last cut to decrease the weight.

(Analogous case if the cut formula appears on the other side of the meet.)

■ Right  $\cap\text{-}r$ :

$$\begin{array}{c}
\vdots \\
\frac{\vdots}{\Delta \rightarrow f} \quad \frac{\Gamma, f, \Sigma \rightarrow e_0 \quad \Gamma, f, \Sigma \rightarrow e_1}{\cap\text{-}r} \\
\frac{\Delta \rightarrow f \quad \Gamma, f, \Sigma \rightarrow e_0 \cap e_1}{\text{cut}} \\
\Gamma, \Delta, \Sigma \rightarrow e_0 \cap e_1
\end{array}
\mapsto
\begin{array}{c}
\vdots \quad \vdots \quad \vdots \quad \vdots \\
\frac{\Delta \rightarrow f \quad \Gamma, f, \Sigma \rightarrow e_0}{\text{cut}} \quad \frac{\Delta \rightarrow f \quad \Gamma, f, \Sigma \rightarrow e_1}{\text{cut}} \\
\frac{\Gamma, \Delta, \Sigma \rightarrow e_0 \quad \Gamma, \Delta, \Sigma \rightarrow e_1}{\cap\text{-}r} \\
\Gamma, \Delta, \Sigma \rightarrow e_0 \cap e_1
\end{array}$$

$$\begin{array}{ccc}
v[\vec{M}, \vec{N}, \vec{O}] & & v[\vec{M}, \vec{N}, \vec{O}] \\
\rightsquigarrow_{\text{cut}} v1[\vec{M}, v0[\vec{N}, \vec{O}]] & & \rightsquigarrow \langle\langle v0[\vec{M}, \vec{N}, \vec{O}], v1[\vec{M}, \vec{N}, \vec{O}] \rangle\rangle \\
\rightsquigarrow^n v1[\vec{M}, N', \vec{O}] & & \\
\rightsquigarrow \langle\langle v10[\vec{M}, N', \vec{O}], v11[\vec{M}, N', \vec{O}] \rangle\rangle & & 
\end{array}$$

As explained in the main text, either the head of  $\pi$  does not touch the produced pair, in which case the length of the head is decreased by  $n + 1$  and the cut has been removed;

or it destructs the pair using a  $\cap\text{-}l_i$  program reduction, at which point the head of  $\pi'$  catches up with  $1 + n$  steps (the blocked cut followed by the reductions yielding  $N'$ ). In the latter case, the length of the run is preserved and the cut is moved many steps towards the end; when operating on the last cut of the head, the other cuts did not move.

■ Left  $\setminus\text{-}l$ :

$$\frac{\frac{\frac{\vdots}{\Delta_2 \rightarrow e} \quad \frac{\vdots}{\Delta_1, f, \Sigma \rightarrow g}}{\text{cut} \frac{\Delta_1, \Delta_2, e \setminus f, \Sigma \rightarrow g}{\Gamma, \Delta_1, \Delta_2, e \setminus f, \Sigma, \Pi \rightarrow h}} \quad \frac{\vdots}{\Gamma, g, \Pi \rightarrow h}}{\setminus\text{-}l} \mapsto \frac{\frac{\vdots}{\Delta_2 \rightarrow e} \quad \frac{\frac{\vdots}{\Delta_1, f, \Sigma \rightarrow g} \quad \frac{\vdots}{\Gamma, g, \Pi \rightarrow h}}{\text{cut} \frac{\Delta_1, f, \Sigma \rightarrow g \quad \Gamma, g, \Pi \rightarrow h}{\Gamma, \Delta_1, f, \Sigma, \Pi \rightarrow h}}}{\setminus\text{-}l} \frac{\Gamma, \Delta_1, \Delta_2, e \setminus f, \Sigma, \Pi \rightarrow h}{\Gamma, \Delta_1, \Delta_2, e \setminus f, \Sigma, \Pi \rightarrow h}}$$

$$\begin{array}{ll} \rightsquigarrow_{\text{cut}} & v[\vec{M}, \vec{N}, \vec{O}, \lambda x.F, \vec{P}, \vec{Q}] \\ \rightsquigarrow & v1[\vec{M}, v0[\vec{N}, \vec{O}, \lambda x.F, \vec{P}], \vec{Q}] \\ \rightsquigarrow^r & v1[\vec{M}, v01[\vec{N}, F\{v00[\vec{O}]/x\}, \vec{P}], \vec{Q}] \\ \rightsquigarrow^{\text{cut}} & v1[\vec{M}, v01[\vec{N}, R, \vec{P}], \vec{Q}] \end{array} \quad \begin{array}{ll} & v[\vec{M}, \vec{N}, \vec{O}, \lambda x.F, \vec{P}, \vec{Q}] \\ \rightsquigarrow & v1[\vec{M}, \vec{N}, F\{v0[\vec{O}]/x\}, \vec{P}, \vec{Q}] \\ \rightsquigarrow^r & v1[\vec{M}, \vec{N}, R, \vec{P}, \vec{Q}] \\ \rightsquigarrow_{\text{cut}} & v11[\vec{M}, v10[\vec{N}, R, \vec{P}], \vec{Q}] \end{array}$$

The length of the run is preserved and the cut is moved  $r + 1$  steps towards the end; when operating on the last cut of the head, the other cuts did not move.

■ Right  $\setminus\text{-}l$ :

$$\frac{\frac{\frac{\vdots}{\Sigma \rightarrow g} \quad \frac{\frac{\vdots}{\Delta_1, g, \Delta_2 \rightarrow e} \quad \frac{\vdots}{\Gamma, f, \Pi \rightarrow h}}{\setminus\text{-}l} \frac{\Gamma, \Delta_1, g, \Delta_2, e \setminus f, \Pi \rightarrow h}{\Gamma, \Delta_1, \Sigma, \Delta_2, e \setminus f, \Pi \rightarrow h}}{\text{cut} \frac{\Sigma \rightarrow g \quad \Gamma, \Delta_1, \Sigma, \Delta_2, e \setminus f, \Pi \rightarrow h}}{\Gamma, \Delta_1, \Sigma, \Delta_2, e \setminus f, \Pi \rightarrow h}} \mapsto \frac{\frac{\frac{\vdots}{\Sigma \rightarrow g} \quad \frac{\frac{\vdots}{\Delta_1, g, \Delta_2 \rightarrow e} \quad \frac{\vdots}{\Gamma, f, \Pi \rightarrow h}}{\setminus\text{-}l} \frac{\Gamma, f, \Pi \rightarrow h}{\Gamma, \Delta_1, \Sigma, \Delta_2, e \setminus f, \Pi \rightarrow h}}{\text{cut} \frac{\Sigma \rightarrow g \quad \Delta_1, g, \Delta_2 \rightarrow e}{\Gamma, \Delta_1, \Sigma, \Delta_2, e \setminus f, \Pi \rightarrow h}}}{\setminus\text{-}l} \frac{\Gamma, \Delta_1, \Sigma, \Delta_2, e \setminus f, \Pi \rightarrow h}{\Gamma, \Delta_1, \Sigma, \Delta_2, e \setminus f, \Pi \rightarrow h}}$$

$$\begin{array}{ll} \rightsquigarrow_{\text{cut}} & v[\vec{M}, \vec{N}, \vec{O}, \vec{P}, \lambda x.F, \vec{Q}] \\ \rightsquigarrow^o & v1[\vec{M}, \vec{N}, v0[\vec{O}], \vec{P}, \lambda x.F, \vec{Q}] \\ \rightsquigarrow & v1[\vec{M}, \vec{N}, O', \vec{P}, \lambda x.F, \vec{Q}] \\ \rightsquigarrow & v11[\vec{M}, F\{v10[\vec{N}, O', \vec{P}]/x\}, \vec{Q}] \\ \rightsquigarrow^f & v11[\vec{M}, F'\{v10[\vec{N}, O', \vec{P}]/x\}, \vec{Q}] \end{array} \quad \begin{array}{ll} & v[\vec{M}, \vec{N}, \vec{O}, \vec{P}, \lambda x.F, \vec{Q}] \\ \rightsquigarrow & v1[\vec{M}, F\{v0[\vec{N}, \vec{O}, \vec{P}]/x\}, \vec{Q}] \\ \rightsquigarrow^f & v1[\vec{M}, F'\{v0[\vec{N}, \vec{O}, \vec{P}]/x\}, \vec{Q}] \\ \rightsquigarrow_{\text{cut}} & v1[\vec{M}, F'\{v01[\vec{N}, v00[\vec{O}], \vec{P}]/x\}, \vec{Q}] \\ \rightsquigarrow^o & v1[\vec{M}, F'\{v01[\vec{N}, O', \vec{P}]/x\}, \vec{Q}] \end{array}$$

In the above reductions, we assumed that  $F$  brings  $x$  to evaluation position after  $f$  steps. If so, then the length of the head is preserved, the cut is moved  $f + 1$  steps forward, and the weight decreases when working on the last visited cut. Otherwise, either the head of  $\pi'$  stops earlier, not visiting the cut anymore, or the surrounding computation eventually brings  $x$  to evaluation position in which case the head of  $\pi'$  catches up, as in the displayed reductions (the length is preserved, the cut is moved forward by many steps, and weight decreases when working on the last visited cut).

Note that linearity of  $\lambda x.F$  is crucial here: if it were to duplicate  $x$  (outside of additive pairs), the head of  $\pi'$  would need to perform the sequence of  $o$  reductions several times.

■ Other right  $\setminus\text{-}l$ :

$$\frac{\frac{\frac{\vdots}{\Delta \rightarrow g} \quad \frac{\frac{\vdots}{\Sigma_2 \rightarrow e} \quad \frac{\vdots}{\Gamma, g, \Sigma_1, f, \Pi \rightarrow h}}{\setminus\text{-}l} \frac{\Gamma, g, \Sigma_1, \Sigma_2, e \setminus f, \Pi \rightarrow h}{\Gamma, \Delta, \Sigma_1, \Sigma_2, e \setminus f, \Pi \rightarrow h}}{\text{cut} \frac{\Delta \rightarrow g \quad \Gamma, \Delta, \Sigma_1, \Sigma_2, e \setminus f, \Pi \rightarrow h}}{\Gamma, \Delta, \Sigma_1, \Sigma_2, e \setminus f, \Pi \rightarrow h}} \mapsto \frac{\frac{\frac{\vdots}{\Sigma_2 \rightarrow e} \quad \frac{\frac{\vdots}{\Delta \rightarrow g} \quad \frac{\vdots}{\Gamma, g, \Sigma_1, f, \Pi \rightarrow h}}{\text{cut} \frac{\Delta \rightarrow g \quad \Gamma, g, \Sigma_1, f, \Pi \rightarrow h}{\Gamma, \Delta, \Sigma_1, f, \Pi \rightarrow h}}}{\setminus\text{-}l} \frac{\Gamma, \Delta, \Sigma_1, \Sigma_2, e \setminus f, \Pi \rightarrow h}{\Gamma, \Delta, \Sigma_1, \Sigma_2, e \setminus f, \Pi \rightarrow h}}$$

$$\begin{array}{ll}
v[\vec{M}, \vec{N}, \vec{O}, \vec{P}, \lambda x.F, \vec{Q}] & v[\vec{M}, \vec{N}, \vec{O}, \vec{P}, \lambda x.F, \vec{Q}] \\
\rightsquigarrow_{cut} v1[\vec{M}, v0[\vec{N}], \vec{O}, \vec{P}, \lambda x.F, \vec{Q}] & \rightsquigarrow v1[\vec{M}, \vec{N}, \vec{O}, F\{v0[\vec{P}]/x\}, \vec{Q}] \\
\rightsquigarrow^n v1[\vec{M}, N', \vec{O}, \vec{P}, \lambda x.F, \vec{Q}] & \rightsquigarrow^r v1[\vec{M}, \vec{N}, \vec{O}, R, \vec{Q}] \\
\rightsquigarrow v11[\vec{M}, N', \vec{O}, F\{v10[\vec{P}]/x\}, \vec{Q}] & \rightsquigarrow_{cut} v11[\vec{M}, v10[\vec{N}], \vec{O}, R, \vec{Q}] \\
\rightsquigarrow^r v11[\vec{M}, N', \vec{O}, R, \vec{Q}] & \rightsquigarrow^n v11[\vec{M}, N', \vec{O}, R, \vec{Q}]
\end{array}$$

The length is preserved, the cut is moved  $r + 1$  steps forward, the weight decreases when working on the last visited cut.

(Last case of a right  $\setminus l$  is similar.)

- Right  $\setminus r$ :

$$\begin{array}{c}
\vdots \\
\vdots \\
\frac{\Delta \rightarrow g}{cut} \quad \frac{e, \Gamma, g, \Sigma \rightarrow f}{\setminus r} \quad \frac{\Gamma, g, \Sigma \rightarrow e \setminus f}{\Gamma, \Delta, \Sigma \rightarrow e \setminus f} \quad \mapsto \quad \frac{\Delta \rightarrow g}{cut} \quad \frac{e, \Gamma, g, \Sigma \rightarrow f}{\setminus r} \quad \frac{e, \Gamma, \Delta, \Sigma \rightarrow f}{\Gamma, \Delta, \Sigma \rightarrow e \setminus f} \\
\vdots \\
\vdots
\end{array}$$

$$\begin{array}{ll}
v[\vec{M}, \vec{N}, \vec{O}] & v[\vec{M}, \vec{N}, \vec{O}] \\
\rightsquigarrow_{cut} v1[\vec{M}, v0[\vec{N}], \vec{O}] & \rightsquigarrow \lambda x.v1[x, \vec{M}, \vec{N}, \vec{O}] \\
\rightsquigarrow^n v1[\vec{M}, N', \vec{O}] & \\
\rightsquigarrow \lambda x.v11[x, \vec{M}, N', \vec{O}] &
\end{array}$$

Like the right  $\cap r$  case, we have two cases:

- either the head of  $\pi$  does not subsequently apply the produced  $\lambda$ -abstraction, in which case the length of the head decreases and the cut is no longer visited;
- or it does, the head of  $\pi'$  catches up at this point in  $1 + n$  steps, the length of the head is preserved, the cut is moved many steps forward, and other the cuts do not move if this was the last visited one.
- Cases for right residuals are symmetrical to those for left residuals.