



HAL
open science

Notes on explicit and inversion formulas for the Chebyshev polynomials of the first two kinds

Feng Qi, Da-Wei Niu, Dongkyu Lim

► **To cite this version:**

Feng Qi, Da-Wei Niu, Dongkyu Lim. Notes on explicit and inversion formulas for the Chebyshev polynomials of the first two kinds: Explicit and inversion formulas for Chebyshev polynomials. *Mathematical Notes*, 2019, 20 (2), pp.1127–1139. 10.18514/MMN.2019.2976 . hal-01705040v2

HAL Id: hal-01705040

<https://hal.science/hal-01705040v2>

Submitted on 19 Jan 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

NOTES ON EXPLICIT AND INVERSION FORMULAS FOR THE CHEBYSHEV POLYNOMIALS OF THE FIRST TWO KINDS

FENG QI, DA-WEI NIU, AND DONGKYU LIM

ABSTRACT. In the paper, starting from the Rodrigues formulas for the Chebyshev polynomials of the first and second kinds, by virtue of the Faà di Bruno formula, with the help of two identities for the Bell polynomials of the second kind, and making use of a new inversion theorem for combinatorial coefficients, the authors derive two nice explicit formulas and their corresponding inversion formulas for the Chebyshev polynomials of the first and second kinds.

1. INTRODUCTION

It is well known [5–7, 29] that the Chebyshev polynomials of the first and second kinds T_n and $U_n(x)$ are very important in mathematical sciences and that, in the study of ordinary differential equations [5, pp. xxxv and 1004], they arise as solutions to the Chebyshev differential equations

$$(1-x^2)y'' - xy' + n^2y = 0 \quad \text{and} \quad (1-x^2)y'' - 3xy' + n(n+2)y = 0$$

for the Chebyshev polynomials of the first and second kinds T_n and U_n respectively.

In [6, Eqs. (4.30) and (4.31)], the Rodrigues formulas for the Chebyshev polynomials of the first and second kinds T_n and U_n read that

$$T_n(x) = (-1)^n \frac{2^n n!}{(2n)!} (1-x^2)^{1/2} \frac{d^n}{dx^n} [(1-x^2)^{n-1/2}] \quad (1.1)$$

and

$$U_n(x) = (-1)^n \frac{2^n (n+1)!}{(2n+1)!} (1-x^2)^{-1/2} \frac{d^n}{dx^n} [(1-x^2)^{n+1/2}]. \quad (1.2)$$

For variants of the Rodrigues formulas for the Chebyshev polynomials of the first and second kinds T_n and U_n , please refer to, for example, [5, pp. 1003–1004], [7, p. 442], [18, Section 4], and [29, pp. 432–433].

In [5, p. 1003], the Rodrigues formulas for $T_n(x)$ and $U_n(x)$ are written in the forms

$$T_n(x) = (-1)^n \frac{\sqrt{1-x^2}}{(2n-1)!} \frac{d^n}{dx^n} [(1-x^2)^{n-1/2}] \quad (1.3)$$

2010 *Mathematics Subject Classification.* Primary 11B38; Secondary 11C08, 26C05, 33C45, 33C47, 33D45.

Key words and phrases. explicit formula; inversion formula; Rodrigues formula; Chebyshev polynomial of the first kind; Chebyshev polynomial of the second kind; Faà di Bruno formula; Bell polynomial of the second kind.

Please cite this article as “Feng Qi, Da-Wei Niu, and Dongkyu Lim, *Notes on explicit and inversion formulas for the Chebyshev polynomials of the first two kinds*, Miskolc Mathematical Notes **20** (2019), no. 2, 1129–1137; available online at <https://doi.org/10.18514/MMN.2019.2976>.”

and

$$U_n(x) = \frac{(-1)^n(n+1)}{\sqrt{1-x^2}(2n+1)!!} \frac{d^n}{dx^n} [(1-x^2)^{n+1/2}]. \quad (1.4)$$

In [5, p. 1004] and [29, pp. 432–433], the Rodrigues formulas for $T_n(x)$ and $U_n(x)$ are formulated as

$$T_n(x) = \frac{(-1)^n\sqrt{\pi}}{2^n\Gamma(n+1/2)} (1-x^2)^{1/2} \frac{d^n}{dx^n} [(1-x^2)^{n-1/2}] \quad (1.5)$$

and

$$U_n(x) = \frac{(-1)^n\sqrt{\pi}(n+1)}{2^{n+1}\Gamma(n+3/2)} (1-x^2)^{-1/2} \frac{d^n}{dx^n} [(1-x^2)^{n+1/2}], \quad (1.6)$$

where $\Gamma(z)$ stands for the classical gamma function which can be defined [?, ?] by

$$\Gamma(z) = \lim_{n \rightarrow \infty} \frac{n!n^z}{\prod_{k=0}^n (z+k)}, \quad z \in \mathbb{C} \setminus \{0, -1, -2, \dots\}$$

or by

$$\Gamma(z) = \int_0^\infty t^{z-1} e^{-t} dt, \quad \Re(z) > 0.$$

In [7, p. 442], the Rodrigues formulas for $T_n(x)$ and $U_n(x)$ are arranged as

$$T_n(x) = \frac{(1-x^2)^{1/2}}{(-2)^n(1/2)_n} \frac{d^n}{dx^n} [(1-x^2)^{n-1/2}] \quad (1.7)$$

and

$$U_n(x) = \frac{(n+1)(1-x^2)^{-1/2}}{(-2)^n(3/2)_n} \frac{d^n}{dx^n} [(1-x^2)^{n+1/2}], \quad (1.8)$$

where $(x)_n$ for $n \geq 0$ and $x \in \mathbb{R}$ denotes the rising factorial which can be defined [22] by

$$(x)_n = \prod_{\ell=0}^{n-1} (x+\ell) = \frac{\Gamma(x+n)}{\Gamma(x)} = \begin{cases} x(x+1)\cdots(x+n-1), & n \geq 1; \\ 1, & n = 0. \end{cases}$$

By virtue of the recurrence relation $\Gamma(x+1) = x\Gamma(x)$, we have

$$\Gamma\left(n + \frac{1}{2}\right) = \prod_{\ell=0}^{n-1} \left(n - \ell - \frac{1}{2}\right) \Gamma\left(\frac{1}{2}\right) = \left(\frac{1}{2}\right)_n \sqrt{\pi} = \frac{(2n-1)!!}{2^n} \sqrt{\pi}$$

and

$$\Gamma\left(n + \frac{3}{2}\right) = \prod_{\ell=0}^n \left(n - \ell + \frac{1}{2}\right) \Gamma\left(\frac{1}{2}\right) = \left(\frac{3}{2}\right)_n \frac{\sqrt{\pi}}{2} = \frac{(2n+1)!!}{2^{n+1}} \sqrt{\pi}.$$

Substituting these into (1.5) and (1.6) respectively leads to (1.3), (1.4), (1.7), and (1.8) which are equivalent to (1.1) and (1.2) respectively.

In [29, pp. 432–433], it was listed that

$$T_n(x) = \frac{n}{2} \sum_{m=0}^{\lfloor n/2 \rfloor} (-1)^m \frac{(n-m-1)!}{m!(n-2m)!} (2x)^{n-2m} \quad (1.9)$$

and

$$U_n(x) = \sum_{m=0}^{\lfloor n/2 \rfloor} (-1)^m \frac{(n-m)!}{m!(n-2m)!} (2x)^{n-2m}, \quad (1.10)$$

where $n \in \mathbb{N}$ and $\lfloor t \rfloor$ denotes the floor function whose value equals the largest integer less than or equal to t .

In this paper, starting from the four formulas (1.1), (1.2), (1.9), and (1.10), by virtue of the Faà di Bruno formula, with the help of two identities for the Bell polynomials of the second kind, and making use of a new inversion theorem [26, Theorem 4.3] for combinatorial coefficients, we will derive the following two nice explicit formulas and their corresponding inversion formulas for the Chebyshev polynomials T_n and U_n .

2. FOUR LEMMAS

For proving our main results, Theorems 1 and 2 below, we need the following four lemmas.

Lemma 1 ([4, pp. 134 and 139]). *For $n \geq k \geq 0$, the Faà di Bruno formula can be described in terms of the Bell polynomials of the second kind $B_{n,k}(x_1, x_2, \dots, x_{n-k+1})$ by*

$$\frac{d^n}{dt^n} f \circ h(t) = \sum_{k=0}^n f^{(k)}(h(t)) B_{n,k}(h'(t), h''(t), \dots, h^{(n-k+1)}(t)). \quad (2.1)$$

Lemma 2 ([4, p. 135]). *For $n \geq k \geq 0$, we have*

$$B_{n,k}(abx_1, ab^2x_2, \dots, ab^{n-k+1}x_{n-k+1}) = a^k b^n B_{n,k}(x_1, x_2, \dots, x_{n-k+1}), \quad (2.2)$$

where a and b are any complex numbers.

Lemma 3 ([13, Theorem 5.1] and [25, Section 3]). *For $n \geq k \geq 0$, the Bell polynomials of the second kind $B_{n,k}(x_1, x_2, \dots, x_{n-k+1})$ satisfy*

$$B_{n,k}(x, 1, 0, \dots, 0) = \frac{1}{2^{n-k}} \frac{n!}{k!} \binom{k}{n-k} x^{2k-n}, \quad (2.3)$$

where $\binom{0}{0} = 1$ and $\binom{p}{q} = 0$ for $q > p \geq 0$.

Lemma 4 ([26, Theorem 4.3]). *For $n \geq k \geq 1$, let s_k and S_k be two sequences independent of n . Then*

$$\frac{s_n}{n!} = \sum_{k=1}^n (-1)^k \binom{k}{n-k} S_k$$

if and only if

$$nS_n = \sum_{k=1}^n \frac{(-1)^k}{(k-1)!} \binom{2n-k-1}{n-1} s_k.$$

3. MAIN RESULTS AND THEIR PROOFS

Now we begin to state and prove our main results, Theorems 1 and 2 below.

Theorem 1. *For $n \geq 0$, the Chebyshev polynomials T_n and U_n can be explicitly computed by*

$$T_n(x) = x^n \sum_{\ell=0}^{\lfloor n/2 \rfloor} \binom{n}{2\ell} \left(1 - \frac{1}{x^2}\right)^\ell \quad (3.1)$$

and

$$U_n(x) = x^n \sum_{\ell=0}^{\lfloor n/2 \rfloor} \binom{n+1}{2\ell+1} \left(1 - \frac{1}{x^2}\right)^\ell. \quad (3.2)$$

Proof. By virtue of the formuals (2.1), (2.2), and (2.3), we have

$$\begin{aligned} \frac{d^n}{dx^n} [(1-x^2)^{n-1/2}] &= \sum_{k=1}^n \frac{d^k u^{n-1/2}}{d u^k} B_{n,k}(-2x, -2, 0, \dots, 0) \\ &= \sum_{k=1}^n \prod_{\ell=0}^{k-1} \left(n - \ell - \frac{1}{2}\right) u^{n-k-1/2} (-2)^k B_{n,k}(x, 1, 0, \dots, 0) \\ &= \sum_{k=1}^n \frac{1}{2^k} \prod_{\ell=0}^{k-1} (2n - 2\ell - 1) (1-x^2)^{n-k-1/2} (-2)^k \frac{1}{2^{n-k}} \frac{n!}{k!} \binom{k}{n-k} x^{2k-n} \\ &= \frac{n!}{(2x)^n} (1-x^2)^{n-1/2} \sum_{k=1}^n (-1)^k \binom{k}{n-k} \frac{(2n-1)!!}{[2(n-k)-1]!!} \frac{2^k}{k!} \left(\frac{x^2}{1-x^2}\right)^k \\ &= \frac{n!(2n-1)!!}{(2x)^n} (1-x^2)^{n-1/2} \sum_{k=1}^n (-1)^k \binom{k}{n-k} \frac{2^k}{k![2(n-k)-1]!!} \left(\frac{x^2}{1-x^2}\right)^k, \end{aligned}$$

where $n \in \mathbb{N}$, $u = u(x) = 1 - x^2$, and the double factorial of negative odd integers $-2n - 1$ is defined by

$$(-2n-1)!! = \frac{(-1)^n}{(2n-1)!!} = (-1)^n \frac{2^n n!}{(2n)!}, \quad n \geq 0.$$

Substituting the above established equality into (1.1) and simplifying lead to

$$T_n(x) = \sum_{k=1}^n \frac{(-1)^{n-k}}{4^{n-k}} \binom{k}{n-k} \binom{n}{k} \frac{[2(n-k)]!!}{[2(n-k)-1]!!} x^{2k-n} (1-x^2)^{n-k}$$

which can be rearranged, by replacing $n - k$ by ℓ , as

$$T_n(x) = x^n \sum_{\ell=0}^{n-1} \frac{(-1)^\ell}{4^\ell} \binom{n}{n-\ell} \binom{n-\ell}{\ell} \frac{(2\ell)!!}{(2\ell-1)!!} \left(\frac{1}{x^2} - 1\right)^\ell.$$

Since

$$\frac{1}{4^\ell} \binom{n}{n-\ell} \binom{n-\ell}{\ell} \frac{(2\ell)!!}{(2\ell-1)!!} = \binom{n}{2\ell},$$

we arrives at the identity (3.1).

Repeating the above process, we can obtain

$$\begin{aligned} \frac{d^n}{dx^n} [(1-x^2)^{n+1/2}] &= \sum_{k=1}^n \frac{d^k u^{n+1/2}}{d u^k} B_{n,k}(-2x, -2, 0, \dots, 0) \\ &= \sum_{k=1}^n \prod_{\ell=0}^{k-1} \left(n - \ell + \frac{1}{2}\right) u^{n-k+1/2} (-2)^k B_{n,k}(x, 1, 0, \dots, 0) \\ &= \sum_{k=1}^n \frac{1}{2^k} \prod_{\ell=0}^{k-1} (2n - 2\ell + 1) (1-x^2)^{n-k+1/2} (-2)^k \frac{1}{2^{n-k}} \frac{n!}{k!} \binom{k}{n-k} x^{2k-n} \end{aligned}$$

$$\begin{aligned}
&= \frac{n!}{(2x)^n} (1-x^2)^{n+1/2} \sum_{k=1}^n (-1)^k \binom{k}{n-k} \frac{(2n+1)!!}{[2(n-k)+1]!!} \frac{2^k}{k!} \left(\frac{x^2}{1-x^2} \right)^k \\
&= \frac{n!(2n+1)!!}{(2x)^n} (1-x^2)^{n+1/2} \sum_{k=1}^n (-1)^k \binom{k}{n-k} \frac{2^k}{k![2(n-k)+1]!!} \left(\frac{x^2}{1-x^2} \right)^k.
\end{aligned}$$

Substituting this into (1.2) and simplifying lead to

$$U_n(x) = \sum_{k=1}^n \frac{(-1)^{n-k}}{2^{2n-2k+1}} \binom{k}{n-k} \binom{n+1}{k} \frac{[2(n-k+1)]!!}{[2(n-k)+1]!!} x^{2k-n} (1-x^2)^{n-k}.$$

Replacing $n-k$ by ℓ reveals that

$$U_n(x) = \sum_{\ell=0}^{n-1} \frac{(-1)^\ell}{2^{2\ell+1}} \binom{n+1}{n-\ell} \binom{n-\ell}{\ell} \frac{[2(\ell+1)]!!}{(2\ell+1)!!} x^{n-2\ell} (1-x^2)^\ell.$$

Due to

$$\frac{1}{2^{2\ell+1}} \binom{n+1}{n-\ell} \binom{n-\ell}{\ell} \frac{[2(\ell+1)]!!}{(2\ell+1)!!} = \binom{n+1}{2\ell+1},$$

we derive (3.2). The proof of Theorem 1 is complete. \square

Theorem 2. For $n \in \mathbb{N}$, we have

$$\sum_{k=1}^n \binom{2n-k-1}{n-1} (2x)^k T_k(x) = \frac{1}{2} (2x)^{2n} \quad (3.3)$$

and

$$\sum_{k=1}^n k \binom{2n-k-1}{n-1} (2x)^k U_k(x) = n(2x)^{2n}. \quad (3.4)$$

Proof. We notice that the formulas (1.9) and (1.10) can be rearranged as

$$T_n(x) = \frac{n}{2} \sum_{m=0}^{\lfloor n/2 \rfloor} (-1)^m \binom{n-m}{m} \frac{(2x)^{n-2m}}{n-m} \quad (3.5)$$

and

$$U_n(x) = \sum_{m=0}^{\lfloor n/2 \rfloor} (-1)^m \binom{n-m}{m} (2x)^{n-2m}. \quad (3.6)$$

The inversion theorem in Lemma 4 can be restated as

$$(-1)^n \frac{s_n}{n!} = \sum_{\ell=0}^{n-1} (-1)^\ell \binom{n-\ell}{\ell} S_{n-\ell} = \sum_{\ell=0}^{\lfloor n/2 \rfloor} (-1)^\ell \binom{n-\ell}{\ell} S_{n-\ell}$$

if and only if

$$nS_n = \sum_{\ell=1}^n \frac{(-1)^\ell}{(\ell-1)!} \binom{2n-\ell-1}{n-1} s_\ell.$$

The formulas (3.5) and (3.6) can be rearranged as

$$\frac{2}{n} (2x)^n T_n(x) = \sum_{\ell=0}^{\lfloor n/2 \rfloor} (-1)^\ell \binom{n-\ell}{\ell} \frac{(2x)^{2(n-\ell)}}{n-\ell}$$

and

$$(2x)^n U_n(x) = \sum_{\ell=0}^{\lfloor n/2 \rfloor} (-1)^\ell \binom{n-\ell}{\ell} (2x)^{2(n-\ell)}.$$

Consequently, we obtain

$$n \frac{(2x)^{2n}}{n} = \sum_{k=1}^n \frac{(-1)^k}{(k-1)!} \binom{2n-k-1}{n-1} (-1)^k 2^k (k-1)! (2x)^k T_k(x)$$

and

$$n(2x)^{2n} = \sum_{k=1}^n \frac{(-1)^k}{(k-1)!} \binom{2n-k-1}{n-1} (-1)^k k! (2x)^k U_k(x)$$

which can be simplified as (3.3) and (3.4). The proof of Theorem 2 is complete. \square

4. REMARKS

In this section, we will list several remarks to explain more about the formula (2.3), Lemma 4, our main results, and other things.

Remark 1. To the best of our knowledge, the nice formula (2.3) was first concluded in [13] and has been extensively applied in the papers [8–16, 18, 19, 21, 23, 25, 27, 28] and closely related references therein. The formula (2.3) has been generalized in the papers [16, 19, 20] and closely related references therein.

Remark 2. To the best of our knowledge, Lemma 4 is a new inversion theorem and has been applied in the paper [8, 17, 18, 24].

Remark 3. Because both the formula (2.3) and Lemma 4 are new, our main results stated in Theorems 1 and 2, or at least their proofs, are also new.

Remark 4. The Chebyshev polynomials are classical, but their study is still very active. As examples, we recommend three newly-published papers [1–3] to readers. Considering the length of this paper, we would not like to detail main results in these three papers and the closely-related references therein.

Remark 5. This paper is a slightly revised version of the preprint [18].

Acknowledgements. The third author was supported by the National Research Foundation of Korea (Grant No. 2018R1D1A1B07041846).

The authors appreciate anonymous referees for their valuable comments on the original version of this paper.

REFERENCES

- [1] C. Cesarano, *Integral representations and new generating functions of Chebyshev polynomials*, Hacet. J. Math. Stat. **44** (2015), no. 3, 535–546; available online at <https://doi.org/10.15672/HJMS.20154610029>.
- [2] C. Cesarano, *Multi-dimensional Chebyshev polynomials: a non-conventional approach*, Commun. Appl. Ind. Math. **10** (2019), no. 1, 1–19; available online at <https://doi.org/10.1515/caim-2019-0008>.
- [3] C. Cesarano and C. Fornaro, Claudio, *A note on two-variable Chebyshev polynomials*, Georgian Math. J. **24** (2017), no. 3, 339–349; available online at <https://doi.org/10.1515/gmj-2016-0034>.

- [4] L. Comtet, *Advanced Combinatorics: The Art of Finite and Infinite Expansions*, Revised and Enlarged Edition, D. Reidel Publishing Co., 1974; available online at <https://doi.org/10.1007/978-94-010-2196-8>.
- [5] I. S. Gradshteyn and I. M. Ryzhik, *Table of Integrals, Series, and Products*, Translated from the Russian, Translation edited and with a preface by Daniel Zwillinger and Victor Moll, Eighth edition, Revised from the seventh edition, Elsevier/Academic Press, Amsterdam, 2015; available online at <https://doi.org/10.1016/B978-0-12-384933-5.00013-8>.
- [6] J. C. Mason and D. C. Handscomb, *Chebyshev Polynomials*, Chapman & Hall/CRC, Boca Raton, FL, 2003.
- [7] F. W. J. Olver, D. W. Lozier, R. F. Boisvert, and C. W. Clark (eds.), *NIST Handbook of Mathematical Functions*, Cambridge University Press, New York, 2010; available online at <http://dlmf.nist.gov/>.
- [8] F. Qi, *A simple form for coefficients in a family of ordinary differential equations related to the generating function of the Legendre polynomials*, Adv. Appl. Math. Sci. **17** (2018), no. 11, 693–700.
- [9] F. Qi, *Integral representations for multivariate logarithmic polynomials*, J. Comput. Appl. Math. **336** (2018), 54–62; available online at <https://doi.org/10.1016/j.cam.2017.11.047>.
- [10] F. Qi and V. Čerňanová, *Some discussions on a kind of improper integrals*, Int. J. Anal. Appl. **11** (2016), no. 2, 101–109.
- [11] F. Qi, V. Čerňanová, X.-T. Shi, and B.-N. Guo, *Some properties of central Delannoy numbers*, J. Comput. Appl. Math. **328** (2018), 101–115; available online at <https://doi.org/10.1016/j.cam.2017.07.013>.
- [12] F. Qi and B.-N. Guo, *Explicit and recursive formulas, integral representations, and properties of the large Schröder numbers*, Kragujevac J. Math. **41** (2017), no. 1, 121–141; available online at <https://doi.org/10.5937/KgJMath1701121F>.
- [13] F. Qi and B.-N. Guo, *Explicit formulas for special values of the Bell polynomials of the second kind and for the Euler numbers and polynomials*, Mediterr. J. Math. **14** (2017), no. 3, Article 140, 14 pages; available online at <https://doi.org/10.1007/s00009-017-0939-1>.
- [14] F. Qi and B.-N. Guo, *Several explicit and recursive formulas for the generalized Motzkin numbers*, Preprints **2017**, 2017030200, 11 pages; available online at <https://doi.org/10.20944/preprints201703.0200.v1>.
- [15] F. Qi and B.-N. Guo, *Some properties of the Hermite polynomials*, AIMS Mathematics **5** (2020), in press; available online at <https://doi.org/10.20944/preprints201611.0145.v1>.
- [16] F. Qi and B.-N. Guo, *Viewing some ordinary differential equations from the angle of derivative polynomials*, Iran. J. Math. Sci. Inform. **15** (2020), no. 2, in press.
- [17] F. Qi, D.-W. Niu, and B.-N. Guo, *Some identities for a sequence of unnamed polynomials connected with the Bell polynomials*, Rev. R. Acad. Cienc. Exactas Fis. Nat. Ser. A Math. RACSAM **113** (2019), no. 2, 557–567; available online at <https://doi.org/10.1007/s13398-018-0494-z>.
- [18] F. Qi, D.-W. Niu, and D. Lim, *Notes on the Rodrigues formulas for two kinds of the Chebyshev polynomials*, HAL preprint (2018), available online at <https://hal.archives-ouvertes.fr/hal-01705040>.
- [19] F. Qi, D.-W. Niu, D. Lim, and B.-N. Guo, *Explicit formulas and identities on Bell polynomials and falling factorials*, ResearchGate Preprint (2018), available online at <https://doi.org/10.13140/RG.2.2.34679.52640>.
- [20] F. Qi, D.-W. Niu, D. Lim, and Y.-H. Yao, *Special values of the Bell polynomials of the second kind for some sequences and functions*, HAL preprint (2018), available online at <https://hal.archives-ouvertes.fr/hal-01766566>.
- [21] F. Qi, X.-T. Shi, and B.-N. Guo, *Two explicit formulas of the Schröder numbers*, Integers **16** (2016), Paper No. A23, 15 pages.
- [22] F. Qi, X.-T. Shi, and F.-F. Liu, *Several identities involving the falling and rising factorials and the Cauchy, Lah, and Stirling numbers*, Acta Univ. Sapientiae Math. **8** (2016), no. 2, 282–297; available online at <https://doi.org/10.1515/ausm-2016-0019>.
- [23] F. Qi, X.-T. Shi, F.-F. Liu, and D. V. Kruchinin, *Several formulas for special values of the Bell polynomials of the second kind and applications*, J. Appl. Anal. Comput. **7** (2017), no. 3, 857–871; available online at <https://doi.org/10.11948/2017054>.

- [24] F. Qi and Y.-H. Yao, *Simplifying coefficients in differential equations for generating function of Catalan numbers*, J. Taibah Univ. Sci. **13** (2019), no. 1, 947–950; available online at <https://doi.org/10.1080/16583655.2019.1663782>.
- [25] F. Qi and M.-M. Zheng, *Explicit expressions for a family of the Bell polynomials and applications*, Appl. Math. Comput. **258** (2015), 597–607; available online at <https://doi.org/10.1016/j.amc.2015.02.027>.
- [26] F. Qi, Q. Zou, and B.-N. Guo, *The inverse of a triangular matrix and several identities of the Catalan numbers*, Appl. Anal. Discrete Math. **13** (2019), no. 2, 518–541; available online at <https://doi.org/10.2298/AADM190118018Q>.
- [27] C.-F. Wei and F. Qi, *Several closed expressions for the Euler numbers*, J. Inequal. Appl. **2015**, 2015:219, 8 pages; available online at <https://doi.org/10.1186/s13660-015-0738-9>.
- [28] J.-L. Zhao and F. Qi, *Two explicit formulas for the generalized Motzkin numbers*, J. Inequal. Appl. **2017**, 2017:44, 8 pages; available online at <https://doi.org/10.1186/s13660-017-1313-3>.
- [29] D. Zwillinger, *CRC Standard Mathematical Tables and Formulae*, Thirty-first edition, CRC Press, Boca Raton, FL, 2003.

Authors' addresses

Feng Qi

INSTITUTE OF MATHEMATICS, HENAN POLYTECHNIC UNIVERSITY, JIAOZUO 454010, HENAN, CHINA;
COLLEGE OF MATHEMATICS, INNER MONGOLIA UNIVERSITY FOR NATIONALITIES, TONGLIAO 028043,
INNER MONGOLIA, CHINA; SCHOOL OF MATHEMATICAL SCIENCES, TIANJIN POLYTECHNIC UNIVERSITY,
TIANJIN 300387, CHINA

Email address: qifeng618@gmail.com, qifeng618@hotmail.com

URL: <https://qifeng618.wordpress.com>

Da-Wei Niu

DEPARTMENT OF SCIENCE, HENAN UNIVERSITY OF ANIMAL HUSBANDRY AND ECONOMY, ZHENGZHOU
450046, HENAN, CHINA

Email address: nnddww@gmail.com

URL: <https://orcid.org/0000-0003-4033-7911>

Dongkyu Lim

DEPARTMENT OF MATHEMATICS EDUCATION, ANDONG NATIONAL UNIVERSITY, ANDONG 36729,
REPUBLIC OF KOREA

Email address: dgrim84@gmail.com, dklim@andong.ac.kr

URL: <https://orcid.org/0000-0002-0928-8480>