

# Simulating the effect of public health interventions using dated virus sequences and geographical data

Samuel Alizon

#### ▶ To cite this version:

Samuel Alizon. Simulating the effect of public health interventions using dated virus sequences and geographical data. Peer Community In Evolutionary Biology, 2018, pp.100046. 10.24072/pci.evolbiol.100046 . hal-01705816

### HAL Id: hal-01705816 https://hal.science/hal-01705816

Submitted on 9 Feb 2018

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NoDerivatives 4.0 International License

## Peer Community In Evolutionary Biology

## Simulating the effect of public health interventions using dated virus sequences and geographical data

## **Open Access**

Copyright: This work is licensed under the Creative Commons Attribution-NoDerivatives 4.0 International License. To view a copy of this license, visit http://creativecommons.org/licen ses/by-nd/4.0/

#### Samuel Alizon

----

MIVEGEC, CNRS -- Montpellier, France Correspondence to Samuel Alizon (samuel.alizon@cnrs.fr) doi: 10.24072/pci.evolbiol.100046

Published: 09th Feb. 2018

Cite as: Alizon S. 2018. Simulating the effect of public health interventions using dated virus sequences and geographical data. Peer Community in Evolutionary Biology, 100046 doi: 10.24072/pci.evolbiol.100046

#### A recommendation – based on reviews by Christian Althaus and Chris Wymant – of

Dellicour S, Baele G, Dudas G, Faria NR, Pybus OG, Suchard MA, Rambaud A and Lemey P. 2018. Phylodynamic assessment of intervention strategies for the West African Ebola virus outbreak. bioRxiv, 163691, ver. 3 peer-reviewed by Peer Community In Evolutionary Biology. doi: https://doi.org/10.1101/163691

Perhaps because of its deadliness, the 2013-2016 Ebola Virus (EBOV) epidemics in West-Africa has led to unprecedented publication and sharing of full virus genome sequences. This was both rapid (90 full genomes were shared within weeks [1]) and important (more than 1500 full genomes have been released overall [2]). Furthermore, the availability of the metadata (especially GPS location) has led to depth analyses of the geographical spread of the epidemics [3].

In their work, Dellicour et al. [4] pursue earlier phylogeographical investigations in an original and yet simple approach to address questions of key public health importance. The originality of the approach is dual. First, from a technical standpoint, they capture the spread of infectious diseases in a continuous framework using a novel model that allows for rare long-distance dispersal events. Second, in a more classical discrete meta-population framework, they simulate the effect of public health interventions by pruning the phylogenetic tree and assessing how this affects key parameters. For instance, to simulate the effect of closing borders they remove subsets of the phylogeny that involved dispersal



between countries and to simulate the effect of protecting a region by quarantine they remove all the leaves (i.e. the infections sampled) from this region. This phylogeny pruning is both original and simple. It is however limited because it currently assumes that policies are 100% effective and earlier modelling work on human influenza showed that long distance travel bans had to be implemented with >99% efficiency in order to slow epidemic growth from a time scale of days to weeks [5].

From a biological standpoint, Dellicour et al. [4] corroborate earlier findings that highly populated locations (>1,000,000 inhabitants) were crucial in explaining the magnitude of the epidemics but also show the importance of the transmission between the three capital cities. They also show that rare long-distance dispersing events of the virus are not key to explaining the magnitude of the epidemics (even though they assume 100% efficiency of suppressing long-distance event). Finally, thanks to their continuous model they estimate the speed of spread of the epidemics and are able to detect the effect of border closing on this speed.

Overall, this study [4], which involves state-of-the-art Bayesian inference methods of infection phylogenies using MCMC, stands out because of its effort to simulate public health interventions. It stands as an encouragement for the development of intervention models with increased realism and for even faster and larger virus sequence data sharing.

#### References

[1] Gire et al. (2014) Genomic surveillance elucidates Ebola virus origin and transmission during the 2014 outbreak. Science 345: 1369–1372. doi: https://doi.org/10.1126/science.1259657

[2] Holmes EC, Dudas G, Rambaut A and Andersen KG (2016) The evolution of Ebola virus: insights from the 2013-2016 epidemic. Nature 538: 193–200. doi: https://doi.org/10.1038/nature19790

[3] Dudas et al. (2017) Virus genomes reveal factors that spread and sustained the Ebola epidemic. Nature 544: 309–315 (2017). doi: https://doi.org/10.1038/nature22040

[4] Dellicour S, Baele G, Dudas G, Faria NR, Pybus OG, Suchard MA, Rambaud A and Lemey P (2018) Phylodynamic assessment of intervention strategies for the West African Ebola virus outbreak. bioRxiv, 163691, ver. 3 peer-reviewed by Peer Community In Evolutionary Biology. doi: https://doi.org/10.1101/163691

[5] Hollingsworth TD, Ferguson NM and Anderson RM (2006) Will travel restrictions control the international spread of pandemic influenza? Nature Medicine 12, 497–499. doi: https://doi.org/10.1038/nm0506-497

### Appendix

Reviews by Christian Althaus and Chris Wymant: https://doi.org/10.24072/pci.evolbiol.100046