



HAL
open science

Rate of convergence for polymers in a weak disorder

Francis Comets, Quansheng Liu

► **To cite this version:**

Francis Comets, Quansheng Liu. Rate of convergence for polymers in a weak disorder. *Journal of Mathematical Analysis and Applications*, 2017, 455 (1), pp.312 - 335. 10.1016/j.jmaa.2017.05.043 . hal-01705859

HAL Id: hal-01705859

<https://hal.science/hal-01705859>

Submitted on 9 Feb 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Rate of convergence for polymers in a weak disorder

Francis Comets¹

Quansheng Liu²

24 mai 2017

¹Université Paris Diderot – Paris 7, Mathématiques, case 7012, F-75205 Paris Cedex 13, France
e-mail : comets@math.univ-paris-diderot.fr

² Corresponding author. Univ. Bretagne-Sud, UMR 6205, LMBA, F-56000 Vannes, France
e-mail : quansheng.liu@univ-ubs.fr

Résumé

We consider directed polymers in random environment on the lattice \mathbb{Z}^d at small inverse temperature and dimension $d \geq 3$. Then, the normalized partition function W_n is a regular martingale with limit W . We prove that $n^{(d-2)/4}(W_n - W)/W_n$ converges in distribution to a Gaussian law. Both the polynomial rate of convergence and the scaling with the martingale W_n are different from those for polymers on trees.

Keywords : Directed polymers, random environment, weak disorder, rate of convergence, regular martingale, central limit theorem for martingales, stable and mixing convergence.

AMS 2010 subject classifications : Primary 60K37. Secondary 60F05, 60G45, 60J80, 82D60.

1 Polymer models and statement of the main result

1.1 Motivation

We consider directed polymers in random environment, given by a simple random walk on the d -dimensional lattice in a space-time random potential. In a seminal paper, Derrida and Spohn [12] perform a detailed analysis of polymers on the Cayley tree, or equivalently, the branching random walk with a fixed branching number. Later the same model has been taken up as an approximation and a toy model with explicit computations : in the physics literature, we mention the pleasant, recent and documented survey [15], and also [11] for the statistics of extremes on the hierarchical tree at zero temperature ; on the mathematical side, the authors of [1] study the near-critical scaling window on the tree, the analogue of the intermediate disorder regime where the rescaled lattice model on line converges to the KPZ continuum random polymer [2, 7]. Not only a source of inspiration and guidance, this model, as well as related random cascades, were also found to provide quantitative bounds on polymer models on the lattice in [9, 26, 27].

In spite of these similarities, the two models behave quite differently in many aspects. In the strong disorder phase, the free energy of the branching process is linear in the inverse temperature β though it is strictly convex for the polymer on the lattice, see Theorem 1.5

in [8] in the case of a Bernoulli environment. Also, the fluctuations are expected to be of a completely different nature in the two models. In this paper, we consider the weak disorder regime, and we show that the martingale convergence takes place at a polynomial rate, whereas it is exponential in the corresponding supercritical Galton-Watson process [16, 17].

More precisely, it is shown in [16, 17] that, for a Galton-Watson process (Z_n) with $Z_0 = 1$, $m = EZ_1 > 1$ and $EZ_1^2 < \infty$, the renormalized population size $W_n = Z_n/m^n$ is a regular martingale with limit W such that

$$m^{n/2}(W - W_n) \rightarrow aW^{1/2}G \quad \text{in distribution} \quad (1)$$

and

$$m^{n/2} \frac{(W - W_n)}{W_n^{1/2}} \rightarrow aG \quad \text{in distribution,} \quad (2)$$

where $a^2 = \frac{\text{Var}Z_1}{m^2 - m}$, G is a Gaussian $\mathcal{N}(0, 1)$ distributed random variable independent of W . Similarly, for branching random walks, the convergence of the Biggins martingale to its limit is exponentially fast [20, 21] in the regular case. Recently the same question was studied for a branching process in a random environment [19, 31], leading to similar conclusions.

In this paper, we consider random polymers on the lattice in a time-space dependent random medium, deep inside the weak disorder regime. Similar to the supercritical case of a branching process, weak disorder can be defined as the regime where the natural martingale is regular [5, 22], or where the polymer is diffusive [10]. It holds in space dimension $d \geq 3$ [23] and at a temperature larger than some critical value which can be estimated by second moment and entropy considerations [4, 6, 18]. In Theorem 1.1 below, we prove that, at large temperature, the *speed of convergence is polynomial* but not exponential, and the limit scales with W or W_n instead of their square root as in (1) and (2). Precisely, we show a central limit theorem for the difference between the martingale and its limit : the ratio of the difference divided by $n^{-(d-2)/4}$ times the martingale is asymptotically normal.

In view of (1) and (2), this limit behavior has two remarkable and unexpected features. The slowdown in the rate of convergence (compared to the branching case) is due to space correlations coming from further intersections between paths on the lattice but not on the tree. Also the unusual linear scaling in the martingale can be understood as coming from fluctuations, and quadratic variations scale like the square of the martingale.

The result helps us for a deeper comprehension of the polymers model, and opens a way for further limit theorems about it.

1.2 Notations

- *The random walk* : $(\{S_n\}_{n \geq 0}, P_x)$ is a nearest neighbor, symmetric simple random walk on the d -dimensional integer lattice \mathbb{Z}^d starting from x , $d \geq 3$. We let $P = P_0$ and we denote by $P[f] = \int f dP$ the expectation of f with respect to P .
- *The random environment* : $\eta = \{\eta(n, x) : n \in \mathbb{N}, x \in \mathbb{Z}^d\}$ is an independent and identically distributed (i.i.d.) sequence of real random variables (r.v.'s), non-constant, such that,

$$\lambda(\beta) := \ln \mathbb{E}[\exp(\beta\eta(0, 0))] < \infty \quad \text{for all } \beta \in \mathbb{R},$$

where we denote by \mathbb{E} the expectation over the environment. The corresponding probability measure will be denoted by \mathbb{P} .

- The *partition function* at inverse temperature $\beta \in \mathbb{R}$

$$W_n := P[\exp(\beta H_n(S) - n\lambda(\beta))] \quad \text{with} \quad H_n(S) = \sum_{0 \leq t \leq n-1} \eta(t, S_t), \quad n \geq 1, \quad (3)$$

is the normalization which makes the Gibbs measure $W_n^{-1} \exp\{\beta H_n(S) - n\lambda(\beta)\} dP$ a probability measure on the path space for a fixed realization of the environment. Note that we use a slightly different definition than usual, including $t = 0$ but not $t = n$ in the Hamiltonian H_n . This makes no fundamental difference in the results (see Remark 1.2 below), but it yields simpler formulas here : for two independent simple random walks $S = (S_t)$ and $\tilde{S} = (\tilde{S}_t)$, we have $\text{Cov}_{\mathbb{P}}(H_n(S), H_n(\tilde{S})) = \text{Var}(\eta(0, 0)) N_n$, where $\text{Cov}_{\mathbb{P}}$ denotes the covariance with respect to \mathbb{P} , N_n is the number of intersections of the paths S, \tilde{S} up to time n :

$$N_n = N_n(S, \tilde{S}) := \sum_{t=0}^{n-1} \mathbf{1}_{S_t = \tilde{S}_t}, \quad (4)$$

whose limit

$$N_\infty = N_\infty(S, \tilde{S}) := \sum_{t=0}^{\infty} \mathbf{1}_{S_t = \tilde{S}_t} \quad (5)$$

has expectation given by the standard Green function (20). The sequence (W_n) depends on the environment, and it is a positive martingale with respect to the filtration

$$\mathcal{G}_n = \sigma\{\eta(t, x); t \leq n-1, x \in \mathbb{Z}^d\}, \quad n \geq 1. \quad (6)$$

It is well known [5, 22] that

$$W = \lim_{n \rightarrow \infty} W_n \quad \text{exists a.s., with} \quad \mathbb{P}(W > 0) = 0 \text{ or } 1.$$

Moreover, the convergence holds in L^2 for β in a neighborhood of 0, defined by

$$\mathbf{(L2)} \quad \lambda_2 := \lambda(2\beta) - 2\lambda(\beta) < \ln(1/\pi_d), \quad (7)$$

where π_d is the return probability of the simple random walk,

$$\pi_d := P\{S_n = 0 \text{ for some } n \geq 1\} \in (0, 1) \quad (8)$$

by transience since $d \geq 3$. Now, we give a short account of the main steps of the computation of [5], which is useful for the sequel. We can express W_n^2 as a sum $W_n^2 = P^{\otimes 2} \left[e^{\beta[H_n(S) + H_n(\tilde{S})] - 2n\lambda(\beta)} \right]$ over independent paths (so-called replicas), and we compute, using Fubini's theorem and independence,

$$\begin{aligned} \mathbb{E}[W_n^2] &= P^{\otimes 2} \left[\prod_{t=0}^{n-1} \mathbb{E} e^{\beta[\eta(t, S_t) + \eta(t, \tilde{S}_t)] - 2\lambda(\beta)} \right] \\ &= P^{\otimes 2} \left[\prod_{t=0}^{n-1} \left(e^{\lambda(2\beta) - 2\lambda(\beta)} \mathbf{1}_{S_t = \tilde{S}_t} + \mathbf{1}_{S_t \neq \tilde{S}_t} \right) \right] \\ &= P^{\otimes 2} \left[\prod_{t=0}^{n-1} e^{\lambda_2 \mathbf{1}_{S_t = \tilde{S}_t}} \right] \\ &= P^{\otimes 2} \left[e^{\lambda_2 N_n} \right], \end{aligned} \quad (9)$$

with N_n as in (4). As $n \rightarrow \infty$, $N_n \nearrow N_\infty = \sum_{t=0}^{\infty} \mathbf{1}_{S_t = \tilde{S}_t}$. Since the process $(S_t - \tilde{S}_t)$ under $P^{\otimes 2}$ has the same law as (S_{2t}) under P and $S_{2t+1} \neq 0$ P -a.s., N_n has the same law as the number of visits to 0 of a simple random walk in time $2n$, and N_∞ is geometrically distributed with "failure" probability π_d :

$$P^{\otimes 2}(N_\infty = k) = \pi_d^{k-1}(1 - \pi_d) \quad \text{for } k \geq 1. \quad (10)$$

Therefore, by the monotone convergence theorem, as $n \rightarrow \infty$,

$$\mathbb{E}[W_n^2] \nearrow P^{\otimes 2} [e^{\lambda_2 N_\infty}] = \begin{cases} \frac{(1 - \pi_d)e^{\lambda_2}}{1 - \pi_d e^{\lambda_2}} & \text{if } \pi_d e^{\lambda_2} < 1, \\ +\infty & \text{if } \pi_d e^{\lambda_2} \geq 1. \end{cases}$$

In this paper, we always assume (7), so that the martingale (W_n) is bounded in L^2 . By Doob's convergence theorem, $W_n \rightarrow W$ in L^2 and then $W > 0$. In particular,

$$\mathbb{E}W^2 = \frac{(1 - \pi_d)e^{\lambda_2}}{1 - \pi_d e^{\lambda_2}} \quad \text{and} \quad \text{Var}(W) = \frac{e^{\lambda_2} - 1}{1 - \pi_d e^{\lambda_2}}. \quad (11)$$

1.3 A Gaussian limit and the rate of convergence

Before coming to our main result, we recall two convergence modes. Let (Y_n) be a sequence of real random variables defined on a common probability space (Ω, \mathcal{F}, P) , converging in distribution to a limit Y .

- This convergence is called *stable* if for all $B \in \mathcal{F}$ with $P(B) > 0$, the conditional law of Y_n given B converges to some probability distribution depending on B .
- This convergence is called *mixing* if it is stable and the limit of conditional laws does not depend on B – and therefore is the law of Y –.

The stable convergence allows to add extra variables : for any fixed r.v. Z on (Ω, \mathcal{F}, P) , the couple (Y_n, Z) converges in law to some coupling of Y and Z on an extended space. The mixing convergence means that Y_n is asymptotically independent of all event $A \in \mathcal{F}$. These convergences were introduced by Rényi [28]; we refer to [3] for a nice presentation with the main consequences, and to [14] pp. 56-57 for an extended account on the connections with martingale central limit theorem.

Theorem 1.1 *For $d \geq 3$, there exists some $\beta_0 > 0$ such that, for $|\beta| < \beta_0$,*

$$n^{\frac{d-2}{4}}(W - W_n) \rightarrow \sigma_1 W G \quad \text{in distribution} \quad (12)$$

and

$$n^{\frac{d-2}{4}} \frac{(W - W_n)}{W_n} \rightarrow \sigma_1 G \quad \text{in distribution}, \quad (13)$$

where σ_1 is from (30), G is a Gaussian r.v. with law $N(0, 1)$, which is independent of W . Moreover, the convergence in (12) is stable, and the convergence in (13) is mixing.

The theorem calls for some comments. The value of β_0 is defined by the conditions in Lemma 4.1, Lemma 2.3 (b) and Lemma 3.4. The result is quite different from (1)–(2). The speed of convergence of W_n to its limit is $n^{-\frac{d-2}{4}}$. The mixing convergence in (13) shows that the random variable G_n defined in the left hand side of (13) is asymptotically independent of each event A of the environmental probability space, in the sense that for all real y , we have $\lim_{n \rightarrow \infty} P(\{G_n \leq y\} \cap A) = P(G \leq y)P(A)$. Our approach relies on a central limit theorem for martingales.

Remark 1.2 (Usual Hamiltonian) For the standard Hamiltonian collecting the environment at times $1, 2, \dots, n$,

$$\bar{W}_n := P \left[\exp \left(\beta \sum_{1 \leq t \leq n} \eta(t, S_t) - n\lambda(\beta) \right) \right] = W_{n+1} \exp\{-\beta\eta(0, 0) + \lambda(\beta)\},$$

it is straightforward to see that, for $d \geq 3$ and $|\beta| < \beta_0$, $\bar{W}_n \rightarrow \bar{W} := W \exp\{-\beta\eta(0, 0) + \lambda(\beta)\}$, that

$$n^{\frac{d-2}{4}} (\bar{W} - \bar{W}_n) \rightarrow \sigma_1 \bar{W} G \quad \text{in distribution}$$

and

$$n^{\frac{d-2}{4}} \frac{(\bar{W} - \bar{W}_n)}{\bar{W}_n} \rightarrow \sigma_1 G \quad \text{in distribution,}$$

with σ_1 and G as above and G independent from \bar{W} .

Organization of the paper : In Section 2 we start with algebraic computations of covariances and use Green function estimates to derive asymptotics. Of independent interest, a specific form of central limit theorem for infinite martingale arrays is given in Lemma 3.1 of Section 3, and used to prove Theorem 1.1. The proofs of intermediate lemmas are postponed to Section 4. A key step consists in controlling the sum of conditional variances in the quadratic norm, so we implement the necessary algebra for a system of 4 replicas at the beginning of the section.

2 The correlation structure

It is useful to introduce

$$W_n(x) = P \left[e^{\beta H_n(S) - n\lambda(\beta)} \mathbf{1}_{S_n=x} \right], \quad (14)$$

and to observe that, for $m \geq 0$ including $m = \infty$ with the convention $W_\infty = W$,

$$W_{n+m} = \sum_x W_n(x) W_m \circ \theta_{n,x}, \quad (15)$$

by Markov property. We view W_n as a function of η , we denote by $\theta_{n,x}$ the shift operator on the environment, $\theta_{n,x}\eta : (t, y) \mapsto \eta(n+t, x+y)$. We use $\theta_x = \theta_{0,x}$ as a short notation. Taking $m = \infty$ we see that

$$W - W_n = \sum_x W_n(x) (W \circ \theta_{n,x} - 1). \quad (16)$$

For nearest-neighbor paths S, \tilde{S} define τ_n the time delay of first intersection after time $n \geq 0$,

$$\tau_n(S, \tilde{S}) = \inf\{k \geq 0 : S_{n+k} = \tilde{S}_{n+k}\},$$

with the convention that $\inf \emptyset = +\infty$. We write $\tau = \tau_0$, and we note that $\pi_d = P_{0,0}^{\otimes 2}(\tau_1 < \infty)$.

2.1 Covariance of the martingale limit and rate of convergence in L^2

Recall that $\text{Var}(W) = \frac{e^{\lambda_2} - 1}{1 - P_{0,0}^{\otimes 2}(\tau_1 < \infty)e^{\lambda_2}}$, cf. (11).

Proposition 2.1 *Assume $\lambda_2 < \frac{2}{d} \ln \frac{1}{\pi_d}$. Then,*

$$\text{Cov}(W, W \circ \theta_x) = \text{Var}(W) \times P_{0,x}^{\otimes 2}(\tau < \infty). \quad (17)$$

Moreover,

$$\|W - W_n\|_2^2 = \text{Var}(W) \times P_{0,0}^{\otimes 2}(e^{\lambda_2 N_n} \mathbf{1}_{\tau_n < \infty}), \quad (18)$$

and, as $n \rightarrow \infty$,

$$\|W - W_n\|_2^2 \sim \sigma^2 \times n^{-(d-2)/2}, \quad (19)$$

with the constant σ^2 from (28).

Proof. We first compute the covariance of W and $W \circ \theta_x$. Denote by \mathcal{F}_n the σ -field generated by S_i, \tilde{S}_i for $0 \leq i \leq n$. By convergence in L^2 , we obtain as in (9),

$$\begin{aligned} \text{Cov}(W, W \circ \theta_x) &= \lim_{m \rightarrow \infty} \mathbb{E}[(W_m - 1)(W_m \circ \theta_x - 1)] \\ &= \lim_{m \rightarrow \infty} P_{0,x}^{\otimes 2} \mathbb{E}[(e^{\beta H_m(S) - m\lambda(\beta)} - 1)(e^{\beta H_m(\tilde{S}) - m\lambda(\beta)} - 1)] \quad (\text{Fubini}) \\ &= \lim_{m \rightarrow \infty} P_{0,x}^{\otimes 2} \mathbb{E}[e^{\beta H_m(S) + \beta H_m(\tilde{S}) - 2m\lambda(\beta)} - 1] \\ &= \lim_{m \rightarrow \infty} P_{0,x}^{\otimes 2} [e^{\lambda_2 N_m}] - 1 \\ &= P_{0,x}^{\otimes 2} [(e^{\lambda_2 N_\infty} - 1)] \\ &= P_{0,x}^{\otimes 2} [\mathbf{1}_{\tau < \infty} P_{0,x}^{\otimes 2} [(e^{\lambda_2 N_\infty} - 1) | \mathcal{F}_\tau]] \\ &= P_{0,x}^{\otimes 2}(\tau < \infty) P_{0,0}^{\otimes 2} [(e^{\lambda_2 N_\infty} - 1)] \quad (\text{strong Markov property}) \end{aligned}$$

which is the first claim since $\text{Var}(W) = P_{0,0}^{\otimes 2} [e^{\lambda_2 N_\infty}] - 1$.

We next calculate the L^2 norm of $W - W_n$. By (16),

$$\begin{aligned} \|W - W_n\|_2^2 &= \mathbb{E} \left(\sum_x W_n(x) (W \circ \theta_{n,x} - 1) \right)^2 \\ &= \mathbb{E} \left(\sum_{x,y} W_n(x) W_n(y) (W \circ \theta_{n,x} - 1) (W \circ \theta_{n,y} - 1) \right) \\ &= \sum_{x,y} \mathbb{E}[W_n(x) W_n(y)] \mathbb{E}[(W \circ \theta_{n,x} - 1) (W \circ \theta_{n,y} - 1)] \quad (\text{independence}) \\ &= \text{Var}(W) \sum_{x,y} \mathbb{E}[W_n(x) W_n(y)] P_{x,y}^{\otimes 2}(\tau < \infty) \quad (\text{by (17)}) \\ &= \text{Var}(W) \sum_{x,y} P_{0,0}^{\otimes 2} [e^{\lambda_2 N_n} \mathbf{1}_{\{S_n=x, \tilde{S}_n=y\}}] P_{x,y}^{\otimes 2}(\tau < \infty) \quad (\text{cf. (9)}) \\ &= \text{Var}(W) \times P_{0,0}^{\otimes 2}(e^{\lambda_2 N_n} \mathbf{1}_{\tau_n < \infty}), \end{aligned}$$

by Markov property. This is (18).

We finally derive (19) using classical estimates for the Green function. Denote by

$$G(x) := P_{0,x}^{\otimes 2} \left[\sum_{n=0}^{\infty} \mathbf{1}_{\{S_n - \tilde{S}_n = 0\}} \right] = P_{0,x}^{\otimes 2} (N_{\infty}) \quad (20)$$

the Green function for the symmetrized walk $(S_n - \tilde{S}_n)_n$, and observe that it is equal on even sites x (i.e., when $\|x\|_1 = 0 \pmod{2}$) to the Green function of the simple random walk : for even sites x

$$G(x) = P_x \left[\sum_{n=0}^{\infty} \mathbf{1}_{S_n=0} \right],$$

since the process $(S_n - \tilde{S}_n)$ under $P_{0,x}^{\otimes 2}$ has the same law as (S_{2n}) under P_x and $S_{2n+1} \neq 0$ P_x a.s. (for even sites x). For odd sites x , $G(x) = 0$, since in this case $S_n - \tilde{S}_n \neq 0$ $P_{0,x}^{\otimes 2}$ a.s.. From the geometric distribution of N_{∞} under $P_{0,0}^{\otimes 2}$, we have $G(0) = (1 - \pi_d)^{-1}$. By Markov property we have

$$P_{0,x}^{\otimes 2}(\tau < \infty) = G(x)/G(0), \quad x \in \mathbb{Z}^d,$$

and can use the classical estimates for the Green function, see e.g. [25, Theorem 4.3.1] : for even sites x ,

$$G(x) = \frac{\mathfrak{K}_d}{|x|^{d-2}} + \mathcal{O}\left(\frac{1}{|x|^d}\right), \quad |x| \rightarrow \infty, \quad (21)$$

where $\mathfrak{K}_d \in (0, \infty)$ is a constant whose value is

$$\mathfrak{K}_d = \frac{d \Gamma(d/2)}{(d-2)\pi^{d/2}}. \quad (22)$$

By the central limit theorem, we have the following convergence in distribution under $P_{0,0}^{\otimes 2}$ to a Gaussian vector :

$$n^{-1/2}(S_n - \tilde{S}_n) \rightarrow Z \quad \text{in distribution,}$$

where Z is a Gaussian vector with mean 0 and covariance matrix $\frac{2}{d} I_d$, I_d being the d -dimensional identity matrix. Thus, with (21),

$$n^{(d-2)/2} G(S_n - \tilde{S}_n) \rightarrow \frac{\mathfrak{K}_d}{|Z|^{d-2}} \quad \text{in distribution.} \quad (23)$$

We shall need the following two lemmas, whose proofs are postponed by the end of the section.

Lemma 2.2 (Asymptotic independence) *Under $P^{\otimes 2}$ we have the following convergence in distribution of random vectors :*

$$\left(N_n, n^{-1/2} S_n, n^{-1/2} \tilde{S}_n \right) \longrightarrow (N_{\infty}, Z_1, \tilde{Z}_1) \quad \text{in distribution,}$$

where $N_{\infty}, Z_1, \tilde{Z}_1$ are independent, and where

- N_{∞} is geometrically distributed with parameter $1 - \pi_d$ (see (10)),
- Z_1 and \tilde{Z}_1 are Gaussian vectors with mean 0 and covariance matrix $\frac{1}{d} I_d$.

Lemma 2.3 (Boundedness in $L^{1+\delta}$) (a) For $a > 0$,

$$\limsup_{n \rightarrow \infty} P^{\otimes 2} \left(\left| \frac{S_n - \tilde{S}_n}{\sqrt{n}} \right|^{-a} \mathbf{1}_{\{S_n - \tilde{S}_n \neq 0\}} \right) < \infty \quad \text{if and only if } a < d. \quad (24)$$

(b) If $\lambda_2 < \frac{2}{d} \ln \frac{1}{\pi_d}$, then for $\delta > 0$ small enough,

$$\limsup_{n \rightarrow \infty} P^{\otimes 2} [e^{\lambda_2 N_n} n^{\frac{d-2}{2}} G(S_n - \tilde{S}_n)]^{1+\delta} < \infty. \quad (25)$$

We first end the proof of Proposition 2.1. From Lemma 2.2, with $Z = Z_1 - \tilde{Z}_1$,

$$e^{\lambda_2 N_n} \times n^{\frac{d-2}{2}} G(S_n - \tilde{S}_n) \xrightarrow{\text{law}} e^{\lambda_2 N_\infty} \times \frac{\mathfrak{K}_d}{|Z|^{d-2}}. \quad (26)$$

By Lemma 2.3(b), the sequence in the left-hand side of (26) is uniformly integrable, so that the convergence in law implies the convergence of expectations, allowing to write

$$\begin{aligned} n^{\frac{d-2}{2}} P_{0,0}^{\otimes 2} (e^{\lambda_2 N_n} \mathbf{1}_{\tau_n < \infty}) &= n^{\frac{d-2}{2}} P_{0,0}^{\otimes 2} [e^{\lambda_2 N_n} P_{0,0}^{\otimes 2} (\tau_n < \infty | \mathcal{F}_n)] \\ &= G(0)^{-1} P_{0,0}^{\otimes 2} [e^{\lambda_2 N_n} \times n^{\frac{d-2}{2}} G(S_n - \tilde{S}_n)] \\ &\longrightarrow G(0)^{-1} P_{0,0}^{\otimes 2} [e^{\lambda_2 N_\infty}] \times E \left[\frac{\mathfrak{K}_d}{|Z|^{d-2}} \right]. \end{aligned} \quad (27)$$

Together with (18), this ends the proof of (19), yielding the value

$$\begin{aligned} \sigma^2 &= \frac{\mathfrak{K}_d \mathfrak{Z}_d}{G(0)} \text{Var}(W) \mathbb{E}(W^2) \\ &= \mathfrak{K}_d \mathfrak{Z}_d (1 - \pi_d)^2 \times (e^{\lambda_2} - 1) e^{\lambda_2} \times \frac{1}{(1 - \pi_d e^{\lambda_2})^2}, \end{aligned} \quad (28)$$

with $\text{Var}(W)$ from (11) and

$$\mathfrak{Z}_d \stackrel{\text{def}}{=} E \left[\frac{1}{|Z|^{d-2}} \right] = \frac{1}{\Gamma(d/2)} \left(\frac{d}{4} \right)^{(d-2)/2} \quad (29)$$

from the chi-square distribution. This proof of Proposition 2.1 is complete. ■

For later purposes, define

$$\sigma_1^2 = \mathfrak{K}_d \mathfrak{Z}_d (1 - \pi_d) \times \text{Var}(W), \quad (30)$$

so that $\sigma^2 = \sigma_1^2 \mathbb{E}W^2$.

2.2 Proof of the lemmas

It remains to prove Lemmas 2.2 and 2.3.

Proof of Lemma 2.2. First observe that

$$\sup_{n \geq m} P^{\otimes 2} (N_n > N_m) \rightarrow 0 \quad \text{as } m \rightarrow \infty,$$

since $N_n \nearrow N_\infty < \infty$ a.s. Fix $m \geq 1$ and f, g, \tilde{g} continuous and bounded. For all $n \geq m$, we write

$$\begin{aligned}
& P^{\otimes 2} \left[f(N_n)g(n^{-1/2}S_n)\tilde{g}(n^{-1/2}\tilde{S}_n) \right] \\
&= P^{\otimes 2} \left[f(N_n)g(n^{-1/2}S_n)\tilde{g}(n^{-1/2}\tilde{S}_n)\mathbf{1}_{N_n=N_m} \right] + \epsilon(n, m) \\
&= P^{\otimes 2} \left[f(N_m)g(n^{-1/2}S_n)\tilde{g}(n^{-1/2}\tilde{S}_n)\mathbf{1}_{N_n=N_m} \right] + \epsilon(n, m) \\
&= P^{\otimes 2} \left[f(N_m)g(n^{-1/2}(S_n - S_m))\tilde{g}(n^{-1/2}(\tilde{S}_n - \tilde{S}_m))\mathbf{1}_{N_n=N_m} \right] + \epsilon'(n, m) \\
&= P^{\otimes 2} \left[f(N_m)g(n^{-1/2}(S_n - S_m))\tilde{g}(n^{-1/2}(\tilde{S}_n - \tilde{S}_m)) \right] + \epsilon''(n, m) \\
&= P^{\otimes 2}[f(N_m)] \times P[g(n^{-1/2}(S_n - S_m))] \times P[\tilde{g}(n^{-1/2}(\tilde{S}_n - \tilde{S}_m))] + \epsilon''(n, m),
\end{aligned}$$

where the equalities define the terms $\epsilon(n, m), \epsilon'(n, m), \epsilon''(n, m)$ on their first occurrence. Here,

$$|\epsilon(n, m)| \leq \|f\|_\infty \|g\|_\infty \|\tilde{g}\|_\infty P(N_n \neq N_m)$$

tends to 0 as $m \rightarrow \infty$ uniformly in $n \geq m$, $\epsilon'(n, m) - \epsilon(n, m) \rightarrow 0$ as $n \rightarrow \infty$ for all fixed m , and $\sup_{n \geq m} \epsilon''(n, m) \rightarrow 0$ as $m \rightarrow \infty$. The last equality comes from independence in the increments of the random walks, and of the two random walks S and \tilde{S} . Hence, letting $n \rightarrow \infty$ and then $m \rightarrow \infty$, we get

$$P^{\otimes 2} \left[f(N_n)g(n^{-1/2}S_n)\tilde{g}(n^{-1/2}\tilde{S}_n) \right] \rightarrow P^{\otimes 2}[f(N_\infty)] \times P[g(Z_1)] \times P[\tilde{g}(\tilde{Z}_1)].$$

Since N_∞ is geometrically distributed with parameter $1 - \pi_d$ (see (10)), this ends the proof of the lemma. \blacksquare

Remark 2.4 *We could have taken another route to prove the lemma. The couple $n^{-1/2}(S_n, \tilde{S}_n)$ converges (mixing) to the Gaussian vector (Z_1, \tilde{Z}_1) , see e.g. Theorem 2 in [3]. On the other hand $N_n \rightarrow N_\infty$ a.s. From the mixing consequence that we mentioned above Theorem 1.1 (which remains valid for random variables with values in \mathbb{R}^d) it follows the convergence in distribution of $(n^{-1/2}S_n, n^{-1/2}\tilde{S}_n, N_\infty)$ to (Z_1, \tilde{Z}_1, N) . It is not difficult to see that the sequence $(n^{-1/2}S_n, n^{-1/2}\tilde{S}_n, N_\nu)$ has the same limit in distribution. However, we have given the above proof, which is short and instructive, for the sake of completeness.*

Proof of Lemma 2.3. (a) The "only if" part is evident by Fatou's lemma since

$$\left| \frac{S_n - \tilde{S}_n}{\sqrt{n}} \right|^{-a} \mathbf{1}_{\{S_n - \tilde{S}_n \neq 0\}} \rightarrow |Z|^{-a} \quad \text{in distribution}$$

and $E|Z|^{-a} < \infty$ if and only if $a < d$.

Let's show the "if" part. Since the L^a -norm is increasing in a , we only need to prove the finiteness of the lim sup for $0 < a < d$ sufficiently close to d . So we fix $a \in (d - 1, d)$. By the local central limit theorem (see e.g. [24, Theorem 1.2.1, p.14]) we know that

$$p_n(x) := P^{\otimes 2}(S_n - \tilde{S}_n = x) = P(S_{2n} = x)$$

satisfies

$$|p_n(x) - \bar{p}_n(x)| \leq c_2 n^{-d/2} |x|^{-2}, \text{ with } \bar{p}_n(x) = c_1 n^{-d/2} \exp\left\{-\frac{d|x|^2}{4n}\right\}, \quad (31)$$

where $c_1, c_2 > 0$ are constants. (In fact we have $c_1 = \mathfrak{C}_d$, with \mathfrak{C}_d defined by (63).) Denote by I_n the integral in (24). Then

$$I_n \leq \sum_{x \neq 0} \left(\frac{|x|}{\sqrt{n}}\right)^{-a} \bar{p}_n(x) + \sum_{x \neq 0} \left(\frac{|x|}{\sqrt{n}}\right)^{-a} c_2 n^{-d/2} |x|^{-2} := I_{n,1} + I_{n,2}. \quad (32)$$

In the following to avoid sums over non-integer valued numbers, we shall use the integer valued L^1 norm $\|x\|_1 = |x_1| + \dots + |x_d|$ for $x = (x_1, \dots, x_d) \in \mathbb{Z}^d$, instead of the Euclidean norm $|x|$, and the elementary inequality $\|x\|_1/d \leq |x| \leq \|x\|_1$ valid for all $x \in \mathbb{Z}^d$.

For the first sum in (32), we have, for some constant $c_3 > 0$,

$$\begin{aligned} I_{n,1} &\leq c_1 d^a n^{-d/2} \sum_{x \neq 0} \left(\frac{\|x\|_1}{\sqrt{n}}\right)^{-a} \exp\left\{-\frac{\|x\|_1^2}{4dn}\right\} \\ &= c_1 d^a n^{-d/2} \sum_{r \geq 1} \sum_{\|x\|_1=r} \left(\frac{r}{\sqrt{n}}\right)^{-a} \exp\left\{-\frac{r^2}{4dn}\right\} \\ &\leq c_1 c_3 d^a n^{-d/2} \sum_{r \geq 1} r^{d-1} \left(\frac{r}{\sqrt{n}}\right)^{-a} \exp\left\{-\frac{r^2}{4dn}\right\} \\ &= \frac{c_1 c_3 d^a}{\sqrt{n}} \sum_{r \geq 1} \left(\frac{r}{\sqrt{n}}\right)^{-(a-d+1)} \exp\left\{-\frac{1}{4d} \left(\frac{r}{\sqrt{n}}\right)^2\right\}, \end{aligned}$$

where the next to last step holds as the number of $x = (x_1, \dots, x_d) \in \mathbb{Z}^d$ with $|x_1| + \dots + |x_d| = r$ is bounded by $2(2r+1)^{d-1} \leq c_3 r^{d-1}$ (notice that each coordinate satisfies $|x_i| \leq r$, and while the first $d-1$ are chosen, the absolute value of the last coordinate is determined by the equation, so that the last coordinate has at most 2 possibilities). As $0 < a-d+1 < 1$, we have, for all $n \geq 1$,

$$\begin{aligned} I_{n,1} &\leq \frac{c_1 c_3 d^a}{\sqrt{n}} \left(\sum_{1 \leq r \leq \sqrt{n}} \left(\frac{r}{\sqrt{n}}\right)^{-(a-d+1)} + \sum_{r > \sqrt{n}} \exp\left\{-\frac{1}{4d} \left(\frac{r}{\sqrt{n}}\right)^2\right\} \right) \\ &\leq c_1 c_3 d^a \left(\sum_{1 \leq r \leq \sqrt{n}} \int_{\frac{r-1}{\sqrt{n}}}^{\frac{r}{\sqrt{n}}} x^{-(a-d+1)} dx + \sum_{r > \sqrt{n}} \int_{\frac{r-1}{\sqrt{n}}}^{\frac{r}{\sqrt{n}}} \exp\left\{-\frac{x^2}{4d}\right\} dx \right) \\ &\leq c_4 := c_1 c_3 d^a \left(\int_0^1 x^{-(a-d+1)} dx + \int_0^\infty \exp\left\{-\frac{x^2}{4d}\right\} dx \right) < \infty. \end{aligned}$$

Similarly, for the second sum in (32), we have,

$$\begin{aligned} I_{n,2} &\leq c_2 c_3 d^{a+2} n^{-d/2} \sum_{r \geq 1} \left(\frac{r}{\sqrt{n}}\right)^{-a} r^{-2} r^{d-1} \\ &= c_2 c_3 d^{a+2} n^{(a-d)/2} \sum_{r \geq 1} r^{-(a+3-d)} \end{aligned}$$

which tends to 0 as $n \rightarrow \infty$ (since $d-1 < a < d$). This ends the proof of Part (a) of Lemma 2.3.

(b) Let $G_n = G_n(S_n - \tilde{S}_n) := n^{\frac{d-2}{2}} G(S_n - \tilde{S}_n)$ and $\lambda'_2 = (1 + \delta)\lambda_2$. Recall from (21) that $G(x) \leq c_5|x|^{2-d}$. Then

$$\begin{aligned} P^{\otimes 2} [(e^{\lambda_2 N_n} G_n)^{1+\delta}] &= \sum_z P^{\otimes 2} [e^{\lambda'_2 N_n}; S_n - \tilde{S}_n = z] G_n(z)^{1+\delta} \\ &\leq c_5 \sum_{z \neq 0} P^{\otimes 2} [e^{\lambda'_2 N_n}; S_n - \tilde{S}_n = z] |n^{-1/2} z|^{-(d-2)(1+\delta)} \\ &\quad + c_5 P^{\otimes 2} [e^{\lambda'_2 N_n}; S_n - \tilde{S}_n = 0] n^{(d-2)(1+\delta)/2}. \end{aligned} \quad (33)$$

Denote by $L_n = \sup\{j = 0, \dots, n : S_j - \tilde{S}_j = 0\}$ and $T = \inf\{j \geq 1 : S_j - \tilde{S}_j = 0\}$ the last (before n) and the first hitting times of 0. Let $\delta > 0$ be small enough such that $\lambda'_2 < \log \frac{1}{\pi_d}$; this is possible thanks to the condition **(L2)** (see (7)). Then

$$P^{\otimes 2} [e^{\lambda'_2 N_n}; S_n - \tilde{S}_n = 0] \sim e^{\lambda'_2} [1 - e^{\lambda'_2 \pi_d}]^{-2} P^{\otimes 2}[T = n] \quad \text{as } n \rightarrow \infty, \quad (34)$$

by Theorem 2.2 (case 2) in [13]. Since $P^{\otimes 2}[T = n] \leq P^{\otimes 2}(S_n - \tilde{S}_n = 0) = O(n^{-d/2})$, this implies that when $(d-2)(1+\delta) < d$, the last term in (33) vanishes as $n \rightarrow \infty$, and yields, for $z \neq 0$,

$$\begin{aligned} P^{\otimes 2} [e^{\lambda'_2 N_n}; S_n - \tilde{S}_n = z] &= \sum_{j=0}^n P^{\otimes 2} [e^{\lambda'_2 N_n}; S_n - \tilde{S}_n = z, L_n = j] \\ &\stackrel{\text{Markov}}{=} \sum_{j=0}^n P^{\otimes 2} [e^{\lambda'_2 N_j}; S_j - \tilde{S}_j = 0] P^{\otimes 2} [S_{n-j} - \tilde{S}_{n-j} = z, L_{n-j} = 0] \\ &\stackrel{(34)}{\leq} c_6 \sum_{j=0}^n P^{\otimes 2}[T = j] P^{\otimes 2} [S_{n-j} - \tilde{S}_{n-j} = z, L_{n-j} = 0] \\ &\leq c_6 \sum_{j=0}^n P^{\otimes 2}[T = j] P^{\otimes 2} [S_{n-j} - \tilde{S}_{n-j} = z] \\ &= c_6 P^{\otimes 2} [S_n - \tilde{S}_n = z]. \end{aligned}$$

where the last step holds because we have $S_j - \tilde{S}_j = 0$ on $T = j$, so that

$$\begin{aligned} P^{\otimes 2} [S_n - \tilde{S}_n = z] &= \sum_{j=0}^n P^{\otimes 2}[T = j, S_n - \tilde{S}_n = z] \\ &= \sum_{j=0}^n P^{\otimes 2}[T = j, (S_n - S_j) - (\tilde{S}_n - \tilde{S}_j) = z] \\ &= \sum_{j=0}^n P^{\otimes 2}[T = j] P^{\otimes 2}[S_{n-j} - \tilde{S}_{n-j} = z], \end{aligned}$$

using the fact that the event $[T = j]$ is independent of $(S_n - S_j) - (\tilde{S}_n - \tilde{S}_j)$ which has the same law as $S_{n-j} - \tilde{S}_{n-j}$.

Inserting the last bound on $P^{\otimes 2} [e^{\lambda_2 N_n}; S_n - \tilde{S}_n = z]$ into (33), we obtain for large n ,

$$P^{\otimes 2} [(e^{\lambda_2 N_n} G_n)^{1+\delta}] = c_5 c_6 \left[1 + \sum_{z \neq 0} P^{\otimes 2} [S_n - \tilde{S}_n = z] \left| n^{-1/2} z \right|^{-(d-2)(1+\delta)} \right],$$

which, according to (24), is bounded provided that $(d-2)(1+\delta) < d$. ■

3 Proof of the Central Limit Theorem

In this section we give the proof of Theorem 1.1. Our proof is based on the following central limit theorem for infinite martingale arrays, which is a slight extension of Corollaries 3.1 and 3.2 of the book by Hall and Heyde [14] (pp. 58-59 and p.64), but we could not find it in the literature.

Lemma 3.1 *For $n \geq 1$, let $\{(S_{n,i}, \mathcal{F}_{n,i}) : i \geq 0\}$ be an array of martingales defined on a probability space (Ω, \mathcal{F}, P) , with $S_{n,0} = 0$ and*

$$\sup_{n,i \geq 1} E S_{n,i}^2 < \infty. \quad (35)$$

Let $X_{n,i} = S_{n,i} - S_{n,i-1}$, $i \geq 1$ be the martingale differences, and $S_{n,\infty} = \lim_{i \rightarrow \infty} S_{n,i}$ be the a.s. limit of $(S_{n,i}, i \geq 0)$. Suppose that :

(a) *the conditional variance converges in probability : for a real random variable $V \in [0, \infty)$,*

$$V_{n,\infty}^2 := \sum_{i=1}^{\infty} E(X_{n,i}^2 | \mathcal{F}_{n,i-1}) \longrightarrow V^2 \text{ in probability}; \quad (36)$$

(b) *the conditional Lindeberg condition holds :*

$$\forall \varepsilon > 0, \quad \sum_{i=1}^{\infty} E(X_{n,i}^2 \mathbf{1}_{|X_{n,i}| > \varepsilon} | \mathcal{F}_{n,i-1}) \longrightarrow 0 \text{ in probability}; \quad (37)$$

(c) *the σ -fields are nested : $\mathcal{F}_{n,i} \subset \mathcal{F}_{n+1,i}$ for all $n, i \geq 1$.*

Then

$$S_{n,\infty} \longrightarrow V G \quad \text{in distribution} \quad (38)$$

where G is a Gaussian variable with law $N(0, 1)$ and independent of V ; if additionally $V \neq 0$ a.s., then

$$\frac{S_{n,\infty}}{V_{n,\infty}} \longrightarrow G \quad \text{in distribution.} \quad (39)$$

Moreover, the convergence in (38) is stable, and the convergence in (39) is mixing.

Lemma 3.1 reduces to Corollaries 3.1 and 3.2 in [14] for a triangular array of martingales differences, that is, when $X_{n,i} = 0$ for all $i > k_n$, for some sequence of integers k_n increasing to ∞ . As in the case of a triangular array, if V is measurable with respect to each $\mathcal{F}_{n,i}$ for $n, i \geq 1$ (e.g. when V is a constant), then the nested condition (c) can be removed, but the convergence (38) may no longer be stable, and the convergence (39) may no longer be mixing

(see the remarks in p.59 and p. 64 of [14] for a triangular array). Lemma 3.1 can be extended in a clear way to two-sided martingale arrays $\{(S_{n,i}, \mathcal{F}_{n,i}) : -\infty < i < \infty\}$; for a version using conditions and norming in terms of $\sum_i X_{n,i}^2$, see Theorem 3.6 of [14] (p.77).

Proof of Lemma 3.1. We will see that Lemma 3.1 can be obtained from the corresponding result for a triangular array of martingales. Let k_n be positive integers increasing to ∞ such that

$$E(S_{n,\infty} - S_{n,k_n})^2 = E \sum_{i>k_n} X_{n,i}^2 \rightarrow 0.$$

Then

$$V_{n,k_n}^2 := \sum_{i=1}^{k_n} E(X_{n,i}^2 | \mathcal{F}_{n,i-1}) \rightarrow V^2 \text{ in probability}$$

since $V_{n,\infty}^2 - V_{n,k_n}^2 = \sum_{i>k_n} E(X_{n,i}^2 | \mathcal{F}_{n,i-1}) \rightarrow 0$ in L^1 . Clearly, by condition (b), the conditional Lindeberg condition for the triangular array $\{(X_{n,i}, \mathcal{F}_{n,i}) : 0 \leq i \leq k_n\}$ holds, that is, (37) holds with $\sum_{i=1}^{\infty}$ replaced by $\sum_{i=1}^{k_n}$. Therefore, by Corollaries 3.1 and 3.2 of [14] (pp.58-59 and p. 64),

$$S_{n,k_n} \rightarrow VG \quad \text{in distribution} \quad (40)$$

and

$$\frac{S_{n,k_n}}{V_{n,k_n}} \rightarrow G \quad \text{in distribution.} \quad (41)$$

Since $S_{n,\infty} - S_{n,k_n} \rightarrow 0$ in L^2 and hence in probability, (40) implies (38). As $V_{n,\infty}^2 - V_{n,k_n}^2 \rightarrow 0$ in probability (in fact in L^1), when $V > 0$ a.s. we have $V_{n,\infty}^2/V_{n,k_n}^2 \rightarrow 1$ in probability. Therefore (41) implies (39). ■

Lemma 3.1 is well suited for studying the rate of convergence of a martingale, as shown in the following

Corollary 3.2 *Let $\{(S_i, \mathcal{F}_i) : i \geq 0\}$ be a martingale defined on a probability space (Ω, \mathcal{F}, P) , with $S_0 = 0$ and $\sup_{i \geq 1} ES_i^2 < \infty$. Let $X_i = S_i - S_{i-1}, i \geq 1$ be the martingale differences, $S_\infty = \lim_{i \rightarrow \infty} S_i$ be the a.s. limit of (S_i) , and let*

$$v_n^2 = E(S_\infty - S_n)^2 = E \sum_{i=n+1}^{\infty} X_i^2.$$

Suppose that $v_n > 0$ and that :

(a) *the conditional variance converges in probability : for a real random variable $V \in [0, \infty)$,*

$$V_n^2 := \frac{1}{v_n^2} \sum_{i=n+1}^{\infty} E(X_i^2 | \mathcal{F}_{i-1}) \rightarrow V^2 \text{ in probability;} \quad (42)$$

(b) *the conditional Lindeberg condition holds :*

$$\forall \varepsilon > 0, \quad \frac{1}{v_n^2} \sum_{i=n+1}^{\infty} E(X_i^2 \mathbf{1}_{|X_i| > \varepsilon v_n} | \mathcal{F}_{i-1}) \rightarrow 0 \text{ in probability.} \quad (43)$$

Then

$$\frac{S_\infty - S_n}{v_n} \rightarrow VG \quad \text{in distribution,} \quad (44)$$

where G is a Gaussian variable with law $N(0, 1)$ and independent of V ; if additionally $V \neq 0$ a.s., then

$$\frac{S_\infty - S_n}{V_n} \rightarrow G \quad \text{in distribution.} \quad (45)$$

Moreover, the convergence in (44) is stable, and the convergence in (45) is mixing.

Proof. Corollary 3.2 is an immediate consequence of Lemma 3.1 applied to $X_{n,i} = \frac{X_{n+i}}{v_n}$ for $i \geq 1$, $\mathcal{F}_{n,i} = \mathcal{F}_{n+i}$ for $i \geq 0$, and $X_{n,0} = 0$. \blacksquare

Proof of Theorem 1.1. We can use Corollary 3.2 with $1/\|W - W_n\|_2$ for the norming, but, since $\|W - W_n\|_2 \sim \sigma n^{-(d-2)/4}$, we prefer to use the more explicit norming $n^{(d-2)/4}$, together with the spirit of the proof of Corollary 3.2. So we rely on the decomposition

$$n^{\frac{d-2}{4}}(W - W_n) = n^{\frac{d-2}{4}} \sum_{k=n}^{\infty} D_{k+1}, \quad (46)$$

where

$$D_{k+1} = W_{k+1} - W_k, \quad k \geq n,$$

forms a sequence of martingale differences. To prove Theorem 1.1, by (46) and Lemma 3.1 applied to $X_{n,i} = n^{\frac{d-2}{4}} D_{n+i}$ for $i \geq 1$, $\mathcal{F}_{n,i} = \mathcal{G}_{n+i}$ for $i \geq 0$ (recall (6)), and $X_{n,0} = 0$, it suffices to prove that :

(a) the following convergence about the conditional variance holds :

$$s_n^2 := n^{\frac{d-2}{2}} \sum_{k \geq n} \mathbb{E}_k D_{k+1}^2 \rightarrow \sigma_1^2 W^2 \quad \text{in probability,} \quad (47)$$

where $\mathbb{E}_k(\cdot) = \mathbb{E}(\cdot | \mathcal{G}_k)$ denotes the conditional expectation given \mathcal{G}_k ;

(b) the following Lindeberg condition holds :

$$\forall \epsilon > 0, \quad n^{\frac{d-2}{2}} \sum_{k \geq n} \mathbb{E}_k \left(D_{k+1}^2 \mathbf{1}_{\{n^{\frac{d-2}{4}} |D_{k+1}| > \epsilon\}} \right) \rightarrow 0 \quad \text{in probability.} \quad (48)$$

Actually, by Lemma 3.1, from (47) and (48) we conclude that (12) and (13) hold with the norming $1/W_n$ in (13) replaced by $1/W$. As $W_n/W \rightarrow 1$ a.s. (and thus in probability), we can change the factor $1/W$ to $1/W_n$ without changing the convergence in distribution.

To show the convergence (47) of the conditional variance, we will prove in the next section the following

Lemma 3.3 *There exists $\beta_0 > 0$ such that for $|\beta| < \beta_0$ and σ_1 from (30), we have, as $n \rightarrow \infty$,*

$$\mathbb{E}(W_n - W)^4 \rightarrow 0, \quad (49)$$

$$\mathbb{E}s_n^4 - \sigma_1^4 \mathbb{E}W_n^4 \rightarrow 0, \quad (50)$$

$$\mathbb{E}(s_n^2 W_n^2) - \sigma_1^2 \mathbb{E}W_n^4 \rightarrow 0. \quad (51)$$

It follows from Lemma 3.3 that

$$\begin{aligned}\mathbb{E}(s_n^2 - \sigma_1^2 \mathbb{E}W_n^2)^2 &= \mathbb{E}s_n^4 - 2\sigma_1^2 \mathbb{E}(s_n^2 W_n^2) + \sigma_1^4 \mathbb{E}W_n^4 \\ &\rightarrow 0.\end{aligned}$$

Therefore, $s_n^2 - \sigma_1^2 W_n^2 \rightarrow 0$ in L^2 . As $W_n^2 \rightarrow W^2$ in L^2 , it follows that $s_n^2 \rightarrow \sigma_1^2 W^2$ in L^2 . We thus obtain (47).

To show Lindeberg's condition (48), we will prove the following convergence rate of $\mathbb{E}D_{k+1}^4$.

Lemma 3.4 *For any $q > 1$, when $|\beta| > 0$ is small enough, we have*

$$\mathbb{E}D_{k+1}^4 = O(k^{-d/q}), \quad k \geq 1. \quad (52)$$

The proof of the Lemma is postponed to the end of the paper.

Note that a sufficient condition for Lindeberg's condition (48) to hold is clearly

$$n^{(d-2)} \sum_{k \geq n} \mathbb{E}_k D_{k+1}^4 \rightarrow 0 \quad \text{in probability,}$$

which is implied by

$$n^{(d-2)} \sum_{k \geq n} \mathbb{E}D_{k+1}^4 \rightarrow 0. \quad (53)$$

By Lemma 3.4, when $|\beta| > 0$ is small enough, the left-hand side of (53) is smaller than $c n^{d-1-d/q}$ for some constant $c > 0$, which tends to 0 by taking $1 < q < d/(d-1)$, so that Lindeberg's condition (48) holds. (In fact one can check that it suffices to take $|\beta| < \beta_0$ with $\beta_0 > 0$ determined in Lemma 4.1.) This ends the proof of Theorem 1.1, using Lemmas 3.3 and 3.4 whose proofs will be given in the next section. ■

Remark 3.5 *The convergence (13) can also be proved using the decomposition*

$$\frac{n^{\frac{d-2}{4}}(W - W_n)}{W_n} = \frac{n^{\frac{d-2}{4}}}{W_n} \sum_{k=n}^{\infty} D_{k+1},$$

where $(n^{\frac{d-2}{4}} D_{k+1}/W_n, k \geq n)$ remains a sequence of martingale differences. But proving (12) requires a different route.

We end this section with a **warm-up calculation** : we recover the value of $\mathbb{E}W_n^2$ and (18) from the martingale decomposition (46), i.e., using the conditional variance. This calculation is instructive and it will be useful in the forthcoming computations. Write for short

$$h_k(S) = \sum_{i=0}^{k-1} [\beta \eta(i, S_i) - \lambda(\beta)].$$

Since $W_{k+1} - W_k = P e^{h_k(S)} (e^{\beta \eta(k, S_k) - \lambda(\beta)} - 1)$,

$$(W_{k+1} - W_k)^2 = P^{\otimes 2} \left[e^{h_k(S)} (e^{\beta \eta(k, S_k) - \lambda(\beta)} - 1) e^{h_k(\tilde{S})} (e^{\beta \eta(k, \tilde{S}_k) - \lambda(\beta)} - 1) \right],$$

using Fubini's theorem we have

$$\mathbb{E}_k D_{k+1}^2 = \kappa_2 P^{\otimes 2} e^{h_k(S)} e^{h_k(\tilde{S})} \mathbf{1}_{\{S_k = \tilde{S}_k\}} \quad (54)$$

with

$$\kappa_2 = \kappa_2(\beta) = e^{\lambda_2} - 1. \quad (55)$$

We compute s_n^2 from its definition in (47) :

$$s_n^2 = \kappa_2 n^{\frac{d-2}{2}} \sum_{k \geq n} P^{\otimes 2} e^{h_k(S)} e^{h_k(\tilde{S})} \mathbf{1}_{S_k = \tilde{S}_k}. \quad (56)$$

Observe that

$$\mathbf{1}_{\{S_k = \tilde{S}_k\}} = \frac{e^{\lambda_2(N_{k+1} - N_k)} - 1}{\kappa_2}. \quad (57)$$

We now check that, with s_n^2 from (56) and σ^2 be defined by (28),

$$\mathbb{E} s_n^2 \rightarrow \sigma^2.$$

This is a remake of (19), but, as we will see, from a different route. By Fubini we have

$$\begin{aligned} \mathbb{E} s_n^2 &= \kappa_2 n^{\frac{d-2}{2}} \sum_{k \geq n} P^{\otimes 2} \mathbb{E} e^{h_k(S)} e^{h_k(\tilde{S})} \mathbf{1}_{S_k = \tilde{S}_k} \\ &= n^{\frac{d-2}{2}} P^{\otimes 2} \sum_{k \geq n} e^{\lambda_2 N_k} \kappa_2 \mathbf{1}_{S_k = \tilde{S}_k} \\ &= n^{\frac{d-2}{2}} P^{\otimes 2} \sum_{k \geq n} e^{\lambda_2 N_k} (e^{\lambda_2(N_{k+1} - N_k)} - 1) \quad (\text{by (57)}) \\ &= n^{\frac{d-2}{2}} P^{\otimes 2} (e^{\lambda_2 N_\infty} - e^{\lambda_2 N_n}) \quad (\text{telescopic sum}) \\ &= n^{\frac{d-2}{2}} P^{\otimes 2} [P^{\otimes 2} (e^{\lambda_2(N_\infty - N_n)} - 1) | \mathcal{F}_n] e^{\lambda_2 N_n} \\ &= n^{\frac{d-2}{2}} P^{\otimes 2} [F(S_n - \tilde{S}_n) e^{\lambda_2 N_n}], \end{aligned}$$

where we can express

$$\begin{aligned} F(x) &= P_{0,x}^{\otimes 2} (e^{\lambda_2 N_\infty} - 1) \\ &= P_{0,x}^{\otimes 2} ((e^{\lambda_2 N_\infty} - 1) \mathbf{1}_{\tau < \infty}) \\ &= P_{0,x}^{\otimes 2} (\tau < \infty) P_{0,0}^{\otimes 2} (e^{\lambda_2 N_\infty} - 1) \\ &= \frac{G(x)}{G(0)} (\mathbb{E}(W^2) - 1). \end{aligned} \quad (58)$$

Therefore, by the same argument as in (27), we derive from the last 2 formulas,

$$\begin{aligned} \mathbb{E} s_n^2 &= \frac{\text{Var}(W)}{G(0)} P^{\otimes 2} [n^{\frac{d-2}{2}} G(S_n - \tilde{S}_n) e^{\lambda_2 N_n}] \\ &\rightarrow \frac{\mathfrak{K}_d}{G(0)} \text{Var}(W) E \frac{1}{|Z|^{d-2}} \times P^{\otimes 2} e^{\lambda_2 N_\infty} \\ &= \mathfrak{K}_d \text{Var}(W) (1 - \pi_d) \mathfrak{Z}_d \times \mathbb{E} W^2 \end{aligned}$$

which is equal to σ^2 .

4 Proof of the instrumental lemmas

In this section we give the proofs of Lemmas 3.3 and 3.4.

Proof of Lemma 3.3. We first give the **proof of** (49). For 4 independent paths $S^{(i)}$, $i = 1, \dots, 4$, we need to count the number of intersections of 2, 3 or four of them. Denote by E_4 the set of elements of $(\mathbb{Z}^d)^4$ with all 4 elements equal, by E_3 the set of those with 3 equal lattice sites and a different fourth, by $E_{2,2}$ the set of those with 2 pairs of equal lattice sites but the two are different, and by $E_{2,0}$ the set of those with one pair being equal and different from the two other ones. Let A be the corresponding set of indices, $A = \{4, 3, (2, 2), (2, 0)\}$, and define for $m \geq 2$, $n < k \leq \infty$ and $a \in A$,

$$\begin{aligned}\lambda_m &= \lambda(m\beta) - m\lambda(\beta), \\ N_{n,k}^{(a)} &= \sum_{n \leq t < k} \mathbf{1}_{\{(S_t^{(1)}, S_t^{(2)}, S_t^{(3)}, S_t^{(4)}) \in E_a\}}, \\ \Sigma_{n,k} &= \lambda_4 N_{n,k}^{(4)} + \lambda_3 N_{n,k}^{(3)} + 2\lambda_2 N_{n,k}^{(2,2)} + \lambda_2 N_{n,k}^{(2,0)}.\end{aligned}$$

Then it is elementary to check that

$$\mathbb{E} \exp\left\{\sum_{i=1}^4 h_n(S^{(i)})\right\} = \exp \Sigma_{0,n}, \quad \mathbb{E} W_n^4 = P^{\otimes 4} (e^{\Sigma_{0,n}}). \quad (59)$$

For all pair S, \tilde{S} of paths and all $0 \leq n < k \leq \infty$, put

$$N_{n,k}(S, \tilde{S}) = \sum_{n \leq t < k} \mathbf{1}_{S_t = \tilde{S}_t}.$$

For simplicity we sometimes just write $N_{n,k}$ for $N_{n,k}(S, \tilde{S})$ when there is no confusion. Note that, for all $a \in A$,

$$N_{n,k}^{(a)} \leq \sum_{1 \leq i < j \leq 4} N_{n,k}(S^{(i)}, S^{(j)}), \quad (60)$$

and that $\lambda_m \searrow 0$ as $|\beta| \searrow 0$. Thus, for each $t > 0$, writing $\lambda := 3\lambda_2 + \lambda_3 + \lambda_4$ and taking $|\beta| > 0$ small enough such that $\lambda < \frac{1}{6t} \log \frac{1}{\pi_d}$, we have

$$\begin{aligned}P^{\otimes 4} (e^{t\Sigma_{0,\infty}}) &\leq P^{\otimes 4} e^{t\lambda \sum_{1 \leq i < j \leq 4} N_{0,\infty}(S^{(i)}, S^{(j)})} \\ &\leq P^{\otimes 2} e^{6t\lambda N_\infty} < \infty\end{aligned} \quad (61)$$

(recall the calculation after (10)), where the second inequality holds by Hölder's inequality $E|f_1 \cdots f_6| \leq (E|f_1|^6)^{1/6} \cdots (E|f_6|^6)^{1/6}$ and the fact that the number of pairs (i, j) with $1 \leq i < j \leq 4$ is equal to 6. Therefore when $|\beta| > 0$ is small enough such that $\lambda < \frac{1}{6} \log \frac{1}{\pi_d}$, we have

$$\sup_{n \geq 1} \mathbb{E} W_n^4 = P^{\otimes 4} (e^{\Sigma_{0,\infty}}) \leq P^{\otimes 2} e^{6\lambda N_\infty} < \infty.$$

This shows that the martingale (W_n) is bounded in L^4 , so that it converges in L^4 , which gives the assertion (49).

We next give the **proof of (50)**. We compute

$$\begin{aligned}\mathbb{E}S_n^4 &= \kappa_2^2 n^{d-2} \sum_{k, \ell \geq n} P^{\otimes 4} \left[\mathbb{E} e^{h_k(S^{(1)}) + h_k(S^{(2)}) + h_\ell(S^{(3)}) + h_\ell(S^{(4)})} \mathbf{1}_{S_k^{(1)}=S_k^{(2)}} \mathbf{1}_{S_\ell^{(3)}=S_\ell^{(4)}} \right] \\ &= T^<(n) + T^=(n) + T^>(n),\end{aligned}$$

where $T^=(n)$ [resp. $T^<(n)$, resp. $T^>(n)$] is the contribution in the sum of the terms with $k = \ell$, [resp. $k < \ell$, resp. $k > \ell$]. Thus,

$$T^=(n) = \kappa_2^2 n^{d-2} \sum_{k \geq n} P^{\otimes 4} \left[e^{\Sigma_{0,k}} \mathbf{1}_{S_k^{(1)}=S_k^{(2)}} \mathbf{1}_{S_k^{(3)}=S_k^{(4)}} \right].$$

By symmetry, $T^<(n) = T^>(n)$. We calculate $T^<(n)$:

$$\begin{aligned}T^<(n) &= \kappa_2^2 n^{d-2} P^{\otimes 4} \left[\sum_{\ell > k \geq n} e^{\Sigma_{0,k} + \lambda_2 N_{k,\ell}(S^{(3)}, S^{(4)})} \mathbf{1}_{S_k^{(1)}=S_k^{(2)}} \mathbf{1}_{S_\ell^{(3)}=S_\ell^{(4)}} \right] \\ &= \kappa_2 n^{d-2} P^{\otimes 4} \left[\sum_{k \geq n} e^{\Sigma_{0,k}} \mathbf{1}_{S_k^{(1)}=S_k^{(2)}} \sum_{\ell > k} e^{\lambda_2 N_{k,\ell}(S^{(3)}, S^{(4)})} \kappa_2 \mathbf{1}_{S_\ell^{(3)}=S_\ell^{(4)}} \right] \\ &\stackrel{(57)}{=} \kappa_2 n^{d-2} P^{\otimes 4} \left[\sum_{k \geq n} e^{\Sigma_{0,k}} \mathbf{1}_{S_k^{(1)}=S_k^{(2)}} \sum_{\ell > k} e^{\lambda_2 N_{k,\ell}(S^{(3)}, S^{(4)})} \left(e^{\lambda_2 N_{\ell,\ell+1}(S^{(3)}, S^{(4)})} - 1 \right) \right] \\ &= \kappa_2 n^{d-2} P^{\otimes 4} \left[\sum_{k \geq n} e^{\Sigma_{0,k}} \mathbf{1}_{S_k^{(1)}=S_k^{(2)}} \sum_{\ell > k} \left(e^{\lambda_2 N_{k,\ell+1}(S^{(3)}, S^{(4)})} - e^{\lambda_2 N_{k,\ell}(S^{(3)}, S^{(4)})} \right) \right] \\ &= \kappa_2 n^{d-2} P^{\otimes 4} \left[\sum_{k \geq n} e^{\Sigma_{0,k}} \mathbf{1}_{S_k^{(1)}=S_k^{(2)}} \left(e^{\lambda_2 N_{k,\infty}(S^{(3)}, S^{(4)})} - e^{\lambda_2 N_{k,k+1}(S^{(3)}, S^{(4)})} \right) \right].\end{aligned}$$

Therefore, using (57) and the expression of $T^=(n)$, we obtain

$$\begin{aligned}T^<(n) &= \kappa_2 n^{d-2} P^{\otimes 4} \left[\sum_{k \geq n} e^{\Sigma_{0,k}} \mathbf{1}_{S_k^{(1)}=S_k^{(2)}} \left(e^{\lambda_2 N_{k,\infty}(S^{(3)}, S^{(4)})} - 1 \right) \right] - T^=(n) \\ &= \kappa_2 n^{d-2} P^{\otimes 4} \left[\sum_{k \geq n} e^{\Sigma_{0,k}} \mathbf{1}_{S_k^{(1)}=S_k^{(2)}} P^{\otimes 2} \left(e^{\lambda_2 N_{k,\infty}(S^{(3)}, S^{(4)})} - 1 \mid \mathcal{F}_k \right) \right] - T^=(n) \\ &\stackrel{(58)}{=} \frac{\text{Var}(W)}{G(0)} \kappa_2 \times n^{d-2} P^{\otimes 4} \left[\sum_{k \geq n} e^{\Sigma_{0,k}} \mathbf{1}_{S_k^{(1)}=S_k^{(2)}} G(S_k^{(3)} - S_k^{(4)}) \right] - T^=(n) \\ &= T_1^<(n) - T^=(n),\end{aligned}$$

which serves also as the definition of $T_1^<(n)$. For $0 \leq m \leq n$ define

$$T_2^<(m, n) = \frac{\text{Var}(W)}{G(0)} \kappa_2 \times n^{d-2} \sum_{k \geq n} P^{\otimes 4} \left[e^{\Sigma_{0,m} + \lambda_2 N_{k-m,k}(S^{(1)}, S^{(2)})} \mathbf{1}_{S_k^{(1)}=S_k^{(2)}} G(S_k^{(3)} - S_k^{(4)}) \right].$$

It is sufficient to study the limit of $T_2^<(m, n)$ for large n, m , since we will prove the following

Lemma 4.1 *Let $\beta_0 > 0$ be such that, for some $\epsilon > 0$ and all $|\beta| < \beta_0$, $P^{\otimes 4}[e^{(d+\epsilon)\Sigma_{0,\infty}}] < \infty$. Then*

$$\lim_{n \rightarrow \infty} T^-(n) = \lim_{m \rightarrow \infty} \limsup_{n \rightarrow \infty} |T_1^<(n) - T_2^<(m, n)| = 0.$$

Remark 4.2 *The occurrence of the term $N_{k-m,k}$ may be surprising. It is reminiscent of a similar phenomenon in the local limit theorem for polymers [29, 30]. The constraint $S_k^{(1)} = S_k^{(2)}$ makes likely intersections between $S^{(1)}$ and $S^{(2)}$ just before time k , whereas it is likely that $S^{(3)}$ and $S^{(4)}$ are far from them and far apart one from another.*

With Lemma 4.1 we continue our proof. To analyse $T_2^<(m, n)$ we condition on the vectors

$$\begin{aligned} \mathcal{S}^{(1,2)} &= \mathcal{S}_{0,m}^{(1,2)} := (S_t^{(i)}; i = 1, 2, t = 0, \dots, m-1), \\ \mathcal{S}^{(3,4)} &= \mathcal{S}_{0,m}^{(3,4)} := (S_t^{(i)}; i = 3, 4, t = 0, \dots, m-1), \end{aligned}$$

and use the independence of the paths $S^{(i)}$ to obtain that for fixed m , as $n \rightarrow \infty$,

$$\begin{aligned} T_2^<(m, n) &= \frac{\text{Var}(W)}{G(0)} \kappa_2 n^{d-2} \sum_{k \geq n} P^{\otimes 4} \left[e^{\Sigma_{0,m}} P^{\otimes 2}(\mathbf{1}_{S_k^{(1)}=S_k^{(2)}} e^{\lambda_2 N_{k-m,k}(S^{(1)}, S^{(2)})} | \mathcal{S}^{(1,2)}) \right. \\ &\quad \left. \times P^{\otimes 2}(G(S_k^{(3)} - S_k^{(4)}) | \mathcal{S}^{(3,4)}) \right] \\ &\sim \frac{\text{Var}(W) \mathfrak{K}_d \mathfrak{J}_d}{G(0)} \kappa_2 n^{d-2} \sum_{k \geq n} \frac{1}{k^{\frac{d-2}{2}}} P^{\otimes 4} \left[e^{\Sigma_{0,m}} P^{\otimes 2}(\mathbf{1}_{S_k^{(1)}=S_k^{(2)}} e^{\lambda_2 N_{k-m,k}(S^{(1)}, S^{(2)})} | \mathcal{S}^{(1,2)}) \right]. \end{aligned} \quad (62)$$

Indeed, using the independence between $S_k^{(i)} - S_{m-1}^{(i)}$ and $S_{m-1}^{(i)}$ and the fact that $S_k^{(i)} - S_{m-1}^{(i)}$ has the same law as $S_{k-m+1}^{(i)}$, we have,

$$\begin{aligned} P^{\otimes 2}(G(S_k^{(3)} - S_k^{(4)}) | \mathcal{S}^{(3,4)}) &= P^{\otimes 2}(G((S_k^{(3)} - S_{m-1}^{(3)}) - (S_k^{(4)} - S_{m-1}^{(4)}) + (S_{m-1}^{(3)} - S_{m-1}^{(4)})) | \mathcal{S}^{(3,4)}) \\ &= g_{k-m+1}(S_{m-1}^{(3)} - S_{m-1}^{(4)}), \end{aligned}$$

where $g_{k-m+1}(x) = P^{\otimes 2}G(S_{k-m+1}^{(3)} - S_{k-m+1}^{(4)} + x)$, $x \in \mathbb{Z}^d$. It can be easily seen that (23), (24), (25) still hold with $S_n - \tilde{S}_n$ replaced by $S_n - \tilde{S}_n + x$ for any fixed $x \in \mathbb{Z}^d$ (with the same argument). This shows that for any $x \in \mathbb{Z}^d$,

$$k^{\frac{d-2}{2}} G(S_k^{(3)} - S_k^{(4)} + x) \rightarrow \frac{\mathfrak{K}_d}{|Z|^{d-2}} \quad \text{in distribution,}$$

and that the sequence $k^{\frac{d-2}{2}} G(S_k^{(3)} - S_k^{(4)} + x)$ is $L^{1+\delta}$ -bounded under $P^{\otimes 2}$ (see (25) with $\lambda_2 = 0$), and is therefore uniformly integrable. With the uniform integrability, the convergence in distribution implies the convergence of expectations. Thus for each fixed $x \in \mathbb{Z}^d$,

$$k^{\frac{d-2}{2}} P^{\otimes 2} G(S_k^{(3)} - S_k^{(4)} + x) \rightarrow \mathfrak{K}_d \mathfrak{J}_d.$$

It follows that as $k \rightarrow \infty$,

$$k^{\frac{d-2}{2}} P^{\otimes 2}(G(S_k^{(3)} - S_k^{(4)}) | \mathcal{S}^{(3,4)}) \rightarrow \mathfrak{K}_d \mathfrak{J}_d.$$

For each fixed m , the convergence is uniform over all possible values of $\mathcal{S}^{(3,4)}$, since the number of possible values is finite due to the fact that $|S_t^{(i)}| \leq m$ for $0 \leq t \leq m-1$. Therefore for

each $\varepsilon > 0$, there is $k_\varepsilon > 0$ such that for all $k \geq k_\varepsilon$, $k^{\frac{d-2}{2}} P^{\otimes 2}(G(S_k^{(3)} - S_k^{(4)}) | \mathcal{S}^{(3,4)})$ lies between $\mathfrak{K}_d \mathfrak{Z}_d - \varepsilon$ and $\mathfrak{K}_d \mathfrak{Z}_d + \varepsilon$. This implies (62).

Now by Markov property of the random walks $(S_t^{(i)})_t$ and time-reversal on the event $\{S_k^{(1)} = S_k^{(2)}\}$, writing

$$\begin{aligned} f_{k-m+1}(s^{(1)}, s^{(2)}) &= P^{\otimes 2}(\mathbf{1}_{\{S_{k-m+1}^{(1)} + s^{(1)} = S_{k-m+1}^{(2)} + s^{(2)}\}} e^{\lambda_2 \sum_{t=1}^m \mathbf{1}_{\{S_t^{(1)} = S_t^{(2)}\}}} \\ &= e^{-\lambda_2} P^{\otimes 2} \mathbf{1}_{\{S_{k-m+1}^{(1)} + s^{(1)} = S_{k-m+1}^{(2)} + s^{(2)}\}} (e^{\lambda_2 N_{0,m+1}}), \end{aligned}$$

we have, when $k - m > m \geq 1$,

$$\begin{aligned} &P^{\otimes 2}(\mathbf{1}_{\{S_k^{(1)} = S_k^{(2)}\}} e^{\lambda_2 N_{k-m,k}(S^{(1)}, S^{(2)})} | \mathcal{S}^{(1,2)}) \\ &= P^{\otimes 2}(\mathbf{1}_{\{S_k^{(1)} = S_k^{(2)}\}} e^{\lambda_2 N_{k-m,k}(S^{(1)}, S^{(2)})} | S_{m-1}^{(i)}, i = 1, 2) \\ &= P^{\otimes 2}(\mathbf{1}_{\{S_k^{(1)} - S_{m-1}^{(1)} + S_{m-1}^{(1)} = S_k^{(2)} - S_{m-1}^{(2)} + S_{m-1}^{(2)}\}} e^{\lambda_2 \sum_{t=k-m}^{k-1} \mathbf{1}_{\{S_t^{(1)} - S_t^{(1)} = S_t^{(2)} - S_t^{(2)}\}}} | S_{m-1}^{(i)}, i = 1, 2) \\ &= f_{k-m+1}(S_{m-1}^{(1)}, S_{m-1}^{(2)}). \end{aligned}$$

where the last step holds because the reverse random walk $\{S_k^{(i)} - S_t^{(i)} : t = k-1, \dots, k-m, \dots, m-1\}$ has the same law as the initial random walk $\{S_t^{(i)} : t = 1, \dots, m, \dots, k-m+1\}$. Now for each possible value $(s^{(1)}, s^{(2)})$ of $(S_{m-1}^{(1)}, S_{m-1}^{(2)})$, by first conditioning on $\{S_t^{(i)} : i = 1, 2, t = 0, \dots, m\}$ and by using the local limit theorem (see (31)), we see that for fixed $m \geq 1$, as $k \rightarrow \infty$,

$$k^{d/2} f_k(s^{(1)}, s^{(2)}) \rightarrow \mathfrak{C}_d P^{\otimes 2}(e^{\lambda_2 N_{0,m+1}})$$

with

$$\mathfrak{C}_d = 2(d/4\pi)^{d/2}. \quad (63)$$

It follows that for fixed $m \geq 1$, as $k \rightarrow \infty$,

$$\begin{aligned} k^{d/2} P^{\otimes 2}(\mathbf{1}_{\{S_k^{(1)} = S_k^{(2)}\}} e^{\lambda_2 N_{k-m,k}(S^{(1)}, S^{(2)})} | \mathcal{S}^{(1,2)}) &= k^{d/2} f_{k-m+1}(S_{m-1}^{(1)}, S_{m-1}^{(2)}) \\ &\rightarrow \mathfrak{C}_d P^{\otimes 2}(e^{\lambda_2 N_{0,m+1}}), \end{aligned} \quad (64)$$

and the convergence is uniform over all possible values of $(S_{m-1}^{(1)}, S_{m-1}^{(2)})$ since the number of possible values is finite. Therefore, for each $\varepsilon > 0$, there is $k_\varepsilon > 0$ such that for all $k \geq k_\varepsilon$, $k^{d/2} f_{k-m+1}(S_{m-1}^{(1)}, S_{m-1}^{(2)})$ lies between $\mathfrak{C}_d P^{\otimes 2}(e^{\lambda_2 N_{0,m+1}}) \pm \varepsilon$. Thus, for fixed $m \geq 1$, as $k \rightarrow \infty$, it holds

$$\begin{aligned} k^{\frac{d}{2}} P^{\otimes 4} \left[e^{\sum_{0,m} P^{\otimes 2}(\mathbf{1}_{\{S_k^{(1)} = S_k^{(2)}\}} e^{\lambda_2 N_{k-m,k}} | \mathcal{S}^{(1,2)})} \right] &= k^{\frac{d}{2}} P^{\otimes 4} \left[e^{\sum_{0,m} f_{k-m+1}(S_{m-1}^{(1)}, S_{m-1}^{(2)})} \right] \\ &\rightarrow e^{-\lambda_2} \mathfrak{C}_d P^{\otimes 4} \left[e^{\sum_{0,m}} \right] P^{\otimes 2}(e^{\lambda_2 N_{0,m+1}}). \end{aligned} \quad (65)$$

It follows that the sum in (62) can be estimated, for $n \rightarrow \infty$, by

$$\begin{aligned} n^{d-2} \sum_{k \geq n} \text{of (62)} &\sim e^{-\lambda_2} P^{\otimes 4} \left[e^{\sum_{0,m}} \right] \mathfrak{C}_d P^{\otimes 2}(e^{\lambda_2 N_{0,m+1}}) n^{d-2} \sum_{k \geq n} \frac{1}{k^{d-1}} \\ &\rightarrow e^{-\lambda_2} \mathfrak{C}_d P^{\otimes 4} \left[e^{\sum_{0,m}} \right] P^{\otimes 2}(e^{\lambda_2 N_{0,m+1}}) \frac{1}{d-2}. \end{aligned}$$

Collecting all this we conclude that the limit $\lim_{n \rightarrow \infty} T_2^<(m, n)$ exists, and further, that the limits $\lim_{m \rightarrow \infty} \lim_{n \rightarrow \infty} T_2^<(m, n)$ and $\lim_{n \rightarrow \infty} T^<(n)$ exist and are equal. Finally,

$$\begin{aligned} \lim_{n \rightarrow \infty} \mathbb{E} s_n^4 &= 2 \lim_{n \rightarrow \infty} T^<(n) \\ &\stackrel{(62)}{=} 2 \frac{\mathfrak{K}_d \mathfrak{J}_d \mathfrak{C}_d (1 - \pi_d)}{d - 2} \kappa_2 e^{-\lambda_2} \times \mathbb{E}(W^4) \mathbb{E}(W^2) \text{Var}(W) \\ &= \sigma_1^4 \mathbb{E}(W^4). \end{aligned} \tag{66}$$

This concludes the proof of (50).

We then give the **proof of (51)**. We estimate the cross term

$$\mathbb{E} s_n^2 W_n^2 = \kappa_2 n^{\frac{d-2}{2}} \sum_{k \geq n} P^{\otimes 4} \left[e^{\Sigma_{0,n} + \lambda_2 N_{n,k}(S^{(1)}, S^{(2)})} \mathbf{1}_{S_k^{(1)} = S_k^{(2)}} \right]$$

Using the identity $\mathbf{1}_{S_k^{(1)} = S_k^{(2)}} = \frac{e^{\lambda_2 N_{k,k+1}(S^{(1)}, S^{(2)})} - 1}{\kappa_2}$ (see (57)), we obtain that

$$\begin{aligned} \mathbb{E} s_n^2 W_n^2 &= n^{\frac{d-2}{2}} P^{\otimes 4} \left[e^{\Sigma_{0,n}} \sum_{k \geq n} \left(e^{\lambda_2 N_{n,k+1}(S^{(1)}, S^{(2)})} - e^{\lambda_2 N_{n,k+1}(S^{(1)}, S^{(2)})} \right) \right] \\ &= n^{\frac{d-2}{2}} P^{\otimes 4} \left[e^{\Sigma_{0,n}} (e^{\lambda_2 N_{n,\infty}(S^{(1)}, S^{(2)})} - 1) \right] \end{aligned}$$

(telescopic sum). By (58) and the argument preceding it, we know that

$$\mathbb{E} \left(e^{\lambda_2 N_{n,\infty}(S^{(1)}, S^{(2)})} - 1 \mid S_t^{(i)}, i = 1, 2, t = 0, \dots, n \right) = \frac{G(S_n^{(1)} - S_n^{(2)})}{G(0)} (\mathbb{E}(W^2) - 1).$$

Therefore, taking first the conditional expectation given $\{S_t^{(i)}, i = 1, 2, t = 0, \dots, n\}$ in the preceding expression of $\mathbb{E} s_n^2 W_n^2$, we get

$$\begin{aligned} \mathbb{E} s_n^2 W_n^2 &= \frac{\mathbb{E}(W^2) - 1}{G(0)} n^{\frac{d-2}{2}} P^{\otimes 4} \left[e^{\Sigma_{0,n}} G(S_n^{(1)} - S_n^{(2)}) \right] \\ &\rightarrow \frac{\mathbb{E}(W^2) - 1}{G(0)} \mathfrak{K}_d \mathfrak{J}_d P^{\otimes 4} e^{\Sigma_{0,\infty}} = \sigma_1^2 \mathbb{E} W^4, \end{aligned}$$

where the last convergence can be proved as follows. Indeed, it can be easily seen that a similar result like Lemma 2.2 holds for the asymptotic independence of $\Sigma_{0,n}$ and $n^{-1/2} \tilde{S}_n^{(i)}, 1 \leq i \leq 4$: under $P^{\otimes 4}$, we have the following convergence of random vectors :

$$\left(\Sigma_{0,n}, n^{-1/2} S_n^{(1)}, n^{-1/2} \tilde{S}_n^{(2)}, n^{-1/2} \tilde{S}_n^{(3)}, n^{-1/2} \tilde{S}_n^{(4)} \right) \longrightarrow (\Sigma_{0,\infty}, Z_1, Z_2, Z_3, Z_4) \quad \text{in distribution,}$$

where $\Sigma_{0,\infty}, Z_1, Z_2, Z_3, Z_4$ are all independent, and each Z_i is a Gaussian vector with mean 0 and covariance matrix $\frac{1}{d} I_d$. This can be proved in a similar way as in the proof of Lemma 2.2, and can also be derived by a general result on the mixing convergence as indicated in Remark 2.4. Also, the assertions (25) and (26) still hold with (N_n, S_n, \tilde{S}_n) replaced by $(\Sigma_{0,n}, S_n^{(1)}, \tilde{S}_n^{(2)})$, yielding that the sequence $\{n^{\frac{d-2}{2}} e^{\Sigma_{0,n}} G(S_n^{(1)} - S_n^{(2)})\}$ converges in law and in mean to $e^{\Sigma_{0,\infty}} \times \frac{\mathfrak{K}_d}{|Z|^{d-2}}$. This gives the desired conclusion on $\mathbb{E} s_n^2 W_n^2$, which ends the proof of (51). ■

We now give the

Proof of Lemma 4.1. For the first limit, we put $q = (d + \epsilon)/(d - 1 + \epsilon)$ and we estimate

$$\begin{aligned}
T^=(n) &= \kappa_2^2 n^{d-2} \sum_{k \geq n} P^{\otimes 4} \left[e^{\Sigma_{0,k}} \mathbf{1}_{S_k^{(1)}=S_k^{(2)}} \mathbf{1}_{S_k^{(3)}=S_k^{(4)}} \right] \\
&\stackrel{\text{H\"older}}{\leq} \kappa_2^2 n^{d-2} \sum_{k \geq n} P^{\otimes 4} \left[e^{(d+\epsilon)\Sigma_{0,k}} \right]^{1/(d+\epsilon)} P^{\otimes 2} [S_k^{(1)} = S_k^{(2)}]^{2/q} \\
&\leq C n^{d-2} \sum_{k \geq n} k^{-d/q}
\end{aligned}$$

with $C \in (0, \infty)$ a constant, by the local limit theorem. Being of order $n^{-\epsilon/(d+\epsilon)}$, the last term vanishes as $n \rightarrow \infty$.

We now prove the second limit, using arguments which are similar to the ones above. Since the difference is non-negative, the term $|T_1^<(n) - T_2^<(m, n)|$ is equal to

$$\begin{aligned}
&C n^{d-2} \sum_{k \geq n} P^{\otimes 4} \left[e^{\Sigma_{0,m} + \lambda_2 N_{k-m,k}(S^{(1)}, S^{(2)})} \left\{ e^{\Sigma_{m,k} - \lambda_2 N_{k-m,k}(S^{(1)}, S^{(2)})} - 1 \right\} \mathbf{1}_{S_k^{(1)}=S_k^{(2)}} G(S_k^{(3)} - S_k^{(4)}) \right] \\
= &C n^{d-2} \sum_{k \geq n} P^{\otimes 4} \left[e^{\Sigma_{0,m}} \mathbf{1}_{S_k^{(1)}=S_k^{(2)}} G(S_k^{(3)} - S_k^{(4)}) \right. \\
&\left. P^{\otimes 4} \left\{ e^{\lambda_2 N_{k-m,k}(S^{(1)}, S^{(2)})} \left(e^{\Sigma_{m,k} - \lambda_2 N_{k-m,k}(S^{(1)}, S^{(2)})} - 1 \right) \mid S_t^{(i)}, 1 \leq i \leq 4, t = m, k \right\} \right] \quad (67)
\end{aligned}$$

with $C = \frac{\text{Var}(W)}{G(0)} \kappa_2$, by first conditioning on the paths at times $t = 1, \dots, m$ and $t = k$. Indeed, in (67) the internal conditional expectation given the paths at times $t = m, k$ is the same as the conditional expectation given the paths at times $t = 1, \dots, m, k$, as shown by the following fact : for any positive function f , writing $S_t = (S_t^{(i)}, 1 \leq i \leq 4)$, we have, for $1 \leq m < k$ and for any $s_t = (s_t^{(i)}, 1 \leq i \leq 4)$ with $s_t^{(i)} \in \mathbb{Z}^d$,

$$\begin{aligned}
&P^{\otimes 4} (f(S_t, m < t \leq k) \mid S_t = s_t, t = 1, \dots, m, k) \\
= &P^{\otimes 4} (f(S_t, m < t \leq k) \mid S_t = s_t, t = m, k) \\
= &P^{\otimes 4} (f(S_t + s_m, 0 < t \leq k - m) \mid S_{k-m} = s_k - s_m).
\end{aligned}$$

This can be easily checked by the definition of the conditional expectation, using the fact that : a) when $S_m = s_m$, then $S_k = s_k$ if and only if $S_k - S_m = s_k - s_m$, b) on the event $\{S_m = s_m\}$, we have $S_t = S_t - S_m + s_m$ for $m < t \leq k$, c) the process $\{S_t - S_m : m < t \leq k\}$ is independent of $\{S_t : t = 1, \dots, m\}$ and has the same law as $\{S_t : 0 < t \leq k - m\}$.

The event

$$B_{m,k} = \left\{ |S_t^{(i)} - S_t^{(j)}| \geq m^{1/4}, \text{ for } i \neq j \text{ and } t = m, \text{ and for } \{i, j\} \neq \{1, 2\} \text{ and } t = k \right\}$$

(where $i, j = 1, \dots, 4$, and for $\{i, j\} \neq \{1, 2\}$ we include additionally the terms with $t = k$) has probability larger than $1 - cm^{-d/4}$ for some constant $c \in (0, \infty)$, by the local limit theorem. On this event, by transience, the intersections between the paths between times m and $k - m$ essentially come, when m and k are large, from those of $S^{(1)}$ and $S^{(2)}$ between times $k - m$ and m . Precisely,

$$\lim_{m \rightarrow \infty} \limsup_{k \rightarrow \infty} \sup_{B_{m,k}} P^{\otimes 4} \left\{ \Sigma_{m,k} \neq \lambda_2 N_{k-m,k}(S^{(1)}, S^{(2)}) \mid S_t^{(i)}, 1 \leq i \leq 4, t = m, k \right\} = 0.$$

(Note that the event $\Sigma_{m,k} = \lambda_2 N_{m-k,k}(S^{(1)}, S^{(2)})$, the contrary of that in the previous line, means that there are no intersections of $S^{(i)}$ and $S^{(j)}$ between times m and k for $i \geq 3$ or $j \geq 3$, and no intersections of $S^{(1)}$ and $S^{(2)}$ between times m and $k - m$.) To see this, recall that, for even $x \in \mathbb{Z}^d$, $P_{0,x}^{\otimes 2}(N_{0,\infty} \geq 1) = G(x)/G(0) = O(|x|^{2-d})$. Thus,

$$\begin{aligned} & \sup_{B_{m,k}} P^{\otimes 4} \left\{ \Sigma_{m,k} \neq \lambda_2 N_{m-k,k}(S^{(1)}, S^{(2)}) \mid S_t^{(i)}, 1 \leq i \leq 4, t = m, k \right\} \\ & \leq \sup_{B_{m,k}} P^{\otimes 4} \left\{ \exists (i, j) \neq (1, 2) : N_{k-m,k}(S^{(i)}, S^{(j)}) \geq 1 \mid S_t^{(i)}, 1 \leq i \leq 4, t = m, k \right\} \\ & \leq \sum_{(i,j) \neq (1,2)} \sup_{B_{m,k}} P^{\otimes 4} \left\{ N_{k-m,k}(S^{(i)}, S^{(j)}) \geq 1 \mid S_t^{(i)}, 1 \leq i \leq 4, t = m, k \right\} \\ & = O(|m|^{-(d-2)/4}), \end{aligned}$$

which implies our claim. (Here, to get the last line, we use time reversal to bound the conditional expectation in the third line by the Green function evaluated at $S_k^{(i)} - S_k^{(j)}$, together with the Green function estimate and the fact that $|S_k^{(i)} - S_k^{(j)}| \geq m^{1/4}$ on $B_{m,k}$.) With the integrability condition (61) and the inequality of Cauchy - Schwarz, this is enough to imply that

$$\lim_{m \rightarrow \infty} \limsup_{k \rightarrow \infty} \sup_{B_{m,k}} P^{\otimes 4} \left\{ e^{\lambda_2 N_{k-m,k}(S^{(1)}, S^{(2)})} (e^{\Sigma_{m,k} - \lambda_2 N_{k-m,k}(S^{(1)}, S^{(2)})} - 1) \mid S_t^{(i)}, 1 \leq i \leq 4, t = m, k \right\} = 0.$$

Plugging this in (67) and using the fact that $n^{d-2} \sum_{k \geq n} P^{\otimes 4} [e^{\Sigma_{0,m} + \lambda_2 N_{k-m,k}(S^{(1)}, S^{(2)})} \mathbf{1}_{S_k^{(1)} = S_k^{(2)}}]$ has a finite limit, we conclude that the right-hand side of (67) vanishes as $n \rightarrow \infty$. \blacksquare

Proof of Lemma 3.4. With the notation introduced in the beginning of this section, we have $D_{k+1} = W_{k+1} - W_k = P[e^{h_k(S)}(e^{\beta \eta(k, S_k) - \lambda(\beta)} - 1)]$,

$$D_{k+1}^4 = P^{\otimes 4} \left[e^{\sum_{i=1}^4 h_k(S^{(i)})} \prod_{i=1}^4 (e^{\beta \eta(k, S_k^{(i)}) - \lambda(\beta)} - 1) \right].$$

Using Fubini's theorem and independence, we obtain

$$\begin{aligned} \mathbb{E} D_{k+1}^4 &= P^{\otimes 4} \left[\mathbb{E} e^{\sum_{i=1}^4 h_k(S^{(i)})} \mathbb{E} \prod_{i=1}^4 (e^{\beta \eta(k, S_k^{(i)}) - \lambda(\beta)} - 1) \right] \\ &= P^{\otimes 4} e^{\Sigma_{0,k}} (\gamma_4 \mathbf{1}_{(S_k^{(1)}, S_k^{(2)}, S_k^{(3)}, S_k^{(4)}) \in E_4} + \gamma_2 \mathbf{1}_{(S_k^{(1)}, S_k^{(2)}, S_k^{(3)}, S_k^{(4)}) \in E_{2,2}}), \end{aligned} \quad (68)$$

where $\gamma_4 = \mathbb{E}(e^{\beta \eta(0,0) - \lambda(\beta)} - 1)^4$ and $\gamma_2 = (\mathbb{E}(e^{\beta \eta(0,0) - \lambda(\beta)} - 1)^2)^2$. Notice that $(S_k^{(1)}, S_k^{(2)}, S_k^{(3)}, S_k^{(4)}) \in E_{2,2}$ if and only if one of the following cases occurs : (a) $S_k^{(1)} = S_k^{(2)} \neq S_k^{(3)} = S_k^{(4)}$, (b) $S_k^{(1)} = S_k^{(3)} \neq S_k^{(2)} = S_k^{(4)}$, (c) $S_k^{(1)} = S_k^{(4)} \neq S_k^{(2)} = S_k^{(3)}$. Therefore by symmetry, we obtain

$$\mathbb{E} D_{k+1}^4 \leq (\gamma_4 + 3\gamma_2) P^{\otimes 4} e^{\Sigma_{0,k}} \mathbf{1}_{S_k^{(1)} = S_k^{(2)}} \mathbf{1}_{S_k^{(3)} = S_k^{(4)}}.$$

Hence, using Hölder's inequality, for $p, q > 1$ with $1/p + 1/q = 1$, we have

$$\mathbb{E} D_{k+1}^4 \leq (\gamma_4 + 3\gamma_2) (P^{\otimes 4} e^{p\Sigma_{0,k}})^{1/p} (P^{\otimes 2}(S_k^{(1)} = S_k^{(2)}))^{2/q}.$$

By the local limit theorem, $P^{\otimes 2}(S_k^{(1)} = S_k^{(2)}) = O(k^{-d/2})$. Thus taking $|\beta| > 0$ small enough such that $P^{\otimes 4}e^{p\Sigma_{0,\infty}} < \infty$, we see that Eq. (52) holds. This ends the proof of Lemma 3.4. ■

Acknowledgements. The authors would like to express their gratitude to the referee for his careful reading and valuable comments which led to an improvement of the paper. The work has been supported by CNRS (UMR 7599 Probabilités et Modèles Aléatoires et UMR 6205 Laboratoire de Mathématique de Bretagne Atlantique); in particular Q. Liu has benefited from a delegation CNRS during the preparation of the paper. It has also been partially supported by the U.S. National Science Foundation (Grant no. NSF PHY11-25915) and the National Natural Science Foundation of China (Grants no. 11571052 and no. 11401590).

Références

- [1] ALBERTS, T. ; ORTGIESE, M. (2013). The near-critical scaling window for directed polymers on disordered trees. *Electron. J. Probab.* **18** no. 19.
- [2] ALBERTS, T. ; KHANIN, K. ; QUASTEL, J. (2014) The intermediate disorder regime for directed polymers in dimension 1+1. *Ann. Probab.* **42**, 1212-1256.
- [3] ALDOUS, D. ; EAGLESON, G. (1978). On mixing and stability of limit theorems. *Ann. Probab.* **6** 325–331.
- [4] BIRKNER, M. (2004) A condition for weak disorder for directed polymers in random environment. *Electron. Comm. Probab.* **9** 22-25.
- [5] BOLTHAUSEN, E. (1989). A note on diffusion of directed polymers in a random environment, *Commun. Math. Phys.* **123**, 529–534.
- [6] CAMANES, A. ; CARMONA, P. (2009) The critical temperature of a directed polymer in a random environment *Markov Process. Related Fields* **15** 105–116.
- [7] CARAVENNA, F. ; SUN, R. ; ZYGOURAS, N. Polynomial chaos and scaling limits of disordered systems. [arXiv: 1312.3357.pdf](#), to appear, *J. Eur. Math. Soc.*
- [8] COMETS, F. ; POPOV, S. ; VACHKOVSKAIA, M. (2008). The number of open paths in an oriented ρ -percolation model. *J. Stat. Phys.* **131** 357–379.
- [9] COMETS, F. ; VARGAS, V. (2006). Majorizing multiplicative cascades for directed polymers in random media. *ALEA Lat. Am. J. Probab. Math. Stat.* **2** 267–277.
- [10] COMETS, F. ; YOSHIDA, N. Directed polymers in random environment are diffusive at weak disorder. *Ann. Probab.* **34** 1746-1770.
- [11] DEAN, D. ; MAJUMDAR, S. (2001). Extreme-value statistics of hierarchically correlated variables deviation from Gumbel statistics and anomalous persistence. *Phys. Rev. E* **64**, 046121.
- [12] DERRIDA, B. ; SPOHN, H. (1988). Polymers on disordered trees, spin glasses, and traveling waves. *J. Statist. Phys.* **51** 817–840.
- [13] GIACOMIN, G. (2007). *Random polymer models*. Imperial College Press.
- [14] HALL, P. ; HEYDE, C.C. (1980). *Martingale limit theory and its application*. Academic press.
- [15] HALPIN-HEALY, T. ; TAKEUCHI, K. (2015). A KPZ cocktail – shaken, not stirred ... Toasting 30 years of kinetically roughened surfaces. *J. Statist. Phys.* **160** 794–814.
- [16] HEYDE, C. (1970). A rate of convergence result for the super-critical Galton-Watson process. *J. Appl. Probab.* **7** 451–454
- [17] HEYDE, C. (1971). Some central limit analogues for supercritical Galton-Watson processes. *J. Appl. Probab.* **8** 52-59

- [18] HU, Y. ; SHAO, Q.-M. A note on directed polymers in Gaussian environments. *Electron. Commun. Probab.* **14** 518-528.
- [19] HUANG, C. ; LIU, Q. (2014). Convergence rates for a supercritical branching process in a random environment. *Markov Processes and Related Fields*, **20** 265-285.
- [20] IKSANOV, O. (2006). On the rate of convergence of a regular martingale related to a branching random walk. *Ukrainian Math. J.* **58** 368–387
- [21] IKSANOV, O. ; KABLUCHKO, Z. (2016). A central limit theorem and a law of the iterated logarithm for the Biggins martingale of the supercritical branching random walk. *Adv. Appl. Prob.*, to appear. <http://arxiv.org/pdf/1507.08458v2.pdf>
- [22] IMBRIE, J.Z. ; SPENCER, T. (1988). Diffusion of directed polymer in a random environment, *J. Stat. Phys.* **52** 609-626.
- [23] LACOIN, H. (2010) New bounds for the free energy of directed polymers in dimension 1+1 and 1+2. *Comm. Math. Phys.* **294** 471–503
- [24] LAWLER, G. (1991). *Intersections of Random Walks*. Birkhäuser Boston.
- [25] LAWLER, G. ; LIMIC, V. (2010). *Random Walk : A Modern Introduction*. Cambridge University Press.
- [26] LIU, Q. ; WATBLED, F. (2009). Exponential inequalities for martingales and asymptotic properties of the free energy of directed polymers in random environment. *Stoch. Proc. Appl.* **119** 3101-3132.
- [27] NGUYEN, V.-L. (2016). Stochastic order and disordered systems. Preprint
- [28] RÉNYI, A. (1963). On stable sequences of events. *Sankhyā Ser. A* **25** 293–302.
- [29] SINAI, Y. (1995). A remark concerning random walks with random potentials *Fund. Math.* **147**, 173–180.
- [30] VARGAS, V. (2006). A Local limit theorem for directed polymers in random media : the continuous and the discrete case. *Ann. Inst. H. Poincaré Probab. Statist.* **42** 521–534.
- [31] WANG, H. ; GAO, Z. ; LIU, Q. (2011). Central limit theorems for a supercritical branching process in a random environment. *Statist. Probab. Lett.* **81** 539–547.