



HAL
open science

NON-WANDERING FATOU COMPONENTS FOR STRONGLY ATTRACTING POLYNOMIAL SKEW PRODUCTS

Zhuchao Ji

► **To cite this version:**

Zhuchao Ji. NON-WANDERING FATOU COMPONENTS FOR STRONGLY ATTRACTING POLYNOMIAL SKEW PRODUCTS. 2018. hal-01710467v1

HAL Id: hal-01710467

<https://hal.science/hal-01710467v1>

Preprint submitted on 16 Feb 2018 (v1), last revised 11 Dec 2018 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

NON-WANDERING FATOU COMPONENTS FOR STRONGLY ATTRACTING POLYNOMIAL SKEW PRODUCTS

ZHUCHAO JI

ABSTRACT. We show a partial generalization of Sullivan’s non-wandering domain theorem in complex dimension two. More precisely, we show the non-existence of wandering Fatou components for polynomial skew products in \mathbb{C}^2 with an invariant attracting fiber, under the assumption that the multiplier λ is small. We actually show a stronger result, namely that every forward orbit of any vertical Fatou disk intersects a fattened Fatou component.

CONTENTS

1. Introduction	1
2. Preliminaries	4
2.1. Local attracting polynomial skew products	4
2.2. Denker-Przytycki-Urbanski’s Lemma	5
3. Structure of fattened Fatou components	6
3.1. Attracting basin case	6
3.2. Parabolic basin case	7
3.3. Siegel disk case	8
3.4. Wandering vertical Fatou disks	9
4. Estimate of horizontal size of fattened Fatou components	9
5. Estimate of size of forward images of vertical Fatou disks	15
6. Proof of the non-wandering domain theorem	21
7. Polynomial skew products	23
References	24

1. INTRODUCTION

Complex dynamics, also known as Fatou-Julia theory, is naturally subdivided according to these two terms. One is focused on the Julia set. This is the set where chaotic dynamics occurs. The other direction of investigation is concerned with the dynamically stable part - the Fatou set. In this paper we will concentrate on the Fatou theory.

In a general setting, let M be a complex manifold, and let $f : M \rightarrow M$ be a holomorphic self map. We consider f as a dynamical system, that is, we study the long-time behavior of the sequence of iterates $\{f^n\}_{n \geq 0}$. The **Fatou set** $F(f)$ is classically defined as the largest

Date: February 16, 2018.

2000 Mathematics Subject Classification. 37F10, 37F50, 32H50.

Key words and phrases. polynomial skew product, Fatou component, non wandering domain theorem.

open subset of M in which the sequence of iterates is normal. Its complement is the **Julia set** $J(f)$. A **Fatou component** is a connected component of $F(f)$.

In one-dimensional case, we study the dynamics of iterated holomorphic self map on a Riemann surface. The classical case of rational functions on Riemann sphere \mathbb{P}^1 occupies an important place and produces a fruitful theory. The Non-Wandering Domain Theorem due to Sullivan [8] asserts that every Fatou component of a rational map is eventually periodic. This result is fundamental in the Fatou theory since it leads to a complete classification of the dynamics in the Fatou set: the orbit of any point in the Fatou set eventually lands in an attracting basin, a parabolic basin, or a rotation domain.

The same question arises in higher dimensions, i.e. to investigate the non-wandering domain theorem for higher dimensional holomorphic endomorphisms on \mathbb{P}^k . A good test class is that of polynomial skew products hence one-dimensional tools can be used.

A polynomial skew product is a map $P : \mathbb{C}^2 \rightarrow \mathbb{C}^2$ of the form

$$P(t, z) = (g(t), f(t, z)),$$

where g is a one variable polynomial and f is a two variable polynomial. See Jonsson [3] for a systematic study of such polynomial skew products.

The first positive result is due to Lilov. Let us assume that $g(0) = 0$ which means that $\{t = 0\}$ is an invariant fiber. Assume that f can be written as a polynomial in z ,

$$f(t, z) = a_0(t) + a_1(t)z + \cdots + a_d(t)z^d,$$

with coefficients $a_i(t)$ holomorphic in t in a neighborhood of 0, and $a_d(0) \neq 0$. In this case the dynamics in $\{t = 0\}$ is given by the polynomial

$$p(z) = f(0, z)$$

and is very well understood. P is called a local polynomial skew product. Under the assumption that $0 \leq |g'(0)| < 1$, Koenigs' Theorem and Böttcher's Theorem tell us that the dynamical system is locally conjugate to

$$P(t, z) = (\lambda t, f(t, z)),$$

when $g'(0) = \lambda \neq 0$, or

$$P(t, z) = (t^m, f(t, z)), \quad m \geq 2,$$

when $g'(0) = 0$. In the first case the invariant fiber is called attracting and in the second case the invariant fiber is called super-attracting. In his unpublished PhD thesis [4], Lilov first showed that every Fatou component of p in the super-attracting invariant fiber is contained in a two-dimensional Fatou component, which is called a fattened Fatou component. We will show that this bulging property of Fatou component of p also holds in attracting case.

Lilov's main result is the non-existence of wandering Fatou components for local polynomial skew products in the basin of a super-attracting invariant fiber. Since this is a local result, it can be stated as follows.

Theorem (Lilov). *For a local polynomial skew product P with a super-attracting invariant fiber,*

$$P(t, z) = (t^m, f(t, z)), \quad m \geq 2,$$

every forward orbit of a vertical Fatou disk intersects a fattened Fatou component. This implies that every Fatou component iterates to a fattened Fatou component. In particular there are no wandering Fatou components.

On the other hand, recently Astorg, Buff, Dujardin, Peters and Raissy [1] constructed a holomorphic endomorphism $h : \mathbb{P}^2 \rightarrow \mathbb{P}^2$, induced by a polynomial skew product $P = (g(t), f(t, z)) : \mathbb{C}^2 \rightarrow \mathbb{C}^2$ with parabolic invariant fiber, possessing a wandering Fatou component, here parabolic invariant fiber means that $g(0) = 0$ and $g'(0) = e^{i\pi p/q}$ with p/q a rational number. Thus the non-wandering domain theorem does not hold for general polynomial skew products.

At this stage it remains an interesting problem to investigate the existence of wandering Fatou components for local polynomial skew products with attracting but not super-attracting invariant fiber. As it is clear from Lilov's theorem, Lilov actually showed a stronger result, namely that every forward orbit of a vertical Fatou disk intersects a fattened Fatou component. Peters and Vivas showed in [6] that there is an attracting local polynomial skew product with a wandering vertical Fatou disk, which shows that Lilov's proof breaks down in the general attracting case. Note that this result does not answer the existence question of wandering Fatou components, but shows that the question is considerably more complicated than in the super-attracting case. On the other hand, by using a different strategy from Lilov's, Peters and Smit in [5] showed that the non-wandering domain theorem holds in the attracting case, under the assumption that the dynamics on the invariant fiber is sub-hyperbolic.

In this paper we prove a non-wandering domain theorem in the attracting local polynomial skew product case without any assumption on the dynamics on the invariant fiber. Actually we show that Lilov's stronger result holds in the attracting case when the multiplier λ is small.

Theorem (Main Theorem). *For a local polynomial skew product P with an attracting invariant fiber,*

$$P(t, z) = (\lambda t, f(t, z)),$$

for any fixed f , there is a constant $\lambda_0 = \lambda_0(f) > 0$ such that if λ satisfies $0 < |\lambda| < \lambda_0$, every Fatou component iterates to a fattened Fatou component. In particular there are no wandering Fatou components.

We can also apply this local result to globally defined polynomial skew products, see Theorem 7.1 for a precise statement..

The proof of the main theorem basically follows Lilov's strategy. The difficulty is that Lilov's argument highly depends on the super-attracting condition and breaks down in the attracting case by [6]. The main idea of this paper is to use and adapt a one-dimensional lemma due to Denker-Przytycki-Urbanski (the DPU Lemma for short) to our case. This will give estimates of the horizontal size of fattened Fatou components and of the size of forward images of a wandering vertical Fatou disk (these concepts will be explained later). We note that some results in our paper already appear in Lilov's thesis [4] (Theorem 3.5, Lemma 4.1, Lemma 5.1 and Lemma 5.2). Since his paper is not easily available, we choose to present the whole proof with all details. On the other hand we believe that the introduction of the DPU Lemma makes the argument conceptually simpler even in the super-attracting case.

The outline of the paper is as follows. In section 2 we start with some definitions, then we present the DPU Lemma and some corollaries. In section 3 we show that every Fatou component of p in the invariant fiber is bulged, i.e. contained in a two-dimensional fattened Fatou component. This result follows classical ideas from normal hyperbolicity theory.

In section 4 we give an estimate of the horizontal size of the fattened Fatou components by applying the one-dimensional DPU Lemma. Let $z \in F(p)$ be a point in a Fatou component of

the invariant fiber and denote by $r(z)$ the supremum radius of a horizontal holomorphic disk (see Definition 2.4 for the precise definition) centered at z that is contained in the fattened Fatou component. We have the following key estimate.

Theorem 4.4. *If λ is chosen sufficiently small, then there are constants $k > 0, l > 0, R > 0$ such that for any point $z \in F(p) \cap \{|z| < R\}$,*

$$r(z) \geq k d(z, J(p))^l,$$

where $J(p)$ is the Julia set in the invariant fiber.

In section 5 we adapt the DPU Lemma to the attracting local polynomial skew product case, to show that the size of forward images of a wandering vertical Fatou disk shrinks slowly, which is also important in the proof of the main theorem.

Proposition 5.5. *Let $\Delta_0 \subset \{t = t_0\}$ be a wandering vertical Fatou disk centered at $x_0 = (t_0, z_0)$. Let $x_n = (t_n, z_n) = P^n(x_0)$. Define a function ρ as follows: for a domain $U \subset \mathbb{C}$, for every $z \in U \subset \mathbb{C}$, define*

$$\rho(z, U) = \sup \{r > 0 \mid D(z, r) \subset U\}.$$

Set $\Delta_n = P^n(\Delta_0)$ for every $n \geq 1$ and let $\rho_n = \rho(z_n, \pi(\Delta_n))$, here π is the projection $\pi : (t, z) \mapsto z$. If λ is chosen sufficiently small, we have

$$\lim_{n \rightarrow \infty} \frac{|\lambda|^n}{\rho_n} = 0.$$

The proof of the main theorem is given in section 6. The main point is to combine Theorem 4.4 and Proposition 5.5 to show wandering vertical Fatou disk can not exist. We finish section 6 with some remarks around the main theorem. We also show how our main theorem can be applied to globally defined polynomial skew products in section 7.

Acknowledgements. I would like to thank Romain Dujardin for drawing my attention to the subject and for his invaluable help. The work is partially supported by ANR-LAMBDA, ANR-13-BS01-0002.

2. PRELIMINARIES

2.1. Local attracting polynomial skew products. In this subsection we make some definitions around local attracting polynomial skew products.

Definition 2.1. *A local attracting polynomial skew product is a map $P : \Delta \times \mathbb{C} \rightarrow \Delta \times \mathbb{C}$, here $\Delta \subset \mathbb{C}$ is a disk centered at 0, such that*

$$P(t, z) = (\lambda t, f(t, z)),$$

here f is a polynomial in z ,

$$f(t, z) = a_0(t) + a_1(t)z + \cdots + a_d(t)z^d$$

with coefficients $a_i(t)$ holomorphic in Δ , and $a_d(0) \neq 0$. We denote by $p(z) = f(0, z)$ the self map of the invariant fiber $\{t = 0\}$.

Definition 2.2. Let π denote the projection to the z -axis, that is

$$\begin{aligned} \pi : \Delta \times \mathbb{C} &\longrightarrow \mathbb{C}, \\ (t, z) &\longmapsto z. \end{aligned}$$

A subset Δ_0 lying in some $\{t = t_0\}$ is called a **vertical disk** if $\pi(\Delta_0)$ is a disk in the complex plane. A vertical disk centered at x_0 with radius r is denoted by $\Delta(x_0, r)$. Δ_0 is called a **vertical Fatou disk** if the restriction of $\{P^n\}_{n \geq 0}$ to Δ_0 is a normal family.

We remark that for a disk on the complex plane centered at z with radius r , we denote it by $D(z, r)$ to distinguish.

Proposition 2.3. A vertical disk contained in a Fatou component of P is a vertical Fatou disk.

Definition 2.4. A **horizontal holomorphic disk** is a subset of the form

$$\{(t, z) \in \Delta \times \mathbb{C} \mid z = \phi(t), |t| < \delta\}$$

where $\phi(t)$ is holomorphic in $\{|t| < \delta\}$ for some $\delta > 0$. δ is called the **size** of the horizontal holomorphic disk.

Example: Let

$$P(t, z) = (g(t), f(t, z)) : \mathbb{C}^2 \rightarrow \mathbb{C}^2$$

be a globally defined polynomial skew product, where g, f are polynomials. Assume that g has an attracting fixed point, without loss of generality we assume that it is 0. When working in a small neighborhood of $\{t = 0\}$ we can change coordinates so that $g(t) = \lambda t$. Now $f(t, z)$ is no longer a polynomial in t and z but a polynomial in z with coefficients holomorphic functions in t . Thus in this local coordinate P is a local attracting polynomial skew product.

2.2. Denker-Przytycki-Urbanski's Lemma. In this subsection we introduce the work of Denker-Przytycki-Urbanski in [2], and give some corollaries. Denker, Przytycki and Urbanski consider rational maps on \mathbb{P}^1 , and study the local dynamical behavior of some neighborhood of a critical point lying in Julia set. As a consequence they deduce an upper bound of the size of the pre-images of a ball centered at a point in Julia set.

In the following let f be a rational map on \mathbb{P}^1 , denote by $C(f)$ the set of critical points of f lying in Julia set. Assume that $\#C(f) = q$. We begin with a definition,

Definition 2.5. For a critical point $c \in C(f)$, define a positive valued function $k_c(x)$ by

$$k_c(x) = \begin{cases} -\log d(x, c), & \text{if } x \neq c \\ \infty, & \text{if } x = c. \end{cases}$$

Define a function $k(x)$ by

$$k(x) = \max_{c \in C(f)} k_c(x).$$

Here the distance is relative to the spherical metric on \mathbb{P}^1 .

Let x_0 be arbitrary and consider the forward orbit $\{x_0, x_1, \dots, x_n, \dots\}$, where $x_n = f^n(x_0)$. We let the function $k(x)$ act on this orbit and the following DPU Lemma gives an asymptotic description of the sum of $k(x)$ on this orbit. Recall that q denote the number of critical points lying in J .

Lemma 2.6 (Denker, Przytycki, Urbanski). *There exist a constant $Q > 0$ such that for every $x \in \mathbb{P}^1$, and $n \geq 0$, there exists a subset $\{j_1, \dots, j_{q'}\} \subset \{0, 1, \dots, n\}$, such that*

$$\sum_{j=0}^n k(x_j) - \sum_{\alpha=1}^q k(x_{j_\alpha}) \leq Qn,$$

here $q' \leq q$ is an integer.

Lemma 2.6 implies that in a sense the orbit of a point can not come close to $C(f)$ very frequently. As a consequence Denker, Przytycki, Urbanski deduce an upper bound of the size of the pre-images of a ball centered at a point in $J(f)$.

Corollary 2.7 (Denker, Przytycki, Urbanski). *There exist $s \geq 1$ and $\rho > 0$ such that for every $x \in J(f)$, for every $\epsilon > 0$, $n \geq 0$, and for every connected component V of $f^{-n}(B(x, \epsilon))$, one has $\text{diam } V \leq s^n \epsilon^\rho$.*

Corollary 2.8. *Let f be a polynomial map in \mathbb{C} . For fixed $R > 0$, there exist $s \geq 1$ and $\rho > 0$ such that for any $n \geq 0$ and any $z \in \mathbb{C}$ satisfying $f^n(z) \in \{|z| < R\}$, we have*

$$d(z, J(f)) \leq s^n d(f^n(z), J(f))^\rho,$$

where the diameter is relative to the Euclidian metric.

Comment: The existence of such a result is intuitive since the Julia set is expected to be repelling in some sense - However the presence of critical points on J makes it non-trivial.

Proof. Since the Euclidian metric and the spherical metric are equivalent on a compact subset of \mathbb{C} , by Corollary 2.7 for fixed $R > 0$, there exist $s \geq 1$ and $\rho > 0$ such that for every z satisfying $z \in J(f)$, $0 < \epsilon \leq R$, $n \geq 0$, and for every connected component V of $f^{-n}(D(z, \epsilon))$, one has $\text{diam } V \leq s^n \epsilon^\rho$.

For any z and n satisfy $f^n(z) \in \{|z| < R\}$, choose $y \in J$ such that $d(f^n(z), J(f)) = d(y, f^n(z))$. Consider the disk $D(y, \epsilon) = D(y, 2d(f^n(z), J(f)))$, for every connected component V of $f^{-n}(D(y, \epsilon))$, one has $\text{diam } V \leq s_1^n \epsilon^\rho$, so that

$$d(z, J(f)) \leq d(z, f^{-n}(y)) \leq \text{diam } V \leq s_1^n \epsilon^\rho = s_1^n 2^\rho d(f^n(z), J(f))^\rho.$$

Set $s = 2^\rho s_1$ and the proof is complete. \square

3. STRUCTURE OF FATTENED FATOU COMPONENTS

In this section we show that every Fatou component of p in the invariant fiber is actually contained in a Fatou component of P , which is called a **fattened Fatou component**, and in this case the Fatou component of p is called **bulged**. By Sullivan's theorem every Fatou component of p is pre-periodic, it is sufficient to show that every periodic Fatou component of p is contained in a Fatou component of P . There are three kinds of periodic Fatou components of p , i.e. attracting basin, parabolic basin and Siegel disk. For all these three kinds we study the structure of the associated fattened Fatou components.

We may iterate P many times to ensure that all periodic Fatou components of p are actually fixed, and all parabolic fixed point has multiplier equals to 1. In the following of this paper the metric referred to is the Euclidian metric.

3.1. Attracting basin case. In the attracting basin case, assume that we have an attracting basin B of p in the invariant fiber. Without loss generality we may assume 0 is the fixed point in B , so that $(0, 0)$ becomes a fixed point of P , and $p'(0) = \lambda'$ with $|\lambda'| < 1$. We have the following well known Theorem ([7]).

Theorem 3.1. *If $P : \Omega \rightarrow \Omega$ is a holomorphic self map, where Ω is an open set of \mathbb{C}^2 and $(0, 0) \in \Omega$ is a fixed point. If all eigenvalues of the derivative $DP(0, 0)$ are less than 1 in absolute value then P has an open attracting basin at the origin.*

In our case we have

$$DP(0, 0) = \begin{pmatrix} \lambda & 0 \\ \frac{\partial f}{\partial t}(0, 0) & \lambda' \end{pmatrix},$$

so that all the all eigenvalues of the derivative $DP(0, 0)$ are less than 1 in absolute value. As a consequence B is contained in a two dimensional attracting basin of $(0, 0)$, say U , so that B is bulged.

3.2. Parabolic basin case. In the parabolic basin case suppose 0 is a parabolic fixed point of p . Assume that p is locally conjugate to $z \mapsto z + az^s + O(z^{s+1})$ for some $s \geq 2$, $a \neq 0$. We first prove that near the fixed point $(0, 0)$, P is locally conjugate to

$$(t, z) \mapsto (\lambda t, z + az^s + O(z^{s+1})).$$

where $O(z^{s+1})$ means there are constant C such that the error term $\leq C|z|^{s+1}$, for all (t, z) in a neighborhood of the origin. Then we prove in this coordinate every parabolic basin of p is bulged.

Lemma 3.2. *Assume $(0, 0)$ is a fixed point of P , and $|p'(0)| = 1$, then there exist a stable manifold through the origin in the horizontal direction. More precisely, there is a holomorphic function $z = \phi(t)$ defined on a small disk $\{|t| < \delta\}$ such that*

$$\phi(0) = 0, \text{ and } f(t, \phi(t)) = \phi(\lambda t).$$

Proof. This is due to the classical Poincaré's theorem since for any integer $m \geq 0$, $\lambda^m \neq p'(0)$. See [9] Theorem 3.1 for the two dimensional version of Poincaré's theorem. \square

Lemma 3.3. *Suppose 0 is a parabolic fixed point of p , then there is a local biholomorphic change of variables such that P is locally conjugate to*

$$(t, z) \mapsto (\lambda t, z + az^s + O(z^{s+1})).$$

where $O(z^{s+1})$ means there are constant C such that the error term $\leq C|z|^{s+1}$, for all (t, z) in a neighborhood of the origin.

Proof. This is a special case of [9, (6.3) _{i, j}], in our case $j = 0$ and $i = s + 1$. \square

By a completely similar argument as in the proof of Leau-Fatou flower Theorem, we can deduce the existence of the two dimensional attracting petals,

Theorem 3.4. *We assume that the local skew product is given by*

$$P(t, z) = (\lambda t, z + az^s + O(z^{s+1})),$$

then there exist a constant $\delta > 0$ and $s-1$ pairwise disjoint simply connected open sets $U_j \subset \{|t| < \delta\} \times \mathbb{C}$, referred to as two dimensional attracting petals, with the following properties:

(1) $P(U_j) \subset U_j$, points in U_j converge to $(0,0)$ locally uniformly.

(2) For any point $x_0 = (t_0, z_0)$ such that $P^n(x_0) \rightarrow (0,0)$, there exist an integer N and j such that for all $n \geq N$ either $P^n(x_0) \in U_j$ or $z_n = 0$.

(3) $U_j = \{|t| < \delta\} \times (U_j \cap \{t = 0\})$.

Proof. This is a special case of [9, §7.2]. □

Thus by Theorem 3.4, for fixed j , all the points x_0 whose orbit finally lands on U_j form an open subset Ω_j , which is contained in the Fatou set of P . It is obvious that every parabolic basin of p is contained in one of such Ω_j , this implies all parabolic basin of p is bulged.

3.3. Siegel disk case. In the Siegel disk case, we assume that 0 is a Siegel point with a Siegel disk $D \subset \{t = 0\}$. We are going to prove that D is contained in a two dimensional Fatou component.

Theorem 3.5. *Assume that p is locally conjugate to $z \mapsto e^{i\theta}z$ with θ a irrational multiple of π , then there is a neighborhood Ω of D such that $D \subset \Omega \subset \mathbb{C}^2$, and there exists a biholomorphic map ψ defined on Ω such that*

$$\psi \circ P \circ \psi^{-1}(t, z) = (\lambda t, e^{i\theta}z).$$

Proof. We may assume that p is conjugate to $z \mapsto e^{i\theta}z$, then by Lemma 3.2 there is a stable manifold $z = \phi(t)$. A change of variables $z \mapsto z + \phi(t)$ straightens the stable manifold so that P is conjugate to

$$(t, z) \rightarrow (\lambda t, e^{i\theta}z + tg(t, z)),$$

where $g(t, z)$ is a holomorphic function. By an abuse of notation we rename this map by P .

Let U be a relatively compact neighborhood of \bar{D} in \mathbb{C}^2 . Set $C = \sup |g(t, z)|$ on U . Let δ be so small that $\frac{C\delta}{1-\delta} < \text{dist}(D, \partial U)$, and then $\Omega = \{|t| < \delta\} \times D$ is an open subset of U . Let (t_0, z_0) be an arbitrary initial point in Ω , and denote $P^n(t_0, z_0)$ by (t_n, z_n) , then

$$\|z_{n+1} - z_n\| \leq |e^{i\theta}z_{n+1} - z_n| = |t_n g(t_n, z_n)| \leq C|\lambda|^n \delta.$$

Then we have

$$\|z_n - z_0\| \leq \frac{C\delta(1 - |\lambda|)^n}{1 - \delta} + |z_0| \leq \frac{C\delta}{1 - \delta} + |z_0| \leq \text{dist}(z_0, \partial U) + |z_0|,$$

so that (t_n, z_n) still lies in U . Thus $\{P^n\}$ is a normal family on Ω , for the reason that $P^n(\Omega)$ is uniformly bounded.

Thus we can select a subsequence $\{n_j\}$ for which the sequence

$$\phi_{t_0}(z_0) = \lim_{j \rightarrow \infty} e^{-in_j\theta} f_{t_{n_j}} \circ f_{t_{n_j-1}} \circ \cdots \circ f_{t_0}(z_0)$$

uniformly converges on compact subset of Ω . Thus $\phi_t(z)$ is a holomorphic function on Ω , and we have

$$\begin{aligned}\phi_{\lambda t_0} \circ f_{t_0}(z_0) &= \lim_{j \rightarrow \infty} e^{-in_j \theta} f_{t_{n_j+1}} \circ f_{t_{n_j}} \circ \cdots \circ f_{t_0}(z_0) \\ &= e^{i\theta} \lim_{j \rightarrow \infty} e^{-i(n_j+1)\theta} f_{t_{n_j}} \circ f_{t_{n_j-1}} \circ \cdots \circ f_{t_0}(z_0) \\ &= e^{i\theta} \phi_{t_0}(z_0)\end{aligned}$$

for every $(t_0, z_0) \in \Omega$. Thus if we let $\psi(t, z) = (t, \phi_t(z))$, since $\phi_0(z) = z$ we can shrink Ω if necessary to make sure that ψ is invertible on Ω , and we have

$$\psi \circ P \circ \psi^{-1}(t, z) = (\lambda t, e^{i\theta} z).$$

For every $(t, z) \in \Omega$. □

It is obvious that Ω is contained in the Fatou set of P . Since $D \subset \Omega$, this implies that every Siegel disk of p is bulged.

3.4. Wandering vertical Fatou disks. We finish section 3 with a definition.

Definition 3.6. *A vertical Fatou disk Δ is called **wandering** if the forward images of Δ does not intersect any fattened Fatou component.*

We note that "wandering" has special meaning in our definition. The definition of wandering vertical Fatou disk we made here is not equivalent to vertical Fatou disks containing wandering points.

Proposition 3.7. *The forward orbit of a wandering vertical Fatou disk clusters only on $J(p)$.*

Proof. This is simply because for every $x = (t, z) \in \Delta$, if $P^n(x)$ tends to $(0, z_0) \in F(p)$ then eventually $P^n((t, z))$ lands in the fattened Fatou component that contains $(0, z_0)$. This contradicts the fact x lying in a wandering Fatou disk. □

4. ESTIMATE OF HORIZONTAL SIZE OF FATTENED FATOU COMPONENTS

In this section we deduce an estimate of the horizontal size of the fattened Fatou components, by applying the one-dimensional DPU Lemma.

In the following we choose $R > 0$ such that if (t, z) satisfies $t \in \Delta$, $|z| > R$, then $|f(t, z)| \geq 2|z|$. This follows the line at infinity is super-attracting. Thus for any holomorphic function $\phi(t)$ on $\{|t| < r\}$ bounded by R , we have for all $|t| < r$,

$$(4.1) \quad |\phi(t) - \phi(0)| \leq 2R \frac{|t|}{r},$$

this follows from the classical Schwarz Lemma.

We begin with a lemma.

Lemma 4.1. *Let $\text{Crit}(P) = \{(t, z) \mid \frac{\partial f}{\partial z}(t, z) = 0\}$, then there exist constants $0 < \delta_1 < 1$ and $K > K_1 > 0$ such that any connected component C_k of $\text{Crit}(P) \cap \{|t| < \delta_1\}$ intersects the line $\{t = 0\}$ in a unique point, say c_k , and for any point $x = (t, z) \in \text{Crit}(P)$, say $x \in C_k$, we have*

$$(4.2) \quad |z - c_k| \leq K_1 |t|^{\frac{1}{d_1}}.$$

and

$$(4.3) \quad |f(t, z) - p(c_k)| \leq K|t|^{\frac{1}{d_1}},$$

where $c_k = C_k \cap \{t = 0\}$, and d_1 is the maximal multiplicity of critical points of p .

Proof. Since $\text{Crit}(P)$ is an analytic variety, by Weierstrass preparation theorem we can let $\delta_1 < 1$ small enough so that $\text{Crit}(P) \cap \{|t| < \delta_1\} = \cup_{k=1}^l C_k$ where C_k are local connected analytic sets, $C_k \cap \{t = 0\} = \{c_k\}$. For each fixed component C intersect $\{t = 0\}$ at c , C is given by the zero set of a Weierstrass polynomial,

$$C = \{(t, z) \in \{|t| < \delta_1\} \times \mathbb{C}, g(t, z) = 0\},$$

where $g(t, z) = (z - c)^m + a_{m-1}(t)(z - c)^{m-1} + \dots + a_0(t)$ is a Weierstrass polynomial, $m \leq d_1$ is an integer, $a_i(t)$ are holomorphic functions in t satisfying $|a_i(t)| \leq M|t|$ for some constant $M > 0$.

We show that

$$|z - c| \leq mM|t|^{\frac{1}{m}}.$$

We argue by contradiction. Suppose there exist a point $(t_0, z_0) \in C$ such that $\frac{|z_0 - c|}{|t_0|^{\frac{1}{m}}} = a > mM$, then we have

$$|z_0 - c|^m = a^m |t_0|,$$

and

$$(4.4) \quad \begin{aligned} |a_{m-1}(t_0)(z_0 - c)^{m-1} + \dots + a_0(t_0)| &\leq |a_{m-1}(t_0)(z_0 - c)^{m-1}| + \dots + |a_0(t_0)| \\ &\leq M|t_0|a^{m-1}|t_0|^{\frac{m-1}{m}} + \dots + M|t_0| \\ &\leq mM a^{m-1} |t_0|. \end{aligned}$$

Thus we have $|z_0 - c|^m > |a_{m-1}(t_0)(z_0 - c)^{m-1} + \dots + a_0(t_0)|$, which contradicts to $(t_0, z_0) \in C$. Setting $K_1 = 2d_1 M$ we get (4.2).

Let Ω be a relatively compact open set that contains $\text{Crit}(P) \cap \{|t| < \delta_1\}$. Let

$$M' = \max \left\{ \left| \frac{\partial f}{\partial z} \right|, \left| \frac{\partial f}{\partial t} \right| : (t, z) \in \Omega \right\}.$$

Then for $(t, z) \in C_k$ we have

$$\begin{aligned} |f(t, z) - p(c_k)| &\leq M'|t| + M'|z - c_k| \\ &\leq CM'(|t| + K_1|t|^{\frac{1}{d_1}}) \\ &\leq M'(1 + K_1)|t|^{\frac{1}{d_1}}. \end{aligned}$$

To get (4.3) we set $K = 2 \max \{M'(1 + K_1), 2R\}$. Thus the proof is complete. \square

Remark 4.2. We note that K_1 and K are invariant under a local coordinate change of the form $t \mapsto \phi(t)$ with $\phi(0) = 0$ and $\phi'(0) = 1$. To see this let $a_i(t)$ be the coefficients of the Weierstrass polynomial, the coordinate change $t \mapsto \phi(t)$ with $\phi'(0) = 1$ takes $a_i(t)$ become $a_i(\phi(t))$. We have $|a_i(\phi(t))| \leq 2M|t|$ by shrinking $\delta_1(\phi)$ if necessary, then we get (4.2) with the same constant K_1 (this is the reason for the 2 in definition of K_1). By shrinking $\delta_1(\phi)$ we see that Ω and R are invariant, and

$$\max \left\{ \left| \frac{\partial f(\phi(t), z)}{\partial z} \right|, \left| \frac{\partial f(\phi(t), z)}{\partial t} \right| : (t, z) \in \Omega \right\} \leq 2M'.$$

By the same reason we get (4.3) with the same constant K .

We define a positive real valued function $r(z)$, which measures the horizontal size of the fattened Fatou components.

Definition 4.3. For $z \in \mathbb{C}$ satisfying $|z| < R$ and z lying in the Fatou set of p , we define $r(z)$ to be the supremum of all positive real numbers r such that there exist a horizontal holomorphic disk passing through z with size $2r$, contained in $F(P) \cap \{|z| \leq R\}$.

We are going to prove the following estimate of $r(z)$ under the assumption that the multiplier λ is sufficiently small.

Theorem 4.4. There exist a constant $\lambda_1 = \lambda_1(f) > 0$ such that for fixed $|\lambda| < \lambda_1$, there are constants $k > 0$, $l > 0$ such that for any point $z \in F(p) \cap \{|z| < R\}$,

$$r(z) \geq k d(z, J(p))^l,$$

here $J(p)$ is the Julia set of p in the invariant fiber. Furthermore l depends only on p .

We would like to give an outline of the proof of Theorem 4.4 first. Since there are only finitely many invariant Fatou components of p , and every Fatou component is pre-periodic to one of them, it is enough to prove Theorem 4.4 for z in the basin of an invariant Fatou component. We first fix an invariant Fatou component U , and we prove Theorem 4.4 holds for $z \in U$, then we use the following Pull Back Lemma to obtain the result for z in the basin of U . We start with the Pull Back Lemma.

Lemma 4.5 (Pull Back lemma). There exist a constant $0 < \epsilon < 1$, such that if we let

$$V = \{z \in F(p), d(z, J(p)) < \epsilon\},$$

then for any $z_0 \in F(p) \cap \{|z| < R\}$ such that $p(z_0) \in V$, at least one of the following holds:

$$(4.5) \quad r(z_0) \geq \frac{\alpha}{|\lambda|} r(p(z_0)) d(z_0, C(p))^{d_1(d_1+1)};$$

or

$$(4.6) \quad r(z_0) \geq \beta d(z_0, J(p))^{d_1(d_1+1)}.$$

Here α, β are positive constants only depending on p and the constant K from Lemma 4.1, and $C(p)$ is the set of critical points lying in $J(p)$.

Proof. Let $\text{Crit}(p)$ be the set of critical points of p . We choose ϵ small such that $p(z) \in V$ implies $d(z, p(C(p))) = d(z, p(\text{Crit}(p)))$. Let ϕ be the associated holomorphic function with respect to $p(z_0)$ with size $r(p(z_0))$. We are going to show that the critical value set of P does not intersect the graph of ϕ when the domain of ϕ is small.

Suppose $x' = (t', z')$ lies in $\text{Crit}(P)$ satisfying $t' < r(z_0)$ and $P(x')$ lying in the graph of ϕ . then by Lemma 4.1 the connected component of $\text{Crit}(P)$ containing x' intersects $\{t = 0\}$ at a unique point c . Then we have

$$(4.7) \quad \begin{aligned} d(p(z_0), p(C(p))) &\leq |p(z_0) - p(c)| \\ &= |\phi(0) - \phi(\lambda t') + f(t', z') - p(c)| \\ &\leq |\phi(0) - \phi(\lambda t')| + |f(t', z') - p(c)| \\ &\leq K \frac{|\lambda t'|}{r(p(z_0))} + K |t'|^{1/d_1}. \end{aligned}$$

(4.7) holds by applying Lemma 4.1 and inequality (4.1).

Now there are two possibilities,

(a) If $\frac{|\lambda t'|}{r(p(z_0))} \geq 1$, then

$$|t'| \geq \frac{r(p(z_0))}{|\lambda|}.$$

(b) If $\frac{|\lambda t'|}{r(p(z_0))} < 1$, then

$$\frac{|\lambda t'|}{r(p(z_0))} \leq \frac{|\lambda t'|^{1/d_1}}{r(p(z_0))^{1/d_1}},$$

so that by (4.7) we have

$$d(p(z_0), p(C(p))) \leq K \frac{|\lambda t'|^{1/d_1}}{r(p(z_0))^{1/d_1}} + K |t'|^{1/d_1}.$$

For case (b), there are two subcases,

(b1) If $r(p(z_0)) \leq |\lambda|$, then

$$d(p(z_0), p(C(p))) \leq 2K \frac{|\lambda t'|^{1/d_1}}{r(p(z_0))^{1/d_1}},$$

by applying the fact that there is a constant $c = c(p) > 0$ such that $d(p(z_0), p(C(p))) \geq c d(z_0, C(p))^{d_1+1}$ we have

$$|t'| \geq \frac{\alpha}{|\lambda|} r(p(z_0)) d(z_0, C(p))^{d_1(d_1+1)},$$

where $\alpha = \left(\frac{c}{2K}\right)^{d_1}$.

(b2) If $r(p(z_0)) > |\lambda|$, then

$$d(p(z_0), p(C(p))) < 2K |t'|^{1/d_1}.$$

Thus we have

$$|t'| > \left(\frac{1}{2K}\right)^{d_1} d(p(z_0), J(p))^{d_1}.$$

By applying the fact that there is a constant $c = c(p) > 0$ such that $d(p(z_0), J(p)) \geq c d(z_0, J(p))^{d_1+1}$, we get

$$|t'| \geq \beta d(z_0, J(p))^{d_1(d_1+1)}.$$

where $\beta = c \left(\frac{c}{2K}\right)^{d_1}$.

We can let α small enough such that actually $\alpha d(z_0, C(p))^{d_1(d_1+1)} < 1$, thus for case (b1) we have

$$|t'| \geq \frac{\alpha}{|\lambda|} r(p(z_0)) d(z_0, C(p))^{d_1(d_1+1)} \geq \frac{r(p(z_0))}{|\lambda|},$$

thus case (a) is actually contained in case (b1).

In either case (b1) or (b2) we get a lower bound on t' . Thus for any t which does not exceed that lower bound, $\phi(\lambda t)$ is not a critical value of f_t and so all branches of f_t^{-1} are well defined and holomorphic in a neighborhood of the graph of ϕ . Therefore, choose g_t to be the branch of f_t^{-1} for which $g_0(f_0(z)) = z$, then the function $\psi(t) = g_t(\phi(\lambda t))$ is well defined from $t = 0$ up to $|t| < \eta$ satisfying $\psi(0) = z_0$ and the graph of ψ containing in the Fatou set, where η is the lowerbound from (4.5) and (4.6). We know that ψ is also bounded by R , since otherwise ϕ would not be bounded by R . To avoid the case $|t'| \geq \delta_1$, we can shrink β such

that $\beta d(z_0, J(p))^{d_1(d_1+1)} < \delta_1$ for all z_0 . Thus $|t'| \geq \delta_1$ implies $|t'| \geq \beta d(z_0, J(p))^{d_1(d_1+1)}$. Thus at least one of (4.5) and (4.6) holds. □

Proof of Theorem 4.4. In the following we fix an invariant Fatou component U of p , denote the basin of U by B (If B is the basin of infinity we let B be contained in $\{|z| < R\}$). We can shrink ϵ to ensure that the set $\{z \in B, d(z, J(p)) < 2\epsilon\}$ is contained in $\{|z| < R\}$. In either case we first construct a subset W of B , satisfies the following conditions,

- (1) W eventually traps the forward orbit of any point in B .
- (2) W contains the compact subset $\{z \in B, d(z, J(p)) \geq \epsilon\}$.
- (3) Theorem 4.4 holds for $z \in W$.

Finally we use the Pull Back Lemma to prove Theorem 4.4 holds for $z \in B$.

Step 1: Construction of W . We split the argument in several cases.

• **U is an immediate attracting basin.** Let $\omega \subset U$ be a compact neighborhood of the attracting fixed point. We set $W = \{z \in B, d(z, J(p)) \geq \epsilon\} \cup \omega$, then W automatically satisfies (1) and (2). Since W is also compact and contained in $F(P)$, there is a lower bound $a > 0$ such that $r(z) \geq a$ for every $z \in W$. So there exist $k > 0$ such that $r(z) \geq k d(z, J(p))$ for $z \in W$.

• **U is the attracting basin of ∞ .** We set $W = \{z \in B, d(z, J(p)) \geq \epsilon\}$, then W automatically satisfies (1) and (2). There is a lower bound $a > 0$ such that $r(z) \geq a$ for every $z \in W$. So there exist $k > 0$ such that $r(z) \geq k d(z, J(p))$.

• **U is an immediate parabolic basin.** Let Q be the associated attracting petal of Theorem 3.4. We set $W = \{z \in B, d(z, J(p)) \geq \epsilon\} \cup Q$, then W automatically satisfies (1) and (2). By Theorem 3.4 there is a lower bound $a > 0$ such that $r(z) \geq a$ for every $z \in P$. Thus there is a lower bound $b > 0$ such that $r(z) \geq b$ for every $z \in W$. So there exist $k > 0$ such that $r(z) \geq k d(z, J(p))$.

• **U is a Siegel disk.** We set $W = U \cup \{z \in B, d(z, J(p)) \geq \epsilon\}$, then W automatically satisfies (1) and (2). To prove (3), it is enough to prove (3) for $z \in U$.

Lemma 4.6. *Let U be a Siegel disk, then there are constants $k > 0, l > 0$ such that for any point $z \in U$,*

$$r(z) \geq k d(z, J(p))^l.$$

Further more l only depends on p .

Proof. Since the technique of the proof is similar to that of Theorem 4.4, we postpone the proof to the end of this subsection. □

Step 2: Pull back argument.

We already have the estimate for $z \in W$. For every $z_0 \in B \setminus W$, let $\{z_i\}_{i \geq 0}$ be its forward orbit, and let n be the smallest integer such that z_n lies in W . Let m be the smallest integer such that case (4.5) does not happen, if this m dose not exist, let $m = n$, in either case we have

$$r(z_m) \geq k d(z_m, J(p))^l,$$

for some $k, l > 0$, and for all z_i , $0 \leq i \leq m-1$, we have

$$(4.8) \quad r(z_i) \geq \frac{\alpha}{|\lambda|} r(z_{i+1}) d(z_i, C(p))^{d_1(d_1+1)}.$$

By (4.8) we have

$$\begin{aligned} \log r(z_i) &\geq \log r(z_{i+1}) + \log d(z_i, C(p))^{d_1(d_1+1)} + \log \frac{\alpha}{|\lambda|} \\ &= \log r(z_{i+1}) - d_1(d_1+1)k(z_i) + \log \frac{\alpha}{|\lambda|}, \end{aligned}$$

for all $0 \leq i \leq m-1$, where $k(z_i)$ is as in Lemma 2.6.

Thus we have

$$\log r(z_0) \geq \log r(z_m) - d_1(d_1+1) \sum_{i=0}^{m-1} k(z_i) + m \log \frac{\alpha}{|\lambda|}.$$

By Lemma 2.6 there exist a subset $\{i_1, \dots, i_{q'}\} \subset \{0, 1, \dots, m-1\}$ such that

$$\sum_{i=0}^{m-1} k(z_i) - \sum_{\alpha=1}^{q'} k(z_{i_\alpha}) \leq Qm.$$

Therefore we have

$$(4.9) \quad \begin{aligned} \log r(z_0) &\geq \log r(z_m) - d_1(d_1+1) \sum_{\alpha=1}^{q'} k(z_{i_\alpha}) - d_1(d_1+1)Qm + m \log \frac{\alpha}{|\lambda|} \\ &\geq \log r(z_m) + d_1(d_1+1) \sum_{\alpha=1}^{q'} \log d(z_{i_\alpha}, J(p)) - d_1(d_1+1)Qm + m \log \frac{\alpha}{|\lambda|}. \end{aligned}$$

By Corollary 2.8 we have for each i_α ,

$$\begin{aligned} \log d(z_{i_\alpha}, J(p)) &\geq \frac{1}{\rho} \log d(z_0, J(p)) - \frac{1}{\rho} i_\alpha \log s \\ &\geq \frac{1}{\rho} \log d(z_0, J(p)) - \frac{1}{\rho} m \log s. \end{aligned}$$

Likewise we have,

$$\begin{aligned} \log r(z_m) &\geq \log k + l \log d(z_m, J(p)) \\ &\geq \log k + \frac{l}{\rho} \log d(z_0, J(p)) - \frac{l}{\rho} m \log s. \end{aligned}$$

Thus applying the estimates of $\log d(z_{i_\alpha}, J(p))$ and $\log r(z_m)$ to (4.9) gives

$$\begin{aligned} \log r(z_0) &\geq \log r(z_m) + d_1(d_1+1) \sum_{\alpha=1}^{q'} \log d(z_{i_\alpha}, J(p)) - d_1(d_1+1)Qm + m \log \frac{\alpha}{|\lambda|} \\ &\geq \log k + \frac{l + qd_1(d_1+1)}{\rho} \log d(z_0, J(p)) - \frac{l + qd_1(d_1+1)}{\rho} m \log s - d_1(d_1+1)Qm + m \log \frac{\alpha}{|\lambda|}. \end{aligned}$$

Let us now fix λ_1 so small such that

$$(4.10) \quad \log \frac{\alpha}{\lambda_1} \geq \frac{l + qd_1(d_1+1)}{\rho} \log s + d_1(d_1+1)Q,$$

then for every $|\lambda| < \lambda_1$ we have

$$\log r(z_0) \geq \log k + \frac{l + qd_1(d_1 + 1)}{\rho} \log d(z_0, J(p)),$$

which is equivalent to

$$r(z_0) \geq k d(z_0, J(p))^{l'},$$

where $l' = \frac{l + qd_1(d_1 + 1)}{\rho}$.

We have shown that there are constants $k > 0$, $l' > 0$ such that $r(z) \geq k d(z, J(p))^{l'}$ for $z \in B$, and l' only depends on p , this finishes the proof of Theorem 4.4. \square

Proof of Lemma 4.6 It is enough to prove the estimate for an invariant subset $U_\epsilon \subset U \setminus \{z \in B, d(z, J(p)) \geq \epsilon\}$. First note that the conclusion of Lemma 4.5 holds for all $z_0 \in U_\epsilon$, since for all $z_0 \in U_\epsilon$ the condition $p(z_0) \in V$ holds. Since U is a Siegel disk, the forward orbit $\{z_n\}_{n \geq 0}$ lies in a compact subset S of U , where $z_n = p^n(z_0)$. Thus there is a lower bound $a > 0$ such that $r(z) \geq a$ for $z \in S$, a depending on S . By Lemma 4.5 there are two cases,

(1) There is no such integer n that $r(z_n) \geq \beta d(z_n, J(p))^{(d_1+1)d_1}$, thus all z_n satisfy $r(z_n) \geq \frac{\alpha}{|\lambda|} r(z_{n+1}) d(z_n, C(p))^{d_1(d_1+1)}$.

(2) There is an integer n such that $r(z_n) \geq \beta d(z_n, J(p))^{d_1(d_1+1)}$.

In case (1) for every $i \geq 0$

$$\begin{aligned} \log r(z_i) &\geq \log r(z_{i+1}) + \log d(z_i, C(p))^{d_1(d_1+1)} + \log \frac{\alpha}{|\lambda|} \\ &= \log r(z_{i+1}) - d_1(d_1 + 1)k(z_i) + \log \frac{\alpha}{|\lambda|}. \end{aligned}$$

Thus we have for every $n \geq 0$,

$$\begin{aligned} \log r(z_0) &\geq \log r(z_n) - d_1(d_1 + 1) \sum_{i=0}^{n-1} k(z_i) + n \log \frac{\alpha}{|\lambda|} \\ &\geq \log a + \frac{qd_1(d_1 + 1)}{\rho} \log d(z_0, J(p)) - \frac{qd_1(d_1 + 1)}{\rho} n \log s - d_1(d_1 + 1)Qn + n \log \frac{\alpha}{|\lambda|}. \end{aligned}$$

Let us now fix $|\lambda_1|$ so small such that

$$(4.11) \quad \log \frac{\alpha}{\lambda_1} \geq \frac{l + qd_1(d_1 + 1)}{\rho} \log s + d_1(d_1 + 1)Q + 1,$$

thus for every $|\lambda| < \lambda_1$ we have

$$\log r(z_0) \geq \log a + \frac{qd_1(d_1 + 1)}{\rho} \log d(z_0, J(p)) + n.$$

Let $n \rightarrow \infty$ then $r(z_0)$ can be arbitrary large, which is a contradiction, thus actually case (1) can not happen.

For the case (2), the proof is same as the proof of Theorem 4.4, thus the proof is complete. \square

Remark 4.7. *The constant λ_1 appearing in Theorem 4.4 is invariant under local coordinate change $t \mapsto \phi(t)$ with $\phi(0) = 0$ and $\phi'(0) = 1$. To see this from Lemma 4.5 and Remark 4.2 we know that α is invariant since it only depends on p and K . By (4.10) and (4.11) λ_1 only depends on α and p , hence $\lambda_1(f)$ is invariant.*

5. ESTIMATE OF SIZE OF FORWARD IMAGES OF VERTICAL FATOU DISKS

In this section we adapt the DPU Lemma to the attracting local polynomial skew product case, to show that the size of forward images of a wandering vertical Fatou disk shrinks slowly. We begin with two classical lemmas. We follow Lilov's presentation.

Lemma 5.1. *There exist $c_0 > 0$ depending only on p and $\delta_2 > 0$ such that if $|t| < \delta_2$ and $\Delta(x, r) \subset \{t = t_0\}$ is an arbitrary vertical disk, then $P(\Delta(x, r))$ contains a disk $\Delta(P(x), r') \subset \{t = \lambda t_0\}$ of radius $\geq c_0 r^d$.*

Proof. For fixed $x = (t, z)$ satisfying $|t| < \delta_2, z \in \mathbb{C}$, and for fixed $r > 0$, define a function

$$f_{t,z,r}(w) = \frac{1}{rM_{t,z,r}} (f_t(z - rw) - f_t(z))$$

which is a polynomial defined on the closed unit disk $\overline{D}(0, 1)$. The positive number $M_{t,z,r}$ is defined by

$$M_{t,z,r} = \sup_{w \in \pi(\overline{\Delta}(x,r))} |f'_t(w)|.$$

Let A be the finite dimensional normed space containing all polynomials with degree $\leq d$ on $\overline{D}(0, 1)$, equipped with the uniform norm. Since $|f'_{t,z,r}(w)| \leq 1$ on $\overline{D}(0, 1)$, the family $\{f'_{t,z,r}\}$ is bounded in A . Notice that $f_{t,z,r}(0) = 0$, so that $\{f_{t,z,r}\}$ is also bounded in A . The closure of $\{f_{t,z,r}\}$ contains no constant map since the derivative of constant map vanishes. but $\sup_{\overline{D}(0,1)} |f'_{t,z,r}(w)| = 1$.

Now suppose that there is a sequence $\{f_{t_n, z_n, r_n}\}$ such that $f_{t_n, z_n, r_n}(D(0, 1))$ does not contains $D(0, \delta_n)$, with $\delta_n \rightarrow 0$. We can take a subsequence $f_{t_n, z_n, r_n} \rightarrow g$, where g is a non-constant polynomial map with $g(0) = 0$. Therefore by open mapping Theorem $g(D(0, \frac{1}{2}))$ contains $D(0, \delta)$ for some $\delta > 0$. Then for n large enough $f_{t_n, z_n, r_n}(D(0, 1))$ also contains $D(0, \delta)$, which is a contradiction. Therefore for all parameter $\{t, z, r\}$, $f_{t,z,r}(D(0, 1))$ contain a ball $D(0, \delta)$, which is equivalent to say that

$$(5.1) \quad \Delta(P(x), \delta r M_{t,z,r}) \subset P(\Delta(x, r)).$$

Next we estimate $M_{t,z,r}$ from below. Let $z_1(t), z_2(t), \dots, z_{d-1}(t)$ be all zeroes of $f'_t(z)$. Then $f'_t(z) = da_d(t)(z - z_1(t)) \cdots (z - z_{d-1}(t))$. We choose δ_2 small such that $c_0 = \inf_{|t| \leq \delta_2} |da_d(t)| > 0$. Then we have

$$\begin{aligned} M_{t,z,r} &= \sup_{w \in \pi(\overline{\Delta}(x,r))} |f'_t(w)| = \sup_{w \in \pi(\overline{\Delta}(x,r))} |da_d(t)(z - z_1(t)) \cdots (z - z_{d-1}(t))| \\ &\geq c_0 r^{d-1} \sin^{d-1} \frac{\pi}{d-1}, \end{aligned}$$

this with (5.1) finishes the proof. \square

Lemma 5.2. *There exist $0 < c < c_0$, $\delta_2 > 0$ such that if a vertical disk $\Delta(x, r) \subset \{t = t_0\}$ satisfies $\Delta(x, r) \subset \{|z| < R\}$, $|t_0| < \delta_2$ and $\eta = d(\Delta(x, r), \{t = t_0\} \cap \text{Crit}(P)) > 0$, then $P(\Delta(x, r))$ contains a disk $\Delta(P(x), r') \subset \{t = \lambda t_0\}$ of radius $\geq c \eta^{2d-2} r$.*

Proof. Let $V = \{x_0 = (t_0, z_0) : |t_0| < \delta_2, |z_0| < R, d(x_0, \{t = t_0\} \cap \text{Crit}(P)) > \eta\}$, and set

$$M_1 = \inf_V \left| \frac{\partial f}{\partial z} \right| > 0, \quad M_2 = \sup_V \left| \frac{\partial^2 f}{\partial z^2} \right| < \infty,$$

here M_1 depends on η but M_2 does not.

Thus for $\Delta(x_0, r) \subset \{t = t_0\}$ satisfying $\Delta(x_0, r) \subset \{|z| < R\}$ and $\eta = d(\Delta(x_0, r), \{t = t_0\} \cap \text{Crit}(P)) > 0$, we have $\Delta = \Delta(x_0, r) \subset V \cap \{t = t_0\}$. Pick an arbitrary a in the interior of $\pi(\Delta)$. Then for all $z \in \partial\pi(\Delta)$, we let

$$h(z) = f_{t_0}(z) - f_{t_0}(a) = f'_{t_0}(z)(z - a) + \frac{1}{2}(z - a)^2 g(z).$$

We know $g(z)$ satisfies $|g(z)| \leq M_2$, so that

$$|f'_{t_0}(z)(z - a)| \geq M_1|z - a| \geq M_1 \frac{|z - a|^2}{2r}.$$

In the case $r \leq \frac{M_1}{2M_2}$ we have

$$|f'_{t_0}(z)(z - a)| \geq M_2|z - a|^2 > \frac{1}{2}|(z - a)^2 g(z)|.$$

Thus by Rouché's Theorem the function $h(z)$ has the same number of zero points as $f'_{t_0}(z)(z - a)$, thus $h(z)$ has exactly one zero point $\{z = a\}$. Since $a \in \pi(\Delta)$ is arbitrary we have f_{t_0} is injective on Δ . The classical Koebe's one-quarter Theorem shows that $P(\Delta(x_0, r))$ contains a disk with radius at least

$$(5.2) \quad \frac{1}{4} \left| \frac{\partial f_{t_0}}{\partial z}(z_0) \right| r.$$

Now we estimate $\left| \frac{\partial f_{t_0}}{\partial z}(z_0) \right|$ from below. Let $z_1(t), z_2(t), \dots, z_{d-1}(t)$ be all zeroes of $f'_t(z)$. Then $f'_t(z) = da_d(t)(z - z_1(t)) \cdots (z - z_{d-1}(t))$. We choose δ_2 such that $c_0 = \inf_{|t| \leq \delta_2} |da_d(t)| > 0$. We have for every $1 \leq i \leq d - 1$, $|z_0 - a_i(t_0)| \geq \eta$. Thus we have

$$\left| \frac{\partial f_{t_0}}{\partial z}(z_0) \right| = |da_d(t_0)(z_0 - z_1(t_0)) \cdots (z_0 - z_{d-1}(t_0))| \geq c_0 \eta^{d-1},$$

this with (5.2) gives

$$r' \geq \frac{1}{4} c_0 \eta^{d-1} r.$$

In the case $r \geq \frac{M_1}{2M_2}$, by the same argument we have

$$(5.3) \quad r' \geq \frac{1}{4} c_0 \eta^{d-1} \frac{M_1}{2M_2} \geq \frac{1}{8M_2} c_0^2 \eta^{2d-2}.$$

Setting $c = \frac{1}{2} \min \left\{ \frac{c_0}{4R^{d-1}}, \frac{1}{8RM_2} c_0^2 \right\}$ we get the conclusion.

Remark 5.3. *We note that c is invariant under a local coordinate change of the form $t \mapsto \phi(t)$ with $\phi(0) = 0$ and $\phi'(0) = 1$. To see this, we know c_0 and R are invariant under a local coordinate change of the form $t \mapsto \phi(t)$ with $\phi(0) = 0$ and $\phi'(0) = 1$, and by shrinking $\delta_2(\phi)$ we have*

$$\sup_V \left| \frac{\partial^2 f(\phi(t), z)}{\partial z^2} \right| \leq 2M_2,$$

thus from (5.3) and $c = \frac{1}{2} \min \left\{ \frac{c_0}{4R^{d-1}}, \frac{1}{8RM_2} c_0^2 \right\}$ we get that c is invariant.

□

Now we show that the size of forward images of a wandering vertical Fatou disk shrinks slowly. We begin with a definition.

Definition 5.4. Define the *inradius* ρ as follows: for a domain $U \subset \mathbb{C}$, for every $z \in U \subset \mathbb{C}$, define

$$\rho(z, U) = \sup \{r > 0 \mid D(z, r) \subset U\},$$

here $D(z, r)$ is a disk centered at z with radius r .

Proposition 5.5. Let $\Delta_0 \subset \{t = t_0\}$ be a wandering vertical Fatou disk centered at $x_0 = (t_0, z_0)$. Let $x_n = (t_n, z_n) = P^n(x_0)$. Set $\Delta_n = P^n(\Delta_0)$ for every $n \geq 1$ and let $\rho_n = \rho(z_n, \pi(\Delta_n))$. There is a constant $\lambda_2(f)$ such that for fixed $|\lambda| < \lambda_2$, we have

$$\lim_{n \rightarrow \infty} \frac{|\lambda|^n}{\rho_n} = 0.$$

Proof. Let λ_3 be a positive constant to be determined. It is sufficient to prove the result by replacing Δ_n by $\Delta_n \cap \Delta(x_n, \lambda_3^{n+1})$. In the following we let Δ_n always be contained in $\Delta(x_n, \lambda_3^{n+1})$.

Without loss generality we can assume that $|t_0| < \min\{\delta_1, \delta_2, \lambda_3\}$, where δ_1 is the constant in Lemma 4.1 and δ_2 is the constant in Lemma 5.1 and Lemma 5.2. Let N be a fixed integer such that $N > d^q + 1$, where q is the number of critical points lying in $J(p)$. Let $K = \{|t| < \min\{\delta_1, \delta_2, \lambda_3\}\} \times \{|z| < R\}$ be a relatively compact subset of \mathbb{C}^2 such that for $(t, z) \notin K$, $|f(t, z)| \geq 2|z|$. Since the orbits of points in Δ_0 cluster only on $J(p)$, we have $\Delta_n \in K$ for every n . We need the following lemma:

Lemma 5.6. There is a constant $M > 0$ such that if $|\lambda| < \lambda_3$, for every n and for every $x' = (t_n, w_n) \in \Delta_n$, for every integer m , letting $(t_{n+m}, w_{n+m}) = P^m(x')$ we have,

$$(5.4) \quad |w_{n+m} - p^m(z_n)| \leq M^m \lambda_3^{n+1}.$$

Proof. We prove it by induction. Let M satisfying for $(t, z) \in K$, $\left|\frac{\partial f(t, z)}{\partial t}\right| \leq \frac{M}{2}$ and $\left|\frac{\partial f(t, z)}{\partial z}\right| \leq \frac{M}{2}$. We can also assume M is larger than the constant K in Lemma 4.1. Thus For $m = 0$ it is obviously true. Assume that when $m = k - 1$ is true, we have

$$\begin{aligned} |w_{n+k} - p^k(z_n)| &= |f(t_{n+k-1}, w_{n+k-1}) - f(0, p^{k-1}(z_n))| \\ &\leq \frac{M}{2}|t_{n+k-1}| + \frac{M}{2}|w_{n+k-1} - p^{k-1}(z_n)| \\ &\leq \frac{M}{2}|\lambda|^{n+1} + \frac{M^k}{2}\lambda_3^{n+1} \\ &\leq \frac{M^k}{2}|\lambda|^{n+1} + \frac{M^k}{2}\lambda_3^{n+1} \\ &\leq M^k \lambda_3^{n+1}. \end{aligned}$$

Thus for every m , (5.4) holds. \square

Remark that when $w_n = z_n$, the same argument gives

$$|z_{n+m} - p^m(z_n)| \leq M^m |\lambda|^{n+1}.$$

Let $C(P)$ be the union of components of $\text{Crit}(P)$ such that meet $C(p) = \text{Crit}(p) \cap J(p)$ in the invariant fiber. For every point $x \in \Delta_n$, we define $k(x) = -\log d(x, C(P) \cap \{t = t_n\})$, and $k_n = \sup_{x \in \Delta_n} k(x)$. (This definition allows $k_n = +\infty$.) Recall that N is a fixed integer such that $N > d^q + 1$. We are going to prove a two dimensional DPU Lemma for attracting polynomial skew products:

Lemma 5.7 (Two Dimensional DPU Lemma). *Let $|\lambda| < \lambda_3$, then for every $N^k \leq n < N^{k+1}$, there is a subset*

$$\{\alpha_1, \dots, \alpha_{q'}\} \subset \{N^k - 1, N^k, \dots, n - 1\}$$

and a constant $Q > 0$ such that

$$(5.5) \quad \sum_{i=N^k-1}^{n-1} k_i - \sum_{i=1}^{q'} k_{\alpha_i} \leq Q(n - N^k + 1),$$

here k is an arbitrary integer, $q' \leq q$ is an integer. Recall that q is the number of critical points lying in $J(p)$.

Proof. Recall that the DPU Lemma implies that there is a subset

$$\{\alpha_1, \dots, \alpha_{q'}\} \subset \{N^k - 1, N^k, \dots, n - 1\}$$

and a constant $Q > 0$ such that

$$(5.6) \quad \sum_{i=N^k-1}^{n-1} k(p^{i-N^k+1}(z_{N^k-1})) - \sum_{j=1}^{q'} k(p^{\alpha_j-N^k+1}(z_{N^k-1})) \leq \frac{Q}{2}(n - N^k + 1).$$

So it is sufficient to prove $k_i \leq 2k(p^{i-N^k+1}(z_{N^k-1}))$ for every i not appearing in $\{\alpha_1, \dots, \alpha_{q'}\}$. This is equivalent to

$$(5.7) \quad d(\Delta_i, C(P) \cap \{t = t_i\}) \geq d(p^{i-N^k+1}(z_{N^k-1}), C(p))^2.$$

To prove (5.7), assume that $d(p^{i-N^k+1}(z_{N^k-1}), C(p)) = d(p^{i-N^k+1}(z_{N^k-1}), c_k)$ for some point $c_k \in C(p)$, let C_k be the component of $C(P)$ which meets c_k at invariant fiber, by (5.4) and Lemma 4.1 we have

$$\begin{aligned} d(\Delta_i, C(P) \cap \{t = t_i\}) &\geq d(p^{i-N^k+1}(z_{N^k-1}), C(p)) - \sup_{x' \in \Delta_i} |\pi(x') - p^{i-N^k+1}(z_{N^k-1})| - |w_i - c_k| \\ &\geq d(p^{i-N^k+1}(z_{N^k-1}), C(p)) - M^{i-N^k+1} \lambda_3^{N^k} - M|\lambda|^{\frac{N^k}{d_1}}, \end{aligned}$$

where w_i is $\pi(C_k \cap \{t = t_i\})$.

By $|\lambda| < \lambda_3$ we have

$$M^{i-N^k+1} \lambda_3^{N^k} + M|\lambda|^{\frac{N^k}{d_1}} = \left(M^{i-N^k+1} + M\right) \lambda_3^{\frac{N^k}{d_1}}.$$

Thus we have

$$d(\Delta_i, C(P) \cap \{t = t_i\}) \geq d(p^{i-N^k+1}(z_{N^k-1}), C(p)) - \left(M^{i-N^k+1} + M\right) \lambda_3^{\frac{N^k}{d_1}}.$$

To prove (5.7) it is sufficient to prove

$$(5.8) \quad \left(M^{i-N^k+1} + M\right) \lambda_3^{\frac{N^k}{d_1}} \leq d(p^{i-N^k+1}(z_{N^k-1}), C(p)) - d(p^{i-N^k+1}(z_{N^k-1}), C(p))^2.$$

By (5.6) we have

$$d(p^{i-N^k+1}(z_{N^k-1}), C(p)) \geq e^{-\frac{Q}{2}(n-N^k+1)},$$

thus it is sufficient to prove

$$(5.9) \quad (M^{n-N^k+1} + M)\lambda_3^{\frac{N^k}{d^1}} \leq e^{-\frac{Q}{2}(n-N^k+1)} - e^{-Q(n-N^k+1)}.$$

We can always choose λ_3 sufficiently small to make (5.9) holds for all $k \geq 0$. This end the proof of the two dimensional DPU Lemma (5.5). \square

By Lemma 5.1 and Lemma 5.2 there is a constant $c > 0$ such that

$$(5.10) \quad \rho_{n+1} \geq ce^{-(2d-2)kn} \rho_n.$$

and

$$(5.11) \quad \rho_{n+1} \geq c\rho_n^d.$$

From the above we can now give some estimates of ρ_n . Recall that ρ_n is assumed smaller than $|\lambda_3|^{n+1}$ otherwise we replace it by $\min\{\rho_n, \lambda_3^{n+1}\}$.

Lemma 5.8. *There is a constant $c_1 > 0$ such that for $N^k \leq n < N^{k+1}$, we have*

$$\rho_n \geq c_1^{N^k} \rho_{N^k-1}^{d^q}.$$

Proof. For $N^k \leq i \leq n$, if $i-1 \in \{\alpha_1, \dots, \alpha_q\}$ we apply inequality (5.10), if $i \notin \{\alpha_1, \dots, \alpha_{q'}\}$ we apply inequality (5.11). Thus we have

$$\begin{aligned} \rho_n &\geq c^{n-\alpha_{q'}+1} \exp\left(- (2d-2) \sum_{j=\alpha_{q'}+1}^n k_j\right) \left(\dots \left(c^{\alpha_1-N^k+1} \exp\left(- (2d-2) \sum_{j=N^k-1}^{\alpha_1-1} k_j\right) \rho_{N^k-1}\right)^d \dots\right)^d \\ &\geq c^{(n-N^k+1)d^q} \exp\left(-d^q \sum_{j=N^k-1}^{n-1} k_i + d^q \sum_{j=1}^{q'} k_{\alpha_j}\right) \rho_{N^k-1}^{d^q} \quad (\text{because } q' \leq q) \\ &\geq c^{(n-N^k+1)d^q} \exp\left(-Qd^q (n-N^k+1)\right) \rho_{N^k-1}^{d^q} \quad (\text{by Lemma 5.7}) \\ &\geq c^{N^{k+1}d^q} \exp\left(-Qd^q N^{k+1}\right) \rho_{N^k-1}^{d^q}. \end{aligned}$$

Setting $c_1 = \min\{c^{N^k} e^{-QN^k}, \lambda_3\}$ we get the desired conclusion. \square

Lemma 5.9. *For $N^k \leq n < N^{k+1}$, $\rho_0 \leq \lambda_3$ we have*

$$\rho_n \geq c_1^{N^{k+1}} \rho_0^{d^{q(k+1)}}.$$

Proof. By iterating Lemma 5.8 we get that

$$\rho_{N^k-1} \geq c_1^{\frac{N^k-d^qk}{N-d^q}} \rho_0^{d^{qk}} \geq c_1^{N^k} \rho_0^{d^{qk}},$$

so that

$$\rho_n \geq c_1^{N^k} \rho_{N^k-1}^{d^q} \geq c_1^{N^{k+1}} \rho_0^{d^{q(k+1)}},$$

this finishes the proof. \square

Now we can conclude the proof of Proposition 5.5. For $N^k \leq n < N^{k+1}$ we get

$$\frac{|\lambda|^n}{\rho_n} \leq \frac{|\lambda|^{N^k}}{c_1^{N^{k+1}} \rho_0^{d^{q(k+1)}}}.$$

Choosing λ_2 small such that

$$(5.12) \quad \lambda_2 < c_1^N,$$

since $N > d^q + 1$ we deduce that for every $|\lambda| < \lambda_0$,

$$\lim_{k \rightarrow \infty} \frac{|\lambda|^{N^k}}{c_1^{N^{k+1}} \rho_0^{d^q(k+1)}} = 0,$$

finally $\lim_{n \rightarrow \infty} \frac{|\lambda|^n}{\rho_n} = 0$, which finishes the proof. □

Remark 5.10. *The constant λ_2 appearing in Proposition 5.5 is invariant under a local coordinate change of the form $t \mapsto \phi(t)$ with $\phi(0) = 0$ and $\phi'(0) = 1$. To see this we know that by (5.9) λ_3 depends only on M and p , M can be dealt with by replacing it everywhere by $2M$ (see Remark 4.2), so that λ_3 is invariant. By putting $c_1 = \min \{c^{Nd^q} e^{-QNd^q}, \lambda_3\}$ we get that c_1 is invariant. Then by (5.12) we get that λ_2 is invariant.*

Corollary 5.11. *In the same setting as Proposition 5.5, for every $l > 0$, if λ is chosen sufficiently small, we have*

$$\lim_{n \rightarrow \infty} \frac{|\lambda|^n}{\rho_n^l} = 0.$$

Proof. By Proposition 5.5 if $|\lambda| < \lambda_2$, then $\lim_{n \rightarrow \infty} \frac{|\lambda|^n}{\rho_n} = 0$ holds. For any $l > 0$, we then let $|\lambda|$ smaller than λ_2^l to make the conclusion holds. □

6. PROOF OF THE NON-WANDERING DOMAIN THEOREM

In this section we prove the non-existence of wandering Fatou components. Let us recall the statement

Theorem 6.1 (No wandering Fatou components). *Let P be a local polynomial skew product with an attracting invariant fiber,*

$$P(t, z) = (\lambda t, f(t, z)).$$

Then for any fixed f , there is a constant $\lambda_0(f) > 0$ such that if λ satisfies $0 < |\lambda| < \lambda_0$, every forward orbit of a vertical Fatou disk intersects a fattened Fatou component. In particular every Fatou component iterates to a fattened Fatou component, and there are no wandering Fatou components.

Proof. We argue by contradiction. Suppose $\Delta_0 \subset \{t = t_0\}$ is a vertical disk lying in a Fatou component which does not iterate to a fattened Fatou component. Without loss generality we may assume $|t_0| < \min \{1, \delta_1, \delta_2, \lambda_3\}$. By Proposition 2.3, Δ_0 is a vertical Fatou disk. Let $x_0 = (t_0, z_0) \in \Delta_0$ be the center of Δ_0 and set $x_n = (t_n, z_n) = P^n(t_0, z_0)$ and $\Delta_n = P^n(\Delta_0)$. We divide the proof into several steps, We set $\rho_n = \rho(z_n, \pi(\Delta_n))$ as before and assume that $\rho_0 \leq \lambda_3$. Notice that Δ_0 can not be contained in the basin of infinity, thus Δ_n is uniformly bounded. Let $\lambda_0 < \min \{\lambda_1, \lambda_2^l\}$, where λ_1 and λ_2 come from Theorem 4.4 and Proposition 5.5. In the course of the proof we will have to shrink λ_0 one more time.

- **Step 1.** By Proposition 3.7, the orbits of points in Δ_0 cluster only on $J(p)$.

• **Step 2.** We show that there exist $N_0 > 0$ such that when $n \geq N_0$, the projection $\pi(\Delta(x_n, \frac{\rho_n}{4}))$ intersects $J(p)$. We determine N_0 in the following. Suppose $\pi(\Delta(x_n, \frac{\rho_n}{4}))$ does not intersect $J(p)$. Thus $z_n \in F(p)$ and Theorem 4.4 implies $r(z_n) \geq k d(z_n, J(p))^l$, then we have

$$\frac{|t_n|}{r(z_n)} \leq \frac{|t_n|}{k d(z_n, J(p))^l} \leq \frac{4^l |t_n|}{k \rho_n^l}.$$

By Corollary 5.11 we can let N_0 large enough so that for all $n \geq N_0$, $\frac{4^l |t_n|}{k \rho_n^l} < 1$. From the definition of $r(z_n)$ we get a horizontal holomorphic disk defined by $\phi(t)$, $|t| < r(z_n)$ contained in the fattened Fatou components, with $\phi(0) = z_n$, and t_n is in the domain of ϕ . Then we have

$$|\phi(t_n) - z_n| = |\phi(t_n) - \phi(0)| \leq 2R \frac{|t_n|}{r(z_n)} \leq 2R \frac{|t_n|}{k d(z_n, J(p))^l} \leq 2R \frac{4^l |t_n|}{k \rho_n^l}.$$

Again by Corollary 5.11, we can let N_0 large enough that for all $n \geq N_0$, $2R \frac{4^l |t_n|}{k \rho_n^l} < \frac{\rho_n}{4}$. Thus $\phi(t_n) \in \Delta(x_n, \frac{\rho_n}{4}) \subset \Delta_n$. Since $\phi(t_n)$ is contained in the fattened Fatou components that contains z_n , this implies Δ_n intersects the fattened Fatou component so it can not be wandering. This contradiction shows that $\pi(\Delta(x_n, \frac{\rho_n}{4}))$ intersects $J(p)$.

Let $y_n \in \Delta_n$ satisfies $\pi(y_n) \in \pi(\Delta(x_n, \frac{\rho_n}{4})) \cap J(p)$, then for all $x \in \Delta(y_n, \frac{\rho_n}{4})$ we have $\rho(\pi(x), \pi(\Delta_n)) \geq \frac{\rho_n}{2}$.

• **Step 3.** We show that there is a integer $N_1 > N_0$ such that for every $x \in \Delta(y_{N_1}, \frac{\rho_{N_1}}{4})$, for every $m \geq 0$, $p^m(\pi(x)) \in \pi(\Delta_{m+N_1})$, here $\pi(y_{N_1}) \in \pi(\Delta(x_{N_1}, \frac{\rho_{N_1}}{4})) \cap J(p)$. This means that the orbit of $\pi(x)$ is always shadowed by the orbit of Δ_{N_1} , which will contradict the fact that $\pi(\Delta_{m+N_1})$ intersects $J(p)$. To show this, we inductively prove the more precise statement that for fixed $N > d^q + 1$, there exist a large $N_1 = N^{k_0} - 1 > N_0$, such that for every $k \geq k_0$, $N^k \leq n < N^{k+1}$, we have

$$(6.1) \quad p^{n-N_1}(\pi(x)) \in \pi(\Delta_n)$$

and

$$(6.2) \quad \rho'_n \geq c_2^{N^{k+1}} \rho_0^{d^q(k+1)},$$

where $\rho'_n = \rho(p^{n-N_1}(\pi(x)), \pi(\Delta_n))$, $c_2 = \frac{c_1}{2}$ comes from Lemma 5.8 and Lemma 5.9. We will determine k_0 in the following.

From Lemma 5.9 we know that (6.1) and (6.2) hold for $n = N_1$. Assume that for some $k \geq k_0$, for all $n \leq N^k - 1$, (6.1) and (6.2) holds. Then for $N^k \leq n < N^{k+1}$, let $y = P^{n-N^k+1}(t_{N^k-1}, p^{N^k-1-N_1}(\pi(x)))$, by Lemma 5.8 we have

$$(6.3) \quad \rho(y, \Delta_n) \geq c_1^{N^k} (\rho'_{N^k-1})^{d^q}.$$

To estimate the distance between $\pi(y)$ and $p^{n-N_1}(\pi(x))$, by Lemma 5.6 we have

$$(6.4) \quad |\pi(y) - p^{n-N_1}(\pi(x))| \leq M^{n-N^k+1} |\lambda|^{N^k}.$$

From (6.3) and (6.4) we have

$$\begin{aligned} \rho'_n &\geq c_1^{N^k} (\rho'_{N^k-1})^{d^q} - M^{n-N^k+1} |\lambda|^{N^k} \\ &\geq c_1^{N^k} (c_2^{N^k} \rho_0^{d^q})^{d^q} - M^{n-N^k+1} |\lambda|^{N^k} \quad (\text{By the induction hypothesis (6.2)}) \\ &\geq c_1^{N^k} c_2^{N^k d^q} \rho_0^{d^q(k+1)} - M^{n-N^k+1} |\lambda|^{N^k}. \end{aligned}$$

By the choice $c_2 = \frac{c_1}{2}$ we have

$$c_1^{N^k} c_2^{N^k d^q} \geq 2c_2^{N^k+1}.$$

Thus

$$\rho'_n \geq 2c_2^{N^k+1} \rho_0^{d^q(k+1)} - M^{n-N^k+1} |\lambda|^{N^k}.$$

To get (6.2) it is sufficient to prove

$$c_2^{N^k+1} \rho_0^{d^q(k+1)} \geq M^{n-N^k+1} |\lambda|^{N^k}.$$

We take λ_0 sufficiently small such that

$$(6.5) \quad c_2^{N^k+1} \geq M^{n-N^k+1} \lambda_0^{\frac{N^k}{2}}$$

($\lambda_0 \leq (\frac{c_2}{M})^{2N}$ is enough). Thus to prove (6.2) it is sufficient to prove that when $|\lambda| < \lambda_0$,

$$(6.6) \quad \rho_0^{d^q(k+1)} \geq |\lambda|^{\frac{N^k}{2}}.$$

Since $N > d^q + 1$, we can choose k_0 large enough such that for every $k > k_0$ (6.6) holds. This finishes the induction.

This shows that (6.1) and (6.2) are true for all $n \geq N_1$.

• **Step 4.** Since for every $x \in \Delta(y_{N_1}, \frac{\rho_{N_1}}{4})$, for every $m \geq 0$, $p^m(\pi(x)) \in \pi(\Delta_{m+N_1})$, and Δ_n is uniformly bounded, the family $\{p^m\}_{m \geq 0}$ restrict on $D(\pi(y_{N_1}), \frac{\rho_{N_1}}{4})$ is a normal family. Thus $\pi(y_{N_1})$ belongs to the Fatou set $F(p)$, this contradicts to $\pi(y_{N_1}) \in J(p)$. Thus the proof is complete. □

Remark 6.2. *The constant λ_0 appearing in Theorem 6.1 is invariant under a local coordinate change of the form $t \mapsto \phi(t)$ with $\phi(0) = 0$ and $\phi'(0) = 1$. To see this we know that the constants $c_2 = \frac{c_1}{2}$, M and N are invariant under a local coordinate change of the form $t \mapsto \phi(t)$ with $\phi(0) = 0$ and $\phi'(0) = 1$ (M can be dealt with by replacing it everywhere by $2M$, see Remark 4.2). Then by (6.5) λ_0 only depends on c_2 , M and N , thus λ_0 is invariant.*

Remark 6.3. *Lilov's Theorem can be seen as a consequence of Theorem 6.1. In fact, for the super-attracting case, the Fatou components of p are bulged for a similar reason. Since when $|t|$ is very small, the contraction to the invariant fiber is stronger than any geometric contraction $t \mapsto \lambda t$, Theorem 4.4 and Proposition 5.5 follows easily. Thus following the argument of Theorem 6.1 gives the result.*

7. POLYNOMIAL SKEW PRODUCTS

In this section we show how the main theorem can be applied to polynomial skew products.

Theorem 7.1. *Let*

$$P(t, z) = (g(t), f(t, z)) : \mathbb{C}^2 \rightarrow \mathbb{C}^2$$

be a globally defined polynomial skew product, where g, f are polynomials. Then there exist a constant $\lambda_0(t_0, f) > 0$ depending only on f and t_0 such that if $g(t_0) = t_0$ and $|g'(t_0)| < \lambda_0$ then there are no wandering Fatou components in $B(t_0) \times \mathbb{C}$, where $B(t_0)$ is the attracting basin of g at t_0 in the t -coordinate.

Proof. First by a coordinate change $\phi_0 : t \mapsto t + t_0$, P is conjugate to

$$P_0 : (t, z) \mapsto (g_0(t), f_0(t, z)),$$

where $g_0(t) = g(t + t_0) - t_0$, and $f_0(t, z) = f(t + t_0, z)$. It is clear that $\{t = 0\}$ becomes an invariant fiber.

By Koenig's Theorem we can introduce a local coordinate change $\phi : t \mapsto \phi(t)$ with $\phi(0) = 0$ and $\phi'(0) = 1$ such that P_0 is locally conjugate to

$$(7.1) \quad (t, z) \mapsto (\lambda t, f_0(\phi(t), z)),$$

where $\lambda = g'(t_0)$.

We have seen in Remark 6.2 that the constant $\lambda_0(f)$ is invariant under a local coordinate change of the form $t \mapsto \phi(t)$ with $\phi(0) = 0$ and $\phi'(0) = 1$. This means that for fixed f , for every such ϕ ,

$$P_\phi : (t, z) \mapsto (\lambda t, f(\phi(t), z))$$

has no wandering Fatou components when $|\lambda| < \lambda_0(f)$. Thus applying this to (7.1) when $|\lambda| = |g'(t_0)| < \lambda_0(f_0)$ we get the local skew product $(t, z) \mapsto (\lambda t, f_0(\phi(t), z))$ has no wandering Fatou components. Thus by conjugation P has no wandering Fatou components in a neighborhood of $\{t = t_0\}$, thus actually P has no wandering Fatou components in $B(t_0) \times \mathbb{C}$, where $B(t_0)$ is the attracting basin of g at t_0 in the t -coordinate. \square

REFERENCES

- [1] Matthieu Astorg, Xavier Buff, Romain Dujardin, Han Peters, and Jasmin Raissy. A two-dimensional polynomial mapping with a wandering fatou component. *Annals of mathematics*, 184(1):263–313, 2016.
- [2] Manfred Denker, Feliks Przytycki, and Mariusz Urbański. On the transfer operator for rational functions on the Riemann sphere. *Ergodic Theory and Dynamical Systems*, 16(02):255–266, 1996.
- [3] Mattias Jonsson. Dynamics of polynomial skew products on \mathbb{C}^2 . *Mathematische Annalen*, 314(3):403–447, 1999.
- [4] Krastio Lilov. Fatou theory in two dimensions. *University of Michigan PhD Thesis*, 2004.
- [5] Han Peters and Iris Marjan Smit. Fatou components of attracting skew-products. *The Journal of Geometric Analysis*, pages 1–27, 2017.
- [6] Han Peters and Liz Raquel Vivas. Polynomial skew-products with wandering Fatou-disks. *Mathematische Zeitschrift*, 283(1-2):349–366, 2016.
- [7] Jean-Pierre Rosay and Walter Rudin. Holomorphic maps from \mathbb{C}^n to \mathbb{C}^n . *Transactions of the American Mathematical Society*, 310(1):47–86, 1988.
- [8] Dennis Sullivan. Quasiconformal homeomorphisms and dynamics I. Solution of the Fatou-Julia problem on wandering domains. *Annals of mathematics*, 122(2):401–418, 1985.
- [9] Tetsuo Ueda et al. Local structure of analytic transformations of two complex variables, I. *Journal of Mathematics of Kyoto University*, 26(2):233–261, 1986.

SORBONNE UNIVERSITÉS, LABORATOIRE DE PROBABILITÉS, STATISTIQUE ET MODÉLISATION (LPSM, UMR 8001), 4 PLACE JUSSIEU, 75252 PARIS CEDEX 05, FRANCE

E-mail address: zhuchao.ji@upmc.fr