



## 10th World Congress of Chemical Engineering

Promoters:



World Chemical Engineering Council



EFCE



ESBES

Organized by EFCE-Spain Group:



ANQUE



Enginvers



SEQUI



A19

[www.wcce10.org](http://www.wcce10.org)



# Experimental determination and modelling of cyclopentane hydrates in presence of NaCl, KCl, CaCl<sub>2</sub>, and NaCl-KCl

Son HO-VAN (3<sup>rd</sup> year PhD student)

Jérôme DOUZET

Baptiste BOUILLOT (supervisor)

Jean-Michel HERRI (supervisor)

Barcelona, 26/02/2018




INSPIRING INNOVATION  INNOVANTE PAR TRADITION



# Outlines

- 1. Introduction and objective**
- 2. Experimental Methodology**
- 3. Experimental results**
- 4. Modelling**
- 5. Conclusions & Perspectives**


# Hydrates can be used as a separation technique



Phase equilibria data in presence of diverse salts is needed.




Objective



1. provide equilibrium data CPH in presence of salts
2. Model CPH

# Experimental methodology

## Equilibrium temperature measurement




schematic diagram

- 1. Reactor
- 2. Cryostat
- 3. Impeller
- 4. Agitator
- 5. Cooling jacket
- 6. Motor
- 7. Temperature transmitter
- 8. Computer
- 9. Temperature probe
- 10. Drying oven
- 11. Ion chromatography
- 12. Camera



Actual system

# Protocol: Two procedures




$$T_e = 7.7^\circ\text{C}$$

**T Estimation**



**T Accurate**

Temperature profile : Quick dissociation



$$T_e = 7.1^\circ\text{C}$$


Temperature profile : Slow dissociation



# Outlines

- 1. Introduction and objective**
- 2. Experimental Methodology**
- 3. Experimental results**
- 4. Modelling of cyclopentane hydrate**
- 5. Conclusions & Perspectives**

## Equilibrium temperatures in the presence of salts



## Equilibrium temperatures (°C) in presence of NaCl, KCl, NaCl-KCl, and CaCl<sub>2</sub>

Our values in the presence of NaCl, 0%, 5%, 10%, 20% and 23% are very close to Zlyftari [1] and Kishimoto [2]

Salinity, % w/w	NaCl, quick	NaCl, slow	NaCl, [1]	NaCl [2]	KCl, slow	NaCl-KCl, slow	CaCl <sub>2</sub> , slow
0	7.7	7.1	7.11	-	7.1	7.1	7.1
1	6.9	6.4	-	-	6.9	6.7	6.8
2	6.3	5.9	-	-	6.1	6.0	6.3
3.5	5.7	5	-	-	5.5	5.4	5.6
5	4.9	4.1	-	4.45	4.9	4.6	4.9
8	3.5	2.4	-	-	3.6	3.0	3.2
10	2	0.9	1.16	1.25	2.4	1.9	1.8
12	0.9	-0.4	-	-	1.4	0.6	0.1
14	-1	-1.8	-	-	0.4	-0.6	-1.9
16	-2.7	-3.8	-	-	-0.5	-2.1	-4
18	-5	-5.3	-	-	-1.9	-3.6	-6.7
20	-7.2	-7.8	-8.00	-	-3	-5.4	-9.6
22	-9.7	-10.2	-	-	-	-7.2	-13.2
23	-11	-11.6	-11.66	-	-	-	-15.1
25							-19.0

[1]. G. Zlyftari, May 2013.  
[2]. M. Kishimoto, Apr. 2012.

# Modelling: Approach n°1


## Standard SLE equation:

$$\ln a_w = \frac{\Delta H_{fm}}{R} \frac{(T_f - T)}{T_f T} + \frac{\Delta C_{fm}}{R} \left[ \frac{(T_f - T)}{T} - \ln \left( \frac{T_f}{T} \right) \right]$$

Geochemical model PHREEQC predicts water activity  $a_w$



No data for  $\Delta C_{fm}$  in the literature for CPH



# Approach n°2 & 3

At thermodynamic equilibrium


Statistical thermodynamics

$$\Delta\mu_w^{\beta-H} = \Delta\mu_w^{\beta-L}$$

Gibbs-Duhem






$$\Delta\mu_w^{\beta-H} = -RT \sum_i \nu_i \ln(1 - \sum_j \theta_j^i)$$



Approach n°2: Kihara potential  
Interaction potential ( $\varepsilon, \sigma, a$ )

Approach n°3: correlation  
 $\theta = \theta(a_w)$



- All models are capable of predicting equilibrium

- n°3 is the most accurate and simplest method



# Outlines


- 1. Introduction and objective**
- 2. Methodology**
- 3. Experimental results**
- 4. Modelling**
- 5. Conclusions & Perspectives**

# Conclusions

- ❖ Equilibrium temperatures CPH with NaCl, KCl, CaCl<sub>2</sub> or NaCl-KCl present were determined
- ❖ Accurate thermodynamic predictions : Occupancy correlation is the best

***PERSPECTIVES: ADD GAS MOLECULES***


**Thank you very much for your  
attention!!!**



## Procedure to optimize the Kihara parameters approach n°2



## Approach n°3 - Correlation $\theta = F(a_w)$

As in the approach n°2, the **van de Waals and Platteeuw** model is used to describe the hydrate phase. The new correlation  $\theta = F(a_w)$  is obtained



Algorithm of equilibrium temperature prediction: **Approach n°3**

## Approach n°2



Typical form of the deviation between the predicted and the experimental data

$\varepsilon/k$  versus  $\sigma$  at the minimum deviation with experimental data

Kihara parameters for cyclopentane hydrate

$a (10^{-10} \text{ m})$	$\sigma (10^{-10} \text{ m})$	$\varepsilon/k$
0.8968	2.72	265.5