



HAL
open science

The real plane Cremona group is an amalgamated product

Susanna Zimmermann

► **To cite this version:**

Susanna Zimmermann. The real plane Cremona group is an amalgamated product. *Annales de l'Institut Fourier*, 2021, 71 (3), pp.1023-1045. 10.5802/aif.3415/. hal-01719064v2

HAL Id: hal-01719064

<https://hal.science/hal-01719064v2>

Submitted on 27 Mar 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

THE REAL PLANE CREMONA GROUP IS AN AMALGAMATED PRODUCT

SUSANNA ZIMMERMANN

ABSTRACT. We show that the real Cremona group of the plane is a non-trivial amalgam of two groups amalgamated along their intersection and give an alternative proof of its abelianisation.

1. INTRODUCTION

The plane Cremona group is the group $\text{Bir}_k(\mathbb{P}^2)$ of birational transformations of \mathbb{P}^2 defined over a field k . For algebraically closed fields k , the Noether-Castelnuovo theorem [5] shows that $\text{Bir}_k(\mathbb{P}^2)$ is generated by $\text{Aut}_k(\mathbb{P}^2)$ and the subgroup preserving the pencil of lines through $[1 : 0 : 0]$. It implies that the normal subgroup generated by $\text{Aut}_k(\mathbb{P}^2)$ is equal to $\text{Bir}_k(\mathbb{P}^2)$. Furthermore, [4, Appendix by CORNULIER] shows that $\text{Bir}_k(\mathbb{P}^2)$ is not isomorphic to a non-trivial amalgam of two groups. However, it is isomorphic to a non-trivial amalgam modulo one simple relation [1, 10, 7], and it is isomorphic to a generalised amalgamated product of three groups, amalgamated along all pairwise intersections [16]. For $k = \mathbb{R}$, the group $\text{Bir}_{\mathbb{R}}(\mathbb{P}^2)$ is generated by $\text{Aut}_{\mathbb{R}}(\mathbb{P}^2)$ and the two subgroups

$$\mathcal{J}_* = \{f \in \text{Bir}_{\mathbb{R}}(\mathbb{P}^2) \mid f \text{ preserves the pencil of lines through } [1 : 0 : 0]\}$$

$$\mathcal{J}_\circ = \{f \in \text{Bir}_{\mathbb{R}}(\mathbb{P}^2) \mid f \text{ preserves the pencil of conics through } p_1, \bar{p}_1, p_2, \bar{p}_2\}$$

where $p_1, p_2 \in \mathbb{P}^2$ are two fixed non-real points such that $p_1, \bar{p}_1, p_2, \bar{p}_2$ are not collinear [3, Theorem 1.1]. Over \mathbb{C} , the analog of \mathcal{J}_\circ is conjugate to \mathcal{J}_* since a pencil of conics through four points in \mathbb{P}^2 in general position can be sent onto a pencil of lines through point.

We define $\mathcal{G}_\circ \subset \text{Bir}_{\mathbb{R}}(\mathbb{P}^2)$ to be the subgroup generated by $\text{Aut}_{\mathbb{R}}(\mathbb{P}^2)$ and \mathcal{J}_\circ , and by $\mathcal{G}_* \subset \text{Bir}_{\mathbb{R}}(\mathbb{P}^2)$ the subgroup generated by $\text{Aut}_{\mathbb{R}}(\mathbb{P}^2)$ and \mathcal{J}_* . Then $\text{Bir}_{\mathbb{R}}(\mathbb{P}^2)$ is generated by $\mathcal{G}_* \cup \mathcal{G}_\circ$, and the intersection $\mathcal{G}_* \cap \mathcal{G}_\circ$ contains the subgroup \mathcal{H} generated by $\text{Aut}_{\mathbb{R}}(\mathbb{P}^2)$ and the involution $[x : y : z] \mapsto [xz : yz : x^2 + y^2]$, which is contained $\mathcal{J}_\circ \cap \mathcal{J}_*$.

Theorem 1.1. *We have $\mathcal{G}_* \cap \mathcal{G}_\circ = \mathcal{H}$, it is a proper subgroup of \mathcal{G}_\circ and \mathcal{G}_* and*

$$\text{Bir}_{\mathbb{R}}(\mathbb{P}^2) \simeq \mathcal{G}_\circ \underset{\mathcal{H}}{*} \mathcal{G}_*.$$

Moreover, both \mathcal{G}_ and \mathcal{G}_\circ have uncountable index in $\text{Bir}_{\mathbb{R}}(\mathbb{P}^2)$.*

The action of $\text{Bir}_{\mathbb{R}}(\mathbb{P}^2)$ on the Bass-Serre tree associated to the amalgamated product yields the following:

2010 *Mathematics Subject Classification.* 14E07; 20F05; 14P99.

During this work, the author was supported by the Swiss National Science Foundation, by Projet PEPS 2018 JC/JC and by ANR Project FIBALGA ANR-18-CE40-0003-01.

Corollary 1.2. *The group $\mathrm{Bir}_{\mathbb{R}}(\mathbb{P}^2)$ acts on a tree, and all its algebraic subgroups are conjugate to a subgroup of \mathcal{G}_* or of \mathcal{G}_\circ .*

For the finite subgroups of odd order Corollary 1.2 can also be verified by checking their classification in [17].

An earlier version of this article used an explicit presentation of $\mathrm{Bir}_{\mathbb{R}}(\mathbb{P}^2)$ and of \mathcal{G}_\circ in terms of generators and generating relations, the first of which is proven in [18, Theorem 4.4], and the second was proven analogously in the earlier version of this article. The present version does not use either presentation. Instead, we look at the groupoid $\mathrm{Sar}_{\mathbb{R}}(\mathbb{P}^2)$ of birational maps between rational real Mori fibre spaces of dimension 2. It contains $\mathrm{Bir}_{\mathbb{R}}(\mathbb{P}^2)$ as subgroupoid, is generated by Sarkisov links and the elementary relations are a set of generating relations [11, Theorem 3.1], see also Theorem 2.5. This information is encoded in a square complex on which $\mathrm{Bir}_{\mathbb{R}}(\mathbb{P}^2)$ acts [11, §2–3], see also § 2. The groups \mathcal{G}_\circ , \mathcal{G}_* and \mathcal{H} will turn out to be stabilisers of the union of elementary discs containing vertices marked with elements of \mathcal{G}_\circ , \mathcal{G}_* and $\mathcal{G}_\circ \cap \mathcal{G}_*$.

This allows us to provide a new proof of the abelianisation of $\mathrm{Bir}_{\mathbb{R}}(\mathbb{P}^2)$ given in [18]. Using the presentation of $\mathrm{Sar}_{\mathbb{R}}(\mathbb{P}^2)$ we can construct a surjective homomorphism of groupoids $\mathrm{Sar}_{\mathbb{R}}(\mathbb{P}^2) \rightarrow \bigoplus_{(0,1]} \mathbb{Z}/2\mathbb{Z}$, whose restriction onto $\mathrm{Bir}_{\mathbb{R}}(\mathbb{P}^2)$ is surjective as well and coincides with its abelianisation by construction. This will give an alternative proof of the following statement.

Theorem 1.3. [18, Theorem 1.1(1)&(3)] *There is a surjective homomorphism of groups*

$$\Phi: \mathrm{Bir}_{\mathbb{R}}(\mathbb{P}^2) \rightarrow \bigoplus_{(0,1]} \mathbb{Z}/2\mathbb{Z}$$

such that its restriction to \mathcal{J}_\circ is surjective and $\mathcal{G}_ \subset \ker(\Phi) = [\mathrm{Bir}_{\mathbb{R}}(\mathbb{P}^2), \mathrm{Bir}_{\mathbb{R}}(\mathbb{P}^2)]$, where the right hand side is also equal to the normal subgroup generated by $\mathrm{Aut}_{\mathbb{R}}(\mathbb{P}^2)$.*

ACKNOWLEDGEMENT: I would like to thank Anne Lonjou for asking me whether the real plane Cremona group is isomorphic to a generalised amalgamated product of several groups, and the interesting discussions that followed. I would also like to thank Stéphane Lamy for discussions on the square complex, and Jérémy Blanc, Yves de Cornulier for helpful remarks, questions and discussions.

2. A SQUARE COMPLEX ASSOCIATED TO THE CREMONA GROUP

In this section we recall the square complex constructed in [11], whose vertices are marked rational surfaces, its edges of length two are Sarkisov links between them and discs relations between Sarkisov links.

By a surface S , we mean a smooth projective surface defined over \mathbb{R} , and by $S_{\mathbb{C}}$ the same surface but defined over \mathbb{C} . We define the Néron-Severi space $N^1(S_{\mathbb{C}})$ as the space of \mathbb{R} -divisors $N^1(S_{\mathbb{C}}) := \mathrm{Div}(S_{\mathbb{C}}) \otimes \mathbb{R} / \equiv$. The action of the Galois group $\mathrm{Gal}(\mathbb{C}/\mathbb{R})$ on $N^1(S_{\mathbb{C}})$ factors through a finite group, and we denote by $N^1(S)$ the subspace of the $\mathrm{Gal}(\mathbb{C}/\mathbb{R})$ -invariant classes. Since we only consider \mathbb{R} -rational surfaces and $\mathbb{C}[S_{\mathbb{C}}]^* = \mathbb{C}^*$, $N^1(S)$ is also the space of classes of divisors defined over \mathbb{R} (see for instance [14, Lemma 6.3(iii)]). The dimension of $N^1(S_{\mathbb{C}})$ is the Picard rank of S and is denoted by $\rho(S)$. If not stated otherwise, all morphisms are defined over \mathbb{R} .

2.1. Rank r fibrations.

Definition 2.1. Let S be a smooth rational surface, and $r \geq 1$ an integer. We say that S is a *rank r fibration* if there exists a morphism $\pi: S \rightarrow B$, where B is a point or $B = \mathbb{P}^1$, with relative Picard number $\rho(S/B) = r$, and such that the anticanonical divisor $-K_S$ is π -ample.

The last condition means that for any curve C contracted to a point by π , we have $K_S \cdot C < 0$. Observe that the condition on the Picard number is that $\rho(S) = r$ if B is a point, and $\rho(S) = r + 1$ if $B = \mathbb{P}^1$. If S is a rank r fibration, we will write S/B if we want to emphasize the basis of the fibration, and S^r when we want to emphasize the rank.

An isomorphism between two fibrations S/B and S'/B' (necessarily of the same rank r) is an isomorphism $S \xrightarrow{\cong} S'$ such that there exists an isomorphism on the bases (necessarily uniquely defined) that makes the following diagram commute:

$$\begin{array}{ccc} S & \xrightarrow{\cong} & S' \\ \pi \downarrow & & \downarrow \pi' \\ B & \xrightarrow{\cong} & B' \end{array}$$

Observe that the definition of a rank r fibration puts together several notions. If B is a point, then S is just a del Pezzo surface of Picard rank r (over the base field \mathbb{R}). If B is a curve, then S is just a conic bundle of relative Picard rank r : a general fiber is isomorphic to a smooth plane conic over \mathbb{R} , and over \mathbb{C} any singular fiber is the union of two (-1) -curves secant at one point. Remark also that being a rank 1 fibrations is equivalent to being a (smooth) rational Mori fibre space of dimension 2.

As the following examples make it clear, there are sometimes several choices for a structure of rank r fibration on a given surface, that may even correspond to distinct ranks.

Example 2.2. (1) \mathbb{P}^2 with the morphism $\mathbb{P}^2 \rightarrow \text{pt}$, or the Hirzebruch surface \mathbb{F}_n with the morphism $\mathbb{F}_n \rightarrow \mathbb{P}^1$, are rank 1 fibrations.

(2) \mathbb{F}_1 with the morphism $\mathbb{F}_1 \rightarrow \text{pt}$ is a rank 2 fibration. Idem for $\mathbb{F}_0 \rightarrow \text{pt}$. The blow-up $S^2 \rightarrow \mathbb{F}_n \rightarrow \mathbb{P}^1$ of a Hirzebruch surface in a real point or a pair of non-real conjugate point such that each of them is in a distinct fiber, is a rank 2 fibration over \mathbb{P}^1 .

(3) The blow-up of two distinct real points on \mathbb{P}^2 , or of two points of \mathbb{F}_n not lying on the same fiber, gives examples of rank 3 fibrations, with morphisms to the point or to \mathbb{P}^1 respectively.

(4) The quadric surface \mathcal{Q} obtained by blowing up \mathbb{P}^2 in a pair of non-real conjugate points and then contracting the strict transform of the line passing through them is a del Pezzo surface of degree 8 of Picard rank $\rho(\mathcal{Q}) = 1$, and so $\mathcal{Q} \rightarrow \text{pt}$ is a rank 1 fibration.

(5) The blow-up \mathcal{CB}_6 of a pair of non-real conjugate points on \mathcal{Q} is a del Pezzo surface of degree 6 of Picard rank $\rho(\mathcal{CB}_6) = 2$. It has a conic bundle structure $\mathcal{CB}_6 \rightarrow \mathbb{P}^1$, and so \mathcal{CB}_6/pt is a rank 2 fibration while $\mathcal{CB}_6/\mathbb{P}^1$ is a rank 1 fibration.

(6) The blow-up of a real point in \mathcal{CB}_6 yields a rank 2 fibration $\mathcal{CB}_5/\mathbb{P}^1$. Note that \mathcal{CB}_5 is the blow-up of \mathbb{P}^2 in two pairs of non-real conjugate points and the fibres of $\mathcal{CB}_5/\mathbb{P}^1$ are the strict transforms of conics passing through the two pairs.

We call *marking* on a rank r fibration S/B a choice of a birational map $\varphi: S \dashrightarrow \mathbb{P}^2$. We say that two marked fibrations $\varphi: S/B \dashrightarrow \mathbb{P}^2$ and $\varphi': S'/B' \dashrightarrow \mathbb{P}^2$ are *equivalent* if $\varphi'^{-1} \circ \varphi: S/B \rightarrow S'/B'$ is an isomorphism of fibrations. We denote by $(S/B, \varphi)$ an equivalence class under this relation.

If S'/B' and S/B are marked fibrations of respective rank r' and r , we say that S'/B' *factorizes* through S/B if the birational map $S' \rightarrow S$ induced by the markings is a morphism, and moreover there exists a (uniquely defined) morphism $B \rightarrow B'$ such that the following diagram commutes:

$$\begin{array}{ccc} S' & \xrightarrow{\pi'} & B' \\ & \searrow & \nearrow \\ & S \xrightarrow{\pi} & B \end{array}$$

In fact if $B' = \text{pt}$ the last condition is empty, and if $B' \simeq \mathbb{P}^1$ it means that $S' \rightarrow S$ is a morphism of fibration over a common basis \mathbb{P}^1 . Note that $r' \geq r$.

2.2. Square complex. We define a 2-dimensional complex \mathcal{X} as follows. Vertices are equivalence classes of marked rank r fibrations, with $3 \geq r \geq 1$. We put an oriented edge from $(S'/B', \varphi')$ to $(S/B, \varphi)$ if S'/B' factorizes through S/B . If $r' > r$ are the respective ranks of S'/B' and S/B , we say that the edge has type r', r . For each triplets of pairwise linked vertices $(S''^3/B'', \varphi'')$, $(S'^2/B', \varphi')$, $(S^1/B, \varphi)$, we glue a triangle. In this way we obtain a 2-dimensional simplicial complex \mathcal{X} .

Lemma 2.3. [11, Lemma 2.3] *For each edge of type 3, 1 from S''/B'' to S/B , there exist exactly two triangles that admit this edge as a side.*

In view of the lemma, by gluing all the pairs of triangles along edges of type 3,1, and keeping only edges of types 3,2 and 2,1, we obtain a square complex that we still denote \mathcal{X} . When drawing subcomplexes of \mathcal{X} we will often drop part of the information which is clear by context, about the markings, the equivalence classes and/or the fibration. For instance S/B must be understood as $(S/B, \varphi)$ for an implicit marking φ , and (\mathbb{P}^2, φ) as $(\mathbb{P}^2/\text{pt}, \varphi)$. The Cremona group $\text{Bir}_{\mathbb{R}}(\mathbb{P}^2)$ acts on \mathcal{X} by

$$f \cdot (S/B, \varphi) := (S/B, f \circ \varphi).$$

2.3. Sarkisov links and elementary relations. In this section we show that the complex \mathcal{X} encodes the notion of Sarkisov links (or *links* for short), and of elementary relation between them.


First we rephrase the usual notion of Sarkisov links between 2-dimensional Mori fiber spaces. Let $(S/B, \varphi)$, $(S'/B', \varphi')$ be two marked rank 1 fibrations. We say that the induced birational map $S \dashrightarrow S'$ is a *Sarkisov link* if there exists a marked rank 2 fibration S''/B'' that factorizes through both S/B and S'/B' . Equivalently, the vertices corresponding to S/B and S'/B' are at distance 2 in the complex \mathcal{X} , with middle vertex S''/B'' :

$$\begin{array}{ccc} & S''/B'' & \\ & \swarrow & \searrow \\ S/B & & S'/B' \end{array}$$

This definition is in fact equivalent to the usual definition of a link of type I, II, III or IV from S/B to S'/B' (see [9, Definition 2.14] for the definition in arbitrary

dimension). Below we recall these definitions in the context of surfaces, in terms of commutative diagrams where each morphism has relative Picard number 1 (such a diagram corresponds to a “two rays game”), and we give some examples. Remark that these diagrams are not part of the complex \mathcal{X} : in each case, the corresponding subcomplex of \mathcal{X} is just a path of two edges, as described above.


- Type I: B is a point, $B' \simeq \mathbb{P}^1$, and $S' \rightarrow S$ is the blow-up of a real point or a pair of non-real conjugate points such that we have a diagram



Then we take $S''/B'' := S'/\text{pt}$. We also refer to the map $S \dashrightarrow S'$ as link of type I.

Examples are given by the blow-up of a real point on $S' = \mathbb{F}_1 \rightarrow \mathbb{P}^2 = S$ or the blow-up of a pair of non-real conjugate points $S' = \mathcal{CB}_6 \rightarrow \mathcal{Q} = S$. The fibration $\mathbb{F}_1/\mathbb{P}^1$ corresponds to the lines through the point, and the fibration $\mathcal{CB}_6/\mathbb{P}^1$ corresponds to a pencil of conics in \mathbb{P}^2 passing through two pairs of non-real conjugate points.

- Type II: $B = B'$, and there exist two blow-ups $S'' \rightarrow S$ and $S'' \rightarrow S'$ that fit into a diagram of the form:




Then we take $S''/B'' := S''/B$. We also refer to the map $S \dashrightarrow S'$ as link of type II.

An example is given by $S = \mathbb{P}^2$, S'' the blow-up of a pair of non-real conjugate points and $S' = \mathcal{Q}$. Other examples are the blow-up of a conic bundle in a real point, or in a pair of non-real conjugate points not contained in the same fibre, followed by the contraction of the strict transform of the fibre(s) containing it (them).

- Type III: symmetric situation of a link of type I. We refer to the map $S' \rightarrow S$ as link of type III.

- Type IV: (S, φ) and (S', φ') are equal as marked surfaces, but the fibrations to B and B' are distinct. In this situation B and B' must be isomorphic to \mathbb{P}^1 , and we have a diagram



Then we take $S''/B'' := S/\text{pt}$.

For rational surfaces, a type IV link always corresponds to the two rulings on $\mathbb{F}_0 = \mathbb{P}^1 \times \mathbb{P}^1$, that is, $S/B = \mathbb{F}_0/\mathbb{P}^1$ is one of the rulings, $S'/B' = \mathbb{F}_0/\mathbb{P}^1$ the other one, and $S''/B'' = \mathbb{F}_0/\text{pt}$. See [8, Theorem 2.6 (iv)] for other examples in the context of non-rational surfaces.

A *path of links* is a finite sequence of marked rank 1 fibrations

$$(S_0/B_0, \varphi_0), \dots, (S_n/B_n, \varphi_n),$$

such that for all $0 \leq i \leq n-1$, the induced map $g_i: S_i/B_i \dashrightarrow S_{i+1}/B_{i+1}$ is a Sarkisov link.

Proposition 2.4. [11, Proposition 2.6] *Let $(S'/B, \varphi)$ be a marked rank 3 fibration. Then there exist finitely many squares in \mathcal{X} with S' as a corner, and the union of these squares is a subcomplex of \mathcal{X} homeomorphic to a disk with center corresponding to S' .*

In the situation of Proposition 2.4, by going around the boundary of the disc we obtain a path of Sarkisov links whose composition is the identity (or strictly speaking, an automorphism). We say that this path is an *elementary relation* between links, coming from S^3/B . More generally, any composition of links that corresponds to a loop in the complex \mathcal{X} is called a *relation* between Sarkisov links.

Theorem 2.5. [11, Proposition 3.14, Proposition 3.15]

- (1) *Any birational map between rank 1 fibrations is a composition of links, and in particular the complex \mathcal{X} is connected.*
- (2) *Any relation between links is a composition of elementary relations, and in particular \mathcal{X} is simply connected.*

The first part of Theorem 2.5 can also be found in [8, Theorem 2.5]. In fact, a relative version can be extracted from the classification of links in [8, Theorem 2.6].

Proposition 2.6. *Let S/\mathbb{P}^1 and S'/\mathbb{P}^1 be rank 1 fibrations. Any birational map $f: S \dashrightarrow S'$ over \mathbb{P}^1 is a composition of links of rank 1 fibrations over \mathbb{P}^1 , that is, a composition of Sarkisov links of type II over \mathbb{P}^1 . In particular:*

- (1) *Any element of \mathcal{J}_* is conjugate to a composition of automorphisms of some Hirzebruch surface and links of type II between them.*
- (2) *Any element of \mathcal{J}_\circ is conjugate to a composition automorphisms of \mathcal{CB}_6 and links $\mathcal{CB}_6 \dashrightarrow \mathcal{CB}_6$ of type II.*

2.4. Elementary discs. We call the disc with center a rank 3 fibration from Proposition 2.4 an *elementary disc*. In this section, we classify them and therewith obtain an explicit list of elementary relations among rank 1 fibrations.

Lemma 2.7. *Any edge of \mathcal{X} is contained in a square. In particular, \mathcal{X} is the union of elementary discs.*

Proof. Let e be an edge of \mathcal{X} . If it is an edge between a rank 2 fibration and a rank 3 fibration, let e' be an edge from the rank 2 fibration to a rank 1 fibration. By [11, Lemma 2.3], there is a unique square in \mathcal{X} with e and e' among its edges. Suppose that e is the edge between a rank 1 fibration S/B and a rank 2 fibration S'/B' . If $S' = S$ (and so $B = \mathbb{P}^1$, $B' = \text{pt}$), then S is a del Pezzo surface with $\rho(S) = 2$ endowed with a conic bundle structure S/\mathbb{P}^1 . In particular, $S = \mathbb{F}_n$, $n \in \{0, 1\}$ or $S = \mathcal{CB}_6$. Let $S'' \rightarrow S$ be the blow-up of a general point. Then S'' is also a del Pezzo surface, and so S''/pt is a rank 3 fibration and the square


$$\begin{array}{ccc} S/\text{pt} & \longleftarrow & S''/\text{pt} \\ \downarrow e & & \downarrow \\ S/\mathbb{P}^1 & \longleftarrow & S''/\mathbb{P}^1 \end{array}$$

contained in \mathcal{X} . Suppose that $S' \rightarrow S$ is the blow-up of a real point or a pair of non-real conjugate points (and so $B = B'$). Note that if $B = \text{pt}$, the surface S' is a del Pezzo surface of degree ≤ 6 . The blow-up $S'' \rightarrow S'$ of a general point yields a rank 3 fibration S''/B which factorises through S/B . The two blow-ups commute and \mathcal{X} contains a square of the form

$$\begin{array}{ccc} S'/B & \longleftarrow & S''/B \\ \downarrow e & & \downarrow \\ S/B & \longleftarrow & \tilde{S}/B \end{array}$$

It follows that \mathcal{X} is the union of squares. The claim now follows from Proposition 2.4. □

In general, Lemma 2.7 is not true for an arbitrary perfect field k . Indeed, if k has an extension of degree 8, then a Bertini involution whose set of base-points is an orbit of 8 conjugate points is a link of type II



with S/pt a del Pezzo surface of degree 1, and whose corresponding two edges in \mathcal{X} are not contained in any square [11, Lemma 4.3].

In what follows, X_d is a del Pezzo surface of degree $d = (K_{X_d})^2$. A arrow marked by 1 is the blow-up of a real point and an arrow marked with 2 is the blow-up of a pair of non-real conjugate points.

Proposition 2.8. *Any elementary disc in \mathcal{X} is one of the discs $\mathcal{D}_1, \dots, \mathcal{D}_6$ in Figures 4 – 1, where the arrows are marked by 1 (or 2) if it is the blow-up of a real point (or a pair of non-real conjugate points).*


Proof. Let \mathcal{D} be an elementary disc and let S/B be the rank 3 fibration that is its center, and let S'/B' be a rank 1 fibration through which the rank 3 fibration S/B factors.

(a) Suppose that $B = \text{pt}$. Then S is a del Pezzo surface with $\rho(S) = 3$, and hence S'/pt is a del Pezzo surface as well.



If $B' = \mathbb{P}^1$, then $S' \simeq \mathbb{F}_n$, $n \in \{0, 1\}$ or $S' \simeq \mathcal{CB}_6$, and $S \rightarrow S' \rightarrow \mathbb{P}^1$ is a rank 2 fibration. Then $S \rightarrow S'$ is the blow-up of one real point or one pair of non-real conjugate points. If $S' = \mathbb{F}_0$, then $\mathcal{D} = \mathcal{D}_5$. If $S' = \mathbb{F}_1$, then $\mathcal{D} = \mathcal{D}_1$ or $\mathcal{D} = \mathcal{D}_5$. If $S' = \mathcal{CB}_6$ then $\mathcal{D} = \mathcal{D}_3$ or $\mathcal{D} = \mathcal{D}_4$.

If $B' = \text{pt}$, then $S' = \mathbb{P}^2$ or $S' = \mathcal{Q}$, and $\pi: S \rightarrow S'$ is the blow-up of two points (two real points, two pairs of non-real conjugate points or a real point and a pair of non-real points) in general position. If $S' = \mathbb{P}^2$ then \mathcal{D} is one of the discs $\mathcal{D}_1, \mathcal{D}_3, \mathcal{D}_5$. Note that in the discs \mathcal{D}_1 and \mathcal{D}_3 we cover also the case where $S' = \mathcal{Q}$ and π blows up two points, at least one of which is a real point. If $S' = \mathcal{Q}$ and π blows up two pairs of non-real conjugate points, $\mathcal{D} = \mathcal{D}_4$.


(b) Suppose that $B = \mathbb{P}^1$. Then $B' = \mathbb{P}^1$, $S' = \mathbb{F}_n$ or $S' = \mathcal{CB}_6$ and $\pi: S \rightarrow S'$ is a blow-up in two points (again two real points, two pairs of non-real conjugate points or a real point and a pair of non-real points). Furthermore, any rank 1

FIGURE 1. Disc \mathcal{D}_1 on the left, \mathcal{D}_2 on the right.

fibration in the elementary disc is of the form S''/\mathbb{P}^1 . If $S' = \mathcal{CB}_6$, then S has two singular fibres whose \mathbb{C} -components are conjugate, and hence all rank 1 fibrations in the elementary discs are isomorphic to \mathcal{CB}_6 . The hypothesis that $-K_S$ is π -ample implies that the \mathbb{C} -components of the points blown up by π are on pairwise distinct smooth fibres. Thus $\mathcal{D} = \mathcal{D}_2$ or $\mathcal{D} = \mathcal{D}_6$. \square

FIGURE 2. Disc \mathcal{D}_3 .FIGURE 3. Disc \mathcal{D}_4 .

Lemma 2.9. *Let $i, j \in \{1, \dots, 6\}$ and $\mathcal{D}_i \neq \mathcal{D}_j$. If they intersect non-trivially, their intersection is a single vertex or is a boundary segment of the form $S/B \leftarrow S''/B'' \rightarrow S'/B'$, where S/B and S'/B' are rank 1 fibrations.*


 FIGURE 4. Disc \mathcal{D}_5 on the left, disc \mathcal{D}_6 on the right.

Proof. Suppose that $\mathcal{D}_i \cap \mathcal{D}_j$ is non-empty and does not consist of one vertex. Then it contains at least one edge e . If it is in the relative interior of \mathcal{D}_i , the center of \mathcal{D}_i is a vertex of e and hence $\mathcal{D}_i = \mathcal{D}_j$. Suppose that e is on the relative boundary of \mathcal{D}_i , i.e. is of the form $S''/B'' \rightarrow S/B$ with vertices S''/B'' and S/B a rank 2 and rank 1 fibration, respectively. Following the two-rays game, there is a unique edge $S''/B'' \rightarrow S'/B'$ to another rank 1 fibration S'/B' , which is hence contained in both \mathcal{D}_i and \mathcal{D}_j . If $\mathcal{D}_i \cap \mathcal{D}_j$ contains another edge in the relative boundary of \mathcal{D}_i , then the list of elementary discs $\mathcal{D}_1, \dots, \mathcal{D}_6$ in Proposition 2.8 implies that $\mathcal{D}_i = \mathcal{D}_j$. \square

3. PROOF OF THE MAIN THEOREMS

3.1. The group $\mathcal{G}_* \cap \mathcal{G}_\circ$. We denote by $\mathcal{H} \subset \text{Bir}_{\mathbb{R}}(\mathbb{P}^2)$ the subgroup generated by $\text{Aut}_{\mathbb{R}}(\mathbb{P}^2)$ and the quadratic involution $\sigma: [x : y : z] \mapsto [xz : yz : x^2 + y^2]$. Equivalently, it is generated by $\text{Aut}_{\mathbb{R}}(\mathbb{P}^2)$ and the quadratic maps in $\mathcal{J}_\circ \cap \mathcal{J}_*$ [18, Lemma 3.2]. In particular, $\mathcal{H} \subseteq \mathcal{G}_\circ \cap \mathcal{G}_*$. We will prove equality in this section.

- Remark 3.1.**
- (1) For any quadratic map $f \in \text{Bir}_{\mathbb{R}}(\mathbb{P}^2)$ with one real base-point and a pair of non-real conjugate base-points there exist $\alpha, \beta \in \text{Aut}_{\mathbb{R}}(\mathbb{P}^2)$ such that $\alpha f \beta = \sigma$.
 - (2) The path along the boundary of an elementary disc \mathcal{D}_1 connecting its two vertices \mathbb{P}^1/pt corresponds to a quadratic map with one real and two non-real conjugate base-points. It is thus contained in \mathcal{H} and hence also in $\mathcal{G}_\circ \cap \mathcal{G}_*$.
 - (3) We can write $\sigma = \varphi_2 \circ \varphi_1$ where $\varphi_1: \mathbb{P}^2 \dashrightarrow \mathcal{Q}$ and $\varphi_2: \mathcal{Q} \dashrightarrow \mathbb{P}^2$ are two links of type II. It follows that any element of \mathcal{H} is a composition of links $\mathbb{P}^2 \dashrightarrow \mathcal{Q}$ and $\mathcal{Q} \dashrightarrow \mathbb{P}^2$ of type II.

Definition 3.2.

- (1) We call $\mathcal{X}_\circ \subset \mathcal{X}$ the union of the \mathcal{G}_\circ -orbits of all discs $\mathcal{D}_1, \dots, \mathcal{D}_4$ containing a vertex $(\mathcal{Q}/\text{pt}, \tau)$ or $(\mathcal{CB}_6/\mathbb{P}^1, \tau \circ \pi)$ with $\tau: \mathcal{Q} \dashrightarrow \mathbb{P}^2$ is a link of type II and $\pi: \mathcal{CB}_6 \rightarrow \mathcal{Q}$ a link of type III.
- (2) We call $\mathcal{X}_* \subset \mathcal{X}$ the union of \mathcal{G}_* -orbits of all discs $\mathcal{D}_1, \mathcal{D}_5, \mathcal{D}_6$ containing a vertex $(\mathbb{F}_n/\mathbb{P}^1, \pi_n)$ for some $n \geq 0$, with $\pi_1: \mathbb{F}_1 \rightarrow \mathbb{P}^2$ a link of type III and $\pi_n = \pi_1 \circ \varphi_n$, $n \neq 1$, where $\varphi_n: \mathbb{F}_n \dashrightarrow \mathbb{F}_1$ is a composition of Sarkisov links of type II between Hirzebruch surfaces.

As the type of elementary disc is invariant by the action of $\text{Bir}_{\mathbb{R}}(\mathbb{P}^2)$, any elementary disc contained in \mathcal{X}_{\circ} is one of type $\mathcal{D}_1, \dots, \mathcal{D}_4$, and any elementary disc contained in \mathcal{X}_{*} is one of type $\mathcal{D}_1, \mathcal{D}_5$ or \mathcal{D}_6 .

Lemma 3.3.

- (1) The set of vertices (\mathbb{P}^2, g) of \mathcal{X}_{\circ} is the \mathcal{G}_{\circ} -orbit of $(\mathbb{P}^2, \text{id})$.
- (2) The set of vertices (\mathbb{P}^2, g) of \mathcal{X}_{*} is the \mathcal{G}_{*} -orbit of $(\mathbb{P}^2, \text{id})$.
- (3) We have

$$\mathcal{X} = \bigcup_{f \in \text{Bir}_{\mathbb{R}}(\mathbb{P}^2)} f(\mathcal{X}_{\circ}) \cup \bigcup_{f \in \text{Bir}_{\mathbb{R}}(\mathbb{P}^2)} f(\mathcal{X}_{*}).$$

- (4) The sets \mathcal{X}_{\circ} and \mathcal{X}_{*} are connected.
- (5) For any $f \in \text{Bir}_{\mathbb{R}}(\mathbb{P}^2)$, the set $\mathcal{X}_{\circ} \cap f(\mathcal{X}_{*})$ is the union of all elementary discs \mathcal{D}_1 containing a vertex (\mathbb{P}^2, g) with $g \in \mathcal{G}_{\circ}$ and a vertex $(\mathbb{P}^2, f \circ g')$ with $g' \in \mathcal{G}_{*}$.
- (6) The set $\mathcal{X}_{\circ} \cap \mathcal{X}_{*}$ is the union of all elementary discs \mathcal{D}_1 containing a vertex (\mathbb{P}^2, g) with $g \in \mathcal{G}_{\circ} \cap \mathcal{G}_{*}$.

Proof. (1) For any $g \in \mathcal{G}_{\circ}$ we have $(\mathbb{P}^2, g) = g(\mathbb{P}^2, \text{id})$. Let $\tau: \mathcal{Q} \dashrightarrow \mathbb{P}^2$ be a link of type II. There is an elementary disc \mathcal{D}_1 containing $(\mathbb{P}^2, \text{id})$ and (\mathcal{Q}, τ) . Then $g\mathcal{D}_1$ is an elementary disc containing (\mathbb{P}^2, g) and $(\mathcal{Q}, g \circ \tau)$, and it follows that $(\mathbb{P}^2, g) \in \mathcal{X}_{\circ}$. For the converse, if (\mathbb{P}^2, f) is contained in \mathcal{X}_{\circ} , it is the vertex of an elementary disc \mathcal{D}_i , $i \in \{1, 3\}$ contained \mathcal{X}_{\circ} . There is $g \in \mathcal{G}_{\circ}$ such that $\mathcal{D}_i = g\mathcal{D}'_i$ is the image by g of another elementary disc of type \mathcal{D}'_i containing a vertex (\mathcal{Q}, τ) where $\tau: \mathcal{Q} \dashrightarrow \mathbb{P}^2$ is a link of type II. We can choose the vertex (\mathcal{Q}, τ) inside \mathcal{D}'_i such that there are two consecutive edges connecting it to the vertex $(\mathbb{P}^2, g^{-1} \circ f) \in \mathcal{D}'_i$. The two edges correspond to a link $\tau': \mathcal{Q} \dashrightarrow \mathbb{P}^2$ of type II. In particular, $\tau \circ \tau' = g^{-1} \circ f$. Since $\tau \circ \tau' \in \mathcal{G}_{\circ}$, it follows that $f \in \mathcal{G}_{\circ}$.

(2) Is proven analogously to (1) with \mathbb{F}_1 instead of \mathcal{Q} and links $\pi_1: \mathbb{F}_1 \rightarrow \mathbb{P}^2$ of type I and their inverse instead of links $\tau: \mathcal{Q} \dashrightarrow \mathbb{P}^2$ of type II.

(3) The complex \mathcal{X} is the union of elementary discs by Lemma 2.7, hence the orbit of \mathcal{X}_{\circ} and \mathcal{X}_{*} by $\text{Bir}_{\mathbb{R}}(\mathbb{P}^2)$ covers \mathcal{X} .

(4) Let $i \in \{*, \circ\}$. It suffices to check that for any vertex (\mathbb{P}^2, f) of \mathcal{X}_i there is a path in \mathcal{X}_i from (\mathbb{P}^2, f) to $(\mathbb{P}^2, \text{id})$. (1)&(2) imply that $f \in \mathcal{G}_i$. We write $f = \alpha_n f_n \alpha_{n-1} \cdots f_1 \alpha_0$, where $\alpha_j \in \text{Aut}_{\mathbb{R}}(\mathbb{P}^2)$, $f_j \in \mathcal{J}_i$. Proposition 2.6 implies that each f_j is conjugate by $\psi_i: \mathbb{P}^2 \dashrightarrow S_i$ to a composition of links of type II preserving the associated conic bundle S_i/\mathbb{P}^1 , where $\psi_*: \mathbb{P}^2 \dashrightarrow \mathbb{F}_1 = S_*$ is the blow-up of $[1 : 0 : 0]$ and $\psi_{\circ}: \mathbb{P}^2 \dashrightarrow \mathcal{Q} \dashrightarrow \mathcal{CB}_6 = S_{\circ}$ the composition a link of type II and of type I blowing up $[1 : i : 0]$, $[0 : 1 : i]$ and their conjugates. This yields a decomposition of f into links which corresponds to a path from $(\mathbb{P}^2, \text{id})$ to (\mathbb{P}^2, f) along the boundary of elementary discs contained in \mathcal{X}_i .

(5) Lemma 2.9 and Proposition 2.8 imply that the intersection $\mathcal{X}_{\circ} \cap f(\mathcal{X}_{*})$ consists of elementary discs of type \mathcal{D}_1 . By (1)&(2) such a disc contains a vertex (\mathbb{P}^2, g) with $g \in \mathcal{G}_{\circ}$ and a vertex $(\mathbb{P}^2, f \circ g')$ with $g' \in \mathcal{G}_{*}$, and any elementary disc \mathcal{D}_1 with this property is contained $\mathcal{X}_{*} \cap f(\mathcal{X}_{\circ})$.

(6) By (5) for $f = \text{id}$, the set $\mathcal{X}_{\circ} \cap \mathcal{X}_{*}$ is the union of elementary discs \mathcal{D}_1 containing a vertex (\mathbb{P}^2, g) with $g \in \mathcal{G}_{\circ}$ and a vertex (\mathbb{P}^2, g') with $g' \in \mathcal{G}_{*}$. Remark 3.1(2) implies that $g, g' \in \mathcal{G}_{*} \cap \mathcal{G}_{\circ}$. \square

Lemma 3.4. *We have the following stabilisers under the action of $\text{Bir}_{\mathbb{R}}(\mathbb{P}^2)$ on \mathcal{X} :*

- (1) $\text{Stab}(\mathcal{X}_{\circ}) = \mathcal{G}_{\circ}$,
- (2) $\text{Stab}(\mathcal{X}_{*}) = \mathcal{G}_{*}$,

$$(3) \text{Stab}(\mathcal{X}_\circ \cap \mathcal{X}_*) = \mathcal{G}_* \cap \mathcal{G}_\circ.$$

Proof. (1)&(2) Let $i \in \{\circ, *\}$. As \mathcal{X}_i is the \mathcal{G}_i -orbit of certain elementary discs, we have $\mathcal{G}_i \subset \text{Stab}(\mathcal{X}_i)$. Let $f \in \text{Stab}(\mathcal{X}_i)$ and $(\mathbb{P}^2, g) \in \mathcal{X}_i$. Then $(\mathbb{P}^2, f \circ g) \in \mathcal{X}_i$, thus $f \circ g \in \mathcal{G}_i$ by Lemma 3.3(1), and so $f \in \mathcal{G}_i$.

(3) The inclusion $\mathcal{G}_\circ \cap \mathcal{G}_* \subset \text{Stab}(\mathcal{X}_\circ \cap \mathcal{X}_*)$ follows from (1)&(2). Let $f \in \text{Stab}(\mathcal{X}_\circ \cap \mathcal{X}_*)$ and $(\mathbb{P}^2, g) \in \mathcal{X}_\circ \cap \mathcal{X}_*$. Then $(\mathbb{P}^2, g \circ f) \in \mathcal{X}_\circ \cap \mathcal{X}_*$. By Lemma 3.3(6), we have $g, g \circ f \in \mathcal{G}_* \cap \mathcal{G}_\circ$, and so $f \in \mathcal{G}_\circ \cap \mathcal{G}_*$. \square

Lemma 3.5. *Let $f \in \text{Bir}_{\mathbb{R}}(\mathbb{P}^2)$. Then the following hold:*

- (1) *Let $i \in \{\circ, *\}$. If \mathcal{X}_i and $f(\mathcal{X}_i)$ intersect, then they are equal.*
- (2) *$\mathcal{X}_* \cap \mathcal{X}_\circ$ is connected.*
- (3) *$\text{Stab}(\mathcal{X}_\circ \cap \mathcal{X}_*) = \mathcal{G}_\circ \cap \mathcal{G}_* = \mathcal{H}$.*

Proof. (1) Suppose that $\mathcal{X}_i \cap f(\mathcal{X}_i)$ is non-empty. As the type of elementary disc is invariant by the action of $\text{Bir}_{\mathbb{R}}(\mathbb{P}^2)$, the set $\mathcal{X}_i \cap f(\mathcal{X}_i)$ contains a vertex (\mathbb{P}^2, h) for some $h \in \text{Bir}_{\mathbb{R}}(\mathbb{P}^2)$. By Lemma 3.3(1)&(2) we have $h \in \mathcal{G}_i$ and there exists $g \in \mathcal{G}_i$ such that $h = f \circ g$. It follows that $f \in \mathcal{G}_i$. As $\mathcal{G}_i \subset \text{Stab}(\mathcal{X}_i)$, we have $f(\mathcal{X}_i) = \mathcal{X}_i$.

(2)&(3) Let $g \in \mathcal{G}_* \cap \mathcal{G}_\circ$ and write $g = g_1 \circ \cdots \circ g_n = f_1 \circ \cdots \circ f_m$ for some $g_i \in \text{Aut}_{\mathbb{R}}(\mathbb{P}^2) \cup \mathcal{J}_*$ and $f_i \in \text{Aut}_{\mathbb{R}}(\mathbb{P}^2) \cup \mathcal{J}_\circ$. Then the relation

$$g_n^{-1} \circ \cdots \circ g_1^{-1} \circ f_1 \circ \cdots \circ f_m = \text{id}$$

induces a loop γ in \mathcal{X} starting at $(\mathbb{P}^2, \text{id})$ and passing through (\mathbb{P}^2, g_n^{-1}) , $(\mathbb{P}^2, g_n^{-1} \circ g_{n-1}^{-1})$, \dots , $(\mathbb{P}^2, g_n^{-1} \circ \cdots \circ g_1^{-1}) = (\mathbb{P}^2, g^{-1})$, $(\mathbb{P}^2, g^{-1} \circ f_1)$, \dots , $(\mathbb{P}^2, g^{-1} \circ f_1 \circ \cdots \circ f_{m-1})$, $(\mathbb{P}^2, g^{-1} \circ f) = (\mathbb{P}^2, \text{id})$. We can suppose that $\gamma(\frac{1}{2}) = (\mathbb{P}^2, g^{-1})$. Since \mathcal{X}_* and \mathcal{X}_\circ are connected by Lemma 3.3(4) and \mathcal{X} is simply connected by Proposition 2.5(2), we can assume that γ that $\gamma([0, \frac{1}{2}]) \subset \mathcal{X}_*$ and $\gamma([\frac{1}{2}, 1]) \subset g^{-1}(\mathcal{X}_\circ) = \mathcal{X}_\circ$ (Lemma 3.4(1)). The loop γ is the boundary of a disc D in \mathcal{X} , and by Proposition 2.5(2) the disc D is a finite union of elementary discs. In particular, $\gamma([0, \frac{1}{2}]) \subset \mathcal{X}_*$ passes through elementary discs contained in \mathcal{X}_* and $\gamma([\frac{1}{2}, 1]) \subset \mathcal{X}_\circ$ passes through elementary discs contained in \mathcal{X}_\circ . The complex \mathcal{X} is covered by its subcomplexes $f(\mathcal{X}_i)$, $i \in \{*, \circ\}$ by Lemma 3.3(3), hence so is the disc D . It follows that D contains a finite connected union U of elementary discs, all contained in the union of sets of the form $f(\mathcal{X}_\circ) \cap f'(\mathcal{X}_*)$, $f, f' \in \text{Bir}_{\mathbb{R}}(\mathbb{P}^2)$, and which contains $(\mathbb{P}^2, \text{id})$ and (\mathbb{P}^2, g^{-1}) . Lemma 3.3(5) implies that U is covered by elementary discs of type \mathcal{D}_1 . The path from $(\mathbb{P}^2, \text{id})$ to (\mathbb{P}^2, g^{-1}) along the boundary of these discs induces a decomposition of g^{-1} into elements of \mathcal{H} (Remark 3.1(2)). This yields $\mathcal{G}_* \cap \mathcal{G}_\circ \subset \mathcal{H}$. Now (3) follows from Lemma 3.4(3). This and Lemma 3.3(6) imply (2). \square

3.2. The index of the subgroup \mathcal{G}_\circ and the properness of \mathcal{H} in \mathcal{G}_* . We now prove that \mathcal{G}_* is not contained in \mathcal{G}_\circ by showing that quadratic maps having three real base-points in \mathbb{P}^2 are in $\mathcal{G}_* \setminus \mathcal{G}_\circ$, and as consequence that the set $\text{Bir}_{\mathbb{R}}(\mathbb{P}^2) / \mathcal{G}_\circ$ is uncountable.

Recall that any quadratic map $f \in \text{Bir}_{\mathbb{R}}(\mathbb{P}^2)$ with three real base-points in \mathbb{P}^2 is contained in \mathcal{G}_* , as there exist $\alpha, \beta \in \text{Aut}_{\mathbb{R}}(\mathbb{P}^2)$ such that $\alpha f \beta$ is given by

$$\alpha f \beta: [x : y : z] \mapsto [yz : xz : xy]$$

and so $\alpha f \beta \in \mathcal{J}_*$.

Lemma 3.6. *The group \mathcal{G}_\circ does not contain any quadratic map with three real base-points in \mathbb{P}^2 . In particular, $\mathcal{H} \subsetneq \mathcal{G}_*$.*

Proof. Let $f \in \text{Bir}_{\mathbb{R}}(\mathbb{P}^2)$ be a quadratic map with three real base-points p_1, p_2, p_3 in \mathbb{P}^2 and suppose that $f \in \mathcal{G}_o$. Then $f \in \mathcal{G}_o \cap \mathcal{G}_* = \mathcal{H}$ by Lemma 3.5(3) and there is a decomposition $f = \psi_r \cdots \psi_1$ into links $\psi_i: \mathbb{P}^2 \dashrightarrow \mathcal{Q}$ and $\psi_j: \mathcal{Q} \dashrightarrow \mathbb{P}^2$ of type II (Remark 3.1(3)). On the other hand, there is a decomposition of $f = \varphi_4 \cdots \varphi_1$ as follows: The link $\varphi_1: \mathbb{P}^2 \dashrightarrow \mathbb{F}_1$ is the blow-up of p_1 of f , $\varphi_2: \mathbb{F}_1 \dashrightarrow \mathbb{F}_0$ is the link of type II blowing up of p_2 and contracting the line strict transform of the line through p_1, p_2 , and $\varphi_3: \mathbb{F}_0 \dashrightarrow \mathbb{F}_1$ is a link of type II blowing up p_3 and contracting the strict transform of the line through p_1, p_3 , and $\varphi_4: \mathbb{F}_1 \dashrightarrow \mathbb{P}^2$ is a link of type III contracting the strict transform of the line through p_2, p_3 . This corresponds to a path along the boundary of the two discs \mathcal{D}_5 and \mathcal{D}'_5 with respective center (X_7, φ) and $(X_7, \varphi' \circ f^{-1})$, where $\varphi: X_7 \dashrightarrow \mathbb{P}^2$ is the blow-up of the points p_1, p_2 and $\varphi': X_7 \dashrightarrow \mathbb{P}^2$ is the blow-up of the two base-points of f^{-1} which are the images of the lines through p_2, p_3 and p_1, p_3 . The relation

$$\text{id} = \psi_1^{-1} \cdots \psi_r^{-1} \varphi_4 \cdots \varphi_1$$

corresponds to a loop γ in \mathcal{X} which is the border of a disc $D \subset \mathcal{X}$ that is a finite union of elementary discs (Theorem 2.5). We can assume that $\gamma(\frac{1}{2}) = (\mathbb{P}^2, f^{-1})$ and that $\gamma([0, \frac{1}{2}]) \subset \mathcal{X}_o \cap \mathcal{X}_*$. It is therefore covered by elementary discs of the form \mathcal{D}_1 (Lemma 3.3(6)). Furthermore, $\gamma([\frac{1}{2}, 1]) \subset \mathcal{D}_5 \cup \mathcal{D}'_5 \subset \mathcal{X}_*$ by construction of γ . The discs \mathcal{D}_5 and \mathcal{D}'_5 intersect only elementary discs of type \mathcal{D}_1 (they do so in a vertex \mathbb{P}^2/pt or in exactly two connected edges corresponding to a link of type I) and elementary discs type \mathcal{D}_6 , which in turn only intersect other discs of type \mathcal{D}_6 or discs of type \mathcal{D}_5 . It follows that there is a connected union $V \subset D$ of discs of type \mathcal{D}_1 that contains $(\mathbb{P}^2, \text{id})$ and (\mathbb{P}^2, f^{-1}) such that the subset of \tilde{D} bounded by V and $\gamma([\frac{1}{2}, 1])$ is only covered by discs of type $\mathcal{D}_1, \mathcal{D}_5$ and \mathcal{D}_6 . We can then assume that $\gamma([0, \frac{1}{2}]) \subset V$ and that the set \tilde{D} is homeomorphic to a disc. As discs of type \mathcal{D}_6 only intersect other discs \mathcal{D}_6 or discs of type \mathcal{D}_5 , it follows that there is a connected union $\tilde{V} \subset \tilde{D}$ of discs \mathcal{D}_5 containing $(\mathbb{P}^2, \text{id})$ and (\mathbb{P}^2, f^{-1}) intersecting the discs of type \mathcal{D}_1 in V along their boundary. We obtain a path along boundaries of discs of type \mathcal{D}_1 of links of type I and type III from $(\mathbb{P}^2, \text{id})$ to (\mathbb{P}^2, f^{-1}) . This is impossible as f^{-1} has not such decomposition. It follows that $f \notin \mathcal{G}_o$. The rest of the claim follows from the fact that $\mathcal{H} = \mathcal{G}_* \cap \mathcal{G}_o$ (Lemma 3.5(3)). \square

Lemma 3.7. *The group \mathcal{G}_o has uncountable index in $\text{Bir}_{\mathbb{R}}(\mathbb{P}^2)$.*

Proof. Denote by $\sigma: [x : y : z] \mapsto [yz : xz : xy]$ the standard quadratic involution of \mathbb{P}^2 , and define the group

$$A := \{[x : y : z] \mapsto [x + az : y + bz : z] \mid a, b \in \mathbb{R}\} \subset \text{Aut}_{\mathbb{R}}(\mathbb{P}^2)$$

Consider the map between sets

$$\psi: A \longrightarrow \text{Bir}_{\mathbb{R}}(\mathbb{P}^2)/\mathcal{G}_o, \quad \alpha \mapsto (\alpha \circ \sigma)\mathcal{G}_o$$

We now prove that it is injective, which will yield the claim. First, note that $\sigma \circ \alpha \circ \sigma$ is of degree ≤ 2 for all $\alpha \in A$ since $\sigma \circ \alpha$ and σ have two common base-points, namely $[1 : 0 : 0]$ and $[0 : 0 : 1]$. Then $\sigma \circ \alpha \circ \sigma \in \text{Aut}_{\mathbb{R}}(\mathbb{P}^2)$ if and only if σ and $\sigma \circ \alpha$ have three common base-points. This is the case if and only if $\alpha = \text{Id}$, as $\alpha \in A$. Moreover, if $\sigma \circ \alpha \circ \sigma$ is of degree 2, it has three real base-points.

Now, let $\beta, \gamma \in A$ such that $(\beta \circ \sigma)\mathcal{G}_o = (\gamma \circ \sigma)\mathcal{G}_o$. Then $\sigma \circ (\beta^{-1} \circ \gamma) \circ \sigma \in \mathcal{G}_o$. It follows from Lemma 3.6 and the above that $\sigma \circ \beta^{-1} \circ \gamma \circ \sigma \in \text{Aut}_{\mathbb{R}}(\mathbb{P}^2)$, which is the case if and only if $\beta^{-1} \circ \gamma = \text{id}$. Thus the map ψ is injective. \square

3.3. The index of the subgroup \mathcal{G}_o and a quotient of $\text{Bir}_{\mathbb{R}}(\mathbb{P}^2)$. We recall the construction of a surjective homomorphism

$$\psi: \mathcal{J}_o \longrightarrow \bigoplus_{(0,1]} \mathbb{Z}/2\mathbb{Z}$$

given in [18, §3.2.1, §3.2.2].

Definition–Lemma 3.8 ([18, §3.2.1, §3.2.2]). Any element $f \in \mathcal{J}_o$ is conjugate to a birational map $\hat{f}: \mathcal{CB}_6/\mathbb{P}^1 \dashrightarrow \mathcal{CB}_6/\mathbb{P}^1$ preserving the conic bundle structure. Such a map \hat{f} has a decomposition $\hat{f} = \varphi_s \cdots \varphi_1$ into links $\mathcal{CB}_6 \dashrightarrow \mathcal{CB}_6$ of type II (Proposition 2.6), and such a link does not blow up any point on a singular fibre of $\mathcal{CB}_6/\mathbb{P}^1$. For $j = 1, \dots, s$, let C_j be a (real or non-real) fibre of $\mathcal{CB}_6/\mathbb{P}^1$ contracted by φ_j and $[a_j + ib_j : 1] \in \mathbb{P}^1$ its image by the projection onto \mathbb{P}^1 . We define $v_j = 1 - \frac{|a_j|}{a_j^2 + b_j^2} \in (0, 1]$ if $a_j + b_j \notin \mathbb{R}$, and $v_j = 0$ otherwise. Then

$$\psi: \mathcal{J}_o \longrightarrow \bigoplus_{(0,1]} \mathbb{Z}/2\mathbb{Z}, \quad f \mapsto \sum_{j=1}^s e_{v_j}$$

is a surjective homomorphism of groups whose kernel contains all elements of \mathcal{J}_o of degree ≤ 4 .

Remark 3.9. A standard quintic transformation $f \in \text{Bir}_{\mathbb{R}}(\mathbb{P}^2)$ is a transformation of degree 5 having three pairs of non-real conjugate base-points in \mathbb{P}^2 , and no three of the six base-points are collinear (see [3, Example] or [13, §1] for equivalent definitions). In particular, the six points are not on one conic, and f sends the pencil of conics through two of the pairs of base-points of f onto a pencil of conics through two pairs of base-points of f^{-1} . As two non-collinear pairs of non-real conjugate points can be sent by an automorphism of \mathbb{P}^2 onto any other two non-collinear pairs of non-real conjugate points, there are $\alpha, \beta \in \text{Aut}_{\mathbb{R}}(\mathbb{P}^2)$ such that $g := \alpha \circ f \circ \beta \in \mathcal{J}_o$. In particular, $f \in \mathcal{G}_o$. Moreover, it is conjugate to a link of type II of the conic bundle $\mathcal{CB}_6/\mathbb{P}^1$. Moreover, the image $\psi(g)$ of g under the homomorphism $\psi: \mathcal{J}_o \longrightarrow \bigoplus_{(0,1]} \mathbb{Z}/2\mathbb{Z}$ from Definition-Lemma 3.8 is a generator of a $\mathbb{Z}/2\mathbb{Z}$.

We denote by $\text{Sar}_{\mathbb{R}}(\mathbb{P}^2)$ the set of birational transformations between rank 1 fibrations. It is a groupoid and contains $\text{Bir}_{\mathbb{R}}(\mathbb{P}^2)$ as subgroupoid.

Remark 3.10. By Theorem 2.5, the groupoid $\text{Sar}_{\mathbb{R}}(\mathbb{P}^2)$ is generated by links and any relation in $\text{Sar}_{\mathbb{R}}(\mathbb{P}^2)$ is a product of conjugates of elementary relations.

Proposition 3.11 (cf. [18, Proposition 5.3]). *The homomorphism $\psi: \mathcal{J}_o \rightarrow \bigoplus_{(0,1]} \mathbb{Z}/2\mathbb{Z}$ lift to a surjective homomorphism*

$$\Psi: \text{Bir}_{\mathbb{R}}(\mathbb{P}^2) \longrightarrow \bigoplus_{(0,1]} \mathbb{Z}/2\mathbb{Z}$$

whose kernel contains \mathcal{G}_* .

Proof. It suffices to construct a homomorphism of groupoids

$$\Psi: \text{Sar}_{\mathbb{R}}(\mathbb{P}^2) \longrightarrow \bigoplus_{(0,1]} \mathbb{Z}/2\mathbb{Z}$$

whose restriction to its subgroup \mathcal{J}_o is exactly $\psi: \mathcal{J}_o \longrightarrow \bigoplus_{(0,1]} \mathbb{Z}/2\mathbb{Z}$ and whose kernel contains \mathcal{G}_* .

Let $\varphi: \mathcal{CB}_6/\mathbb{P}^1 \dashrightarrow \mathcal{CB}_6/\mathbb{P}^1$ be a link of type II blowing up a pair of non-real conjugate points. Let $\tau_1: \mathbb{P}^2 \dashrightarrow \mathcal{Q}$ be a link of type II blowing up $[1 : i : 0], [1 : -i : 0]$, and $\tau_2: \mathcal{Q} \dashrightarrow \mathcal{CB}_6$ the link of type I blowing up $\tau_1([0 : 1 : i]), \tau_1([0 : 1 : -i])$. Then $f := \tau_1^{-1} \circ \tau_2^{-1} \circ \varphi \circ \tau_2 \circ \tau_1: \mathbb{P}^2 \dashrightarrow \mathbb{P}^2$ is a standard quintic transformation contained in \mathcal{J}_o and we define $\Psi(\varphi) := \psi(f)$. For any other link φ we define $\Psi(\varphi) := 0$. Note that by Definition–Lemma 3.8 and Remark 3.9, we have $\Psi(f) = \psi(f)$ for any $f \in \mathcal{J}_o$. By Remark 3.10 it remains to check that any relation in $\text{Sar}_{\mathbb{R}}(\mathbb{P}^2)$ is sent onto zero by Ψ . Any relation in $\text{Sar}_{\mathbb{R}}(\mathbb{P}^2)$ is the composition of conjugates of elementary relations. Let $\varphi_r \circ \dots \circ \varphi_1$ be an elementary relation. We can assume that one of the φ_i is a link of type II of $\mathcal{CB}_6/\mathbb{P}^1$, as otherwise the relation is sent onto zero by definition of Ψ . The elementary relation corresponds to an elementary disc \mathcal{D} in \mathcal{X} (Theorem 2.5(2)) and φ_i corresponds to a segment on the boundary of the elementary disc. The list of elementary discs in Proposition 2.8 yields that \mathcal{D} is one of $\mathcal{D}_2, \mathcal{D}_3$ or \mathcal{D}_4 . The boundary of each such disc is conjugate via τ_1 or $\tau_1\tau_2$ to a relation inside the group \mathcal{J}_o , so its image by ψ is zero (as ψ is a homomorphism of groups). Thus its image by Ψ is zero. Note that \mathcal{G}_* is contained in $\ker(\Psi)$ by definition of Ψ . \square

The proof of Proposition 3.11 uses the principal idea of the construction of the quotients constructed in [2]; instead of working with a presentation of $\text{Bir}_{\mathbb{R}}(\mathbb{P}^2)$ in terms of generators and generating relations, we pass to such a presentation of the groupoid of birational transformations between rational rank 1 fibrations (rational Mori fibre spaces) and construct a homomorphism from this groupoid to $\bigoplus_{(0,1]} \mathbb{Z}/2\mathbb{Z}$. In [2] one has then to prove in a second step that it is surjective. Here, this is not necessary, as we know that by construction its restriction to \mathcal{J}_o is surjective. The proof of Proposition 3.11 given in [18, Proposition 5.2] uses a presentation of $\text{Bir}_{\mathbb{R}}(\mathbb{P}^2)$ in terms of generators and relations. The proof given here is independent from it.

For a perfect field with an extension of degree 8 (for instance $k = \mathbb{Q}$), there is a surjective homomorphism $\text{Bir}_k(\mathbb{P}^2) \rightarrow *_I \mathbb{Z}/2\mathbb{Z}$ with $|I| \geq |k|$ [11]. It is induced by the free product structure $\text{Bir}_k(\mathbb{P}^2) \simeq G * (*_I \mathbb{Z}/2\mathbb{Z})$, where the factors $\mathbb{Z}/2\mathbb{Z}$ are generated by Bertini involutions whose set of base-points consist of one orbit of eight conjugate points and G is generated by the other generators given in [6] and $\text{Aut}_k(\mathbb{P}^2)$. The free product structure is proven by constructing a quotient of \mathcal{X} on which $\text{Bir}_k(\mathbb{P}^2)$ acts and by showing that it is its Bass-Serre tree. So at its heart, also the construction of the homomorphism from $\text{Bir}_k(\mathbb{P}^2) \rightarrow *_I \mathbb{Z}/2\mathbb{Z}$ uses the presentation of $\text{Sar}_k(\mathbb{P}^2)$ in terms of generators and generating relations.

Corollary 3.12. *The group \mathcal{G}_* does not contain any standard quintic transformations (in particular, $\mathcal{H} \subsetneq \mathcal{G}_o$) and the index of \mathcal{G}_* in $\text{Bir}_{\mathbb{R}}(\mathbb{P}^2)$ is uncountable.*

Proof. Let $\Psi: \text{Bir}_{\mathbb{R}}(\mathbb{P}^2) \longrightarrow \bigoplus_{(0,1]} \mathbb{Z}/2\mathbb{Z}$ be the surjective homomorphism from Proposition 3.11 and $f \in \mathcal{G}_o$ a standard quintic transformation. It is conjugate to a standard quintic transformation in \mathcal{J}_o (Remark 3.9), so $\Psi(f) = \psi(f) \neq 0$ by Proposition 3.11 and Lemma 3.8. Therefore $f \notin \ker \Psi$, while $\mathcal{G}_* \subset \ker \Psi$. It follows that $\mathcal{G}_* \cap \mathcal{G}_o$ is a proper subgroup of \mathcal{G}_o . Moreover, Ψ induces a surjective map $\text{Bir}_{\mathbb{R}}(\mathbb{P}^2)/\mathcal{G}_* \longrightarrow \bigoplus_{(0,1]} \mathbb{Z}/2\mathbb{Z}$ and hence the quotient $\text{Bir}_{\mathbb{R}}(\mathbb{P}^2)/\mathcal{G}_*$ is uncountable. \square

3.4. Proof of the main results. We finally assemble the proofs of Theorem 1.1, Corollary 1.2 and Theorem 1.3.

Proof of Theorem 1.1. The group $\text{Bir}_{\mathbb{R}}(\mathbb{P}^2)$ is generated by \mathcal{G}_* and \mathcal{G}_o . Let $f_n \circ \dots \circ f_1 = \text{id}_{\mathbb{P}^2}$ be a relation in $\text{Bir}_{\mathbb{R}}(\mathbb{P}^2)$ with $f_i \in \mathcal{G}_* \cup \mathcal{G}_o$. By Theorem 2.5(1) for each $i = 1, \dots, r$ there are links $\varphi_{i1}, \dots, \varphi_{ir_i}$ such that $f_i = \varphi_{ir_i} \circ \dots \circ \varphi_{i1}$. Then

$$\varphi_{nr_n} \circ \dots \circ \varphi_{n1} \circ \dots \circ \varphi_{1r_1} \circ \dots \circ \varphi_{11} = \text{id}_{\mathbb{P}^2}$$

is a relation inside the groupoid $\text{Sar}_{\mathbb{R}}(\mathbb{P}^2)$ and is thus a composition of conjugates of elementary relations (Remark 3.10), and each elementary relation corresponds to a loop around the boundary of an elementary disc (Theorem 2.5). Elementary discs are classified in Proposition 2.8. The boundary of discs $\mathcal{D}_1, \mathcal{D}_3$ and \mathcal{D}_5 respectively corresponds to a relation in $\mathcal{G}_o \cap \mathcal{G}_*$, in \mathcal{G}_o and in \mathcal{G}_* , respectively. We attach to the boundary of a disc \mathcal{D}_2 a segment e of the form $\mathcal{Q}/\text{pt} \leftarrow X_7/\text{pt} \rightarrow \mathbb{P}^2/\text{pt}$. To the boundary of a disc \mathcal{D}_4 we attach first the segment $\mathcal{CB}_6/\mathbb{P}^1 \leftarrow \mathcal{CB}_6/\text{pt} \rightarrow \mathcal{Q}/\text{pt}$ and at that one the segment e . This yields that the elementary relation associate to the disc is conjugate to a relation in \mathcal{G}_o . To the boundary of a disc \mathcal{D}_6 we attach the segment corresponding to a composition of links of type II (or an isomorphism if $n = 1$) and a link of type III $\mathbb{F}_n \dashrightarrow \mathbb{F}_1 \rightarrow \mathbb{P}^2$. Then the elementary relation associated to \mathcal{D}_6 is conjugate to a relation in \mathcal{G}_* .

It follows that the relation $f_n \circ \dots \circ f_1 = \text{id}$ is conjugate to a composition of conjugates of relations in \mathcal{G}_* or \mathcal{G}_o or $\mathcal{G}_o \cap \mathcal{G}_*$. This implies that

$$\text{Bir}_{\mathbb{R}}(\mathbb{P}^2) = \mathcal{G}_* \underset{\mathcal{G}_* \cap \mathcal{G}_o}{*} \mathcal{G}_o.$$

Moreover, we have $\mathcal{G}_* \cap \mathcal{G}_o = \mathcal{H}$ by Lemma 3.5(3), and it is a proper subgroup of \mathcal{G}_* and \mathcal{G}_o by Lemma 3.6 and Corollary 3.12. So the amalgamated structure on $\text{Bir}_{\mathbb{R}}(\mathbb{P}^2)$ is nontrivial. Finally, the index of \mathcal{G}_* is uncountable by Lemma 3.7 and the index of \mathcal{G}_o is uncountable by Corollary 3.12. \square

Theorem 1.3. The homomorphism $\Psi: \text{Bir}_{\mathbb{R}}(\mathbb{P}^2) \rightarrow \bigoplus_{(0,1]} \mathbb{Z}/2\mathbb{Z}$ from Proposition 3.11 coincides with the one given in [18, Proposition 5.3] since its restriction to \mathcal{J}_o is the surjective homomorphism $\psi: \mathcal{J}_o \rightarrow \bigoplus_{(0,1]} \mathbb{Z}/2\mathbb{Z}$ and its kernel contains \mathcal{G}_* by construction, hence it also contains $\text{Aut}_{\mathbb{R}}(\mathbb{P}^2)$ and \mathcal{J}_* . The kernel of Ψ is computed in [18, §6] by using [18, §2-3] and is equal to $[\text{Bir}_{\mathbb{R}}(\mathbb{P}^2), \text{Bir}_{\mathbb{R}}(\mathbb{P}^2)]$ and to the normal subgroup generated by $\text{Aut}_{\mathbb{R}}(\mathbb{P}^2)$. \square

Proof of Corollary 1.2. By Theorem 1.1, the group $\text{Bir}_{\mathbb{R}}(\mathbb{P}^2)$ acts on the Bass-Serre tree T of the amalgamated product $\mathcal{G}_* \underset{\mathcal{G}_* \cap \mathcal{G}_o}{*} \mathcal{G}_o$. Then every element of $\text{Bir}_{\mathbb{R}}(\mathbb{P}^2)$ of finite order has a fixed point on T . It follows that every finite subgroup of $\text{Bir}_{\mathbb{R}}(\mathbb{P}^2)$ has a fixed point on T [15, §I.6.5, Corollary 3], and is in particular conjugate to a subgroup of \mathcal{G}_* or of \mathcal{G}_o . For infinite algebraic subgroups of $\text{Bir}_{\mathbb{R}}(\mathbb{P}^2)$, it suffices to check the claim for the maximal algebraic subgroups of $\text{Bir}_{\mathbb{R}}(\mathbb{P}^2)$. By [12, Theorem 1.1], the infinite maximal algebraic subgroups of $\text{Bir}_{\mathbb{R}}(\mathbb{P}^2)$ are conjugate to the group $\text{Aut}_{\mathbb{R}}(X)$ of real automorphisms of a real rational surface X from the following list:

- (1) $X = \mathbb{P}^2$,
- (2) $X = \mathcal{Q}$,
- (3) $X = \mathbb{F}_n$, $n = 0$, $n \geq 2$,
- (4) X is a del Pezzo surface of degree 6 with a birational morphism $X \rightarrow \mathbb{F}_0$ blowing-up a pair of non-real conjugate points.

- (5) X is a del Pezzo surface of degree 6 with a birational morphism $X \rightarrow \mathbb{F}_0$ blowing-up two real points on \mathbb{F}_0 ,
- (6) There is a birational morphism $X/\mathbb{P}^1 \rightarrow \mathcal{CB}_6/\mathbb{P}^1$ of conic bundles blowing up $n \geq 1$ pairs of non-real conjugate points on non-real fibres on the pair of non-real conjugate disjoint (-1) -curves of \mathcal{CB}_6 (the exceptional divisors of the link $\mathcal{CB}_6 \rightarrow \mathcal{Q}$ of type III),
- (7) There is a birational morphism $X \rightarrow \mathbb{F}_n$ of conic bundles blowing up $2n \geq 4$ points on the zero section of self-intersection n .

(1)&(2)&(3) We have $\text{Aut}_{\mathbb{R}}(\mathbb{P}^2) \subset \mathcal{H} = \mathcal{G}_* \cap \mathcal{G}_o$. The group $\text{Aut}_{\mathbb{R}}(\mathcal{Q})$ is conjugate to a subgroup of \mathcal{G}_o via a link $f: \mathcal{Q} \dashrightarrow \mathbb{P}^2$ of type II, and $\text{Aut}_{\mathbb{R}}(\mathbb{F}_n)$ is conjugate to a subgroup of \mathcal{G}_* via a (possibly empty) composition of links of type II and one link of type III $\mathbb{F}_n \dashrightarrow \mathbb{F}_1 \rightarrow \mathbb{P}^2$.

(4) The surface X contains exactly three pairs of non-real conjugate disjoint (-1) -curves. The group $\text{Aut}_{\mathbb{R}}(X)$ is generated by the lift of a subgroup of $\text{Aut}_{\mathbb{R}}(\mathbb{F}_0)$ and two elements, each of which descends via the contraction $X \rightarrow \mathbb{F}_0$ of one pair of (-1) -curves to birational maps of \mathbb{F}_0 preserving one of the two fibrations $\mathbb{F}_0/\mathbb{P}^1$ [12, Proposition 3.5(2)&(3)]. It follows that $\text{Aut}_{\mathbb{R}}(X)$ is conjugate to a subgroup of \mathcal{G}_* .

(5) The surface X contains exactly six real (-1) -curves. Via the blow-down $\eta: X \rightarrow \mathbb{P}^2$ of three disjoint ones, the group $\text{Aut}_{\mathbb{R}}(X)$ is conjugate to a subgroup of $\text{Bir}_{\mathbb{R}}(\mathbb{P}^2)$ generated by the of $\text{Aut}_{\mathbb{R}}(\mathbb{P}^2)$ preserving the images of the (-1) -curves and by $[x : y : z] \mapsto [yz : xz : xy]$ [12, Proposition 3.6(2)&(3)]. So, $\eta \text{Aut}_{\mathbb{R}}(X) \eta^{-1} \subset \mathcal{G}_*$.

(6) The group $\text{Aut}_{\mathbb{R}}(X)$ is generated by the lift of a subgroup of $\text{Aut}_{\mathbb{R}}(\mathcal{Q})$ by $X \rightarrow \mathcal{CB}_6 \rightarrow \mathcal{Q}$ and by elements descending via $X \rightarrow \mathcal{CB}_6$ to birational maps of \mathcal{CB}_6 preserving the conic bundle structure [12, Proposition 4.5(1)&(2)]. It follows that $\text{Aut}_{\mathbb{R}}(X)$ is conjugate to a subgroup of \mathcal{G}_o .

(7) The group $\text{Aut}_{\mathbb{R}}(X)$ is generated by the lift of a subgroups of $\text{Aut}_{\mathbb{R}}(\mathbb{F}_n)$ by $X \rightarrow \mathbb{F}_n$ and by elements descending via $X \rightarrow \mathbb{F}_n$ to birational maps of \mathbb{F}_n preserving the conic bundle structure [12, Proposition 4.8(1)&(2)]. It follows that $\text{Aut}_{\mathbb{R}}(X)$ is conjugate to a subgroup of \mathcal{G}_* . \square

REFERENCES

- [1] J. Blanc. Simple relations in the Cremona group. *Proc. Amer. Math. Soc.*, 140(5):1495–1500, 2012. [1](#)
- [2] J. Blanc, S. Lamy, and S. Zimmermann. Quotients of higher dimensional cremona groups. 01 2019. [14](#)
- [3] J. Blanc and F. Mangolte. Cremona groups of real surfaces. In *Automorphisms in birational and affine geometry*, volume 79 of *Springer Proc. Math. Stat.*, pages 35–58. Springer, Cham, 2014. [1](#), [13](#)
- [4] S. Cantat and S. Lamy. Normal subgroups in the Cremona group. *Acta Math.*, 210(1):31–94, 2013. With an appendix by Yves de Cornulier. [1](#)
- [5] G. Castelnuovo. Le trasformazioni generatrici del gruppo cremoniano nel piano. *Atti della R. Accad. delle Scienze di Torino*, (36):861–874, 1901. [1](#)
- [6] V. Iskovskikh. Generators in the two-dimensional Cremona group over a nonclosed field. Translation of the 1991 paper from Trudy Mat. Inst. Steklov. pages 173–188. 1991. [14](#)
- [7] V. A. Iskovskikh. Proof of a theorem on relations in the two-dimensional Cremona group. *Uspekhi Mat. Nauk*, 40(5(245)):255–256, 1985. [1](#)
- [8] V. A. Iskovskikh. Factorization of birational mappings of rational surfaces from the point of view of Mori theory. *Uspekhi Mat. Nauk*, 51(4(310)):3–72, 1996. [5](#), [6](#)
- [9] A.-S. Kaloghiros. Relations in the Sarkisov program. *Compos. Math.*, 149(10):1685–1709, 2013. [4](#)

- [10] S. Lamy. Groupes de transformations birationnelles de surfaces. Mémoire d'habilitation à diriger des recherches, Université Claude Bernarde Lyon 1, 2010. [1](#)
- [11] S. Lamy and S. Zimmermann. Signature morphisms from the cremona group over a non-closed field. 07 2017. [2](#), [4](#), [6](#), [7](#), [14](#)
- [12] M. Robayo and S. Zimmermann. Infinite algebraic subgroups of the real cremona group. *Osaka J. of Math*, 55(4):681–712, 2018. [15](#), [16](#)
- [13] F. Ronga and T. Vust. Birational diffeomorphisms of the real projective plane. *Comment. Math. Helv.*, 80(3):517–540, 2005. [13](#)
- [14] J.-J. Sansuc. Groupe de Brauer et arithmétique des groupes algébriques linéaires sur un corps de nombres. *J. Reine Angew. Math.*, 327:12–80, 1981. [2](#)
- [15] J.-P. Serre. *Trees*. Springer Monographs in Mathematics. Springer-Verlag, Berlin, 2003. Translated from the French original by John Stillwell, Corrected 2nd printing of the 1980 English translation. [15](#)
- [16] D. Wright. Two-dimensional Cremona groups acting on simplicial complexes. *Trans. Amer. Math. Soc.*, 331(1):281–300, 1992. [1](#)
- [17] E. Yasinsky. Subgroups of odd order in the real plane Cremona group. *J. Algebra*, 461:87–120, 2016. [2](#)
- [18] S. Zimmermann. The abelianisation of the real Cremona group. *arXiv:1510.08705*, 2015. [2](#), [9](#), [13](#), [14](#), [15](#)

SUSANNA ZIMMERMANN, LABORATOIRE ANGEVIN DE RECHERCHE EN MATHÉMATIQUES (LAREMA),
CNRS, UNIVERSITÉ D'ANGERS, 49045 ANGERS CEDEX 1, FRANCE
E-mail address: susanna.zimmermann@univ-angers.fr