



HAL
open science

Real wave propagation in the isotropic-relaxed micromorphic model

Patrizio Neff, Angela Madeo, Gabriele Barbagallo, Marco Valerio d'Agostino, Rafael Abreu, Ionel-Dumitrel Ghiba

► **To cite this version:**

Patrizio Neff, Angela Madeo, Gabriele Barbagallo, Marco Valerio d'Agostino, Rafael Abreu, et al.. Real wave propagation in the isotropic-relaxed micromorphic model. Proceedings of the Royal Society A: Mathematical, Physical and Engineering Sciences, 2017, 473 (2197), 10.1098/rspa.2016.0790 . hal-01720523

HAL Id: hal-01720523

<https://hal.science/hal-01720523>

Submitted on 1 Mar 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Real wave propagation in the isotropic relaxed micromorphic model

Patrizio Neff¹ and Angela Madeo² and Gabriele Barbagallo³
and Marco Valerio d'Agostino⁴ and Rafael Abreu⁵ and Ionel-Dumitrel Ghiba⁶

February 16, 2017

Abstract

For the recently introduced isotropic relaxed micromorphic generalized continuum model, we show that under the assumption of positive definite energy, planar harmonic waves have real velocity. We also obtain a necessary and sufficient condition for real wave velocity which is weaker than positive-definiteness of the energy. Connections to isotropic linear elasticity and micropolar elasticity are established. Notably, we show that strong ellipticity does not imply real wave velocity in micropolar elasticity, while it does in isotropic linear elasticity.

Keywords: ellipticity, positive-definiteness, real wave velocity, planar harmonic waves, rank-one convexity, acoustic tensor, generalized continuum, micropolar, Cosserat, micromorphic

AMS 2010 subject classification: 74A10 (stress), 74A30 (nonsimple materials), 74A60 (micromechanical theories), 74E15 (crystalline structure), 74M25 (micromechanics), 74Q15 (effective constitutive equations)

1 Introduction

Investigations of real wave propagation and ellipticity are not new in principle. Indeed, it is textbook knowledge for linear elasticity that positive definiteness of the elastic energy implies real wave velocities (phase velocities) $v = \omega/k$ where ω [rad/s] is the angular frequency and k [rad/m] $\in \mathbb{R}$ is the wavenumber of planar propagating waves. In classical elasticity, having real wave velocities is equivalent to rank-one convexity (strong ellipticity or Legendre-Hadamard ellipticity). Moreover, ellipticity is equivalent to the positive definiteness of the acoustic tensor. For anisotropic linear elasticity we mention [7], while for anisotropic nonlinear elasticity we refer the reader to [2, 30, 48, 49].

The same question of ellipticity and real wave velocities in generalized continuum mechanics has been discussed for micropolar models, e.g. in [50] and for elastic materials with voids in [8]. For the isotropic micromorphic model results can be found with respect to positive definite energy and/or real wave velocity [46, 51], Mindlin [31, 32] and Eringen's book [11, pp. 277-280]. These latter results present conditions which are neither easily verifiable nor are truly transparent. This is due to the very high number of material coefficients of the Eringen-Mindlin theory that are strongly reduced in the relaxed micromorphic model [29]. Indeed,

¹Patrizio Neff, corresponding author, patrizio.neff@uni-due.de, Head of Chair for Nonlinear Analysis and Modelling, Fakultät für Mathematik, Universität Duisburg-Essen, Mathematik-Carrée, Thea-Leymann-Straße 9, 45127 Essen

²Angela Madeo, angela.madeo@insa-lyon.fr, LGCIE, INSA-Lyon, Université de Lyon, 20 avenue Albert Einstein, 69621, Villeurbanne cedex, France and IUF, Institut universitaire de France, 1 rue Descartes, 75231 Paris Cedex 05, France

³Gabriele Barbagallo, gabriele.barbagallo@insa-lyon.fr, LaMCoS-CNRS & LGCIE, INSA-Lyon, Université de Lyon, 20 avenue Albert Einstein, 69621, Villeurbanne cedex, France

⁴Marco Valerio d'Agostino, marco-valerio.dagostino@insa-lyon.fr, LGCIE, INSA-Lyon, Université de Lyon, 20 avenue Albert Einstein, 69621, Villeurbanne cedex, France

⁵Rafael Abreu, abreu@uni-muenster.de, Institut für Geophysik, Westfälische Wilhelms-Universität Münster, Corrensstraße 24, 48149, Münster, Germany

⁶Ionel-Dumitrel Ghiba, dumitrel.ghiba@uni-due.de, dumitrel.ghiba@uaic.ro, Lehrstuhl für Nichtlineare Analysis und Modellierung, Fakultät für Mathematik, Universität Duisburg-Essen, Thea-Leymann Str. 9, 45127 Essen, Germany; Alexandru Ioan Cuza University of Iași, Department of Mathematics, Blvd. Carol I, no. 11, 700506 Iași, Romania; and Octav Mayer Institute of Mathematics of the Romanian Academy, Iași Branch, 700505 Iași.

the implication that positive definiteness of the energy always implies real wave velocities is not directly established and demonstrated. In this paper we investigate the relaxed micromorphic model in terms of conditions for real wave velocities for planar waves and establish a necessary and sufficient conditions for this to happen.

This paper is organized as follows. We shortly recall the basics of the relaxed micromorphic model and discuss the wave propagation problem for propagating planar waves. Since we deal with an isotropic model, we can, without loss of generality, assume wave propagation in one specific direction only. The dispersion relations are then obtained and real wave-velocities under assumption of uniform-positiveness of the elastic energy are established.

We next present a set of necessary and sufficient conditions for real wave-velocities in the relaxed micromorphic model which is weaker than positivity of the energy, as the strong ellipticity condition is with respect to positive definiteness of the energy in the case of linear elasticity. Then, for didactic purposes, we repeat the analysis for isotropic linear elasticity in order to see relations of our necessary and sufficient condition to the strong ellipticity condition in linear elasticity. Similarly, we discuss micropolar elasticity and establish necessary and sufficient conditions for real wave propagation. We finally show that strong ellipticity in micropolar and micromorphic models is **not** sufficient for having real wave velocities, when dealing with plane waves.

2 The relaxed micromorphic model

The relaxed micromorphic model has been recently introduced into continuum mechanics in [40]. In subsequent works [20,26–28], the model has shown its wider applicability compared to the classical Mindlin-Eringen micromorphic model in diverse areas [11, 17, 31, 32].

The dynamic relaxed micromorphic model counts only 8 constitutive parameters in the (simplified) isotropic case, namely 5 elastic moduli $\mu_e, \lambda_e, \mu_{\text{micro}}, \lambda_{\text{micro}}, \mu_c$ [Pa], one characteristic length L_c [m], the average macroscopic inertia ρ [kg], and the micro-inertia η [kg/m]. The simplification consists in assuming one scalar micro-inertia parameter η and a uni-constant curvature expression. The characteristic length L_c is intrinsically related to non-local effects due to the fact that it weights a suitable combination of first order space derivatives of the microdistortion tensor in the strain energy density (1). For a general presentation of the features of the relaxed micromorphic model in the anisotropic setting, we refer to [3].

2.1 Elastic energy density

The relaxed micromorphic model couples the macroscopic displacement $u \in \mathbb{R}^3$ and an affine substructure deformation attached at each macroscopic point encoded by the **micro-distortion** field $P \in \mathbb{R}^{3 \times 3}$. Our novel relaxed micromorphic model endows Mindlin-Eringen's representation of linear micromorphic models with the second order **dislocation density tensor** $\alpha = -\text{Curl}P$ instead of the full gradient ∇P .⁷ In the isotropic hyperelastic case the elastic energy density reads

$$\begin{aligned}
W &= \mu_e \|\text{sym}(\nabla u - P)\|^2 + \frac{\lambda_e}{2} (\text{tr}(\nabla u - P))^2 + \mu_c \|\text{skew}(\nabla u - P)\|^2 \\
&\quad + \mu_{\text{micro}} \|\text{sym}P\|^2 + \frac{\lambda_{\text{micro}}}{2} (\text{tr}P)^2 + \frac{\mu_e L_c^2}{2} \|\text{Curl}P\|^2 \\
&= \underbrace{\mu_e \|\text{dev sym}(\nabla u - P)\|^2 + \frac{2\mu_e + 3\lambda_e}{3} (\text{tr}(\nabla u - P))^2}_{\text{isotropic elastic - energy}} + \underbrace{\mu_c \|\text{skew}(\nabla u - P)\|^2}_{\text{rotational elastic coupling}} \\
&\quad + \underbrace{\mu_{\text{micro}} \|\text{dev sym}P\|^2 + \frac{2\mu_{\text{micro}} + 3\lambda_{\text{micro}}}{3} (\text{tr}P)^2}_{\text{micro - self - energy}} + \underbrace{\frac{\mu_e L_c^2}{2} \|\text{Curl}P\|^2}_{\text{simplified isotropic curvature}},
\end{aligned} \tag{1}$$

where the parameters and the elastic stress are analogous to the standard Mindlin-Eringen micromorphic model. The model is well-posed in the statical and dynamical case even for zero **Cosserat couple modulus**

⁷The dislocation tensor is defined as $\alpha_{ij} = -(\text{Curl}P)_{ij} = -P_{ih,k}\epsilon_{jkh}$, where ϵ is the Levi-Civita tensor.

$\mu_c = 0$, see [14, 39]. In that case, it is non-redundant in the sense of [47]. Well-posedness results for the static and dynamical cases have been provided in [40] making decisive use of recently established new coercive inequalities, generalizing Korn's inequality to incompatible tensor fields [4, 5, 43–45].

Decisive for the relaxed micromorphic formulation is the definition of the elastic energy in terms of suitable strain tensors. Since ∇u is the macroscopic displacement gradient and P is the micro-distortion there appears the possibility to use the non-symmetric relative (elastic) strain tensor $\nabla u - P$ as basic building block in the energy. Using the Cartan-Lie orthogonal orthogonal decomposition we may introduce:

$$\mu_e \|\operatorname{dev sym} (\nabla u - P)\|^2 + \frac{2\mu_e + 3\lambda_e}{3} (\operatorname{tr} (\nabla u - P))^2 + \mu_c \|\operatorname{skew} (\nabla u - P)\|^2. \quad (2)$$

The microstructure contribution based on P alone is restricted, by infinitesimal frame-indifference to

$$\mu_{\text{micro}} \|\operatorname{dev sym} P\|^2 + \frac{2\mu_{\text{micro}} + 3\lambda_{\text{micro}}}{3} (\operatorname{tr} P)^2 + \frac{\mu_e L_c^2}{2} \|\operatorname{Curl} P\|^2. \quad (3)$$

Strict positive definiteness of the potential energy is equivalent to the following simple relations for the introduced parameters [40]:

$$\mu_e > 0, \quad \mu_c > 0, \quad 2\mu_e + 3\lambda_e > 0, \quad \mu_{\text{micro}} > 0, \quad 2\mu_{\text{micro}} + 3\lambda_{\text{micro}} > 0, \quad L_c > 0. \quad (4)$$

As for the kinetic energy density, we consider that it takes the following (simplified) form

$$J = \frac{\rho}{2} \|u_{,t}\|^2 + \underbrace{\frac{\eta}{2} \|P_{,t}\|^2}_{\text{simplified micro - inertia}}, \quad (5)$$

where $\rho > 0$ is the value of the averaged macroscopic mass density of the considered material, while $\eta > 0$ is its micro-inertia density.

For very large sample sizes, a scaling argument shows easily that the relative characteristic length scale L_c of the micromorphic model must vanish. Therefore, we have a way of comparing a classical first gradient formulation with the relaxed micromorphic model and to offer an a priori relation between the microscopic parameters $\lambda_e, \lambda_{\text{micro}}, \mu_e, \mu_{\text{micro}}$ on the one side and the resulting macroscopic parameters $\lambda_{\text{macro}}, \mu_{\text{macro}}$ on the other side [3, 35, 38]. We have

$$(2\mu_{\text{macro}} + 3\lambda_{\text{macro}}) = \frac{(2\mu_e + 3\lambda_e)(2\mu_{\text{micro}} + 3\lambda_{\text{micro}})}{(2\mu_e + 3\lambda_e) + (2\mu_{\text{micro}} + 3\lambda_{\text{micro}})}, \quad \mu_{\text{macro}} = \frac{\mu_e \mu_{\text{micro}}}{\mu_e + \mu_{\text{micro}}}, \quad (6)$$

where $\mu_{\text{macro}}, \lambda_{\text{macro}}$ are the moduli obtained for $L_c \rightarrow 0$.

For future use we define the elastic bulk modulus κ_e , the microscopic bulk modulus κ_{micro} and the macroscopic bulk modulus κ_{macro} , respectively:

$$\kappa_e = \frac{2\mu_e + 3\lambda_e}{3}, \quad \kappa_{\text{micro}} = \frac{2\mu_{\text{micro}} + 3\lambda_{\text{micro}}}{3}, \quad \kappa_{\text{macro}} = \frac{2\mu_{\text{macro}} + 3\lambda_{\text{macro}}}{3}. \quad (7)$$

In terms of these moduli, strict positive-definiteness of the energy is equivalent to:

$$\mu_e > 0, \quad \mu_c > 0, \quad \kappa_e > 0, \quad \mu_{\text{micro}} > 0, \quad \kappa_{\text{micro}} > 0, \quad L_c > 0. \quad (8)$$

If strict positive-definiteness (8) holds we can write the macroscopic consistency conditions as:

$$\kappa_{\text{macro}} = \frac{\kappa_e \kappa_{\text{micro}}}{\kappa_e + \kappa_{\text{micro}}}, \quad \mu_{\text{macro}} = \frac{\mu_e \mu_{\text{micro}}}{\mu_e + \mu_{\text{micro}}}, \quad (9)$$

and, again under condition (8)

$$\kappa_e = \frac{\kappa_{\text{micro}} \kappa_{\text{macro}}}{\kappa_{\text{micro}} - \kappa_{\text{macro}}}, \quad \kappa_{\text{micro}} = \frac{\kappa_e \kappa_{\text{macro}}}{\kappa_e - \kappa_{\text{macro}}}, \quad \mu_e = \frac{\mu_{\text{micro}} \mu_{\text{macro}}}{\mu_{\text{micro}} - \mu_{\text{macro}}}, \quad \mu_{\text{micro}} = \frac{\mu_e \mu_{\text{macro}}}{\mu_e - \mu_{\text{macro}}}. \quad (10)$$

Here, strict positivity (8) implies that:

$$\begin{aligned} \kappa_e + \kappa_{\text{micro}} > 0, \quad \mu_e + \mu_{\text{micro}} > 0, \quad \kappa_e > \kappa_{\text{macro}}, \quad \kappa_{\text{micro}} > \kappa_{\text{macro}}, \quad \mu_e > \mu_{\text{macro}}, \\ \mu_{\text{micro}} > \mu_{\text{macro}}. \end{aligned} \quad (11)$$

Since it is useful in what follows we explicitly remark that:

$$2\mu_e + \lambda_e = \frac{4}{3}\mu_e + \frac{2\mu_e + 3\lambda_e}{3} = \frac{4}{3}\mu_e + \kappa_e = \frac{4\mu_e + 3\kappa_e}{3}, \quad 2\mu_{\text{micro}} + \lambda_{\text{micro}} = \frac{4\mu_{\text{micro}} + 3\kappa_{\text{micro}}}{3}. \quad (12)$$

With these relations, it is easy to show how $\mu_e > 0$ and $\kappa_e > 0$ imply $2\mu_e + \lambda_e > 0$. Moreover, as shown in the appendix (equations (94) and (95)), we note here that if only $\mu_e + \mu_{\text{micro}} > 0$ and $\kappa_e + \kappa_{\text{micro}} > 0$, then the macroscopic parameters are less or equal than respective microscopic parameters, namely:

$$\kappa_e \geq \kappa_{\text{macro}}, \quad \kappa_{\text{micro}} \geq \kappa_{\text{macro}}, \quad \mu_e \geq \mu_{\text{macro}}, \quad \mu_{\text{micro}} \geq \mu_{\text{macro}}, \quad (13)$$

and moreover the following inequalities are satisfied:

$$2\mu_e + \lambda_e \geq 2\mu_{\text{macro}} + \lambda_{\text{macro}}, \quad 2\mu_{\text{micro}} + \lambda_{\text{micro}} \geq 2\mu_{\text{macro}} + \lambda_{\text{macro}}, \quad \frac{4\mu_{\text{macro}} + 3\kappa_e}{3} \geq 2\mu_{\text{macro}} + \lambda_{\text{macro}}. \quad (14)$$

Note that the Cosserat couple modulus μ_c [36] does not appear in the introduced scale between micro and macro.

2.2 Dynamic formulation

The dynamical formulation is obtained defining a joint Hamiltonian and assuming stationary action. The dynamical equilibrium equations are:

$$\begin{aligned} \rho u_{,tt} &= \text{Div} [2\mu_e \text{sym} (\nabla u - P) + 2\mu_c \text{skew} (\nabla u - P) + \lambda_e \text{tr} (\nabla u - P) \mathbf{1}], \\ \eta P_{,tt} &= -\mu_e L_c^2 \text{Curl Curl } P + 2\mu_e \text{sym} (\nabla u - P) + 2\mu_c \text{skew} (\nabla u - P) \\ &\quad + \lambda_e \text{tr} (\nabla u - P) \mathbf{1} - [2\mu_{\text{micro}} \text{sym} P + \lambda_{\text{micro}} \text{tr} (P) \mathbf{1}]. \end{aligned} \quad (15)$$

We note here that the presence of the $\text{Curl } P$ in the energy generates a non-local term $\text{Curl Curl } P$ in the equation of motion, while the possibility of band-gaps is still present, see [26]. The presence of the $\text{Curl } P$ term is essential to simultaneously allow for the description of non localities and band gap in an enriched continuum mechanics framework.

Sufficiently far from a source, dynamic wave solutions may be treated as planar waves. Therefore, we now want to study harmonic solutions traveling in an infinite domain for the differential system (15). To do so, we define:

$$\begin{aligned} P^S &:= \frac{1}{3} \text{tr} (P), & P_{[ij]} &:= (\text{skew } P)_{ij} = \frac{1}{2} (P_{ij} - P_{ji}), \\ P^D &:= P_{11} - P^S, & P_{(ij)} &:= (\text{sym } P)_{ij} = \frac{1}{2} (P_{ij} + P_{ji}), \\ P^V &:= P_{22} - P_{33} \end{aligned} \quad (16)$$

and we introduce the unknown vectors

$$\underbrace{v_1 = (u_1, P^D, P^S)}_{\text{longitudinal}} \quad \underbrace{v_\tau = (u_\tau, P_{(1\tau)}, P_{[1\tau]})}_{\text{transversal}}, \quad \tau = 2, 3, \quad \underbrace{v_4 = (P_{(23)}, P_{[23]}, P^V)}_{\text{uncoupled}}. \quad (17)$$

The definition of the unknown vectors was made considering the coupling of the variables in the equations of motion, see [9, 20, 21, 23–28]. More particularly, it has been shown in these previous works that three sets of equations can be isolated: one involving only longitudinal quantities, one involving only transverse quantities and one of three completely uncoupled equations. We suppose that the space dependence of all

introduced kinematic fields are limited to a direction defined by a unit vector $\tilde{\xi} \in \mathbb{R}^3$, which is the direction of propagation of the wave and which is assumed given. Hence, we look for solutions of (15) in the form:

$$v_1 = \beta^1 e^{i(k\langle\tilde{\xi}, x\rangle_{\mathbb{R}^3} - \omega t)}, \quad v_\tau = \beta^\tau e^{i(k\langle\tilde{\xi}, x\rangle_{\mathbb{R}^3} - \omega t)}, \quad \tau = 2, 3, \quad v_4 = \beta^4 e^{i(k\langle\tilde{\xi}, x\rangle_{\mathbb{R}^3} - \omega t)}. \quad (18)$$

where $\beta^1 = (\beta_1^1, \beta_2^1, \beta_3^1)^T \in \mathbb{C}^3$, $\beta^\tau = (\beta_1^\tau, \beta_2^\tau, \beta_3^\tau)^T \in \mathbb{C}^3$ and $\beta^4 = (\beta_1^4, \beta_2^4, \beta_3^4)^T \in \mathbb{C}^3$ are the unknown amplitudes of the considered waves, \mathbb{C}^3 is the space of complex constant three-dimensional vectors⁸, k is the wavenumber and ω is the wave-frequency. Since our formulation is isotropic, we can, without loss of generality, specify the propagation direction $\tilde{\xi} = e_1$. Then $X = \langle e_1, x \rangle_{\mathbb{R}^3} = x_1$, and we obtain that the space dependence of all introduced kinematic fields are limited to the component X which is now the direction of propagation of the wave⁹. This means that we look for solutions in the form:

$$v_1 = \beta^1 e^{i(kX - \omega t)}, \quad v_\tau = \beta^\tau e^{i(kX - \omega t)}, \quad \tau = 2, 3, \quad v_4 = \beta^4 e^{i(kX - \omega t)}, \quad (19)$$

Replacing these expressions in equations (15), it is possible to express the system (see [26, 27]) as:

$$A_1 \cdot \beta^1 = 0, \quad A_\tau \cdot \beta^\tau = 0, \quad \tau = 2, 3, \quad A_4 \cdot \beta^4 = 0, \quad (20)$$

with

$$A_1(\omega, k) = \begin{pmatrix} -\omega^2 + c_p^2 k^2 & i k 2\mu_e / \rho & i k (2\mu_e + 3\lambda_e) / \rho \\ -i k \frac{4}{3} \mu_e / \eta & -\omega^2 + \frac{1}{3} k^2 c_m^2 + \omega_s^2 & -\frac{2}{3} k^2 c_m^2 \\ -\frac{1}{3} i k (2\mu_e + 3\lambda_e) / \eta & -\frac{1}{3} k^2 c_m^2 & -\omega^2 + \frac{2}{3} k^2 c_m^2 + \omega_p^2 \end{pmatrix}, \quad (21)$$

$$A_2(\omega, k) = A_3(\omega, k) = \begin{pmatrix} -\omega^2 + k^2 c_s^2 & i k 2\mu_e / \rho & -i k \frac{2}{\rho} \omega_r^2 \\ -i k \mu_e / \eta, & -\omega^2 + \frac{c_m^2}{2} k^2 + \omega_s^2 & \frac{c_m^2}{2} k^2 \\ \frac{i}{2} \omega_r^2 k & \frac{c_m^2}{2} k^2 & -\omega^2 + \frac{c_m^2}{2} k^2 + \omega_r^2 \end{pmatrix}, \quad (22)$$

$$A_4(\omega, k) = \begin{pmatrix} -\omega^2 + c_m^2 k^2 + \omega_s^2 & 0 & 0 \\ 0 & -\omega^2 + c_m^2 k^2 + \omega_r^2 & 0 \\ 0 & 0 & -\omega^2 + c_m^2 k^2 + \omega_s^2 \end{pmatrix}. \quad (23)$$

Here, we have defined:

$$\boxed{\begin{aligned} c_m &= \sqrt{\frac{\mu_e L_c^2}{\eta}}, & c_s &= \sqrt{\frac{\mu_e + \mu_c}{\rho}}, & c_p &= \sqrt{\frac{2\mu_e + \lambda_e}{\rho}}, \\ \omega_s &= \sqrt{\frac{2(\mu_e + \mu_{\text{micro}})}{\eta}}, & \omega_p &= \sqrt{\frac{(2\mu_e + 3\lambda_e) + (2\mu_{\text{micro}} + 3\lambda_{\text{micro}})}{\eta}}, & \omega_r &= \sqrt{\frac{2\mu_c}{\eta}}, \\ \omega_l &= \sqrt{\frac{2\mu_{\text{micro}} + \lambda_{\text{micro}}}{\eta}}, & \omega_t &= \sqrt{\frac{\mu_{\text{micro}}}{\eta}}. \end{aligned}}$$

⁸Here, we understand that having found the (in general, complex) solutions of (19) only the real or imaginary parts separately constitute actual wave solutions which can be observed in reality.

⁹In an isotropic model it is clear that there is no direction dependence. More specifically, let us consider an arbitrary direction $\tilde{\xi} \in \mathbb{R}^3$. Now we consider an orthogonal spatial coordinate change $Q \cdot e_1 = \tilde{\xi}$ with $Q \in \text{SO}(3)$. In the rotated variables, the ensuing system of PDE's (15) is form-invariant, see [33].

Let us next define the diagonal matrix:

$$\text{diag}_1 = \begin{pmatrix} \sqrt{\rho} & 0 & 0 \\ 0 & i\frac{\sqrt{6\eta}}{2} & 0 \\ 0 & 0 & i\sqrt{3\eta} \end{pmatrix}. \quad (24)$$

Considering $\gamma = \text{diag}_1 \cdot \beta$ and the matrix $\bar{A}_1(\omega, k) = \text{diag}_1 \cdot A_1(\omega, k) \cdot \text{diag}_1^{-1}$, it is possible to formulate the problem (20) equivalently as¹⁰:

$$\bar{A}_1 \cdot \gamma = \begin{pmatrix} -\omega^2 + c_p^2 k^2 & \frac{2\sqrt{6}}{3} k \mu_e / \sqrt{\rho\eta} & \frac{\sqrt{3}}{3} k (2\mu_e + 3\lambda_e) / \sqrt{\rho\eta} \\ \frac{2\sqrt{6}}{3} k \mu_e / \sqrt{\rho\eta} & -\omega^2 + \frac{1}{3} k^2 c_m^2 + \omega_s^2 & -\frac{\sqrt{2}}{3} k^2 c_m^2 \\ \frac{\sqrt{3}}{3} k (2\mu_e + 3\lambda_e) / \sqrt{\rho\eta} & -\frac{\sqrt{2}}{3} k^2 c_m^2 & -\omega^2 + \frac{2}{3} k^2 c_m^2 + \omega_p^2 \end{pmatrix} \begin{pmatrix} \gamma_1 \\ \gamma_2 \\ \gamma_3 \end{pmatrix} = 0. \quad (25)$$

Analogously considering

$$\text{diag}_2 = \begin{pmatrix} \sqrt{\rho} & 0 & 0 \\ 0 & i\sqrt{2\eta} & 0 \\ 0 & 0 & i\sqrt{2\eta} \end{pmatrix}, \quad (26)$$

it is possible to obtain $\bar{A}_2(\omega, k) = \bar{A}_3(\omega, k) = \text{diag}_2 \cdot A_2(\omega, k) \cdot \text{diag}_2^{-1}$

$$\bar{A}_2(\omega, k) = \bar{A}_3(\omega, k) = \begin{pmatrix} -\omega^2 + k^2 c_s^2 & k \sqrt{2} \mu_e / \sqrt{\rho\eta} & -k \sqrt{2} \mu_c / \sqrt{\rho\eta}, \\ k \sqrt{2} \mu_e / \sqrt{\rho\eta}, & -\omega^2 + \frac{c_m^2}{2} k^2 + \omega_s^2 & \frac{c_m^2}{2} k^2 \\ -k \sqrt{2} \mu_c / \sqrt{\rho\eta} & \frac{c_m^2}{2} k^2 & -\omega^2 + \frac{c_m^2}{2} k^2 + \omega_r^2 \end{pmatrix}. \quad (27)$$

In order to have non-trivial solutions of the algebraic systems (20), one must impose that

$$\det \bar{A}_1(\omega, k) = 0, \quad \det \bar{A}_2(\omega, k) = \det \bar{A}_3(\omega, k) = 0, \quad \det A_4(\omega, k) = 0, \quad (28)$$

the solution of which allow us to determine the so-called dispersion relations $\omega = \omega(k)$ for the longitudinal and transverse waves in the relaxed micromorphic continuum, see Figure 1¹¹. The solutions of the eigenvalue problem obtained via the proposed decomposition are the same as the ones obtained via the standard formulation shown in the Appendix 7.2 with the full 12×12 matrix, for more details see [9]. For estimates on the isotropic moduli, we refer [20, 21] and, for a comparison with other micromorphic models, to [24, 25].

For solutions $\omega = \omega(k)$ of (28) we define the

$$\text{phase velocity: } v = \frac{\omega}{k}, \quad \text{group velocity: } \frac{d\omega(k)}{dk}. \quad (29)$$

Real wave numbers $k \in \mathbb{R}$ correspond to propagating waves, while complex values of k are associated with waves whose amplitude either grows or decays along the coordinate X . In linear elasticity, phase velocity and group velocity coincide since there is no dispersion and both are real, see section 3.

Since in this paper we are only interested in real k (outside the band gap region), the wave velocity (phase velocity) is real if and only if ω is real.

¹⁰It is possible to face the problem in two more equivalent ways. The first one is to consider from the start that the amplitudes of the micro-distortion field are multiplied by the imaginary unit i , i.e. $\beta = (\beta_1, i\beta_2, i\beta_3)^T \in \mathbb{C}^3$, as done in [31, p. 24, eq. 8.6].

Doing so, we obtaining a real matrix that can be symmetrized with $\text{diag}_1 = \begin{pmatrix} \sqrt{\rho} & 0 & 0 \\ 0 & \frac{\sqrt{6\eta}}{2} & 0 \\ 0 & 0 & \sqrt{3\eta} \end{pmatrix}$. On the other hand, it is

also possible to consider from the beginning $\beta = (\sqrt{\rho}\beta_1, i\frac{\sqrt{6\eta}}{2}\beta_2, i\sqrt{3\eta}\beta_3)^T \in \mathbb{C}^3$ obtaining directly a real symmetric matrix.

¹¹The formal limit $\eta \rightarrow +\infty$ shows no dispersion at all giving two pseudo-acoustic linear curves, longitudinal and transverse with slopes $c_p = \sqrt{(2\mu_e + \lambda_e)/\rho}$ and $c_s = \sqrt{(\mu_e + \mu_c)/\rho}$, respectively.




Figure 1: Dispersion relations $\omega = \omega(k)$ [20] for the **relaxed micromorphic model** with non-vanishing Cosserat couple modulus $\mu_c > 0$. Uncoupled waves (a), longitudinal waves (b) and transverse waves (c). TRO: transverse rotational optic, TSO: transverse shear optic, TCVO: transverse constant-volume optic, LA: longitudinal acoustic, LO₁-LO₂: 1st and 2nd longitudinal optic, TA: transverse acoustic, TO₁-TO₂: 1st and 2nd transverse optic.

Since ω^2 appears on the diagonal only, the problem (28) can be analogously expressed as an eigenvalue-problem:

$$\begin{aligned} \det (B_1(k) - \omega^2 \mathbf{1}) &= 0, & \det (B_2(k) - \omega^2 \mathbf{1}) &= 0, \\ \det (B_3(k) - \omega^2 \mathbf{1}) &= 0, & \det (B_4(k) - \omega^2 \mathbf{1}) &= 0, \end{aligned} \quad (30)$$

where

$$B_1(k) = \begin{pmatrix} c_p^2 k^2 & \frac{2\sqrt{6}}{3} k \mu_e / \sqrt{\rho\eta} & \frac{\sqrt{3}}{3} k (2\mu_e + 3\lambda_e) / \sqrt{\rho\eta} \\ \frac{2\sqrt{6}}{3} k \mu_e / \sqrt{\rho\eta} & \frac{1}{3} k^2 c_m^2 + \omega_s^2 & -\frac{\sqrt{2}}{3} k^2 c_m^2 \\ \frac{\sqrt{3}}{3} k (2\mu_e + 3\lambda_e) / \sqrt{\rho\eta} & -\frac{\sqrt{2}}{3} k^2 c_m^2 & +\frac{2}{3} k^2 c_m^2 + \omega_p^2 \end{pmatrix}, \quad (31)$$

$$B_2(k) = B_3(k) = \begin{pmatrix} k^2 c_s^2 & k \sqrt{2} \mu_e / \sqrt{\rho\eta} & -k \sqrt{2} \mu_c / \sqrt{\rho\eta}, \\ k \sqrt{2} \mu_e / \sqrt{\rho\eta}, & \frac{c_m^2}{2} k^2 + \omega_s^2 & \frac{c_m^2}{2} k^2 \\ -k \sqrt{2} \mu_c / \sqrt{\rho\eta} & \frac{c_m^2}{2} k^2 & \frac{c_m^2}{2} k^2 + \omega_r^2 \end{pmatrix}, \quad (32)$$

$$B_4(k) = \begin{pmatrix} c_m^2 k^2 + \omega_s^2 & 0 & 0 \\ 0 & c_m^2 k^2 + \omega_r^2 & 0 \\ 0 & 0 & c_m^2 k^2 + \omega_s^2 \end{pmatrix}. \quad (33)$$

Note that $B_1(k)$, $B_2(k)$, $B_3(k)$ and $B_4(k)$ are real symmetric matrices and therefore the resulting eigenvalues ω^2 are real. Obtaining real wave velocities is tantamount to having $\omega^2 \geq 0$ for all solutions of (30).

2.3 Necessary and sufficient conditions for real wave propagation

We will show next that all the eigenvalues ω^2 of $B_1(k)$, $B_2(k)$ and $B_3(k)$ are real and positive for every $k \neq 0$ and non-negative for $k = 0$ provided certain conditions on the material coefficients are satisfied. Sylvester's criterion states that a Hermitian matrix M is positive-definite if and only if the leading principal minors are positive [15]. For the matrix B_1 the three principal minors are:

$$(B_1)_{11} = \frac{2\mu_e + \lambda_e}{\rho}, \quad (34)$$

$$(\text{Cof}(B_1))_{33} = \frac{k^2}{3\eta\rho} [6(2\mu_e + \lambda_e)\mu_{\text{micro}} + 6\mu_e\kappa_e + (2\mu_e + \lambda_e)\mu_e L_c^2 k^2] \quad (35)$$

$$= \frac{k^2}{3\eta\rho} [2(4\mu_{\text{macro}} + 3\kappa_e)(\mu_e + \mu_{\text{micro}}) + (2\mu_e + \lambda_e)\mu_e L_c^2 k^2],$$

$$\det(B_1) = \frac{k^2}{\eta^2\rho} \left[6\kappa_e\kappa_{\text{micro}}(\mu_e + \mu_{\text{micro}}) + 8\mu_e\mu_{\text{micro}}(\kappa_e + \kappa_{\text{micro}}) + (2\mu_e + \lambda_e)(2\mu_{\text{micro}} + \lambda_{\text{micro}})\mu_e L_c^2 k^2 \right] \quad (36)$$

$$= \frac{k^2}{\eta^2\rho} \left[6(\kappa_e + \kappa_{\text{micro}})(\mu_e + \mu_{\text{micro}})(2\mu_{\text{macro}} + \lambda_{\text{macro}}) + (2\mu_e + \lambda_e)(2\mu_{\text{micro}} + \lambda_{\text{micro}})\mu_e L_c^2 k^2 \right].$$

The three principal minors of B_1 are clearly positive for $k \neq 0$ if¹²:

$$\begin{aligned} \mu_e > 0, & \quad \mu_{\text{micro}} > 0, & \quad \kappa_e + \kappa_{\text{micro}} > 0, & \quad 2\mu_{\text{macro}} + \lambda_{\text{macro}} > 0, \\ 4\mu_{\text{macro}} + 3\kappa_e > 0, & \quad 2\mu_e + \lambda_e > 0, & \quad 2\mu_{\text{micro}} + \lambda_{\text{micro}} > 0. \end{aligned} \quad (37)$$

Similarly, for the matrix B_2 the three principal minors are:

$$(B_2)_{11} = \frac{\mu_e + \mu_c}{\rho}, \quad (38)$$

$$(\text{Cof}(B_2))_{33} = \frac{k^2}{2\eta\rho} \left[4(\mu_e\mu_c + \mu_{\text{micro}}(\mu_e + \mu_c) + (\mu_e + \mu_c)\mu_e L_c^2 k^2) \right]. \quad (39)$$

$$\det(B_2) = \frac{k^2}{\eta^2\rho} \left[4\mu_{\text{micro}}\mu_c\mu_e + (\mu_e + \mu_c)\mu_{\text{micro}}\mu_e L_c^2 k^2 \right]. \quad (40)$$

For the matrix $B_2(k) = B_3(k)$, considering positive η , ρ and separating terms in the brackets by looking at large and small values of k , we can state **necessary** and **sufficient** conditions for strict positive-definiteness of $B_2(k)$ at arbitrary $k \neq 0$:

$$\mu_e > 0, \quad \mu_{\text{micro}} > 0, \quad \mu_c \geq 0. \quad (41)$$

Since $B_4(k)$ is diagonal, it easy to show that positive definiteness is tantamount to the set of **necessary** and **sufficient** conditions for $k \neq 0$:

$$\mu_e > 0, \quad \mu_e + \mu_{\text{micro}} > 0, \quad \mu_c \geq 0. \quad (42)$$

On the other hand, considering the case $k = 0$, we obtain that the matrices reduce to:

$$B_1(0) = \begin{pmatrix} 0 & 0 & 0 \\ 0 & \omega_s^2 & 0 \\ 0 & 0 & \omega_p^2 \end{pmatrix}, \quad B_2(0) = B_3(0) = \begin{pmatrix} 0 & 0 & 0 \\ 0 & \omega_s^2 & 0 \\ 0 & 0 & \omega_r^2 \end{pmatrix}, \quad B_4(0) = \begin{pmatrix} \omega_s^2 & 0 & 0 \\ 0 & \omega_r^2 & 0 \\ 0 & 0 & \omega_s^2 \end{pmatrix}. \quad (43)$$

Since the matrices are diagonal for $k = 0$, it easy to show that positive semi-definiteness is tantamount to the set of **necessary** and **sufficient** conditions :

$$\mu_e \geq 0, \quad \mu_e + \mu_{\text{micro}} \geq 0, \quad \mu_c \geq 0, \quad \kappa_e + \kappa_{\text{micro}} \geq 0. \quad (44)$$

¹²We note here that $4\mu_{\text{macro}} + 3\kappa_e > 0 \iff 2\mu_e + \lambda_e > \frac{4}{3}(\mu_e - \mu_{\text{macro}}) \iff 2\mu_{\text{macro}} + \lambda_{\text{macro}} > \kappa_{\text{macro}} - \kappa_e$. Furthermore, if $\mu_e + \mu_{\text{micro}} > 0$ and $\kappa_e + \kappa_{\text{micro}} > 0$, we have $3(2\mu_e + \lambda_e) \geq 4\mu_{\text{macro}} + 3\kappa_e \geq 3(2\mu_{\text{macro}} + \lambda_{\text{macro}})$, see Appendix.

Hence, we can state a simple **sufficient** condition for real wave velocities for all real k :

$$\begin{aligned} \mu_e > 0, & \quad \mu_{\text{micro}} > 0, & \quad \kappa_e + \kappa_{\text{micro}} > 0, & \quad 2\mu_{\text{macro}} + \lambda_{\text{macro}} > 0, \\ 4\mu_{\text{macro}} + 3\kappa_e > 0, & \quad 2\mu_e + \lambda_e > 0, & \quad 2\mu_{\text{micro}} + \lambda_{\text{micro}} > 0. \end{aligned} \quad (45)$$

In order to see a set of global necessary conditions for positivity at arbitrary $k \neq 0$ we consider first large and small values of $k \neq 0$ separately. For $k \rightarrow +\infty$ we must have:

$$2\mu_e + \lambda_e > 0, \quad (2\mu_e + \lambda_e)\mu_e L_c^2 > 0, \quad (2\mu_e + \lambda_e)(2\mu_{\text{micro}} + \lambda_{\text{micro}})\mu_e L_c^2 > 0, \quad (46)$$

or analogously:

$$2\mu_e + \lambda_e > 0, \quad \mu_e L_c^2 > 0, \quad 2\mu_{\text{micro}} + \lambda_{\text{micro}} > 0, \quad (47)$$

while for $k \rightarrow 0$ we must have:

$$2\mu_e + \lambda_e > 0, \quad (4\mu_{\text{macro}} + 3\kappa_e)(\mu_e + \mu_{\text{micro}}) > 0, \quad (\kappa_e + \kappa_{\text{micro}})(\mu_e + \mu_{\text{micro}})(2\mu_{\text{macro}} + \lambda_{\text{macro}}) > 0. \quad (48)$$

Since from (41) we have necessarily $\mu_e > 0$, $\mu_{\text{micro}} > 0$, and from (44) we get $\kappa_e + \kappa_{\text{micro}} \geq 0$ and considering together the two limits for k we obtain the necessary condition:

$$\begin{aligned} 2\mu_e + \lambda_e > 0, & \quad 2\mu_{\text{micro}} + \lambda_{\text{micro}} > 0, & \quad 4\mu_{\text{macro}} + 3\kappa_e > 0, & \quad \kappa_e + \kappa_{\text{micro}} > 0, \\ \mu_e > 0, & \quad \mu_{\text{micro}} > 0, & \quad \mu_c \geq 0, & \quad 2\mu_{\text{macro}} + \lambda_{\text{macro}} > 0. \end{aligned} \quad (49)$$

Inspection shows that (49) is our proposed sufficient condition (37). From $\mu_e > 0$ and $\mu_{\text{micro}} > 0$, it follows that $\mu_{\text{macro}} > 0$. Therefore condition (49) is **necessary** and **sufficient**. We have shown our main proposition:

Proposition (real wave velocities). *The dynamic relaxed micromorphic model (eq. (15)) admits real planar waves if and only if*

$$\begin{aligned} \mu_c \geq 0, & \quad \mu_e > 0, & \quad 2\mu_e + \lambda_e > 0, \\ & \quad \mu_{\text{micro}} > 0, & \quad 2\mu_{\text{micro}} + \lambda_{\text{micro}} > 0, \\ & \quad (\mu_{\text{macro}} > 0), & \quad 2\mu_{\text{macro}} + \lambda_{\text{macro}} > 0, \\ & \quad \kappa_e + \kappa_{\text{micro}} > 0, & \quad 4\mu_{\text{macro}} + 3\kappa_e > 0. \end{aligned} \quad (50) \quad \blacksquare$$

In (50) the requirement $\mu_{\text{macro}} > 0$ is redundant, since it is already assumed that $\mu_e, \mu_{\text{micro}} > 0$. It is clear that positive definiteness of the elastic energy (4) implies (50). We remark that, as shown in the appendix 7.1, the set of inequalities (50) is already implied by:

$$\boxed{\mu_e > 0, \quad \mu_{\text{micro}} > 0, \quad \mu_c \geq 0, \quad \kappa_e + \kappa_{\text{micro}} > 0, \quad 2\mu_{\text{macro}} + \lambda_{\text{macro}} > 0.} \quad (51)$$

Letting finally $\mu_{\text{micro}} \rightarrow +\infty$ and $\kappa_{\text{micro}} \rightarrow +\infty$ (or $\mu_{\text{micro}} \rightarrow +\infty$ and $\lambda_{\text{micro}} > \text{const.}$) generates the limit condition for real wave velocities ($\mu_e \rightarrow \mu_{\text{macro}}$)

$$\mu_{\text{macro}} > 0, \quad \mu_c \geq 0, \quad 2\mu_{\text{macro}} + \lambda_{\text{macro}} > 0. \quad (52)$$

which coincides, up to μ_c , with the strong ellipticity condition in isotropic linear elasticity, see section 3, and it coincides fully with the condition for real wave velocities in micropolar elasticity, see section 4. A condition similar to (52) can be found in [31, eq. 8.14 p. 26] where Mindlin requires that $\mu_{\text{macro}} > 0$, $2\mu_{\text{macro}} + \lambda_{\text{macro}} > 0$ ¹³ (in our notation) which are obtained from the requirement of positive **group velocity** at $k = 0$

$$\frac{d\omega_{\text{acoustic, long}}(0)}{dk} > 0, \quad \frac{d\omega_{\text{acoustic, trans}}(0)}{dk} > 0. \quad (53)$$

¹³Mindlin explains that such parameters “are less than those that would be calculated from the strain-stiffnesses [of the unit cell]. This phenomenon is due to the compliance of the unit cell and has been found in a theory of crystal lattices by Gazis and Wallis [12]”.

Let us emphasize that our method is not easily generalized to two immediate extensions. First, one could be interested in the isotropic relaxed micromorphic model with weighted inertia contributions and weighted curvatures [9]. Second, one could be interested in the anisotropic setting [3]. In the second cases the block-structure of the problem will be lost and one has to deal with the full 12×12 case, see equation (115) in the Appendix. Nonetheless, we expect positive-definiteness to always imply real wave propagation.

In [9] we show that the tangents of the acoustic branches in $k = 0$ in the dispersion curves are

$$c_t = \frac{d\omega_{\text{acoustic, long}}(0)}{dk} = \sqrt{\frac{2\mu_{\text{macro}} + \lambda_{\text{macro}}}{\rho}}, \quad c_t = \frac{d\omega_{\text{acoustic, trans}}(0)}{dk} = \sqrt{\frac{\mu_{\text{macro}}}{\rho}}. \quad (54)$$

The tangents coincide with the classical linear elastic response if the latter has Lamé constants μ_{macro} and λ_{macro} , as it is shown in Figure 2.




Figure 2: Dispersion relations $\omega = \omega(k)$ for the longitudinal acoustic wave LA, and the transverse acoustic TA in the **relaxed micromorphic model** (a) and in a classical Cauchy medium (b).

3 A comparison: classical isotropic linear elasticity

For classical linear elasticity with isotropic energy density and kinetic energy density:

$$W(\nabla u) = \mu_{\text{macro}} \|\text{sym } \nabla u\|^2 + \frac{\lambda_{\text{macro}}}{2} (\text{tr}(\nabla u))^2, \quad J = \frac{\rho}{2} \|u_t\|^2. \quad (55)$$

The positive definiteness of the energy is equivalent to:

$$\mu_{\text{macro}} > 0, \quad 2\mu_{\text{macro}} + 3\lambda_{\text{macro}} > 0. \quad (56)$$

It is easy to see that our homogenization formula (6) implies (56) under condition of positive definiteness of the relaxed micromorphic model.

The dynamical formulation is obtained defining a joint Hamiltonian and assuming stationary action. The dynamical equilibrium equations are:

$$\rho u_{,tt} = \text{Div} [2\mu_{\text{macro}} \text{sym}(\nabla u) + \lambda_{\text{macro}} \text{tr}(\nabla u) \mathbf{1}]. \quad (57)$$

As before, in our study of wave propagation in micromorphic media we limit ourselves to the case of plane waves traveling in an infinite domain. We suppose that the space dependence of all introduced kinematic fields are limited to a direction defined by a unit vector $\tilde{\xi} \in \mathbb{R}^3$ which is the direction of propagation of the wave. Therefore, we look for solutions of (57) in the form:

$$u(x, t) = \hat{u} e^{i(k\langle \tilde{\xi}, x \rangle_{\mathbb{R}^3} - \omega t)}, \quad \hat{u} \in \mathbb{C}^3, \quad \|\tilde{\xi}\|^2 = 1. \quad (58)$$

Since our formulation is isotropic, we can, without loss of generality, specify the direction $\tilde{\xi} = e_1$. Then $X = \langle e_1, x \rangle_{\mathbb{R}^3} = x_1$, and we obtain:

$$u(x, t) = \hat{u} e^{i(kX - \omega t)}, \quad \hat{u} \in \mathbb{C}^3. \quad (59)$$

With this ansatz, it is possible to write (57) as:

$$A_5(e_1, \omega, k) \cdot \hat{u} = 0 \iff (\mathcal{B}(e_1, k) - \omega^2 \mathbb{1}) \cdot \hat{u} = 0, \quad (60)$$

where:

$$A_5(e_1, \omega, k) = \begin{pmatrix} \frac{2\mu_{\text{macro}} + \lambda_{\text{macro}}}{\rho} k^2 - \omega^2 & 0 & 0 \\ 0 & \frac{\mu_{\text{macro}}}{\rho} k^2 - \omega^2 & 0 \\ 0 & 0 & \frac{\mu_{\text{macro}}}{\rho} k^2 - \omega^2 \end{pmatrix}, \quad (61)$$

$$\mathcal{B}(e_1, k) = \frac{k^2}{\rho} \begin{pmatrix} 2\mu_{\text{macro}} + \lambda_{\text{macro}} & 0 & 0 \\ 0 & \mu_{\text{macro}} & 0 \\ 0 & 0 & \mu_{\text{macro}} \end{pmatrix}. \quad (62)$$

Here, we observe that $A_5(e_1, \omega, k)$ is already diagonal and real. Requesting real wave velocities means $\omega^2 \geq 0$. For $k \neq 0$, this leads to the classical so-called **strong ellipticity condition**:

$$\mu_{\text{macro}} > 0, \quad 2\mu_{\text{macro}} + \lambda_{\text{macro}} > 0, \quad (63)$$

which is implied by positive definiteness of the energy (56).

In classical (linear or nonlinear) elasticity, the condition of real wave propagation (63) is equivalent to **strong ellipticity** and **rank-one convexity**. Indeed, rank-one convexity amounts to set $(\xi = k\tilde{\xi}$ with $\|\xi\|^2 = 1)$:

$$\left. \frac{d^2}{dt^2} \right|_{t=0} W(\nabla u + t\hat{u} \otimes \xi) \geq 0 \iff \langle \mathbb{C}(\hat{u} \otimes \xi), \hat{u} \otimes \xi \rangle_{\mathbb{R}^{3 \times 3}} \geq 0, \quad (64)$$

where \mathbb{C} is the fourth-order elasticity tensor. Condition (64) reads then:

$$0 \leq 2\mu_{\text{macro}} \|\text{sym}(\hat{u} \otimes \xi)\|^2 + \lambda_{\text{macro}} (\text{tr}(\hat{u} \otimes \xi))^2 = \mu_{\text{macro}} \|\hat{u}\|^2 \|\xi\|^2 + (\mu_{\text{macro}} + \lambda_{\text{macro}}) \langle \hat{u}, \xi \rangle_{\mathbb{R}^3}^2.$$

We may express (65) given $\xi \in \mathbb{R}^3$ as a quadratic form in $\hat{u} \in \mathbb{R}^3$, which results in:

$$\mu_{\text{macro}} \|\hat{u}\|^2 \|\xi\|^2 + (\mu_{\text{macro}} + \lambda_{\text{macro}}) \langle \hat{u}, \xi \rangle_{\mathbb{R}^3}^2 = \langle \mathcal{D}(\xi) \hat{u}, \hat{u} \rangle_{\mathbb{R}^3}, \quad (65)$$

where the components of the symmetric and real 3×3 matrix $\mathcal{D}(\xi)$ read

$$\mathcal{D}(\xi) = \begin{pmatrix} (2\mu_{\text{macro}} + \lambda_{\text{macro}})\xi_1^2 + \mu_{\text{macro}}(\xi_2^2 + \xi_3^2) & (\lambda_{\text{macro}} + \mu_{\text{macro}})\xi_1 \xi_2 & \\ (\lambda_{\text{macro}} + \mu_{\text{macro}})\xi_1 \xi_2 & (2\mu_{\text{macro}} + \lambda_{\text{macro}})\xi_2^2 + \mu_{\text{macro}}(\xi_1^2 + \xi_3^2) & \\ (\lambda_{\text{macro}} + \mu_{\text{macro}})\xi_1 \xi_3 & (\lambda_{\text{macro}} + \mu_{\text{macro}})\xi_1 \xi_2 & (\lambda_{\text{macro}} + \mu_{\text{macro}})\xi_1 \xi_2 \\ (\lambda_{\text{macro}} + \mu_{\text{macro}})\xi_1 \xi_3 & & \\ (\lambda_{\text{macro}} + \mu_{\text{macro}})\xi_2 \xi_3 & & \\ 2\mu_{\text{macro}} + \lambda_{\text{macro}} \xi_3^2 + \mu_{\text{macro}}(\xi_1^2 + \xi_2^2) & & \end{pmatrix}. \quad (66)$$

The three principal invariants are independent of the direction ξ due to isotropy and are given by:

$$\begin{aligned} \text{tr}(\mathcal{D}(\xi)) &= \|\xi\|^2 (4\mu_{\text{macro}} + \lambda_{\text{macro}}) = k^2 (4\mu_{\text{macro}} + \lambda_{\text{macro}}), \\ \text{tr}(\text{Cof} \mathcal{D}(\xi)) &= \|\xi\|^4 \mu_{\text{macro}} (5\mu_{\text{macro}} + 2\lambda_{\text{macro}}) = k^4 \mu_{\text{macro}} (5\mu_{\text{macro}} + 2\lambda_{\text{macro}}), \\ \det(\mathcal{D}(\xi)) &= \|\xi\|^6 \mu_{\text{macro}}^2 (2\mu_{\text{macro}} + \lambda_{\text{macro}}) = k^6 \mu_{\text{macro}}^2 (2\mu_{\text{macro}} + \lambda_{\text{macro}}). \end{aligned} \quad (67)$$

Since $\mathcal{D}(\xi)$ is real and symmetric, its eigenvalues are real. The eigenvalues of the matrix $\mathcal{D}(\xi)$ are $k^2(2\mu_{\text{macro}} + \lambda_{\text{macro}})$ and $k^2\mu_{\text{macro}}$ (of multiplicity 2) such that positivity at $k \neq 0$ is satisfied if and only if¹⁴:

$$\mu_{\text{macro}} > 0, \quad 2\mu_{\text{macro}} + \lambda_{\text{macro}} > 0, \quad (68)$$

¹⁴The eigenvalues of $\mathcal{D}(\xi)$ are independent of the propagation direction $\xi \in \mathbb{R}^3$ which makes sense for the isotropic formulation at hand.

which are the usual strong ellipticity conditions. We note here that the latter calculations also show that $B(e_1) = \frac{1}{\rho} k^2 \mathcal{D}(e_1)$. Alternatively, one may directly form the so-called **acoustic tensor** $B(\xi) \in \mathbb{R}^{3 \times 3}$ by

$$B(\xi) \cdot \hat{u} := [\mathbb{C}(\hat{u} \otimes \xi)] \cdot \xi, \quad \forall \hat{u} \in \mathbb{R}^3, \quad (69)$$

in indices we have $(B(\xi))_{ij} = \mathbb{C}_{ikjl} \hat{u}_k \hat{u}_l \neq \mathbb{C}(\xi \otimes \xi)$. With (69) we obtain¹⁵:

$$\begin{aligned} \langle \hat{u}, B(\xi) \cdot \hat{u} \rangle_{\mathbb{R}^3} &= \langle \underbrace{[\mathbb{C}(\hat{u} \otimes \xi)] \xi}_{=: \hat{B} \in \mathbb{R}^{3 \times 3}}, \hat{u} \rangle_{\mathbb{R}^3} = \langle \hat{B} \cdot \xi, \hat{u} \rangle_{\mathbb{R}^3} = \langle \hat{B} \cdot (\xi \otimes \hat{u}), \mathbb{1} \rangle_{\mathbb{R}^{3 \times 3}} = \langle \hat{B}, (\xi \otimes \hat{u})^T \rangle_{\mathbb{R}^{3 \times 3}} \\ &= \langle \hat{B}, \hat{u} \otimes \xi \rangle_{\mathbb{R}^{3 \times 3}} = \langle \mathbb{C}(\hat{u} \otimes \xi), \hat{u} \otimes \xi \rangle_{\mathbb{R}^{3 \times 3}}, \end{aligned} \quad (70)$$

and we see that strong ellipticity $\langle \mathbb{C}(\hat{u} \otimes \xi), \hat{u} \otimes \xi \rangle_{\mathbb{R}^{3 \times 3}} > 0$ is equivalent to the positive definiteness of the acoustic tensor $B(\xi)$.

4 A further comparison: the linear Cosserat model

In the isotropic hyperelastic case the elastic energy density and the kinetic energy density of the Cosserat model read:

$$\begin{aligned} W &= \mu_{\text{macro}} \|\text{sym } \nabla u\|^2 + \mu_c \|\text{skew}(\nabla u - A)\|^2 + \frac{\lambda_{\text{macro}}}{2} (\text{tr}(\nabla u))^2 + \frac{\mu_{\text{macro}} L_c^2}{2} \|\text{Curl} A\|^2, \\ J &= \frac{\rho}{2} \|u_t\|^2 + \frac{\eta}{2} \|A_t\|^2. \end{aligned} \quad (71)$$

Introducing the canonical identification of R^3 with $\mathfrak{so}(3)$, A can be expressed as a function of $a \in \mathbb{R}^3$ as:

$$A = \text{anti}(a) = \begin{pmatrix} 0 & -a_3 & a_2 \\ a_3 & 0 & -a_1 \\ -a_2 & a_1 & 0 \end{pmatrix}. \quad (72)$$

Here, we assume for clarity a uni-constant curvature expression in terms of only $\|\text{Curl} A\|^2$. Strict positive definiteness of the potential energy is equivalent to the following simple relations for the introduced parameters

$$2\mu_{\text{macro}} + 3\lambda_{\text{macro}} > 0, \quad \mu_{\text{macro}} > 0, \quad \mu_c > 0, \quad L_c > 0. \quad (73)$$

The dynamical formulation is obtained defining a joint Hamiltonian and assuming stationary action. The dynamical equilibrium equations are:

$$\begin{aligned} \rho u_{,tt} &= \text{Div} [2\mu_{\text{macro}} \text{sym}(\nabla u - A) + 2\mu_c \text{skew}(\nabla u - A) + \lambda_{\text{macro}} \text{tr}(\nabla u - A) \mathbb{1}], \\ \eta A_{,tt} &= -\mu_{\text{macro}} L_c^2 \text{skew}(\text{Curl} \text{Curl} A) + 2\mu_c \text{skew}(\nabla u - A), \end{aligned} \quad (74)$$

see also [18, 19, 41, 42] for formulations in terms of axial vectors. Note that for zero Cosserat couple modulus $\mu_c = 0$ the coupling of the two fields (u, A) is absent, in opposition to the relaxed micromorphic model (eq. (15)). Considering plane and stationary waves of amplitudes \hat{u} and \hat{a} , it is possible to express this system as:

$$A_6(\omega, k) \cdot \begin{pmatrix} \hat{u}_1 & \hat{a}_1 \end{pmatrix}^T = 0, \quad A_7(\omega, k) \cdot \begin{pmatrix} \hat{u}_2 & -\hat{a}_3 \end{pmatrix}^T = 0, \quad A_7(\omega, k) \cdot \begin{pmatrix} \hat{u}_3 & \hat{a}_2 \end{pmatrix}^T = 0, \quad (75)$$

where

$$A_6(\omega, k) = \begin{pmatrix} k^2(2\mu_{\text{macro}} + \lambda_{\text{macro}})/\rho - \omega^2 & 0 \\ 0 & (2\mu_{\text{macro}} L_c^2 k^2 + 2\mu_c)/\eta - \omega^2 \end{pmatrix}, \quad (76)$$

$$A_7(\omega, k) = \begin{pmatrix} k^2(\mu_{\text{macro}} + \mu_c)/\rho - \omega^2 & -2ik\mu_c/\rho \\ ik\mu_c/\eta & (k^2\mu_{\text{macro}} L_c^2 + 4\mu_c)/(2\eta) - \omega^2 \end{pmatrix}. \quad (77)$$

¹⁵The term $[\mathbb{C}(\hat{u} \otimes \xi)] \cdot (\hat{u} \otimes \xi)$ that in index notation reads $\mathbb{C}_{ijkl} \hat{u}_k \xi_l \hat{u}_j \xi_m$, is different from $\mathbb{C}[(\hat{u} \otimes \xi) \cdot (\hat{u} \otimes \xi)]$, i.e. $\mathbb{C}_{ijkl} \hat{u}_k \xi_m \hat{u}_m \xi_l$.

As done in the case of the relaxed micromorphic model, it is possible to express equivalently the problem with $A_6(\omega, k)$ and the following symmetric matrix:

$$\bar{A}_7(k) = \text{diag}_7 \cdot A_7(\omega, k) \cdot \text{diag}_7^{-1} = \begin{pmatrix} k^2(\mu_{\text{macro}} + \mu_c)/\rho - \omega^2 & \sqrt{2}k\mu_c/\sqrt{\rho\eta} \\ \sqrt{2}k\mu_c/\sqrt{\rho\eta} & (k^2\mu_{\text{macro}}L_c^2 + 4\mu_c)/(2\eta) - \omega^2 \end{pmatrix}, \quad (78)$$

where

$$\text{diag}_7 = \begin{pmatrix} \sqrt{\rho} & 0 \\ 0 & i\sqrt{2\eta} \end{pmatrix}. \quad (79)$$

Since ω^2 appears only on the diagonal, the problem can be analogously expressed as the following eigenvalue-problems:

$$\det (B_6(k) - \omega^2 \mathbb{1}) = 0, \quad \det (B_7(k) - \omega^2 \mathbb{1}) = 0, \quad (80)$$

where

$$B_6(k) = \begin{pmatrix} k^2(2\mu_{\text{macro}} + \lambda_{\text{macro}})/\rho & 0 \\ 0 & (2\mu_{\text{macro}}L_c^2k^2 + 2\mu_c)/\eta^2 \end{pmatrix}, \quad (81)$$

$$B_7(k) = \begin{pmatrix} k^2(\mu_{\text{macro}} + \mu_c)/\rho & \sqrt{2}k\mu_c/\sqrt{\rho\eta} \\ \sqrt{2}k\mu_c/\sqrt{\rho\eta} & (k^2\mu_{\text{macro}}L_c^2 + 4\mu_c)/(2\eta) \end{pmatrix}, \quad (82)$$

are the blocks of the **acoustic tensor** B

$$B(k) = \begin{pmatrix} B_6 & 0 & 0 \\ 0 & B_7 & 0 \\ 0 & 0 & B_7 \end{pmatrix}. \quad (83)$$

The eigenvalues of the matrix $B_6(k)$ are simply the elements of the diagonal, therefore we have:

$$\omega_{\text{acoustic, long}}(k) = k \sqrt{\frac{2\mu_{\text{macro}} + \lambda_{\text{macro}}}{\rho}}, \quad \omega_{\text{optic, long}}(k) = \sqrt{\frac{2\mu_{\text{macro}}L_c^2k^2 + 2\mu_c}{\eta}}, \quad (84)$$

while for $B_7(k)$ it is possible to find:

$$\omega_{\text{acoustic, trans}}(k) = \sqrt{a(k) - \sqrt{a(k)^2 - b^2k^2}}, \quad \omega_{\text{optic, trans}}(k) = \sqrt{a(k) + \sqrt{a(k)^2 - b^2k^2}}, \quad (85)$$

where we have set:

$$a(k) = \frac{4\mu_c + \mu_{\text{macro}}L_c^2k^2}{\eta} + 2\frac{\mu_{\text{macro}} + \mu_c}{\rho}k^2, \quad b^2 = 8\frac{\mu_{\text{macro}}(4\mu_c + k^2L_c^2(\mu_{\text{macro}} + \mu_c))}{\rho\eta}. \quad (86)$$

The acoustic branches are those curves $\omega = \omega(k)$ as solutions of (79) that satisfy $\omega(0) = 0$. We note here that the acoustic branches of the longitudinal and transverse dispersion curves have as tangent in $k = 0$ ¹⁶

$$c_l = \frac{d\omega_{\text{acoustic, long}}(0)}{dk} = \sqrt{\frac{2\mu_{\text{macro}} + \lambda_{\text{macro}}}{\rho}}, \quad c_t = \frac{d\omega_{\text{acoustic, trans}}(0)}{dk} = \sqrt{\frac{\mu_{\text{macro}}}{\rho}}, \quad (87)$$

respectively. Moreover, the longitudinal acoustic branch is non-dispersive, i.e. a straight line with slope (87)₁. The matrix $B_6(k)$ is positive-definite for arbitrary $k \neq 0$ if:

$$2\mu_{\text{macro}} + \lambda_{\text{macro}} > 0, \quad \mu_{\text{macro}} > 0, \quad \mu_c \geq 0, \quad (88)$$

¹⁶To obtain the slopes in 0 it is possible to search for a solution of the type $\omega = ak$ and then evaluate the limit for $a \rightarrow 0$, see [9] for a thorough explanation in the relaxed micromorphic case.

Using the Sylvester criterion, $B_7(k)$ is positive-definite if and only if the principal minors are positive, namely:

$$\begin{aligned} (B_7)_{11} &= k^2 \frac{(\mu_{\text{macro}} + \mu_c)}{\rho} > 0, \\ \det(B_7) &= \frac{k^2}{2\eta\rho} (4\mu_{\text{macro}}\mu_c + k^2\mu_{\text{macro}}L_c^2(\mu_{\text{macro}} + \mu_c)) > 0, \end{aligned} \tag{89}$$

from which we obtain the condition:

$$\mu_{\text{macro}} + \mu_c > 0, \quad \mu_{\text{macro}} > 0, \quad \mu_c \geq 0. \tag{90}$$

Considering these two sets of conditions, it is possible to state a **necessary** and **sufficient** condition for the positive definiteness of $B_6(k)$ and $B_7(k)$ and therefore of the acoustic tensor $B(k)$:

$$2\mu_{\text{macro}} + \lambda_{\text{macro}} > 0, \quad \mu_{\text{macro}} > 0, \quad \mu_c \geq 0. \tag{91}$$

which are implied by the positive-definiteness of the energy (73). Eringen [11, p.150] also obtains correctly (88) **and** (90) (in his notation $\mu_c = \kappa/2$, $\mu_{\text{macro}} = \mu_{\text{Eringen}} + \kappa/2$).

In [1, 10] strong ellipticity for the Cosserat-micropolar model is defined and investigated. In this respect we note that ellipticity is connected to acceleration waves while our investigation concerns real wave velocities for planar waves. Similarly to [37] it is established in [1, 10] that strong ellipticity for the micropolar model holds if and only if (the uni-constant curvature case in our notation):

$$2\mu_{\text{macro}} + \lambda_{\text{macro}} > 0, \quad \mu_{\text{macro}} + \mu_c > 0. \tag{92}$$

We conclude that for micropolar material models, (and therefore also for micromorphic materials) strong ellipticity (92) is too weak to ensure real planar waves since it is implied by, but does not imply (91). This fact seems to have been appreciated also in the study of the Cosserat model [6, 13, 16, 22, 34].

5 Conclusion

In this paper we derive the set of necessary and sufficient conditions that have to be imposed on the constitutive parameters of the relaxed micromorphic model in order to guarantee

- positive definiteness,
- real wave velocity ,
- Legendre-Hadamard strong ellipticity condition.

We show that, if, on the one hand, definite positiveness implies real wave propagation, on the other hand, real wave propagation is not guaranteed by the strong ellipticity condition.

We conclude that, in strong contrast to the case of classical isotropic linear elasticity, where the three concepts are known to be equivalent, in the case of the relaxed micromorphic continua only definite positiveness of the strain energy density can be considered to be a good criterion to guarantee real wave speeds in the considered media. The proposed considerations can be extended to all generalized continua where the equivalence between the three notions is far from being straightforward.

Data accessibility

This work does not have any experimental data.

Competing interests

The authors have no conflict of interests to declare.

Author's contribute

All the authors contributed equally to this work.

Acknowledgments

We thank Victor A. Eremeyev for helpful clarification. The work of Ionel-Dumitrel Ghiba was supported by a grant of the Romanian National Authority for Scientific Research and Innovation, CNCS-UEFISCDI, project number PN-II-RU-TE-2014-4-1109. Angela Madeo thanks INSA-Lyon for the funding of the BQR 2016 "Caractérisation mécanique inverse des métamatériaux: modélisation, identification expérimentale des paramètres et évolutions possibles", as well as the PEPS CNRS-INSIS.

Funding statement

This research received no specific grant.

Ethics statement

This work did not involve any research on humans or animals.

6 References

- [1] Holm Altenbach, Victor A. Eremeyev, Leonid P. Lebedev, and Leonardo A. Rendón. Acceleration waves and ellipticity in thermoelastic micropolar media. *Archive of Applied Mechanics*, 80(3):217–227, 2010.
- [2] Daniel Balzani, Patrizio Neff, Jörg Schröder, and Gerhard A. Holzapfel. A polyconvex framework for soft biological tissues. Adjustment to experimental data. *International Journal of Solids and Structures*, 43(20):6052–6070, 2006.
- [3] Gabriele Barbagallo, Marco Valerio d’Agostino, Rafael Abreu, Ionel-Dumitrel Ghiba, Angela Madeo, and Patrizio Neff. Transparent anisotropy for the relaxed micromorphic model: macroscopic consistency conditions and long wave length asymptotics. *Preprint ArXiv*, 1601.03667, 2016.
- [4] Sebastian Bauer, Patrizio Neff, Dirk Pauly, and Gerhard Starke. New Poincaré-type inequalities. *Comptes Rendus Mathématique*, 352(2):163–166, 2014.
- [5] Sebastian Bauer, Patrizio Neff, Dirk Pauly, and Gerhard Starke. Dev-Div- and DevSym-DevCurl-inequalities for incompatible square tensor fields with mixed boundary conditions. *ESAIM: Control, Optimisation and Calculus of Variations*, 22(1):112–133, 2016.
- [6] Davide Bigoni and Panos A. Gougiotis. Folding and faulting of an elastic continuum. *Proceedings of the Royal Society A*, 472(2187):20160018, 2016.
- [7] Stan Chiriță, Alexandre Danescu, and Michele Ciarletta. On the strong ellipticity of the anisotropic linearly elastic materials. *Journal of Elasticity*, 87(1):1–27, 2007.
- [8] Stan Chiriță and Ionel-Dumitrel Ghiba. Strong ellipticity and progressive waves in elastic materials with voids. *Proceedings of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 466(2114):439–458, 2009.
- [9] Marco Valerio d’Agostino, Gabriele Barbagallo, Ionel-Dumitrel Ghiba, Angela Madeo, and Patrizio Neff. A panorama of dispersion curves for the weighted isotropic relaxed micromorphic model. *Preprint ArXiv*, 1610.03296, 2016.
- [10] Victor A. Eremeyev. Acceleration waves in micropolar elastic media. *Doklady Physics*, 50(4):204–206, 2005.
- [11] Ahmed Cemal Eringen. *Microcontinuum field theories*. Springer-Verlag, New York, 1999.
- [12] Denos C. Gazis and Richard F. Wallis. Extensional waves in cubic crystals plates. In *Proceedings of the 4th U.S. National Congress in Applied Mechanics*, pages 161–168, 1962.
- [13] Ionel-Dumitrel Ghiba, Patrizio Neff, Angela Madeo, and Ingo Münch. A variant of the linear isotropic indeterminate couple-stress model with symmetric local force-stress, symmetric nonlocal force-stress, symmetric couple-stresses and orthogonal boundary conditions. *Mathematics and Mechanics of Solids*, 2016.
- [14] Ionel-Dumitrel Ghiba, Patrizio Neff, Angela Madeo, Luca Placidi, and Giuseppe Rosi. The relaxed linear micromorphic continuum: existence, uniqueness and continuous dependence in dynamics. *Mathematics and Mechanics of Solids*, 20(10):1171–1197, 2014.
- [15] George T. Gilbert. Positive definite matrices and Sylvester’s criterion. *The American Mathematical Monthly*, 98(1):44–46, 1991.
- [16] Panos A. Gougiotis and Davide Bigoni. Stress channelling in extreme couple-stress materials Part I: Strong ellipticity, wave propagation, ellipticity, and discontinuity relations. *Journal of the Mechanics and Physics of Solids*, 88:150–168, 2016.

- [17] Steven Greene, Stefano Gonella, and Wing Kam Liu. Microelastic wave field signatures and their implications for microstructure identification. *International Journal of Solids and Structures*, 49(22):3148–3157, 2012.
- [18] Jena Jeong and Patrizio Neff. Existence, uniqueness and stability in linear Cosserat elasticity for weakest curvature conditions. *Mathematics and Mechanics of Solids*, 15(1):78–95, 2010.
- [19] Jena Jeong, Hamidréza Ramézani, Ingo Münch, and Patrizio Neff. A numerical study for linear isotropic Cosserat elasticity with conformally invariant curvature. *Zeitschrift für Angewandte Mathematik und Mechanik*, 89(7):552–569, 2009.
- [20] Angela Madeo, Gabriele Barbagallo, Marco Valerio d’Agostino, Luca Placidi, and Patrizio Neff. First evidence of non-locality in real band-gap metamaterials: determining parameters in the relaxed micromorphic model. *Proceedings of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 472(2190):20160169, 2016.
- [21] Angela Madeo, Manuel Collet, Marco Miniaci, Kévin Billon, Morvan Ouisse, and Patrizio Neff. Modeling real phononic crystals via the weighted relaxed micromorphic model with free and gradient micro-inertia. *Preprint ArXiv*, 1610.03878, 2016.
- [22] Angela Madeo, Ionel-Dumitrel Ghiba, Patrizio Neff, and Ingo Münch. A new view on boundary conditions in the Grioli–Koiter–Mindlin–Toupin indeterminate couple stress model. *European Journal of Mechanics - A/Solids*, 59:294–322, 2016.
- [23] Angela Madeo, Patrizio Neff, Elias C. Aifantis, Gabriele Barbagallo, and Marco Valerio d’Agostino. On the role of micro-inertia in enriched continuum mechanics. *Preprint ArXiv*, 1607.07385, 2016.
- [24] Angela Madeo, Patrizio Neff, Gabriele Barbagallo, Marco Valerio d’Agostino, and Ionel-Dumitrel Ghiba. A review on wave propagation modeling in band-gap metamaterials via enriched continuum models. *Preprint ArXiv*, 1609.01073, 2016.
- [25] Angela Madeo, Patrizio Neff, Marco Valerio d’Agostino, and Gabriele Barbagallo. Complete band gaps including non-local effects occur only in the relaxed micromorphic model. *Comptes Rendus Mécanique*, 2016.
- [26] Angela Madeo, Patrizio Neff, Ionel-Dumitrel Ghiba, Luca Placidi, and Giuseppe Rosi. Band gaps in the relaxed linear micromorphic continuum. *Zeitschrift für Angewandte Mathematik und Mechanik*, 95(9):880–887, 2014.
- [27] Angela Madeo, Patrizio Neff, Ionel-Dumitrel Ghiba, Luca Placidi, and Giuseppe Rosi. Wave propagation in relaxed micromorphic continua: modeling metamaterials with frequency band-gaps. *Continuum Mechanics and Thermodynamics*, 27(4-5):551–570, 2015.
- [28] Angela Madeo, Patrizio Neff, Ionel-Dumitrel Ghiba, and Giuseppe Rosi. Reflection and transmission of elastic waves in non-local band-gap metamaterials: a comprehensive study via the relaxed micromorphic model. *Journal of the Mechanics and Physics of Solids*, 95:441–479, 2016.
- [29] Gérard A. Maugin. *Non-Classical Continuum Mechanics*, volume 51. Springer, 2016.
- [30] Jose Merodio and Patrizio Neff. A note on tensile instabilities and loss of ellipticity for a fiber-reinforced nonlinearly elastic solid. *Archives of Mechanics*, 58(3):293–303, 2006.
- [31] Raymond David Mindlin. Microstructure in linear elasticity. Technical report, Office of Naval Research, 1963.
- [32] Raymond David Mindlin. Micro-structure in linear elasticity. *Archive for Rational Mechanics and Analysis*, 16(1):51–78, 1964.
- [33] Ingo Münch and Patrizio Neff. Rotational invariance conditions in elasticity, gradient elasticity and its connection to isotropy. *Preprint ArXiv*, 1603.06153, 2016.
- [34] Ingo Münch, Patrizio Neff, Angela Madeo, and Ionel-Dumitrel Ghiba. The modified indeterminate couple stress model: Why Yang et al.’s arguments motivating a symmetric couple stress tensor contain a gap and why the couple stress tensor may be chosen symmetric nevertheless. *Preprint ArXiv*, 1512.02053, 2015.
- [35] Patrizio Neff. On material constants for micromorphic continua. In *Trends in Applications of Mathematics to Mechanics, STAMM Proceedings, Seeheim*, pages 337–348. Shaker–Verlag, 2004.
- [36] Patrizio Neff. The Cosserat couple modulus for continuous solids is zero viz the linearized Cauchy-stress tensor is symmetric. *Zeitschrift für Angewandte Mathematik und Mechanik*, 86(11):892–912, 2006.
- [37] Patrizio Neff. Relations of constants for isotropic linear Cosserat elasticity (http://www.uni-due.de/%7ehm0014/Cosserat_files/web_coss_relations.pdf, typos in equations 2.8,2.9,2.10). Technical report, Fachbereich Mathematik, Technische Universität Darmstadt, Darmstadt, Germany, 2008.
- [38] Patrizio Neff and Samuel Forest. A geometrically exact micromorphic model for elastic metallic foams accounting for affine microstructure. Modelling, existence of minimizers, identification of moduli and computational results. *Journal of Elasticity*, 87(2-3):239–276, 2007.
- [39] Patrizio Neff, Ionel-Dumitrel Ghiba, Markus Lazar, and Angela Madeo. The relaxed linear micromorphic continuum: well-posedness of the static problem and relations to the gauge theory of dislocations. *The Quarterly Journal of Mechanics and Applied Mathematics*, 68(1):53–84, 2015.
- [40] Patrizio Neff, Ionel-Dumitrel Ghiba, Angela Madeo, Luca Placidi, and Giuseppe Rosi. A unifying perspective: the relaxed linear micromorphic continuum. *Continuum Mechanics and Thermodynamics*, 26(5):639–681, 2014.
- [41] Patrizio Neff and Jena Jeong. A new paradigm: the linear isotropic Cosserat model with conformally invariant curvature energy. *Zeitschrift für Angewandte Mathematik und Mechanik*, 89(2):107–122, 2009.
- [42] Patrizio Neff, Jena Jeong, and Andreas Fischle. Stable identification of linear isotropic Cosserat parameters: bounded stiffness in bending and torsion implies conformal invariance of curvature. *Acta Mechanica*, 211(3-4):237–249, 2010.

- [43] Patrizio Neff, Dirk Pauly, and Karl-Josef Witsch. A canonical extension of Korn's first inequality to $H(\text{Curl})$ motivated by gradient plasticity with plastic spin. *Comptes Rendus Mathematique*, 349(23):1251–1254, 2011.
- [44] Patrizio Neff, Dirk Pauly, and Karl-Josef Witsch. Maxwell meets Korn: A new coercive inequality for tensor fields in $R^{n \times n}$ with square-integrable exterior derivative. *Mathematical Methods in the Applied Sciences*, 35(1):65–71, 2012.
- [45] Patrizio Neff, Dirk Pauly, and Karl-Josef Witsch. Poincaré meets Korn via Maxwell: Extending Korn's first inequality to incompatible tensor fields. *Journal of Differential Equations*, 258(4):1267–1302, 2015.
- [46] Witold Nowacki. *Theory of Asymmetric Elasticity*. Pergamon Press, 1985.
- [47] Giovanni Romano, Raffaele Barretta, and Marina Diaco. Micromorphic continua: non-redundant formulations. *Continuum Mechanics and Thermodynamics*, 2016.
- [48] Jörg Schröder and Patrizio Neff. Application of polyconvex anisotropic free energies to soft tissues. *5th World Congress on Computational Mechanics*, 2002.
- [49] Jörg Schröder, Patrizio Neff, and Vera Ebbing. Anisotropic polyconvex energies on the basis of crystallographic motivated structural tensors. *Journal of the Mechanics and Physics of Solids*, 56(12):3486–3506, 2008.
- [50] A.C. Smith. Waves in micropolar elastic solids. *International Journal of Engineering Science*, 5(10):741–746, 1967.
- [51] A.C. Smith. Inequalities between the constants of a linear micro-elastic solid. *International Journal of Engineering Science*, 6(2):65–74, 1968.

7 Appendix

7.1 Inequality relations between material parameters

The formulas in section 2.1 are based on the harmonic mean of two numbers κ_e and κ_{micro} (or μ_e and μ_{micro}). If the two numbers are positive, it is easy to see that:

$$\kappa_{\text{macro}} \leq \min(\kappa_e, \kappa_{\text{micro}}). \quad (93)$$

Here, we show that the same conclusion still holds if we merely assume that $\kappa_e + \kappa_{\text{micro}} > 0$. This allows for either $\kappa_e < 0$ or $\kappa_{\text{micro}} < 0$. Therefore, considering that $\kappa_e + \kappa_{\text{micro}} > 0$, even if the energy is not strictly positive, it is possible to derive that:

$$\begin{aligned} \kappa_{\text{macro}} &= \frac{\kappa_{\text{micro}} \kappa_e}{\kappa_e + \kappa_{\text{micro}}} = \frac{\kappa_{\text{micro}} \kappa_e + \kappa_e^2 - \kappa_e^2}{\kappa_e + \kappa_{\text{micro}}} = \kappa_e \frac{\kappa_{\text{micro}} + \kappa_e}{\kappa_e + \kappa_{\text{micro}}} - \frac{\kappa_e^2}{\kappa_e + \kappa_{\text{micro}}} = \kappa_e - \underbrace{\frac{\kappa_e^2}{\kappa_e + \kappa_{\text{micro}}}}_{\leq 0} \leq \kappa_e, \quad (94) \\ \kappa_{\text{macro}} &= \frac{\kappa_{\text{micro}} \kappa_e}{\kappa_e + \kappa_{\text{micro}}} = \frac{\kappa_{\text{micro}} \kappa_e + \kappa_{\text{micro}}^2 - \kappa_{\text{micro}}^2}{\kappa_e + \kappa_{\text{micro}}} = \kappa_{\text{micro}} \frac{\kappa_{\text{micro}} + \kappa_e}{\kappa_e + \kappa_{\text{micro}}} - \frac{\kappa_{\text{micro}}^2}{\kappa_e + \kappa_{\text{micro}}} = \kappa_{\text{micro}} - \underbrace{\frac{\kappa_{\text{micro}}^2}{\kappa_e + \kappa_{\text{micro}}}}_{\leq 0} \leq \kappa_{\text{micro}}. \end{aligned}$$

Considering similarly $\mu_e + \mu_{\text{micro}} > 0$, it is possible to obtain:

$$\begin{aligned} \mu_{\text{macro}} &= \frac{\mu_{\text{micro}} \mu_e}{\mu_e + \mu_{\text{micro}}} = \frac{\mu_{\text{micro}} \mu_e + \mu_e^2 - \mu_e^2}{\mu_e + \mu_{\text{micro}}} = \mu_e \frac{\mu_{\text{micro}} + \mu_e}{\mu_e + \mu_{\text{micro}}} - \frac{\mu_e^2}{\mu_e + \mu_{\text{micro}}} = \mu_e - \underbrace{\frac{\mu_e^2}{\mu_e + \mu_{\text{micro}}}}_{\leq 0} \leq \mu_e, \quad (95) \\ \mu_{\text{macro}} &= \frac{\mu_{\text{micro}} \mu_e}{\mu_e + \mu_{\text{micro}}} = \frac{\mu_{\text{micro}} \mu_e + \mu_{\text{micro}}^2 - \mu_{\text{micro}}^2}{\mu_e + \mu_{\text{micro}}} = \mu_{\text{micro}} \frac{\mu_{\text{micro}} + \mu_e}{\mu_e + \mu_{\text{micro}}} - \frac{\mu_{\text{micro}}^2}{\mu_e + \mu_{\text{micro}}} = \mu_{\text{micro}} - \underbrace{\frac{\mu_{\text{micro}}^2}{\mu_e + \mu_{\text{micro}}}}_{\leq 0} \leq \mu_{\text{micro}}. \end{aligned}$$

Therefore, if $\mu_e + \mu_{\text{micro}} > 0$ and $\kappa_e + \kappa_{\text{micro}} > 0$, the macroscopic parameters are less or equal than respective microscopic parameters, namely:

$$\kappa_e \geq \kappa_{\text{macro}}, \quad \kappa_{\text{micro}} \geq \kappa_{\text{macro}}, \quad \mu_e \geq \mu_{\text{macro}}, \quad \mu_{\text{micro}} \geq \mu_{\text{macro}}, \quad (96)$$

and it is possible to show that:

$$\begin{aligned} 2\mu_e + \lambda_e &= \frac{1}{3}(4\mu_e + 3\kappa_e) \geq \frac{1}{3}(4\mu_{\text{macro}} + 3\kappa_{\text{macro}}) = 2\mu_{\text{macro}} + \lambda_{\text{macro}} > 0, \\ 2\mu_{\text{micro}} + \lambda_{\text{micro}} &= \frac{1}{3}(4\mu_{\text{micro}} + 3\kappa_{\text{micro}}) \geq \frac{1}{3}(4\mu_{\text{macro}} + 3\kappa_{\text{macro}}) = 2\mu_{\text{macro}} + \lambda_{\text{macro}} > 0, \quad (97) \\ (2\mu_e + \lambda_e) + (2\mu_{\text{micro}} + \lambda_{\text{micro}}) &\geq 2(2\mu_{\text{macro}} + \lambda_{\text{macro}}) > 0, \\ 4\mu_{\text{macro}} + 3\kappa_e &\geq 4\mu_{\text{macro}} + 3\kappa_{\text{macro}} = 3(2\mu_{\text{macro}} + \lambda_{\text{macro}}) > 0. \end{aligned}$$

Therefore, the set of inequalities (50) is implied from the smaller set:

$$\boxed{\mu_e > 0, \quad \mu_{\text{micro}} > 0, \quad \mu_c \geq 0, \quad \kappa_e + \kappa_{\text{micro}} > 0, \quad 2\mu_{\text{macro}} + \lambda_{\text{macro}} > 0.} \quad (98)$$

We note here that $3(2\mu_e + \lambda_e) \geq 4\mu_{\text{macro}} + 3\kappa_e \geq 3(2\mu_{\text{macro}} + \lambda_{\text{macro}})$ because:

$$3(2\mu_e + \lambda_e) = 4\mu_e + 3\kappa_e \geq 4\mu_{\text{macro}} + 3\kappa_e \geq 4\mu_{\text{macro}} + 3\kappa_{\text{macro}} = 3(2\mu_{\text{macro}} + \lambda_{\text{macro}}). \quad (99)$$

7.2 The 12×12 acoustic tensor for arbitrary direction

We suppose that the space dependence of all introduced kinematic fields are limited to a direction defined by a unit vector ξ which is the direction of propagation of the wave. Therefore, we look for solutions of:

$$\begin{aligned} \rho u_{,tt} &= \text{Div} [2\mu_e \text{sym}(\nabla u - P) + 2\mu_c \text{skew}(\nabla u - P) + \lambda_e \text{tr}(\nabla u - P)\mathbb{1}], \\ \eta P_{,tt} &= -\mu_e L_c^2 \text{Curl} \text{Curl} P + 2\mu_e \text{sym}(\nabla u - P) + 2\mu_c \text{skew}(\nabla u - P) \\ &\quad + \lambda_e \text{tr}(\nabla u - P)\mathbb{1} - [2\mu_{\text{micro}} \text{sym} P + \lambda_{\text{micro}} \text{tr}(P)\mathbb{1}], \end{aligned} \quad (100)$$

in the form:

$$\begin{aligned} u(x, t) &= \widehat{u} \underbrace{e^{i(k(\xi, x)_{\mathbb{R}^3} - \omega t)}}_{s(x, t) \in \mathbb{R}/\mathbb{C} \text{ scalar}}, & \widehat{u} &\in \mathbb{C}^3, & \|\xi\|^2 &= 1, \\ P(x, t) &= \widehat{P} \underbrace{e^{i(k(\xi, x)_{\mathbb{R}^3} - \omega t)}}_{s(x, t) \in \mathbb{R}/\mathbb{C} \text{ scalar}}, & \widehat{P} &\in \mathbb{C}^{3 \times 3}, \end{aligned} \quad (101)$$

where \widehat{u} is the polarization vector and \widehat{P} is the polarization matrix. We start by remarking that considering $A, B \in \mathbb{R}^{3 \times 3}$ we have that:

$$\text{Curl}(A \cdot B) = L_B(\nabla A) + A \cdot \text{Curl}(B), \quad (102)$$

where $L_B : \mathbb{R}^{27} \rightarrow \mathbb{R}^{3 \times 3}$ is a linear operator with constant coefficients defined by the appropriate product rule of differentiation. Therefore we obtain:

$$\text{Curl}(\widehat{P} s(x, t)) = \text{Curl}(\widehat{P} \cdot \mathbb{1} s(x, t)) = \widehat{P} \cdot \text{Curl}(\mathbb{1} s(x, t)), \quad (103)$$

where:

$$\text{Curl}(\mathbb{1} s(x, t)) = \begin{pmatrix} 0 & \partial_3 s(x, t) & \partial_2 s(x, t) \\ -\partial_3 s(x, t) & 0 & \partial_1 s(x, t) \\ \partial_2 s(x, t) & -\partial_1 s(x, t) & 0 \end{pmatrix} \in \mathfrak{so}(3). \quad (104)$$

The derivatives of $s(x, t)$ can be evaluated considering:

$$\nabla_x s(x, t) = \begin{pmatrix} \partial_1 s(x, t) \\ \partial_2 s(x, t) \\ \partial_3 s(x, t) \end{pmatrix} = e^{i(k(\xi, x)_{\mathbb{R}^3} - \omega t)} \begin{pmatrix} i k \xi_1 \\ i k \xi_2 \\ i k \xi_3 \end{pmatrix} = e^{i(k(\xi, x)_{\mathbb{R}^3} - \omega t)} i k \xi = i k \xi s(x, t). \quad (105)$$

It can be noticed that:

$$\text{Curl}(s(x, t)\mathbb{1}) = \text{anti}(\nabla s(x, t)) = e^{i(k(\xi, x)_{\mathbb{R}^3} - \omega t)} i k \text{anti}(\xi) = s(x, t) i k \text{anti}(\xi). \quad (106)$$

Therefore, it is possible to evaluate the $\text{Curl} \text{Curl} P$ as:

$$\begin{aligned} \text{Curl} \text{Curl}(\widehat{P} s(x, t)) &= \text{Curl}(\underbrace{\widehat{P} \cdot \text{anti}(\xi)}_{\in \mathfrak{so}(3)} i k s(x, t)) = i k \text{Curl}([\widehat{P} \cdot \text{anti}(\xi)] \cdot \mathbb{1} s(x, t)) = i k \widehat{P} \cdot \text{anti}(\xi) \text{Curl}(\mathbb{1} s(x, t)) \\ &= i k i k \widehat{P} \cdot \text{anti}(\xi) \cdot \text{anti}(\xi) s(x, t) = -k^2 \widehat{P} \cdot \text{anti}(\xi) \cdot \text{anti}(\xi) e^{i(k(\xi, x)_{\mathbb{R}^3} - \omega t)}. \end{aligned} \quad (107)$$

On the other hand, the second derivative of P with respect to time is:

$$P_{,tt} = \partial_t^2(\widehat{P} e^{i(k(\xi, x)_{\mathbb{R}^3} - \omega t)}) = -\omega^2 \widehat{P} e^{i(k(\xi, x)_{\mathbb{R}^3} - \omega t)} = -\omega^2 \widehat{P} s(x, t). \quad (108)$$

Analogously for u it is possible to evaluate the gradient and the derivatives with respect to time as:

$$\nabla_x u = i k s(x, t) \widehat{u} \otimes \xi, \quad u_{,tt} = -\omega^2 \widehat{u} s(x, t). \quad (109)$$

The sym, skew and tr of $\nabla u - P$ can then be expressed as:

$$\begin{aligned} \text{sym}(\nabla u - P) &= \text{sym}(i k \widehat{u} \otimes \xi - \widehat{P}) s(x, t) = (i k \text{sym}(\widehat{u} \otimes \xi) - \text{sym} \widehat{P}) s(x, t), \\ \text{skew}(\nabla u - P) &= \text{skew}(i k \widehat{u} \otimes \xi - \widehat{P}) s(x, t) = (i k \text{skew}(\widehat{u} \otimes \xi) - \text{skew} \widehat{P}) s(x, t), \\ \text{tr}(\nabla u - P) &= \text{tr}(i k \widehat{u} \otimes \xi - \widehat{P}) s(x, t) = (i k \langle \widehat{u}, \xi \rangle_{\mathbb{R}^3} - \text{tr} \widehat{P}) s(x, t). \end{aligned} \quad (110)$$

Therefore, we have:

$$\begin{aligned} \text{Div} \text{sym}(\nabla u - P) &= \text{Div} \left[(i k \text{sym}(\widehat{u} \otimes \xi) - \text{sym} \widehat{P}) s(x, t) \right] = (i k \text{sym}(\widehat{u} \otimes \xi) - \text{sym} \widehat{P}) \cdot \nabla_x s(x, t) \\ &= (i k \text{sym}(\widehat{u} \otimes \xi) - \text{sym} \widehat{P}) \cdot (i k \xi s(x, t)) = -(k^2 \text{sym}(\widehat{u} \otimes \xi) \cdot \xi + i k \text{sym} \widehat{P} \cdot \xi) s(x, t), \\ \text{Div} \text{skew}(\nabla u - P) &= \text{Div} \left[(i k \text{skew}(\widehat{u} \otimes \xi) - \text{skew} \widehat{P}) s(x, t) \right] = (i k \text{skew}(\widehat{u} \otimes \xi) - \text{skew} \widehat{P}) \cdot \nabla_x s(x, t) \\ &= (i k \text{skew}(\widehat{u} \otimes \xi) - \text{skew} \widehat{P}) \cdot (i k \xi s(x, t)) = -(k^2 \text{skew}(\widehat{u} \otimes \xi) \cdot \xi + i k \text{skew} \widehat{P} \cdot \xi) s(x, t), \\ \text{Div}(\text{tr}(\nabla u - P)\mathbb{1}) &= \text{Div} \left[(i k \langle \widehat{u}, \xi \rangle_{\mathbb{R}^3} - \text{tr} \widehat{P}) s(x, t) \right] = (i k \langle \widehat{u}, \xi \rangle_{\mathbb{R}^3} - \text{tr} \widehat{P}) \mathbb{1} \cdot \nabla_x s(x, t) \\ &= (i k \langle \widehat{u}, \xi \rangle_{\mathbb{R}^3} - \text{tr} \widehat{P}) \mathbb{1} \cdot (i k \xi s(x, t)) = -(k^2 \langle \widehat{u}, \xi \rangle_{\mathbb{R}^3} + i k \text{tr} \widehat{P}) \xi s(x, t). \end{aligned} \quad (111)$$

Here, we have considered that, given a generic $B \in \mathbb{R}^{3 \times 3}$ and a scalar $s(x, t)$, we have:

$$\text{Div}[B s(x, t)] = \underbrace{\text{Div}[B]}_{=0} s(x, t) + B \cdot \nabla_x s(x, t), \quad (112)$$

With all the formulas obtained it is possible to write (100) simplifying $s(x, t)$ everywhere as:

$$\begin{aligned} -\rho\omega^2 \hat{u} &= - [2\mu_e (k^2 \text{sym}(\hat{u} \otimes \xi) \cdot \xi + ik \text{sym} \hat{P} \cdot \xi) + 2\mu_c (k^2 \text{skew}(\hat{u} \otimes \xi) \cdot \xi + ik \text{skew} \hat{P} \cdot \xi) \\ &\quad + \lambda_e (k^2 \langle \hat{u}, \xi \rangle_{\mathbb{R}^3} + ik \text{tr} \hat{P}) \xi], \\ -\eta\omega^2 \hat{P} &= \mu_e L_c^2 k^2 \hat{P} \text{anti}(\xi) \cdot \text{anti}(\xi) + 2\mu_e (ik \text{sym}(\hat{u} \otimes \xi) - \text{sym} \hat{P}) + 2\mu_c (ik \text{skew}(\hat{u} \otimes \xi) - \text{skew} \hat{P}) \\ &\quad + \lambda_e (ik \langle \hat{u}, \xi \rangle_{\mathbb{R}^3} - \text{tr} \hat{P}) \mathbf{1} - [2\mu_{\text{micro}} \text{sym} \hat{P} + \lambda_{\text{micro}} \text{tr}(\hat{P}) \mathbf{1}], \end{aligned} \quad (113)$$

or analogously:

$$\begin{aligned} -\rho\omega^2 \hat{u} + k^2 (2\mu_e \text{sym}(\hat{u} \otimes \xi) \cdot \xi + 2\mu_c \text{skew}(\hat{u} \otimes \xi) \cdot \xi + \lambda_e \langle \hat{u}, \xi \rangle_{\mathbb{R}^3} \xi) \\ + ik (2\mu_e \text{sym} \hat{P} \cdot \xi + 2\mu_c \text{skew} \hat{P} \cdot \xi + \lambda_e \text{tr} \hat{P} \xi) = 0, \\ -\eta\omega^2 \hat{P} - \mu_e L_c^2 k^2 \hat{P} \text{anti}(\xi) \cdot \text{anti}(\xi) + 2(\mu_e + \mu_{\text{micro}}) \text{sym} \hat{P} + 2\mu_c \text{skew} \hat{P} + (\lambda_e + \lambda_{\text{micro}}) \text{tr}(\hat{P}) \mathbf{1} \\ - 2\mu_e ik \text{sym}(\hat{u} \otimes \xi) - 2\mu_c ik \text{skew}(\hat{u} \otimes \xi) - \lambda_e ik \langle \hat{u}, \xi \rangle_{\mathbb{R}^3} \mathbf{1} = 0. \end{aligned} \quad (114)$$

At given $\xi \in \mathbb{R}^3$, this is a linear system in $(\hat{u}, \hat{P}) \in \mathbb{C}^{12}$ which can be written in 12×12 matrix format as:

$$\begin{pmatrix} \tilde{A}(\xi, \omega, k) \end{pmatrix} \begin{pmatrix} \hat{u}_1 \\ \hat{u}_2 \\ \hat{u}_3 \\ \hat{P}_{11} \\ \hat{P}_{12} \\ \hat{P}_{13} \\ \hat{P}_{21} \\ \hat{P}_{22} \\ \hat{P}_{23} \\ \hat{P}_{31} \\ \hat{P}_{32} \\ \hat{P}_{33} \end{pmatrix} = 0, \quad \begin{pmatrix} \tilde{B}(\xi, k) - \omega^2 \mathbf{1} \end{pmatrix} \begin{pmatrix} \hat{u}_1 \\ \hat{u}_2 \\ \hat{u}_3 \\ \hat{P}_{11} \\ \hat{P}_{12} \\ \hat{P}_{13} \\ \hat{P}_{21} \\ \hat{P}_{22} \\ \hat{P}_{23} \\ \hat{P}_{31} \\ \hat{P}_{32} \\ \hat{P}_{33} \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \end{pmatrix}. \quad (115)$$

Here, $\tilde{B}(\xi, k)$ is the 12×12 acoustic tensor. The columns of \tilde{A} are:

$$\begin{aligned} \tilde{A}_{i1} &= \begin{pmatrix} \rho\omega^2 - k^2(\lambda_e + 2\mu_e)\xi_1^2 - k^2(\mu_c + \mu_e)(\xi_2^2 + \xi_3^2) \\ -k^2(\lambda_e - \mu_c + \mu_e)\xi_1\xi_2 \\ -k^2(\lambda_e - \mu_c + \mu_e)\xi_1\xi_3 \\ ik(\lambda_e + 2\mu_e)\xi_1 \\ ik(\mu_c + \mu_e)\xi_2 \\ ik(\mu_c + \mu_e)\xi_3 \\ -ik(\mu_c - \mu_e)\xi_2 \\ ik\lambda_e\xi_1 \\ 0 \\ -ik(\mu_c - \mu_e)\xi_3 \\ 0 \\ ik\lambda_e\xi_1 \end{pmatrix}, \quad \tilde{A}_{i2} = \begin{pmatrix} -k^2(\lambda_e - \mu_c + \mu_e)\xi_1\xi_2 \\ \rho\omega^2 - k^2(\lambda_e + 2\mu_e)\xi_2^2 - k^2(\mu_c + \mu_e)(\xi_1^2 + \xi_3^2) \\ -k^2(\lambda_e - \mu_c + \mu_e)\xi_2\xi_3 \\ ik\lambda_e\xi_2 \\ -ik(\mu_c - \mu_e)\xi_1 \\ 0 \\ ik(\mu_c + \mu_e)\xi_1 \\ ik(\lambda_e + 2\mu_e)\xi_2 \\ ik(\mu_c + \mu_e)\xi_3 \\ 0 \\ -ik(\mu_c - \mu_e)\xi_3 \\ ik\lambda_e\xi_2 \end{pmatrix}, \\ \tilde{A}_{i3} &= \begin{pmatrix} -k^2(\lambda_e - \mu_c + \mu_e)\xi_1\xi_3 \\ -k^2(\lambda_e - \mu_c + \mu_e)\xi_2\xi_3 \\ \rho\omega^2 - k^2(\lambda_e + 2\mu_e)\xi_3^2 - k^2(\mu_c + \mu_e)(\xi_1^2 + \xi_2^2) \\ ik\lambda_e\xi_3 \\ 0 \\ -ik(\mu_c - \mu_e)\xi_1 \\ 0 \\ ik\lambda_e\xi_3 \\ -ik(\mu_c - \mu_e)\xi_2 \\ ik(\mu_c + \mu_e)\xi_1 \\ ik(\mu_c + \mu_e)\xi_2 \\ ik(\lambda_e + 2\mu_e)\xi_3 \end{pmatrix}, \quad \tilde{A}_{i4} = \begin{pmatrix} -ik(\lambda_e + 2\mu_e)\xi_1 \\ -ik\lambda_e\xi_2 \\ -ik\lambda_e\xi_3 \\ \eta\omega^2 - (2(\mu_e + \mu_{\text{micro}}) + \lambda_e + \lambda_{\text{micro}}) - k^2\mu_e L_c^2(\xi_2^2 + \xi_3^2) \\ k^2\mu_e L_c^2\xi_1\xi_2 \\ k^2\mu_e L_c^2\xi_1\xi_3 \\ 0 \\ -(\lambda_e + \lambda_{\text{micro}}) \\ 0 \\ 0 \\ 0 \\ -(\lambda_e + \lambda_{\text{micro}}) \end{pmatrix}, \end{aligned}$$

$$\begin{aligned}
\tilde{A}_{i5} &= \begin{pmatrix} -ik(\mu_c + \mu_e)\xi_2 \\ ik(\mu_c - \mu_e)\xi_1 \\ 0 \\ k^2\mu_e L_c^2\xi_1\xi_2 \\ \eta\omega^2 - (\mu_c + \mu_e + \mu_{\text{micro}}) - k^2\mu_e L_c^2(\xi_1^2 + \xi_3^2) \\ k^2\mu_e L_c^2\xi_1\xi_2 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \end{pmatrix}, & \tilde{A}_{i6} &= \begin{pmatrix} -ik(\mu_c + \mu_e)\xi_3 \\ 0 \\ ik(\mu_c - \mu_e)\xi_1 \\ k^2\mu_e L_c^2\xi_1\xi_3 \\ k^2\mu_e L_c^2\xi_2\xi_3 \\ \eta\omega^2 - (\mu_c + \mu_e + \mu_{\text{micro}}) - k^2\mu_e L_c^2(\xi_1^2 + \xi_2^2) \\ 0 \\ 0 \\ 0 \\ \mu_c - \mu_e - \mu_{\text{micro}} \\ 0 \\ 0 \end{pmatrix}, \\
\tilde{A}_{i7} &= \begin{pmatrix} ik(\mu_c - \mu_e)\xi_2 \\ -ik(\mu_c + \mu_e)\xi_1 \\ 0 \\ 0 \\ \mu_c - \mu_e - \mu_{\text{micro}} \\ 0 \\ \eta\omega^2 - (\mu_c + \mu_e + \mu_{\text{micro}}) - k^2\mu_e L_c^2(\xi_2^2 + \xi_3^2) \\ k^2\mu_e L_c^2\xi_1\xi_2 \\ k^2\mu_e L_c^2\xi_1\xi_3 \\ 0 \\ 0 \\ 0 \end{pmatrix}, & \tilde{A}_{i8} &= \begin{pmatrix} -ik\lambda_e\xi_1 \\ -ik(2\mu_e + \lambda_e)\xi_2 \\ -ik\lambda_e\xi_3 \\ -\lambda_e - \lambda_{\text{micro}} \\ 0 \\ k^2\mu_e L_c^2\xi_1\xi_2 \\ \eta\omega^2 - (\mu_c + \mu_e + \mu_{\text{micro}}) - k^2\mu_e L_c^2(\xi_1^2 + \xi_3^2) \\ k^2\mu_e L_c^2\xi_2\xi_3 \\ 0 \\ 0 \\ -\lambda_e - \lambda_{\text{micro}} \end{pmatrix}, \\
\tilde{A}_{i9} &= \begin{pmatrix} 0 \\ -ik(\mu_c + \mu_e)\xi_3 \\ ik(\mu_c + \mu_e)\xi_2 \\ 0 \\ 0 \\ 0 \\ k^2\mu_e L_c^2\xi_1\xi_3 \\ k^2\mu_e L_c^2\xi_2\xi_3 \\ \eta\omega^2 - (\mu_c + \mu_e + \mu_{\text{micro}}) - k^2\mu_e L_c^2(\xi_1^2 + \xi_2^2) \\ 0 \\ \mu_c - \mu_e - \mu_{\text{micro}} \\ 0 \end{pmatrix}, & \tilde{A}_{i10} &= \begin{pmatrix} ik(\mu_c - \mu_e)\xi_3 \\ 0 \\ -ik(\mu_c + \mu_e)\xi_1 \\ 0 \\ 0 \\ \mu_c - \mu_e - \mu_{\text{micro}} \\ 0 \\ 0 \\ 0 \\ \eta\omega^2 - (\mu_c + \mu_e + \mu_{\text{micro}}) - k^2\mu_e L_c^2(\xi_2^2 + \xi_3^2) \\ k^2\mu_e L_c^2\xi_1\xi_2 \\ k^2\mu_e L_c^2\xi_1\xi_3 \end{pmatrix}, \\
\tilde{A}_{i11} &= \begin{pmatrix} 0 \\ ik(\mu_c - \mu_e)\xi_3 \\ -ik(\mu_c + \mu_e)\xi_2 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ \mu_c - \mu_e - \mu_{\text{micro}} \\ k^2\mu_e L_c^2\xi_1\xi_2 \\ \eta\omega^2 - (\mu_c + \mu_e + \mu_{\text{micro}}) - k^2\mu_e L_c^2(\xi_1^2 + \xi_3^2) \\ k^2\mu_e L_c^2\xi_2\xi_3 \end{pmatrix}, & \tilde{A}_{i12} &= \begin{pmatrix} -ik\lambda_e\xi_1 \\ -ik\lambda_e\xi_2 \\ -ik(\lambda_e + 2\mu_e)\xi_3 \\ -\lambda_e - \lambda_{\text{micro}} \\ 0 \\ 0 \\ 0 \\ -\lambda_e - \lambda_{\text{micro}} \\ 0 \\ k^2\mu_e L_c^2\xi_1\xi_3 \\ k^2\mu_e L_c^2\xi_2\xi_3 \\ \eta\omega^2 - (\mu_c + \mu_e + \mu_{\text{micro}}) - k^2\mu_e L_c^2(\xi_1^2 + \xi_2^2) \end{pmatrix}.
\end{aligned}$$

It is clear that even with the aid of up-to-date computer algebra systems, it is practically impossible to determine positive-definiteness of the 12×12 acoustic tensor $\tilde{\mathcal{B}}$ in dependence of the given material parameters. In the main body of our paper we succeed by choosing immediately the propagation direction $\xi = e_1$ and by considering a set of new variables (16). This allows us to obtain a certain pre-factorization of $\tilde{\mathcal{B}}(e_1, k)$ in 3×3 blocks. Since the formulation is isotropic, choosing $\xi = e_1$ is no restriction, as argued before.