



HAL
open science

About Non-Fickian Hyperbolic Diffusion

Jean-Louis Auriault, Jolanta Lewandowska, Pascale Royer

► **To cite this version:**

Jean-Louis Auriault, Jolanta Lewandowska, Pascale Royer. About Non-Fickian Hyperbolic Diffusion. CFM 2007 - 18ème Congrès Français de Mécanique, Aug 2007, Grenoble, France. hal-01720704

HAL Id: hal-01720704

<https://hal.science/hal-01720704>

Submitted on 1 Mar 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

About Non-Fickian Hyperbolic Diffusion

Jean-Louis Auriault, Jolanta Lewandowska & Pascale Royer

UJF, INPG, CNRS
Laboratoire Sols Solides Structures-Risques
Domaine Universitaire, BP 53, 38041 GRENOBLE Cedex, France
jean-louis.auriault@hmg.inpg.fr

Abstract :

Fick's law expresses the proportionality of solute flux with respect to concentration gradient. Similar relations are Darcy's law for the fluid flow in porous media, Ohm's law for the electric flux and Fourier's law for heat transfers. When introduced in the corresponding balance equations, these flux laws yield diffusion equations of parabolic character. Different attempts have been made to obtain hyperbolic equations so as to point out propagative phenomena. This was done by adding a time derivative flux term to the flow law. In the paper we focus on solute transport. Two possible non-Fickian diffusion cases are addressed. We firstly investigate diffusion in fluids by a mechanistic approach. Secondly, we study the macroscopic diffusion law in composite materials with large contrast of diffusion coefficient. We show that the obtained diffusion law yields hyperbolicity for drastically small characteristic times or non-propagative waves, respectively.

Résumé :

La loi de Fick exprime la proportionnalité du flux de soluté au gradient de concentration. Les lois de Darcy pour les écoulements en milieu poreux, d'Ohm en électricité et de Fourier pour la chaleur sont de même nature. Introduites dans les équations de bilan correspondantes, elles conduisent à des équations de diffusion paraboliques. Afin d'obtenir des phénomènes propagatifs, un terme de dérivée temporelle du flux est souvent introduit dans les lois d'écoulement. Dans la présente étude, nous étudions deux cas possibles de lois non-fickiennes : la diffusion dynamique de soluté dans un liquide et la diffusion dans un composite. Nous montrons qu'un comportement propagatif est obtenu dans le premier cas, mais pour des temps caractéristiques extrêmement courts, et ne peut être mis en évidence dans le deuxième cas.

Key-words :

Non-Fickian diffusion; propagation; composite material

1 Introduction

Isotropic Fick's law for diffusion of solute in fluids is written

$$\vec{J} = -D\vec{\nabla}c, \quad (1)$$

where \vec{J} is the solute flux, D is the molecular diffusion and c is the solute concentration. When introduced in the solute balance equation

$$\vec{\nabla} \cdot \vec{J} = -\frac{\partial c}{\partial t}, \quad (2)$$

equation (1) yields a diffusion equation of parabolic character

$$\vec{\nabla} \cdot D\vec{\nabla}c = \frac{\partial c}{\partial t}. \quad (3)$$

In such parabolic second order differential equations, information propagates at an infinite velocity, a property which is common to all diffusion processes, i.e., to Fick's, Fourier's, Ohm's or Darcy's processes. To overcome this "non-physical" behaviour, modifications are introduced in these laws so as to render equation (3) hyperbolic. First attempts concerned Fourier's law, see Maxwell (1867); Cattaneo (1958); Vernotte (1958a,b); Chester (1963); Volz *et al.* (1997); Moyne and Degiovanni (2003) and Duhamel (2001) for a more complete bibliography. Ohm's law is also addressed in a similar way, see, e.g., Cuevas *et al.* (1999). Darcy's law poses less difficulty since it is a momentum balance equation whose extension to dynamics is easily demonstrated, Levy (1979); Auriault (1980). That yields to Biot's law, Biot (1956), which shows added mass phenomenon. For solute transport, studies are not numerous. Scheidegger (1960) proposed a hyperbolic diffusion equation, which corresponds to a modified Fick's law, Hassanizadeh (1996). Let us consider the isotropic form of this hyperbolic diffusion law

$$\vec{J} + A \frac{\partial \vec{J}}{\partial t} = -D \vec{\nabla} c, \quad (4)$$

where time $A > 0$ characterizes together with D the solute transport. Relation (4) is similar to Cattaneo-Vernotte law for heat flow, Cattaneo (1958); Vernotte (1958b). When introduced in the solute balance equation, such a non-Fickian law yields a hyperbolic equation which enables to point out finite velocity propagation phenomena: time A is responsible of inertial effects

$$\vec{\nabla} \cdot D \vec{\nabla} c = \frac{\partial c}{\partial t} + A \frac{\partial^2 c}{\partial t^2}. \quad (5)$$

The form of transport equation (4) is often justified as being a consequence of material heterogeneity, Duhamel (2001). Anyhow, Moyne and Degiovanni (2003) have pointed out that a transport equation of propagative type could not be recovered in 1D composite materials. In the paper, we present two attempts to recovering such a non-Fickian law (4). In part 2 we investigate transient solute diffusion in liquids by a mechanistic approach following Einstein's diffusion theory. We demonstrate that such a relation (4) is valid, but for very small characteristic transport characteristic times. Finally we discuss in part 3 the possible non-Fickian character of solute flow through a composite material with high contrast of diffusion coefficients. For this purpose, we revisit the paper by Auriault (1983) where the macroscopic equivalent heat flow law is obtained by upscaling the physics at the heterogeneity scale.

2 Non-Fickian hyperbolic diffusion in liquids

The molecular diffusion is related to the mobility b by the Einstein relation

$$D = kTb, \quad (6)$$

where k is the Boltzmann constant, $k \approx 1.38 \cdot 10^{-23} \text{ J K}^{-1}$ and T is the absolute temperature.

2.1 Quasi-static diffusion coefficient

Consider a solute particle moving at a constant velocity v in a stagnant fluid. The fluid exerts on the particle a force F which is related to v by

$$F = \frac{v}{b}. \quad (7)$$

In case of spherical particle, b is given by the Stokes formula

$$b = \frac{1}{6\pi\mu a}, \quad (8)$$

where μ is the viscosity of the fluid and a is the particle radius. Therefore, we obtain the quasi-static diffusion coefficient in the form

$$D_0 = \frac{kT}{6\pi\mu a}. \quad (9)$$

2.2 Transient diffusion

When the particle velocity is not a constant and depends on time t , relation (7) becomes (see Landau and Lifchitz (1959), §24)

$$F = 2\pi\rho a^3 \left(\frac{1}{3} \frac{dv}{dt} + \frac{3\mu}{\rho a^2} v + \frac{3}{a} \sqrt{\frac{\mu}{\pi\rho}} \int_{-\infty}^t \frac{dv}{d\tau} \frac{d\tau}{\sqrt{t-\tau}} \right). \quad (10)$$

Using (7) to obtain the value of b in the transient case and (6) for D yields the following non-Fickian law

$$-D_0 \nabla \vec{c} = \frac{\rho a^2}{3\mu} \left(\frac{1}{3} \frac{d\vec{J}}{dt} + \frac{3\mu}{\rho a^2} \vec{J} + \frac{3}{a} \sqrt{\frac{\mu}{\pi\rho}} \int_{-\infty}^t \frac{d\vec{J}}{d\tau} \frac{d\tau}{\sqrt{t-\tau}} \right). \quad (11)$$

Relation (11) represents the correct generalization of quasi-static Fick's law (1) to dynamics. It is different from (4), due to the presence of memory effects that are represented by the last term in the right hand member. The memory function is in the form

$$M(t) = \sqrt{\frac{\rho a^2}{\pi\mu t}}, \quad (12)$$

and its characteristic time is given by

$$T_m = \frac{\rho a^2}{\pi\mu}. \quad (13)$$

2.3 Estimation of the domain of validity of non-Fickian relation (11)

Assume that the fluid is water, $\mu = 10^{-3} \text{ kg m}^{-1} \text{ s}^{-1}$, $\rho = 10^3 \text{ kg m}^{-3}$. For $a = 10 \text{ \AA}$, we obtain $T_m \approx 3 \times 10^{-13} \text{ s}$. For $a = 10^{-6} \text{ m}$, we have $T_m \approx 3 \times 10^{-7} \text{ s}$.

On an other hand, the right hand member of relation (11) introduces two dimensionless numbers. Noting T_c the characteristic time of the process yields

$$\mathcal{P} = \frac{\left| \frac{1}{3} \frac{d\vec{J}}{dt} \right|}{\left| \frac{3}{a} \sqrt{\frac{\mu}{\pi\rho}} \int_{-\infty}^t \frac{d\vec{J}}{d\tau} \frac{d\tau}{\sqrt{t-\tau}} \right|} = \frac{\pi}{9} \sqrt{\frac{T_m}{T_c}},$$

$$\mathcal{Q} = \frac{\left| \frac{1}{3} \frac{d\vec{J}}{dt} \right|}{\left| \frac{3\mu}{\rho a^2} \vec{J} \right|} = \frac{\pi T_m}{9 T_c}.$$

We can then point out three different behaviours :

- $T_c \gg T_m$. Relation (11) reduces to Fick's law (1).
- $T_c = \mathcal{O}(T_m)$. The flow law is given by (11) with its three terms in the right hand member. This is a non-Fickian law, but different from (4).

- $T_c \ll T_m$. The two last terms in (11) are negligible. This is a non-Fickian law which yields a hyperbolic behaviour as in (4), but without the first term in the right hand member.

However, as seen above, T_m is very small. Therefore, the two last behaviours are likely not to exist. Remark that the hyperbolic character in the two last behaviours comes from relation (10) where dynamics is considered.

3 Memory effects in composite materials

Consider a periodic composite material of period Ω , made of two constituents of diffusion coefficients D_1 and D_2 which occupy volumes Ω_1 and Ω_2 , respectively. The interface is Γ . Medium 1 is connected whereas medium 2 is connected or not (see Figure 1 where medium 2 is not connected). Both constituents follow Fick's law (1) and equation (3). The characteristic size of the heterogeneities or of the period is noted l , whereas the characteristic size of the macroscopic sample or of the macroscopic solute flux is L and we assume a separation of scales: $l \ll L$. When $D_1 = \mathcal{O}(D_2)$, the macroscopic equivalent behaviour is Fickian (see Auriault (1983) where heat flow in composite materials is investigated): the macroscopic equivalent description is given by

$$\vec{\nabla} \cdot \mathbf{D}^{eff} \vec{\nabla} c = \frac{\partial c}{\partial t},$$

where \mathbf{D}^{eff} is the effective diffusion tensor which depends on both constituents 1 and 2 and on the geometry at the fine scale. $D_2 = \mathcal{O}(D_1 l/L)$ or $D_2 = \mathcal{O}(D_1 (l/L)^3)$ are particular cases of the previous one.

We consider here the case where medium 2 is much less conductive than medium 1

$$\frac{D_2}{D_1} = \mathcal{O}\left(\frac{l}{L}\right)^2 \ll 1.$$

The problem of a macroscopic equivalent behaviour was investigated in Auriault (1983) for




Figure 1: Schematic view of a period (2D case)

heat transfer by using the method of multiscale asymptotic expansions. The reader is referred to this paper for details of the upscaling process.

The upscaling process yields the following equivalent macroscopic behaviour, at the first order of approximation (the model is valid within a relative error $\mathcal{O}(l/L)$)

$$\vec{\nabla} \cdot \mathbf{D}_1^{eff} \vec{\nabla} c = \frac{\partial c}{\partial t} - \int_{-\infty}^t K^{eff}(t - \tau) \frac{\partial^2 c(\tau)}{\partial \tau^2} d\tau. \quad (14)$$

In the above relation, c represents the concentration in medium 1 and \mathbf{D}_1^{eff} is the effective diffusion tensor of the composite material, when considering medium 2 as non diffusive. The macroscopic solute flux is shown to be

$$\vec{J} = -\mathbf{D}_1^{eff} \vec{\nabla} c. \quad (15)$$

The memory function $K^{eff}(t)$ is obtained as the inverse Fourier transform of $\bar{k}/i\omega$, where

$$\bar{k}(\omega) = \frac{1}{|\Omega|} \int_{\Omega_2} k(\omega) dV, \quad (16)$$

and the complex periodic function $k(\omega)$ is the solution of the following boundary value problem in Ω_2

$$\begin{aligned} \vec{\nabla} \cdot D_2 \vec{\nabla} k &= i\omega(k - 1), \\ k &= 0 \quad \text{on } \Gamma. \end{aligned} \quad (17)$$

Illustrations for \bar{k} are given in Auriault (1983) for layered media and for circular cylindrical or spherical inclusions. Important properties of the real and imaginary parts of \bar{k} are that both are positive and verify, Auriault (1983), the following inequalities

$$\begin{aligned} \bar{k} &= \bar{k}_1 + i\bar{k}_2, \\ 0 < \bar{k}_1 &\leq n, \quad 0 < \bar{k}_2 \leq n, \quad n = \frac{|\Omega_2|}{|\Omega|}, \end{aligned} \quad (18)$$

and

$$\lim_{\omega \rightarrow 0} \frac{\bar{k}_2}{\omega} > 0.$$

Relation (14) can be seen as describing a Fickian process in presence of a source term represented by the second term in the right member: the macroscopic equivalent medium is described by solute concentration in medium 1 and Fick's law (15). Medium 2 plays the role of a solute reservoir connected to medium 1. Medium 1 is responsible of the macroscopic diffusion flux. However, the form of the source term in (14) incites us to look for an equivalent description in the form of non-Fickian relation (5). This can be done by considering solute flow at constant frequency ω . Relation (14) then becomes

$$\vec{\nabla} \cdot \mathbf{D}_1^{eff} \vec{\nabla} c = (1 - \bar{k}_1) \frac{\partial c}{\partial t} - \frac{\bar{k}_2}{\omega} \frac{\partial^2 c}{\partial t^2}. \quad (19)$$

We recover (5), but with a different solute capacity $1 - \bar{k}_1 < 1$ and with $A = -\bar{k}_2/\omega < 0$. Therefore, differential equation (19) is elliptic, of different nature from (5) which is hyperbolic. There is no finite velocity propagation in the composite material. This result was also obtained by a different approach in Moyne and Degiovanni (2003) for 1-D heat flow. On the contrary to section 2, no dynamics is present at the heterogeneity scale nor is introduced later. That explain why we do not recover a propagative process at the macroscopic scale. It is likely that some dynamics should be introduced somewhere so as to obtain a propagative process. Remark that the upscaling method in use is rigorous in the sense that the macroscopic description is obtained from the fine scale description, without any prerequisite as it is usual with other methods.

4 Conclusion

Two attempts to justify non-Fickian Cattaneo-Vernotte law for solute diffusion have been conducted, in a view to enable finite velocity propagation of solute perturbations. By analyzing molecular diffusion through its mechanistic Einstein definition, it was shown that such a hyperbolic description is valid, but for transient solute flows with unrealistic very small characteristic times. In composite material with diffusion coefficients of similar or of very different order of magnitude, we did not recover Cattaneo-Vernotte law, but either a parabolic or an elliptic phenomenon.

References

- Auriault, J.-L. 1980 Dynamic behaviour of a porous medium saturated by a Newtonian fluid. *Int. J. Engng. Sc.* **18** 775-785
- Auriault, J.-L. 1983 Effective macroscopic description for heat conduction in periodic composites. *Int. J. Heat Mass Transfer* **26**(6) 861-869
- Biot, M.A. 1956 Theory of propagation of elastic waves in a fluid-saturated porous solid. I Low frequency range *J.A.S.A.* **28** 168-178, II Higher frequency range *J.A.S.A.* **28** 179-191
- Cattaneo, C. 1993 Sur une forme de l'équation de la chaleur éliminant le paradoxe d'une propagation instantanée. *C.R.A.S. Paris* **247** 431-433
- Chester, M. 1963 Second sound in solids. *Phys. Rev.* **131**(5) 2013-2015
- Cuevas, S., del Rio, J.A., Lòpez de Haro, M. 1999 Consequences of a generalized Ohm's law for magnetic transport in conducting media. *J. Phys. D: Appl. Phys.* **32** 639-643
- Duhamel, P. 2001 A new finite integral transform pair for hyperbolic conduction problems in heterogeneous media. *Int. J. Heat Mass Transfer* **44** 3307-3320
- Hassanizadeh, S.M. 1996 On the transient non-Fickian dispersion theory. *Transport in Porous Media* **23** 107-124
- Landau, L., Lifchitz, E. 1959 Fluid Mechanics. Pergamon Press, London.
- Levy, T. 1979 Propagation of waves in a fluid saturated porous elastic solid. *Int. J. Engng. Sc.* **217** 1005-1014
- Maxwell, J.C. 1867 On the dynamical theory of gases. *Phil. Trans. Roy. Soc.* **157** 49-88
- Moyne, C., Degiovanni, A. 2003 Représentation non locale d'un milieu hétérogène en diffusion pure. *C.R.A.S. Paris* **42** 931-942
- Scheidegger, A.E. 1960 The Physics of Flow Through Porous Media. Univ. of Toronto Press, Toronto.
- Vernotte, P. 1958a Les paradoxes de la théorie continue de l'équation de la chaleur. *C.R.A.S. Paris* **246** 3154-3155
- Vernotte, P. 1958b La véritable equation de la chaleur. *C.R.A.S. Paris* **247** 2103-2105
- Volz, S., Lallemand, M., Saulnier, J.-B. 1997 Analyse de la conduction de la chaleur aux temps ultra-courts dans un solide par la thermodynamique irréversible étendue et la dynamique moléculaire. *Rev. Gén. Therm.* **36** 826-835