



**HAL**  
open science

## Coverability Synthesis in Parametric Petri Nets

Nicolas David, Claude Jard, Didier Lime, Olivier Henri Roux

► **To cite this version:**

Nicolas David, Claude Jard, Didier Lime, Olivier Henri Roux. Coverability Synthesis in Parametric Petri Nets. 28th International Conference on Concurrency Theory (CONCUR 2017), Sep 2017, Berlin, Germany. 10.4230/LIPIcs.CONCUR.2017.10 . hal-01724302

**HAL Id: hal-01724302**

**<https://hal.science/hal-01724302v1>**

Submitted on 6 Mar 2018

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Coverability Synthesis in Parametric Petri Nets <sup>\*</sup>

Nicolas David<sup>1</sup>, Claude Jard<sup>2</sup>, Didier Lime<sup>3</sup>, and Olivier H. Roux<sup>4</sup>

- 1 University of Nantes, LS2N, Nantes, France  
nicolas.david1@univ-nantes.fr
- 2 University of Nantes, LS2N, Nantes, France  
claude.jard@univ-nantes.fr
- 3 École Centrale de Nantes, LS2N, Nantes, France  
didier.lime@ec-nantes.fr
- 4 École Centrale de Nantes, LS2N, Nantes, France  
olivier-h.roux@ircyn.ec-nantes.fr

---

## Abstract

We study Parametric Petri Nets (PPNs), i.e., Petri nets for which some arc weights can be parameters. In that setting, we address a problem of parameter synthesis, which consists in computing the exact set of values for the parameters such that a given marking is coverable in the instantiated net.

Since the emptiness of that solution set is already undecidable for general PPNS, we address a special case where parameters are used only as input weights (preT-PPNs), and consequently for which the solution set is downward-closed. To this end, we invoke a result for the representation of upward closed set from Valk and Jantzen. To use this procedure, we show we need to decide universal coverability, that is decide if some marking is coverable for every possible values of the parameters. We therefore provide a proof of its EXPSpace-completeness, thus settling the previously open problem of its decidability.

We also propose an adaptation of this reasoning to the case of parameters used only as output weights (postT-PPNs). In this case, the condition to use this procedure can be reduced to the decidability of the existential coverability, that is decide if there exists values of the parameters making a given marking coverable. This problem is known decidable but we provide here a cleaner proof, providing its EXPSpace-completeness, by reduction to  $\omega$ -Petri Nets.

**1998 ACM Subject Classification** F.1.1 Models of Computation

**Keywords and phrases** Petri net, Parameters, Coverability, Unboundedness, Synthesis

**Digital Object Identifier** 10.4230/LIPIcs.CONCUR.2017.10

## 1 Introduction

The introduction of parameters in models aims to improve genericity. It also allows the designer to leave unspecified aspects, such as those related to the modeling of the environment. This increase in modeling power usually results in greater complexity in the analysis and verification of the model. Beyond verification of properties, the use of parameters opens the way to very relevant issues in design, such as the computation of the parameters values ensuring satisfaction of the expected properties. This is the synthesis problem: given a property, compute the exact set of all parameter values such that, instantiated with these values, the system satisfies this property. This notably permits an estimation of the *robustness*

---

<sup>\*</sup>Work partially supported by ANR project PACS (ANR-14-CE28-0002) and Pays de la Loire research project AFSEC.



of a given instance of a model. Indeed, in full knowledge of “good values” for the parameters, we may be able to quantify the distance from a “bad value” providing an idea of how reliable is the system. Thus parameterised systems are of particular interest both in allowing the handling of more realistic classes of models and addressing more realistic verification issues. We therefore address here the case of parameterised concurrent systems modelled as parametric Petri nets.

**Related work.** The study of parameterised models and more specifically the synthesis has been studied in different parametric settings. Parameters representing delays in timed systems modeled as timed automata have been particularly studied, but with very few decidability results [1]. Synthesis for such systems is only possible in very particular settings, such as bounded integer parameters computed symbolically in timed automata [15] or integer parameters in timed automata with parameters used only as upper bounds, or only as lower bounds, in timing constraints [4]. We focus here on a different type of parameters which represent discrete values. In Petri nets, this corresponds to parameterising the initial marking [5, 17] and the weights of the arcs in transitions [7]. The latter work deals with two decidability problems induced by the use of parameters: The *existential coverability*: does there exist an integer valuation  $v$  on the set of parameters such that  $m$  is coverable in the marked Petri net where parameters are replaced by the value given by  $v$ ? And the *universal coverability*: is  $m$  is coverable in such a net for every possible valuation  $v$ ? Those problems are both undecidable in the most general case, and syntactic subclasses restricting the use of parameters have been introduced, for which the different problems are decidable.

**Contributions.** We focus on computing the exact solution set to the synthesis problem for coverability in parametric Petri nets, i.e., the set of all parameter values such that in the net instantiated with these values, a given marking is coverable.

The emptiness and universality of the solution set being undecidable in general, computing this set can only be done in a restricted setting. We thus focus on the case when parameters are used only as input weights (preT-PPNs) or only as output weights (postT-PPNs). These assumptions give some structure to the solution set: we prove that it is then downward-closed wrt. the usual order on integer vectors for preT-PPNs, and upward-closed for postT-PPNs.

We show how a procedure by Valk and Jantzen from [20] can be used for computing a finite minimal basis of the solution set for postT-PPNs or its complement for preT-PPNs. This requires deciding universal coverability in preT-PPNs and existential coverability in postT-PPNs. The former is an open problem: our main result is it is EXPSpace-complete, which we prove by considering the more generic property of simultaneous unboundedness studied by Demri in [8]. The latter is known decidable but we provide a cleaner proof and additionally prove its EXPSpace-completeness. These results interestingly allow us to carry over a Rackoff upper bound into this parametric setting.

Finally, we prove that in what is called distinctT-PPNs in [7], i.e., when the set of parameters appearing as input weights, and the set of parameters appearing as output weights are disjoint, the solution set cannot be represented using any formalism for which the emptiness of the intersection with equality constraints is decidable.

**Organization of the Paper.** Section 2 gives basic notations and recalls useful mathematical results on orders and Petri nets. Section 3 presents Parametric Petri Nets and recalls some decidability problems. There, we also study the structure of the solution sets for preT- and postT-PPNs and show under which condition Valk and Jantzen’s algorithm can be used to construct finite representation of those sets. In section 4 we give our construction for proving the EXPSpace-completeness of the universal coverability for preT-PPNs. Section 5 revisits the proof of the decidability of existential coverability in postT-PPNs and proves its

EXPSpace-completeness. We also discuss the case of distinct T-PPNs. Finally, in Section 6, we conclude and present future work. Due to space constraints, all proofs can be found in appendix.

## 2 Background

### 2.1 Notations

We denote by  $\mathbb{Z}$  the set of integers, and by  $\mathbb{N}$  the set of natural numbers. As usual,  $\mathbb{N}_\omega$  is the union  $\mathbb{N} \cup \{\omega\}$  where for each  $n \in \mathbb{N}$ ,  $n + \omega = \omega$ ,  $\omega - n = \omega$  and  $\omega \leq \omega$ . Moreover, if  $n \in \mathbb{N}$ ,  $n < \omega$ . Let  $X$  be a finite set. We denote by  $2^X$  the powerset of  $X$  and  $|X|$  the size of  $X$ . If  $X \subseteq \mathbb{N}^k$ ,  $\neg X$  denotes its complement in  $\mathbb{N}^k$ . Given a finite set  $X$ ,  $S_X$  denotes the symmetric group on  $X$  (i.e. the set of all permutations of elements of  $X$ ). Given a set  $X$ , we define a *linear expression* on  $X$  by the following grammar:  $\lambda ::= k \mid k * x \mid \lambda + \lambda$  where  $k \in \mathbb{Z}$ ,  $x \in X$ . We denote by  $\mathcal{L}(X)$  the set of linear expressions on  $X$ .

Let  $V \subseteq \mathbb{N}$ , a  $V$ -valuation for  $X$  is a function from  $X$  to  $V$ . We denote by  $V^X$  the set of  $V$ -valuations on  $X$ . Considering  $v \in V^X$ , we write  $\text{dom}(v)$  the domain of  $v$  ( $X$  in this case) and  $\text{im}(v)$  its image. We refer to  $\mathbb{N}_\omega$ -valuations as *extended valuations* and to  $\mathbb{N}$ -valuations simply as *valuations*. The set  $V^\emptyset$  is reduced to a singleton  $\{\emptyset_V\}$  where  $\emptyset_V$  is the empty function. If  $X$  is finite, considering some arbitrary order on  $X$ , an (extended) valuation can be seen as a vector of size  $|X|$ . For any subset  $X' \subseteq X$  and valuation  $v \in V^X$ , we define the restriction  $v|_{X'}$  of  $v$  to  $X'$  as the unique  $V$ -valuation on  $X'$  such that  $v|_{X'}(x) = v(x)$  for all  $x \in X'$ . We extend this notation to sets of valuations: given  $Y \subseteq V^X$ ,  $Y|_{X'}$  denotes its projection on  $X'$  that is to say  $Y|_{X'} = \{v|_{X'} \mid v \in Y\}$ . Given a value  $a$  of  $\mathbb{N}_\omega$ , we denote as  $\mathbf{a}$  the uniform (extended) valuation that maps every element of  $X$  to  $a$ . Given an extended valuation  $v$ , we write  $\omega(v)$  for the subset of  $X$  such that  $x \in \omega(v)$  iff  $v(x) = \omega$ . We write  $\mathbb{N}(v)$  for the subset of  $X$  such that  $x \in \mathbb{N}(v)$  iff  $v(x) \in \mathbb{N}$ .

Given a linear expression  $\lambda$  on  $X$  and an extended valuation  $v$  on  $X' \subseteq X$ ,  $v(\lambda)$  is the linear expression obtained when substituting each element  $x$  in  $X'$  from  $\lambda$ , by the corresponding value  $v(x)$ . If  $X' = X$  we obtain an element of  $\mathbb{N}_\omega$ .

Given a set  $R$ , finite sets  $S, A, B$  such that  $S = A \cup B$  and  $A \cap B = \emptyset$ , and functions  $f \in R^A$  and  $g \in R^B$ , we write  $f \cup g \in R^S$  the function defined by  $(f \cup g)|_A = f$  and  $(f \cup g)|_B = g$ . We call  $f \cup g$  the *union* of  $f$  and  $g$ . Note that given  $x$  in  $A$ ,  $y = f(x)$  is called the image of  $x$  and when there is no ambiguity (i.e. when  $f$  is injective),  $x$  is called the fiber of  $y$  by  $f$ .

Finally, let  $A$  be an alphabet and  $A^*$  be the free monoid over  $A$ . Let  $w \in A^*$  be a word. We write  $|w|$  the length of  $w$ . Given  $a \in A$ ,  $|w|_a$  is the number of occurrences of  $a$  in  $w$ . We define  $\epsilon$  as the identity element of  $A^*$ . We write  $t \sqsubseteq s$  when  $t$  is a prefix of  $s$ . We denote by  $\text{Pref}(L)$  the prefix closure of a language  $L$ , i.e.  $\text{Pref}(L) = \bigcup_{s \in L} \{t \mid t \sqsubseteq s\}$ .

### 2.2 Order

A *quasi order* (qo for short)  $\lesssim$  on some set  $S$  is a reflexive and transitive binary relation on  $S$ . The pair  $(S, \lesssim)$  is then called a quasiordered set. For  $x, y \in S$  and given a qo  $\lesssim$  on  $S$ ,  $x$  and  $y$  are said *comparable* if either  $x \lesssim y$  or  $y \lesssim x$ . A relation  $<$  is a *strict order* on a set  $S$  if it is irreflexive and transitive (which implies asymmetry).

Given any quasi order  $\lesssim$  on a set  $S$  we can define: (i) a strict order  $<$  given by  $x < y$  iff  $x \lesssim y \wedge \neg(y \lesssim x)$ , (ii) an equivalence relation  $\sim$  given by  $x \sim y$  iff  $x \lesssim y \wedge y \lesssim x$ , (iii) its dual quasi order  $\gtrsim$  given by  $y \gtrsim x$  iff  $x \lesssim y$ . A *well-quasi-ordering* (wqo for short) is a qo  $\lesssim$  on a

set  $S$  such that, for any infinite sequence  $s = x_0, x_1, x_2, \dots$  in  $S$ , there exists indexes  $i < j$  with  $x_i \lesssim x_j$ . Now consider  $\leq$ , the qo on  $\mathbb{N}^k$  component-wise. Formally, for every  $x, y \in \mathbb{N}^k$ , we write that  $x \leq y$  iff for every component  $i$  of  $x$  and  $y$ ,  $x(i) \leq y(i)$ . Dickson's Lemma [9] states that  $(\mathbb{N}^k, \leq)$  is a well-quasi-ordered set. Let us also recall the following lemma<sup>1</sup>:

► **Lemma 1** ([9]). *Let  $p_0, p_1, \dots, p_n, \dots$  be an infinite sequence of elements of  $(\mathbb{N}_\omega)^k$ . Then, there is an infinite sequence  $p_{i_1}, p_{i_2}, \dots, p_{i_n}, \dots$  such that  $p_{i_1} \leq p_{i_2} \leq \dots \leq p_{i_n} \leq \dots$  (with  $i_1 < i_2 < \dots < i_n < \dots$ ).*

### 2.3 Downward and Upward closed sets

We reuse definitions and concepts from [10, 11] which are summed up in [12]. An upward closed set of the well quasi ordered set  $(\mathbb{N}^k, \leq)$  is a subset  $U$  of  $\mathbb{N}^k$  such that if  $x \in U$ ,  $y \in \mathbb{N}^k$  and  $x \leq y$  then  $y \in U$ . The upward closure of a vector  $u$ , written  $\uparrow u$  is the set  $\{m \in \mathbb{N}^k \mid u \leq m\}$ . Given a set  $U$ , we write  $\uparrow U$  for the upward closure of  $U$ , defined as  $\uparrow U = \bigcup_{u \in U} \uparrow u$ . This implies that  $\uparrow U$  is the least upward closed set in which  $U$  is included. Any upward closed set  $U$  can be represented by a finite set  $F$ , called basis, such that  $U = \uparrow F$ . The minimal elements of  $F$  still form a basis of  $U$  independently of  $F$ . This basis is minimal for inclusion among all bases and is thus called the minimal upward basis of  $F$ .

A downward closed set of the well quasi ordered set  $(\mathbb{N}^k, \leq)$  is a subset  $D$  of  $\mathbb{N}^k$  such that if  $x \in D$ ,  $y \in \mathbb{N}^k$  and  $y \leq x$  then  $y \in D$ . The downward closure of a vector  $d$ , written  $\downarrow d$  is the set  $\{m \in \mathbb{N}^k \mid m \leq d\}$ . Given a set  $D$ , we write  $\downarrow D$  the downward closure of  $D$ , defined as  $\downarrow D = \bigcup_{d \in D} \downarrow d$ . This implies that  $\downarrow D$  is the least downward closed set in which  $D$  is included. Moreover, the downward closure of a finite set is finite. To symbolically represent downward closed sets, we use the extension  $\mathbb{N}_\omega^k$ . The definitions remain otherwise the same. If  $D$  is a downward closed set, we can write  $D = \mathbb{N}^k \cap \downarrow F$  where  $F$  is a finite set of  $\mathbb{N}_\omega^k$ . We call  $F$  a downward basis of  $D$ . The maximal elements of  $F$  still form a basis of  $D$  independently of  $F$ . This basis is minimal for the inclusion among all bases and is thus called the minimal downward basis of  $D$ .

We also recall results from [3]: the union and the intersection of two upward (resp. downward) closed sets is an upward (resp. downward) closed set. The complement of an upward closed set is a downward closed set and vice-versa. Given the basis of an upward closed set, it is possible to compute the basis of its complement using for instance the procedure suggested in Example 5 of [14], and vice versa by adapting this procedure.

Finally, Valk and Jantzen proposed in [20] a necessary and sufficient condition, recalled in Lemma 2, to ensure that a finite basis of an upward closed set is effectively computable.

► **Lemma 2** ([20]). *Given an upward closed set  $U \subseteq \mathbb{N}^k$ , a finite basis of  $U$  is effectively computable iff for each  $v \in \mathbb{N}_\omega^k$ , the emptiness of  $\downarrow v \cap U$  is decidable, which is also equivalent to ask whether for all element  $v \in \mathbb{N}_\omega^k$ , it is decidable to answer whether  $\downarrow v \cap \mathbb{N}^k \subseteq \neg U$ .*

### 2.4 Petri Nets

► **Definition 3** (Marked Petri Net). A marked Petri Net (PN) is a pair  $\mathcal{S} = (\mathcal{N}, m_0)$  where  $\mathcal{N} = (P, T, Pre, Post)$  is a Petri net such that  $P$  is a finite set of places of  $\mathcal{S}$ ,  $T$  is a finite set of transitions of  $\mathcal{S}$ ,  $Pre$  and  $Post$  are functions from  $P \times T$  to  $\mathbb{N}$ . A *marking* of  $\mathcal{N}$  is an  $\mathbb{N}$ -valuation on  $P$  and  $m_0 \in \mathbb{N}^P$  is the *initial* marking of  $\mathcal{N}$ .

<sup>1</sup>The existence of such increasing subsequences can also be used as a definition for wqo, which leads to an equivalent notion. Note that a proof can also be found in [16].

Given a transition  $t$  of  $T$ , we define  $Pre(t)$  as the univariate function on  $P$  at the point  $t$  which associates to each  $p$  of  $P$  the weight  $Pre(p, t)$ . We define  $Post(t)$  in a similar way. A transition  $t \in T$  is said *enabled* by a marking  $m$  when  $m \geq Pre(t)$ .

► **Definition 4** (PN Semantics). The semantics of a PN is a transition system  $\mathcal{S}_{\mathcal{T}} = (Q, q_0, \rightarrow)$  where,  $Q = \mathbb{N}^P$ ,  $q_0 = m_0$ ,  $\rightarrow \subseteq Q \times T \times Q$  such that for all  $t \in T$ ,  $m \xrightarrow{t} m' \Leftrightarrow m \geq Pre(t)$  and  $m' = m - Pre(t) + Post(t)$

This relation can be extended to sequences of transitions as follows: (i)  $m \xrightarrow{c} m'$  if  $m = m'$  (ii)  $m \xrightarrow{wt} m'$  if  $\exists m'', m \xrightarrow{w} m'' \wedge m'' \xrightarrow{t} m'$  where  $w \in T^*$  and  $t \in T$ . We write  $\xrightarrow{*}$  the reflexive transitive closure of  $\rightarrow$ , i.e.,  $m \xrightarrow{*} m'$  when there exists  $w \in T^*$  such that  $m \xrightarrow{w} m'$ .

► **Definition 5** (Reachability). Let  $\mathcal{S} = (\mathcal{N}, m_0)$ , where  $\mathcal{N} = (P, T, Pre, Post)$ , a marking  $m$  of  $\mathbb{N}^P$  is reachable in  $\mathcal{S}$  iff  $m_0 \xrightarrow{*} m$ .

The reachability set  $\mathcal{RS}(\mathcal{S})$  of  $\mathcal{S}$  is the set of all reachable markings of  $\mathcal{S}$ .

► **Definition 6** (Coverability). Let  $\mathcal{S} = (\mathcal{N}, m_0)$ , where  $\mathcal{N} = (P, T, Pre, Post)$ , and  $m$  a marking of  $\mathbb{N}^P$ ,  $m$  is coverable in  $\mathcal{S}$  iff  $\exists m' \in \mathcal{RS}(\mathcal{S}), m' \geq m$ .

The coverability set  $\mathcal{CS}(\mathcal{S})$  of  $\mathcal{S}$  is the set of markings coverable in  $\mathcal{S}$ . Coverability is decidable in marked Petri nets [16]. The coverability set is an over approximation of the reachability set in the sense that  $\mathcal{CS}(\mathcal{S}) = \downarrow \mathcal{RS}(\mathcal{S})$ . Given a marked PN  $\mathcal{S}$ , and a marking  $m$ , we denote by  $cov(\mathcal{S}, m) \in \{True, False\}$  the coverability of  $m$  in  $\mathcal{S}$ . In Petri nets, coverability allows to verify safety properties. We recall that the coverability set of a marked Petri net is computable in the sense that its minimal downward basis is computable (see, e.g., [12]).

## 3 Monotonicity in Parametric Petri Nets

### 3.1 Parametric Petri Nets and Parametric Problems

Following [7], we recall the definitions related to marked Parametric Petri Nets (PPNs). We omit the case of parametric initial markings which is a subcase of parametric output weights.

► **Definition 7** (Parametric Petri Net). A marked parametric Petri Net (PPN) is a pair  $\mathcal{S} = (\mathcal{N}, m_0)$  where  $\mathcal{N} = (P, T, Pre, Post, \mathbb{P})$  such that  $P$  is a finite set of places of  $\mathcal{N}$ ,  $T$  is a finite set of transitions of  $\mathcal{N}$ ,  $\mathbb{P}$  is a finite set of parameters of  $\mathcal{N}$ ,  $Pre$  and  $Post$  are functions from  $P \times T$  to  $\mathbb{N} \cup \mathbb{P}$ ,  $m_0$  is the initial marking of  $\mathcal{N}$  belonging to  $\mathbb{N}^P$ .

We define the *parametric transitions* of  $\mathcal{S}$ ,  $\Theta \subseteq T$  as the set of transitions with at least one parameter on an input or output arc:  $\Theta = \{t \in T \mid \exists p \in P \text{ s.t. } Pre(p, t) \in \mathbb{P} \vee Post(p, t) \in \mathbb{P}\}$ . We refer to  $T \setminus \Theta$  as the set of plain transitions in echo to the notations of [13].

Considering an arbitrary ordering on places, parametric markings can be represented as vectors of linear combinations on the set of parameters i.e. from  $\mathcal{L}(\mathbb{P})^{|P|}$ . Similarly,  $Pre$  and  $Post$  can be seen as matrices of  $(\mathbb{N} \cup \mathbb{P})^{|P| \times |T|}$ .

Given a marked PPN  $\mathcal{S} = (\mathcal{N}, m_0)$ , where  $\mathcal{N} = (P, T, Pre, Post, \mathbb{P})$ , for any  $\mathbb{N}$ -valuation  $v$  on a subset  $X$  of  $\mathbb{P}$ , we define the  $v$ -instance of  $\mathcal{S}$  as the marked PPN  $v(\mathcal{S}) = (v(\mathcal{N}), m_0)$  where  $v(\mathcal{N}) = (P, T, v(Pre), v(Post), \mathbb{P} \setminus X)$ . By  $v(Pre)$  and  $v(Post)$  we denote the functions/matrices obtained by replacing in their entries each parameter  $\lambda$  in  $\text{dom}(v)$  by  $v(\lambda)$ . If  $X = \mathbb{P}$ ,  $v(\mathcal{N})$  and  $v(\mathcal{S})$  are respectively a Petri net and a marked Petri net. We also recall subclasses introduced in [7]. Given a PPN,  $\mathcal{N} = (P, T, Pre, Post, \mathbb{P})$ , if  $Pre \in P \times T \rightarrow \mathbb{N}$ , we call it a postT-PPN, whereas if  $Post \in P \times T \rightarrow \mathbb{N}$ , we call it a preT-PPN.

Given a PPN  $\mathcal{S}$ , and a marking  $m$ , we define two basic parametric decision problems: Does there exist a valuation  $v$  such that  $m$  is coverable in  $v(\mathcal{S})$  (Existential coverability) ? Is  $m$  coverable in  $v(\mathcal{S})$  for all valuations  $v$  (Universal coverability) ? Those problems were partly studied in [7]. In particular both *Existential coverability* and *Universal coverability* are proved to be undecidable for the generic class of PPNs. In this paper, we are interested in a more general question:

► **Definition 8** (*coverability-Synthesis problem*). Compute all the valuations  $v$ , such that  $\text{cov}(v(\mathcal{S}), m)$  is true.

We call this set of valuations the coverability synthesis set of a marked PPN  $\mathcal{S}$  and a marking  $m$ , denoted by  $\mathcal{CV}(\mathcal{S}, m)$ . We also call it the solution set to the synthesis problem.

► **Remark.** From any PN  $\mathcal{S}$ , we can build a PPN  $\mathcal{S}'$  by adding an unused parameter. Then checking existential or universal coverability on  $\mathcal{S}'$  is equivalent to checking coverability on  $\mathcal{S}$ . Those parametric problems are therefore EXPSPACE-hard. The same reasoning applies for other properties such as (simultaneous) unboundedness.

### 3.2 Special Structure of the Coverability Synthesis Set for PreT-PPNs and PostT-PPNs

When we restrict the use of parameters to input arcs, we ensure that any marking coverable in a  $v$ -instance remains coverable for any  $v'$ -instance such that  $v' \leq v$ . Intuitively, decreasing the valuation leads to a more permissive firing condition. Symmetrically, when we restrict the use of parameters to output arcs, we ensure that any marking coverable in a  $v$ -instance, remains coverable for any  $v'$ -instance such that  $v \leq v'$ . Intuitively, firing the same parametric transition while increasing the valuation leads to greater (and thus more permissive) markings. Those results are formalized in Lemma 9.

► **Lemma 9.** Let  $\mathcal{S}_{pre}$  and  $\mathcal{S}_{post}$  be respectively a marked preT-PPN and postT-PPN of initial marking  $m_0$  and  $s_0$ .

- For every transitions sequence  $w$  of  $\mathcal{S}_{pre}$  and for every valuation  $v$ , if  $m_0 \xrightarrow{w} m$  in  $v(\mathcal{S}_{pre})$ , then for every valuation  $v' \leq v$ , there exists  $m' \geq m$  such that  $m_0 \xrightarrow{w} m'$  in  $v'(\mathcal{S}_{pre})$ .
- For every transitions sequence  $w$  of  $\mathcal{S}_{post}$  and for every valuation  $v$ , if  $s_0 \xrightarrow{w} s$  in  $v(\mathcal{S}_{post})$ , then for every valuation  $v' \geq v$ , there exists  $s' \geq s$  such that  $s_0 \xrightarrow{w} s'$  in  $v'(\mathcal{S}_{post})$ .


Note that those properties of monotonicity directly provide a notable structure for the solution set of the synthesis for those two subclasses presented in Corollary 10.

► **Corollary 10.** Given  $\mathcal{S}_{pre}$ ,  $\mathcal{S}_{post}$  a marked preT-PPN and a marked postT-PPN respectively and goal markings  $m$  and  $s$  for each of those nets,


- $\mathcal{CV}(\mathcal{S}_{pre}, m)$  is downward closed.
- $\mathcal{CV}(\mathcal{S}_{post}, s)$  is upward closed.

### 3.3 Reduction of Valk and Jantzen Condition for PreT-PPNs and PostT-PPNs

Given a preT-PPN  $\mathcal{S}$  and a marking  $m$ , one way to compute  $\mathcal{CV}(\mathcal{S}, m)$  is thus to find its finite minimal basis. A naive enumeration is not possible however since this set may be infinite. In particular, the strategy that consists in testing for universality and, in the negative case, enumerating until a witness of non coverability is found would in general provide only a



■ **Figure 1**  $\mathcal{S}_1$  where  $\mathbb{P} = \{a, b\}$



■ **Figure 2** Reduction of coverability to the place boundedness

subset of  $\mathcal{CV}(\mathcal{S}, m)$ . In fact, the main difficulty here resides in the fact that the elements of the minimal basis have to be found among the complete lattice induced by  $\leq$  on  $\mathbb{N}_\omega^\mathbb{P}$ .

In order to represent a finite basis of a downward closed set of valuations, we need to extend valuations to  $\mathbb{N}_\omega$ . Given a preT-PPN and an extended valuation  $v$ , we extend the predicate  $\text{cov}(v(\mathcal{S}), m)$  to extended valuations as follows:  $\text{cov}(v(\mathcal{S}), m) \stackrel{\text{def}}{\Leftrightarrow} \forall v' \in \mathbb{N}^{\omega(v)}, \text{cov}(v'(v|_{\mathbb{N}(v)}(\mathcal{S})), m)$ .

Figure 1 presents a preT-PPN with two parameters  $a$  and  $b$ . If we consider the valuation  $v$  defined by  $v(a) = 1$  and  $v(b) = \omega$ ,  $\text{cov}(v(\mathcal{S}), m)$  is therefore equivalent to the universal coverability of  $m$  in  $v|_{\{a\}}(\mathcal{S})$  where  $v|_{\{a\}}$  is a valuation defined by  $v(a) = 1$ , that is to say “can we cover  $m$  in  $\mathbf{1}|_{\{a\}}(\mathcal{S})$  for any value of  $b$ ?”. Note that this extension of  $\text{cov}(v(\mathcal{S}), m)$  is consistent with the classic behavior: if  $\mathbb{N}(v) = \mathbb{P}$ , then  $\text{cov}(v(\mathcal{S}), m)$  asks the coverability of  $m$  in the marked Petri Net  $v(\mathcal{S})$ .

We recall that in postT-PPNs, universal coverability is true iff  $\text{cov}(\mathbf{0}(\mathcal{S}), m)$ . In a similar manner to preT-PPN, we extend the notation of  $\text{cov}$ , as follows: given a postT-PPN and an extended valuation  $v$ , we extend the predicate  $\text{cov}(v(\mathcal{S}), m)$  to extended valuations as follows:  $\neg \text{cov}(v(\mathcal{S}), m) \stackrel{\text{def}}{\Leftrightarrow} \forall v' \in \mathbb{N}^{\omega(v)}, \neg \text{cov}(v'(v|_{\mathbb{N}(v)}(\mathcal{S})), m)$ . This definition is similar to the definition extended for preT-PPNs where coverability has been replaced by non-coverability.

With those extended notations, we now wonder if it is possible to compute a finite basis of  $\mathcal{CV}(\mathcal{S}, m)$  where  $\mathcal{S}$  is a preT-PPN or a postT-PPN and  $m$  a goal marking. To this end, we suggest to use an algorithm by Valk and Jantzen [20] to compute a finite representation of those sets. Nevertheless, to ensure that this algorithm is suitable to our context and that those basis are effectively computable, we need to clarify two points:

- First, this algorithm is used to compute bases of upward closed sets.
- Second, the necessary and sufficient condition recalled in Lemma 2 must be satisfied.

To address the first point, by Corollary 10 notice that in the case of postT-PPNs, the set  $\mathcal{CV}(\mathcal{S}, m)$  is upward closed, and the procedure could be applied directly on it. In the case of preT-PPNs, since  $\mathcal{CV}(\mathcal{S}, m)$  is downward closed, we need to consider  $\neg \mathcal{CV}(\mathcal{S}, m)$ , which is upward closed as recalled in Section 2.3. We also recalled in that section that given a finite basis of an upward closed set, it is possible to compute a finite basis of its complement. It is therefore equivalent to being able to compute a finite basis of the set of valuations for which it is not possible to cover  $m$  or to be able to compute a finite basis of the set of valuations for which it is possible to cover  $m$ . Thus, a finite basis of  $\neg \mathcal{CV}(\mathcal{S}, m)$  is effectively computable iff a finite basis of  $\mathcal{CV}(\mathcal{S}, m)$  is computable. Considering this reasoning, we address the second point through the following Lemma:

- **Lemma 11.** *The Valk and Jantzen condition can be reduced to the following criteria:*
1. *we can compute a finite representation of the coverability synthesis set in preT-PPNs iff universal coverability is decidable in preT-PPNs*
  2. *we can compute a finite representation of the coverability synthesis set in postT-PPNs iff*



*existential coverability is decidable in postT-PPNs*

We now start by focusing on universal coverability in preT-PPNs.

## 4 Universal Coverability in preT-PPNs

We address the problem of universal coverability through that of the more general universal simultaneous unboundedness. We will prove that both are EXPSPACE-complete.

### 4.1 Simultaneous Unboundedness

► **Definition 12** (Simultaneous Unboundedness [8]). Given  $\mathcal{N} = (P, T, Pre, Post)$ , and  $\mathcal{S} = (\mathcal{N}, m_0)$ , considering a subset  $X \subseteq P$ ,  $\mathcal{S}$  is simultaneous  $X$  unbounded if for any  $B \geq 0$ , there is a run  $w$  such that  $m_0 \xrightarrow{w} m$  and for all  $i \in X$ , we have  $m(i) \geq B$ . If  $X$  is reduced to a singleton  $\{p\}$ ,  $\mathcal{S}$  is said  $p$ -unbounded.

► **Remark.** Notice that coverability can be easily reduced to simultaneous unboundedness by the use of an observer as depicted in Figure 2. The transition  $t_{obs}$  has an input condition equal to the marking we want to cover  $m$ . Its output effect provides a token in a place  $p_{obs}$ , that, once is marked, can become unbounded through the firing of  $t_{cumul}$ . With this construction,  $m$  is coverable in the net iff it is simultaneous  $p_{obs}$ -unbounded.

Since there exist polynomial translations from VASS to VAS and PN and from PN to VAS (and VASS) [18, 2], we have the following Theorem, initially stated with VASS in [8].

► **Theorem 13** ([8]). *Simultaneous unboundedness for PNs is EXPSPACE-complete.*

### 4.2 Notion of Incremental Model

To prove the decidability of universal coverability in preT-PPNs, we will prove the decidability of universal simultaneous unboundedness. We will also prove that the latter belong to EXPSPACE. Together with Remark of Section 3.1, we can then conclude that both problems are EXPSPACE-complete.

Formally, given a parametric Petri net, and a set of places  $X$ , the parametric net is *universally simulatenous  $X$  unbounded* iff for every possible valuation  $v$  of its parameters, the  $v$ -instance of this net is simultaneous  $X$  unbounded.

We first show that it is sufficient for a net to be simultaneous unbounded on a set of places in infinitely many instances (under uniform valuations) of the parametric Petri net to be universally simultaneous unbounded on this set of places. Indeed, for any valuation  $v$ , we can find a uniform valuation  $\mathbf{k}$  such that  $v \leq \mathbf{k}$  and apply Lemma 9.

► **Lemma 14.** *Given a marking  $m$  and a marked preT-PPN  $\mathcal{S}$  and  $X$  a subset of places of  $\mathcal{S}$ , the two following propositions are equivalent :*

1.  $(\mathcal{S}, m_0)$  is universally simultaneous  $X$  unbounded
2.  $\{k \in \mathbb{N} \mid (\mathbf{k}(\mathcal{S}), m_0) \text{ is simultaneous } X \text{ unbounded}\}$  is infinite

Note that this Lemma is not directly used in this paper, but is necessary to proof the upcoming Lemma 15. It is indeed an important result since it reduces the infinite set of valuations over which we should investigate to the infinite set of uniform valuation that is totally ordered *i.e.* two elements of this set are always comparable.

We can now address the problem of universal simultaneous unboundedness. To solve this problem, we reduce it to the existence of a classic Petri net built upon our parametric model

satisfying an adequately chosen simultaneous unboundedness property. The classic Petri net is in fact obtained by evaluating a preT-PPN, called incremental net, under the uniform valuation  $\mathbf{0}$ . The incremental net has a polynomial size in the original preT-PPN and it directly depends on the original preT-PPN and a sequence of distinct parametric transitions. This Section is thus driven by the idea that universal simultaneous  $X$  unboundedness on a preT-PPN  $\mathcal{S}$  is equivalent to the existence of a sequence  $\sigma$  of distinct parametric transitions of  $\mathcal{S}$ , such that the incremental model build on  $\mathcal{S}$  and  $\sigma$  evaluated under  $\mathbf{0}$  satisfies a simultaneous unboundedness property depending on  $X$  and  $\sigma$ .

Before providing the theoretical definition, let us consider the main intuition of our construction. If a net is universally simultaneous unbounded on a set of places  $X$ , two main cases are possible: we can either find a path such that the places of  $X$  are unbounded without using any parametric transition, and then the corresponding run works for any valuation, or we need at least one parametric transition.

In the latter case, since there is an infinite number of valuations and a finite number of parametric transitions, using the pigeonhole principle, there is at least one such transition that must be used as the first parametric transition in the run for an infinite number of valuations. The input places of its parametric arcs are therefore not bounded. Thus, the valuation of the input parametric arcs of this transition is not limiting anymore since we can generate an arbitrary large amount of tokens in the corresponding places. Therefore, we will later evaluate<sup>2</sup> those parameters to 0 in order to perform the verification on a classic Petri net.

Nevertheless, we need to ensure that the set of input places of the parametric arcs are not bounded (without using that transition). This is exactly the goal of this incremental model. Indeed, once fired, we could then consider a new net where the first parametric transition can be involved as well as non parametric transitions and investigate for the newly unbounded places. Either we can unbound the places of the goal set  $X$  or we can reuse previous reasoning and choose a new parametric transition that has to be involved in infinitely many instances. What is important to note here is that at each firing of a new parametric transition, that never occurred in the run, we need to ensure that its input places of parametric input arcs were unbounded using only previous transitions of the run and to remember what are the new places that can be unbounded through the use of this new transition. We will now formalize how it is possible to remember the boundedness of the input places of parametric arcs by presenting exhaustively the model of incremental nets.

Given a preT-PPN  $\mathcal{N} = (P, T', Pre, Post, \mathbb{P})$  and a partition of its transitions  $T' = T \cup \Theta$  between its plain and parametric transition, we denote by  $\mathcal{N}_p$  the Petri Net obtained from  $\mathcal{N}$  by removing all transitions of  $\Theta$  from  $\mathcal{N}$ . An example is given in Figure 3. Let us now consider a finite sequence  $\sigma \in Pref(S_\Theta)$  where  $S_\Theta$  is the symmetric group over  $\Theta$  seen as a language. Let  $|T| = m$ ,  $|P| = n$  and  $|\sigma| = k$ . We define the incremental model  $\mathcal{I}$  of  $\mathcal{N}$  along  $\sigma$ . We write  $\mathcal{I} = \text{incr}(\mathcal{N}, \sigma)$  to denote this preT-PPN. This model is illustrated by the example<sup>3</sup> at the right hand side of Figure 3. Its construction consists of the following main steps:

- (i) Consider  $\mathcal{N}_p$  and  $k$  copies of  $\mathcal{N}_p$ , where to each of those  $k + 1$  subnets is associated a global lock place, ensuring that exactly one copy is active at any given instant. The


<sup>2</sup>Note that any other finite valuation would be suitable since the input places of the parametric arcs are unbounded.

<sup>3</sup>The exact meaning of the notations used to refer to the different components of this example will be provided after this informal intuition on the construction.

copies correspond to the black subnet of this example, whereas the global locks  $p_0^i$ 's are depicted in blue and dotted arcs.

- (ii) Add a copy of the first  $i$  transitions of  $\sigma$  to each  $i^{\text{th}}$  copy of  $\mathcal{N}_p$ , for  $1 \leq i \leq k$ .
- (iii) Between the  $i-1^{\text{th}}$  and  $i^{\text{th}}$  subnets, add a copy of the  $i+1^{\text{th}}$  transition of  $\sigma$ , depicted in plain green arcs in Figure 3, for  $1 \leq i \leq k$ . Notice that this copy presents a special behaviour: its input effect impacts the  $i-1^{\text{th}}$  subnet and its lock  $p_0^{i-1}$  whereas its output effect impacts the  $i^{\text{th}}$  subnet and its lock  $p_0^i$ . This ensures that we change of active subnet only after the firing of a the first occurrence of a precise parametric transition.
- (iv) Finally, we ensure that given every copy of a transition, including the intermediate copies that allow to change the active copy, it modifies simultaneously the places in the associated copy as explained above, but also all copies of greater index (*i.e.* those that have not been activated yet). Those arcs ensure that every later subnet always has the “same” marking as the active copy. They are depicted by dashed red arcs in Figure 3. Note that we synchronise the different copies and do not merge them because we use them to remember the order in which the different input places to parametric transitions become unbounded.

Let us suppose we evaluated this incremental model in order to perform an execution. At the beginning of any execution, given a precise subnet, let us say the  $i^{\text{th}}$  subnet, it follows the behavior of the subnets with lower index because of synchronisations introduced in item (iv). Then, once this copy become active, after the firing of a given parametric transition introduced in item (iii), it will dictate the behavior of the global net (and thus of the subnets with greater index through synchronisations). Once the next copy becomes active, our original  $i^{\text{th}}$  subnet cannot change its state anymore. It is now literally an historic state of the global run of the incremental net.



■ **Figure 3** Construction of an Incremental Net3

More formally, the incremental net  $\text{incr}(\mathcal{N}, \sigma)$  is the preT-PPN  $(\mathcal{P}, \mathcal{T}, \text{PRE}, \text{POST}, \mathbb{P})$  such that  $\mathcal{P} = \{p_j^i \mid 0 \leq i \leq k \wedge 0 \leq j \leq n\}$  where  $p_0^i$  represents the lock related to the  $i^{\text{th}}$  copy of  $\mathcal{N}$  whereas  $p_j^i$  with  $j > 0$  represents the copy of the place  $p_j$  of  $\mathcal{P}$  in the  $i^{\text{th}}$  subnet and  $\mathcal{T} = \{t_j^i \mid 0 \leq i \leq k \wedge 1 \leq j \leq m\} \cup \{\theta_j^i \mid 1 \leq j \leq i \leq k\} \cup (\cup_{1 \leq i \leq k} \{\theta_i^0\})$  where  $t_j^i$  represents the copy of the transition  $t_j$  of  $\mathcal{T}$  in the  $i^{\text{th}}$  subnet,  $\theta_j^i$  represents the copy of the transition  $\theta_j$  of  $\Theta$  in the  $i^{\text{th}}$  subnet,  $\theta_i^0$  represents a copy of the transition  $\theta_i$  from  $\sigma$  which is used to change the active copy (from the  $i-1^{\text{th}}$  to the  $i^{\text{th}}$ ).

We define with this construction a net mapping to relate places and transitions from both

models. Given the two nets  $\mathcal{N}$  and  $\mathcal{I}$  defined as above, considering previous notations, for each  $0 \leq i \leq |\sigma|$  we define the application  $f_{\mathcal{N} \rightarrow \mathcal{I}}^i$  that links the original net  $\mathcal{N}$  to its  $i^{\text{th}}$  copy in  $\mathcal{I}$  (except the corresponding lock). We define  $f_{\mathcal{N} \rightarrow \mathcal{I}}^i : T \cup \{\theta_j \in \sigma \mid j \leq i\} \cup P \rightarrow \mathcal{T} \cup \mathcal{P}$  such that for  $t_j \in T$  (resp.  $\theta_j \in \sigma$  and  $p_j \in P$ ),  $f_{\mathcal{N} \rightarrow \mathcal{I}}^i(t_j) = t_j^i$  (resp.  $f_{\mathcal{N} \rightarrow \mathcal{I}}^i(\theta_j) = \theta_j^i$  and  $f_{\mathcal{N} \rightarrow \mathcal{I}}^i(p_j) = p_j^i$ ). We can then define  $f^{-1}$  the function that maps components of the copies of  $\mathcal{N}$  in  $\mathcal{I}$  to their original fiber by the previous application. Formally,  $f^{-1}$  is defined by:  $f^{-1} : \cup_{0 \leq i \leq k} \text{im}(f_{\mathcal{N} \rightarrow \mathcal{I}}^i) \rightarrow T \cup P \cup \Theta$  and associates to  $t_j^i \in \mathcal{T}$  (resp.  $\theta_j^i \in \mathcal{T}$  and  $p_j^i \in \mathcal{P}$ )  $f^{-1}(t_j^i) = t_j$  (resp.  $f^{-1}(\theta_j^i) = \theta_j$  and  $f^{-1}(p_j^i) = p_j$ ). Finally, we define the application  $h_{\mathcal{I} \rightarrow \mathcal{N}} : \cup_{1 \leq i \leq k} \{\theta_i^0\} \subseteq \mathcal{T} \rightarrow \Theta$  that maps the intermediate parametric transitions between each copies of  $\mathcal{N}$  in  $\mathcal{I}$ ,  $\theta_i^0$  to their original fiber from  $\mathcal{N}$  and occurring in  $\sigma$ , that is to say the  $i^{\text{th}}$  transition of  $\sigma$ .

Those applications allow us to define formally the functions PRE and POST. Given  $i'$  and  $j'$ , let  $x_{j'}^{i'}$  denote either  $t_j^i$  or  $\theta_j^i$  from  $\{t_j^i \mid 0 \leq i \leq k \wedge 1 \leq j \leq m\} \cup \{\theta_j^i \mid 1 \leq j \leq i \leq k\}$  in the following expressions:

$$\begin{aligned} \blacksquare \text{PRE}(p_j^i, x_{j'}^{i'}) (\text{resp. POST}(p_j^i, x_{j'}^{i'})) &= \begin{cases} 0 & \text{if } (i < i') \text{ or } (i > i' \text{ and } j = 0) \\ 1 & \text{if } i = i' \text{ and } j = 0 \\ \text{Pre}(f^{-1}(p_j^i), f^{-1}(x_{j'}^{i'})) & \text{otherwise} \\ (\text{resp. Post}(f^{-1}(p_j^i), f^{-1}(x_{j'}^{i'}))) & \text{otherwise} \end{cases} \\ \blacksquare \text{PRE}(p_j^i, \theta_{i'}^0) &= \begin{cases} 0 & \text{if } (i + 1 < i') \text{ or } (i + 1 > i' \text{ and } j = 0) \\ 1 & \text{if } i + 1 = i' \text{ and } j = 0 \\ \text{Pre}(f^{-1}(p_j^i), h^{-1}(\theta_{i'}^0)) & \text{otherwise} \end{cases} \\ \blacksquare \text{POST}(p_j^i, \theta_{i'}^0) &= \begin{cases} 0 & \text{if } (i < i') \text{ or } (i > i' \text{ and } j = 0) \\ 1 & \text{if } i = i' \text{ and } j = 0 \\ \text{Post}(f^{-1}(p_j^i), h^{-1}(\theta_{i'}^0)) & \text{otherwise} \end{cases} \end{aligned}$$

Given a net  $\mathcal{N}$  and the function  $f_{\mathcal{N} \rightarrow \mathcal{I}}^i$  we extend the definition of  $f_{\mathcal{N} \rightarrow \mathcal{I}}^i$  to sets by  $f_{\mathcal{N} \rightarrow \mathcal{I}}^i(X) = \{f^i(x) \mid x \in X\}$  and nets by defining  $f_{\mathcal{N} \rightarrow \mathcal{I}}^i(\mathcal{N})$  as  $(f^i(P), f^i(T), \text{PRE}_{f^i(P) \times f^i(T)}, \text{POST}_{f^i(P) \times f^i(T)}, \mathbb{P})$ . When the context is clear, we omit the subscript  $\mathcal{N} \rightarrow \mathcal{I}$ . As examples,  $f^0(\mathcal{N})$  and  $f^1(\mathcal{N})$  are provided in Figure 3. Finally, we associate to a marking  $m$  of  $\mathcal{N}$  the marking  $f(m)$  defined by for all  $p$  of  $\cup_{0 \leq i \leq k} (f^i(P))$ ,  $f(m)(p) = m(f^{-1}(p))$ . Notice that this ignores the locks introduced in the net. Given the initial marking  $m_0$ , we thus define the initial marking of the incremental net  $\mu_0$  as  $f(m_0)$  for the copies of the places, and 0 in all locks except the first one which receives 1. Formally,  $\mu_0(p_j^i) = m_0(p_j)$  if  $j \neq 0$ , 1 if  $i = j = 0$  and 0 otherwise.

The idea behind this construction is double. First, we can enforce the order of the first occurrence of a parametric transition which is dictated by the sequence  $\sigma$ . Second, we can access the exact amount of tokens stored in a place before the firing of the first occurrence of a parametric transition and thus keep an historic of the state of a run, just before the firing of this parametric transition, through the copies of the original net. Based on those two observations, we will be able to observe if the input places of the parametric arcs of the first occurrence of a parametric transition in a run are bounded or not.

### 4.3 Complexity of Universal Simultaneous Unboundedness

We will now see that universal simultaneous unboundedness can be reduced to the existence of a sequence  $\sigma$  of distinct parametric transitions such that the incremental net built upon this sequence is simultaneous unbounded on an adequately defined set of places. We first provide the intuition behind this statement before providing its formal version. We must ensure that each input place of a parametric arc of the first occurrence of a parametric

transition is unbounded. Based on the previous construction, one can notice that given a parametric transition  $\theta_i$  occurring in  $\sigma$ , its input places are only impacted by the transitions occurring in the first  $i$  copies of the net. Thus, once  $\theta_i$  is fired in the incremental net, the new feasible transitions will not impact the amount of tokens stored in the  $i$  first subnets. We will thus be able to verify if the input places were bounded or not before its firing, by observing the places of the copy that occurs just before the first firing of this transition. For each parametric transition of  $\sigma$ , we should thus verify that the input places in the corresponding copies are unbounded, and finally verify that the places that should be unbounded as part of the original property are indeed unbounded in the last copy of the net and that for each instance of the incremental net under a uniform valuation. Nevertheless, since the corresponding input places of parametric transitions are unbounded, it is sufficient to verify this property for only one instance of the incremental net, and in particular we will later choose the  $\mathbf{0}$ -instance. Indeed, if such a property is verified for any  $\mathbf{k}$ -instance, then, it could be verified for any  $\mathbf{k}'$ -instance (with  $k' > k$ ) by exhibiting the witness run and performing more loops.

► **Lemma 15.** *Let  $\mathcal{N} = (P, T', Pre, Post, \mathbb{P})$  be a preT-PPN, such that  $T' = T \cup \Theta$  where  $\Theta$  represents the parametric transitions of  $\mathcal{N}$  and  $T$  its plain transitions. For every a set of places of  $X \subseteq P$ , the following propositions are equivalent:*

1.  $(\mathcal{N}, m_0)$  is universally simultaneous  $X$  unbounded
2.  $\exists \sigma = t_1, \dots, t_l \in Pref(S_\Theta)$ , considering the incremental model  $\mathcal{I}$  of  $\mathcal{N}$  along  $\sigma$ ,  $\mathcal{I} = \mathit{incr}(\mathcal{N}, \sigma)$ ,  $\exists k \in \mathbb{N}$ ,  $(\mathbf{k}(\mathcal{I}), \mu_0)$  is simultaneous  $f_{\mathcal{N} \rightarrow \mathcal{I}}^l(X) \cup (\bigcup_{t_i \in \sigma} f_{\mathcal{N} \rightarrow \mathcal{I}}^{i-1}(\Pi(t_i)))$  unbounded.
3.  $\exists \sigma = t_1, \dots, t_l \in Pref(S_\Theta)$ , considering the incremental model  $\mathcal{I}$  of  $\mathcal{N}$  along  $\sigma$ ,  $\mathcal{I} = \mathit{incr}(\mathcal{N}, \sigma)$ ,  $\forall k \in \mathbb{N}$ ,  $(\mathbf{k}(\mathcal{I}), \mu_0)$  is simultaneous  $f_{\mathcal{N} \rightarrow \mathcal{I}}^l(X) \cup (\bigcup_{t_i \in \sigma} f_{\mathcal{N} \rightarrow \mathcal{I}}^{i-1}(\Pi(t_i)))$  unbounded.

Following the notations,  $Pref(S_\Theta)$  corresponds to the finite set of sequences of distinct parametric transitions. Remark that  $f_{\mathcal{N} \rightarrow \mathcal{I}}^l(X)$  represents the copy of the places of  $X$  in the last subnet of the  $\mathcal{I}$ . Set  $\bigcup_{t_i \in \sigma} f_{\mathcal{N} \rightarrow \mathcal{I}}^{i-1}(\Pi(t_i))$  is a bit more complex: for each transition  $t_i \in \sigma$ ,  $\Pi(t_i)$  represents the input places of the parametric arcs. We therefore address here the unboundedness of the copies of those places in the corresponding subnet of the incremental net.

**Proof.** We provide here the sketch of the direct implication  $(1) \Rightarrow (3)$  proof. The goal is to find the sequence of parametric transitions along which we construct a Petri net, called the incremental model, handling the *universal simultaneous  $X$  unboundedness* all at once. This proof is done by induction on the number of parametric transitions in the preT-PPN  $\mathcal{N}$ .

- In the base case,  $\mathcal{N}$  is a PN. Therefore the incremental model considered is isomorphic to  $\mathcal{N}$  and the property is immediate.
- In the inductive step, we filter the case where it is possible that the places from  $X$  are unbounded without using parametric transitions. Then, we show that for an infinity of uniform valuation  $\mathbf{k}$ , there is a parametric transition  $\theta$  that can be used as the first parametric transition occurring in a run leading to some simultaneous  $X$  unbounded markings in the coverability tree of  $(\mathbf{k}(\mathcal{N}), m_0)$ .
  1. We then prove that the input places of the parametric input arcs of  $\theta$  must be unbounded in  $(\mathcal{N}_p, m_0)$ , that is to say, there is a marking  $z$  with some  $\omega$ 's on those components in the basis of the coverability set of this net.
  2. We now consider the net where those places have been removed. The transition  $\theta$  is now a plain transition and it is possible to call the induction assumption. We can thus

build an incremental model  $\mathcal{J}$  along a sequence  $\sigma$  for this reduced net.

3. The remaining work is now to build the incremental model  $\mathcal{I}$  along the sequence  $\theta\sigma$  using several mappings and to construct the set  $Y$  accordingly. ◀

From Lemma 15 we can observe that answering the universal simultaneous  $X$  unboundedness on a preT-PPN can be reduced to guessing an element  $\sigma$  of the finite set  $Pref(S_\Theta)$  such that  $(\mathbf{0}(\mathcal{I}), \mu_0)$  is simultaneous  $Y$  unbounded. Since checking simultaneous  $X$  unboundedness can be done in EXPSPACE as recalled in Theorem 13, we obtain a NEXPSPACE procedure. Then, a well-known theorem by Savitch [19] stating that there is therefore a EXPSPACE deterministic procedure answering this problem and the Remark from Section 3.1 allow us to deduce Theorem 16. Note that following Corollary directly comes from Theorem 16 and Theorem 11.

► **Theorem 16.** *The Universal Simultaneous Unboundedness problem for preT-PPNs is EXPSPACE-complete.*

► **Corollary 17.** *Given a marked preT-PPN  $\mathcal{S}$  and a marking  $m$ , we can compute a finite representation of  $\mathcal{CV}(\mathcal{S}, m)$ .*

## 5 Synthesis in postT-PPNs and distinctT-PPNs

### 5.1 Complexity of Existential Coverability in postT-PPNs

We propose here a cleaner proof for the decidability of the existential coverability in postT-PPNs, and provide its complexity. We use a polynomial time transformation<sup>4</sup> from postT-PPN to  $\omega$ PN (see [13]) which preserves existential coverability and invoke a transformation from  $\omega$ PN to PN underlined in [13]. First, we recall definitions and results from [13].

► **Definition 18** ( $\omega$ -Petri Net [13]). An  $\omega$ -Petri Net ( $\omega$ PN) is a tuple  $(P, T, Pre, Post)$  where  $P$  and  $T$  are respectively a finite set of places and transitions.  $Pre$  (resp.  $Post$ ) is a function of  $P \times T$  to  $\mathbb{N}_\omega$  that gives the input (resp. output) effect of a transition  $t$  on a place  $p$ .

► **Definition 19** ( $\omega$ PN Semantics). Given a marking  $m$ , and a transition  $t$  such that  $m \geq Pre(t)$ , firing  $t$  from  $m$  gives a new marking  $m'$  s.t.  $\forall p \in P, m'(p) = m(p) - Pre(p, t) + o$  where  $o = Post(t, p)$  if  $Post(p, t) \in \mathbb{N}$  and  $o \geq 0$  if  $Post(p, t) = \omega$ . We denote this by  $m \xrightarrow{t} m'$ . Thus  $Post(p, t) = \omega$  means that an arbitrary number of tokens are generated in  $p$ .

From this semantics, we can notice that  $\omega$ 's play a role not unlike output parameters, with the crucial difference that their "value" can change along the execution of the net. Let consider a postT-PPN  $\mathcal{N}$  and let us associate to this model the  $\omega$ OPN  $\mathcal{N}'$  such that we replace each parametric arc of  $\mathcal{N}$  by an  $\omega$  arc.

► **Lemma 20** (postT-PPNs to  $\omega$ OPNs). *Let  $\mathcal{N}$  be a postT-PPN (which involves parameters of a set  $\mathbb{P}$ ) and let  $\mathcal{N}'$  be its corresponding  $\omega$ OPN (with the same set of places and transitions) and let  $m_0$  be their common initial marking. Given a marking  $m \in \mathcal{RS}(\mathcal{N}', m_0)$ , there exists a valuation  $v$  such that there exists a marking  $m' \geq m$  with  $m' \in \mathcal{RS}(v(\mathcal{N}), m_0)$ . Moreover<sup>5</sup>,  $\cup_{v \in \mathbb{N}^{\mathbb{P}}} \mathcal{RS}(v(\mathcal{N}), m_0) \subseteq \mathcal{RS}(\mathcal{N}', m_0)$ .*

<sup>4</sup>More specifically we obtain an  $\omega$ -output-PN or  $\omega$ OPN for short, which corresponds to the natural subclass of  $\omega$ PNs where  $Pre \in P \times T \mapsto \mathbb{N}$ .

<sup>5</sup>Notice that this is only an inclusion. Indeed, contrarily to postT-PPNs, in  $\omega$ PNs, the effect of an arc can change along the same execution.

We can thus directly deduce Theorem 21 by reducing existential coverability in postT-PPNs to coverability in  $\omega$ PN which belongs to EXPSPACE by [13]. Note that following Corollary comes from Theorem 21 and Theorem 11.

► **Theorem 21** (Complexity of Existential Coverability). *The existential coverability problem on postT-PPNs is EXPSPACE-complete.*

► **Corollary 22.** *Given a marked postT-PPN  $\mathcal{S}$  and a marking  $m$ , we can compute a finite representation of  $\mathcal{CV}(\mathcal{S}, m)$ .*

## 5.2 Representing the Coverability Synthesis Set for DistinctT-PPNs

Let us finally consider the case of PPNs in which the set of parameters used as input weights, and the set of parameters used as output weights, are disjoint. For this class, called distinctT-PPNs, the emptiness of the solution set to the synthesis problem for coverability is decidable [7]. Interestingly, we can adapt an idea originally used for L/U-automata in [15] to prove that the structure of this set is however much more complex than for preT-PPNs or postT-PPNs. In particular, one cannot represent this set with a finite set, a finite union of downward and/or upward closed sets or a finite union of convex polyhedra.

► **Lemma 23.** *If it can be computed, the solution of the synthesis of coverability in distinctT-PPN cannot, in general, be represented using any formalism for which emptiness of the intersection with equality constraints is decidable.*

## 6 Conclusion

It can be challenging to find meaningful parametric infinite state systems with decidable decision problems. We achieved here to prove a powerful result for two strict syntactical subclasses of parametric Petri nets: interestingly, the set of all valid valuations of parameters, allowing to cover a given marking, is effectively computable for parametric Petri nets where parameters are restricted to only input arcs or only output arcs.

Indeed, we have shown how the computability of the synthesis set for coverability in preT-PPNs and postT-PPNs can be reduced to a decision problem, respectively, universal coverability and existential coverability, which is then used in Valk and Jantzen's procedure. We proved that these two decision problems are both EXPSPACE-complete. Putting the two types of parameters together while forbidding any parameter to be used as both an input and output weight preserves the decidability of the emptiness of the solution set. However, we have proved that, even with this restriction, the solution set can in general not be represented using any formalism for which emptiness of the intersection with equality constraints is decidable, which seems a big restriction in practice.

Future work includes studying (simultaneous) unboundedness for classes other than preT-PPNs that is to say postT-PPNs and distinctT-PPNs. Most problems (such as universal simultaneous unboundedness for postT-PPNs and distinctT-PPNs) can be settled easily by adapting the proofs of [7], except for existential simultaneous unboundedness for postT-PPNs. The translation to  $\omega$ PNs proposed here is not sufficient to conclude its decidability either since we have to ensure that the increasing markings are all reached for a common parameter valuation.

**Acknowledgements.** This work was partially supported by ANR project PACS (ANR-14-CE28-0002) and Pays de la Loire research project AFSEC. We are grateful to anonymous commenters whose helpful remarks have allowed us to improve this paper.

---

**References**

---

- 1 Rajeev Alur, Thomas A. Henzinger, and Moshe Y. Vardi. Parametric real-time reasoning. In *ACM Symposium on Theory of Computing*, pages 592–601, 1993.
- 2 Florent Avellaneda and Rémi Morin. Vector Addition Systems with States vs. Petri Nets. report, Laboratoire d’informatique Fondamentale de Marseille - LIF, October 2012.
- 3 Ahmed Bouajjani and Richard Mayr. Model checking lossy vector addition systems. In *16th Annual Symposium on Theoretical Aspects of Computer (STACS’09)*, pages 323–333, Trier, Germany, March 1999. Springer.
- 4 Laura Bozzelli and Salvatore La Torre. Decision problems for lower/upper bound parametric timed automata. *Formal Methods in System Design*, 35(2):121–151, 2009.
- 5 Giovanni Chiola, Susanna Donatelli, and Giuliana Franceschinis. On Parametric PT nets and their Modelling Power. *12th International Conference Application and Theory of Petri Nets (ICATPN’91)*, page 206, 1991.
- 6 Jean-Michel Couvreur, Denis Poitrenaud, and Pascal Weil. *Branching Processes of General Petri Nets*, pages 129–148. Springer Berlin Heidelberg, Berlin, Heidelberg, 2011.
- 7 Nicolas David, Claude Jard, Didier Lime, and Olivier H. Roux. Discrete parameters in Petri nets. In *The 36th International Conference on Application and Theory of Petri Nets and Concurrency (Petri Nets 2015)*, volume 9115 of *LNCS*, pages 137–156. Springer, Brussels, Belgium, June 2015.
- 8 Stéphane Demri. On selective unboundedness of VASS. *Journal of Computer and System Sciences*, 79(5):689–713, August 2013.
- 9 Leonard Eugene Dickson. Finiteness of the odd perfect and primitive abundant numbers with  $n$  distinct prime factors. *American Journal of Mathematics*, 35(4):413–422, 1913. URL: <http://www.jstor.org/stable/2370405>.
- 10 Alain Finkel and Jean Goubault-Larrecq. Forward analysis for WSTS, part I: Completions. In *26th Annual Symposium on Theoretical Aspects of Computer Science (STACS’09)*, volume 3 of *Leibniz International Proceedings in Informatics*, pages 433–444, Freiburg, Germany, February 2009. Leibniz-Zentrum für Informatik.
- 11 Alain Finkel and Jean Goubault-Larrecq. Forward analysis for WSTS, part II: complete WSTS. *Logical Methods in Computer Science*, 8(3), 2012.
- 12 Alain Finkel and Jérôme Leroux. Recent and simple algorithms for Petri nets. *Software & Systems Modeling*, 14(2):719–725, 2015.
- 13 Gilles Geeraerts, Alexander Heußner, M. Praveen, and Jean-François Raskin.  $\omega$ -Petri nets: Algorithms and complexity. *Fundam. Inform.*, 137(1):29–60, 2015.
- 14 Jean Goubault-Larrecq. On a Generalization of a Result by Valk and Jantzen. Technical report, LSV, 2009.
- 15 Aleksandra Jovanović, Didier Lime, and Olivier H. Roux. Integer Parameter Synthesis for Real-Time Systems. *IEEE Transactions on Software Engineering*, 41(5):445–461, May 2015.
- 16 Richard M. Karp and Raymond E. Miller. Parallel program schemata. *Journal of Computer and System Sciences*, 3(2):147–195, 1969.
- 17 Marco Ajmone Marsan, Gianfranco Balbo, Gianni Conte, Susanna Donatelli, and Giuliana Franceschinis. *Modelling with Generalized Stochastic Petri Nets*. John Wiley & Sons, Inc., New York, NY, USA, 1st edition, 1994.
- 18 Christophe Reutenauer. *The mathematics of Petri nets*. Prentice-Hall, Inc., April 1990.
- 19 Walter J. Savitch. Relationships between nondeterministic and deterministic tape complexities. *Journal of Computer and System Sciences*, 4(2):177–192, April 1970.
- 20 Rüdiger Valk and Matthias Jantzen. The residue of vector sets with applications to decidability problems in Petri nets. *Acta Informatica*, 21(6):643–674, 1985.



In the sequel, given a marked Petri net  $\mathcal{S}$ , we write  $\mathcal{KM}(\mathcal{S})$  to denote its coverability graph obtained by a procedure *à la* Karp and Miller. We write  $\mathcal{BCS}(\mathcal{S})$  for the minimal downward basis of  $\mathcal{CS}(\mathcal{S})$ .

## A Proof of Lemma 9

We only provide the proof for the preT-PPN. The case of postT-PPN can be adapted easily.

Let  $\mathcal{N} = (P, T, Pre, Post, \mathbb{P})$  be a preT-PPN and  $\mathcal{S} = (\mathcal{N}, m_0)$  be a marked preT-PPN. Let  $w \in T^*$  be a transitions sequence such that  $m_0 \xrightarrow{w} m$  in  $v(\mathcal{S})$ . Then for any valuation  $v' \leq v$ ,  $m_0 \xrightarrow{w} m'$  in  $v'(\mathcal{S})$  with  $m' \geq m$ .

**Proof.** Given  $w_v = t_1, t_2, \dots, t_i, \dots, t_n$ , we proceed by induction on the length of  $w_v$

■ *Base case:  $n=0$*

If the sequence is empty, then  $m_0 \geq m_0$  and by definition  $m_0$  is reachable in  $v$  and  $v'$  instances.

■ *Inductive step: Let us assume it is true for a sequence of length  $n$*

Given  $w = t_1, t_2, \dots, t_i, \dots, t_n, t_{n+1}$  We apply the induction hypothesis on  $w_n = t_1, t_2, \dots, t_n$  such that  $m_0 \xrightarrow{w_n} m_n \xrightarrow{t_{n+1}} m_{n+1}$  in  $v(\mathcal{S})$ . By the induction hypothesis, for  $v' \leq v$  we have  $m_0 \xrightarrow{w_n} m'_n \geq m_n$  in  $v'(\mathcal{S})$ . Moreover  $t_{n+1}$  is firable in  $v(\mathcal{S})$  so  $m_n \geq v(Pre(t_{n+1})) \geq v'(Pre(t_{n+1}))$ . So  $m'_n \geq v'(Pre(t_{n+1}))$  and  $m_0 \xrightarrow{w} m'_{n+1}$  in  $v'(\mathcal{S})$ .

Moreover as  $v(Pre(t_{n+1})) \geq v'(Pre(t_{n+1}))$  and  $Post(t_{n+1}) \geq \mathbf{0}$ , then  $m'_{n+1} \geq m_{n+1}$ , so the proposition is true at rank  $n + 1$ .

By induction, it is true for every sequence of length  $n$  which proves Lemma 9. ◀

## B Proof of Lemma 11

### B.1 Proof of Lemma 11 for PreT-PPNs

The two following propositions are equivalent

1. we can compute a finite representation of the coverability synthesis set in preT-PPNs
2. universal coverability is decidable in preT-PPNs

**Proof.** (1)  $\Rightarrow$  (2): Let us consider a marked preT-PPN  $\mathcal{S}$  and a marking  $m$ . We assume a finite basis of  $\mathcal{CV}(\mathcal{S}, m)$  is effectively computable. Thus, we can get a representation of its complement,  $-\mathcal{CV}(\mathcal{S}, m)$  using a procedure similar to the one of Example 5 in [14]. Since  $-\mathcal{CV}(\mathcal{S}, m)$  is upward closed, as the complement of the downward closed set  $\mathcal{CV}(\mathcal{S}, m)$  from Lemma 10, we can invoke the result of Valk and Jantzen from [20] which implies that for all element  $v \in \mathbb{N}_\omega^k$ , it is decidable to answer whether  $\downarrow v \cap -\mathcal{CV}(\mathcal{S}, m) \neq \emptyset$ . This is equivalent to decide whether  $\downarrow v \cap \mathbb{N}^k \subseteq \mathcal{CV}(\mathcal{S}, m)$ . Let us consider  $v = \omega$ ,  $\downarrow v \cap \mathbb{N}^k = \mathbb{N}^k$ . This exactly means that universal coverability of  $m$  in  $\mathcal{S}$  is decidable.

(2)  $\Rightarrow$  (1): We suppose universal coverability in preT-PPNs is decidable. Let us consider a preT-PPN  $\mathcal{S}$  and a marking  $m$ . Then, for any valuation  $v \in \mathbb{N}_\omega^k$ , coverability of  $m$  in  $v(\mathcal{S})$  is decidable. Indeed, if  $v \in \mathbb{N}^k$  it is trivially decidable, otherwise, it means that universal coverability of  $m$  in  $v_{|\mathbb{N}(v)}(\mathcal{S})$  is decidable, which is true by assumption. Thus, for any  $v \in \mathbb{N}_\omega^k$ , it is decidable to answer whether  $\text{cov}(v(\mathcal{S}), m)$  is true, which directly gives whether  $\downarrow v \cap \mathbb{N}^k \subseteq \mathcal{CV}(\mathcal{S}, m)$  by Lemma 9. This is equivalent to answering for all element  $v \in \mathbb{N}_\omega^k$  whether  $\downarrow v \cap -\mathcal{CV}(\mathcal{S}, m) \neq \emptyset$  which is exactly the condition given by Lemma 2 to use Valk and Jantzen algorithm to compute a finite basis of  $-\mathcal{CV}(\mathcal{S}, m)$ . Now, as recalled in the paper, we can use a procedure adapted from the one in Example 5 of [14], to compute a finite basis of  $-(\mathcal{CV}(\mathcal{S}, m))$ , that is to say  $\mathcal{CV}(\mathcal{S}, m)$ .




## B.2 Proof of Lemma 11 for PostT-PPNs

The two following propositions are equivalent

1. we can compute a finite representation of the coverability synthesis set in postT-PPNs
2. existential coverability is decidable in postT-PPNs

For the case of PostT-PPNs, we can carry out a dual development, by directly considering Lemma 2 with the synthesis set as  $U$  and noticing that universal non coverability is equivalent to non existential coverability.

**Proof.** (1)  $\Rightarrow$  (2): Let us consider a marked postT-PPN  $\mathcal{S}$  and a marking  $m$ . Set  $\mathcal{CV}(\mathcal{S}, m)$  being upward closed, we can invoke Valk and Jantzen's result and deduce that for all  $v \in \mathbb{N}_{\omega}^k$ , it is decidable to answer whether  $\downarrow v \cap \mathbb{N}^k \subseteq \neg \mathcal{CV}(\mathcal{S}, m)$ . Let us now notice that non coverability is monotonic in postT-PPNs in the sense that if  $m$  is not coverable in  $v(\mathcal{S})$ , then it is not coverable in any  $v'(\mathcal{S})$  for all valuation  $v' \leq v$ . This is indeed exactly the contrapositive of Lemma 9. Thus  $\downarrow v \cap \mathbb{N}^k \subseteq \neg \mathcal{CV}(\mathcal{S}, m)$  is equivalent to  $\neg \text{cov}(v(\mathcal{S}), m)$ . In particular,  $\neg \text{cov}(\omega(\mathcal{S}), m)$  is decidable, that is to say universal non coverability is decidable in postT-PPNs. We can now conclude by remarking that universal non coverability is equivalent to non existential coverability.


(2)  $\Rightarrow$  (1): We suppose existential coverability in postT-PPNs is decidable. Since universal non coverability is equivalent to non existential coverability, we can now directly adapt the proof of Lemma 11 to get this result. 

## C Proof of Lemma 14

Given a marking  $m$  and a marked preT-PPN  $\mathcal{S}$ , the two following propositions are equivalent:

1.  $(\mathcal{S}, m_0)$  is universally simultaneous  $X$  unbounded
2.  $\{k \in \mathbb{N} \mid (\mathbf{k}(\mathcal{S}), m_0) \text{ is simultaneous } X \text{ unbounded}\}$  is infinite

**Proof.** (1)  $\Rightarrow$  (2): Let  $\mathcal{V} = \{v \in \mathbb{N}^P \mid \exists k \in \mathbb{N}, v = \mathbf{k}\}$ . Set  $\mathcal{V}$  is infinite. Suppose that for all valuations  $v \in \mathbb{N}^P$ ,  $v(\mathcal{S})$  is simultaneous  $X$  unbounded. As  $\mathcal{V} \subseteq \mathbb{N}^P$ , we have in particular that for any  $v \in \mathcal{V}$ ,  $v(\mathcal{S})$  is simultaneous  $X$  unbounded *i.e.* (2).

(2)  $\Rightarrow$  (1): Let  $\mathcal{V} = \{\mathbf{k} \mid k \in \mathbb{N} \text{ and } (\mathbf{k}(\mathcal{S}), m_0) \text{ is simultaneous } X \text{ unbounded}\}$ . Suppose  $\mathcal{V}$  is infinite. Let us consider  $v \in \mathbb{N}^P$ . We define  $k_1 = \max_{\lambda \in \mathbb{P}}(v(\lambda))$  *i.e.* the maximum of  $v$  componentwise. Clearly the number of elements in  $\mathcal{V}$  that are less or equal to  $\mathbf{k}_1$  is finite (they form a bounded subset of  $\mathbb{N}^P$ ), so there exists  $\mathbf{k}_2 \in \mathcal{V}$  such that  $\mathbf{k}_2 \geq \mathbf{k}_1$ . So  $\mathbf{k}_2(\mathcal{S})$  is simultaneous  $X$  unbounded. This means that, for every value  $B > 0$  we can consider a transitions sequence  $w$  s.t. in  $\mathbf{k}_2(\mathcal{S}), m_0 \xrightarrow{w} m_{k_2}$  such that for all  $p \in X$ ,  $m_{k_2}(p) \geq B$ . Therefore by Lemma 9, in  $v(\mathcal{S}), m_0 \xrightarrow{w} m_v \geq m_{k_2} \geq B$ . So  $v(\mathcal{S})$  is simultaneous  $X$  unbounded and we have (1). 

## D Proof of Lemma 15

Let  $\mathcal{N} = (P, T', \text{Pre}, \text{Post}, \mathbb{P})$  be a preT-PPN, such that  $T' = T \cup \Theta$  where  $\Theta$  represents the parametric transitions of  $\mathcal{N}$  and  $T$  its plain transitions. For every a set of places of  $X \subseteq P$ , the following propositions are equivalent:

1.  $(\mathcal{N}, m_0)$  is universally simultaneous  $X$  unbounded

2.  $\exists \sigma = t_1, \dots, t_l \in \text{Pref}(S_\Theta)$ , considering the incremental model  $\mathcal{I}$  of  $\mathcal{N}$  along  $\sigma$ ,  $\mathcal{I} = \text{incr}(\mathcal{N}, \sigma)$ ,  $\exists k \in \mathbb{N}$ ,  $(\mathbf{k}(\mathcal{I}), \mu_0)$  is simultaneous  $X$  unbounded where  $Y = f_{\mathcal{N} \rightarrow \mathcal{I}}^l(X) \cup \left( \bigcup_{t_i \in \sigma} f_{\mathcal{N} \rightarrow \mathcal{I}}^{i-1}(\Pi(t_i)) \right)$ .
3.  $\exists \sigma = t_1, \dots, t_l \in \text{Pref}(S_\Theta)$ , considering the incremental model  $\mathcal{I}$  of  $\mathcal{N}$  along  $\sigma$ ,  $\mathcal{I} = \text{incr}(\mathcal{N}, \sigma)$ ,  $\forall k \in \mathbb{N}$ ,  $(\mathbf{k}(\mathcal{I}), \mu_0)$  is simultaneous  $Y$  unbounded where  $Y = f_{\mathcal{N} \rightarrow \mathcal{I}}^l(X) \cup \left( \bigcup_{t_i \in \sigma} f_{\mathcal{N} \rightarrow \mathcal{I}}^{i-1}(\Pi(t_i)) \right)$ .

Let us first introduce a few definitions. In the course of the proof we will relate parts of the construction using isomorphisms.

► **Definition 24** (extension of [6]). Let  $\mathcal{N} = (P, T, \text{Pre}, \text{Post}, \mathbb{P})$  and  $\mathcal{N}' = (P', T', \text{Pre}', \text{Post}', \mathbb{P}')$  be two PPNs and  $m_0, m'_0$  their respective initial markings. Let  $h : P \cup T \rightarrow P' \cup T'$  be a mapping such that  $h(P) \subseteq P'$  and  $h(T) \subseteq T'$ . We say that  $h$  is a isomorphism (of structure) from  $\mathcal{N}$  to  $\mathcal{N}'$  if  $h$  is a bijection and for each transition  $t \in T$ , we have  $\text{Pre}'(h(t)) = h(\text{Pre}(t))$  and  $\text{Post}'(h(t)) = h(\text{Post}(t))$ .

We extend  $h$  to markings as follows: for all  $p \in \text{im}(h)$ ,  $h(m)(p) = m(h^{-1}(p))$ . If  $m'_0 = h(m_0)$  we call it an isomorphism of net.

Considering the notations of Section 4.2,  $(f_{\mathcal{N} \rightarrow \mathcal{I}}^0(\mathcal{N}), f_{\mathcal{N} \rightarrow \mathcal{I}}^0(m_0))$  and  $(\mathcal{N}_p, m_0)$  are isomorphic, which is immediate by construction.

Given  $t \in \Theta$ ,  $\Pi(t) \subseteq P$  is used to represent the places  $p$  of  $P$  such that  $\text{Pre}(p, t)$  or  $\text{Post}(p, t)$  belongs to  $\mathbb{P}$ .

Finally, note that the notion of simultaneous unboundedness can be easily related to the notion of Coverability Set through the following Lemma:


► **Lemma 25** ([16, 8]). *Given a marked Petri Net  $\mathcal{S}$ , the following propositions are equivalent:*

1.  $\mathcal{S}$  is simultaneous  $X$  unbounded
2.  $\mathcal{KM}(\mathcal{S})$  contains a node labelled by a marking  $z$  such that  $X \subseteq \omega(z)$
3.  $\text{BCS}(\mathcal{S})$  contains a marking  $z$  such that  $X \subseteq \omega(z)$

**Proof.** (1)  $\Rightarrow$  (3): We proceed by strong induction on the number  $|\Theta|$  of parametric transitions in  $\mathcal{N}$ .

- base case: if  $|\Theta| = 0$ ,  $\mathcal{N}$  is a classic Petri Net, which means  $(\mathcal{N}, m_0)$  is simultaneous  $X$  unbounded. We recall that  $\mathcal{I}$  denotes the incremental model of  $\mathcal{N}$  along a sequence  $\sigma$  of parametric transitions. and  $\mu_0$  its associated marking. Since  $|\Theta| = 0$ ,  $\sigma$  must be  $\epsilon$ . Thus,  $\mathcal{I}$  only contains  $f^0(\mathcal{N})$ , which is isomorphic to  $\mathcal{N}_p$ , plus a lock. So for all  $k$ ,  $\mathbf{k}(f_{\mathcal{N} \rightarrow \mathcal{I}}^0(X))$  is simultaneous  $f_{\mathcal{N} \rightarrow \mathcal{I}}^0(X)$  unbounded. Thus (3) holds.
- inductive step: we consider  $n \in \mathbb{N}$  and we assume that given a preT-PPN with  $i \leq n$  parametric transitions, (3) holds. Let us assume  $|\Theta| = n + 1$ . Since  $(\mathcal{N}, m_0)$  is universally simultaneous  $X$  unbounded, two cases arise:
  - either  $(\mathcal{N}_p, m_0)$  is simultaneous  $X$  unbounded. Thus  $\sigma = \epsilon$  satisfies the property (3) as explained in the base case.
  - otherwise,  $(\mathcal{N}_p, m_0)$  is not simultaneous  $X$  unbounded. Nevertheless, by definition of universal simultaneous unboundedness, for every integer  $k \geq 0$ ,  $(\mathbf{k}(\mathcal{N}), m_0)$  is simultaneous  $X$  unbounded. Then, using Lemma 25, there exists a marking  $z_k$  reachable in  $\mathcal{KM}(\mathbf{k}(\mathcal{N}), m_0)$  such that  $X \subseteq \omega(z_k)$ . This means that, in the coverability graph of  $(\mathbf{k}(\mathcal{N}), m_0)$ , there is a run  $\rho_k$  leading to a node labelled by  $z_k$ . This run contains at least one parametric transition, otherwise  $(\mathcal{N}_p, m_0)$  would be simultaneous  $X$  unbounded. We can thus decompose  $\rho_k$  as

$$m_0 \xrightarrow{*} m_k \xrightarrow{\theta_k} m'_k \xrightarrow{*} z_k$$



For the sake of clarity, remark that we do not most of the arcs of the incremental net detailed in Figure 3.

■ **Figure 4** Operations

where  $\theta_k$  is a parametric transition and the prefix  $m_0 \xrightarrow{*} m_k$  contains no parametric transition. Therefore,  $m_k$  belongs to  $CS(\mathcal{N}_p, m_0)$ . Moreover, we know that its basis  $BCS(\mathcal{N}_p, m_0)$  is finite. Thus, using the pigeon hole's principle, there exists  $z$  in  $BCS(\mathcal{N}_p, m_0)$  such that  $z \geq m_k$  for infinitely many  $k$ . We define  $V = \{\mathbf{k} \mid z \geq m_k\}$ . This set is infinite. To each  $\mathbf{k}$  of  $V$ , we can associate a parametric transition  $\theta_k$  which is the first parametric transition involved in the run  $\rho_k$ . Nevertheless, there is a finite number of parametric transitions. We can thus invoke again the pigeon hole's principle and define an infinite subset of  $V$ , called  $V_1$ , such that for all  $k \in V_1$ , the  $\theta_k$ 's are equal, and without loss of genericity, we assume they are all equal to  $\theta_1$ . For every  $k$  of  $V_1$ , for every  $p \in P$ , we know that  $z(p) \geq m_k(p)$ . In particular, we can now assert that, for every  $p \in \Pi(\theta_1)$ ,  $z(p) \geq m_k(p) \geq \mathbf{k}(Pre(p, \theta_1)) = k$  by definition of  $\Pi(\theta_1)$ . Therefore,  $\Pi(\theta_1) \subseteq \omega(z)$ . Notice that since  $\theta_1 \in \Theta$ ,  $\Pi(\theta_1) \neq \emptyset$ .

Let  $\alpha$  be the projection that associates to  $\mathcal{N}$  the Petri net where the places of  $\omega(z)$  have been removed. Through this operation<sup>6</sup>, every arc that start or ends in a place

<sup>6</sup>The relations between the different nets involved in this proof:  $\mathcal{N}$ ,  $\alpha(\mathcal{N})$ ,  $\mathcal{I}$ ,  $\mathcal{J}$ , etc. are summarized in Figure 4.

of  $\omega(z)$  is also removed. This projection can be extended to markings in the sense that given  $m$ ,  $\alpha(m) = m|_{\mathbb{N}(z)}$ . We now consider the new preT-PPN  $\alpha(\mathcal{N})$ . It is in fact equal to  $(\mathbb{N}(z), T', Pre|_{\mathbb{N}(z) \times T'}, Post|_{\mathbb{N}(z) \times T'}, \mathbb{P})$ .

In this setting, we can partition  $T'$  into the set of plain and parametric transitions:  $T' = T_1 \cup \Theta_1$ . Remark that  $\theta_1$  is now an element of  $T_1$  since its parametric arcs have been removed through the restriction to  $\mathbb{N}(z)$ . Thus,  $\alpha(\mathcal{N})$  has strictly less than  $n + 1$  parametric transitions. We also define the operation inverse  $\alpha_\omega^{-1}$  operating on a marking of  $\alpha(\mathcal{N})$  which reintroduces the places removed and valuate them to  $\omega$ . The result is thus an extended marking of  $\mathcal{N}$ .

Let  $\alpha(y)$  be the marking obtained by firing  $\theta_1$  from  $\alpha(z)$ . This operation is well defined since  $\theta_1$  is a plain transition in  $\alpha(\mathcal{N})$ . We can define  $y$  such that  $y = \alpha_\omega^{-1}(\alpha(y))$ .

Moreover, for every  $k \in V_1$ ,  $y \geq m'_k$  by construction and we can fire the suffix of  $\rho_k$ , and obtain by monotonicity, in the coverability graph of  $(\mathbf{k}(\mathcal{N}), y)$ , that there is a run such that  $y \xrightarrow{*} y_k \geq z_k$ . Thus  $X \subseteq \omega(y_k)$ , and  $X \cap \mathbb{N}(z) \subseteq \omega(\alpha(y_k))$ . Therefore by Lemma 25 for every uniform valuation  $\mathbf{k}$  of  $V_1$ ,  $(\mathbf{k}(\alpha(\mathcal{N})), y)$  is simultaneous  $X \cap \mathbb{N}(z)$  unbounded. We can invoke Lemma 14 to deduce that  $(\alpha(\mathcal{N}), \alpha(y))$  is universally simultaneous  $X \cap \mathbb{N}(z)$  unbounded.

As explained previously, this preT-PPN  $\alpha(\mathcal{N})$  has at most  $n$  parametric transitions. By applying the induction hypothesis, there exists a sequence  $\sigma$  of parametric transitions from  $\Theta_1 \subseteq (\Theta \setminus \{\theta_1\})$  satisfying item (3). We define the associated incremental net as  $\mathcal{J}$  and its initial marking  $\zeta_0$ . Following the definition of Section 4.2 we define  $g^0, g^1, \dots, g^{|\sigma|}$  from  $\alpha(\mathcal{N})$  to  $\mathcal{J}$  and given a marking  $m$  of  $\alpha(\mathcal{N})$ , we denote by  $g(m)$  the marking of  $\mathcal{J}$  which corresponds to duplicate  $m$  on each copies of  $\alpha(\mathcal{N})$ . Note that since we start from  $\alpha(\mathcal{N})$  with an initial marking equal to  $\alpha(y)$ , the initial marking  $\zeta_0$  of  $\mathcal{J}$  coincides with  $g(\alpha(y))$  on the copies of the places of  $\alpha(\mathcal{N})$  (*i.e.*, if we omit the locks).

Now, let us consider the sequence  $\sigma' = \theta_1 \sigma$ . We can define the incremental model  $\mathcal{I}$  of  $\mathcal{N}$  along  $\sigma'$  and its associated marking  $\mu_0$  which coincide with  $f(m_0)$  on the copies of the places of  $\mathcal{N}$ . Following the definition of Section 4.2 we define  $f^0, f^1, \dots, f^{|\sigma'|}$  from  $\mathcal{N}$  to  $\mathcal{I}$  and given a marking  $m$  of  $\mathcal{N}$ , we denote by  $f(m)$  the marking of  $\mathcal{I}$  which corresponds to duplicate  $m$  on each copies of  $\mathcal{N}$ . We also define  $h$  that maps the intermediate parametric transitions of  $\mathcal{I}$  that allows to change the active subnet to the original ones of  $\sigma'$ . Remark that  $h^{-1}$  is well defined.

Given  $0 \leq i \leq |\sigma'|$ ,  $f^i(\mathcal{N})$  is isomorphic to a subnet of  $\mathcal{N}$  (indeed, only some parametric transitions are being removed). We can thus extend the definition of  $\alpha$  to each of those nets. We now extend this projection to  $\mathcal{I}$  as the union of the projections on each subnets of  $\mathcal{I}$ . By abuse of notation, we keep calling this operation  $\alpha$ . We also extend its inverse  $\alpha_\omega^{-1}$  operating on a marking of  $\alpha(\mathcal{I})$  which reintroduces the places removed and valuate them to  $\omega$ . The result is thus a marking of  $\mathcal{I}$ .

We define  $\beta$  as the projection that associates to  $\mathcal{I}$  a copy of this net where the following components have been removed:

- \* the places and transitions of  $f^0(\mathcal{N})$
- \* the lock corresponding to  $f^0(\mathcal{N})$
- \* the parametric transition that changes the active subnet from  $f^0(\mathcal{N})$  to  $f^1(\mathcal{N})$ .

As previously we define its inverse  $\beta_0^{-1}$  operating on a marking of  $\beta(\mathcal{I})$  which reintroduces the places removed by  $\beta$  and valuate them to 0, including the lock.

Let us now consider  $\alpha(\beta(\mathcal{I}))$ . This net is isomorphic<sup>7</sup> to  $\mathcal{J}$ . Indeed, given  $0 \leq i \leq |\sigma|$ ,  $g^i(\alpha(\mathcal{N}))$  is isomorphic to  $\alpha(f^{i+1}(\mathcal{N}))$ . In particular for  $i = 0$ , notice that  $\theta_1$  belongs to  $\alpha(\mathcal{N})$  and therein is not a parametric transition since all its parametric arcs have been removed through  $\alpha$ . Therefore, by construction this transition has a copy in  $g^0(\alpha(\mathcal{N}))$ . Moreover, for  $i > 1$  it also has a copy in every  $f^i(\mathcal{N})$  by construction of  $\mathcal{I}$ . Now remark that  $f^0(\mathcal{N})$ , the only subnet that is not isomorphic to one of the  $g^i(\alpha(\mathcal{N}))$ , has been removed in  $\beta(\mathcal{I})$ . Then through  $\alpha$  the subset  $\omega(z)$  of places of  $\mathcal{N}$  that were copied in  $\mathcal{I}$  (and therefore  $\beta(\mathcal{I})$ ) are removed in  $\alpha(\beta(\mathcal{I}))$  to match the set of places of  $\mathcal{J}$ , which is built upon  $\alpha(\mathcal{N})$ .

We can thus denote by  $\iota$  the isomorphism between  $\alpha(\beta(\mathcal{I}))$  and  $\mathcal{J}$ . Given a marking  $m$  of  $\mathcal{N}$ , we thus have that:

$$\iota \circ \alpha \circ \beta \circ f(m) = g \circ \alpha(m)$$

Let us now study the markings of those nets. Given a valuation  $\mathbf{k}$ , we consider  $(\mathbf{k}(\mathcal{J}), \zeta_0)$ . It is simultaneous  $g^{|\sigma|}(X) \cup (\bigcup_{t_i \in \sigma} g^{i-1}(\Pi(t_i)))$  unbounded by the induction hypothesis. Thus, there exists a sequence of transitions  $w$  in its coverability graph leading to a marking  $\eta$  such that  $g^{|\sigma|}(X) \cup (\bigcup_{t_i \in \sigma} g^{i-1}(\Pi(t_i))) \subseteq \omega(\eta)$ . Let us consider  $(\alpha(\beta(\mathcal{I}), \iota^{-1}(\zeta_0))$ , then  $\iota^{-1}(w)$  leads to a marking  $\iota^{-1}(\eta)$  with  $\iota^{-1}(g^{|\sigma|}(X) \cup (\bigcup_{t_i \in \sigma} g^{i-1}(\Pi(t_i)))) \subseteq \omega(\iota^{-1}(\eta))$  by definition of  $\iota^{-1}$ .

Now let us consider  $(\mathbf{k}(\mathcal{I}), \beta_0^{-1}(\alpha_\omega^{-1}(\iota^{-1}(\zeta_0))))$ . Sequence  $\iota^{-1}(w)$  is still firable. Indeed,  $\alpha_\omega^{-1}$  consists in adding  $\omega$ 's which are not blocking for the firing of a transition and will stay unchanged through the firing of the sequence. Moreover  $\beta_0^{-1}$  adds some zeros in the places of  $f^0(\mathcal{N})$  but no transitions of  $f^0(\mathcal{N})$  are used in this sequence, thus no places of  $f^0(\mathcal{N})$  are inputs to a transition of  $\iota^{-1}(w)$ . From those facts, we can conclude that the marking obtained in the coverability graph of  $\mathbf{k}(\mathcal{I})$  after firing  $\iota^{-1}(w)$  is exactly  $\beta_0^{-1}(\alpha_\omega^{-1}(\iota^{-1}(\eta)))$ .

Let us now focus on  $\zeta_0$ . It can be divided in two parts, first the locks, and second the copies of the places of  $\alpha(\mathcal{N})$ . On this second part, it is exactly equal to  $g(\alpha(y))$ . Thus the marking  $\beta_0^{-1}(\alpha_\omega^{-1}(\iota^{-1}(\zeta_0)))$  can be as well divided in two parts: the locks and the copies of the places of  $\mathcal{N}$ . On the first part, it is equal to  $\zeta_0$  completed with 0 in the initial lock. On the second part, it is exactly equal to  $\beta_0^{-1}(\alpha_\omega^{-1}(\iota^{-1}(g(\alpha(y))))$ .

Now, let  $u = \beta_0^{-1}(\alpha_\omega^{-1}(\iota^{-1}(g(\alpha(y))))$  and recall that  $g \circ \alpha = \iota \circ \alpha \circ \beta \circ f$ . We have:

$$\begin{aligned} u &= \beta_0^{-1}(\alpha_\omega^{-1}(\iota^{-1}(\iota(\alpha(\beta(f(y))))))) \\ &= \beta_0^{-1}(\alpha_\omega^{-1}(\alpha(\beta(f(y)))))) && \text{Because } \iota \circ \iota^{-1} \text{ is the identity function} \\ &= \beta_0^{-1}((\beta(f(y)))) && \text{Because } \alpha_\omega^{-1}(\alpha((x)) = x \text{ iff } x \text{ equal } \omega \text{ on every} \\ & && \text{place removed by } \alpha. \text{ Here } y_{|\omega(z)} = \omega \text{ thus } f(y) \\ & && \text{is equal to } \omega \text{ on every copy of } \omega(z). \\ &\leq f(y) && \text{Because } \beta_0^{-1}(\beta((x)) \leq x \text{ since } \beta_0^{-1} \text{ replaces by} \\ & && \text{0 the values of the places affected by } \beta. \end{aligned}$$

Now, recall that  $f^0$  is an isomorphism between  $\mathcal{N}_p$  and  $f^0(\mathcal{N})$ . In  $\mathcal{KM}(\mathcal{N}_p, m_0)$ , there is a run to a marking  $z'$  greater than  $z$  since  $z \in BCS(\mathcal{N}_p, m_0)$ . Therefore, there exists

<sup>7</sup>Informally, considering the notation of the Section 4.2, the application between  $\mathcal{J}$  and  $\alpha(\beta(\mathcal{I}))$  is defined by increasing all superscript of places and transitions by 1.

a run in  $f^0(\mathcal{N})$  starting from  $f^0(m_0)$  leading to a marking  $f^0(z')$ . This run can also be seen as a run  $w_1$  of  $\mathbf{k}(\mathcal{I})$ , going from the initial marking  $\mu_0$  to a marking  $\psi_0$  equal to  $f(z')$  completed with zeros on the locks corresponding to the subnets  $f^i(\mathcal{N})$  for  $1 \leq i \leq |\sigma'|$  and 1 on the lock corresponding to  $f^0(\mathcal{N})$ . From this marking, we can fire  $h^{-1}(\theta_1)$  since  $f^0(\Pi(\theta_1)) \subseteq \omega(\psi_0)$  and thus have  $\mu_0 \xrightarrow{w_1} \psi_0 \xrightarrow{h^{-1}(\theta_1)} \psi_1$  where  $\psi_1$  can be decomposed in two parts, first the locks, and second the copies of the places of  $\mathcal{N}$ . This second part is greater or equal to  $f(y)$  by construction. Thus,  $\psi_1 \geq u$ . We can thus consider monotonicity, and conclude that  $\iota^{-1}(w)$  is firable from  $\psi_1$  such that in the coverability graph of  $\mathbf{k}(\mathcal{I})$ :  $\psi_0 \xrightarrow{w_1} \psi_0 \xrightarrow{h^{-1}(\theta_1)} \psi_1 \xrightarrow{\iota^{-1}(w)} \psi_2$  where  $\psi_2 \geq \iota^{-1}(\eta)$ . Therefore,  $f^{|\sigma'|}(X) \subseteq \omega(\psi_2)$  and  $(\bigcup_{t_i \in \sigma} f^{i-1}(\Pi(t_i))) \subseteq \omega(\psi_2)$ . Moreover,  $f^0(\Pi(\theta_1)) \subseteq \omega(\psi_2)$  since  $f^0(\Pi(\theta_1)) \subseteq \omega(\psi_0)$  and  $\psi_2$  is a successor of  $\psi_0$ . Thus, since  $\sigma' = \theta_1\sigma$  we obtain exactly that  $f^{|\sigma'|}(X) \cup (\bigcup_{t_i \in \sigma'} f_{\mathcal{N} \rightarrow \mathcal{I}}^{i-1}(\Pi(t_i))) \subseteq \omega(\gamma)$  which concludes the induction.

(3)  $\Rightarrow$  (1): Intuitively, the last copy  $f^{|\sigma|}(\mathcal{N})$  of the incremental net  $\mathcal{I}$  of  $\mathcal{N}$  along  $\sigma$  is synchronized with every previous copy. So every sequence performed on  $\mathcal{I}$  can be folded into the last copy of this net. Moreover, this last copy is isomorphic to a subnet of  $\mathcal{N}$  itself. Therefore, the run will be firable in  $\mathcal{N}$  itself. In the sequel, we omit the subscript  $\mathcal{N} \rightarrow \mathcal{I}$  of the functions  $f_{\mathcal{N} \rightarrow \mathcal{I}}^i$  and  $h_{\mathcal{I} \rightarrow \mathcal{N}}$ . We denote by  $P$  the places of  $\mathcal{N}$ .

More formally, we reason on the coverability graph and reuse the notations introduced in Section 4.2. From (3) we can deduce that, given  $\mathbf{k}$ , we can exhibit a run

$$\mu_0 = y_0 \xrightarrow{\pi_0} z_0 \xrightarrow{\tau_1} y_1 \xrightarrow{\pi_1} z_1 \xrightarrow{\tau_2} y_2 \xrightarrow{\pi_2} z_2 \dots z_{l-1} \xrightarrow{\tau_l} y_l \xrightarrow{\pi_l} z_l \text{ where } \begin{cases} \forall 1 \leq i \leq l, \tau_i \in h^{-1}(\Theta) \\ \forall 0 \leq i \leq l, \pi_i \in f^i(T \cup \Theta) \\ f^l(X) \cup (\bigcup_{t_i \in \sigma} f^{i-1}(\Pi(t_i))) \subseteq \omega(z_l) \end{cases}$$

Moreover, by construction of the incremental net, the tokens in places  $f^j(P)$  are created by transitions of the set  $\bigcup_{i \leq j} (f^i(T))$ . Since for each parametric transition  $\theta_j$ ,  $f^{j-1}(\Pi(\theta_j))$  is equal to  $\omega$ , the  $\omega$ 's of  $f^{j-1}(\Pi(\theta_j))$  have to appear between  $\pi_0$  and  $\pi_{j-1}$ . We can thus deduce that

$$\forall 1 \leq j \leq l, f^{j-1}(\Pi(\theta_j)) \subseteq \omega(z_{j-1})$$

Each time a new parametric transition is involved in the run, the input places of its parametric arcs already contain  $\omega$  in this run in the coverability graph. Since  $f^{|\sigma|}(\mathcal{N})$  is synchronized on all the previous copies to mimic their behaviour along the run, it is possible to simply consider the run folded into  $f^{|\sigma|}(\mathcal{N})$ :

$$f^l(m_0) = y_0|_{f^l(P)} \xrightarrow{f^l(f^{-1}(\pi_0))} z_0|_{f^l(P)} \xrightarrow{f^l(h(\tau_1))} y_1|_{f^l(P)} \dots z_{l-1}|_{f^l(P)} \xrightarrow{f^l(h(\tau_l))} y_l|_{f^l(P)} \xrightarrow{f^l(f^{-1}(\pi_l))} z_l|_{f^l(P)}$$

Where  $\omega((z_l)|_{f^l(P)})$  contains  $f^l(X)$  and, as explained above, before each first occurrence of a parametric transition,  $f^l(h(\tau_i)), f^l(\Pi(h(\tau_i))) \subseteq \omega((z_{i-1})|_{f^l(P)})$ .

Now let us recall that  $f^l$  is an isomorphism between a subnet of  $\mathcal{N}$  and  $f^l(\mathcal{N})$ . Therefore, there exists an isomorphic run in the coverability graph of  $(\mathbf{k}(\mathcal{N}), m_0)$  which provides that same set of places equal to  $\omega$ . In particular, since  $X$  is included in this set, we can deduce that  $\mathbf{k}(\mathcal{N})$  is simultaneous  $X$  unbounded. Moreover, by invoking Lemma 14, we can conclude that  $\mathcal{N}$  is universally simultaneous  $X$  unbounded.

(3)  $\Rightarrow$  (2): trivial

(2)  $\Rightarrow$  (3): We reuse here the decomposition used in (3)  $\Rightarrow$  (1) but push it a bit further. Indeed, since the input places of parametric transitions are unbounded, it is sufficient to

verify this property for only one instance of the incremental net. If such a property is verified for any  $\mathbf{k}$ -instance, then, it could be verified for any  $\mathbf{k}''$ -instance (with  $k'' > k$ ) by exhibiting the witness run and performing more loops.

Formally, from (3) we know that, there exists  $\mathbf{k}$ , such that we can exhibit a run

$$\mu_0 = y_0 \xrightarrow{\pi_0} z_0 \xrightarrow{\tau_1} y_1 \xrightarrow{\pi_1} z_1 \xrightarrow{\tau_2} y_2 \xrightarrow{\pi_2} z_2 \dots z_{l-1} \xrightarrow{\tau_l} y_l \xrightarrow{\pi_l} z_l \text{ where } \begin{cases} \forall 1 \leq i \leq l, \tau_i \in h^{-1}(\Theta) \\ \forall 0 \leq i \leq l, \pi_i \in f^i(T \cup \Theta) \\ f^l(X) \cup (\bigcup_{t_i \in \sigma} f^{i-1}(\Pi(t_i))) \subseteq \omega(z_l) \end{cases}$$

Moreover, as explained in (3)  $\Rightarrow$  (1), by construction of the incremental net, the tokens in places  $f^j(P)$  are created by transitions of the set  $\bigcup_{i \leq j} (f^i(T))$ , which was summed up as:

$$\forall 1 \leq j \leq l, f^{j-1}(\Pi(\theta_j)) \subseteq \omega(z_{j-1})$$

Each time a new parametric transition is involved in the run, the input places of its parametric arcs already contain  $\omega$  in this run in the coverability graph (for the active copy and for all the following one, thus for all the input places of this transition if we consider the synchronisations). Therefore, if we consider the same run under a different valuation  $k'' > k$ , we can assert that there will be no difference for the coverability graph since the only effect of the valuation would be to consume more tokens from places that already contains  $\omega$ . Using Lemma 25 we thus directly obtain that  $\forall k'' > k$ ,  $(\mathbf{k}''(\mathcal{I}), \mu_0)$  is simultaneous  $f_{\mathcal{N} \rightarrow \mathcal{I}}^l(X) \cup (\bigcup_{t_i \in \sigma} f_{\mathcal{N} \rightarrow \mathcal{I}}^{i-1}(\Pi(t_i)))$  unbounded.

Finally, supposing that  $\exists k \in \mathbb{N}$ ,  $(\mathbf{k}(\mathcal{I}), \mu_0)$  is simultaneous  $f_{\mathcal{N} \rightarrow \mathcal{I}}^l(X) \cup (\bigcup_{t_i \in \sigma} f_{\mathcal{N} \rightarrow \mathcal{I}}^{i-1}(\Pi(t_i)))$  unbounded, we can easily adapt the result of Lemma 9 to prove that  $\forall k' \leq k$ ,  $(\mathbf{k}'(\mathcal{I}), \mu_0)$  is simultaneous  $f_{\mathcal{N} \rightarrow \mathcal{I}}^l(X) \cup (\bigcup_{t_i \in \sigma} f_{\mathcal{N} \rightarrow \mathcal{I}}^{i-1}(\Pi(t_i)))$  unbounded.

We have thus proved that (2)  $\Rightarrow$  (3).  $\blacktriangleleft$

## E Proof of Theorem 16

*The Universal Simultaneous Unboundedness problem for preT-PPNs is EXPSPACE-complete.*

**Proof.** The size of  $\mathcal{N}$  is defined as usual, plus the number of parameters. The input consists of  $\mathcal{N}$  plus a subset of its places  $X$ . The EXPSPACE-hardness lower bound is a consequence of Remark from Section 3.1.

We now establish the upper bound for the universal simultaneous unboundedness problem. Given a preT-PPN  $\mathcal{N}$ , we construct a non-deterministic Turing machine which guesses an element  $\sigma$  of  $Pref(S_\Theta)$ . Note that there are  $2^{|\mathbb{P}|+1}$  elements possible in  $Pref(S_\Theta)$ .

We then ask whether given  $\mathcal{I} = \text{incr}(\mathcal{N}, \sigma)$  and  $\mu_0$  its associated initial marking,  $(\mathbf{0}(\mathcal{I}), \mu_0)$  is simultaneous  $Y$  unbounded, where  $\mathcal{I}$  is the incremental net of  $\mathcal{N}$  along  $\sigma$  with the initial marking associated  $\mu_0$ , and  $Y$  is a set<sup>8</sup> of size at most polynomial in the size of the input.

Constructing the incremental net can be done in polynomial time because it consists only in filling matrices of size polynomial in the size of the input. It is therefore also only polynomially larger than  $\mathcal{N}$ .

Checking simultaneous  $X$  unboundedness can be done in EXPSPACE as recalled in Theorem 13.

We thus obtain a NEXPSPACE procedure. By a well-known theorem by Savitch [19], there is therefore a EXPSPACE deterministic algorithm for the universal simultaneous unboundedness problem. This problem is thus EXPSPACE-complete.  $\blacktriangleleft$

<sup>8</sup>The exact formula of  $Y$  is given in Lemma 15.



**F Proof of Lemma 20**

Let  $\mathcal{N}$  be a postT-PPN let  $\mathcal{N}'$  be its corresponding  $\omega$ OPN (with the same set of places and transitions) and let  $m_0$  be their initial marking. Given a marking  $m \in \mathcal{RS}(\mathcal{N}', m_0)$ , there exists a valuation  $v$  such that there exists a marking  $m' \geq m$  with  $m' \in \mathcal{RS}(v(\mathcal{N}), m_0)$ . Moreover,  $\cup_{v \in \mathbb{N}^P} \mathcal{RS}(v(\mathcal{N}), m_0) \subseteq \mathcal{RS}(\mathcal{N}', m_0)$ .

**Proof.** We provide here a simple sketch of this proof.

The first point comes directly from the semantics of  $\omega$  Petri nets. Given  $m \in \mathcal{RS}(\mathcal{N}', m_0)$ , there exists a run  $m_0 \xrightarrow{w} m$ . If  $w$  does not contain any  $\omega$ -transition, then  $m \in \mathcal{RS}(\mathcal{N}_p, m_0)$  thus  $m \in \mathcal{RS}(\mathcal{N}, m_0)$ . Otherwise, we consider all  $\omega$ -transitions involved in  $w$  and compute their output effect on this particular run. We can then consider the valuation  $v$  setting all parameters to the maximum of those effects. In  $v(\mathcal{N})$ , the effect of  $w$  is greater or equal than the effect of  $w$  observed previously, by definition of  $v$ . Thus,  $m_0 \xrightarrow{w} m' \geq m$  in  $v(\mathcal{N})$ .

For the second point, given a valuation and a marking reachable within  $v(\mathcal{N})$  through a run  $w$ , it is easy to construct a run  $w'$  composed of the same transitions in the same order in  $\mathcal{N}'$  with the same effect. Indeed, each time an  $\omega$  transition is fired in  $w'$ , the effect of the  $\omega$  output arc should be equal to the value associate through  $v$  to the parameter of the corresponding parametric output arc. ◀

**G Proof of Lemma 21**

*The existential coverability problem on postT-PPNs is EXPSPACE-complete.*

**Proof.** Given  $\mathcal{N}$  a postT-PPN we construct  $\mathcal{N}'$  its translation (in polynomial time) in an  $\omega$ OPN, by replacing each parametric arc by an  $\omega$  arc. By Lemma 20 we can deduce that  $m$  coverable in  $\mathcal{N}'$  iff  $m$  is existentially coverable in  $\mathcal{N}$ : if  $m$  is coverable in  $\mathcal{N}'$  then there exists a valuation  $v$  such that  $m$  is coverable in  $v(\mathcal{N})$  by the first point of Lemma 20. If  $m$  is existentially coverable in  $\mathcal{N}$ , there exists a valuation  $v$  such that there exists  $m' \in \mathcal{RS}(v(\mathcal{N}), m_0)$  with  $m' \geq m$ . Thus,  $m' \in \mathcal{RS}(\mathcal{N}', m_0)$  by the second point of Lemma 20..

So we need only answer coverability of  $m$  in  $\mathcal{N}'$  which is in EXPSPACE. It follows that existential coverability for postT-PPNs is in EXPSPACE.

Moreover, Remark of Section 3.1 provides the EXPSPACE-hardness of this problem, so it is EXPSPACE-complete. ◀

**H Proof of Lemma 23**

*If it can be computed, the solution of the synthesis of coverability in distinctT-PPN cannot, in general, be represented using any formalism for which emptiness of the intersection with equality constraints is decidable.*

**Proof.** We adapt an idea originally used for L/U-automata in [15]. We recall that following [7] existential coverability is undecidable for PPNs. Let us suppose we can compute the set of valuations under which a given marking of  $\mathbb{N}^P$  is coverable in a distinctT-PPN. Then, let us consider a general PPN  $\mathcal{S}$ . For each parameter  $\lambda$  used on both input and output arcs, we replace its occurrence on input arcs by  $\lambda_i$  and on output arcs by  $\lambda_o$ . We have therefore constructed a distinctT-PPN. We then solve the synthesis problem and compute the set of valuations *Good*. Let  $K$  be the set of equality constraints  $\lambda_i = \lambda_o$  for each parameter  $\lambda$  that

was replaced. Clearly deciding whether there exists a valuation  $v$  such that  $m$  is coverable in  $v(\mathcal{S})$  is equivalent to deciding whether  $Good \cap K$  is empty or not. ◀