



HAL
open science

Fixed points in smooth Calogero-Moser spaces

Cédric Bonnafé, Ruslan Maksimau

► **To cite this version:**

Cédric Bonnafé, Ruslan Maksimau. Fixed points in smooth Calogero-Moser spaces. Annales de l'Institut Fourier, 2019, 71, pp.643-678. 10.5802/aif.3404 . hal-01728667v3

HAL Id: hal-01728667

<https://hal.science/hal-01728667v3>

Submitted on 18 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

FIXED POINTS IN SMOOTH CALOGERO-MOSER SPACES

by

CÉDRIC BONNAFÉ & RUSLAN MAKSIMAU

Abstract. — We prove that every irreducible component of the fixed point variety under the action of μ_d in a smooth Calogero-Moser space is isomorphic to a Calogero-Moser space associated with another reflection group.

Consider a finite subgroup $W \subset \mathbf{GL}_{\mathbb{C}}(V)$ of automorphisms of a finite dimensional complex vector space V , generated by reflections and let c be a complex valued function on the set of conjugacy classes of reflections of W . To the triple (V, W, c) , Etingof and Ginzburg [10] have associated an affine variety $\mathcal{Z}_c(V, W)$, the *Calogero-Moser space*, which is defined as the spectrum of the center of the rational Cherednik algebra \mathbf{H}_c “at $t = 0$ ” (see Section 1 for a precise definition). The algebra \mathbf{H}_c carries a \mathbb{Z} -grading (i.e., a \mathbb{C}^{\times} -action), which induces a \mathbb{C}^{\times} -action on the Calogero-Moser space $\mathcal{Z}_c(V, W)$.

We define a *reflection subquotient* of (V, W) to be a pair (V', W') where V' is a subspace of V , $W' \subset \mathbf{GL}_{\mathbb{C}}(V')$ is generated by reflections and there exists a subgroup H of W which stabilizes V' and whose image in $\mathbf{GL}_{\mathbb{C}}(V')$ is exactly W' .

Note that any element in $\mathbf{GL}_{\mathbb{C}}(V)$ normalizing W acts on $\mathcal{Z}_c(V, W)$. In [4, Conjecture FIX], Rouquier and the first author proposed the following conjecture:

Conjecture F. *Let $\sigma \in \mathbf{GL}_{\mathbb{C}}(V)$ be an element of finite order and normalizing W and let \mathcal{X} be an irreducible component of $\mathcal{Z}_c(V, W)^{\sigma}$ (endowed with its reduced scheme structure). Then there exists a reflection subquotient (V', W') of W and a complex valued function c' on the set of conjugacy classes of reflections in W' such that $\mathcal{X} \simeq \mathcal{Z}_{c'}(V', W')$, as varieties endowed with a \mathbb{C}^{\times} -action.*

Note that the conjecture stated in [4, Conjecture FIX] does not give an explicit description of c' in terms of c : it is just mentioned that the map $c \mapsto c'$ should be linear. Note also that, as stated, it might be a little bit optimistic: maybe one should replace \mathcal{X} by its normalization (as it is not clear whether \mathcal{X} is normal or not). A special case of an element σ normalizing W is when σ is a root of unity, viewed as the corresponding homothety on V . The aim of this paper is to prove the following result:

Theorem. *Conjecture F holds if $\mathcal{Z}_c(V, W)$ is smooth and σ is a root of unity.*

The first author is partly supported by the ANR (Project No ANR-16-CE40-0010-01 GeRepMod).

The proof is by case-by-case analysis, as the triples (V, W, c) such that $\mathcal{Z}_c(V, W)$ is smooth are classified (see the works of Etingof-Ginzburg [10], Gordon [12], Bellamy [1]). The classification and the conjecture can be easily reduced to the case where W acts irreducibly on V/V^W and in this case, the smoothness of $\mathcal{Z}_c(V, W)$ implies that W is of type $G(l, 1, n)$ or G_4 in Shephard-Todd classification. The case of G_4 can be handled by computer calculations (see Section 5), while the infinite family case will be handled by using an isomorphism between $\mathcal{Z}_c(V, W)$ and a quiver variety.

Commentary. The motivation for Conjecture F comes from the modular representation theory of finite reductive groups and conjectures of Broué-Malle-Michel about the endomorphism algebra of some Deligne-Lusztig variety (see [7, §1.A] and [8, Conj. 5.7]). Experimentally, there is an astonishing analogy between the combinatorics involved in Broué-Malle-Michel conjectures and the one involved in the geometry of $\mathcal{Z}_c(V, W)^{\mu_a}$.

The results of this paper confirm this analogy in the particular case of the general linear group $G = \mathbf{GL}_n(\mathbb{F}_q)$ (whose Weyl group W is the symmetric group \mathfrak{S}_n). Let us be more precise. Let ℓ be a prime number not dividing q and let d denote the order of q modulo ℓ . Then (see §4):

- The ℓ -blocks of unipotent characters of G are parametrized by the possible d -cores of partitions of n , as well as the irreducible components of $\mathcal{Z}_1(\mathbb{C}^n, \mathfrak{S}_n)^{\mu_a}$.
- Unipotent characters lying in the block \mathcal{B}_γ parametrized by the d -core γ are in bijection with the irreducible characters of $G(d, 1, r)$ (where $r = (n - |\gamma|)/d$). On the other side, the irreducible component \mathcal{X}_γ of $\mathcal{Z}_1(\mathbb{C}^n, \mathfrak{S}_n)^{\mu_a}$ parametrized by γ is isomorphic to $\mathcal{Z}_c(\mathbb{C}^r, G(d, 1, r))$ and so its \mathbb{C}^\times -fixed points are also parametrized by $\text{Irr } G(d, 1, r)$.
- The conjectures of Broué-Malle-Michel say that the unipotent characters lying in \mathcal{B}_γ are exactly the ones appearing in the cohomology of some Deligne-Lusztig variety \mathbf{X}_γ , and that the endomorphism algebra of the cohomology of \mathbf{X}_γ is isomorphic to some Hecke algebra $\mathcal{H}(G(d, 1, r), c'')$ for some explicitly given parameter c'' . Example 4.19 show that $c'' = c'$ (!).

1. Notation and main result

All along this note, we will abbreviate $\otimes_{\mathbb{C}}$ as \otimes . By an algebraic variety, we mean a reduced scheme of finite type over \mathbb{C} .

Set-up. We fix in this paper a \mathbb{C} -vector space V of finite dimension n and a finite subgroup W of $\mathbf{GL}_{\mathbb{C}}(V)$. We set

$$\text{Ref}(W) = \{s \in W \mid \dim_{\mathbb{C}} V^s = n - 1\}$$

and we assume that

$$W = \langle \text{Ref}(W) \rangle.$$

We also fix an element $\sigma \in \mathbf{GL}_{\mathbb{C}}(V)$ of finite order normalizing W .

We set $\varepsilon : W \rightarrow \mathbb{C}^\times$, $w \mapsto \det(w)$. If $s \in \text{Ref}(W)$, we denote by α_s^\vee and α_s two elements of V and V^* respectively such that $V^s = \text{Ker}(\alpha_s)$ and $V^{*s} = \text{Ker}(\alpha_s^\vee)$, where α_s^\vee is viewed as a linear form on V^* .

1.A. Rational Cherednik algebra at $t = 0$. — All along this note, we fix a function $c : \text{Ref}(W) \rightarrow \mathbb{C}$ which is invariant under conjugacy. We define the \mathbb{C} -algebra \mathbf{H}_c to be the quotient of the algebra $T(V \oplus V^*) \rtimes W$ (the semi-direct product of the tensor algebra $T(V \oplus V^*)$ with the group W) by the relations

$$(\mathcal{H}_c) \quad \begin{cases} [x, x'] = [y, y'] = 0, \\ [y, x] = \sum_{s \in \text{Ref}(W)} (\varepsilon(s) - 1) c_s \frac{\langle y, \alpha_s \rangle \langle \alpha_s^\vee, x \rangle}{\langle \alpha_s^\vee, \alpha_s \rangle} s, \end{cases}$$

for all $x, x' \in V^*$, $y, y' \in V$. The algebra \mathbf{H}_c is called the *rational Cherednik algebra at $t = 0$* .

The first commutation relations imply that we have morphisms of algebras $\mathbb{C}[V] \rightarrow \mathbf{H}_c$ and $\mathbb{C}[V^*] \rightarrow \mathbf{H}_c$. Recall [10, Corollary 4.4] that we have an isomorphism of \mathbb{C} -vector spaces

$$(1.1) \quad \mathbb{C}[V] \otimes \mathbb{C}W \otimes \mathbb{C}[V^*] \xrightarrow{\sim} \mathbf{H}_c$$

induced by multiplication (this is the so-called *PBW-decomposition*).

We denote by \mathbf{Z}_c the center of \mathbf{H}_c : it is well-known [10, Lemma 3.5] that \mathbf{Z}_c is an integral domain, which is integrally closed and contains $\mathbb{C}[V]^W$ and $\mathbb{C}[V^*]^W$ as subalgebras (so it contains $\mathbf{P} = \mathbb{C}[V]^W \otimes \mathbb{C}[V^*]^W$), and which is a free \mathbf{P} -module of rank $|W|$. We denote by \mathcal{X}_c the algebraic variety whose ring of regular functions $\mathbb{C}[\mathcal{X}_c]$ is \mathbf{Z}_c : this is the *Calogero-Moser space* associated with the datum (V, W, c) . If necessary, we will write $\mathcal{X}_c(V, W)$ for \mathcal{X}_c .

Example 1.2. — Let $l \geq 1$, let ζ be a primitive l -th root of unity and let $\mu_l = \langle \zeta \rangle$. We assume in this example that $n = 1$ and $W = \langle t \rangle \simeq \mu_l$, where $t(v) = \zeta v$ for all $v \in V$. Then $\text{Ref}(W) = \{t, t^2, \dots, t^{l-1}\}$ and we set for simplification

$$k_j = \frac{1}{l} \sum_{i=1}^{l-1} \zeta^{-i(j-1)} c_{t^i}$$

($0 \leq j \leq l-1$). Note that $k_0 + k_1 + \dots + k_{l-1} = 0$. Then

$$\mathcal{X}_c \simeq \{(x, y, e) \in \mathbb{C}^3 \mid \prod_{i=0}^{l-1} (e - l k_i) = x y\}$$

(see for instance [4, Theorem 18.2.4]). In particular, \mathcal{X}_c is smooth if and only if

$$\prod_{0 \leq i < j \leq l-1} (k_i - k_j) \neq 0.$$

Note also that the inclusion $\mathbf{P} \subset \mathbf{Z}_c$ corresponds to the morphism of algebraic varieties $\mathcal{X}_c \rightarrow \mathbb{C}^2$, $(x, y, e) \mapsto (x, y)$.

The \mathbb{C}^\times -action is given by the formula

$$\xi \cdot (x, y, e) = (\xi^d x, \xi^{-d} y, e),$$

where $\xi \in \mathbb{C}^\times$ and $(x, y, e) \in \mathcal{X}_c$. ■

1.B. Action of the normalizer. — The normalizer $N_{\mathbf{GL}_C(V)}(W)$ of W in $\mathbf{GL}_C(V)$ acts on V naturally, on V^* by the contragredient action and on W by conjugation. This endows $T(V \oplus V^*) \rtimes W$ with an action of $N_{\mathbf{GL}_C(V)}(W)$ and it is easily checked that the bunch of relations (\mathcal{H}_c) is stable under this action. So \mathbf{H}_c inherits an action of $N_{\mathbf{GL}_C(V)}(W)$. In particular, its centre \mathbf{Z}_c also inherits such an action, and this defines an action of $N_{\mathbf{GL}_C(V)}(W)$ on the Calogero-Moser space \mathcal{X}_c .

Theorem 1.3. — *Assume that \mathcal{X}_c is smooth and that σ is a root of unity, and let \mathcal{X} be an irreducible component of \mathcal{X}_c^σ . Then there exists a reflection subquotient (V', W') of (V, W) and a complex-valued map c' on the set of conjugacy classes of reflections of W' such that*

$$\mathcal{X} \simeq \mathcal{X}_{c'}(V', W'),$$

as varieties endowed with a \mathbb{C}^\times -action.

Remark 1.4. — Note that if \mathcal{X} is reduced to a point, then the theorem is easy because in this case, $\mathcal{X} \simeq \mathcal{X}_0(0, 1)$. ■

This Theorem will be proved in Sections 3 and 5.

1.C. Filtration of the group algebra. — If $w \in W$, we set $\text{cod}(w) = \text{codim}_C(V^w)$. We define a filtration $\mathcal{F}_\bullet(\mathbb{C}W)$ of the group algebra of W as follows: let

$$\mathcal{F}_i(\mathbb{C}W) = \bigoplus_{\text{cod}(w) \leq i} \mathbb{C}w.$$

Then

$$\mathbb{C}\text{Id}_V = \mathcal{F}_0(\mathbb{C}W) \subset \mathcal{F}_1(\mathbb{C}W) \subset \dots \subset \mathcal{F}_n(\mathbb{C}W) = \mathbb{C}W = \mathcal{F}_{n+1}(\mathbb{C}W) = \dots$$

is a filtration of $\mathbb{C}W$. For any subalgebra A of $\mathbb{C}W$, we set $\mathcal{F}_i(A) = A \cap \mathcal{F}_i(\mathbb{C}W)$, so that

$$\mathbb{C}\text{Id}_V = \mathcal{F}_0(A) \subset \mathcal{F}_1(A) \subset \dots \subset \mathcal{F}_n(A) = A = \mathcal{F}_{n+1}(A) = \dots$$

is also a filtration of A .

Now, assume that $\mathcal{X}_c(V, W)$ is smooth, let σ be a root of unity and let \mathcal{X} be an irreducible component of $\mathcal{X}_c(V, W)^\sigma$. According to Theorem 1.3, there exists a reflection subquotient (V', W') of (V, W) , a complex-valued map c' on the set of conjugacy classes of reflections of W' and a \mathbb{C}^\times -equivariant isomorphism $i : \mathcal{X}_{c'}(V', W') \xrightarrow{\sim} \mathcal{X}$. We will view i as a closed immersion $i : \mathcal{X}_{c'}(V', W') \hookrightarrow \mathcal{X}_c(V, W)$. By [12, Corollary 5.8], the smoothness of $\mathcal{X}_c(V, W)$ implies that there exists a bijection between $\text{Irr}(W)$ and $\mathcal{X}_c(V, W)^{\mathbb{C}^\times}$. As the fixed point variety of a finite group in a smooth complex algebraic variety is still smooth, this means that \mathcal{X} is smooth, and so $\mathcal{X}_{c'}(V', W')$ is smooth. Applying again [12, Corollary 5.8], we get another bijection between $\text{Irr}(W')$ and $\mathcal{X}_{c'}(V', W')^{\mathbb{C}^\times}$. As $\mathcal{X}^{\mathbb{C}^\times} \subset \mathcal{X}_c(V, W)^{\mathbb{C}^\times}$, this gives an injective map $i_{\mathcal{X}} : \text{Irr}(W') \hookrightarrow \text{Irr}(W)$, depending on \mathcal{X} and the choice of i . This allows to define a surjective morphism of algebras

$$i_{\mathcal{X}}^* : Z(\mathbb{C}W) \longrightarrow Z(\mathbb{C}W')$$

as follows: if $\chi \in \text{Irr}(W)$, let e_χ^W denote the corresponding primitive central idempotent of $\mathbb{C}W$ and set

$$i_{\mathcal{X}}^*(e_\chi^W) = \begin{cases} e_{\chi'}^{W'} & \text{if } \chi' \in \text{Irr}(W') \text{ is such that } i_{\mathcal{X}}(\chi') = \chi, \\ 0 & \text{if } \chi \text{ does not belong to the image of } i_{\mathcal{X}}. \end{cases}$$

Using results of Shan and the first author [5], Theorem 1.3 has the following consequence:

Corollary 1.5. — *If $i \geq 0$, then $i_{\mathfrak{z}}^*(\mathcal{F}_i(Z(CW))) \subset \mathcal{F}_i(Z(CW'))$.*

Proof. — If \mathcal{Y} is a complex algebraic variety endowed with a \mathbb{C}^\times -action, we denote by $H_{\mathbb{C}^\times}^i(\mathcal{Y})$ the i -th group of equivariant cohomology, with coefficients in \mathbb{C} . Let \hbar be an indeterminate and identify $H_{\mathbb{C}^\times}(\mathbf{pt})$ with $\mathbb{C}[\hbar]$ as usual, with \hbar homogeneous of degree 2. Since $\mathcal{Z}_c(V, W)$ is smooth, it follows from [5, Theorem A] that

$$\begin{cases} H_{\mathbb{C}^\times}^{2i+1}(\mathcal{Z}_c(V, W)) = 0 & \text{if } i \geq 0, \\ H_{\mathbb{C}^\times}^{2\bullet} \simeq \text{Rees}_{\mathcal{F}}(Z(CW)) & \text{as } \mathbb{C}[\hbar]\text{-algebras,} \end{cases}$$

where $\text{Rees}_{\mathcal{F}}(Z(CW)) = \bigoplus_{i \geq 0} \hbar^i \mathcal{F}_i(Z(CW)) \subset \mathbb{C}[\hbar] \otimes_{\mathbb{C}} Z(CW)$ is the Rees algebra associated with the filtration $\mathcal{F}_\bullet(Z(CW))$. Using the analogous result for $\mathcal{Z}_{c'}(V', W')$ and the functoriality of equivariant cohomology, we get a morphism of algebras $i^* : \text{Rees}_{\mathcal{F}}(Z(CW)) \rightarrow \text{Rees}_{\mathcal{F}}(Z(CW'))$.

Now, we will use not only [5, Theorem A] but also its proof: the proof goes by restriction to the fixed point subvariety $\mathcal{Z}_c(V, W)^{\mathbb{C}^\times}$ and identifying $H_{\mathbb{C}^\times}^{2\bullet}(\mathcal{Z}_c(V, W)^{\mathbb{C}^\times})$ with $\mathbb{C}[\hbar] \otimes Z(CW)$ using [12, Corollary 5.8]. As the same strategy holds for $\mathcal{Z}_{c'}(V', W')$, the functoriality of equivariant cohomology implies that the map i^* fits into a commutative diagram

$$\begin{array}{ccc} \text{Rees}_{\mathcal{F}}(Z(CW)) & \xrightarrow{i^*} & \text{Rees}_{\mathcal{F}}(Z(CW')) \\ \downarrow & & \downarrow \\ \mathbb{C}[\hbar] \otimes_{\mathbb{C}} Z(CW) & \xrightarrow{\text{Id}_{\mathbb{C}[\hbar]} \otimes_{\mathbb{C}} i_{\mathfrak{z}}^*} & \mathbb{C}[\hbar] \otimes_{\mathbb{C}} Z(CW'). \end{array}$$

This shows the corollary. □

In Remark 4.18 and Theorem 4.21, we will describe combinatorially the injective map $i_{\mathfrak{z}} : \text{Irr}(W') \hookrightarrow \text{Irr}(W)$ whenever W is of type $G(l, 1, n)$.

2. Preliminaries on quiver varieties

2.A. Quiver varieties. — Let Q_l denote the cyclic quiver with l vertices, defined as follows:

- Vertices: $i \in \mathbb{Z}/l\mathbb{Z}$;
- Arrows: $y_i : i \rightarrow i+1, i \in \mathbb{Z}/l\mathbb{Z}$.

We denote by \overline{Q}_l the double quiver of Q_l that is, the quiver obtained from Q_l by adding an arrow $x_i : i+1 \rightarrow i$ for all $i \in \mathbb{Z}/l\mathbb{Z}$ (see Figure 1).




Fig. 1. The quiver \bar{Q}_l

Now, if $d = (d_i)_{i \in \mathbb{Z}/l\mathbb{Z}}$ is a family of elements of \mathbb{N} , we denote by $\mathbf{GL}(d)$ the direct product

$$\mathbf{GL}(d) = \prod_{i \in \mathbb{Z}/l\mathbb{Z}} \mathbf{GL}_{d_i}(\mathbb{C}),$$

by $\Delta\mathbb{C}^\times$ the image of \mathbb{C}^\times in $\mathbf{GL}(d)$ through the diagonal embedding $\Delta : \mathbb{C}^\times \hookrightarrow \mathbf{GL}(d)$ and we set

$$\mathbf{PGL}(d) = \mathbf{GL}(d) / \Delta\mathbb{C}^\times.$$

The group $\mathbf{PGL}(d)$ acts on the variety $\text{Rep}(\bar{Q}_l, d)$ of representations of \bar{Q}_l in the family of vector spaces $(\mathbb{C}^{d_i})_{i \in \mathbb{Z}/l\mathbb{Z}}$. The orbits are the isomorphism classes of representations of \bar{Q}_l of dimension vector d . We denote by

$$\begin{aligned} \mu_d : \text{Rep}(\bar{Q}_l, d) &\longrightarrow \bigoplus_{i \in \mathbb{Z}/l\mathbb{Z}} \text{Mat}_{d_i}(\mathbb{C}) \\ (X_i, Y_i)_{i \in \mathbb{Z}/l\mathbb{Z}} &\longmapsto (X_i Y_i - Y_{i-1} X_{i-1})_{i \in \mathbb{Z}/l\mathbb{Z}} \end{aligned}$$

the corresponding *moment map*. Finally, if $\theta = (\theta_i)_{i \in \mathbb{Z}/l\mathbb{Z}}$ is a family of complex numbers, we denote by $I_\theta(d)$ the family $(\theta_i \text{Id}_{\mathbb{C}^{d_i}})_{i \in \mathbb{Z}/l\mathbb{Z}}$ and by $\mathcal{O}_\theta(d)$ the closed subvariety of $\text{Mat}_{d_0}(\mathbb{C})$ consisting of matrices of rank ≤ 1 and trace $-\sum_{i \in \mathbb{Z}/l\mathbb{Z}} \theta_i d_i$. Finally, we set

$$\mathcal{Y}_\theta(d) = \mu_d^{-1}(I_\theta(d) + \mathcal{O}_\theta(d)) \quad \text{and} \quad \mathcal{X}_\theta(d) = \mathcal{Y}_\theta(d) // \mathbf{PGL}(d).$$

We will usually denote an element $(X_i, Y_i)_{i \in \mathbb{Z}/l\mathbb{Z}}$ of $\mathcal{Y}_\theta(d)$ by (X, Y) , where $X = (X_i)_{i \in \mathbb{Z}/l\mathbb{Z}}$ and $Y = (Y_i)_{i \in \mathbb{Z}/l\mathbb{Z}}$. Note that $\mathcal{Y}_\theta(d)$ is endowed with a \mathbb{C}^\times -action: if $\xi \in \mathbb{C}^\times$, we set

$$\xi \cdot (X, Y) = (\xi^{-1} X, \xi Y).$$

This action commutes with the action of $\mathbf{GL}(d)$ and the moment map is constant on \mathbb{C}^\times -orbits, so it induces a \mathbb{C}^\times -action on $\mathcal{X}_\theta(d)$.

Remark 2.1. — We extend the definition of $\mathcal{X}_\theta(d)$ to the case where $d \in \mathbb{Z}^{\mathbb{Z}/l\mathbb{Z}}$ by the convention that $\mathcal{X}_\theta(d) = \emptyset$ whenever at least one of the d_i 's is negative. ■

2.B. Action of the affine Weyl group. — If $l \geq 2$, let W_l^{aff} denote the affine Weyl group of type \tilde{A}_{l-1} . It is the Coxeter group with associated Coxeter system $(W_l^{\text{aff}}, S_l^{\text{aff}})$, where $S_l^{\text{aff}} = \{s_i \mid i \in \mathbb{Z}/l\mathbb{Z}\}$ and the Coxeter graph is given by




Fig. 2. Coxeter graph of $(W_l^{\text{aff}}, S_l^{\text{aff}})$

We extend this notation to the case where $l = 1$ by setting $W_1^{\text{aff}} = 1$.

Consider the Lie algebra $\mathfrak{g}_l = \mathfrak{sl}_l(\mathbb{C})$ and its affine version $\widehat{\mathfrak{g}}_l = \widehat{\mathfrak{sl}}_l(\mathbb{C}) = \mathfrak{sl}_l(\mathbb{C})[t, t^{-1}] \oplus \mathbb{C}\mathbf{1} \oplus \mathbb{C}\partial$. Let $\mathfrak{h} \subset \mathfrak{g}_l$ be the Cartan subalgebra formed by the diagonal matrices and set $\widehat{\mathfrak{h}} = \mathfrak{h} \oplus \mathbb{C}\mathbf{1} \oplus \mathbb{C}\partial$.

The \mathbb{C} -vector space $\widehat{\mathfrak{h}}^*$ has a basis $(\alpha_0, \alpha_1, \dots, \alpha_{l-1}, \Lambda_0)$, where $\alpha_0, \alpha_1, \dots, \alpha_{l-1}$ are the simple roots of $\widehat{\mathfrak{g}}_l$ and Λ_0 is such that Λ_0 annihilates \mathfrak{h} and ∂ and $\Lambda_0(\mathbf{1}) = 1$. Denote by R_l^{aff} and R_l the affine and the non-affine root lattices respectively (i.e., R_l^{aff} is the \mathbb{Z} -lattice generated by $\alpha_0, \alpha_1, \dots, \alpha_{l-1}$ and R_l is the sublattice generated by $\alpha_1, \dots, \alpha_{l-1}$.)

Following [19], we define two actions of W_l^{aff} : a non-linear one on $\mathbb{Z}^{\mathbb{Z}/l\mathbb{Z}}$, and a linear one on $\mathbb{C}^{\mathbb{Z}/l\mathbb{Z}}$. If $l = 1$, there is nothing to define so we may assume that $l \geq 2$. If $d = (d_i)_{i \in \mathbb{Z}/l\mathbb{Z}} \in \mathbb{Z}^{\mathbb{Z}/l\mathbb{Z}}$ and if $j \in \mathbb{Z}/l\mathbb{Z}$, we set $s_j(d) = (d'_i)_{i \in \mathbb{Z}/l\mathbb{Z}}$, where

$$d'_i = \begin{cases} d_i & \text{if } i \neq j, \\ \delta_{j0} + d_{i+1} + d_{i-1} - d_i & \text{if } i = j. \end{cases}$$

Remark 2.2. — We can identify $\mathbb{Z}^{\mathbb{Z}/l\mathbb{Z}}$ with the root lattice R_l^{aff} by

$$d \mapsto \sum_{i \in \mathbb{Z}/l\mathbb{Z}} d_i \alpha_i.$$

Beware, the action considered here is not the usual action of W_l^{aff} on the root lattice. When we have $w(d) = d'$ with respect to the action defined above, this corresponds to $w(\Lambda_0 - d) = \Lambda_0 - d'$ for the usual action of W_l^{aff} on $\widehat{\mathfrak{h}}^*$. ■

If $\theta = (\theta_i)_{i \in \mathbb{Z}/l\mathbb{Z}} \in \mathbb{C}^{\mathbb{Z}/l\mathbb{Z}}$, we set $s_j(\theta) = (\theta'_i)_{i \in \mathbb{Z}/l\mathbb{Z}}$, where

$$\theta'_i = \begin{cases} \theta_i & \text{if } i \notin \{j-1, j, j+1\}, \\ \theta_j + \theta_i & \text{if } i \in \{j-1, j+1\}, \\ -\theta_i & \text{if } i = j. \end{cases}$$

It is readily seen that these definitions on generators extend to an action of the whole group W_l^{aff} . We also define a pairing $\mathbb{Z}^{\mathbb{Z}/l\mathbb{Z}} \times \mathbb{C}^{\mathbb{Z}/l\mathbb{Z}} \rightarrow \mathbb{C}$, $(d, \theta) \mapsto d \cdot \theta$, where

$$d \cdot \theta = \sum_{i \in \mathbb{Z}/l\mathbb{Z}} d_i \theta_i.$$

Then

$$(2.3) \quad s_j(d) \cdot s_j(\theta) = (d \cdot \theta) - \delta_{j0} \theta_0.$$

It is proved in [19, Corollary 3.6] that

$$(2.4) \quad \mathcal{X}_{s_j(\theta)}(s_j(d)) \simeq \mathcal{X}_\theta(d) \quad \text{if } \theta_j \neq 0.$$

Note that this isomorphism takes into account the convention of Remark 2.1.

The isomorphism above motivates to consider the following equivalence relation on the set $\mathbb{Z}^{\mathbb{Z}/l\mathbb{Z}} \times \mathbb{C}^{\mathbb{Z}/l\mathbb{Z}}$. Let \sim be the transitive closure of

$$(d, \theta) \sim (s_i(d), s_i(\theta)), \quad \theta_i \neq 0.$$

The isomorphism (2.4) implies that if $(d, \theta) \sim (d', \theta')$, then we have an isomorphism of algebraic varieties $\mathcal{X}_\theta(d) \simeq \mathcal{X}_{\theta'}(d')$.

Recall that the affine Weyl group has another presentation. We have $W_l^{\text{aff}} = W_l \ltimes R_l$, where $W_l = \mathfrak{S}_l$ is the non-affine Weyl group, that is, the subgroup generated by s_1, \dots, s_{l-1} . For each $\alpha \in R_l$, denote by t_α the image of α in W_l^{aff} . Each element of W_l^{aff} can be written in a unique way in the form $w \cdot t_\alpha$, where $w \in W_l$ and $\alpha \in R_l$. Let δ_l denote the constant family $\delta_l = (1)_{i \in \mathbb{Z}/l\mathbb{Z}} \in \mathbb{Z}^{\mathbb{Z}/l\mathbb{Z}}$.

Recall also from [10, §11] the following result, which follows from the fact that $\mathcal{X}_\theta(n\delta_l)$ is isomorphic to some Calogero-Moser space (we will use this fact later, and make this statement more precise):

Lemma 2.5. — *If $n \geq 0$, then $\mathcal{X}_\theta(n\delta_l)$ is normal and irreducible of dimension $2n$.*

Definition 2.6. — *We say that the pair (d, θ) is smooth if it is equivalent to a pair of the form $(n\delta_l, \theta')$ such that $n \geq 0$ and $\mathcal{X}_{\theta'}(n\delta_l)$ is smooth. In particular, in this case, the variety $\mathcal{X}_\theta(d)$ is smooth and non-empty.*

In the following lemma we identify $\mathbb{Z}^{\mathbb{Z}/l\mathbb{Z}}$ with R_l^{aff} .

Lemma 2.7. — *Assume $\alpha \in R_l$ and $d \in \mathbb{Z}^{\mathbb{Z}/l\mathbb{Z}}$. Then we have $t_\alpha(d) \equiv d - \alpha \pmod{\mathbb{Z}\delta_l}$.*

Proof. — This statement is a partial case of [15, (6.5.2)] (see also Remark 2.2). □

Lemma 2.8. — *For each $d \in \mathbb{Z}^{\mathbb{Z}/l\mathbb{Z}}$, there exists a unique $n \in \mathbb{Z}$ such that d and $n\delta_l$ are in the same W_l^{aff} -orbit.*

Proof. — Let us prove the existence of n . Set $\alpha = d - d_0\delta_l$. Then α is clearly an element of R_l . Then by Lemma 2.7, the element $t_\alpha(d)$ is of the form $n\delta_l$.

Let us prove the uniqueness of n . Assume that $n_1\delta_l$ and $n_2\delta_l$ are in the same W_l^{aff} -orbit. Since $n_1\delta_l$ is W_l -stable, the W_l^{aff} -orbit of $n_1\delta_l$ coincides with the R_l -orbit of $n_1\delta_l$. So there exists $\alpha \in R_l$ such that $t_\alpha(n_1\delta_l) = n_2\delta_l$. By Lemma 2.7, this forces $\alpha = 0$ and $n_1 = n_2$. \square

Consider the \mathbb{Z} -linear map

$$R_l^{\text{aff}} \rightarrow \mathbb{C}^{\mathbb{Z}/l\mathbb{Z}}, \quad d \mapsto \bar{d},$$

given by

$$(\bar{\alpha}_r)_i = 2\delta_{i,r} - \delta_{i,r+1} - \delta_{i,r-1}.$$

The kernel of this map is $\mathbb{Z}\delta_l$. Set

$$\Sigma(\theta) = \sum_{i \in \mathbb{Z}/l\mathbb{Z}} \theta_i.$$

Lemma 2.9. — *For each $\alpha \in R_l$ and $\theta \in \mathbb{C}^{\mathbb{Z}/l\mathbb{Z}}$, we have $t_\alpha(\theta) = \theta + \Sigma(\theta)\bar{\alpha}$.*

Proof. — The W_l^{aff} -action on $\mathbb{C}^{\mathbb{Z}/l\mathbb{Z}}$ defined above coincides with the (usual) action of W_l^{aff} on the dual of the span of $\alpha_0, \alpha_1, \dots, \alpha_{l-1}$ in $\hat{\mathfrak{h}}^*$. The statement follows from [15, (6.5.2)]. \square

We denote by $(\mathbb{Z}^{\mathbb{Z}/l\mathbb{Z}})_+$ the set of $d \in \mathbb{Z}^{\mathbb{Z}/l\mathbb{Z}}$ which belong to the orbit of some $n\delta_l$, with $n \geq 0$. This set has a more precise combinatorial description. For example, it will follow from Proposition 4.4 that an element $d \in \mathbb{Z}^{\mathbb{Z}/l\mathbb{Z}}$ is in $(\mathbb{Z}^{\mathbb{Z}/l\mathbb{Z}})_+$ if and only if d is a residue of some Young diagram. In particular, the set $(\mathbb{Z}^{\mathbb{Z}/l\mathbb{Z}})_+$ is contained in $\mathbb{N}^{\mathbb{Z}/l\mathbb{Z}}$.

Lemma 2.10. — *Assume that $d \in \mathbb{Z}^{\mathbb{Z}/l\mathbb{Z}}$ is such that there exists a simple representation in $\mathcal{Y}_\theta(d)$. Then*

- (a) *the variety $\mathcal{X}_\theta(d)$ is normal and irreducible,*
- (b) *we have $d \in (\mathbb{Z}^{\mathbb{Z}/l\mathbb{Z}})_+$.*

Proof. — Let $n \in \mathbb{Z}$ and $w \in W_l^{\text{aff}}$ be such that $n\delta_l = w(d)$. Let us show that (2.4) implies an isomorphism $\mathcal{X}_\theta(d) \simeq \mathcal{X}_{w(\theta)}(n\delta_l)$. (In particular, we must have $n \geq 0$.) The only difficulty that we have is caused by the condition $\theta_i \neq 0$ in (2.4). But [9, Lemma 7.2] implies that we can find a sequence of reflections such that $s_{i_1} s_{i_2} \dots s_{i_r}(d) = n\delta_l$ and such that we can apply a sequence of isomorphisms (2.4) with no danger to get $\theta_{i_j} = 0$ while applying s_{i_j} . \square

2.C. Smoothness. — The following result is proved in [13, Lemma 4.3 and its proof] (which is based on [17, Theorem 1.2] and [9, Theorem 1.2]):

Theorem 2.11. — *Let $n \geq 0$. Then the following are equivalent:*

- (1) *The variety $\mathcal{X}_\theta(n\delta_l)$ is smooth.*
- (2) *Every element of $\mathcal{Y}_\theta(n\delta_l)$ defines a simple representation of the quiver \overline{Q}_l .*
- (3) *The family θ satisfies*

$$\Sigma(\theta) \prod_{\substack{1 \leq i \leq j \leq l-1 \\ -(n-1) \leq k \leq n-1}} ((\theta_i + \theta_{i+1} + \cdots + \theta_j) + k\Sigma(\theta)) \neq 0.$$

If these equivalent conditions hold, then $\mathbf{PGL}(n\delta_l)$ acts freely on $\mathcal{Y}_\theta(n\delta_l)$, and so, as a set, $\mathcal{X}_\theta(n\delta_l)$ identifies with $\mathbf{GL}(n\delta_l)$ -orbits in $\mathcal{Y}_\theta(n\delta_l)$.

2.D. Fixed points. — For every $m \geq 1$, the group of m -th roots of unity μ_m acts on $\mathcal{X}_\theta(n\delta_l)$ (this is the restriction of the \mathbb{C}^\times -action defined in §2.A). We aim to compute the fixed point variety $\mathcal{X}_\theta(n\delta_l)^{\mu_m}$ whenever $\mathcal{X}_\theta(n\delta_l)$ is smooth. Note first that μ_l acts trivially on $\mathcal{X}_\theta(n\delta_l)$: indeed, if $\zeta \in \mu_l$ and if $(X, Y) \in \mathcal{Y}_\theta(n\delta_l)$, then $\zeta \cdot (X, Y) = g_0(X, Y)$, where $g_0 = (\zeta^i \text{Id}_{\mathbb{C}^n})_{i \in \mathbb{Z}/l\mathbb{Z}} \in \mathbf{GL}(n\delta_l)$. This means that, in order to describe $\mathcal{X}_\theta(n\delta_l)^{\mu_m}$, we may replace μ_m by $\langle \mu_m, \mu_l \rangle$ or, in other words, we may assume that l divides m .

Hypothesis. *We fix in this subsection a non-zero natural number k and we set $m = kl$.*

If $\theta = (\theta_i)_{i \in \mathbb{Z}/l\mathbb{Z}} \in \mathbb{C}^{\mathbb{Z}/l\mathbb{Z}}$, we define $\theta[k]$ to be the element $(\theta[k]_j)_{j \in \mathbb{Z}/m\mathbb{Z}}$ such that

$$\theta[k]_j = \theta_i$$

if $j \equiv i \pmod{l}$. Roughly speaking, $\theta[k]$ is the concatenation of k copies of θ .

We denote by $\mathbb{Z}^{\mathbb{Z}/m\mathbb{Z}}[n\delta_l]$ the set of $d = (d_j)_{j \in \mathbb{Z}/m\mathbb{Z}} \in \mathbb{Z}^{\mathbb{Z}/m\mathbb{Z}}$ such that

$$n = \sum_{\substack{j \in \mathbb{Z}/m\mathbb{Z} \\ j \equiv i \pmod{l}}} d_j$$

for all $i \in \mathbb{Z}/l\mathbb{Z}$. Set also $\mathcal{E}(k, l, n) = \mathbb{Z}^{\mathbb{Z}/m\mathbb{Z}}[n\delta_l] \cap (\mathbb{Z}^{\mathbb{Z}/m\mathbb{Z}})_+$. If $d \in \mathcal{E}(k, l, n)$ and $(X, Y) \in \text{Rep}(\overline{Q}_m, d)$, we set

$$i_k^{(d)}(X, Y) = (X', Y'),$$

where $X'_i = \bigoplus_{\substack{j \in \mathbb{Z}/m\mathbb{Z} \\ j \equiv i \pmod{l}}} X_j$ and $Y'_i = \bigoplus_{\substack{j \in \mathbb{Z}/m\mathbb{Z} \\ j \equiv i \pmod{l}}} Y_j$. By the definition of $\mathcal{E}(k, l, n)$, we have

$X'_i, Y'_i \in \text{Mat}_n(\mathbb{C})$. In other words, $(X', Y') \in \text{Rep}(\overline{Q}_l, n\delta_l)$: it is clear that $i_k^{(d)} : \text{Rep}(\overline{Q}_m, d) \hookrightarrow \text{Rep}(\overline{Q}_l, n\delta_l)$ is a closed immersion.

Lemma 2.12. — *If $d \in \mathcal{E}(k, l, n)$ and $(X, Y) \in \mathcal{Y}_{\theta[k]}(d)$, then $i_k^{(d)}(X, Y) \in \mathcal{Y}_\theta(n\delta_l)$.*

Proof. — Write $i_k^{(d)}(X, Y) = (X', Y')$ and let $i \in \mathbb{Z}/l\mathbb{Z}$. Write p_0 for the rank one matrix $X_0 Y_0 - Y_{-1} X_{-1} - \theta[k]_0 \text{Id}_{\mathbb{C}^{d_0}}$. Then

$$\begin{aligned} X'_i Y'_i - Y'_{i-1} X'_{i-1} &= \bigoplus_{\substack{j \in \mathbb{Z}/m\mathbb{Z} \\ j \equiv i \pmod{l}}} (X_j Y_j - Y_{j-1} X_{j-1}) \\ &= \bigoplus_{\substack{j \in \mathbb{Z}/m\mathbb{Z} \\ j \equiv i \pmod{l}}} \theta[k]_j \text{Id}_{\mathbb{C}^{d_j}} + \delta_{0i} p_0 \\ &= \theta_i \text{Id}_{\mathbb{C}^n} + \delta_{0i} p_0, \end{aligned}$$

as desired. \square

It is also clear that, if (X, Y) and (\tilde{X}, \tilde{Y}) are two elements of $\mathcal{Y}_{\theta[k]}(d)$ which are conjugate under $\mathbf{GL}(d)$, then $i_k^{(d)}(X, Y)$ and $i_k^{(d)}(\tilde{X}, \tilde{Y})$ are conjugate under $\mathbf{GL}(n\delta_l)$. This means that $i_k^{(d)}$ induces a morphism of algebraic varieties still denoted by

$$i_k^{(d)} : \mathcal{X}_{\theta[k]}(d) \longrightarrow \mathcal{X}_{\theta}(n\delta_l).$$

For $d \in \mathcal{E}(k, l, n)$, set

$$\mathcal{X}_d = i_k^{(d)}(\mathcal{X}_{\theta[k]}(d)) \subset \mathcal{X}_{\theta}(n\delta_l).$$

For the moment, we allow the possibility that the varieties $\mathcal{X}_{\theta[k]}(d)$ and \mathcal{X}_d are empty. Set

$$\mathcal{E}(k, l, n)_{\neq \emptyset} = \{d \in \mathcal{E}(k, l, n); \mathcal{X}_d \neq \emptyset\}.$$

We will show in Corollary 4.13 that we have $\mathcal{E}(k, l, n)_{\neq \emptyset} = \mathcal{E}(k, l, n)$.

The next result describes the fixed point variety $\mathcal{X}_{\theta}(n\delta_l)^{\mu_m}$: it is the main step towards the proof of Theorem 1.3.

Theorem 2.13. — *Assume that $\mathcal{X}_{\theta}(n\delta_l)$ is smooth. Then:*

- (a) $\mathcal{X}_{\theta}(n\delta_l)^{\mu_m}$ is smooth.
- (b) We have

$$\mathcal{X}_{\theta}(n\delta_l)^{\mu_m} = \coprod_{d \in \mathcal{E}(k, l, n)_{\neq \emptyset}} \mathcal{X}_d$$

and \mathcal{X}_d are exactly the irreducible components of $\mathcal{X}_{\theta}(n\delta_l)^{\mu_m}$.

- (c) If $d \in \mathcal{E}(k, l, n)_{\neq \emptyset}$, then $i_k^{(d)} : \mathcal{X}_{\theta[k]}(d) \rightarrow \mathcal{X}_d$ is an isomorphism.

Proof. — (a) follows from the fact already mentioned in the proof of Corollary 1.5 that the fixed point variety under the action of a finite group on a smooth variety is again smooth.

We will now prove (b). Let ξ be a primitive m -th root of unity and let $\zeta = \xi^k$ (it is a primitive l -th root of unity). Let g_0 denote the element $(\zeta^i \text{Id}_{\mathbb{C}^n})_{i \in \mathbb{Z}/l\mathbb{Z}}$ in $\mathbf{GL}(n\delta_l)$. This element satisfies

$$(*) \quad g_0(X, Y) = (\zeta^{-1} X, \zeta Y)$$

for all $(X, Y) \in \mathcal{Y}_{\theta}(n\delta_l)$. We will use the following variety:

$$\mathcal{Z} = \{(X, Y, g) \in \mathcal{Y}_{\theta}(n\delta_l) \times \mathbf{GL}(n\delta_l) \mid g(X, Y) = (\zeta^{-1} X, \zeta Y) \text{ and } g p_{X, Y} = p_{X, Y}\},$$

where $p_{X,Y} = ([X_i, Y_i] - \theta_i)_{i \in \mathbb{Z}/l\mathbb{Z}}$: it is 0 except on the 0-th component, where it is equal to a rank one matrix of trace $-n\Sigma(\theta)$ (recall from Theorem 2.11 that $\Sigma(\theta) \neq 0$ because $\mathcal{X}_\theta(n\delta_l)$ is assumed to be smooth).

We denote by $p' : \mathcal{Z} \rightarrow \mathcal{Y}_\theta(n\delta_l)$ the projection on the first two factors, by $p : \mathcal{Z} \rightarrow \mathcal{X}_\theta(n\delta_l)$ the map induced by p' and by $q : \mathcal{Z} \rightarrow \mathbf{GL}(n\delta_l)$ the projection on the third factor. Note that $\mathbf{PGL}(n\delta_l)$ acts on \mathcal{Z} by conjugacy on the three factors, and so the maps p' and q are equivariant for this action.

First, it is clear that $p(\mathcal{Z}) \subset \mathcal{X}_\theta(n\delta_l)^{\mu_m}$. Conversely, if $(X, Y) \in \mathcal{Y}_\theta(n\delta_l)$ is a representative of an element of $\mathcal{X}_\theta(n\delta_l)^{\mu_m}$, then there exists $g \in \mathbf{PGL}(n\delta_l)$ such that ${}^g(X, Y) = (\xi^{-1}X, \xi Y)$. In particular, ${}^g p_{X,Y} = p_{X,Y}$. So g stabilizes the image of $p_{X,Y}$, which is of dimension 1: this means that g acts by a non-zero scalar ω on $\text{Im}(p_{X,Y})$. Setting $g' = \omega^{-1}g$, we see that $(X, Y, g) \in \mathcal{Z}$. So we have proved the following fact:

Fact 1. $p(\mathcal{Z}) = \mathcal{X}_\theta(n\delta_l)^{\mu_m}$.

The next fact follows from the freeness of the action of $\mathbf{PGL}(n\delta_l)$ on $\mathcal{Y}_\theta(n\delta_l)$ (see Theorem 2.11).

Fact 2. *If (X, Y, g) and (X, Y, g') belong to \mathcal{Z} , then $g = g'$ and $g^k = g_0$.*

Indeed, the hypothesis implies that ${}^g(X, Y) = {}^{g'}(X, Y)$ and ${}^g(X, Y) = {}^{g^k}(X, Y)$. So there exists two non-zero complex numbers α and β such that $g' = \alpha g$ and $g^k = \beta g_0$. But the condition $g p_{X,Y} = g' p_{X,Y} = g_0 p_{X,Y}$ forces $\alpha = \beta = 1$ because $p_{X,Y} \neq 0$.

Now, let \mathcal{C} denote the set of elements $g \in \mathbf{GL}(n\delta_l)$ such that $g^k = g_0$. Since g_0 is central in $\mathbf{GL}(n\delta_l)$, \mathcal{C} is a disjoint union of finitely many semisimple conjugacy classes (which are the irreducible components of \mathcal{C}). We fix $g \in \mathcal{C}$ such that $q^{-1}(g) \neq \emptyset$ and we denote by \mathcal{C}_g its conjugacy class in $\mathbf{GL}(n\delta_l)$. We will describe $q^{-1}(g)$. For this, let $E = \bigoplus_{i \in \mathbb{Z}/l\mathbb{Z}} \mathbb{C}^n$ and let $E_j = E^{\xi^{-j}g}$ (for $j \in \mathbb{Z}/m\mathbb{Z}$). In other words, E_j is the ξ^j -eigenspace of g in E : it is contained in the i -th component of E , where i is the element of $\mathbb{Z}/l\mathbb{Z}$ such that $j \equiv i \pmod{l}$. Let $d_j = \dim_{\mathbb{C}}(E_j)$ and let $d = (d_j)_{j \in \mathbb{Z}/m\mathbb{Z}}$.

If $(X, Y, g) \in \mathcal{Z}$, then X sends E_j to E_{j-1} (we denote by $X_j : E_j \rightarrow E_{j-1}$ the induced map) while Y sends E_j to E_{j+1} (we denote by $Y_j : E_j \rightarrow E_{j+1}$ the induced map). By choosing a basis of every E_j , the family $(X_j, Y_j)_{j \in \mathbb{Z}/m\mathbb{Z}}$ defines a representation of the quiver \overline{Q}_m , of dimension vector d . This means that we have defined a map $q^{-1}(g) \rightarrow \text{Rep}(\overline{Q}_m, d)$, which is clearly a closed immersion, and whose image is $\mathcal{Y}_{\theta[k]}(d)$ (because $\text{Im}(p_{X,Y})$ is contained in E_0).

Note that by construction we have $d \in \mathbb{Z}^{\mathbb{Z}/m\mathbb{Z}}[n\delta_l]$. Moreover, by Lemma 2.10(b) and Theorem 2.11 we also have $d \in (\mathbb{Z}^{\mathbb{Z}/m\mathbb{Z}})_+$. This means that d is an element of $\mathcal{E}(k, l, n) = \mathbb{Z}^{\mathbb{Z}/m\mathbb{Z}}[n\delta_l] \cap (\mathbb{Z}^{\mathbb{Z}/m\mathbb{Z}})_+$. Moreover, it is clear that the set of elements d that can appear as above (from some $g \in \mathcal{C}$ such that $q^{-1}(g) \neq \emptyset$) is $\mathcal{E}(k, l, n)_{\neq \emptyset}$.

This shows the following fact.

Fact 3. *With the above notation, $q^{-1}(g) \simeq \mathcal{Y}_{\theta[k]}(d)$. In particular, $q^{-1}(g)$ is irreducible.*

Fact 3 and its proof imply the following.

Fact 4. *We have $p(q^{-1}(\mathcal{C}_g)) = \mathcal{X}_d$.*

Denote by π the projection $\pi : \mathcal{Y}_\theta(n\delta_l) \rightarrow \mathcal{X}_\theta(n\delta_l)$. Fact 2 implies that the map $p' : \mathcal{Z} \rightarrow \pi^{-1}(\mathcal{X}_\theta(n\delta_l)^{\mu_m})$ is bijective. Moreover, since $\pi^{-1}(\mathcal{X}_\theta(n\delta_l)^{\mu_m})$ is smooth (the group $\mathbf{PGL}(n\delta_l)$ acts freely on $\mathcal{Y}_\theta(n\delta_l)$), this bijection is an isomorphism of algebraic varieties. Let

$$(p')^{-1} : \pi^{-1}(\mathcal{X}_\theta(n\delta_l)^{\mu_m}) \rightarrow \mathcal{Z}$$

be its inverse.

Consider the morphism $\tilde{\tau}: \pi^{-1}(\mathcal{X}_\theta(n\delta_l)^{\mu_m}) \rightarrow \mathbf{GL}(n\delta_l)$ given by $\tilde{\tau} = q \circ (p')^{-1}$. Since we have $\pi^{-1}(\mathcal{X}_\theta(n\delta_l)^{\mu_m}) // \mathbf{GL}(n\delta_l) = \mathcal{X}_\theta(n\delta_l)^{\mu_m}$, we get a morphism $\tau: \mathcal{X}_\theta(n\delta_l)^{\mu_m} \rightarrow \mathbf{GL}(n\delta_l) // \mathbf{GL}(n\delta_l)$, where the group $\mathbf{GL}(n\delta_l)$ acts on itself by conjugation.

Moreover, since we clearly have $\tau^{-1}(\mathcal{C}_g) = p(q^{-1}(\mathcal{C}_g))$, Fact 4 implies the following.

Fact 5. *We have $\tau^{-1}(\mathcal{C}_g) = \mathcal{X}_d$. In particular \mathcal{X}_d is closed.*

This completes the proof of (b). Now we prove (c). The group $\mathbf{GL}(n\delta_l)$ acts transitively on \mathcal{C}_g . So we get an isomorphism of varieties

$$q^{-1}(g) // C_{\mathbf{GL}(n\delta_l)}(g) \xrightarrow{\sim} q^{-1}(\mathcal{C}_g) // \mathbf{GL}(n\delta_l).$$

Now, if $d = (\dim_{\mathbb{C}}(E^{\xi^{-j}g}))_{j \in \mathbb{Z}/m\mathbb{Z}}$, $q^{-1}(g)$ identifies with $\mathcal{Y}_{\theta[k]}(d)$ and $C_{\mathbf{GL}(n\delta_l)}(g)$ identifies with $\mathbf{GL}(d)$. Glueing this with Facts 3 and 4, we get an isomorphism of varieties

$$\mathcal{X}_{\theta[k]}(d) \xrightarrow{\sim} \mathcal{X}_d,$$

which is the map $i_k^{(d)}$. This proves (c). \square

Corollary 2.14. — *If $k > n$, then we have $\mathcal{X}_\theta(n\delta_l)^{\mu_m} = \mathcal{X}_\theta(n\delta_l)^{\mathbb{C}^\times}$.*

Proof. — Take $d \in \mathcal{E}(k, l, n)_{\neq \emptyset}$. Since $k > n$, some component of d must be zero. This implies that \mathbb{C}^\times acts trivially on $\mathcal{X}_{\theta[k]}(d)$ because the action of \mathbb{C}^\times is induced by elements of $\mathbf{GL}(d)$.

Since $i_k^{(d)}$ is \mathbb{C}^\times -equivariant and

$$\mathcal{X}_\theta(n\delta_l)^{\mu_m} = \coprod_{d \in \mathcal{E}(k, l, n)_{\neq \emptyset}} i_k^{(d)}(\mathcal{X}_{\theta[k]}(d))$$

(by Theorem 2.13(b)), the \mathbb{C}^\times -action on the variety $\mathcal{X}_\theta(n\delta_l)^{\mu_m}$ is also trivial. \square

3. Proof of Theorem 1.3

First, the explicit description of \mathcal{Z}_c whenever $n = 1$ (see Example 1.2) allows to prove easily Theorem 1.3 in this case. Also, let us decompose V as

$$V = V^W \oplus V_1 \oplus \cdots \oplus V_r,$$

where the V_i are the non-trivial irreducible components of V as a CW-module. Then W decomposes as $W = W_1 \times \cdots \times W_r$, where (V_i, W_i) is then a reflection subquotient (subgroup) of W . Let c_i denote the restriction of c to W_i . Then

$$\mathcal{Z}_c(V, W) \simeq V^W \times V^{*W} \times \mathcal{Z}_{c_1}(V_1, W_1) \times \cdots \times \mathcal{Z}_{c_r}(V_r, W_r)$$

so that the proof of Theorem 1.3 is easily reduced to the case where W acts irreducibly on V/V^W .

In this case, the smoothness of $\mathcal{Z}_c(V, W)$ implies that W is of type G_4 or $G(l, 1, n)$ for some $l \geq 1$. The case of type G_4 will be handled by computer calculations: see Section 5 for details. This means that we can work under the following hypothesis:

Hypothesis. From now on, and until the end of this section, we assume that $n \geq 2$, that $V = \mathbb{C}^n$ and that $W = G(l, 1, n)$. We also assume that σ is an m -th root of unity, where $m = kl$ for some $k \geq 1$.

Recall that $G(l, 1, n)$ is the group of monomial matrices with coefficients in μ_l (the group of l -th root of unity in \mathbb{C}^\times) and recall that $n \geq 2$.

3.A. Quiver varieties vs Calogero-Moser spaces. — We fix a primitive l -th root of unity ζ . We denote by s the permutation matrix corresponding to the transposition $(1, 2)$ and we set

$$t = \text{diag}(\zeta, 1, \dots, 1) \in W.$$

Then $s, t, t^2, \dots, t^{l-1}$ is a set of representatives of conjugacy classes of reflections of W . We set for simplification

$$a = c_s \quad \text{and} \quad k_j = \frac{1}{l} \sum_{i=1}^{l-1} \zeta^{-i(j-1)} c_{t^i}$$

for $j \in \mathbb{Z}/l\mathbb{Z}$. Then

$$(3.1) \quad k_0 + \dots + k_{l-1} = 0 \quad \text{and} \quad c_{t^i} = \sum_{j \in \mathbb{Z}/l\mathbb{Z}} \zeta^{i(j-1)} k_j$$

for $1 \leq i \leq l-1$. Finally, if $i \in \mathbb{Z}/l\mathbb{Z}$, we set

$$(3.2) \quad \theta_i = \begin{cases} k_{-i} - k_{1-i} & \text{if } i \neq 0, \\ -a + k_0 - k_1 & \text{if } i = 0. \end{cases}$$

and $\theta = (\theta_i)_{i \in \mathbb{Z}/l\mathbb{Z}}$.

The following result is proved in [13, Theorem 3.10]. (Note that our k_i is related with Gordon's H_i via $H_i = k_{-i} - k_{1-i}$.)

Theorem 3.3. — *With the above notation, there is a \mathbb{C}^\times -equivariant isomorphism of varieties*

$$\mathcal{X}_c \xrightarrow{\sim} \mathcal{X}_\theta(n\delta_l).$$

The theorem above is also true for $n = 1$. But in this case, the variety \mathcal{X}_c has no parameter a . On the other hand, the variety $\mathcal{X}_\theta(\delta_l)$ is independent of θ_0 .

Putting Theorems 2.11 and 3.3 together, one gets:

Corollary 3.4. — *The variety \mathcal{X}_c is smooth if and only if*

$$a \prod_{\substack{0 \leq i \neq j \leq l-1 \\ 0 \leq r \leq n-1}} (k_i - k_j - ra) \neq 0.$$

Proof of Theorem 1.3. — Assume now that \mathcal{Z}_c is smooth and let \mathcal{X} be an irreducible component of $\mathcal{Z}_c^{\mu_m}$. Using the isomorphism of Theorem 3.3, we see that

$$\mathcal{Z}_c^{\mu_m} \xrightarrow{\sim} \mathcal{X}_{\theta}(n\delta_l)^{\mu_m}.$$

So, by Theorem 2.13, there exists $d \in \mathcal{E}(k, l, n)_{\neq \emptyset}$ such that $\mathcal{X} \simeq \mathcal{X}_{\theta[k]}(d)$. By Lemma 2.8, there exists $r \geq 0$ and $w \in W_m^{\text{aff}}$ such that $w(d) = r\delta_m$. So it follows from the isomorphism (2.4) that

$$\mathcal{X} \simeq \mathcal{X}_{w(\theta[k])}(r\delta_m)$$

(see also Lemma 2.10 and its proof). Using Theorem 3.3 in the other way, we see that there exists a complex valued function c' on the set of conjugacy classes of reflections of $W' = G(m, 1, r)$ such that $\mathcal{X}_{w(\theta[k])}(r\delta_m) \simeq \mathcal{Z}_{c'}(W')$. Moreover, as the action of W_m^{aff} on $\mathbb{C}^{\mathbb{Z}/m\mathbb{Z}}$ is linear, this implies that the map $c \mapsto c'$ is linear.

This proves almost every statement of Theorem 1.3, except that W' can be realized as a reflection subquotient of (V, W) . For this, we need the following fact.

Fact. *We have $kr \leq n$.*

Indeed, the element $d - r\delta_m$ is in the W_m^{aff} -orbit of 0. Then Proposition 4.4 implies that $d - r\delta_m$ is a residue of some m -core. In particular, each coordinates of $d - r\delta_m$ must be positive. Since the sum of the coordinates of d is nl , this gives $nl - rkl \geq 0$. This implies $kr \leq n$.

Now, using the fact proved above, let x denotes the matrix

$$x = \text{diag}(\underbrace{\zeta, 1, \dots, 1}_{k \text{ terms}}) \cdot M_{(1,2,\dots,k)} \in \mathbf{GL}_k(\mathbb{C}),$$

where $M_{(1,2,\dots,k)}$ is the permutation matrix associated with the cycle $(1, 2, \dots, k)$. Now, let g be the matrix

$$g = \text{diag}(\underbrace{x, \dots, x}_{r \text{ terms}}, \text{Id}_{\mathbb{C}^{n-rk}}) \in G(l, 1, n).$$

Let V' denote the ξ -eigenspace of g , where ξ is a primitive m -th root of unity such that $\xi^k = \zeta$. Then $C_W(g)$ acts on V' , V' is of dimension r and, if we denote by K the kernel of this action, then $C_W(g)/K \simeq G(m, 1, r)$. So $(V', G(m, 1, r))$ is a reflection subquotient of (V, W) . \square

Remark 3.5. — The formula (3.2) above yields a bijection between $\mathbb{C}^{\mathbb{Z}/l\mathbb{Z}}$ and the set of $(l+1)$ -tuples of complex numbers (a, k_0, \dots, k_{l-1}) such that $k_0 + k_1 + \dots + k_{l-1} = 0$. Moreover, the W_l^{aff} -action on $\mathbb{C}^{\mathbb{Z}/l\mathbb{Z}}$ (see Section 2.B) translates to the action on the $(l+1)$ -tuples in the following way

$$s_r(a, k_0, \dots, k_{l-r}, k_{l-r+1}, \dots, k_{l-1}) = (a, k_0, \dots, k_{l-r+1}, k_{l-r}, \dots, k_{l-1}), \quad \text{for } r = 1, 2, \dots, l-1$$

and

$$s_0(a, k_0, k_1, k_2, \dots, k_{l-1}) = (a, k_1 + a, k_0 - a, k_2, \dots, k_{l-1}).$$

In particular, we see that the subgroup W_l of W_l^{aff} acts by permutation of the parameters k_0, k_1, \dots, k_{l-1} . \blacksquare

The next result follows from [2], but we provide here a different proof (which works only in type $G(d, 1, n)$).

Corollary 3.6. — *A permutation of parameters k_0, k_1, \dots, k_{l-1} does not change the Calogero-Moser space \mathcal{X}_c (up to an isomorphism of algebraic varieties).*


Proof. — The subgroup W_l of W_l^{aff} stabilizes δ_l . Then by (2.4), for each $w \in W_l$, $n \in \mathbb{Z}_{>0}$ and $\theta \in \mathbb{C}^{\mathbb{Z}/l\mathbb{Z}}$, we have $\mathcal{X}_\theta(n\delta_l) \simeq \mathcal{X}_{w(\theta)}(n\delta_l)$. This proves the statement by Theorem 3.3 and Remark 3.5 \square

4. Combinatorics

In this subsection, we aim to make the statement of Theorem 1.3 more precise in the case where $W = G(l, 1, n)$: we wish to describe precisely the map $c \mapsto c'$ as well as the map $i_{\mathcal{X}}^* : \text{Irr}(W') \leftrightarrow \text{Irr}(W)$ in terms of the combinatorics of partitions, cores, l -quotients, etc.

4.A. Partitions and cores. — Let l and n be positive integers. A *partition* is a tuple $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_r)$ of positive integers (with no fixed length) such that $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_r$. Set $|\lambda| = \sum_{i=1}^r \lambda_i$. If $|\lambda| = n$, we say that λ is a partition of n .

Denote by \mathcal{P} (resp. $\mathcal{P}[n]$) be the set of all partitions (resp. the set of all partitions of n). By convention, $\mathcal{P}[0]$ contains one (empty) partition. We will identify partitions with Young diagrams. The partition λ corresponds to a Young diagram with r lines such that the i th line contains λ_i boxes. For example the partition $(4, 2, 1)$ corresponds to the Young diagram




Let b be a box of a Young diagram in the line r and column s . The l -*residue* of the box b is the number $s - r$ modulo l . (We also say that the integer $s - r$ is the ∞ -residue of the box b .) Then we obtain a map

$$\text{Res}_l : \mathcal{P} \rightarrow \mathbb{Z}^{\mathbb{Z}/l\mathbb{Z}}, \quad \lambda \mapsto \text{Res}_l(\lambda),$$

such that for each $i \in \mathbb{Z}/l\mathbb{Z}$ the number of boxes with residue i in λ is $(\text{Res}_l(\lambda))_i$. (Similarly, we obtain a map $\text{Res}_\infty : \mathcal{P} \rightarrow \mathbb{Z}^{\mathbb{Z}}$.)

Example 4.1. — For the partition $\lambda = (4, 2, 1)$ and $l = 3$ the residues of the boxes are



In this case we have $\text{Res}_l(\lambda) = (3, 2, 2)$ because there are three boxes with residue 0, two boxes with residue 1 and two boxes with residue 2. \blacksquare

We say that a box of a Young diagram is *removable* if it has no box on the right and on the bottom. For $\lambda, \mu \in \mathcal{P}$, we write $\mu \leq \lambda$ if the Young diagram of μ can be obtained from the Young diagram of λ by removing a sequence of removable boxes.

Definition 4.2. — We say that the partition λ is an l -core if there is no partition $\mu \leq \lambda$ such that the Young diagram of μ differs from the Young diagram of λ by l boxes with l different l -residues.

See [3] for more details about the combinatorics of l -cores. Let $\mathcal{C}_l \subset \mathcal{P}$ be the set of l -cores. Set $\mathcal{C}_l[n] = \mathcal{P}[n] \cap \mathcal{C}_l$.

If a partition λ is not an l -core, then we can get a smaller Young diagram from its Young diagram by removing l boxes with different l -residues. We can repeat this operation again and again until we get an l -core. It is well-known that the l -core that we get is independent of the choice of the boxes that we remove. Then we get an application

$$\text{Core}_l: \mathcal{P} \rightarrow \mathcal{C}_l.$$

If $\mu = \text{Core}_l(\lambda)$, we will say that the partition μ is the l -core of the partition λ .

Example 4.3. — The partition $(4, 2, 1)$ from the previous example is not a 3-core because it is possible to remove three bottom boxes. We get

$$\boxed{0} \boxed{1} \boxed{2} \boxed{0}$$

But this is still not a 3-core because we can remove three more boxes and we get

$$\boxed{0}$$

This shows that the partition (1) is the 3-core of the partition $(4, 2, 1)$. ■

As mentioned in Lemma 2.8, for each $d \in \mathbb{Z}^{\mathbb{Z}/l\mathbb{Z}}$, the W_l^{aff} -orbit of d contains an element of the form $n\delta_l$ for a unique $n \in \mathbb{Z}$.

Proposition 4.4. — Let d be an element of $\mathbb{Z}^{\mathbb{Z}/l\mathbb{Z}}$. The following statements are equivalent.

- (a) d is of the form $d = \text{Res}_l(\lambda)$ for some l -core λ ,
- (b) d is in the W_l^{aff} -orbit of 0 .

Proof. — Consider the W_l^{aff} -action on \mathcal{C}_l as in [3, Sec. 3]. By construction, the map $\text{Res}_l: \mathcal{C}_l \rightarrow \mathbb{Z}^{\mathbb{Z}/l\mathbb{Z}}$ is W_l^{aff} -invariant. Moreover, the residue of the empty partition is zero. The stabilizer of the empty partition in W_l^{aff} is W_l and the stabilizer of $0 \in \mathbb{Z}^{\mathbb{Z}/l\mathbb{Z}}$ in W_l^{aff} is also W_l . This shows that the map Res_l yields a bijection between the set of l -cores and the W_l^{aff} -orbit of 0 in $\mathbb{Z}^{\mathbb{Z}/l\mathbb{Z}}$. □

Denote by π the obvious surjection $\pi: \mathbb{Z}^{\mathbb{Z}/l\mathbb{Z}} \rightarrow \mathbb{Z}^{\mathbb{Z}/l\mathbb{Z}} / \mathbb{Z}\delta_l$.

Proposition 4.5. — The chain of maps

$$\mathcal{C}_l \xrightarrow{\text{Res}_l} \mathbb{Z}^{\mathbb{Z}/l\mathbb{Z}} \xrightarrow{\pi} \mathbb{Z}^{\mathbb{Z}/l\mathbb{Z}} / \mathbb{Z}\delta_l$$

yields a bijection between \mathcal{C}_l and $\mathbb{Z}^{\mathbb{Z}/l\mathbb{Z}} / \mathbb{Z}\delta_l$.

Proof. — We have seen in the proof of Proposition 4.4 that Res_l yields a bijection between \mathcal{C}_l and the W_l^{aff} -orbit of $0 \in \mathbb{Z}^{\mathbb{Z}/l\mathbb{Z}}$. This proves the statement because the restriction of π to the W_l^{aff} -orbit of $0 \in \mathbb{Z}^{\mathbb{Z}/l\mathbb{Z}}$ is bijective by Lemma 2.8. □

For $d = (d_i)_{i \in \mathbb{Z}/l\mathbb{Z}} \in \mathbb{Z}^{\mathbb{Z}/l\mathbb{Z}}$ we set $|d| = \sum_{i \in \mathbb{Z}/l\mathbb{Z}} d_i$. Set $\mathbb{Z}^{\mathbb{Z}/l\mathbb{Z}}[n] = \{d \in \mathbb{Z}^{\mathbb{Z}/l\mathbb{Z}}; |d| = n\}$. Denote by $\mathcal{C}_l[\equiv n]$ the subset of \mathcal{C}_l that contains only l -cores with the number of boxes congruent to n modulo l . Proposition 4.5 yields the following bijection $\varepsilon: \mathbb{Z}^{\mathbb{Z}/l\mathbb{Z}}[n] \xrightarrow{\sim} \mathcal{C}_l[\equiv n]$.

Lemma 4.6. — Assume $d \in \mathbb{Z}^{\mathbb{Z}/l\mathbb{Z}}[n]$. Then d is in $(\mathbb{Z}^{\mathbb{Z}/l\mathbb{Z}})_+$ if and only if $|\varepsilon(d)| \leq n$.

Proof. — By construction, $\varepsilon(d)$ is the unique l -core $\lambda \in \mathcal{C}_l[\equiv n]$ such that $\text{Res}_l(\lambda)$ is congruent to d modulo δ_l . Write $d = \text{Res}_l(\lambda) + r\delta_l$. Since by Proposition 4.4, the element $\text{Res}_l(\lambda)$ is in the W_l^{aff} -orbit of 0, then d is in the W_l^{aff} -orbit of $r\delta_l$. Then, by definition, the element d is in $(\mathbb{Z}^{\mathbb{Z}/l\mathbb{Z}})_+$ if and only if $r \geq 0$. Now, let us count the sum of the coordinates in the sides of the equality $d = \text{Res}_l(\lambda) + r\delta_l$. We obtain $n = |\lambda| + rl$. This shows that we have $r \geq 0$ if and only if $|\lambda| \leq n$. \square

Let ν be an l -core. Denote by $\mathcal{P}_{l,\nu}$ the set of partitions with the l -core ν . In particular we denote by $\mathcal{P}_{l,\emptyset}$ the set of partitions with trivial l -core. Let \mathcal{P}^l be the set of l -partitions (i.e., the set of l -tuples of partitions). If λ is a partition, we denote by $\beta_l(\lambda)$ its l -quotient (see for example [18, Sec. 2.2]).

Lemma 4.7. — If ν is an l -core, then β_l restricts to a bijection $\beta_{l,\nu}: \mathcal{P}_{l,\nu} \xrightarrow{\sim} \mathcal{P}^l$.

4.B. Description of $\mathcal{E}(k, l, n)$ in terms of cores. — Let k be a positive integer. Set $m = kl$. Set also $(\mathcal{C}_m)_{l,\nu} = \mathcal{P}_{l,\nu} \cap \mathcal{C}_m$.

Recall that we have a bijection $\varepsilon: \mathbb{Z}^{\mathbb{Z}/m\mathbb{Z}}[nl] \xrightarrow{\sim} \mathcal{C}_m[\equiv nl]$. Assume $d \in \mathbb{Z}^{\mathbb{Z}/m\mathbb{Z}}[nl]$.

Lemma 4.8. — We have $d \in \mathbb{Z}^{\mathbb{Z}/m\mathbb{Z}}[n\delta_l]$ if and only if the l -core of $\varepsilon(d)$ is trivial, where the set $\mathbb{Z}^{\mathbb{Z}/m\mathbb{Z}}[n\delta_l]$ is as in Section 2.D.

Proof. — By definition, $\varepsilon(d)$ is the unique m -core $\lambda \in \mathcal{C}_m[\equiv nl]$ such that $\text{Res}_m(\lambda)$ is equal to d modulo δ_m . On the other hand, by Proposition 4.4, the partition λ is an l -core if and only if $\text{Res}_l(\lambda)$ is a multiple of δ_l . By commutativity of the following diagram,

$$\begin{array}{ccc} & \mathcal{P} & \\ \text{Res}_m \swarrow & & \searrow \text{Res}_l \\ \mathbb{Z}^{\mathbb{Z}/m\mathbb{Z}} & \xrightarrow{\quad} & \mathbb{Z}^{\mathbb{Z}/l\mathbb{Z}} \end{array}$$

this condition is clearly equivalent to $d \in \mathbb{Z}^{\mathbb{Z}/m\mathbb{Z}}[n\delta_l]$. \square

Recall that the set $\mathcal{E}(k, l, n)$ was defined as the intersection $\mathcal{E}(k, l, n) = \mathbb{Z}^{\mathbb{Z}/m\mathbb{Z}}[n\delta_l] \cap (\mathbb{Z}^{\mathbb{Z}/m\mathbb{Z}})_+$. Combining Lemmas 4.6 and 4.8, we obtain the following.

Lemma 4.9. — The bijection $\varepsilon: \mathbb{Z}^{\mathbb{Z}/m\mathbb{Z}}[nl] \simeq \mathcal{C}_m[\equiv nl]$ restricts to a bijection

$$\varepsilon': \mathcal{E}(k, l, n) \rightarrow \mathcal{C}_m[\equiv nl] \cap \mathcal{C}_m[\leq nl] \cap (\mathcal{C}_m)_{l,\emptyset}.$$

For $\lambda = (\lambda^0, \dots, \lambda^{l-1}) \in \mathcal{P}^l$ we set $|\lambda| = |\lambda^0| + \dots + |\lambda^{l-1}|$. Set also $\mathcal{P}^l[n] = \{\lambda \in \mathcal{P}^l; |\lambda| = n\}$. We say that an element of $\mathcal{P}^l[n]$ is an l -partition of n .

Now recall the bijection $\beta_{l,v}: \mathcal{P}_{l,v} \xrightarrow{\sim} \mathcal{P}^l$ obtained by restriction from β_l as in Lemma 4.7. By construction, it has the following properties.

Lemma 4.10. — (a) For each $\lambda \in \mathcal{P}_{l,\emptyset}$, we have $\lambda \in \mathcal{C}_m$ if and only if $\beta_{l,\emptyset}(\lambda) \in (\mathcal{C}_k)^l$.
 (b) For each $\lambda \in \mathcal{P}_{l,v}$, we have $|\lambda| = l \cdot |\beta_{l,v}(\lambda)| + |v|$.

Part (a) of the lemma above shows that $\beta_{l,\emptyset}$ can be restricted to a bijection $(\mathcal{C}_m)_{l,\emptyset} \simeq (\mathcal{C}_k)^l$. Moreover, part (b) shows that new bijection restricts to a bijection

$$\mathcal{C}_m[\equiv nl] \cap \mathcal{C}_m[\leq nl] \cap (\mathcal{C}_m)_{l,\emptyset} \simeq (\mathcal{C}_k)^l[\equiv n, \leq n],$$

where

$$(\mathcal{C}_k)^l[\equiv n, \leq n] = \{\lambda \in (\mathcal{C}_k)^l; |\lambda| \leq n \text{ and } |\lambda| \equiv n \pmod{k}\}.$$

Combining this with Lemma 4.9, we get a bijection

$$\delta: \mathcal{E}(k, l, n) \rightarrow (\mathcal{C}_k)^l[\equiv n, \leq n].$$

4.C. Parametrization of the μ_m -fixed points. — We have proved in Theorem 2.13 that the subset $\mathcal{E}(k, l, n)_{\neq \emptyset}$ of $\mathcal{E}(k, l, n)$ parametrizes the irreducible components of $\mathcal{X}_\theta(n\delta_l)^{\mu_m}$. (But we will show in Corollary 4.13 that we have $\mathcal{E}(k, l, n)_{\neq \emptyset} = \mathcal{E}(k, l, n)$.) Then the bijection δ above gives another parametrization of the irreducible components of $\mathcal{X}_\theta(n\delta_l)^{\mu_m}$ in terms of $(\mathcal{C}_k)^l[\equiv n, \leq n]$. For each $\lambda \in (\mathcal{C}_k)^l[\equiv n, \leq n]$, we set $\mathcal{X}_\lambda = \mathcal{X}_{\delta^{-1}(\lambda)}$.

Recall the map $\text{Core}_k: \mathcal{P} \rightarrow \mathcal{C}_k$. It yields a map $\text{Core}_k: \mathcal{P}^l \rightarrow (\mathcal{C}_k)^l$ (component by component). Note that the set $(\mathcal{C}_k)^l[\equiv n, \leq n]$ considered above is nothing else but the image of $\mathcal{P}^l[n]$ by Core_k . In other words, the set $(\mathcal{C}_k)^l[\equiv n, \leq n]$ is the same thing as the set of k -cores of l -partitions of n .

This new parametrization has the following nice property.

Lemma 4.11. — Fix positive integers k_1 and k_2 such that k_1 divides k_2 . Set $m_1 = k_1 l$ and $m_2 = k_2 l$. Fix $\lambda_1 \in (\mathcal{C}_{k_1})^l[\equiv n, \leq n]$ and $\lambda_2 \in (\mathcal{C}_{k_2})^l[\equiv n, \leq n]$. Then we have $\mathcal{X}_{\lambda_2} \subset \mathcal{X}_{\lambda_1}$ if and only if $\text{Core}_{k_1}(\lambda_2) = \lambda_1$.

Proof. — Consider the map

$$p: \mathbb{Z}^{\mathbb{Z}/m_2\mathbb{Z}} \rightarrow \mathbb{Z}^{\mathbb{Z}/m_1\mathbb{Z}}, \quad p(d)_i = \sum_{j \equiv i \pmod{m_1}} d_j.$$

Let $d^1 \in \mathcal{E}(k_1, l, n)$ and $d^2 \in \mathcal{E}(k_2, l, n)$ be such that $\delta(d^1) = \lambda_1$ and $\delta(d^2) = \lambda_2$. We have $\mathcal{X}_{d^1} = \mathcal{X}_{\lambda_1}$ and $\mathcal{X}_{d^2} = \mathcal{X}_{\lambda_2}$. It is clear from the definition that we have $\mathcal{X}_{d^2} \subset \mathcal{X}_{d^1}$ if and

only if $p(d^2) = d^1$. Then the statement follows from the commutativity of the following diagram

$$\begin{array}{ccc}
\mathcal{E}(k_2, l, n) & \xrightarrow{p} & \mathcal{E}(k_1, l, n) \\
\downarrow & & \downarrow \\
\mathcal{C}_{m_2}[\equiv nl] \cap \mathcal{C}_{m_2}[\leq nl] \cap (\mathcal{C}_{m_2})_{l, \emptyset} & \xrightarrow{\text{Core}_{m_1}} & \mathcal{C}_{m_1}[\equiv nl] \cap \mathcal{C}_{m_1}[\leq nl] \cap (\mathcal{C}_{m_1})_{l, \emptyset} \\
\downarrow & & \downarrow \\
(\mathcal{C}_{k_2})^l[\equiv n, \leq n] & \xrightarrow{\text{Core}_{k_1}} & (\mathcal{C}_{k_1})^l[\equiv n, \leq n],
\end{array}$$

where the vertical maps are the bijections discussed above. \square

Proposition 4.12. — *The \mathbb{C}^\times -fixed points in $\mathcal{X}_\theta(n\delta_l)$ are parametrized by the set $\mathcal{P}^l[n]$. This parametrization $\mathcal{X}_\theta(n\delta_l)^{\mathbb{C}^\times} = \{p_\mu; \mu \in \mathcal{P}^l[n]\}$ can be chosen in such a way that for each $k \in \mathbb{Z}$, each $\lambda \in (\mathcal{C}_k)^l[\equiv n, \leq n]$ and each $\mu \in \mathcal{P}^l[n]$ we have $p_\mu \in \mathcal{X}_\lambda$ if and only if λ is the k -core of μ .*

Proof. — Assume $k > n$. Then by Corollary 2.14, we have $\mathcal{X}_\theta(n\delta_l)^{\mu_m} = \mathcal{X}_\theta(n\delta_l)^{\mathbb{C}^\times}$. Moreover, the set $\mathcal{X}_\theta(n\delta_l)^{\mathbb{C}^\times}$ is finite. We already know that $\mathcal{X}_\theta(n\delta_l)^{\mu_m}$ is in bijection with a subset of $(\mathcal{C}_k)^l[\equiv n, \leq n]$. Moreover, $k > n$ also implies that we have $(\mathcal{C}_k)^l[\equiv n, \leq n] = \mathcal{P}^l[n]$. This shows that $\mathcal{X}_\theta(n\delta_l)^{\mu_m} = \mathcal{X}_\theta(n\delta_l)^{\mathbb{C}^\times}$ is in bijection with a subset of $\mathcal{P}^l[n]$. Moreover, by Lemma 4.11, this bijection is independent of the choice of k . But it is well-known (see [12]) that $\mathcal{X}_\theta(n\delta_l)^{\mu_m}$ and $\mathcal{P}^l[n]$ have the same cardinalities. This shows that the bijection above is a bijection between $\mathcal{X}_\theta(n\delta_l)^{\mu_m}$ and $\mathcal{P}^l[n]$ (not just a subset of $\mathcal{P}^l[n]$).

The second statement follows from Lemma 4.11. \square

Corollary 4.13. — *For each $\lambda \in (\mathcal{C}_k)^l[\equiv n, \leq n]$, we have $\mathcal{X}_\lambda \neq \emptyset$. In particular, we have $\mathcal{E}(k, l, n) = \mathcal{E}(k, l, n)_{\neq \emptyset}$.*

Proof. — Assume $\lambda \in (\mathcal{C}_k)^l[\equiv n, \leq n]$. Then there exists $\mu \in \mathcal{P}^l[n]$ such that $\text{Core}_k(\mu) = \lambda$. Then \mathcal{X}_λ is not empty because we have $p_\mu \in \mathcal{X}_\lambda$ by Proposition 4.12. \square

4.D. \mathbb{C}^\times -fixed points. — In the previous section we have constructed a parametrization of the \mathbb{C}^\times -fixed points in $\mathcal{X}_\theta(n\delta_l)$ by $\mathcal{P}^l[n]$. Another parametrization of the \mathbb{C}^\times -fixed points in $\mathcal{X}_\theta(n\delta_l)$ by the same set is done by Gordon [12]. This parametrization is given in terms of baby Verma modules.

It is not obvious at all how to compare these two parametrizations. But this can be deduced from the main theorem in [21].

For $\lambda = (\lambda^0, \dots, \lambda^{l-1}) \in \mathcal{P}^l$ we set $\lambda^\flat = (\lambda^{l-1}, \dots, \lambda^0)$.

Proposition 4.14. — *The parametrization of the \mathbb{C}^\times -fixed points in $\mathcal{X}_\theta(n\delta_l)$ given in Proposition 4.12 differs from Gordon's parametrization by the twist*

$$\mathcal{P}^l[n] \rightarrow \mathcal{P}^l[n], \quad \lambda \mapsto \lambda^b.$$

Proof. — We have constructed in Lemma 4.9 a parametrization of the irreducible components of $\mathcal{X}_\theta(n\delta_l)^{\mu_m}$ by the set $\mathcal{C}_m[\equiv nl] \cap \mathcal{C}_m[\leq nl] \cap (\mathcal{C}_m)_{l,\emptyset}$. Similarly to the proof of Proposition 4.12, this yields a bijection between $\mathcal{X}_\theta(n\delta_l)^{\mathbb{C}^\times}$ and $\mathcal{P}[nl] \cap \mathcal{P}_{l,\emptyset}$.

Indeed, if $k > n$, then we have $\mathcal{X}_\theta(n\delta_l)^{\mathbb{C}^\times} = \mathcal{X}_\theta(n\delta_l)^{\mu_m}$ by Corollary 2.14 and

$$\mathcal{C}_m[\equiv nl] \cap \mathcal{C}_m[\leq nl] \cap (\mathcal{C}_m)_{l,\emptyset} = \mathcal{P}[nl] \cap \mathcal{P}_{l,\emptyset}.$$

Moreover, an argument similar to the proof of Proposition 4.12 shows that the obtained bijection $\mathcal{X}_\theta(n\delta_l)^{\mathbb{C}^\times} \simeq \mathcal{P}[nl] \cap \mathcal{P}_{l,\emptyset}$ is independent of k .

On the other hand, [21, Theorem 1.1 a)] also constructs a bijection between $\mathcal{X}_\theta(n\delta_l)^{\mathbb{C}^\times}$ and $\mathcal{P}[nl] \cap \mathcal{P}_{l,\emptyset}$. It is clear from the construction that this bijection coincides with ours.

Theorem [21, Theorem 1.2] claims that the bijection

$$\mathcal{P}[nl] \cap \mathcal{P}_{l,\emptyset} \rightarrow \mathcal{P}^l[n], \quad \lambda \mapsto (\beta_{l,\emptyset}(\lambda))^b$$

identifies the parametrization of the \mathbb{C}^\times -fixed points by the set $\mathcal{P}[nl] \cap \mathcal{P}_{l,\emptyset}$ with Gordon's parametrization of the \mathbb{C}^\times -fixed points by the set $\mathcal{P}^l[n]$.

On the other hand, the parametrization of the \mathbb{C}^\times -fixed points by $\mathcal{P}[nl] \cap \mathcal{P}_{l,\emptyset}$ composed with the l -quotient bijection $\beta_{l,\emptyset}$ is nothing else but the parametrization of the \mathbb{C}^\times -fixed points by $\mathcal{P}^l[n]$ from Proposition 4.12. \square

Remark 4.15. — The parametrization of the points of $\mathcal{X}_\theta(n\delta_l)^{\mathbb{C}^\times}$ given in Proposition 4.12 is constructed as a parametrization of the irreducible components of $\mathcal{X}_\theta(n\delta_l)^{\mu_m}$ for a very big m . This parametrization can be seen in an " m -independent way" if we replace the quiver \overline{Q}_m with a very big m by an infinite quiver. This can be done in the following way.

Let \overline{Q}_∞ be the quiver defined in the same way as \overline{Q}_l with respect to the vertex set \mathbb{Z} instead of $\mathbb{Z}/l\mathbb{Z}$. The dimension vectors of the quiver \overline{Q}_∞ are in $\mathbb{Z}^{\mathbb{Z}}$. Set

$$\mathbb{Z}_{\text{fin}}^{\mathbb{Z}} = \{\hat{d} \in \mathbb{Z}^{\mathbb{Z}}; \hat{d} \text{ has finitely many non-zero components}\}.$$

Consider the map

$$p: \mathbb{Z}_{\text{fin}}^{\mathbb{Z}} \rightarrow \mathbb{Z}^{\mathbb{Z}/l\mathbb{Z}}, \quad (p(\hat{d}))_i = \sum_{j \equiv i \pmod{l}} \hat{d}_j.$$

For each $\hat{d} \in \mathbb{Z}_{\text{fin}}^{\mathbb{Z}}$, we have a linear map $i_\infty^{(\hat{d})}: \text{Rep}(\overline{Q}_\infty, \hat{d}) \rightarrow \text{Rep}(\overline{Q}_l, p(\hat{d}))$, defined in the same way as in Section 2.D. Now, for each $\theta \in \mathbb{C}^{\mathbb{Z}/l\mathbb{Z}}$ we consider the element $\theta[\infty] \in \mathbb{C}^{\mathbb{Z}}$ given by $\theta[\infty]_i = \theta_{i \pmod{l}}$. Then we obtain a \mathbb{C}^\times -invariant morphism of algebraic varieties $i_\infty^{(\hat{d})}: \mathcal{X}_{\theta[\infty]}(\hat{d}) \rightarrow \mathcal{X}_\theta(p(\hat{d}))$. But the variety $\mathcal{X}_{\theta[\infty]}(\hat{d})$ is obviously \mathbb{C}^\times -stable because the \mathbb{C}^\times action is induced by elements of $\mathbf{GL}(\hat{d})$. Since $\mathcal{X}_{\theta[\infty]}(\hat{d})$ is connected, it is a singleton (if it is not empty). Set

$$\mathcal{E}(\infty, l, d) = \{\hat{d} \in \mathbb{Z}_{\text{fin}}^{\mathbb{Z}}; p(\hat{d}) = d, \mathcal{X}_{\theta[\infty]}(\hat{d}) \neq \emptyset\}.$$

If $\hat{d} \in \mathcal{E}(\infty, l, d)$, the image of $i_\infty^{(\hat{d})}$ is a \mathbb{C}^\times -fixed point in $\mathcal{X}_\theta(d)$. Let us call this point $p_{\hat{d}}$.

Now, we assume $d = n\delta_l$, $n \geq 0$ and that $\mathcal{X}_\theta(d)$ is smooth. The construction of the parametrization of the \mathbb{C}^\times -fixed points in $\mathcal{X}_\theta(n\delta_l)$ given in Proposition 4.12 implies that each \mathbb{C}^\times -fixed point p_μ , $\mu \in \mathcal{P}^l[n]$, in $\mathcal{X}_\theta(n\delta_l)$ is of the form $p_{\hat{d}}$ for a unique $\hat{d} \in \mathcal{E}(\infty, l, n\delta_l)$. Moreover, this \hat{d} is given by $\hat{d} = \text{Res}_\infty(\beta_{l,\emptyset}^{-1}(\mu))$. ■

Lemma 4.16. — *Let $d \in \mathbb{Z}^{\mathbb{Z}/l\mathbb{Z}}$ be such that the pair (d, θ) is smooth (see Definition 2.6). Then by (2.4), the variety $\mathcal{X}_\theta(d)$ is smooth and non-empty. Let ν be the l -core such that we have $d = n\delta_l + \text{Res}_l(\nu)$ (see Proposition 4.4). Then we have the following.*

- (a) *The \mathbb{C}^\times -fixed points in $\mathcal{X}_\theta(d)$ are exactly $p_{\hat{d}}$, $\hat{d} \in \mathcal{E}(\infty, l, d)$ (without repetition).*
- (b) *We have $\mathcal{E}(\infty, l, d) = \{\text{Res}_\infty(\lambda); \lambda \in \mathcal{P}_{l,\nu}^l[|d|]\}$.*

Proof. — The case $d = n\delta_l$ follows from Remark 4.15. But the proof in the general case is completely the same. It is just important to know that under the given assumption on (d, θ) , each representation in $\mathcal{X}_\theta(d)$ is simple. □

The lemma above yields parametrizations of $\mathcal{X}_\theta(d)^{\mathbb{C}^\times}$ by $\mathcal{E}(\infty, l, d)$ and by $\mathcal{P}_{l,\nu}^l[|d|]$. Then the l -quotient bijection $\beta_{l,\nu}: \mathcal{P}_{l,\nu}^l[|d|] \rightarrow \mathcal{P}^l[n]$ (see Lemma 4.10(b)) yields also a parametrization of $\mathcal{X}_\theta(d)^{\mathbb{C}^\times}$ by $\mathcal{P}^l[n]$. For $\mu \in \mathcal{P}^l[n]$ we denote by p_μ the corresponding \mathbb{C}^\times -fixed point of $\mathcal{X}_\theta(d)$ (i.e., we have $p_\mu = p_{\hat{d}}$ if $\hat{d} = \text{Res}_\infty(\beta_{l,\nu}^{-1}(\mu))$). We can write p_μ^d instead of p_μ to stress that it is a point in $\mathcal{X}_\theta(d)^{\mathbb{C}^\times}$.

The group W_l^{aff} acts on $\mathcal{P}^l[n]$ by permutation of components. More precisely, for each $i \in \mathbb{Z}/l\mathbb{Z}$, the element s_i exchanges the components μ^i and μ^{i+1} of $\mu = (\mu^0, \dots, \mu^{l-1})$. It is clear that the kernel of the canonical surjection $W_l^{\text{aff}} \rightarrow W_l$ acts trivially.

Lemma 4.17. — *Let (d, θ) be a smooth pair (see Definition 2.6) in $\mathbb{Z}^{\mathbb{Z}/l\mathbb{Z}} \times \mathbb{C}^{\mathbb{Z}/l\mathbb{Z}}$. Assume that $\theta_i \neq 0$. Then for every $\mu \in \mathcal{P}^l[n]$, the bijection $\mathcal{X}_\theta(d) \simeq \mathcal{X}_{s_i(\theta)}(s_i(d))$ given by (2.4) sends the \mathbb{C}^\times -point $p_\mu^d \in \mathcal{X}_\theta(d)$ to $p_{s_i(\mu)}^{s_i(d)} \in \mathcal{X}_{s_i(\theta)}(s_i(d))$.*

Proof. — There is an action of W_l^{aff} on $\mathbb{Z}_{\text{fin}}^{\mathbb{Z}}$ given by

$$(s_i(\hat{d}))_j = \begin{cases} \delta_{j0} + \hat{d}_{j+1} + \hat{d}_{j-1} - \hat{d}_j & \text{if } i \equiv j \pmod{l}, \\ \hat{d}_j & \text{if } i \not\equiv j \pmod{l}. \end{cases}$$

It is clear that this action lifts the W_l^{aff} -actions on $\mathbb{Z}^{\mathbb{Z}/l\mathbb{Z}}$ with respect to the map $p: \mathbb{Z}_{\text{fin}}^{\mathbb{Z}} \rightarrow \mathbb{Z}^{\mathbb{Z}/l\mathbb{Z}}$.

Let ν and ν' be the l -cores such that we have $d = \text{Res}_l(\nu) + n\delta_l$ and $s_i(d) = \text{Res}_l(\nu') + n\delta_l$. For each $\hat{d} \in \mathcal{E}(\infty, l, d)$, the following diagram is commutative

$$\begin{array}{ccc} \mathcal{X}_{\theta[\infty]}(\hat{d}) & \longrightarrow & \mathcal{X}_{s_i(\theta)[\infty]}(s_i(\hat{d})) \\ \downarrow i_\infty^{(\hat{d})} & & \downarrow i_\infty^{(s_i(\hat{d}))} \\ \mathcal{X}_\theta(d) & \longrightarrow & \mathcal{X}_{s_i(\theta)}(s_i(d)). \end{array}$$

where the horizontal maps are the isomorphisms given by (2.4). This implies that the isomorphism (2.4) identifies the \mathbb{C}^\times -fixed point $p_{\widehat{d}}$ with the \mathbb{C}^\times -fixed point $p_{s_i(\widehat{d})}$. Moreover, it follows from [18, Proposition 4.1.3] that the following diagram is commutative

$$\begin{array}{ccccc}
 \mathcal{P}^l & \xrightarrow{\beta_{l,\nu}^{-1}} & \mathcal{P}_{l,\nu} & \xrightarrow{\text{Res}_\infty} & \mathbb{Z}_{\text{fin}}^{\mathbb{Z}} \\
 \downarrow s_i & & & & \downarrow s_i \\
 \mathcal{P}^l & \xrightarrow{\beta_{l,\nu'}^{-1}} & \mathcal{P}_{l,\nu'} & \xrightarrow{\text{Res}_\infty} & \mathbb{Z}_{\text{fin}}^{\mathbb{Z}}
 \end{array}$$

This implies that (2.4) identifies the \mathbb{C}^\times -fixed point p_μ^d with the \mathbb{C}^\times -fixed point $p_{s_i(\mu)}^{s_i(d)}$. \square

Remark 4.18. — Fix $d \in \mathcal{E}(k, l, n)$. Let r be such that d is in the W_m^{aff} -orbit of $r\delta_m$. Let ν be the m -core such that $d = r\delta_m + \text{Res}_m(\nu)$. Consider the irreducible component \mathcal{X}_d of $\mathcal{X}(n\delta_l)^{\mu_m}$. By Theorem 2.13 and the proof of Theorem 1.3, we have an isomorphism

$$\mathcal{X}_d \simeq \mathcal{X}_{\theta'}(r\delta_m)$$

for some θ' . This isomorphism depends on a choice of an element $w \in W_m^{\text{aff}}$ such that $w(d) = r\delta_m$ (see the proof of Theorem 1.3) and of a choice of a reduced expression of w that allows to apply a sequence of isomorphisms (2.4). We will assume that the element w is chosen as in the proof of Lemma 2.8. In particular w is in the kernel of the canonical surjection $W_m^{\text{aff}} \rightarrow W_m$. This choice implies that the bijection $\mathcal{X}_{\theta[k]}(d) \simeq \mathcal{X}_{w(\theta[k])}(r\delta_m)$ yields the following bijection on the \mathbb{C}^\times -fixed points (see Lemma 4.17):

$$\mathcal{X}_{\theta[k]}^{\mathbb{C}^\times}(d) \rightarrow \mathcal{X}_{w(\theta[k])}^{\mathbb{C}^\times}(r\delta_m), \quad p_\mu^d \mapsto p_\mu^{r\delta_m}.$$

The inclusion $\mathcal{X}_{\theta'}(r\delta_m)^{\mathbb{C}^\times} \subset \mathcal{X}_\theta(n\delta_l)^{\mathbb{C}^\times}$ yields a map

$$(4.18) \quad \mathcal{P}^m[r] \rightarrow \mathcal{P}^l[n].$$

Let us describe this map.

We have the following bijections obtained from the l -quotient (or m -quotient) map (see also Lemma 4.10(b))

$$\mathcal{P}^m[r] \simeq \mathcal{P}_{m,\nu}[mr + |\nu|], \quad \mathcal{P}^l[n] \simeq \mathcal{P}_{l,\emptyset}[nl].$$

Note that we have $mr + |\nu| = |d| = nl$. Moreover, we have $\mathcal{P}_{m,\nu}[mr + |\nu|] \subset \mathcal{P}_{l,\emptyset}[nl]$ because every partition with m -core ν has a trivial l -core (since the l -core of ν is trivial). Then under the isomorphisms above, the inclusion $\mathcal{P}_{m,\nu}[mr + |\nu|] \subset \mathcal{P}_{l,\emptyset}[nl]$ is exactly the map (4.18). Indeed, this follows from Lemma 4.17 (and the choice of w above) and from the commutativity of the following diagram

$$\begin{array}{ccc}
 & \mathcal{X}_{\theta[\infty]}(\widehat{d}) & \\
 i_\infty^{(\widehat{d})} \swarrow & & \searrow i_\infty^{(\widehat{d})} \\
 \mathcal{X}_{\theta[k]}(d) & \xrightarrow{i_k^{(d)}} & \mathcal{X}_\theta(n\delta_l)
 \end{array}$$

for each $\widehat{d} \in \mathcal{E}(\infty, m, d)$.

Now, let us describe the inclusion map $\mathcal{P}_{m,\nu}[mr + |\nu|] \rightarrow \mathcal{P}_{l,\emptyset}[nl]$ as a map $\mathcal{P}^m[r] \rightarrow \mathcal{P}^l[n]$. Set $\gamma = \beta_{l,\emptyset}(\nu)$. Then, by Lemma 4.10, we have $\gamma = (\gamma^0, \dots, \gamma^{l-1}) \in (\mathcal{C}_k)^l$. For each $\lambda \in \mathcal{P}^l[n]$ such that $\text{Core}_k(\lambda) = \gamma$, we define $\beta_{k,\gamma}(\lambda)$ to be the m -partition $\mu = (\mu^0, \mu^1, \dots, \mu^{m-1})$ defined by

$$\beta_{k,\gamma^i}(\lambda^i) = (\mu^i, \mu^{i+l}, \dots, \mu^{i+(k-1)l})$$

for all $i \in \{0, 1, \dots, l-1\}$. The map $\beta_{k,\gamma}$ is an isomorphism between $\mathcal{P}_{k,\gamma}^l[n]$ and $\mathcal{P}^m[r]$, where $\mathcal{P}_{k,\gamma}^l[n] = \{\lambda \in \mathcal{P}^l[n]; \text{Core}_k(\lambda) = \gamma\}$. The following diagram is commutative

$$\begin{array}{ccc} \mathcal{P}_{m,\nu}[mr + |\nu|] & \longrightarrow & \mathcal{P}_{l,\emptyset}[nl] \\ \beta_{m,\nu} \downarrow & & \downarrow \beta_{l,\emptyset} \\ \mathcal{P}^m[r] & \xrightarrow{\beta_{k,\gamma}^{-1}} & \mathcal{P}^l[n] \end{array}$$

This implies that the map $\mathcal{P}^m[r] \rightarrow \mathcal{P}^l[n]$ obtained from the inclusion $\mathcal{X}_{\theta'}(r\delta_m)^{\mathbb{C}^\times} \subset \mathcal{X}_{\theta}(n\delta_l)^{\mathbb{C}^\times}$ is given by $\beta_{k,\gamma}^{-1}$. ■

4.E. Explicit construction of c' . — We know from Theorem 2.13 that each irreducible component \mathcal{X} of $\mathcal{X}_{\theta}(n\delta_l)^{\mu_m}$ is of the form $\mathcal{X} = \mathcal{X}_d$. Moreover, by Theorem 1.3 there exists a reflection subquotient (V', W') of (V, W) and a complex-valued map $c': W' \rightarrow \mathbb{C}$ such that $\mathcal{X} \simeq \mathcal{X}_{c'}(V', W')$. The reflection subquotient was constructed in the proof of Theorem 1.3. Now we give an explicit construction of c' .

Recall from §1.A that the parameter c is determined by the sequence of parameters $(a, k_0, k_1, \dots, k_{l-1})$. Similarly, the parameter c' is determined by $(a', k'_0, k'_1, \dots, k'_{m-1})$. We are going to give explicit expressions for $(a', k'_0, k'_1, \dots, k'_{m-1})$ in terms of $d \in \mathcal{E}(k, l, m)$ and $(a, k_0, k_1, \dots, k_{l-1})$.

First, we recall the construction of c' in the proof of Theorem 1.3. There is an element $w \in W_m^{\text{aff}}$ such that $w(d)$ is a multiple of δ_m . Then we put $\theta' = w(\theta[k])$. Finally, we can obtain $(a', k'_0, \dots, k'_{m-1})$ from θ' using (3.1) and (3.2).

This construction of c' is not canonical because there is no preferable choice of an element w . Note that δ_m is stable by the subgroup W_m of W_m^{aff} . So the element w can be replaced by any elements of $W_m w$. (By Remark 3.5, a different choice of w yields just a permutation of the parameters $k'_0, k'_1, \dots, k'_{m-1}$.) The affine Weyl group W_m^{aff} is of the form $W_m \ltimes R_m$, where R_m is the root lattice for \mathfrak{sl}_m . In particular, there is an element of $w' \in W_m w$ that is in R_m . More precisely, we have $w' = t_a$, where $a = d - d_0 \delta_m$ (see the proof of Lemma 2.8).

Then, by Lemma 2.9, we have $w'(\theta[k]) = t_a(\theta[k]) = \theta[k] + \Sigma(\theta[k]) \cdot \bar{a} = \theta[k] + k\Sigma(\theta) \cdot \bar{d}$. (Where the notation \bar{a} and \bar{d} is as in Section 2.B.) This means that we have

$$\theta'_i = (\theta[k])_i + k\Sigma(\theta)(2d_i - d_{i-1} + d_{i+1}) = \theta_{(i \bmod l)} - ka(2d_i - d_{i-1} + d_{i+1})$$

Passing back to the k_i -notations, we get $a' = ka$ and

$$k'_i = k_{(i \bmod l)} + a \left(\left\lfloor \frac{i-1}{l} \right\rfloor - \frac{k-1}{2} + k(d_{1-i} - d_{-i}) \right)$$

for $i = 1, \dots, m-1, m$. (Here $k'_0 = k'_m$.)

Example 4.19. — Let us make the above formula explicit in the case $l = 1$. In this case we have $m = k$. Then we get $\theta = \theta_0 = -a$ and $\theta[k] = (-a, -a, \dots, -a)$ (k times). This gives

$$\theta'_i = -a - ak(2d_i - d_{i-1} - d_{i+1}),$$

for $i \in \mathbb{Z}/k\mathbb{Z}$.

$$k'_i = a \left(i - \frac{k+1}{2} + k(d_{1-i} - d_{-i}) \right)$$

for $i = 1, \dots, k-1, k$. (Here $k_0 = k_k$.) ■

Example 4.20. — Assume $l = 2$, $k = 2$ and $d = (0, 0)$. In this case c is determined by (a, k_0, k_1) such that $k_0 + k_1 = 0$. Put $k_1 = b$ (and then $k_0 = -b$). Let us make the formulas for k'_i explicit in this case. We have

$$\begin{cases} k'_0 &= -b + a/2, \\ k'_1 &= b - a/2, \\ k'_2 &= -b - a/2, \\ k'_3 &= b + a/2. \end{cases}$$

w

4.F. Recollection. — We will rewrite here the statement of Theorem 2.13 with the help of the combinatorics developed in this section. First, we will denote by $\mathcal{Z}_c(l, 1, n)$ the Calogero-Moser space associated with the complex reflection group $G(l, 1, n)$ and with the parameter c . Here, we label \mathbb{C}^\times -fixed points using Gordon's convention, which differs from our usual convention by a twist (see Proposition 4.14). Recall that $(\mathcal{C}_k)^l \equiv [n, \leq n]$ is the set of l -tuples $\gamma = (\gamma^0, \dots, \gamma^{l-1})$ of k -cores such that

$$|\gamma| \leq n \text{ and } |\gamma| \equiv n \pmod{k}.$$

If $\lambda = (\lambda^0, \dots, \lambda^{l-1}) \in \mathcal{D}^l[n]$ and $\text{Core}_k(\lambda) = \gamma = (\gamma^0, \dots, \gamma^{l-1})$, then we define $\beta_{k,\gamma}^b(\lambda)$ to be the m -partition $\mu = (\mu^0, \mu^1, \dots, \mu^{m-1})$ defined by

$$\beta_{k,\gamma^i}(\lambda^i) = (\mu^{i+(k-1)l}, \dots, \mu^{i+l}, \mu^i)$$

for all $i \in \{0, 1, \dots, l-1\}$. (In other words, we have $\beta_{k,\gamma}^b(\lambda) = (\beta_{k,\gamma}(\lambda^b))^b$.) Also, if $\lambda \in \mathcal{D}^l[n]$, we denote by $\chi_\lambda^{(l)}$ the associated irreducible character of $G(l, 1, n)$ and by $z_\lambda^{(l)}$ the \mathbb{C}^\times -fixed point in $\mathcal{Z}_c(l, 1, n)$ associated with $\chi_\lambda^{(l)}$ by [12, Corollary 5.8].

Theorem 4.21. — Assume that $\mathcal{Z}_c(l, 1, n)$ is smooth and let $(a, k_0, k_1, \dots, k_{l-1})$ be the family of parameters associated with c as in §3.A. Then $\mathcal{Z}_c(l, 1, n)^{\mu_{k_i}}$ is smooth (so its irreducible components coincide with its connected components) and:

- (a) There is a bijection $\gamma \mapsto \mathcal{Z}(\gamma)$ between the set $(\mathcal{C}_k)^l \equiv [n, \leq n]$ and the set of irreducible components of $\mathcal{Z}_c(l, 1, n)^{\mu_{k_i}}$ such that $z_\lambda^{(l)} \in \mathcal{Z}(\gamma)$ if and only if $\text{Core}_k(\lambda) = \gamma$ (for any $\lambda \in \mathcal{D}^l[n]$).

(b) Let $\gamma \in (\mathcal{C}_k)^l [\equiv n, \leq n]$ and let $r = (n - |\gamma|)/k$. Then there is an isomorphism of varieties

$$i_\gamma : \mathcal{Z}_{c'}(lk, 1, r) \xrightarrow{\sim} \mathcal{Z}(\gamma),$$

where $c' : \text{Ref}(G(lk, 1, r)) \rightarrow \mathbb{C}$ is the parameter associated with the family $(a', k'_0, k'_1, \dots, k'_{k-1})$ such that

$$\begin{cases} a' = ka, \\ k'_j = k_{(j \bmod l)} + a \left(\left\lfloor \frac{j-1}{l} \right\rfloor - \frac{k-1}{2} + k(d_{1-j} - d_{-j}) \right), \end{cases} \quad 1 \leq j \leq m \text{ and } k'_0 = k'_m,$$

which satisfies $z_{(\beta_{k,\gamma}^b)^{-1}(\mu)}^{(l)} = i_\gamma(z_\mu^{(kl)})$ for all $\mu \in \mathcal{D}^{kl}[r]$. Here, $d = (d_i)_{i \in \mathbb{Z}/k\mathbb{Z}}$ is defined by

$$d = \text{res}_m(v) + r\delta_m,$$

where $v = (\beta_{l,\emptyset})^{-1}(\gamma)$ is an m -core.

Corollary 4.22. — Let $\gamma \in (\mathcal{C}_k)^l [\equiv n, \leq n]$ and let $r = (n - |\gamma|)/k$.

$$i_\gamma^* : Z(\text{CG}(l, 1, n)) \longrightarrow Z(\text{CG}(kl, 1, r))$$

be the (surjective) morphism of algebras defined by

$$i_\gamma^*(e_{\lambda\lambda}) = \begin{cases} e_{\chi_{\beta_{k,\gamma}^b}(\lambda)} & \text{if } \text{Core}_k(\lambda) = \gamma, \\ 0 & \text{otherwise.} \end{cases}$$

Then $i_\gamma^*(\mathcal{F}_i(Z(\text{CG}(l, 1, n)))) \subset \mathcal{F}_i(Z(\text{CG}(kl, 1, r)))$ for all i .

Remark 4.23. — Note that the statement of Corollary 4.22 involves only the character theory of groups of type $G(l, 1, n)$ and the combinatorics of partitions. However, our proof relies on the geometric results on quiver varieties proved here as well as the computation of the equivariant cohomology of smooth Calogero-Moser spaces done in [5].

It is a natural question to ask whether such a result can be proved directly by combinatorial methods. ■

5. Type G_4

Hypothesis. In this section, and only in this section, we assume that $V = \mathbb{C}^2$, we identify $\mathbf{GL}_{\mathbb{C}}(V)$ with $\mathbf{GL}_2(\mathbb{C})$, we fix a primitive third root of unity ω , we set

$$s = \begin{pmatrix} \omega & 0 \\ \omega^2 & 1 \end{pmatrix} \quad \text{and} \quad t = \begin{pmatrix} 1 & -\omega^2 \\ 0 & \omega \end{pmatrix}$$

and we assume that $W = \langle s, t \rangle$, so that W is of type G_4 in Shephard-Todd classification. We denote by (y_1, y_2) the canonical basis of V and by (x_1, x_2) its dual basis. We also set $\rho = 1 + 2\omega$, so that $\rho^2 = -3$.

In this case, we will check Conjecture F even if \mathcal{X}_c is not smooth (and for all elements $\sigma \in \mathbf{GL}_C(V)$ normalizing W , as any such elements belong to $\mathbb{C}^\times \cdot W$, as we will see in the proof below).

Theorem 5.1. — *If W is of type G_4 , then Conjecture F holds.*

Proof. — Let $\sigma \in \mathbf{GL}_2(\mathbb{C})$ be of finite order and normalizing W . The conjugacy class of s is

$$\{s, t, sts^{-1}, tst^{-1}\},$$

and is the unique conjugacy class of reflections of determinant ω . Since $\sigma s \sigma^{-1}$ is a reflection of W of determinant ω , there exists $w \in W$ such that $w \sigma$ commutes with s . Since W acts trivially on \mathcal{X}_c , we may, and we will, assume that $\sigma s \sigma^{-1} = s$. Now, $\sigma t \sigma^{-1}$ belongs to this conjugacy class and is different from s , so there exists $i \in \mathbb{Z}$ such that $s^i \sigma$ commutes with t . So, replacing σ by $s^i \sigma$, we may, and we will, assume that σ commutes with s and t . In other words, σ is a root of unity.

Since $Z(W) \simeq \mu_2$, we may also assume that the order of σ is even, equal to $2d$ for some $d \geq 2$ (because the case $d = 1$ is obvious). Let I_d denote the ideal of \mathbf{Z}_c generated by the $\sigma(z) - z$, where z runs over a set of generators of \mathbf{Z}_c . Then

$$(\clubsuit) \quad \mathbb{C}[\mathcal{X}_c^\sigma] = \mathbf{Z}_c / \sqrt{I_d}.$$

Let ζ denote the root of unity by which σ acts: then $\sigma(z) - z = (\zeta^l - 1)z$ if z is homogeneous of \mathbb{Z} -degree l . This shows that I_d is generated by the generators of \mathbf{Z}_c of degree not divisible by $2d$. As we will see below, there is a set of generators of \mathbf{Z}_c whose \mathbb{Z} -degrees belong to the interval $[-6, 6]$. So, if $d \geq 4$, then $\mathcal{X}_c^\sigma = \mathcal{X}_c^{\mathbb{C}^\times}$, so it is a finite set and then Conjecture F is checked in this case. So we may assume that $d \in \{2, 3\}$.

Note that there is only one W -orbit of reflecting hyperplanes. So, as in Example 1.2, we set

$$k_j = \frac{1}{l} \sum_{i=1}^2 \omega^{-i(j-1)} c_{si},$$

so that $k_0 + k_1 + k_2 = 0$.

In [6], Thiel and the first author have provided algorithms for computing presentations of the algebra \mathbf{Z}_c , and such algorithms have been included in the MAGMA package CHAMP developed by Thiel [22] (about the MAGMA software, see [20]). These algorithms provide an explicit presentation of the center \mathbf{Z}_c given as follows:

Generators:

$$Y_1, Y_2, X_1, X_2, \text{eu}, A, B, D;$$

Relations:

$$\left\{ \begin{array}{l}
\text{(Z1)} \quad eu^4 + 4\rho Y_1 X_1 - AB + 3\rho euD + 18(k_1^2 + k_1 k_2 + k_2^2)eu^2 \\
\quad + 756(k_1^2 k_2 + k_1 k_2^2)eu = 0 \\
\text{(Z2)} \quad \rho eu^3 A - 4X_1 euB - DA - \rho Y_1 X_2 - 30\rho(k_1^2 + k_1 k_2 + k_2^2)euA \\
\quad - 108\rho(k_1^2 k_2 + k_1 k_2^2)A = 0 \\
\text{(Z3)} \quad Y_2 A - euB^2 + \rho Y_1 D + 54(k_1^2 + k_1 k_2 + k_2^2)Y_1 eu - 324(k_1^2 k_2 + k_1 k_2^2)Y_1 = 0 \\
\text{(Z4)} \quad \rho eu^3 B - 4X_1 Y_2 - \rho Y_1 euA - BD - 30\rho(k_1^2 + k_1 k_2 + k_2^2)euB \\
\quad - 108\rho(k_1^2 k_2 + k_1 k_2^2)B = 0 \\
\text{(Z5)} \quad euA^2 - 4X_1 D - X_2 B + 72\rho(k_1^2 + k_1 k_2 + k_2^2)X_1 eu - 432\rho(k_1^2 k_2 + k_1 k_2^2)X_1 = 0 \\
\text{(Z6)} \quad Y_2 eu^3 + 3Y_1^2 A - 3Y_1 eu^2 B - B^3 + 3\rho Y_2 D + 144(k_1^2 + k_1 k_2 + k_2^2)Y_1 B \\
\quad + 18(k_1^2 + k_1 k_2 + k_2^2)Y_2 eu + 756(k_1^2 k_2 + k_1 k_2^2)Y_2 = 0 \\
\text{(Z7)} \quad 4\rho X_1 eu^2 A - A^3 - 16X_1^2 B + X_2 eu^3 + 3\rho X_2 D + 18(k_1^2 + k_1 k_2 + k_2^2)X_2 eu \\
\quad - 192\rho(k_1^2 + k_1 k_2 + k_2^2)X_1 A + 756(k_1^2 k_2 + k_1 k_2^2)X_2 = 0 \\
\text{(Z8)} \quad eu^6 + 4\rho Y_1 X_1 eu^2 + 2\rho D eu^3 - Y_2 X_2 + D^2 - 36(k_1^2 + k_1 k_2 + k_2^2)eu^4 \\
\quad + 12\rho(k_1^2 + k_1 k_2 + k_2^2)euD + 216\rho(k_1^2 k_2 + k_1 k_2^2)D + 1080(k_1^2 k_2 + k_1 k_2^2)eu^3 \\
\quad + 1620(k_1^2 + k_1 k_2 + k_2^2)^2 eu^2 - 3888(k_1^2 k_2 + k_1 k_2^2)(k_1^2 + k_1 k_2 + k_2^2)eu \\
\quad - 34992(k_1^2 k_2 + k_1 k_2^2)^2 = 0 \\
\text{(Z9)} \quad \rho Y_1 A^2 + 4X_1 B^2 + 4eu^3 D - \rho Y_2 X_2 + 4\rho D^2 - 24\rho(k_1^2 + k_1 k_2 + k_2^2)eu^4 \\
\quad - 288(k_1^2 + k_1 k_2 + k_2^2)euD - 576(k_1^2 + k_1 k_2 + k_2^2)Y_1 X_1 \\
\quad + 2160\rho(k_1^2 + k_1 k_2 + k_2^2)^2 eu^2 + 432\rho(k_1^2 k_2 + k_1 k_2^2)eu^3 - 864(k_1^2 k_2 + k_1 k_2^2)D \\
\quad + 20736\rho(k_1^2 k_2 + k_1 k_2^2)(k_1^2 + k_1 k_2 + k_2^2)eu + 46656\rho(k_1^2 k_2 + k_1 k_2^2)^2 = 0
\end{array} \right.$$

The generators have \mathbb{Z} -degrees given by the following map

$$(Y_1, Y_2, X_1, X_2, eu, A, B, D) \longmapsto (-4, -6, 4, 6, 0, 2, -2, 0).$$

Moreover, in this presentation, $\mathbb{C}[V]^W = \mathbb{C}[X_1, X_2]$ and $\mathbb{C}[V^*]^W = \mathbb{C}[Y_1, Y_2]$.

We can deduce from this that \mathcal{X}_c is smooth if and only if

$$(\diamond) \quad k_0 k_1 k_2 (k_0 - k_1)(k_0 - k_2)(k_1 - k_2) \neq 0.$$

First case: assume that $d = 2$. In this case, the above presentation shows that, whenever (\diamond) holds, then \mathcal{X}_c^σ has four irreducible components \mathcal{X} , p , q and r where \mathcal{X} has dimension 2 and is isomorphic to

$$(\heartsuit) \quad \mathcal{X} \simeq \{(x, y, e) \in \mathbb{C}^3 \mid e(e - 12k_0)(e - 12k_1)(e - 12k_2) = xy\},$$

and p , q and r are three points (which belong to $\mathcal{X}_c^{\mathbb{C}^\times}$). If (\diamond) does not hold, then \mathcal{X}_c^σ still contains an irreducible component of dimension 2 with the same description as above, and a few other isolated points (the number depends on the exact values of the k_i). By Remark 1.4, we only need to consider the irreducible component \mathcal{X} . But it follows from (\heartsuit) and Example 1.2 that then \mathcal{X} is isomorphic to the Calogero-Moser associated with a cyclic group of order 4 acting on a one-dimensional space: as there is an element w of order 4 in W , one can choose for V' an eigenspace of w and for W' the cyclic group $\langle w \rangle$ and Conjecture F is checked in this case (note also that the map $c \mapsto c'$ is linear).

Second case: assume that $d = 3$. In this case, the above presentation shows that, whenever (\diamond) holds, then \mathcal{X}_c^σ has two irreducible components \mathcal{X} and p where \mathcal{X} has dimension 2 and is isomorphic to

$$(\spadesuit) \quad \mathcal{X} \simeq \{(x, y, e) \in \mathbb{C}^3 \mid (e + 6k_0)(e + 6k_1)(e + 6k_2)(e - 12k_0)(e - 12k_1)(e - 12k_2) = xy\},$$

and p is a point (which belongs to $\mathcal{X}_c^{\mathbb{C}^\times}$). If (\diamond) does not hold, then \mathcal{X}_c^σ still contains an irreducible component of dimension 2 with the same description as above, and maybe one extra point according to the values of the k_i . By Remark 1.4, we only need to consider the irreducible component \mathcal{X} . But it follows from (\spadesuit) and Example 1.2 that then \mathcal{X} is isomorphic to the Calogero-Moser associated with a cyclic group of order 6 acting on a one-dimensional space: as $-s \in W$ is of order 6, one can choose for V' the ω -eigenspace of s and for W' the cyclic group $\langle -s \rangle$ and Conjecture F is checked in this case (note also that the map $c \mapsto c'$ is linear). \square

References

- [1] G. BELLAMY, *On singular Calogero-Moser spaces*, Bull. LMS **41** (2009), 315-326.
- [2] G. BELLAMY, T. SCHEDLER & U. THIEL, *Hyperplane arrangements associated to symplectic quotient singularities*, Banach Center Publications **116** (2018), 25-45.
- [3] C. BERG, B. JONES & M. VAZIRANI, *A bijection on core partitions and a parabolic quotient of the affine symmetric group*, Journal of Combinatorial Theory, Series A, **116(8)**, (2009), 1344-1360.
- [4] C. BONNAFÉ & R. ROUQUIER, *Cherednik algebras and Calogero-Moser cells*, preprint (2017), arXiv:1708.09764.
- [5] C. BONNAFÉ & P. SHAN, *On the cohomology of Calogero-Moser spaces*, Int. Math. Res. Not. (2018).
- [6] C. BONNAFÉ & U. THIEL, *Calogero-Moser families and cellular characters: computational aspects*, in preparation.
- [7] M. BROUÉ & G. MALLE, *Zyklotomische Heckealgebren*, Astérisque **212** (1993), 119-189.
- [8] M. BROUÉ & J. MICHEL, *Sur certains éléments réguliers des groupes de Weyl et les variétés de Deligne-Lusztig associées*, Progress in Math. **141**, Birkhauser, 1996, 73-139.
- [9] W. CRAWLEY-BOEVEY, *Geometry of the moment map for representations of quivers*, Compo. Math. **126** (2001), 257-293.
- [10] P. ETINGOF & V. GINZBURG, *Symplectic reflection algebras, Calogero-Moser space, and deformed Harish-Chandra homomorphism*, Inventiones Mathematicae **147**, no. 2 (2002), 243-348.
- [11] V. GINZBURG & D. KALEDIN, *Poisson deformations of symplectic quotient singularities*, Adv. in Math. **186** (2004), 1-57.
- [12] I. GORDON, *Baby Verma modules for rational Cherednik algebras*, Bull. London Math. Soc. **35** (2003), 321-336.
- [13] I. GORDON, *Quiver varieties, category \mathcal{O} for rational Cherednik algebras, and Hecke algebras*, Int. Math. Res. Papers (2008), 69 pages.
- [14] I. GORDON & M. MARTINO, *Calogero-Moser space, restricted rational Cherednik algebras, and two-sided cells*, Math. Res. Lett. **16** (2009), 255-262.
- [15] V. G. KAC, *Infinite Dimensional Lie Algebras - An Introduction*, Birkhäuser, 1983
- [16] D. KALEDIN, *On crepant resolutions of symplectic quotient singularities*, Selecta Math. **9** (2003), 529-555.
- [17] L. LE BRUYN, *Noncommutative smoothness and coadjoint orbits*, J. of Algebra **258** (2002), 60-70.
- [18] M. VAN LEEUWEN, *Edge sequences, ribbon tableaux, and an action of affine permutations*, Europ. J. of Comb. **20** (1999), 179-195.
- [19] G. LUSZTIG, *Quiver varieties and Weyl group actions*, Ann. Inst. Fourier **50** (2000), 461-489.
- [20] W. BOSMA, J. CANNON & C. PLAYOUST, *The Magma algebra system. I. The user language*, J. Symbolic Comput. **24** (1997), 235-265.
- [21] T. PRZEZDZIECKI, *The combinatorics of \mathbb{C}^\times -fixed points in generalized Calogero-Moser spaces and Hilbert schemes*, J. Algebra **556** (2020), 936-992.

- [22] U. THIEL, *Champ: a Cherednik algebra Magma package*, LMS J. Comput. Math. **18** (2015), 266-307.
- [23] É. VASSEROT, *Sur l'anneau de cohomologie du schéma de Hilbert de \mathbb{C}^2* , C. R. Acad. Sci. **332** (2001), 7-12.

June 18, 2020

CÉDRIC BONNAFÉ, Institut Montpellierain Alexander Grothendieck (CNRS: UMR 5149), Université Montpellier 2, Case Courrier 051, Place Eugène Bataillon, 34095 MONTPELLIER Cedex, FRANCE
E-mail: `cedric.bonnafe@umontpellier.fr`

RUSLAN MAKSIMAU, Institut Montpellierain Alexander Grothendieck (CNRS: UMR 5149), Université Montpellier 2, Case Courrier 051, Place Eugène Bataillon, 34095 MONTPELLIER Cedex, FRANCE
E-mail: `ruslan.maksimau@umontpellier.fr`