



HAL
open science

Impacts that cause the highest direct mortality of individuals do not necessarily have the greatest influence on temperate eel escapement

Maria Mateo, Patrick Lambert, Stéphane Tétard, Hilaire Drouineau

► To cite this version:

Maria Mateo, Patrick Lambert, Stéphane Tétard, Hilaire Drouineau. Impacts that cause the highest direct mortality of individuals do not necessarily have the greatest influence on temperate eel escapement. *Fisheries Research*, 2017, 193, pp.51-59. 10.1016/j.fishres.2017.03.024 . hal-01728986

HAL Id: hal-01728986

<https://hal.science/hal-01728986v1>

Submitted on 12 Mar 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 Impacts that cause the highest direct mortality of individuals do not necessarily have the greatest
2 influence on temperate eel escapement

3 Maria Mateo¹, Patrick Lambert¹, Stéphane Tétard², and Hilaire Drouineau¹

4
5 ¹ Irstea, UR EABX Ecosystèmes aquatiques et changements globaux, HYNES Irstea-EDF R&D, 50
6 avenue de Verdun, 33612 Cestas, France

7 ² EDF R&D, HYNES Irstea-EDF R&D, Laboratoire National d'Hydraulique et Environnement
8 quai Watier, 78401 Chatou, France

9 **Abstract**

10 Temperate eels are three panmictic catadromous species with a long period of oceanic passive larval
11 drift and large distribution areas in contrasting environments. Spatial patterns of life history traits
12 have been observed, and are correlated with environmental gradients, and may arise from both
13 adaptive phenotypic plasticity and genetic polymorphism. This raises the question of the effect of
14 spatially heterogeneous anthropogenic pressures on these populations. In this context, we used
15 Geneveel, an individual-based optimization model that includes both phenotypic plasticity and
16 genetic polymorphism, to explore the effects of different kinds of anthropogenic pressures: glass eel
17 and silver eel fisheries, obstacles to upstream migration, and turbine mortality. More specifically,
18 we analyzed the effects of these pressures on five output variables: the number of escapees, the
19 proportion of females, the proportion of slow growers, the mean length-at-silvering and the
20 resulting egg production. Our results suggest that phenotypic plasticity could act as a compensatory
21 mechanism that mitigates the effects of some pressures (glass eel fishery and obstacles to upstream
22 migration) and could be a source of resilience for the population, while other pressures did not show
23 any compensatory effect (silver eel fishery and turbine mortality). Therefore, global impacts are
24 very hard to assess, and the pressure that kills the most individuals does not necessarily have the
25 biggest impact on the spawning biomass.

26 **Keywords:**

27 Temperate eels; anthropogenic pressures; adaptive mechanisms; life history traits; modeling

28 Corresponding author: Maria Mateo

29 Phone number: +33 (0)5 57 89 09 98

30 Email: maria.mateo@irstea.fr

31 **1. Introduction**

32 Temperate eels (*Anguilla rostrata*, *A. anguilla*, *A. japonica*) are three catadromous species that
33 reproduce at sea and grow in continental waters. They display remarkable similarities in life history
34 traits (Daverat et al., 2006; Edeline, 2007). The three populations are panmictic (Als et al., 2011;
35 Han et al., 2010; Pujolar, 2013). Reproduction occurs in the Sargasso Sea for *A. anguilla* and *A.*
36 *rostrata* (McCleave, 1993; Schmidt, 1923) and west of the Mariana Islands for *A. japonica*
37 (Tsukamoto, 1992). Larvae (known as leptocephali) are subject to a long and passive trans-oceanic
38 drift. When arriving on the continental shelves, leptocephali metamorphose into young and
39 transparent eels, called glass eels (Tesch, 2003), and enter continental waters, where they become
40 pigmented yellow eels, the immature adult stage. After a variable period, generally lasting from
41 three to 15 years, yellow eels metamorphose again into silver eels (this metamorphosis is generally
42 called “silvering”), the seaward migration stage. They achieve their sexual maturation while
43 migrating back to spawning grounds. Because of the long larval drift, their distribution area is very
44 large, and the growth phase can occur in very heterogeneous river basins, from Morocco to Norway
45 (Tesch, 2003) for the European eel, from Venezuela to Greenland for *A. rostrata* (Helfman et al.,
46 1987), and from the northern Philippines to Korea for *A. japonica* (Tsukamoto, 1992).

47 Concomitantly to this environmental heterogeneity, temperate eels display remarkable life
48 history trait patterns at both distribution and river catchment scales (Vélez-Espino and Koops,
49 2009). Among them, the sex ratio is highly variable at different spatial scales: female-biased sex
50 ratios are generally observed in the northern part of the distribution area, while male-biased sex
51 ratios are observed in the southern part (Kettle et al., 2011). At the catchment scale, sex ratios are
52 male biased in downstream habitats (Oliveira and McCleave, 2000; Tesch, 2003). This question of
53 sex ratio is relevant since eel sex is not genetically determined, but depends on environmental
54 conditions (Davey and Jellyman, 2005; Geffroy, 2012; Geffroy and Bardonnet, 2015), and males
55 and females display very different life history tactics (Helfman et al., 1987). Males are assumed to
56 have a time-minimizing strategy; since their reproductive success does not depend on their size,
57 they are assumed to leave continental waters as soon as they reach the minimal size required to
58 successfully migrate back to spawning grounds (Oliveira, 1999; Van Den Thillart et al., 2007;
59 Vollestad, 1992). On the other hand, female reproductive success is assumed to be a trade-off
60 between size at maturity (known as length-at-silvering) and survival, called a size-maximizing
61 strategy. This size-maximizing strategy assumption is supported by the observation of larger
62 females in the northern part of the distribution area (slow growth but lower mortality) than in the
63 southern part (Davey and Jellyman, 2005; Helfman et al., 1984).

64 Indeed, eels display a large range of tactics in terms of habitat use for growth, with some
65 settling in estuarine waters, while others moving far upstream in river catchments (Arai and Chino,
66 2012; Daverat et al., 2006; Tsukamoto et al., 1998). Higher densities are observed in downstream
67 habitats, which are the most accessible and the most favorable in terms of growth rates (Daverat et
68 al., 2012; Helfman et al., 1984; Melià et al., 2006). However, other eels also settle in the upper parts
69 of river catchments. Edeline (2007) proposed that more limited intraspecific competition may
70 overcome migratory energy costs, such as loss in growth rate. This assumption was challenged by
71 Cairns et al. (2009), who did not observe a decrease in natural mortality in upstream habitats that
72 would outweigh the cost of migration and the decrease in growth rate.

73 Few studies have explored whether the observed spatial phenotypic patterns are the result of
74 adaptive mechanisms to environmental variability, and they therefore remain poorly understood.
75 The panmixia and long and passive larval drifts impair the possibility of local adaptation.
76 Phenotypic plasticity has been proposed as an adaptive response to environment variability for
77 many species (Gotthard and Nylin, 1995; Levins, 1963; Pigliucci, 2005), and Ernande and
78 Dieckmann (2004) demonstrated that density dependence favors the selection of plastic phenotypes.
79 In this context, assuming that phenotypic plasticity is an adaptive response to environmental
80 variability, Drouineau et al. (2014) developed an optimization model that was able to mimic most of

81 the observed patterns at both the distribution area and river catchment scales. The existence of
82 correlations between genotypic patterns and environmental conditions have been observed (Boivin
83 et al., 2015; Côté et al., 2015, 2014, 2009; Gagnaire et al., 2012; Pujolar et al., 2014; Ulrik et al.,
84 2014). Côté et al. (2014, 2009) and Boivin et al. (2015) observed differences in growth rates
85 depending on eel origin that were preserved after several months in common garden experiments.
86 Pujolar et al. (2014), Gagnaire et al. (2012) and Ulrik et al. (2014) observed patterns in single-
87 nucleotide polymorphisms correlated with environmental conditions. They assumed that individual
88 genetic differences contribute to the emergence of phenotypic spatial patterns because of differential
89 selection by the environment, but that these differences are reshuffled in each generation because of
90 panmixia. In this context, Mateo et al. (in press) developed a new model, called GenEveel, that was
91 able to mimic all observed patterns assuming the existence of genetic polymorphism in growth rate
92 and adaptive phenotypic plasticity. It suggested that the genetic polymorphism and phenotypic
93 plasticity may have been selected by natural selection as an adaptation to environmental
94 heterogeneity and density dependence. In this context, selection of growth habitats, length-at-
95 silvering and sex determination would be plastic traits that enable individuals to optimize their
96 fitness in a wide range of environments.

97 This issue of adaptation to environmental variability is presently crucial because temperate eels
98 have suffered dramatic collapses (Dekker, 2009; Dekker and Casselman, 2014; Jacoby et al., 2015),
99 and *A. anguilla* is classified as critically endangered by the IUCN (Jacoby and Gollock, 2014a),
100 while *A. rostrata* and *A. japonica* are classified as endangered (Jacoby et al., 2014; Jacoby and
101 Gollock, 2014b). Several factors have been proposed to explain these declines (Jacoby et al., 2015),
102 including changes in oceanic conditions (Castonguay et al., 1994), contamination and habitat
103 degradation (Belpaire et al., 2016; Byer et al., 2015), parasitism (Feunteun, 2002; Kirk, 2003),
104 fishing pressure (Dekker, 2003a), fragmentation including massive habitat loss (Kettle et al., 2011),
105 and hydroelectricity-induced mortality (Castonguay et al., 1994). In view of this situation, the
106 European Commission introduced European Regulation N° 1100/2007, imposing a new set of
107 measures designed to reverse the decline. Since eel management is under the responsibility of
108 member states, each member state was required to implement Eel Management Plans, enforcing
109 management measures to decrease all sources of anthropogenic mortalities. Because of the
110 heterogeneity in anthropogenic pressures, these measures are quite heterogeneous, targeting
111 different types of pressures among countries and regions. The impact of anthropogenic pressures is
112 indeed generally assessed by quantifying the induced mortality rates. Indeed, the European
113 Regulation uses the biomass of escapees as a management target (the Regulation required that
114 management measures should be implemented to ensure that silver eel escapement is at least 40%
115 of the escapement in pristine conditions). However, in the presence of genetic polymorphism and
116 phenotypic plasticity, anthropogenic pressures can have a wide range of effects in terms of life
117 history traits. They can be a selective pressure advantaging fast or slow growers or have
118 consequences in terms of the sex ratio, length-at-silvering or spatial distribution by affecting plastic
119 traits.

120 In view of this, we decided to use GenEveel (Mateo et al., in press) to assess the impact of
121 anthropogenic pressures on silver eel escapement, not only by quantifying the number of escapees,
122 but also by assessing their effect on the sex ratio, the proportions of slow and fast growers, the
123 length-at-silvering and the resulting egg production after a generation in continental waters exposed
124 to different kinds of anthropogenic pressures. We chose to focus on four kinds of anthropogenic
125 pressures: (i) the glass eel fishery, (ii) obstacles to upstream migration, (iii) turbine mortality during
126 downstream migration, and (iv) the silver eel fishery. (i) Glass eel fishing is a widespread activity in
127 France, the UK, Spain, Portugal and Italy (ICES, 2016), in some North American rivers (Cairns et
128 al., 2008; Dutil et al., 2009; Jessop, 1998) and in Asian waters (Tatsukawa, 2003). In the Bay of
129 Biscay, the glass eel fishery used to be the most important fishery in France in terms of turnover
130 (Castelnaud, 2000), and the Bay of Biscay is assumed to receive the main part of the total

131 recruitment (Dekker, 2000). These fisheries harvest young, sexually undetermined individuals
132 entering continental waters. In some river basins, the glass eel fishery can catch nearly all
133 individuals entering the basins, such as in the Vilaine River, where the recruitment rate is less than
134 5% some years (Briand et al., 2005). (ii) Fragmentation by human-induced obstacles can impact
135 upstream migration (Drouineau et al., 2015; Tremblay et al., 2016). By blocking individuals during
136 their upstream migration, they confine them into restricted parts of river basins (Kettle et al., 2011).
137 (iii) During downstream migration, they can impair migration to the sea (Drouineau et al., 2017;
138 Tremblay et al., 2016), especially because of mortality induced by passage through hydropower
139 facilities (Pedersen et al., 2012; Winter et al., 2006). (iv) Finally, silver eel fishing is a widespread
140 activity (Aalto et al., 2016; Amilhat et al., 2008; Bernotas et al., 2016; Verreault et al., 2012;
141 Westerberg and Wickström, 2016) that targets large silvers when they migrate back to sea.

142 We adapted the GenEveel model and developed five output indicators (number of escapees,
143 proportion of females, egg production, length-at-silvering, and proportion of slow growers) to
144 consider the impact of anthropogenic pressures on population dynamics. We carried out a numerical
145 exploration of the model and fitted statistical models to assess the effects of the anthropogenic
146 pressures on the outputs. This enables us to quantify the impact of pressures on the different
147 components and to discuss the implications.

148 2. Materials and methods

149 2.1. *GenEveel*

150 GenEveel is an individual-based model. It postulates that the population is composed of two
151 types of individuals (slow and fast growing) based on a genetic polymorphism and that individuals
152 determine their sex and select their growth habitat and length-at-silvering to optimize their fitness.
153 Assuming that males follow a time-minimizing strategy, their fitness is assumed to be proportional
154 to their survival rate until a constant length-at-silvering is reached. On the other hand, females are
155 assumed to follow a size-maximizing strategy, and their fitness is constrained by a trade-off
156 between fecundity at length-at-silvering and the survival rate to this length, with the length-at-
157 silvering selected to optimize this trade-off. Growth is assumed to be a combination of intrinsic
158 growth rates (slow and fast growers) modulated by environmental effects, with faster growth rates
159 in downstream habitats. Survival is a product of environmental effects and density dependence.

160 The environment, a single river catchment, is represented by 30 contiguous cells (growth
161 habitat) with prespecified effects on individual growth and mortality rates. One by one, individuals
162 determine their sex and select their growth habitat and length-at-silvering based on the combination
163 that maximizes fitness.

164 A complete description of the model can be found in Mateo et al. (in press). We parameterized
165 the model as in the reference simulation in Mateo et al. (in press) for this study, which consisted of
166 the best set of values found in the literature for the European eel. Seventeen parameters were used
167 in the model to characterize the environment (as cells in the river catchment) and individuals
168 (demographic, growth and fitness parameters).

169 2.2. *Impacts of anthropogenic pressures*

170 2.2.1. Glass eel fishery

171 The glass eel fishery was characterized by the catch rate, g , i.e., the proportion of individuals
172 caught by the fishery. The glass eel fishery was assumed to harvest individuals before sex
173 determination and habitat selection. Consequently, if N eels entered the catchment, glass eel catches
174 were assumed to follow a binomial distribution, $\text{binomial}(N, g)$, and the number of survivors was
175 the complement.

176 2.2.2. Obstacles to upstream migration

177 We used two obstacles (0.66 and 0.1, respectively, according to the relative distance along the
178 river and referring to the 20th and 3th cells of the 30 that represent the river catchment) characterized

179 by their blockage rates ($b_{0.66}$ and $b_{0.1}$, respectively) and locations ($po_{0.66}$ and $po_{0.1}$, respectively,
180 which represent cells of the catchments). For each individual, we restricted the possible
181 combinations of growth habitat, length-at-silvering and sex to cells near the river mouth, $po_{0.1}$, with
182 a probability $b_{0.1}$ and to cells near the river source, $po_{0.66}$, with probability $(1-b_{0.1}) \cdot (1-b_{0.66})$.

183 2.2.3. Turbine mortality

184 The two obstacles also had turbine mortality rates ($t_{0.66}$ and $t_{0.1}$, respectively). Each individual
185 located upstream of the obstacle and that survived until length-at-silvering were killed by turbines,
186 with a probability corresponding to the mortality rates.

187 2.2.4. Silver eel fishery

188 The silver eel fishery was assumed to affect eels in a similar way to the mortality due to
189 turbines. We assumed that two successive silver eel fisheries occur, one located near the river source
190 at position $ps_{0.66}$ and one near the river mouth at position $ps_{0.1}$. These fisheries were characterized
191 by catch rates of $s_{0.66}$ and $s_{0.1}$, respectively. Each individual located upstream of the obstacle and
192 that survived until length-at-silvering were caught by silver eel fisheries with a probability
193 corresponding to the catch rates.

194 2.3. Model sequence

195 The glass eel fishery occurred first, as it is the first source of mortality encountered by young
196 eels. Then, GenEveel was used to position individuals in the catchment with restrictions due to
197 obstacles to determine their sex, length-at-silvering and survival probability to length-at-silvering.
198 The survivors that reached maturity in the upper parts of the river migrated downstream and were
199 impacted by the silver eel fishery and the turbine located near the river source. The remaining fish
200 were then impacted by the silver eel fishery and the turbine located near the river mouth.

201 Fig. 1 represents the model sequence. A proportion, g , of recruits was first harvested by the glass
202 eel fishery. Among the $1-g$ individuals that escaped the glass eel fishery, a proportion, $b_{0.1}$, was
203 constrained to settle downstream of the first obstacle, and only $(1-b_{0.1}) \cdot (1-b_{0.66})$ were free to settle
204 in the whole catchment. After sex determination and habitat selection (the combination that
205 maximized the expected fitness), individuals underwent downstream migration and were harvested
206 with a given probability if they passed a silver eel fishery or a turbine.

207 We computed five indicators based on the survivors (escapees):

- 208 1. number of escapees (Ns);
- 209 2. mean length-at-silvering (Ls);
- 210 3. proportion of females (sex ratio) (SR);
- 211 4. egg production (sum of female fecundity among survivors) (E); and
- 212 5. proportion of slow growers (SI)

213 The first indicator is the traditional indicator for quantifying mortality and is generally used as a
214 proxy of the spawning biomass, although not accounting for trait modifications. The next two
215 indicators aim to measure the effect of anthropogenic pressures on two plastic traits. The fourth
216 indicator quantifies the total effects of pressures on egg production and is a more direct proxy than
217 the number of escapees because it combines both induced mortality and trait modifications. The last
218 indicator is used to quantify the selective pressure induced by each anthropogenic pressure.

219 2.4. Numerical exploration of the model and results analysis

220 2.4.1. Experimental design

221 Our objective was to compare the intensity of effects of anthropogenic pressures rather than
222 locations. Therefore, we chose a set of parameters that were similar among pressures. For locations,
223 pressure positions were set at the same cells in the catchment: $ps_{0.66}=po_{0.66}=20$ and $ps_{0.1}=po_{0.1}=3$. To
224 explore the effect of pressure intensity, we used Latin hypercube sampling (LHS) (Iman and
225 Conover, 1980; McKay, 1988; McKay et al., 2000). LHS is classical tool used to carry out global

226 sensitivity analysis of models (Blower and Dowlatabadi, 1994; Helton et al., 2005; Helton and
227 Davis, 2003; Manache and Melching, 2004). It allows an efficient exploration of the input
228 parameter space; estimation of output statistics, such as sensitivity indices; and variance
229 decomposition. It subdivides each parameter into X intervals and then samples those intervals with
230 probability $1/X$ (random sampling within intervals), ensuring that each parameter interval is
231 sampled only once and that two parameters are not sampled in the same interval in the same
232 experiment. By doing this, LHS ensures a random association of parameter values and allows their
233 respective effects to be disentangled.

234 In our LHS, the resulting design matrix is made up of all anthropogenic pressures (g , $b_{0.66}$, $b_{0.1}$,
235 $s_{0.66}$, $s_{0.1}$, $t_{0.66}$, $t_{0.1}$) in columns, defined at an interval of 0 (no pressure) to 1 (impact all individuals).
236 Intervals were then subdivided into 100 subintervals, corresponding to simulations in rows. We built
237 100 independent LHSs to account for model stochasticity, resulting in 10,000 simulations.

238 2.4.2. Results analysis

239 For each output Y , we fitted a generalized additive model on the 10,000 simulations using Y as a
240 dependent variable and the anthropogenic pressures (g , $b_{0.66}$, $b_{0.1}$, $s_{0.66}$, $s_{0.1}$, $t_{0.66}$, $t_{0.1}$) as explanatory
241 variables:

$$242 \quad Y \sim s(g) + s(s_{0.1}) + s(s_{0.66}) + s(b_{0.1}) + s(b_{0.66}) + s(t_{0.1}) + s(t_{0.66})$$

243 with $s()$ as a smoother function.

244 The Gaussian family was used for the number of escapees (Ns), length-at-silvering (Ls) and egg
245 production (E) (a log transformation was used for Ns and E to normalize the distribution), while the
246 binomial family was used for the proportion of females (SR) and slow growers (Sl).

247 Generalized additive models (GAMs) were fitted using the package `mgcv` (Wood, 2011) in R (R
248 Development Core Team, 2011). Plots of marginal effects were used to explore the relations
249 between pressures and outputs. The signs of the regression coefficients indicated whether the
250 pressure has a positive or negative impact on the indicator. The proportion of explained variance
251 was used to compare the strength of the effects of anthropogenic pressures.

252 We computed the partial rank correlation coefficients (PRCC) (Saltelli et al., 2000) to assess the
253 global effects of pressures on each output variable using the package `sensitivity` (Pujol et al., 2016).
254 These coefficients assessed the degree of association between an output variable and a predictor
255 after removing the effects of other predictors.

256 3. Results

257 The PRCC are summarized in Table 1. The glass eel fishery had the most important influence; it
258 had negative effects on the number of escapees and positive effects on the length-at-silvering and
259 the proportion of females. Its influence was more limited on egg production (negative effects) and
260 the proportion of slow growers (positive effects). Regarding egg production, the silver eel fishery
261 and turbine located near the river mouth had negative effects that were similar to that of the glass
262 eel fishery. Unsurprisingly, they also had important negative effects on the proportion of slow
263 growers, the mean length-at-silvering and the number of escapees. Obstacles located near the river
264 mouth had a great influence on all indicators except egg production; they had negative effects on
265 the length-at-silvering and the proportion of females and positive effects on the number of escapees
266 and the proportion of slow growers. The turbine and silver eel fishery located near the river source
267 had lesser influences on the model outputs than the rest of the pressures.

268 The GAM summaries are presented in Table A1. All pressures had a negative impact on the
269 number of escapees (Ns) except for obstacles located near the river mouth ($b_{0.1}$), which had a
270 positive impact (Table 1 - Fig. 2). Indeed, they blocked individuals in a zone with faster growth
271 rates, decreasing the time to length-at-silvering and increasing the survival until escapement.
272 Interestingly, the slope of the glass eel fishery effect was more moderate for small intensities than
273 for strong intensities (Fig. 2). At low intensities, fisheries led to a decrease in density and therefore

274 to lower density-dependent mortality and a higher survival rate of the remaining individuals, which
275 balanced the effect of the fishery.

276 Regarding the proportion of females (Fig. 2 - Table 1), the glass eel fishery had the greatest
277 impact by removing individuals. It drastically decreased the density and consequently the density-
278 dependent mortality, favoring females, especially at high intensity (Fig. 2). The patterns were less
279 obvious for the other pressures, except for obstacles located near the river mouth ($b_{0,1}$); after a given
280 level, an increasing blockage rate increased the density in downstream habitats, increasing the
281 density-dependent mortality and favoring males (Fig. 2).

282 The effects of other pressures (turbines and the silver eel fishery) was not due to density
283 dependence, but rather the distribution of individuals in the river catchment; there was a female-
284 biased sex ratio in the upstream habitat, and females were consequently more impacted by
285 mortalities directed toward downstream migrants.

286 Regarding the length-at-silvering (Table 1 - Fig. 2), the results were very similar to the
287 proportion of females since males have a constant and low length-at-silvering, while females have a
288 variable and larger length-at-silvering. However, the effects of the silver eel fishery and the turbine
289 located near the river mouth ($s_{0,1}$ and $t_{0,1}$, respectively) were much more obvious.

290 Although obstacles located near the river mouth and the glass eel fishery ($b_{0,1}$ and g ,
291 respectively) were the pressures most impacting the three first indicators, the silver eel fishery and
292 the turbine located near the river mouth ($s_{0,1}$ and $t_{0,1}$, respectively) had similar influences on egg
293 production, and they were more important than the glass eel fishery (g) (Table 1). Indeed, the effect
294 of g on the number of escapees was mitigated by the lower competition and the production of
295 females in downstream zones of the catchment, explaining the flat effects observed for low g (Fig.
296 2). For obstacles located near the river mouth ($b_{0,1}$), the increase in the number of escapees
297 explained an increase in egg production when blockage was low. However, for high blockage rates,
298 the increase in escapees (Fig. 2) was outweighed by a higher proportion of males (Fig. 2), leading to
299 a decrease in egg production (Fig. 2). For the silver eel fishery and the turbine located near the river
300 mouth ($s_{0,1}$ and $t_{0,1}$, respectively), their impacts on both the proportion of females and the number of
301 escapees were negative, decreasing the effect on egg production.

302 Finally, the silver eel fishery and the turbine located near the river mouth decreased the
303 proportion of slow growers because those individuals tended to settle in upstream habitats and were
304 consequently more impacted than fast growers (Table 1 - Fig. 2). On the other hand, obstacles
305 located near the river mouth tended to increase the proportion of slow growers; those individuals
306 were blocked in downstream habitats where growth was faster, resulting in a better survival until
307 length-at-silvering (as observed in the number of escapees (N_s), Fig. 2). However, blocked
308 individuals were “constrained” to remain in downstream habitats, resulting in a decrease in their
309 fitness and to strong competition with fast growers. Interestingly, the glass eel fishery, which
310 harvested slow and fast growers equally, tended to favor fast growers at low intensities and slow
311 growers at high intensities (Fig. 2; at high intensities, the very low density-dependent mortality
312 strongly benefited slow growers by removing fast growers, which were dominant competitors).

313 4. Discussion

314 4.1. Assessing the impact of anthropogenic pressures: why consider other indicators than only the 315 number of escapees?

316 The collapse in temperate eel populations required immediate actions (Dekker, 2009) to
317 decrease all sources of anthropogenic pressures. This is the spirit of the European Regulation on the
318 European eel and of subsequent eel management plans in European Union Member States.
319 Logically, the first criteria used to assess the impact of anthropogenic pressures are mortality rates
320 and escapement. However, given the large phenotypic variability observed at the distribution scale
321 (Vélez-Espino and Koops, 2009), likely due to both phenotypic plasticity (Drouineau et al., 2014;
322 Mateo et al., in press) and the consideration of genetic polymorphism (Côté et al., 2015), the

323 number of escapees may not be sufficient, and it is worthwhile to explore the effects of
324 anthropogenic pressures on other components. Thus, the aim of this study was to use the GenEveel
325 model to explore the effects of different types of anthropogenic pressures on the number of escapees
326 as well as other indicators accounting for life history traits and genetic polymorphism.

327 *4.2. Using a theoretical model to explore the possible consequences of anthropogenic pressures*

328 GenEveel is a theoretical model, but is parameterized with the best biological information
329 available in the literature. Regarding the intensity, high levels have been observed for each type of
330 pressure. For example, the exploitation rates of glass eels were found to be very high in some
331 catchments: 13-30% in the Adour River (France) (Prouzet, 2002), 6.2-48.7% in the Oria River
332 (Spain) (Aranburu et al., 2016), 30.8-51.8% in Nova Scotia (Jessop, 2000), 44.1-75% in Shang-Chi
333 (Taiwan) (Tzeng, 1984) and even 78.1-99.7% in the Vilaine River (France) (Briand et al., 2005).
334 The passability for upstream migration was estimated to range between 19.8 and 49.1% for an
335 obstacle in the Canal des Etangs (France) (Drouineau et al., 2015). A similar result was obtained by
336 Briand et al. (2005), who calculated an efficiency of the trapping ladder as 30% in the Vilaine
337 catchment (France). Passability is probably even lower for obstacles not equipped with an eel
338 ladder. Turbine impacts appear to vary considerably depending on study sites. On the Meuse River,
339 Winter et al. (2007) estimated that the two hydropower plants induced an overall mortality of
340 approximately 20-30%. In a Swedish river, Calles et al. (2010) found mortality rates of 40% and
341 60% induced by two plants. At a Polish site, Dębowski et al. (2016) estimated a mortality rate of
342 55%. Mortality due to silver eel fisheries also varies from 22-26% in the Meuse River (Netherlands)
343 (Winter et al., 2006) to 82% mortality in the Gudena River and Randers Fjord (Denmark)
344 (Aarestrup et al., 2008) and can reach even higher levels in some Mediterranean lagoons (Amilhat
345 et al., 2008).

346 Given the large variability in the pressure intensities, and the fact that they vary at the
347 distribution area scale, we used a Latin hypercube sampling design to carry out the global
348 exploration of their effects in a theoretical catchment in GenEveel. Consequently, the exercise is
349 clearly theoretical, and our raw results should not be used to classify the effects of anthropogenic
350 pressures. However, they provide valuable insights on the type of effects induced by the different
351 pressures and indicate that the significance of these effects greatly depends on the considered output
352 indicator.

353 We did not consider the effects of yellow eel fisheries in this exercise for various reasons. First,
354 the GenEveel model did not describe the growth phase precisely, so it is difficult to incorporate such
355 mortalities. Moreover, most of the pressures occur in specific parts of the catchments (the estuary
356 for glass eel fisheries, obstacles one after another) and at a specific times (migration), while yellow
357 eel fisheries are more widespread and occur all along the growth phase, so comparisons of these
358 pressures would be more uncertain.

359 *4.3. The effect of anthropogenic pressures depends on their location within the catchment*

360 Among the most obvious results, the impacts of anthropogenic pressures clearly depend on their
361 location in the river catchment, and pressures located in the very upper parts of the catchment,
362 where densities are low, have limited impacts. Although this result is obvious, its consequence at the
363 population scale is less obvious because anthropogenic activities are not uniformly distributed
364 across the distribution area. For example, in Europe, some hydropower facilities can be found in
365 downstream areas of river catchments, similar to in Scandinavia, while they are located upstream in
366 France. Silver eel fisheries are mainly located in Scandinavia and the Mediterranean Sea (Dekker,
367 2003b), while glass eel fisheries are dominant in France and Spain. Since life history patterns and
368 differences in densities are also observed at the distribution area scale, it is very difficult to draw up
369 an overall picture of the effect of each individual pressure at the population scale.

370 *4.4. Effect of pressures on escapement: existence of compensatory mechanisms*

371 Another important message is that the number of escapees and the resulting egg production are
372 not necessarily correlated. In our simulations, glass eel fisheries and migratory obstacles had the
373 greatest influence on the number of escapees, the sex ratio and the length-at-silvering (although the
374 effects were different depending on pressures). Nevertheless, the silver eel fishery and turbine
375 mortality had similar negative impacts on egg production, which were greater than that of the glass
376 eel fishery and were more significant than those of obstacles. This is explained by compensatory
377 mechanisms that mitigate the negative effects of glass eel fisheries and obstacles to upstream
378 migration, while there are no such mechanisms for those of silver eel fisheries and turbine mortality.
379 For example, the decrease in escapement induced by the glass eel fishery can be compensated by
380 the higher production of females. For obstacles to upstream migration, the decrease in female
381 production is compensated by the better survival of individuals and consequently an increase in
382 escapement. In other words, the pressure inducing the highest direct mortality does not necessarily
383 have the greatest influence on silver eel escapement. Even pressures that do not induce any direct
384 mortality, such as migratory obstacles, can influence some demographic attributes of escapees, such
385 as the proportion of females or length-at-silvering. For example, obstacles close to the river mouth
386 blocked individuals during their upstream migration in downstream habitats. As a consequence,
387 slow growers that usually tend to settle in upstream habitats are forced to settle in downstream
388 habitats with a faster growth rate and higher mortality due to competition. In these conditions, slow
389 growers become males with a time-minimizing strategy, while they would have turned into females
390 with a size-maximizing strategy in upstream habitats; this is why the proportion of females
391 decreased. Since growth is fast in downstream habitats and those individuals mature very early (at
392 male length-at-silvering), they have a higher survival rate until silvering than they would have
393 known in upstream habitats, leading to an increase in the proportion of slow growers. However,
394 they would have had higher fitness as a female in upstream habitats since the loss in survival would
395 have been compensated by high fecundity.

396 Interestingly, by decreasing the number of individuals that enter the river catchment, a
397 simulation with a glass eel fishery “mimics” a simulation with a depleted population. It is
398 interesting to note that in such a situation, a decrease in competition enables individuals to settle in
399 downstream habitats, where growth is faster. More females are also produced because of these
400 lower densities. As a result, egg production is less impacted than the number of escapees. As such,
401 the phenotypic plasticity of eels can be considered as a resilience factor in a depleted population.

402 *4.5. Anthropogenic pressures as selective pressures*

403 The final output of the model is the impact of anthropogenic pressures on slow and fast growers.
404 Most pressures, except for the glass eel fishery, tend to favor fast growers over slow growers, and as
405 such, anthropogenic pressures can be selective pressures. The selective pressure induced by
406 obstacles on elver migration has already been observed at the gene expression level (Podgorniak et
407 al., 2015a, 2015b, 2016). If most anthropogenic activities act as several pressures favoring fast
408 growers, we can wonder how slow growers were preserved by natural selection. A possible
409 explanation may be due to gender differences. Males are predominantly fast growers, while females
410 are both slow and fast growers (Côté et al., 2015; Mateo et al., in press), but slow-growing females
411 have a larger length-at-silvering and greater fecundity than fast-growing females (Mateo et al., in
412 press). In a high-abundance context, we can assume that males are fast growers, females are
413 predominantly slow growers, and the two types are necessary. In a collapsed population,
414 anthropogenic pressures tend to favor males and fast growers (Table 1), and egg production can
415 become limiting. Consequently, the few remaining slow-growing females (which are less impacted
416 by competition and can settle in downstream habitats) can become even more important, explaining
417 their preservation to some extent. To validate this assumption, it would be interesting to carry out a
418 similar simulation exercise at the population scale and with multiple generations to explore the
419 impacts of anthropogenic pressures on the two types of individuals. Moreover, this would allow us

420 to explore the assumption of spatially varying selection proposed by Gagnaire et al. (2012) and Côté
421 et al. (2014) by simulating the arrival of slow and fast growers in random and contrasting river
422 catchments, where they would suffer different mortalities and would consequently be more or less
423 adapted.

424 Selective pressure due to fishing activities on exploited marine stocks has been demonstrated for
425 many fish stocks (Law, 2000; Pinsky and Palumbi, 2014). A major collapse in abundance is required
426 to reduce the overall genetic diversity. Since current recruitment is now less than 5% of historical
427 recruitment and, for the American eel, recruitment has ceased in Lake Ontario and Upper Saint-
428 Lawrence (Casselman, 2003), a loss of genetic diversity is not impossible. This question of diversity
429 at the individual level is crucial for eel conservation since it may contribute to population resilience
430 through a storage and portfolio effect (Secor, 2015).

431 The misunderstanding of the effects of anthropogenic pressures also means that we poorly
432 understand the effects of some mitigating measures. Stocking programs have been widely used
433 (Couillard et al., 2014) and are still widely used by European countries (Brämick et al., 2016; Josset
434 et al., 2016; Simon et al., 2013). However, by modifying densities and moving slow or fast growers
435 into habitats where they are not adapted, such measures can have unexpected effects on escapement.
436 Such observations were made in Canada, where eels were stocked in the Great Lakes, and the
437 escapees produced were more similar to escapees from their original sites than to native silver eels
438 from the Great Lakes (Couillard et al., 2014; Stacey et al., 2015). Exploring this issue with
439 GenEeel by artificially “moving” individuals after their sex and growth habitat have been
440 determined will bring new insights into the ways stocking programs can affect the whole
441 population.

442 4.6. Conclusion

443 Our theoretical exercise provided new insights on the impact of anthropogenic pressures, both direct
444 and indirect. Since this simulation exercise is purely theoretical and because of the spatial
445 heterogeneity in both anthropogenic pressures and eel life history traits, it is impossible to promote
446 any specific management measures, which should be chosen according to the local conditions.
447 However, our results imply that managers and scientists should not only assess the quantity of
448 escapees but also their quality. Quantifying human-induced mortalities and the effects of
449 anthropogenic pressures on the number of escapees is a main priority, given the dramatic situation
450 of the three temperate eel populations. However, further research to investigate the possible impacts
451 of anthropogenic pressures on phenotypic changes in life history traits and consequently the
452 reproductive output of the spawning stock will be required to achieve sustainable management of
453 the species.

454 5. Acknowledgements

455 This study was supported by the Hynes project between Irstea and EDF R&D. We would like to
456 thank Christian Rigaud, Bruno Ernande, Martin Castonguay and Laurent Beaulaton for their
457 participation in fruitful discussions. We acknowledge the two referees for their helpful comments.

458 6. References

- 459 Aalto, E., Capoccioni, F., Mas, J.T., Schiavina, M., Leone, C., Leo, G.D., Ciccotti, E., 2016. Quantifying 60
460 years of declining European eel (*Anguilla anguilla* L., 1758) fishery yields in Mediterranean coastal
461 lagoons. ICES J. Mar. Sci. J. Cons. 73, 101–110. doi:10.1093/icesjms/fsv084
462 Aarestrup, K., Thorstad, E., Koed, A., Jepsen, N., Svendsen, J., Pedersen, M., Skov, C., Økland, F., 2008.
463 Survival and behaviour of European silver eel in late freshwater and early marine phase during
464 spring migration. Fish. Manag. Ecol. 15, 435–440.
465 Als, T.D., Hansen, M.M., Maes, G.E., Castonguay, M., Riemann, L., Aarestrup, K., Munk, P., Sparholt, H.,
466 Hanel, R., Bernatchez, L., 2011. All roads lead to home: panmixia of European eel in the Sargasso
467 Sea. Mol. Ecol. 20, 1333–1346. doi:10.1111/j.1365-294X.2011.05011.x
468

- 469 Amilhat, E., Farrugio, H., Lecomte-Finiger, R., Simon, G., Sasal, P., 2008. Silver eel population size and
470 escapement in a Mediterranean lagoon: Bages-Sigean, France. *Knowl. Manag. Aquat. Ecosyst.* 390–
471 391, 5p1-5p11.
- 472 Arai, T., Chino, N., 2012. Diverse migration strategy between freshwater and seawater habitats in the
473 freshwater eel genus *Anguilla*. *J. Fish Biol.* 81, 442–455.
- 474 Aranburu, A., Diaz, E., Briand, C., 2016. Glass eel recruitment and exploitation in a South European estuary
475 (Oria Bay of Biscay). *ICES J. Mar. Sci.* 73, 111–121. doi:10.1093/icesjms/fsv116
- 476 Belpaire, C., Pujolar, J.M., Geeraerts, C., Maes, G.E., 2016. Contaminants in Eels and their Role in the
477 Collapse of the Eel Stocks. *Biol. Ecol. Anguillid Eels* 225.
- 478 Bernotas, P., Vetemaa, M., Saks, L., Eschbaum, R., Verliin, A., Järvalt, A., 2016. Dynamics of European eel
479 landings and stocks in the coastal waters of Estonia. *ICES J. Mar. Sci. J. Cons.* 73, 84–90.
480 doi:10.1093/icesjms/fsv245
- 481 Blower, S.M., Dowlatabadi, H., 1994. Sensitivity and uncertainty analysis of complex models of disease
482 transmission: an HIV model, as an example. *Int. Stat. Rev. Int. Stat.* 229–243.
- 483 Boivin, B., Castonguay, M., Audet, C., Pavey, S.A., Dionne, M., Bernatchez, L., 2015. How does salinity
484 influence habitat selection and growth in juvenile American eels *Anguilla rostrata*? *J. Fish Biol.* 86,
485 765–784. doi:10.1111/jfb.12604
- 486 Brämick, U., Fladung, E., Simon, J., 2016. Stocking is essential to meet the silver eel escapement target in a
487 river system with currently low natural recruitment. *ICES J. Mar. Sci. J. Cons.* 73, 91–100.
488 doi:10.1093/icesjms/fsv113
- 489 Briand, C., Fatin, D., Feunteun, E., Fontenelle, G., 2005. Estimating the stock of glass eels in an estuary by
490 mark-recapture experiments using vital dyes. *Bull. Fr. Pêche Prot. Milieux Aquat.* 378–379, 23–46.
491 doi:10.1051/kmae:2005002
- 492 Byer, J.D., Lebeuf, M., Trottier, S., Raach, M., Alae, M., Stephen Brown, R., Backus, S., Casselman, J.M.,
493 Hodson, P.V., 2015. Trends of persistent organic pollutants in American eel (*Anguilla rostrata*) from
494 eastern Lake Ontario, Canada, and their potential effects on recruitment. *Sci. Total Environ.* 529,
495 231–242. doi:10.1016/j.scitotenv.2015.05.054
- 496 Cairns, D.K., Secor, D.A., Morrison, W.E., Hallett, J.A., 2009. Salinity-linked growth in anguillid eels and
497 the paradox of temperate-zone catadromy. *J. Fish Biol.* 74, 2094–2114.
- 498 Cairns, D.K., Tremblay, V., Caron, F., Casselman, J.M., Verreault, G., Jessop, B.M., de Lafontaine, Y.,
499 Bradford, R.G., Verdon, R., Dumont, P., Mailhot, Y., Zhu, J., Mathers, A., Oliveira, K., Benhalima,
500 K., Dietrich, J.P., Hallett, J.A., Lagacé, M., 2008. American eel abundance indicators in Canada
501 (Canadian Data Report of Fisheries and Aquatic Sciences 1207 No. Canadian Data Report of
502 Fisheries and Aquatic Sciences 1207). Oceans and Science Branch, Fisheries and Oceans Canada.
- 503 Calles, O., Olsson, I., Comoglio, C., Kemp, P., Blunden, L., Schmitz, M., Greenberg, L., 2010. Size-
504 dependent mortality of migratory silver eels at a hydropower plant, and implications for escapement
505 to the sea. *Freshw. Biol.* 55, 2167–2180.
- 506 Casselman, J.M. (2003) Dynamics of Resources of the American Eel, *Anguilla rostrata*: Declining
507 Abundance in the 1990s. In: *Eel Biology*. (eds K. Aida, K. Tsukamoto and K. Yamauchi). Springer
508 Japan, pp 255–274.
- 509 Castelnaud, G., 2000. Localisation de la pêche, effectifs de pêcheurs et production des espèces amphihalines
510 dans les fleuves français. *Bull. Fr. Pêche Piscic.* 357/358, 439–460.
- 511 Castonguay, M., Hodson, P.V., Couillard, C.M., Eckersley, M.J., Dutil, J.D., Verreault, G., 1994. Why is
512 recruitment of the American eel, *Anguilla rostrata*, declining in the St. Lawrence River and Gulf?
513 *Can. J. Fish. Aquat. Sci.* 51, 479–488.
- 514 Côté, C.L., Castonguay, M., McWilliam, K.S., Gordon, C., Bernatchez, L., 2014. In absence of local
515 adaptation, plasticity and spatially varying selection rule: a view from genomic reaction norms in a
516 panmictic species (*Anguilla rostrata*). *BMC Genomics* 15, 403. doi:10.1186/1471-2164-15-403
- 517 Côté, C.L., Castonguay, M., Verreault, G., Bernatchez, L., 2009. Differential effects of origin and salinity
518 rearing conditions on growth of glass eels of the American eel *Anguilla rostrata*: implications for
519 stocking programmes. *J. Fish Biol.* 74, 1934–1948. doi:10.1111/j.1095-8649.2009.02291.x
- 520 Côté, C.L., Pavey, S.A., Stacey, J.A., Pratt, T.C., Castonguay, M., Audet, C., Bernatchez, L., 2015. Growth,
521 Female Size, and Sex Ratio Variability in American Eel of Different Origins in Both Controlled
522 Conditions and the Wild: Implications for Stocking Programs. *Trans. Am. Fish. Soc.* 144, 246–257.
523 doi:10.1080/00028487.2014.975841

- 524 Couillard, C.M., Verreault, G., Dumont, P., Stanley, D., Threader, R.W., 2014. Assessment of Fat Reserves
525 Adequacy in the First Migrant Silver American Eels of a Large-Scale Stocking Experiment. North
526 Am. J. Fish. Manag. 34, 802–813. doi:10.1080/02755947.2014.920738
- 527 Daverat, F., Beaulaton, L., Poole, R., Lambert, P., Wickstrom, H., Andersson, J., Aprahamian, M., Hizem, B.,
528 Elie, P., Yalcin-Ozdilek, S., Gumus, A., 2012. One century of eel growth: changes and implications.
529 Ecol. Freshw. Fish 21, 325–336.
- 530 Daverat, F., Limburg, K., Thibault, I., Shiao, J.-C., Dodson, J., Caron, F., Tzeng, W.-N., Iizuka, Y.,
531 Wickström, H., 2006. Phenotypic plasticity of habitat use by three temperate eel species, *Anguilla*
532 *anguilla*, *A. japonica* and *A. rostrata*. Mar. Ecol. Prog. Ser. 308, 231–241.
- 533 Davey, A., Jellyman, D., 2005. Sex determination in freshwater eels and management options for
534 manipulation of sex. Rev. Fish Biol. Fish. 15, 37–52.
- 535 Dębowski, P., Bernaś, R., Skóra, M., Morzuch, J., 2016. Mortality of silver eel (*Anguilla anguilla*) migrating
536 downstream through a small hydroelectric plant on the Drawa River in northern Poland. Arch. Pol.
537 Fish. 24, 69–75.
- 538 Dekker, W., 2009. Worldwide decline of eel resources necessitates immediate action. Quebec Declaration of
539 Concern.
- 540 Dekker, W., 2003a. Did lack of spawners cause the collapse of the European eel, *Anguilla anguilla*? Fish.
541 Manag. Ecol. 10, 365–376. doi:10.1111/j.1365-2400.2003.00352.x
- 542 Dekker, W., 2003b. Status of the European eel stock and fisheries, in: Eel Biology. Springer, pp. 237–254.
- 543 Dekker, W., 2000. A Procrustean assessment of the European eel stock. ICES J. Mar. Sci. 57, 938–947.
544 doi:10.1006/jmsc.2000.0581
- 545 Dekker, W., Casselman, J.M., 2014. The 2003 Québec Declaration of Concern About Eel Declines—11 Years
546 Later: Are Eels Climbing Back up the Slippery Slope? Fisheries 39, 613–614.
547 doi:10.1080/03632415.2014.979342
- 548 Drouineau, H., Bau, F., Alric, A., Deligne, N., Gomes, P., Sagnes, P., 2017. Silver eel downstream migration
549 in fragmented rivers: use of a Bayesian model to track movements triggering and duration. Aquat.
550 Living Resour. 30, 1–19. doi: 10.1051/alr/2017003
- 551 Drouineau, H., Rigaud, C., Daverat, F., Lambert, P., 2014. EvEel (evolutionary ecology-based model for eel):
552 a model to explore the role of phenotypic plasticity as an adaptive response of three temperate eels to
553 spatially structured environments. Can. J. Fish. Aquat. Sci. 71, 1561–1571. doi:10.1139/cjfas-2014-
554 0090
- 555 Drouineau, H., Rigaud, C., Laharanne, A., Fabre, R., Alric, A., Baran, P., 2015. Assessing the efficiency of an
556 elver ladder using a multi-state mark-recapture model. River Res. Appl. 31, 291–300.
557 doi:10.1002/rra.2737
- 558 Dutil, J.-D., Dumont, P., Cairns, D.K., Galbraith, P.S., Verreault, G., Castonguay, M., Proulx, S., 2009.
559 *Anguilla rostrata* glass eel migration and recruitment in the estuary and Gulf of St Lawrence. J. Fish
560 Biol. 74, 1970–1984. doi:10.1111/j.1095-8649.2009.02292.x
- 561 Edeline, E., 2007. Adaptive phenotypic plasticity of eel diadromy. Mar. Ecol.-Prog. Ser. 341, 229–232.
- 562 Ernande, B., Dieckmann, U., 2004. The evolution of phenotypic plasticity in spatially structured
563 environments: Implications of intraspecific competition, plasticity costs and environmental
564 characteristics. J Evol Biol 17, 613–628.
- 565 Feunteun, E., 2002. Management and restoration of European eel population (*Anguilla anguilla*): An
566 impossible bargain. Ecol Eng 18, 575–591. doi:10.1016/S0925-8574(02)00021-6
- 567 Gagnaire, P.-A., Normandeau, E., Côté, C., Hansen, M.M., Bernatchez, L., 2012. The Genetic Consequences
568 of Spatially Varying Selection in the Panmictic American Eel (*Anguilla rostrata*). Genetics 190, 725–
569 736. doi:10.1534/genetics.111.134825
- 570 Geffroy, B., 2012. Déterminisme environnemental du sexe chez l’Anguille Européenne *Anguilla anguilla*
571 (Thèse de doctorat - spécialité physiologie et biologie des organismes-populations-interactions).
572 Université de Pau et des pays de l’Adour - École doctorale 211 sciences exactes et leurs applications.
- 573 Geffroy, B., Bardonnnet, A., 2015. Sex differentiation and sex determination in eels: consequences for
574 management. Fish Fish. n/a-n/a. doi:10.1111/faf.12113
- 575 Gotthard, K., Nylin, S., 1995. Adaptive Plasticity and Plasticity as an Adaptation: A Selective Review of
576 Plasticity in Animal Morphology and Life History. Oikos 74, 3–17. doi:10.2307/3545669
- 577 Han, Y., Hung, C., Liao, Y., Tzeng, W., 2010. Population genetic structure of the Japanese eel *Anguilla*
578 *japonica*: panmixia at spatial and temporal scales. Mar. Ecol. Prog. Ser. 401, 221–232.

- 579 doi:10.3354/meps08422
- 580 Helfman, G., Bozeman, E., Brothers, E., 1984. Size, age, and sex of American eels in a Georgia river. *Trans.*
581 *Am. Fish. Soc.* 113, 132–141.
- 582 Helfman, G., Facey, D.E., Stanton Hales Jr., L., Bozeman Jr., E.L., 1987. Reproductive ecology of the
583 American eel. *Am. Fish. Soc. Symp.* 1, 42–56.
- 584 Helton, J.C., Davis, F.J., 2003. Latin hypercube sampling and the propagation of uncertainty in analyses of
585 complex systems. *Reliab. Eng. Syst. Saf.* 81, 23–69.
- 586 Helton, J.C., Davis, F.J., Johnson, J.D., 2005. A comparison of uncertainty and sensitivity analysis results
587 obtained with random and Latin hypercube sampling. *Reliab. Eng. Syst. Saf.* 89, 305–330.
- 588 ICES. 2016. Report of the Working Group on Eels (WGEEL), 15–22 September 2016, Cordoba, Spain. ICES
589 CM 2016/ACOM:19. 107 pp.
- 590 Iman, R.L., Conover, W.J., 1980. Small sample sensitivity analysis techniques for computer models. with an
591 application to risk assessment. *Commun. Stat.-Theory Methods* 9, 1749–1842.
- 592 Jacoby, D., Casselman, J.M., DeLucia, M., Hammerson, G.A., Gollock, M., 2014. *Anguilla rostrata*, in: The
593 IUCN Red List of Threatened Species. Version 2014. <http://www.iucnredlist.org>.
- 594 Jacoby, D., Gollock, M., 2014a. *Anguilla anguilla*, in: The IUCN Red List of Threatened Species. Version
595 2014.2. <http://www.iucnredlist.org>.
- 596 Jacoby, D., Gollock, M., 2014b. *Anguilla japonica*, in: The IUCN Red List of Threatened Species. Version
597 2014.2. <http://www.iucnredlist.org>.
- 598 Jacoby, D.M.P., Casselman, J.M., Crook, V., DeLucia, M.-B., Ahn, H., Kaifu, K., Kurwie, T., Sasal, P.,
599 Silfvergrip, A.M.C., Smith, K.G., Uchida, K., Walker, A.M., Gollock, M.J., 2015. Synergistic
600 patterns of threat and the challenges facing global anguillid eel conservation. *Glob. Ecol. Conserv.* 4,
601 321–333. doi:10.1016/j.gecco.2015.07.009
- 602 Jessop, B.M., 2000. Size, and exploitation rate by dip net fishery, of the run of American eel, *Anguilla*
603 *rostrata* (LeSueur), elvers in the East River, Nova Scotia. *Dana* 12, 43–57.
- 604 Jessop, B.M., 1998. Geographic and seasonal variation in biological characteristics of American eel elvers in
605 the Bay of Fundy area and on the Atlantic coast of Nova Scotia. *Can. J. Zool.* 76, 2172–2185.
606 doi:10.1139/z98-169
- 607 Josset, Q., Trancart, T., Mazel, V., Charrier, F., Frotté, L., Acou, A., Feunteun, E., 2016. Pre-release processes
608 influencing short-term mortality of glass eels in the French eel (*Anguilla anguilla*, Linnaeus 1758)
609 stocking programme. *ICES J. Mar. Sci. J. Cons.* 73, 150–157. doi:10.1093/icesjms/fsv074
- 610 Kettle, A.J., Asbjørn Vøllestad, L., Wibig, J., 2011. Where once the eel and the elephant were together:
611 decline of the European eel because of changing hydrology in southwest Europe and northwest
612 Africa? *Fish Fish.* 12, 380–411. doi:10.1111/j.1467-2979.2010.00400.x
- 613 Kirk, R.S., 2003. The impact of *Anguillicola crassus* on European eels. *Fish. Manag. Ecol.* 10, 385–394.
- 614 Law, R. 2000. Fishing, selection, and phenotypic evolution. *ICES J. Mar. Sci.* 57, 659–668.
615 doi:10.1006/jmsc.2000.0731
- 616 Levins, R., 1963. Theory of fitness in a heterogeneous environment. II. Developmental flexibility and niche
617 selection. *Amercian Nat.* 47, 75–90.
- 618 Manache, G., Melching, C.S., 2004. Sensitivity analysis of a water-quality model using Latin hypercube
619 sampling. *J. Water Resour. Plan. Manag.* 130, 232–242.
- 620 Mateo, M., Lambert, P., Tétard, S., Castonguay, M., Ernande, B., Drouineau, H., in press. Exploring
621 phenotypic plasticity and spatially varying selection for the European eel: an evolutionary ecology
622 based model. *Can. J. Fish. Aquat. Sci.* doi: 10.1139/cjfas-2016-0214
- 623 McCleave, J., 1993. Physical and behavioural controls on the oceanic distribution and migration of
624 leptocephali. *J Fish Biol* 43, 243–273.
- 625 McKay, M.D., 1988. Sensitivity and uncertainty analysis using a statistical sample of input values.
626 *Uncertain. Anal.* 145–186.
- 627 McKay, M.D., Beckman, R.J., Conover, W.J., 2000. A comparison of three methods for selecting values of
628 input variables in the analysis of output from a computer code. *Technometrics* 42, 55–61.
- 629 Melià, P., Bevacqua, D., Crivelli, A.J., De Leo, G.A., Panfili, J., Gatto, M., 2006. Age and growth of *Anguilla*
630 *anguilla* in the Camargue lagoons. *J. Fish Biol.* 68, 876–890. doi:10.1111/j.0022-1112.2006.00975.x
- 631 Oliveira, K., 1999. Life history characteristics and strategies of the American eel, *Anguilla rostrata* 56, 795–
632 802.
- 633 Oliveira, K., McCleave, J., 2000. Variation in population and life history traits of the American eel, *Anguilla*

- 634 *rostrata*, in four rivers in Maine. *Env. Biol Fishes* 59, 141–151.
- 635 Pedersen, M.I., Jepsen, N., Aarestrup, K., Koed, A., Pedersen, S., Okland, F., 2012. Loss of European silver
636 eel passing a hydropower station. *J. Appl. Ichthyol.* 28, 189–193.
- 637 Pigliucci, M., 2005. Evolution of phenotypic plasticity: Where are we going now? *Trends Ecol Evol* 20, 481–
638 486.
- 639 Pinsky, M.L., Palumbi, S.T., 2014. Meta-analysis reveals lower genetic diversity in overfished populations.
640 *Mol. Ecol.* 23, 29–39. doi: 10.1111/mec.12509
- 641 Podgorniak, T., Angelini, A., Blanchet, S., de Oliveira, E., Pierron, F., Daverat, F., 2015. Climbing
642 experience in glass eels: A cognitive task or a matter of physical capacities? *Physiol. Behav.* 151,
643 448–455. doi:10.1016/j.physbeh.2015.08.001
- 644 Podgorniak, T., Blanchet, S., de, O., Daverat, F., Pierron, F., 2016. To boldly climb: Behavioural and
645 cognitive differences in migrating European glass eels. *R. Soc. Open Sci.* 3. doi:10.1098/rsos.150665
- 646 Podgorniak, T., Milan, M., Pujolar, J.M., Maes, G.E., Bargelloni, L., De Oliveira, E., Pierron, F., Daverat, F.,
647 2015. Differences in brain gene transcription profiles advocate for an important role of cognitive
648 function in upstream migration and water obstacles crossing in European eel. *BMC Genomics* 16,
649 378. doi:10.1186/s12864-015-1589-y
- 650 Prouzet, P., 2002. Historique des captures de civelles, intensité actuelle de leur exploitation, variation de leur
651 capturabilité par la pêche professionnelle maritime et indice de colonisation sur le bassin versant de
652 l'Adour (Rapport final, contrat EC/DG FISH (DGXIV) N No. Rapport final, contrat EC/DG FISH
653 (DGXIV) N).
- 654 Pujol, G., Iooss, B., Boumhaout, A.J. with contributions from K., Veiga, S.D., Fruth, J., Gilquin, L.,
655 Guillaume, J., Gratiet, L.L., Lemaitre, P., Ramos, B., Touati, T., Weber, F., 2016. sensitivity: Global
656 Sensitivity Analysis of Model Outputs.
- 657 Pujolar, J.M., 2013. Conclusive evidence for panmixia in the American eel. *Mol. Ecol.* 22, 1761–1762.
658 doi:10.1111/mec.12143
- 659 Pujolar, J.M., Jacobsen, M.W., Als, T.D., Frydenberg, J., Munch, K., Jónsson, B., Jian, J.B., Cheng, L., Maes,
660 G.E., Bernatchez, L., others, 2014. Genome-wide single-generation signatures of local selection in
661 the panmictic European eel. *Mol. Ecol.* 23, 2514–2528.
- 662 R Development Core Team, 2011. R: A Language and Environment for Statistical Computing.
- 663 Saltelli, A., Chan, K., Scott, E.M., others, 2000. Sensitivity analysis. Wiley New York.
- 664 Schmidt, J., 1923. Breeding places and migrations of the eel. *Nature* 111, 51–54.
- 665 Secor, D.H., 2015. American Eel: When Does Diversity Matter? *Fisheries* 40, 462–463.
666 doi:10.1080/03632415.2015.1073152
- 667 Simon, J., Dörner, H., Scott, R.D., Schreckenbach, K., Knösche, R., 2013. Comparison of growth and
668 condition of European eels stocked as glass and farm sourced eels in lakes in the first 4 years after
669 stocking. *J. Appl. Ichthyol.* 29, 323–330. doi:10.1111/jai.12078
- 670 Stacey, J.A., Pratt, T.C., Verreault, G., Fox, M.G., 2015. A caution for conservation stocking as an approach
671 for recovering Atlantic eels. *Aquat. Conserv. Mar. Freshw. Ecosyst.* 25, 569–580.
672 doi:10.1002/aqc.2498
- 673 Tatsukawa, K., 2003. Eel Resources in East Asia, in: Aida, K., Tsukamoto, K., Yamauchi, K. (Eds.), *Eel*
674 *Biology*. Springer Japan, pp. 293–298.
- 675 Tesch, F.W., 2003. *The Eel*. Blackwell Publishing.
- 676 Tremblay, V., Cossette, C., Dutil, J.-D., Verreault, G., Dumont, P., 2016. Assessment of upstream and
677 downstream passability for eel at dams. *ICES J. Mar. Sci. J. Cons.* 73, 22–32.
678 doi:10.1093/icesjms/fsv106
- 679 Tsukamoto, K., 1992. Discovery of the spawning area for Japanese eel. *Nature* 356, 789–791.
680 doi:10.1038/356789a0
- 681 Tsukamoto, K., Nakai, I., Tesch, W.-V., 1998. Do all freshwater eels migrate? [3]. *Nature* 396, 635–636.
- 682 Tzeng, W.-N., 1984. An estimate of the exploitation rate of *Anguilla japonica* elvers immigrating into the
683 coastal waters off Shuang-Chi River, Taiwan. *Bull. Inst. Zool. Acad. Sin.* 23, 173–180.
- 684 Ulrik, M.G., Pujolar, J.M., Ferchaud, A.-L., Jacobsen, M.W., Als, T.D., Gagnaire, P.A., Frydenberg, J.,
685 Bøcher, P.K., Jónsson, B., Bernatchez, L., Hansen, M.M., 2014. Do North Atlantic eels show parallel
686 patterns of spatially varying selection? *BMC Evol. Biol.* 14, 138. doi:10.1186/1471-2148-14-138
- 687 Van Den Thillart, G., Palstra, A., Van Ginneken, V., 2007. Simulated migration of European silver eel; swim
688 capacity and cost of transport. *J. Mar. Sci. Technol.* 15, 1–16.

- 689 Vélez-Espino, L.A., Koops, M.A., 2009. A synthesis of the ecological processes influencing variation in life
690 history and movement patterns of American eel: towards a global assessment. *Rev. Fish Biol. Fish.*
691 20, 163–186. doi:10.1007/s11160-009-9127-0
- 692 Verreault, G., Mingelbier, M., Dumont, P., 2012. Spawning migration of American eel *Anguilla rostrata* from
693 pristine (1843-1872) to contemporary (1963-1990) periods in the St Lawrence Estuary, Canada. *J.*
694 *Fish Biol.* 81, 387–407.
- 695 Vollestad, L.A., 1992. Geographic variation in age and length at metamorphosis of maturing European eel -
696 Environmental effects and phenotypic plasticity 61, 41–48.
- 697 Westerberg, H., Wickström, H., 2016. Stock assessment of eels in the Baltic: reconciling survey estimates to
698 achieve quantitative analysis. *ICES J. Mar. Sci. J. Cons.* 73, 75–83. doi:10.1093/icesjms/fsv049
- 699 White, J.W., Rassweiler, A., Samhuri, J.F., Stier, A.C., White, C., 2014. Ecologists should not use statistical
700 significance tests to interpret simulation model results. *Oikos* 123, 385–388. doi:10.1111/j.1600-
701 0706.2013.01073.x
- 702 Winter, H., Jansen, H., Breukelaar, A., 2007. Silver eel mortality during downstream migration in the River
703 Meuse, from a population perspective. *ICES J. Mar. Sci.* 64, 1444–1449.
- 704 Winter, H., Jansen, H., Bruijs, M., 2006. Assessing the impact of hydropower and fisheries on downstream
705 migrating silver eel, *Anguilla anguilla*, by telemetry in the River Meuse. *Ecol. Freshw. Fish* 15, 221–
706 228.
- 707 Wood, S.N., 2011. Fast stable restricted maximum likelihood and marginal likelihood estimation of
708 semiparametric generalized linear models. *J. R. Stat. Soc. Ser. B Stat. Methodol.* 73, 3–36.

709 **Tables**


710 **Table 1.** Partial rank correlation coefficients between output variables (columns) and pressures (rows). A
711 positive coefficient indicates a positive correlation. *g*: glass eel fishery, *s*_{0.66}: silver eel fishery located near
712 the river source, *s*_{0.1}: silver eel fishery located near the river mouth, *t*_{0.66}: turbine located near the river source,
713 *t*_{0.1}: turbine located near the river mouth, *b*_{0.66}: obstacle to upstream migration, *b*_{0.1}: obstacle to downstream
714 migration.

	<i>Ns</i>	<i>Ls</i>	<i>SR</i>	<i>E</i>	<i>SI</i>
<i>g</i>	-0.95	0.94	0.95	-0.49	0.22
<i>b</i> _{0.1}	0.77	-0.75	-0.57	-0.01	0.88
<i>b</i> _{0.66}	0.02	0	0.01	0.01	0.02
<i>s</i> _{0.1}	-0.39	-0.33	-0.14	-0.42	-0.35
<i>s</i> _{0.66}	-0.02	0	0.01	-0.01	-0.03
<i>t</i> _{0.1}	-0.39	-0.33	-0.12	-0.4	-0.35
<i>t</i> _{0.66}	-0.01	-0.02	-0.01	-0.01	-0.01

715


716

717 **Figures**



718

719 **Figure 1.** Conceptual representation of the catchment, indicating the seven anthropogenic pressures
720 considered in the study: 1) g : catch rate by the glass eel fishery, 2) $b_{0.66}$ and 3) $b_{0.1}$: blockage rates (located
721 near the river source and the river mouth, respectively), 4) $t_{0.66}$ and 6) $t_{0.1}$: turbine mortality rates (located
722 near the river source and the river mouth, respectively), 5) $s_{0.66}$ and 7) $s_{0.1}$: catch rates by the silver eel
723 fisheries (located near the river source and the river mouth, respectively).
724



725

726 **Figure 2.** Effects of the pressures on the five indicators estimated by the GAMs. g : glass eel fishery, $s_{0.66}$ and
 727 $s_{0.1}$: silver eel fisheries located near the river source and the river mouth, respectively, $t_{0.66}$ and $t_{0.1}$: turbines
 728 located near the river source and the river mouth, respectively, $b_{0.66}$ and $b_{0.1}$: obstacles located near the river
 729 source and the river mouth, respectively.

730

731 **Supplementary data**

732 **Table A1.** Summary of the GAM model. edf corresponds to the estimated degrees of freedom for each model
 733 parameter. F corresponds to the Fisher statistics when the Gaussian family is used and Chis.sq to the χ^2
 734 statistics when the binomial family is used. P-values are not presented (although all are smaller than 0.05)
 735 because significance tests are not recommended when analyzing simulation model results (White et al.,
 736 2014).

	<i>Ns</i>		<i>Ls</i>		<i>SR</i>		<i>E</i>		<i>Sl</i>	
	edf	F	edf	F	edf	Chi.sq	edf	F	edf	Chi.sq
<i>s(g)</i>	9	18517.8	8.6	14036.4	9.0	3849436.9	6.6	1257.1	9.0	23175.6
<i>s(s_{0.1})</i>	5.0	266.2	3.3	451.8	8.8	56266.4	1.0	3906.1	8.7	52527.1
<i>s(s_{0.66})</i>	1.0	7.7	2.2	1.6	8.8	485.7	6.8	2.1	8.1	592.6
<i>s(b_{0.1})</i>	3.4	3066.9	5.2	3311.3	9.0	2598861.4	4.0	194.8	8.7	1001753.5
<i>s(b_{0.66})</i>	1.0	5.3	1.0	0.3	8.7	515.9	1.0	4.3	8.9	783.7
<i>s(t_{0.1})</i>	2.5	545.4	2.9	497.5	8.9	56169.5	1.0	3727.6	8.9	54605.7
<i>s(t_{0.66})</i>	1.0	1.3	2.0	2.0	8.9	541.1	2.5	3.3	8.0	357.9

737