



**HAL**  
open science

## Principal Process Analysis of dynamic GlucoCEST MRI data

Stefano Casagrande, Marco Pizzolato, Francisco Torrealdea, Xavier Golay,  
Timothé Boutelier

► **To cite this version:**

Stefano Casagrande, Marco Pizzolato, Francisco Torrealdea, Xavier Golay, Timothé Boutelier. Principal Process Analysis of dynamic GlucoCEST MRI data. ISMRM 2018, Jun 2018, Paris, France. hal-01729755

**HAL Id: hal-01729755**

**<https://hal.science/hal-01729755>**

Submitted on 12 Mar 2018

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**Synopsis.** GlucoCEST is an MRI contrast enhancement technique sensitive to the concentration of sugar in the tissue. Because of a difference in metabolism, it is thought that tumors consume more sugar than normal tissue. However, glucose metabolism is complex and depends on many processes, which are all important to understand the origin of the measured signal. To achieve this goal we apply here a process analysis method to a deterministic system describing the metabolism of glucose in the tissue.

**Introduction.** Chemical Exchange Saturation Transfer (CEST) is an MRI contrast enhancement technique that enables the indirect detection of molecules with exchangeable protons [1]. GlucoCEST is a CEST technique that measures a signal related to the concentration of injected glucose and its derivatives [1]. It is expected that metabolic anomalies due to the presence of cancerous tissue could be measured or characterized by means of glucoCEST, and in particular to its dynamic characteristics. There the signal is analyzed as a function of time following glucose metabolism, which is paramount to pathological tissue assessment. Different subvoxel compartments contribute to the signal. Particularly, a simplified model made as a system composed by 19 Ordinary Differential Equations (ODEs) can be used in a first step, where different parameters can be set to simulate healthy or tumor-like conditions [2]. The system accounts for the presence of vascular glucose, its equilibrium within the interstitial environment, and the transport into the intracellular space. The purpose of this work is to identify the processes that contribute more to the signal. This analysis allows designing a reduced model, dedicated to the fitting of experimental data, and quantification of physiological parameters. To do so we apply Principal Process Analysis (PPA) [3, 4], a numerical method for the analysis and reduction of biological systems designed with ODE formalism. PPA allows associating a dynamic weight to all involved processes and inferring their importance during the acquisition time.

**Methods.** We consider each ODE of the system as a sum of biological processes. Let

$$\dot{x}_i = \sum_j f_{i,j}(x, p) \quad (1)$$

where  $x = (x_1, x_2, \dots, x_n) \in \mathbb{R}^n$  is the vector of concentrations of glucose in different compartments and  $p \in \mathbb{R}^b$  is the vector of parameters.  $f_{i,j}(x, p)$  is the  $j^{\text{th}}$  process involved in the dynamical evolution of the  $i^{\text{th}}$  variable of the system over the acquisition time  $[0, T]$ . A relative weight  $W_{i,j}(t, p)$  is associated to each process  $f_{i,j}(x(t), p)$ , to study its influence in the evolution of the variable  $x_i$  over time  $t$ :

$$W_{i,j}(t, p) = \frac{|f_{i,j}(x(t), p)|}{\sum_j |f_{i,j}(x(t), p)|} \quad (2)$$

where  $0 \leq W_{i,j}(t, p) \leq 1$  and  $\sum_j W_{i,j}(t, p) = 1$ . We use the thresholds  $\delta$  and  $\nu$  to detect not only the *inactive* processes but also to make a further distinction between processes with a *moderate activity* and a *fully activity*. We call a process  $f_{i,j}(x(t), p)$  *inactive* at time  $t$  when  $W_{i,j}(t, p) < \delta$ , *moderately active* at time  $t$  when  $\delta \leq W_{i,j}(t, p) < \nu$ , *fully active* at time  $t$  when  $W_{i,j}(t, p) \geq \nu$ . Furthermore, because the glucoCEST signal  $y$  is seen as a sum of the glucose concentration in different compartments (described by six variables of the model in [2]) we also compute the contribution of each variable  $x_i(t)$  for the system output  $y(t) = \sum_i x_i(t)$ :

$$M_i(t) = \frac{x_i(t)}{\sum_i x_i(t)} \quad (3)$$

where  $0 \leq M_i(t) \leq 1$  and  $\sum_i M_i(t) = 1$ . Their *activity* is dictated in the same manner by the thresholds  $\delta$  and  $\nu$ , set at 0.1 and 0.4. The outcome of the analysis is summarized with an extended *Boolean Process Map* [3] called *3-Level Process Map* with three possible *activity* outcomes.

**Results.** Fig. 1 shows the two different CEST signals simulated in healthy and cancerous tissue after an IV bolus, using the parameters described in [2]. The two *3-Level Process Maps* in Fig. 2 reports the *activity* of the system variables that contribute to the CEST signal and of their relative processes, in both healthy and cancerous tissue.

**Discussion.** The map in Fig. 2A shows that in a healthy tissue the highest contribution in the generation of CEST signal is given by the glucose in the interstitial compartment as seen during an IP bolus situation [5]. Fig. 2B shows that in the tumor case the interstitial contribution becomes *inactive* because the glucose carriers bring the glucose faster into the cell. For the first minutes after the injection the vascular compartment is *fully active* and then it becomes *moderately active* because most of the glucose bolus has passed through the tissue. From our theoretical developments, the intracellular compartment seems to be dominated by glucose and fructose-1,6-biphosphate within healthy tissue (Fig. 2A), while glucose-6-phosphate and fructose-6-phosphate dominates in cancerous tissue (Fig. 2B).

**Conclusion.** We have presented the PPA of a glucose metabolism model, in the context of dynamic glucoCEST imaging. It shows that dominant processes are different depending on the properties of the tissue. It would be interesting to apply the technique for different kind of tumors and glucose administration. Based on this analysis we can build different reduced models where the processes that remain *inactive* have been removed.

**Acknowledgments.** This project has received funding from the European Union’s Horizon 2020 research and innovation programme under grant agreement No 667510 and the Department of Health’s NIHR-funded Biomedical Research Centre at University College London.

*Reference to the original publication.* In ISMRM 27th Annual Meeting. Proc. Int. Soc. Mag. Reson. Med. 27 (2018).

## References

- [1] Kogan, F., Hariharan, H., & Reddy, R. (2013). Chemical exchange saturation transfer (CEST) imaging: description of technique and potential clinical applications. *Current radiology reports*, 1(2), 102-114.
- [2] Torrealdea, F. (2016). Investigation of brain tumour metabolism using naturally occurring chemical exchange saturation transfer agents with magnetic resonance imaging (Doctoral dissertation, UCL (University College London)).
- [3] Casagranda, S., Ropers, D., & Gouzé, J. L. (2015, June). Model reduction and process analysis of biological models. In 2015 23rd Mediterranean Conference on Control and Automation (MED), IEEE, 1132-1139.
- [4] Casagranda, S., & Gouzé, J. L. (2017). Principal Process Analysis and reduction of biological models with order of magnitude. *IFAC-PapersOnLine*, 50(1), 12661-12666.

<sup>1</sup>Olea Medical, La Ciotat, France

<sup>2</sup>EPFL, Lausanne, Switzerland

<sup>3</sup>Centre for Medical Imaging, UCL, London, United Kingdom

<sup>4</sup>Institute of Neurology, UCL, London, United Kingdom

- [5] Walker-Samuel, S., Ramasawmy, R., Torrealdea, F., Rega, M., Rajkumar, V., Johnson, S. P., Richardson, S., Goncalves, M., Parkes, H. G., Arstad, E., Thomas, D. L., Pedley, B., Lynthgoe, M.F., & Golay X. (2013). In vivo imaging of glucose uptake and metabolism in tumors. *Nature medicine*, 19(8), 1067-1072.




Figure 1: CEST signal in a healthy tissue (blue color) and cancerous tissue (red color) during an acquisition time of 150 minutes.




Figure 2: *Activity* of six model variables of glucoCEST model and of their processes during a 150-minute period in an healthy (A) and cancerous tissue (B). Processes are listed in the first column (white background) ordered by variable (white background, in bold). Their *activity* is depicted in the second column between 0 and 150 minutes. A horizontal dark blue (black) bar is displayed when the variable (process) is *fully active*, blue (grey) when it is *moderately active* and light blue (white) when it is *inactive*.