



HAL
open science

Auxiliary information : the raking-ratio empirical process

Mickael Albertus, Philippe Berthet

► **To cite this version:**

Mickael Albertus, Philippe Berthet. Auxiliary information : the raking-ratio empirical process. 2018. hal-01736276v1

HAL Id: hal-01736276

<https://hal.science/hal-01736276v1>

Preprint submitted on 16 Mar 2018 (v1), last revised 12 Dec 2018 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Auxiliary information : the raking-ratio empirical process.

Mickael Albertus*and Philippe Berthet†

March 16, 2018

Institut de Mathématiques de Toulouse ; UMR5219
Université de Toulouse ; CNRS

Abstract

We study the empirical measure associated to a sample of size n and modified by N iterations of the raking-ratio method. The empirical measure is adjusted to match the true probability of sets in a finite partition which changes each step. We establish asymptotic properties of the raking-ratio empirical process indexed by functions as $n \rightarrow +\infty$, for N fixed. A closed-form expression of the limiting covariance matrices is derived as $N \rightarrow +\infty$. The nonasymptotic Gaussian approximation we use also yields uniform Berry-Esseen type bounds in n, N and sharp estimates of the uniform quadratic risk reduction. In the two-way contingency table formulas characterizing the limiting process are very simple.

Contents

1	Introduction	2
1.1	The raking-ratio method	2
1.2	An auxiliary information viewpoint	5
1.3	Information from N finite partitions	6
1.4	The raking-ratio empirical and Gaussian processes	8
2	Main results	9
2.1	General properties	9
2.2	Limiting variance and risk reduction	12
2.3	The two margins case	13

*mickael.albertus@math.univ-toulouse.fr

†philippe.berthet@math.univ-toulouse.fr

3	Proofs of general results	16
3.1	Raking formulas	16
3.2	Proof of Proposition 1.	17
3.3	Proof of Proposition 2.	18
3.4	Proof of Proposition 3	19
3.5	Proof of Proposition 4 and Theorem 2.1.	21
3.6	Proof of Proposition 5	26
3.7	Proof of Proposition 6.	29
4	Proofs concerning the limiting process	30
4.1	Proof of Proposition 7	30
4.2	Proof of Propositions 8 and 9	34
4.3	Proof of Proposition 10	35
4.4	Proof of Proposition 11	36
4.5	Proof of Theorem 2.2	39
.1	Elementary example	40
.2	Counterexample of Remark J	41
.3	Raked empirical means over a class	42

1 Introduction

1.1 The raking-ratio method

In survey analysis, statistics, economics and computer sciences the raking-ratio iterative procedure aims to exploit the knowledge of several marginals of a discrete multivariate distribution. Despite many papers from the methodological and algorithmic viewpoint, and chapters in classical textbooks for statisticians, economists or engineers, no probabilistic study is available when the entries of the algorithm are random, or when the discrete measure is empirical.

A natural method. It has been re-invented many times in various fields, and was probably used long time ago. Depending on the setting it may be viewed as stratification, calibration, iterating proportional fitting or matrix scaling. Fitting after sampling is also used to deal with missing data. We initially called it auxiliary information of partitions as we re-invented it as a generic special case of the general nonparametric partial information problem stated later.

Representative sample. In a survey analysis context, the raking-ratio method modifies weights of a contingency table built from a sample of size n in order to fit exactly given marginals. Such a strict margin correction is justified when a few properties of the finite population under study are known, like the size of subpopulations. The modified sample frequencies then reflect the marginal structure of the whole population. If the population is large or infinite the information may come from previous and independent statistical inference, from structural properties of the model or from various experts.

Remark A. Making the sample representative of the population is an ad hoc approach based on common sense. The mathematical impact is twofold. On

the one hand all statistics are affected by the new weights in terms of bias, variance and limit law so that practitioners may very well be using estimators, tests or confident bands that have lost their usual properties. On the other hand, replacing marginal frequencies with the true ones may smooth sample fluctuations of statistics correlated to them while leaving the uncorrelated ones rather unaffected. We would like to quantify these statements more precisely.

The algorithm. A sample is drawn from a population for which $k \geq 2$ marginal finite discrete laws are explicitly known. Initially, each data point has a weight $1/n$. The *ratio* part of the algorithm consists in computing new weights in such a way that the modified joint law has the currently desired marginal. The *raking* part of the method consists in iterating the correction according to another known marginal law, changing again all the weights of the previous step. The k margin constraints are usually treated in a periodic order, only one being fulfilled at the same time. The raking-ratio method stops after N iterations with the implicit hope that the previous constraints are still almost satisfied. As a matter of fact, the k margins converge to the desired ones as $N \rightarrow +\infty$. See Section .1 for a numerical example in a two-way contingency table, $k = 2$.

The limit. This algorithm was called *iterative proportions* by Deming and Stephan [8] who first introduced it. They claimed that if the frequencies of a multiway contingency table are raked periodically as $N \rightarrow +\infty$ they converge to the frequencies minimizing the chi-square distance to the initial frequencies, under the margin constraints. It is wrong, and Stephan [19] modified the algorithm to achieve this minimization. Lewis [14] and Brown [6] studied the case of Bernoulli marginals from the Shannon entropy minimization viewpoint. When $k = 2$ Sinkhorn [17, 18] proved that a unique doubly stochastic matrix can be obtained from each positive square matrix A by alternately normalizing the rows and columns of A , which shows that the algorithm converges in this special case. Finally Ireland and Kullback [12] generalized previous arguments to rigorously justify that the raking-ratio converges to the unique projection of the empirical measure in Shannon-Kullback-Leibler relative entropy on the set of discrete measures satisfying the constraints. From a numerical viewpoint, the rate of convergence of the algorithm is geometric, see Franklin and Lorentz [11].

Remark B. When the sample size $n \rightarrow +\infty$ it is equivalent to minimize chi-square, likelihood or discrimination information contrasts since they share the same first order expansion and the empirical frequencies are uniformly close to the true ones. However, none of these minimizers is explicit due to nonlinearity of sums of ratios showing up. This is why iterative proportions are very useful in practice. They are converging, intuitive and each step is easily computed.

Non explicit bias and variance. The initial values being empirical frequencies the converged solution of the algorithm still deviates from the true probability measure, and moreover in a rather complicated way. The modified empirical distribution satisfying only the marginal constraint of the current iteration, there is a permanent bias with respect to other margins. The exact covariance matrix and bias vector of the random weights after N iterations are tedious to compute. For instance, estimates for the variance of cell probabilities

in the case of a two-way contingency table are given by Brackstone and Rao [5] for $N \leq 4$, Konijn [13] or Choudhry and Lee [7] for $N = 2$. Bankier [2] propounded a recursive linearization technique providing an estimator of the asymptotic variance. In Binder and Th  berge [4] the variance of the converged solution requires to calculate weights at each iteration.

Remark C. Exact computations lead to intractable bias and variance formulas. We would like to identify leading terms when n is large compared to N and provide comprehensive explicit formulas as $n \rightarrow +\infty$, then $N \rightarrow +\infty$.

Quadratic risk reduction. Modifying marginals frequencies of a small sample may have serious drawbacks. Typically, a statistic has less variance but more bias when sample weights are changed by using raking, calibration or stratification methods after sampling. In the spirit of Remark A, a risk reduction is expected whenever the statistic of interest is well correlated to the k known discrete marginal variables. Exact computation of moments being ineffective, evaluating the quadratic risk of a specific statistic requires bias expansions and asymptotics for correlations of weights. No global risk reduction property has been established, and no multivariate or uniform central limit theorem as $n \rightarrow +\infty$.

Remark D. To control simultaneously large classes of statistics one needs to work at the empirical measure level. We feel that a general study of empirical measures modified through auxiliary information is lacking to further analyze and unify calibration type methods. In particular, a weak or strong invariance principle for the raked empirical process could provide joint limiting laws and uniform quadratic risk evaluations, directly for additive statistics like empirical means as well as indirectly for non linear estimators like empirical minimizers.

The small population case. An asymptotic approach is no more relevant in survey analysis when the underlying population is rather small. In this case, the way the sample is drawn has a so deep impact that it may even become the main topic. A study of calibration methods for finite population can be found in Deville and Sarndal [9, 10]. This is beyond the scope of our work.

Goals. In this paper we consider classes of empirical means raked N times, sampled jointly from any population. We look for closed-form expressions of their Gaussian limits and limiting covariances as $n \rightarrow +\infty$ then $N \rightarrow +\infty$. We also intend to quantify the uniform risk reduction phenomenon and provide sharp statistical estimation tools such as uniform Berry-Esseen type bounds.

Organization of the paper. In Section 1.2 we relate the raking-ratio problem to nonparametric auxiliary information. The raking-ratio empirical process $\alpha_n^{(N)}$ is defined at Section 1.3 then usual assumptions on a class \mathcal{F} of additive statistics are given at Section 1.4. In Section 2.1 we state results for $\alpha_n^{(N)}(\mathcal{F})$ when N is fixed. A nonasymptotic strong approximation bound yields the uniform central limit theorem with rate as $n \rightarrow +\infty$, as well as a uniform control of the bias and the covariances for fixed n . The approximating Gaussian process is studied at Section 2.2, which establishes the uniform risk reduction phenomenon provided the iterations are stopped properly. At Section 2.3, in the two partitions case

we characterize explicitly the limiting process as $N \rightarrow +\infty$. All statements are proved in Sections 3 and 4. The Appendix provides a few examples.

1.2 An auxiliary information viewpoint

Let X_1, \dots, X_n be independent random variables with unknown law P on some measurable space $(\mathcal{X}, \mathcal{A})$. Let $\mathbb{P}_n = n^{-1} \sum_{i=1}^n \delta_{X_i}$ be the standard empirical measure on $\mathcal{P}(\mathcal{X})$ hence on \mathcal{A} . Our interest for the raking-ratio method came while investigating how to exploit various kinds of partial information on P to make \mathbb{P}_n closer to P . Usually such an information is hidden in assumptions and mixed with technical requirements, sometimes it is clearly isolated out.

Motivation. A parametric model provides a tremendous amount of information by specifying $P = P_\theta$ up to a finite dimensional parameter θ , so that \mathbb{P}_n can be replaced with the most likely $P_{\theta_n(X_1, \dots, X_n)}$ among the model. Clearly \mathbb{P}_n is used to minimize the empirical likelihood, but the resulting $P_{\theta_n(X_1, \dots, X_n)}$ is of a very different nature, far from the initial Dirac measures δ_{X_i} . On the opposite, in a purely nonparametric setting the information mainly comes from the sample itself, so that only slight modifications of \mathbb{P}_n are induced by weak hypotheses on P – like support, regularity, symmetry, logconcavity, Bayesian model, semi-parametric model, etc. In between, we would like to formalize a notion of auxiliary information on P based on partial but concrete clues to be combined with \mathbb{P}_n . Clues may come from experts, models or statistical learning. The simplest generic situation one can start with is when the probabilities $P(A_j)$ of a finite number of sets $A_j \in \mathcal{A}$ are known – which in a parametric setting already determines θ then P . Interestingly, the nonparametric redesign $\mathbb{P}_n^{(1)}$ of \mathbb{P}_n follows easily if the A_j form a finite partition of \mathcal{X} whereas it is not obvious how to optimally combine information from overlapping sets, and hence several partitions. It turns out that the raking-ratio method is a natural way.

Information from one partition. Let $A_1^{(1)}, \dots, A_{m_1}^{(1)} \subset \mathcal{A}$ be a partition of \mathcal{X} such that $P(\mathcal{A}^{(1)}) = (P(A_1^{(1)}), \dots, P(A_{m_1}^{(1)}))$ is known. The random measure

$$\mathbb{P}_n^{(1)} = \frac{1}{n} \sum_{j=1}^{m_1} \frac{P(A_j^{(1)})}{\mathbb{P}_n(A_j^{(1)})} \sum_{X_i \in A_j^{(1)}} \delta_{X_i} \quad (1.1)$$

is called the *strong belief* empirical measure in a companion work in which $P(\mathcal{A}^{(1)})$ is itself estimated at a fast rate and the relative cost of the two types of data is used to balance a risk. In (1.1) $P(\mathcal{A}^{(1)})$ is known exactly thus $\mathbb{P}_n^{(1)}$ is the minimum entropy projection of \mathbb{P}_n on this set of constraints. We establish below, in the special case $N = 1$, that $\mathbb{P}_n^{(1)}$ indexed by functions satisfies classical properties like Donsker theorem, deviation bounds and strong approximation. The main fact is that the limiting Gaussian process of $\sqrt{n}(\mathbb{P}_n^{(1)} - P)$ has a smaller variance than the P -Brownian bridge, despite the bias induced by the random ratios in (1.1).

Extension to N partitions. The next step is to combine several sources of information on P of this type. Unfortunately there is no explicit modification of \mathbb{P}_n matching simultaneously several finite discrete marginals while minimizing the relative entropy to \mathbb{P}_n . We will study elsewhere a centered process that is no more a discrete measure but fulfils the margin constraints. An alternative consists in recursively updating the current modification $\mathbb{P}_n^{(N-1)}$ of \mathbb{P}_n onto $\mathbb{P}_n^{(N)}$ according to the next known marginal $P(\mathcal{A}^{(N)}) = (P(A_1^{(N)}), \dots, P(A_{m_N}^{(N)}))$ exactly as in (1.1) for $\mathbb{P}_n^{(1)}$ from $\mathbb{P}_n = \mathbb{P}_n^{(0)}$. This coincides with the Deming and Stephan's iterative procedure for a multiple-way contingency table.

1.3 Information from N finite partitions

The raking-ratio empirical measure. For all $N \in \mathbb{N}_*$ let $m_N \geq 2$ and $\mathcal{A}^{(N)} = \{A_1^{(N)}, \dots, A_{m_N}^{(N)}\} \subset \mathcal{A}$ be a partition of \mathcal{X} for which we are given the auxiliary information $P(\mathcal{A}^{(N)}) = (P(A_1^{(N)}), \dots, P(A_{m_N}^{(N)}))$ to be exploited. Assume that

$$\delta_N = \min_{1 \leq j \leq m_N} P(A_j^{(N)}) > 0, \quad N \in \mathbb{N}_*, \quad (1.2)$$

and $\mathcal{A}^{(N_1)} \neq \mathcal{A}^{(N_2)}$ if $|N_1 - N_2| = 1$, otherwise $\mathcal{A}^{(N_1)} = \mathcal{A}^{(N_2)}$ is allowed. For $N = 0$ the information is empty, $m_0 = 1$, $\mathcal{A}^{(0)} = \{\mathcal{X}\}$, $P(\mathcal{A}^{(0)}) = \{1\}$ and $\delta_0 = 1$. For any measurable real function f write $\mathbb{P}_n^{(0)}(f) = \mathbb{P}_n(f) = n^{-1} \sum_{i=1}^n f(X_i)$, $P(f) = \int_{\mathcal{X}} f dP$ and $\alpha_n^{(0)}(f) = \sqrt{n}(\mathbb{P}_n^{(0)}(f) - P(f))$. In (1.1) $\mathbb{P}_n^{(1)}$ allocates the random weight $P(A_j^{(1)})/n\mathbb{P}_n(A_j^{(1)})$ to each $X_i \in A_j^{(1)}$. Hence

$$\begin{aligned} \mathbb{P}_n^{(1)}(f) &= \sum_{i=1}^n \mathbb{P}_n^{(1)}(\{X_i\}) f(X_i) = \sum_{j=1}^{m_1} \sum_{X_i \in A_j^{(1)}} \frac{P(A_j^{(1)})}{n\mathbb{P}_n(A_j^{(1)})} f(X_i) \\ &= \sum_{j=1}^{m_1} \frac{P(A_j^{(1)})}{\mathbb{P}_n(A_j^{(1)})} \left(\frac{1}{n} \sum_{X_i \in A_j^{(1)}} f(X_i) \right) = \sum_{j=1}^{m_1} \frac{P(A_j^{(1)})}{\mathbb{P}_n^{(0)}(A_j^{(1)})} \mathbb{P}_n^{(0)}(f 1_{A_j^{(1)}}). \end{aligned}$$

Let define recursively, for $N \in \mathbb{N}_*$, the N -th raking-ratio empirical measure

$$\mathbb{P}_n^{(N)}(f) = \sum_{j=1}^{m_N} \frac{P(A_j^{(N)})}{\mathbb{P}_n^{(N-1)}(A_j^{(N)})} \mathbb{P}_n^{(N-1)}(f 1_{A_j^{(N)}}) \quad (1.3)$$

and the N -th raking-ratio empirical process

$$\alpha_n^{(N)}(f) = \sqrt{n}(\mathbb{P}_n^{(N)}(f) - P(f)). \quad (1.4)$$

For $A \in \mathcal{A}$ we also write $\alpha_n^{(N)}(A) = \alpha_n^{(N)}(1_A)$. By (1.3) we have for all $N \in \mathbb{N}_*$

$$\mathbb{P}_n^{(N)}(A_j^{(N)}) = P(A_j^{(N)}), \quad \alpha_n^{(N)}(A_j^{(N)}) = 0, \quad 1 \leq j \leq m_N, \quad (1.5)$$

as desired. Both weights and support of the discrete probability measure $\mathbb{P}_n^{(N)}$ are random since (1.3) also reads

$$\mathbb{P}_n^{(N)}(\{X_i\}) = \mathbb{P}_n^{(N-1)}(\{X_i\}) \frac{P(A_j^{(N)})}{\mathbb{P}_n^{(N-1)}(A_j^{(N)})}, \quad \text{for } X_i \in A_j^{(N)}.$$

A few more formulas concerning $\alpha_n^{(N)}$ and $\mathbb{P}_n^{(N)}$ are derived in Section 3.1. The main question is whether or not transforming $\mathbb{P}_n^{(0)}$ iteratively per (1.3) could be misleading in terms of estimation risk as well as Donsker type convergence.

Iterated Kullback projections. The formula (1.3) means that the raking-ratio method is an iterated maximum likelihood procedure. The random discrete measures $\mathbb{P}_n^{(1)}, \dots, \mathbb{P}_n^{(N)}$ are well defined provided that

$$\min_{1 \leq k \leq N} \min_{1 \leq j \leq m_k} \mathbb{P}_n(A_j^{(k)}) > 0 \quad (1.6)$$

which almost surely holds for all n large enough and N fixed, by (1.2) and the law of large numbers. In this case, the N -th iteration $\mathbb{P}_n^{(N)}$ is the Shannon-Kullback-Leibler projection of $\mathbb{P}_n^{(N-1)}$ under the constraint $P(\mathcal{A}^{(N)})$. Given probability measures Q_n and Q supported by $\{X_1, \dots, X_n\}$ write

$$d_K(Q_n \parallel Q) = \sum_{i=1}^n Q_n(\{X_i\}) \log \left(\frac{Q_n(\{X_i\})}{Q(\{X_i\})} \right).$$

Proposition 1. *If (1.6) holds then*

$$\mathbb{P}_n^{(N)} = \arg \min \left\{ d_K(\mathbb{P}_n^{(N-1)} \parallel Q) : Q(\mathcal{A}^{(N)}) = P(\mathcal{A}^{(N)}), \text{ supp}(Q) = \{X_1, \dots, X_n\} \right\}.$$

A mixture of conditional empirical processes. By introducing, for $A \in \mathcal{A}$ such that $P(A) > 0$ and $\mathbb{P}_n^{(N)}(A) > 0$, the conditional expectations

$$\mathbb{E}_n^{(N)}(f|A) = \frac{\mathbb{P}_n^{(N)}(f1_A)}{\mathbb{P}_n^{(N)}(A)}, \quad \mathbb{E}(f|A) = \frac{P(f1_A)}{P(A)}, \quad (1.7)$$

we see that (1.3) further reads

$$\mathbb{P}_n^{(N)}(f) = \sum_{j=1}^{m_N} P(A_j^{(N)}) \mathbb{E}_n^{(N-1)}(f|A_j^{(N)}), \quad P(f) = \sum_{j=1}^{m_N} P(A_j^{(N)}) \mathbb{E}(f|A_j^{(N)}).$$

Therefore (1.4) can also be formulated into

$$\begin{aligned} \alpha_n^{(N)}(f) &= \sum_{j=1}^{m_N} P(A_j^{(N)}) \alpha_{n,j}^{(N-1)}(f), \\ \alpha_{n,j}^{(N-1)}(f) &= \sqrt{n} \left(\mathbb{E}_n^{(N-1)}(f|A_j^{(N)}) - \mathbb{E}(f|A_j^{(N)}) \right). \end{aligned} \quad (1.8)$$

Each $\alpha_{n,j}^{(N-1)}$ is the conditional empirical process of $\mathbb{P}_n^{(N-1)}$ on a set $A_j^{(N)}$ of the new partition $\mathcal{A}^{(N)}$. Their mixture with weights $P(\mathcal{A}^{(N)})$ is $\alpha_n^{(N)}$. In view of (1.7) and (1.8) we have to study the consequences of (1.5) on $\mathbb{P}_n^{(N-1)}(f1_{A_j^{(N)}})$ and $\mathbb{E}_n^{(N-1)}(f|A_j^{(N)})$ as $n \rightarrow +\infty$, for $f \neq 1_{A_j^{(N)}}$.

Bias problem. The processes $\alpha_{n,j}^{(N-1)}$ are not centered due to the factors $1/\mathbb{P}_n^{(N-1)}(A_j^{(N)})$ in (1.3) and (1.7). In general we have

$$\mathbb{E} \left(\mathbb{E}_n^{(N-1)}(f|A) - \mathbb{E}(f|A) \right) = \mathbb{E} \left(\mathbb{P}_n^{(N-1)}(f1_A) \left(\frac{1}{\mathbb{P}_n^{(N-1)}(A)} - \frac{1}{P(A)} \right) \right) \neq 0$$

except for $(A, f) = (A_j^{(N-1)}, 1_{A_j^{(N-1)}})$ hence $\alpha_n^{(N)}$ is no more centered if $N \geq 1$. This unavoidable bias is induced by (1.5) to globally compensate for the local cancellation of the variance of $\mathbb{P}_n^{(N)}(A_j^{(N)})$. The bias tends to spread through (1.3) since the auxiliary information $P(\mathcal{A}^{(N)})$ is applied to the biased $\mathbb{P}_n^{(N-1)}$ instead of the unbiased \mathbb{P}_n . However, our results show that the bias process

$$\mathbb{E} \left(\alpha_n^{(N)}(f) \right) = \sqrt{n} \left(\mathbb{E}(\mathbb{P}_n^{(N)}(f)) - Pf \right)$$

vanishes faster than $1/\sqrt{n}$ uniformly over reasonable classes of functions.

Variance problem. In view of (1.5) and Proposition 1 one may expect that $\mathbb{P}_n^{(N)}(f)$ has less variance than $\mathbb{P}_n(f)$ at least whenever f is sufficiently well approximated by the step functions 1_A for $A \in \mathcal{A}^{(k)}$, $k \leq N$. Our results confirm that $\mathbb{V}(\alpha_n^{(N)}(f)) \leq \mathbb{V}(\alpha_n^{(0)}(f))$ for n large and that finite dimensional covariance matrices decrease faster than $1/n$ uniformly over many f . The quadratic risk may still increase, $\mathbb{E}((\mathbb{P}_n^{(N)}(f) - P(f))^2) > \mathbb{E}((\mathbb{P}_n^{(0)}(f) - P(f))^2)$ because of the bias process. Our main result shows that it is asymptotically not the case.

1.4 The raking-ratio empirical and Gaussian processes

Let \mathcal{M} denote the set of measurable real valued functions on $(\mathcal{X}, \mathcal{A})$. Consider a class $\mathcal{F} \subset \mathcal{M}$ such that $\sup_{f \in \mathcal{F}} |f| \leq M < +\infty$ and satisfying the pointwise measurability condition often used to avoid measurability problems. Namely, $\lim_{m \rightarrow +\infty} f_m(x) = f(x)$ for all $x \in \mathcal{X}$ and $f \in \mathcal{F}$ where $\{f_m\} \subset \mathcal{F}_*$ depends on f and $\mathcal{F}_* \subset \mathcal{F}$ is countable. With no loss of generality also assume that

$$1_A f \in \mathcal{F}, \quad A \in \mathcal{A}_\cup^{(N)} = \mathcal{A}^{(1)} \cup \dots \cup \mathcal{A}^{(N)}, \quad f \in \mathcal{F}. \quad (1.9)$$

In addition \mathcal{F} is assumed to have a small uniform entropy, like VC-classes, or a small P -bracketing entropy, such as many classes of smooth functions. For a probability measure Q on $(\mathcal{X}, \mathcal{A})$ and $f, g \in \mathcal{M}$ define $d_Q^2(f, g) = \int_{\mathcal{X}} (f - g)^2 dQ$. Let $N(\mathcal{F}, \varepsilon, d_Q)$ be the minimum number of balls having d_Q -radius ε needed to cover \mathcal{F} . Let $N_{[\cdot]}(\mathcal{F}, \varepsilon, d_P)$ be the least number of ε -brackets necessary to cover \mathcal{F} , of the form $[g_-, g_+] = \{f : g_- \leq f \leq g_+\}$ with $d_P(g_-, g_+) < \varepsilon$.

Hypothesis (VC). For $c_0 > 0$, $\nu_0 > 0$ it holds $\sup_Q N(\mathcal{F}, \varepsilon, d_Q) \leq c_0/\varepsilon^{\nu_0}$ where the supremum is taken over all discrete probability measures Q on $(\mathcal{X}, \mathcal{A})$.

Hypothesis (BR). For $b_0 > 0$, $r_0 \in (0, 1)$ it holds $N_{[\cdot]}(\mathcal{F}, \varepsilon, d_P) \leq \exp(b_0^2/\varepsilon^{2r_0})$.

Only constants c_0 and b_0 change if one extends an initial \mathcal{F} to satisfy (1.9). Uniform boundedness is the less crucial assumption and could be replaced by a moment condition allowing truncation arguments, adding technicalities.

Let $\ell^\infty(\mathcal{F})$ denote the set of real-valued functions bounded on \mathcal{F} , endowed with the supremum norm $\|\cdot\|_{\mathcal{F}}$. The raking-ratio empirical process $\alpha_n^{(N)}$ defined at (1.4) is now denoted $\alpha_n^{(N)}(\mathcal{F}) = \{\alpha_n^{(N)}(f) : f \in \mathcal{F}\}$. Under (VC) or (BR) \mathcal{F} is a P -Donsker class. Thus $\alpha_n^{(0)}(\mathcal{F})$ converges weakly in $\ell^\infty(\mathcal{F})$ to the P -Brownian bridge indexed by \mathcal{F} , that we denote $\mathbb{G}(\mathcal{F}) = \{\mathbb{G}(f) : f \in \mathcal{F}\}$. Remind (1.7) and that $\mathbb{G}(\mathcal{F})$ is a Gaussian linear process such that, for any $f, g \in \mathcal{F}$,

$$\mathbb{E}(\mathbb{G}(f)) = 0, \quad \text{Cov}(\mathbb{G}(f), \mathbb{G}(g)) = P(fg) - P(f)P(g). \quad (1.10)$$

As for $\alpha_n^{(N)}$ we write $\mathbb{G}^{(0)}(\mathcal{F}) = \mathbb{G}(\mathcal{F})$ and, for short, $\mathbb{G}(A) = \mathbb{G}(1_A)$ if $A \in \mathcal{A}$. The N -th raking-ratio P -Brownian bridge $\mathbb{G}^{(N)}(\mathcal{F})$ is the centered Gaussian process indexed by \mathcal{F} defined recursively, for any $N \in \mathbb{N}_*$ and $f \in \mathcal{F}$, by

$$\mathbb{G}^{(N)}(f) = \mathbb{G}^{(N-1)}(f) - \sum_{j=1}^{m_N} \mathbb{E}(f|A_j^{(N)})\mathbb{G}^{(N-1)}(A_j^{(N)}). \quad (1.11)$$

Lastly, the following notation will be useful,

$$\sigma_f^2 = \mathbb{V}(f(X)) = P(f^2) - P(f)^2, \quad \sigma_{\mathcal{F}}^2 = \sup_{f \in \mathcal{F}} \sigma_f^2, \quad (1.12)$$

where $\sigma_f^2 = \mathbb{V}(\alpha_n^{(0)}(f)) = \mathbb{V}(\mathbb{G}^{(0)}(f))$.

2 Main results

2.1 General properties

We now state asymptotic and nonasymptotic properties that always hold after raking N_0 times. The i.i.d. sequence $\{X_n\}$ is defined on a probability space $(\Omega, \mathcal{T}, \mathbb{P})$ so that \mathbb{P} implicitly leads all convergences when $n \rightarrow +\infty$ and $(\mathcal{X}, \mathcal{A})$ is endowed with $P = \mathbb{P}^{X_1}$. For all $N \leq N_0$ the information $P(\mathcal{A}^{(N)})$ satisfies (1.2). Most of the subsequent constants can be bounded by using only N_0 and

$$\delta_{(N_0)} = \min_{0 \leq N \leq N_0} \delta_N = \min_{0 \leq N \leq N_0} \min_{1 \leq j \leq m_N} P(A_j^{(N)}) > 0. \quad (2.1)$$

Write $L(x) = \log(\max(e, x))$ and define $\kappa_{N_0} = \prod_{N=1}^{N_0} (1 + Mm_N)$.

Proposition 2. *If \mathcal{F} satisfies (VC) or (BR) then for all $N_0 \in \mathbb{N}$ it holds*

$$\limsup_{n \rightarrow +\infty} \frac{1}{\sqrt{2L \circ L(n)}} \sup_{0 \leq N \leq N_0} \left\| \alpha_n^{(N)} \right\|_{\mathcal{F}} \leq \kappa_{N_0} \sigma_{\mathcal{F}} \quad a.s.$$

Remark E. The limiting constant $\kappa_{N_0} \leq (1 + M/\delta_{(N_0)})^{N_0}$ is large, and possibly rough, except for $N_0 = 0$ where $\kappa_0 = 1$ coincides with the classical law of the iterated logarithm – from which the proposition follows.

The nonasymptotic deviation bound for $\alpha_n^{(N)}(\mathcal{F})$ also has the usual magnitude. However, since the partition changes at each step the constants are severely penalized by factors similar to κ_{N_0} above, involving

$$\Delta_{N_0} = \prod_{N=1}^{N_0} \delta_N, \quad M_{N_0} = \prod_{N=1}^{N_0} m_N, \quad S_{N_0} = \sum_{N=1}^{N_0} m_N. \quad (2.2)$$

Proposition 3. *If \mathcal{F} is pointwise measurable, bounded by M then for $N_0 \in \mathbb{N}$ and $n \in \mathbb{N}_*$ we have*

$$\mathbb{P} \left(\sup_{0 \leq N \leq N_0} \left\| \alpha_n^{(N)} \right\|_{\mathcal{F}} \geq \lambda \right) \leq 2^{N_0} M_{N_0} \mathbb{P} \left(\left\| \alpha_n^{(0)} \right\|_{\mathcal{F}} \geq \frac{\lambda \Delta_{N_0}}{(1 + M + \lambda/\sqrt{n})^{N_0}} \right) + S_{N_0} (1 - \delta_{(N_0)})^n.$$

Under (BR) it holds, for $n > n_0$ and $\lambda_0 < \lambda < D_0 \sqrt{n}$,

$$\mathbb{P} \left(\sup_{0 \leq N \leq N_0} \left\| \alpha_n^{(N)} \right\|_{\mathcal{F}} \geq \lambda \right) \leq D_1 \exp(-D_2 \lambda^2) + S_{N_0} (1 - \delta_{(N_0)})^n$$

where the constants D_0, D_1, D_2 are defined at (3.8) and n_0, λ_0 just before. Under (VC) it holds, for $n > n_0$ and $\lambda_0 < \lambda < 2M \sqrt{n}$,

$$\mathbb{P} \left(\sup_{0 \leq N \leq N_0} \left\| \alpha_n^{(N)} \right\|_{\mathcal{F}} \geq \lambda \right) \leq D_3 \lambda^{v_0} \exp(-D_4 \lambda^2) + S_{N_0} (1 - \delta_{(N_0)})^n$$

where the constants D_3, D_4 are defined at (3.9) and n_0, λ_0 just before.

Remark F. Clearly, to avoid drawbacks N_0 should be fixed as n increases, and \mathcal{F} limited to the most crucial variables $f(X)$. In this case, Proposition 3 shows that $\|\alpha_n^{(N_0)}\|_{\mathcal{F}}$ is of order $C \sqrt{\log n}$ with polynomial probability and $C > 0$ not too large. We also deduce that $\mathbb{E}(\|\alpha_n^{(N_0)}\|_{\mathcal{F}}) < D$ with an explicit $D < +\infty$. Concentration of measure type bounds are more difficult to handle due to the mixture (1.8) of processes $\alpha_{n,j}^{(N-1)}$ involving unbounded random coefficients.

The raking-ratio empirical process $\alpha_n^{(N)}(\mathcal{F})$ converges weakly in $\ell^\infty(\mathcal{F})$ to the N -th raking-ratio P -Brownian bridge $\mathbb{G}^{(N)}(\mathcal{F})$ defined at (1.11) and studied at Section 2.2. The \mathbb{R}^{N_0} -valued version is as follows.

Proposition 4. *If \mathcal{F} satisfies (VC) or (BR) then for all $N_0 \in \mathbb{N}$, as $n \rightarrow +\infty$ the sequence $(\alpha_n^{(0)}(\mathcal{F}), \dots, \alpha_n^{(N_0)}(\mathcal{F}))$ converges weakly to $(\mathbb{G}^{(0)}(\mathcal{F}), \dots, \mathbb{G}^{(N_0)}(\mathcal{F}))$ on $\ell_\infty(\mathcal{F} \rightarrow \mathbb{R}^{N_0})$.*

By using Berthet and Mason [3] we obtain the following upper bound v_n for the speed of weak convergence in Lévy-Prokhorov distance d_{LP} and the speed of almost sure convergence in $\|\cdot\|_{\mathcal{F}}$ distance. Powers given by Berthet and Mason are $\alpha = 1/(2 + 5\nu_0)$, $\beta = (4 + 5\nu_0)/(4 + 10\nu_0)$ and $\gamma = (1 - r_0)/2r_0$. They can be slightly improved and conditions (VC) and (BR) generalized.

Theorem 2.1. Let $\theta_0 > 0$. If \mathcal{F} satisfies (VC) then write $v_n = (\log n)^\beta/n^\alpha$. If \mathcal{F} satisfies (BR) then write $v_n = 1/(\log n)^\gamma$. In both cases, one can define on the same probability space $(\Omega, \mathcal{T}, \mathbb{P})$ a sequence $\{X_n\}$ of independent random variables with law P and a sequence $\{\mathbb{G}_n\}$ of versions of \mathbb{G} satisfying the following property. For any $N_0 \geq 0$ there exists $n_0 \in \mathbb{N}$ and $d_0 > 0$ depending on $N_0, M, M_{N_0}, S_{N_0}, \Delta_{N_0}, \delta_{(N_0)}, \theta_0, \nu_0, c_0, r_0, b_0$ such that we have, for all $n \geq n_0$,

$$\mathbb{P} \left(\sup_{0 \leq N \leq N_0} \left\| \alpha_n^{(N)} - \mathbb{G}_n^{(N)} \right\|_{\mathcal{F}} \geq d_0 v_n \right) < \frac{1}{n^{\theta_0}}$$

where $\mathbb{G}_n^{(N)}$ is the version of $\mathbb{G}^{(N)}$ derived from $\mathbb{G}_n^{(0)} = \mathbb{G}_n$ through (1.11).

Remark G. Applied with $\theta_0 > 1$, Theorem 2.1 makes the study of weak convergence of functions of $\alpha_n^{(N)}(\mathcal{F})$ easier by substituting $\mathbb{G}_n^{(N)}$ to $\alpha_n^{(N)}$ through

$$\limsup_{n \rightarrow +\infty} \frac{1}{v_n} \sup_{0 \leq N \leq N_0} \left\| \alpha_n^{(N)} - \mathbb{G}_n^{(N)} \right\|_{\mathcal{F}} \leq d_0 < +\infty \quad a.s.$$

then exploiting the properties induced by (1.11) as in Section 2.2. For instance the finite dimensional laws of $\mathbb{G}^{(N)}$ are computed explicitly at Proposition 7. For nonasymptotic applications, given a class \mathcal{F} of interest it is possible to compute crude bounds for n_0 and d_0 since most constants are left explicit in the proofs. From Theorem 2.1 we derive estimates of the rate of uniform convergence for the bias and the variance. The covariance of $\mathbb{G}^{(N)}$ is given at Section 2.1.

Proposition 5. If \mathcal{F} satisfies (VC) or (BR) then for $N_0 \in \mathbb{N}$ it holds

$$\limsup_{n \rightarrow +\infty} \frac{\sqrt{n}}{v_n} \max_{0 \leq N \leq N_0} \sup_{f \in \mathcal{F}} \left| \mathbb{E} \left(\mathbb{P}_n^{(N)}(f) \right) - P(f) \right| \leq d_0$$

where $v_n \rightarrow 0$ and d_0 are as in Theorem 2.1, and

$$\begin{aligned} & \limsup_{n \rightarrow +\infty} \frac{n}{v_n} \sup_{f, g \in \mathcal{F}} \left| \mathbb{E} \left((\mathbb{P}_n^{(N)}(f) - P(f))(\mathbb{P}_n^{(N)}(g) - P(g)) \right) - \frac{1}{n} \text{Cov} \left(\mathbb{G}^{(N)}(f), \mathbb{G}^{(N)}(g) \right) \right| \\ &= \limsup_{n \rightarrow +\infty} \frac{n}{v_n} \sup_{f, g \in \mathcal{F}} \left| \text{Cov} \left(\mathbb{P}_n^{(N)}(f), \mathbb{P}_n^{(N)}(g) \right) - \frac{1}{n} \text{Cov} \left(\mathbb{G}^{(N)}(f), \mathbb{G}^{(N)}(g) \right) \right| \leq \sqrt{\frac{8}{\pi}} d_0 \sigma_{\mathcal{F}}. \end{aligned}$$

A second consequence of Theorem 2.1 is uniform Berry-Esseen type bounds. Let Φ denote the distribution function of the centered standardized normal law.

Proposition 6. Assume that \mathcal{F} satisfies (VC) or (BR), fix $N_0 \in \mathbb{N}$ and let $d_0 > 0$, $v_n \rightarrow 0$ be defined as above. If $\mathcal{F}_0 \subset \mathcal{F}$ is such that

$$\sigma_0^2 = \inf \left\{ \mathbb{V} \left(\mathbb{G}^{(N)}(f) \right) : f \in \mathcal{F}_0, 0 \leq N \leq N_0 \right\} > 0$$

then for any $d_1 > d_0$ there exists $n_1 \in \mathbb{N}$ such that for all $n \geq n_1$,

$$\max_{0 \leq N \leq N_0} \sup_{f \in \mathcal{F}_0} \sup_{x \in \mathbb{R}} \left| \mathbb{P} \left(\frac{\sqrt{n} \mathbb{P}_n^{(N)}(f) - P(f)}{\sqrt{\mathbb{V} \left(\mathbb{G}^{(N)}(f) \right)}} \leq x \right) - \Phi(x) \right| \leq \frac{d_1}{\sqrt{2\pi} \sigma_0} v_n.$$

Let \mathcal{L} be a collection of real valued Lipschitz functions φ defined on $\ell_\infty(\mathcal{F})$ with Lipschitz constant bounded by $C_1 < +\infty$ and such that $\varphi(\mathbb{G}^{(N)})$ has a density bounded by $C_2 < +\infty$ for all $0 \leq N \leq N_0$ and $\varphi \in \mathcal{L}$. Then for all $n \geq n_1$,

$$\max_{0 \leq N \leq N_0} \sup_{\varphi \in \mathcal{L}} \sup_{x \in \mathbb{R}} \left| \mathbb{P} \left(\varphi(\alpha_n^{(N)}) \leq x \right) - \mathbb{P} \left(\varphi(\mathbb{G}^{(N)}) \leq x \right) \right| \leq d_1 C_1 C_2 v_n.$$

Remark H. The first statement is a special case of the second and also reads

$$\max_{0 \leq N \leq N_0} \sup_{f \in \mathcal{F}_0} \sup_{x \in \mathbb{R}} \left| \mathbb{P} \left(\alpha_n^{(N)}(f) \leq x \right) - \mathbb{P} \left(\mathbb{G}^{(N)}(f) \leq x \right) \right| \leq \frac{d_1}{\sqrt{2\pi}\sigma_0} v_n.$$

The functions $f \in \mathcal{F}$ overdetermined by the knowledge of $P(\mathcal{A}^{(N)})$ have a small $\mathbb{V}(\mathbb{G}^{(N)}(f))$ and are excluded from \mathcal{F}_0 . Proposition 6 is especially useful under (VC) since v_n is polynomially decreasing, thus allowing larger $C_1 C_2$ and \mathcal{L} . An example is given at Section .3.

2.2 Limiting variance and risk reduction

In this section we study the covariance structure of $\mathbb{G}^{(N)}(\mathcal{F})$ from (1.11), for N fixed. The following matrix notation is introduced to shorten formulas. The brackets $[\cdot]$ refer to column vectors built from the partition $\mathcal{A}^{(k)}$ appearing inside. Let V^t denote the transpose of a vector V . For $k \leq N$ write

$$\mathbb{E} [f | \mathcal{A}^{(k)}] = \left(\mathbb{E}(f | A_1^{(k)}), \dots, \mathbb{E}(f | A_{m_k}^{(k)}) \right)^t, \quad \mathbb{G} [\mathcal{A}^{(k)}] = \left(\mathbb{G}(A_1^{(k)}), \dots, \mathbb{G}(A_{m_k}^{(k)}) \right)^t$$

and, for $l \leq k \leq N$ define the matrix $\mathbf{P}_{\mathcal{A}^{(k)} | \mathcal{A}^{(l)}}$ to be

$$\left(\mathbf{P}_{\mathcal{A}^{(k)} | \mathcal{A}^{(l)}} \right)_{i,j} = P(A_j^{(k)} | A_i^{(l)}) = \frac{P(A_j^{(k)} \cap A_i^{(l)})}{P(A_i^{(l)})}, \quad 1 \leq i \leq m_l, 1 \leq j \leq m_k.$$

Write Id_k the identity matrix $k \times k$. Remind that $\mathbb{V}(\mathbb{G}(f)) = P(f^2) - (P(f))^2$, $P(\mathcal{A}^{(k)})^t = P[\mathcal{A}^{(k)}]$ and $P(A_i^{(k)} \cap A_j^{(k)}) = 0$ if $i \neq j$. The covariance matrix of the Gaussian vector $\mathbb{G}[\mathcal{A}^{(k)}]$ is $\mathbb{V}(\mathbb{G}[\mathcal{A}^{(k)}]) = \text{diag}(P(\mathcal{A}^{(k)})) - P(\mathcal{A}^{(k)})^t P(\mathcal{A}^{(k)})$. Let \cdot denote a product between a square matrix and a vector. Finally define

$$\begin{aligned} \Phi_k^{(N)}(f) &= \mathbb{E} [f | \mathcal{A}^{(k)}] + \\ &\sum_{\substack{1 \leq L \leq N-k \\ k < l_1 < l_2 < \dots < l_L \leq N}} (-1)^L \mathbf{P}_{\mathcal{A}^{(l_1)} | \mathcal{A}^{(k)}} \mathbf{P}_{\mathcal{A}^{(l_2)} | \mathcal{A}^{(l_1)}} \cdots \mathbf{P}_{\mathcal{A}^{(l_L)} | \mathcal{A}^{(l_{L-1})}} \cdot \mathbb{E} [f | \mathcal{A}^{(l_L)}]. \end{aligned} \tag{2.3}$$

The closed form expression for the covariance function of $\mathbb{G}^{(N)}(\mathcal{F})$ is as follows.

Proposition 7. *For all $N \in \mathbb{N}$ the process $\mathbb{G}^{(N)}(\mathcal{F})$ is Gaussian, centered and linear with covariance function defined to be, for $(f, g) \in \mathcal{F}^2$,*

$$\text{Cov} \left(\mathbb{G}^{(N)}(f), \mathbb{G}^{(N)}(g) \right) = \text{Cov} (\mathbb{G}(f), \mathbb{G}(g)) - \sum_{k=1}^N \Phi_k^{(N)}(f)^t \cdot \mathbb{V} \left(\mathbb{G}[\mathcal{A}^{(k)}] \right) \cdot \Phi_k^{(N)}(g)$$

Proposition 7 implies the following variance reduction phenomenon.

Proposition 8. *For any $\{f_1, \dots, f_m\} \subset \mathcal{F}$ and $N \in \mathbb{N}$ the covariance matrices $\Sigma_m^{(N)} = \mathbb{V}((\mathbb{G}^{(N)}(f_1), \dots, \mathbb{G}^{(N)}(f_k)))$ are such that $\Sigma_m^{(0)} - \Sigma_m^{(N)}$ is positive definite.*

Remark I. In particular we have $\mathbb{V}(\mathbb{G}^{(N)}(f)) \leq \mathbb{V}(\mathbb{G}^{(0)}(f)) = \sigma_f^2$, $f \in \mathcal{F}$. The asymptotic risk reduction after raking is quantified by combining Propositions 5 and 8. Given $\varepsilon_0 > 0$ and $0 < \sigma_0 < \sigma_{\mathcal{F}}$ there exists some $n_0 = n_0(\varepsilon_0, \mathcal{F})$ such that if $n > n_0$ then any $f \in \mathcal{F}$ with initial quadratic risk $\sigma_f^2/n > \sigma_0/n$ has a new risk, after raking N times, equal to

$$\mathbb{E} \left((\mathbb{P}_n^{(N)}(f) - P(f))^2 \right) = \frac{\sigma_f^2}{n} (\Delta(f) + e(f)v_n)$$

where $v_n \rightarrow 0$ and d_0 are as in Theorem 2.1 and

$$\begin{aligned} \Delta(f) &= \frac{\mathbb{V}(\mathbb{G}^{(N)}(f))}{\sigma_f^2} \in [0, 1] \\ \sup_{f \in \mathcal{F}, \sigma_f \geq \sigma_0} |e(f)| &< (1 + \varepsilon_0) \sqrt{\frac{8}{\pi}} d_0 \frac{\sigma_{\mathcal{F}}}{\sigma_0}. \\ \mathbb{V} \left(\mathbb{G}^{(N)}(f) \right) &= \sigma_f^2 - \sum_{k=1}^N \Phi_k^{(N)}(f)^t \cdot \mathbb{V} \left(\mathbb{G}[\mathcal{A}^{(k)}] \right) \cdot \Phi_k^{(N)}(f). \end{aligned} \quad (2.4)$$

so that the risk reduction holds whenever $\Delta(f) < 1$ and n is large enough.

When $N_1 > N_0 > 0$ there is no guaranty that the covariance structure of $\alpha_n^{(N_1)}(\mathcal{F})$ decreases compared to that of $\alpha_n^{(N_0)}(\mathcal{F})$. A simple sufficient condition is to rake two times along the same cycle of partitions.

Proposition 9. *Let $N_0, N_1 \in \mathbb{N}$ be such that $N_1 \geq 2N_0$ and*

$$\mathcal{A}^{(N_0-k)} = \mathcal{A}^{(N_1-k)}, \text{ for } 0 \leq k < N_0.$$

Then it holds $\mathbb{V}(\mathbb{G}^{(N_1)}(f)) \leq \mathbb{V}(\mathbb{G}^{(N_0)}(f))$ for all $f \in \mathcal{F}$ and $\Sigma_m^{(N_0)} - \Sigma_m^{(N_1)}$ is positive definite for all $\{f_1, \dots, f_m\} \subset \mathcal{F}$.

Remark J. In Appendix .2 a counter-example with $N_1 = N_0 + 1$ shows that the variance is not decreasing for all functions at each iteration. This case is excluded from Proposition 9 since $N_1 = N_0 + 1 < 2N_0$ if $N_0 > 1$ and, whenever $N_0 = 1$ and $N_1 = 2$ the requirement $\mathcal{A}^{(N_0)} = \mathcal{A}^{(N_1)}$ is not allowed.

2.3 The two margins case

We now consider the original method where k partitions are raked in a periodic order. Let us focus on the case $k = 2$ of the two-way contingency table case. The Deming and Stephan algorithm coincides with the Sinkhorn algorithm for matrix scaling. Denote $\mathcal{A} = \mathcal{A}^{(1)} = \{A_1, \dots, A_{m_1}\}$ and $\mathcal{B} = \mathcal{A}^{(2)} = \{B_1, \dots, B_{m_2}\}$ the

two known margins, thus $\mathcal{A}^{(2m+1)} = \mathcal{A}$ and $\mathcal{A}^{(2m)} = \mathcal{B}$. Likewise for $1 \leq i \leq m_1$ and $1 \leq j \leq m_2$ rewrite $(\mathbf{P}_{\mathcal{A}|\mathcal{B}})_{i,j} = P(A_j|B_i)$, $(\mathbf{P}_{\mathcal{B}|\mathcal{A}})_{i,j} = P(B_j|A_i)$ and

$$\begin{aligned}\mathbb{G}[\mathcal{A}] &= (\mathbb{G}(A_1), \dots, \mathbb{G}(A_{m_1}))^t, & \mathbb{E}[f|\mathcal{A}] &= (\mathbb{E}[f|A_1], \dots, \mathbb{E}[f|A_{m_1}])^t, \\ \mathbb{G}[\mathcal{B}] &= (\mathbb{G}(B_1), \dots, \mathbb{G}(B_{m_2}))^t, & \mathbb{E}[f|\mathcal{B}] &= (\mathbb{E}[f|B_1], \dots, \mathbb{E}[f|B_{m_2}])^t.\end{aligned}$$

The size of $\mathbf{P}_{\mathcal{A}|\mathcal{B}}\mathbf{P}_{\mathcal{B}|\mathcal{A}}$ is $m_1 \times m_1$ and that of $\mathbf{P}_{\mathcal{B}|\mathcal{A}}\mathbf{P}_{\mathcal{A}|\mathcal{B}}$ is $m_2 \times m_2$. A sum with a negative upper index is null, a matrix with negative power is also null, a square matrix with power zero is the identity matrix. For $N \in \mathbb{N}^*$ define

$$S_{1,\text{even}}^{(N)}(f) = \sum_{k=0}^N (\mathbf{P}_{\mathcal{B}|\mathcal{A}}\mathbf{P}_{\mathcal{A}|\mathcal{B}})^k \cdot (\mathbb{E}[f|\mathcal{A}] - \mathbf{P}_{\mathcal{B}|\mathcal{A}} \cdot \mathbb{E}[f|\mathcal{B}]) \text{ is } m_1 \times 1, \quad (2.5)$$

$$S_{2,\text{odd}}^{(N)}(f) = \sum_{k=0}^N (\mathbf{P}_{\mathcal{A}|\mathcal{B}}\mathbf{P}_{\mathcal{B}|\mathcal{A}})^k \cdot (\mathbb{E}[f|\mathcal{B}] - \mathbf{P}_{\mathcal{A}|\mathcal{B}} \cdot \mathbb{E}[f|\mathcal{A}]) \text{ is } m_2 \times 1, \quad (2.6)$$

$$S_{1,\text{odd}}^{(N)}(f) = S_{1,\text{even}}^{(N)} + (\mathbf{P}_{\mathcal{B}|\mathcal{A}}\mathbf{P}_{\mathcal{A}|\mathcal{B}})^{N+1} \cdot \mathbb{E}[f|\mathcal{A}] \text{ is } m_1 \times 1, \quad (2.7)$$

$$S_{2,\text{even}}^{(N)}(f) = S_{2,\text{odd}}^{(N)} + (\mathbf{P}_{\mathcal{A}|\mathcal{B}}\mathbf{P}_{\mathcal{B}|\mathcal{A}})^{N+1} \cdot \mathbb{E}[f|\mathcal{B}] \text{ is } m_2 \times 1. \quad (2.8)$$

Proposition 10. *Let $m \in \mathbb{N}$. We have*

$$\mathbb{G}^{(2m)}(f) = \mathbb{G}(f) - S_{1,\text{even}}^{(m-1)}(f)^t \cdot \mathbb{G}[\mathcal{A}] - S_{2,\text{even}}^{(m-2)}(f)^t \cdot \mathbb{G}[\mathcal{B}], \quad (2.9)$$

$$\mathbb{G}^{(2m+1)}(f) = \mathbb{G}(f) - S_{1,\text{odd}}^{(m-1)}(f)^t \cdot \mathbb{G}[\mathcal{A}] - S_{2,\text{odd}}^{(m-1)}(f)^t \cdot \mathbb{G}[\mathcal{B}]. \quad (2.10)$$

Remark K. The limiting process $\mathbb{G}^{(N)}$ evaluated at f is then simply $\mathbb{G}(f)$ with a correction depending on the Gaussian vectors $\mathbb{G}[\mathcal{A}]$ and $\mathbb{G}[\mathcal{B}]$ through the two deterministic matrices $\mathbf{P}_{\mathcal{A}|\mathcal{B}}$ and $\mathbf{P}_{\mathcal{B}|\mathcal{A}}$ carrying the information and operating on the conditional expectation vectors $\mathbb{E}[f|\mathcal{A}]$ and $\mathbb{E}[f|\mathcal{B}]$.

The following assumption simplifies the limits and ensures a geometric rate of convergence for matrices $S_{i,\text{even}}^{(N)}$ and $S_{1,\text{odd}}^{(N)}$ as $N \rightarrow +\infty$.

Hypothesis (ER). *The matrices $\mathbf{P}_{\mathcal{A}|\mathcal{B}}\mathbf{P}_{\mathcal{B}|\mathcal{A}}$ and $\mathbf{P}_{\mathcal{B}|\mathcal{A}}\mathbf{P}_{\mathcal{A}|\mathcal{B}}$ are ergodic.*

Remark L. Notice that (ER) holds whenever the matrices have strictly positive coefficients. This is true for $\mathbf{P}_{\mathcal{A}|\mathcal{B}}\mathbf{P}_{\mathcal{B}|\mathcal{A}}$ if $\sum_{j=1}^{m_2} P(A \cap B_j)P(B_j \cap A') > 0$ for all $A, A' \in \mathcal{A}$ hence if each pair $A, A' \in \mathcal{A}$ is intersected by some $B \in \mathcal{B}$ with positive probability. The later requirement is for instance met if $\mathcal{X} = \mathbb{R}^d$, P has a positive density and the partitions concern two distinct coordinates.

Proposition 11. *Under (ER) the matrices $S_{l,\text{even}}^{(N)}(f)$ and $S_{l,\text{odd}}^{(N)}(f)$ for $l = 1, 2$ converge uniformly on \mathcal{F} to $S_{l,\text{even}}(f)$ and $S_{l,\text{odd}}(f)$ satisfying*

$$S_{1,\text{odd}}(f) = S_{1,\text{even}}(f) + P_1[f], \quad S_{2,\text{even}}(f) = S_{2,\text{odd}}(f) + P_2[f],$$

where $P_l[f] = (P(f), \dots, P(f))^t$ are $m_l \times 1$ vectors. More precisely, given any vector norms $\|\cdot\|_{m_l}$ for $l = 1, 2$, there exists $c_l > 0$ and $0 < \lambda_l < 1$ such that

$$\begin{aligned} \sup_{f \in \mathcal{F}} \left\| S_{l, \text{even}}^{(N)}(f) - S_{l, \text{even}}(f) \right\|_{m_l} &\leq c_l \lambda_l^N \\ \sup_{f \in \mathcal{F}} \left\| S_{l, \text{odd}}^{(N)}(f) - S_{l, \text{odd}}(f) \right\|_{m_l} &\leq c_l \lambda_l^N. \end{aligned}$$

The main result of this section is the simple expression of the limiting process for a two partitions raking procedure. Let d_{LP} denote the Lévy-Prokhorov distance. The matrices $S_{1, \text{even}}(f), S_{2, \text{odd}}(f)$ and scalars λ_1, λ_2 are as in Proposition 11.

Theorem 2.2. *Under (ER) the sequence $\{\mathbb{G}^{(N)}(\mathcal{F})\}$ defined at (1.11) converges almost surely to the centered Gaussian process $\mathbb{G}^{(\infty)}(\mathcal{F})$ defined to be*

$$\mathbb{G}^{(\infty)}(f) = \mathbb{G}(f) - S_{1, \text{even}}(f)^t \cdot \mathbb{G}[\mathcal{A}] - S_{2, \text{odd}}(f)^t \cdot \mathbb{G}[\mathcal{B}], \quad f \in \mathcal{F}.$$

Moreover we have, for all N large and $c_3 > 0$ depending on $\lambda_1, \lambda_2, P(\mathcal{A}), P(\mathcal{B})$,

$$d_{LP}(\mathbb{G}^{(N)}, \mathbb{G}^{(\infty)}) \leq c_3 \sqrt{N} \max(\lambda_1, \lambda_2)^{N/2}.$$

Theorem 2.2 may be viewed as a stochastic counterpart of the deterministic rate obtained by Franklin and Lorentz [11] for the Sinkhorn algorithm. Mixing both approaches could strengthen the following two remarks.

Remark M. The matrices $\mathbf{P}_{\mathcal{A}|\mathcal{B}}, \mathbf{P}_{\mathcal{B}|\mathcal{A}}$ and the vectors $\mathbb{E}[f|\mathcal{A}], \mathbb{E}[f|\mathcal{B}]$ are not known without additional information. They can be estimated uniformly over \mathcal{F} as $n \rightarrow +\infty$ to evaluate the distribution of $\mathbb{G}^{(N)}$ and $\mathbb{G}^{(\infty)}$, thus giving access to adaptative tests or estimators. Since λ_1, λ_2 and c_3 are related to eigenvalues of $\mathbf{P}_{\mathcal{A}|\mathcal{B}}\mathbf{P}_{\mathcal{B}|\mathcal{A}}$ and $\mathbf{P}_{\mathcal{B}|\mathcal{A}}\mathbf{P}_{\mathcal{A}|\mathcal{B}}$ they can be estimated adaptively at rate $1/\sqrt{n}$ in probability. This in turn provides an evaluation of $d_{LP}(\mathbb{G}^{(N)}, \mathbb{G}^{(\infty)})$.

Remark N. In the case of an auxiliary information reduced to $P(\mathcal{A}), P(\mathcal{B})$ one should use $\mathcal{A} = \{A, A^c\}, \mathcal{B} = \{B, B^c\}$, estimate the missing $P(A \cap B)$ in $\mathbf{P}_{\mathcal{A}|\mathcal{B}}, \mathbf{P}_{\mathcal{B}|\mathcal{A}}$ and the conditional expectations on the four sets, then $S_{1, \text{even}}, S_{2, \text{odd}}$. On the opposite, the above characterization of the limiting process $\mathbb{G}^{(\infty)}(\mathcal{F})$ can be generalized to a recursive raking among k partitions in the same order.

As a conclusion the above $\mathbb{G}^{(\infty)}$ is the intrinsic limiting process when $n \rightarrow +\infty$ then $N \rightarrow +\infty$. Even if n is large it is wise to choose a very small N to guaranty good constants in previous sections, then estimate $d_{LP}(\mathbb{G}^{(N)}, \mathbb{G}^{(\infty)})$ from the sample as in Remark M. If it is sufficiently small then $\mathbb{G}^{(N)}$ may serve to evaluate risks and confident regions, by estimating $S_{l, \text{even}}^{(N)}, S_{l, \text{odd}}^{(N)}$ at required f . Otherwise N may increase.

3 Proofs of general results

3.1 Raking formulas

Write

$$B_{n,N_0} = \left\{ \min_{0 \leq N \leq N_0} \min_{1 \leq j \leq m_N} \mathbb{P}_n \left(A_j^{(N)} \right) > 0 \right\} \quad (3.1)$$

and $B_{n,N_0}^c = \Omega \setminus B_{n,N_0}$. The probability that $\alpha_n^{(N_0)}$ is undefined is

$$P(B_{n,N_0}^c) \leq \sum_{N=1}^{N_0} m_N (1 - \delta_N)^n \leq S_{N_0} (1 - \delta_{(N_0)})^n.$$

On B_{n,N_0} we have, by (1.3),

$$\begin{aligned} & \alpha_n^{(N)}(f) \\ &= \sqrt{n}(\mathbb{P}_n^{(N)}(f) - P(f)) \\ &= \sqrt{n} \left(\sum_{j=1}^{m_N} \frac{P(A_j^{(N)})}{\mathbb{P}_n^{(N-1)}(A_j^{(N)})} \mathbb{P}_n^{(N-1)}(f1_{A_j^{(N)}}) - \sum_{j=1}^{m_N} P(f1_{A_j^{(N)}}) \right) \\ &= \sum_{j=1}^{m_N} \left(\frac{P(A_j^{(N)})}{\mathbb{P}_n^{(N-1)}(A_j^{(N)})} \alpha_n^{(N-1)}(f1_{A_j^{(N)}}) - \frac{P(f1_{A_j^{(N)}})}{\mathbb{P}_n^{(N-1)}(A_j^{(N)})} \alpha_n^{(N-1)}(A_j^{(N)}) \right) \\ &= \sum_{j=1}^{m_N} \frac{P(A_j^{(N)})}{\mathbb{P}_n^{(N-1)}(A_j^{(N)})} \alpha_n^{(N-1)} \left((f - \mathbb{E}(f|A_j^{(N)}))1_{A_j^{(N)}} \right). \end{aligned} \quad (3.2)$$

By (1.7) this implies (1.8) since for any $A \in \mathcal{A}$,

$$\alpha_n^{(N-1)}(f1_A) - \mathbb{E}(f|A)\alpha_n^{(N-1)}(A) = \mathbb{P}_n^{(N-1)}(f1_A) - \mathbb{E}(f|A)\mathbb{P}_n^{(N-1)}(A).$$

Define $\mathcal{A}_\cap^{(0)} = \{\Omega\}$ and, for $N \geq 1$,

$$\mathcal{A}_\cap^{(N)} = \left\{ A : A = A_{j_1}^{(1)} \cap A_{j_2}^{(2)} \cap \dots \cap A_{j_N}^{(N)}, j_k \leq m_k, k \leq N \right\}. \quad (3.3)$$

For any $A \in \mathcal{A}_\cap^{(N)}$, $\mathbb{P}_n^{(N)}$ associates to each $X_i \in A$ the weight

$$\omega_n^{(N)}(A) = \frac{1}{n} \prod_{k=1}^N \frac{P(A_{j_k}^{(k)})}{\mathbb{P}_n^{(k-1)}(A_{j_k}^{(k)})}. \quad (3.4)$$

The case $N = 1$ is (1.1). By induction on (1.3), (3.4) and since $\mathcal{A}_\cap^{(N)}$ is a refined finite partition of \mathcal{X} , we get

$$\begin{aligned}
\mathbb{P}_n^{(N+1)}(f) &= \sum_{j=1}^{m_{N+1}} \frac{P(A_j^{(N+1)})}{\mathbb{P}_n^{(N)}(A_j^{(N+1)})} \mathbb{P}_n^{(N)}(f \mathbf{1}_{A_j^{(N+1)}}) \\
&= \sum_{j=1}^{m_{N+1}} \frac{P(A_j^{(N+1)})}{\mathbb{P}_n^{(N)}(A_j^{(N+1)})} \sum_{A \in \mathcal{A}_\cap^{(N)}} \sum_{i=1}^n \omega_n^{(N)}(A) f(X_i) \mathbf{1}_{A \cap A_j^{(N+1)}}(X_i) \\
&= \sum_{i=1}^n f(X_i) \sum_{A \in \mathcal{A}_\cap^{(N)}} \sum_{j=1}^{m_{N+1}} \mathbf{1}_{A \cap A_j^{(N+1)}}(X_i) \omega_n^{(N+1)}(A \cap A_j^{(N+1)}) \\
&= \sum_{i=1}^n f(X_i) \sum_{A \in \mathcal{A}_\cap^{(N+1)}} \mathbf{1}_A(X_i) \omega_n^{(N+1)}(A).
\end{aligned}$$

3.2 Proof of Proposition 1.

The partition $\mathcal{A}_\cap^{(N)}$ is defined at (3.3). By using (3.4) it holds, for $N \geq 1$,

$$\begin{aligned}
d_K \left(\mathbb{P}_n^{(N-1)} \parallel \mathbb{P}_n^{(N)} \right) &= \sum_{i=1}^n \mathbb{P}_n^{(N-1)}(\{X_i\}) \log \left(\frac{\mathbb{P}_n^{(N-1)}(\{X_i\})}{\mathbb{P}_n^{(N)}(\{X_i\})} \right) \\
&= \sum_{i=1}^n \mathbb{P}_n^{(N-1)}(\{X_i\}) \sum_{A \in \mathcal{A}_\cap^{(N-1)}} \sum_{j=1}^{m_N} \mathbf{1}_{A \cap A_j^{(N)}}(X_i) \log \left(\frac{\omega_n^{(N-1)}(A)}{\omega_n^{(N)}(A \cap A_j^{(N)})} \right) \\
&= \sum_{j=1}^{m_N} \log \left(\frac{\mathbb{P}_n^{(N-1)}(A_j^{(N)})}{P(A_j^{(N)})} \right) \sum_{i=1}^n \mathbb{P}_n^{(N-1)}(\{X_i\}) \sum_{A \in \mathcal{A}_\cap^{(N-1)}} \mathbf{1}_{A \cap A_j^{(N)}}(X_i) \\
&= \sum_{j=1}^{m_N} \mathbb{P}_n^{(N-1)}(A_j^{(N)}) \log \left(\frac{\mathbb{P}_n^{(N-1)}(A_j^{(N)})}{P(A_j^{(N)})} \right) \tag{3.5}
\end{aligned}$$

since $\mathcal{A}_\cap^{(N-1)}$ is a partition of \mathcal{X} . Hence the contrast between $\mathbb{P}_n^{(N-1)}$ and $\mathbb{P}_n^{(N)}$ is the same on $\{X_1, \dots, X_n\}$ as on $\mathcal{A}^{(N)}$. Now, by convexity of $-\log(x)$ it follows,

for any probability law Q supported by $\{X_1, \dots, X_n\}$,

$$\begin{aligned}
& d_K \left(\mathbb{P}_n^{(N-1)} \parallel Q \right) \\
&= - \sum_{i=1}^n \log \left(\frac{Q(\{X_i\})}{\mathbb{P}_n^{(N-1)}(\{X_i\})} \right) \sum_{j=1}^{m_N} \mathbb{P}_n^{(N-1)}(\{X_i\}) 1_{A_j^{(N)}}(X_i) \\
&= - \sum_{j=1}^{m_N} \mathbb{P}_n^{(N-1)}(A_j^{(N)}) \sum_{i=1}^n \frac{\mathbb{P}_n^{(N-1)}(\{X_i\})}{\mathbb{P}_n^{(N-1)}(A_j^{(N)})} 1_{A_j^{(N)}}(X_i) \log \left(\frac{Q(\{X_i\})}{\mathbb{P}_n^{(N-1)}(\{X_i\})} \right) \\
&\geq - \sum_{j=1}^{m_N} \mathbb{P}_n^{(N-1)}(A_j^{(N)}) \log \left(\sum_{i=1}^n \frac{Q(\{X_i\})}{\mathbb{P}_n^{(N-1)}(A_j^{(N)})} 1_{A_j^{(N)}}(X_i) \right) \\
&= \sum_{j=1}^{m_N} \mathbb{P}_n^{(N-1)}(A_j^{(N)}) \log \left(\frac{\mathbb{P}_n^{(N-1)}(A_j^{(N)})}{P(A_j^{(N)})} \right) = d_K \left(\mathbb{P}_n^{(N-1)} \parallel \mathbb{P}_n^{(N)} \right)
\end{aligned}$$

where the final identification relies on (3.5) and $\mathbb{P}_n^{(N)} = P$ on $\mathcal{A}^{(N)}$.

3.3 Proof of Proposition 2.

The classical law of the iterated logarithm holds for the empirical process $\alpha_n^{(0)}$ indexed by \mathcal{F} under (VC) and (BR). See Alexander [1], in particular Theorem 2.12 for (BR) and Theorem 2.13 based on Theorem 2.8 that uses in its proof the consequence of Lemma 2.7, which is indeed (VC). Namely, for any $\varepsilon > 0$, with probability one there exists $n(\omega)$ such that, for all $n > n(\omega)$,

$$u_n \|\mathbb{P}_n^{(0)} - P\|_{\mathcal{F}} < 1 + \varepsilon, \quad u_n = \sqrt{\frac{n}{2\sigma_{\mathcal{F}}^2 L \circ L(n)}} \quad (3.6)$$

where $\sigma_{\mathcal{F}}^2 = \sup_{\mathcal{F}} \mathbb{V}(f) \leq M^2$. Let $1 \leq N \leq N_0$. By (1.3) it holds

$$\begin{aligned}
\mathbb{P}_n^{(N)}(f) - \mathbb{P}_n^{(N-1)}(f) &= \sum_{j=1}^{m_N} \mathbb{P}_n^{(N-1)}(f 1_{A_j^{(N)}}) \frac{P(A_j^{(N)})}{\mathbb{P}_n^{(N-1)}(A_j^{(N)})} - \sum_{j=1}^{m_N} \mathbb{P}_n^{(N-1)}(f 1_{A_j^{(N)}}) \\
&= \sum_{j=1}^{m_N} \mathbb{E}_n^{(N-1)}(f \mid A_j^{(N)}) \left(P(A_j^{(N)}) - \mathbb{P}_n^{(N-1)}(A_j^{(N)}) \right).
\end{aligned}$$

Since $\mathbb{P}_n^{(N)}$ is a probability measure we have $\|\mathbb{P}_n^{(N)}(f 1_A)\|_{\mathcal{F}} \leq M \mathbb{P}_n^{(N)}(A)$ hence $\|\mathbb{E}_n^{(N-1)}(f \mid A_j^{(N)})\|_{\mathcal{F}} \leq M$ and $|\mathbb{P}_n^{(N)}(f) - \mathbb{P}_n^{(N-1)}(f)| \leq M m_N \|\mathbb{P}_n^{(N-1)} - P\|_{\mathcal{F}}$. Also observe that (1.2) combined to the fact that $\mathcal{A}^{(N)}$ is a partition implies $m_N \leq 1/\delta_N$ and $\delta_N \geq \delta_{(N)} \geq \delta_{(N_0)}$. Therefore

$$\begin{aligned}
u_n \|\mathbb{P}_n^{(N)} - P\|_{\mathcal{F}} &\leq u_n \|\mathbb{P}_n^{(N-1)} - P\|_{\mathcal{F}} + u_n \|\mathbb{P}_n^{(N)} - \mathbb{P}_n^{(N-1)}\|_{\mathcal{F}} \\
&\leq u_n (1 + M m_N) \|\mathbb{P}_n^{(N-1)} - P\|_{\mathcal{F}} \\
&\leq u_n \kappa_N \|\mathbb{P}_n^{(0)} - P\|_{\mathcal{F}}
\end{aligned}$$

where

$$\kappa_N = \prod_{k=1}^N (1 + Mm_N) \leq \prod_{k=1}^N \left(1 + \frac{M}{\delta_k}\right) \leq \left(1 + \frac{M}{\delta_{(N_0)}}\right)^{N_0}$$

which by (3.6) remains true for $N = N_0 = 0$. This proves that, given $N_0 \in \mathbb{N}$ and for all $\varepsilon > 0$,

$$\limsup_{n \rightarrow +\infty} \sqrt{\frac{n}{2L \circ L(n)}} \sup_{0 \leq N \leq N_0} \left\| \mathbb{P}_n^{(N)} - P \right\|_{\mathcal{F}} \leq (1 + \varepsilon) \kappa_N \sigma_{\mathcal{F}} \quad a.s.$$

and Proposition 2 follows.

3.4 Proof of Proposition 3

Step 1. We work on B_{n, N_0} from (3.1), which is unwritten but is in intersection with all subsequent events. By (1.8) and (3.2) we have, for $N \geq 1$,

$$\alpha_{n,j}^{(N-1)}(f) = \frac{1}{\mathbb{P}_n^{(N-1)}(A_j^{(N)})} \alpha_n^{(N-1)} \left((f - \mathbb{E}(f|A_j^{(N)})) \mathbf{1}_{A_j^{(N)}} \right)$$

with $|\mathbb{E}(f|A_j^{(N)})| \leq M$, and

$$\begin{aligned} \mathbb{P} \left(\left\| \alpha_n^{(N)} \right\|_{\mathcal{F}} \geq \lambda \right) &\leq \mathbb{P} \left(\sum_{j=1}^{m_N} P(A_j^{(N)}) \left\| \alpha_{n,j}^{(N-1)} \right\|_{\mathcal{F}} \geq \sum_{j=1}^{m_N} P(A_j^{(N)}) \lambda \right) \\ &\leq \sum_{j=1}^{m_N} \mathbb{P} \left(\left\| \alpha_{n,j}^{(N-1)} \right\|_{\mathcal{F}} \geq \lambda \right). \end{aligned}$$

Each term in the latter sum satisfies, for $K \leq P(A_j^{(N)})$ and $K' \leq P(A_j^{(N)}) - K$,

$$\begin{aligned} &\mathbb{P} \left(\frac{1}{\mathbb{P}_n^{(N-1)}(A_j^{(N)})} \left\| \alpha_n^{(N-1)}(f) - \mathbb{E}(f|A_j^{(N)}) \alpha_n^{(N-1)}(A_j^{(N)}) \right\|_{\mathcal{F}} \geq \lambda \right) \\ &\leq \mathbb{P} \left((1 + M) \left\| \alpha_n^{(N-1)} \right\|_{\mathcal{F}} \geq K\lambda \right) + \mathbb{P} \left(\mathbb{P}_n^{(N-1)}(A_j^{(N)}) \leq K \right) \\ &\leq \mathbb{P} \left(\left\| \alpha_n^{(N-1)} \right\|_{\mathcal{F}} \geq \frac{\lambda K}{1 + M} \right) + \mathbb{P} \left(\alpha_n^{(N-1)}(A_j^{(N)}) \leq -K' \sqrt{n} \right) \\ &\leq 2\mathbb{P} \left(\left\| \alpha_n^{(N-1)} \right\|_{\mathcal{F}} \geq \frac{\lambda K}{1 + M} \right) \end{aligned} \tag{3.7}$$

where the last bound holds provided that $K' \sqrt{n} \geq \lambda K / (1 + M)$. Define

$$\beta = \frac{1}{1 + \lambda / (1 + M) \sqrt{n}} \in (0, 1), \quad K = \beta \delta_N, \quad K' = 1 - \delta_N (1 + \beta).$$

Since $\delta_N \leq 1/2$ for any $N \geq 1$ it holds $K' > 0$ and, as required,

$$\frac{\lambda}{(1 + M) \sqrt{n}} = \frac{1 - \beta}{\beta} = \frac{1 - (1 + \beta)/2}{\beta/2} \leq \frac{1 - \delta_N (1 + \beta)}{\beta \delta_N} = \frac{K'}{K}.$$

We have shown that for any $N \geq 1$,

$$\mathbb{P} \left(\left\| \alpha_n^{(N)} \right\|_{\mathcal{F}} \geq \lambda \right) \leq 2m_N \mathbb{P} \left(\left\| \alpha_n^{(N-1)} \right\|_{\mathcal{F}} \geq \frac{\lambda \beta \delta_N}{1+M} \right).$$

Applying (3.7) again with λ turned into the smaller $\lambda \beta \delta_N / (1+M)$ then iterating from N_0 we get, for $\Delta_{N_0} = \prod_{N=1}^{N_0} \delta_N$ and $M_{N_0} = \prod_{N=1}^{N_0} m_N \leq 1/\Delta_{N_0}$,

$$\mathbb{P} \left(\left\| \alpha_n^{(N_0)} \right\|_{\mathcal{F}} \geq \lambda \right) \leq 2^{N_0} M_{N_0} \mathbb{P} \left(\left\| \alpha_n^{(0)} \right\|_{\mathcal{F}} \geq \frac{\lambda \Delta_{N_0}}{(1+M + \lambda/\sqrt{n})^{N_0}} \right).$$

The latter upper bound being increasing with N_0 we conclude that

$$\mathbb{P} \left(\max_{0 \leq N \leq N_0} \left\| \alpha_n^{(N)} \right\|_{\mathcal{F}} \geq \lambda \right) \leq \sum_{N=1}^{N_0} \mathbb{P} \left(\left\| \alpha_n^{(N)} \right\|_{\mathcal{F}} \geq \lambda \right) \leq N_0 \mathbb{P} \left(\left\| \alpha_n^{(N_0)} \right\|_{\mathcal{F}} \geq \lambda \right).$$

Step 2. By Talagrand [20], for $n \geq 1$, $t > 0$, independent Rademacher random variables $\epsilon_1, \dots, \epsilon_n$ also independent of X_1, \dots, X_n we have, for some universal constants $D'_1 > 0$, $D'_2 > 0$, $C > 0$,

$$\mathbb{P} \left(\left\| \alpha_n^{(0)} \right\|_{\mathcal{F}} > D'_1 (\mu_n + t) \right) \leq \exp \left(-D'_2 \min \left(\frac{t^2}{\sigma_{\mathcal{F}}^2}, \frac{t\sqrt{n}}{M} \right) \right)$$

where, by the last maximal inequality in Theorem 2.14.2 of van der Vaart and Wellner [21] applied to \mathcal{F} with envelop function constant to M , it holds

$$\mu_n = \mathbb{E} \left\| \frac{1}{\sqrt{n}} \sum_{i=1}^n \epsilon_i f(X_i) \right\|_{\mathcal{F}} \leq C \int_0^1 \sqrt{1 + \log N_{[\cdot]}(\mathcal{F}, M\varepsilon, d_P)} d\varepsilon$$

Under (BR), we have $\mu_n < C'$ with $C' = C(1 + b_0/(1 - r_0))$. For $\lambda_0 = 2D'_1 C'$ we get, for any $n > 0$ and $\lambda_0 < \lambda < 2D'_1 \sigma_{\mathcal{F}}^2 \sqrt{n}/M$ we get, for all n ,

$$\mathbb{P} \left(\left\| \alpha_n^{(0)} \right\|_{\mathcal{F}} > \lambda \right) \leq \mathbb{P} \left(\left\| \alpha_n^{(0)} \right\|_{\mathcal{F}} > D'_1 \left(\mu_n + \frac{\lambda}{2D'_1} \right) \right) \leq \exp(-D'_2 \lambda^2)$$

where $D'_2 = D'_2/4(D'_1)^2 \sigma_{\mathcal{F}}^2$. Therefore, according to step 2, taking

$$D_0 = \frac{2D'_1 \sigma_{\mathcal{F}}^2}{M}, \quad D_1 = N_0 2^{N_0} M_{N_0}, \quad D_2 = \frac{D'_2 \Delta_{N_0}^2}{(1+M + D_0)^{2N_0}} \quad (3.8)$$

yields $\mathbb{P}(\max_{0 \leq N \leq N_0} \left\| \alpha_n^{(N)} \right\|_{\mathcal{F}} \geq \lambda) \leq D_1 \exp(-D_2 \lambda^2)$ for $\lambda_0 < \lambda < D_0 \sqrt{n}$.

Step 3. By Theorem 2.14.9 in [21], under (VC) there exists a constant $D(c_0)$ such that, for t_0 large enough and all $t \geq t_0$,

$$\mathbb{P} \left(\left\| \alpha_n^{(0)} \right\|_{\mathcal{F}} \geq t \right) \leq \left(\frac{D(c_0)t}{M\sqrt{v_0}} \right)^{v_0} \exp \left(-\frac{2t^2}{M^2} \right).$$

Denote $\lambda_{1,n}$ and $\lambda_{2,n}$ the two solutions of $\lambda\Delta_{N_0} = t_0(1 + M + \lambda/\sqrt{n})^{N_0}$. Notice that, for n large, $\lambda_{1,n}$ is close to $t_0(1 + M)^{N_0}/\Delta_{N_0}$ and $\lambda_{2,n} = O(n^{N_0/2(N_0-1)})$. It ensues that for some n_0, λ_0 it holds, for all $n > n_0$ and $\lambda_0 < \lambda < 2M\sqrt{n}$, the bound $\mathbb{P}(\max_{0 \leq N \leq N_0} \|\alpha_n^{(N)}\|_{\mathcal{F}} \geq \lambda) \leq D_3\lambda^{v_0} \exp(-D_4\lambda^2)$ where

$$D_3 = \frac{N_0 2^{N_0} M_{N_0}}{(1+M)^{v_0 N_0}} \left(\frac{D(c_0)\Delta_{N_0}}{M\sqrt{v_0}} \right)^{v_0}, \quad D_4 = \frac{2\Delta_{N_0}^2}{M^2(3M+1)^{2N_0}}. \quad (3.9)$$

Finally, at each step, add $S_{N_0} (1 - \delta_{(N_0)})^n$ to take B_{n,N_0}^c from (3.1) into account.

3.5 Proof of Proposition 4 and Theorem 2.1.

Theorem 2.1 implies Proposition 4 since the weak convergence on $(\ell^\infty(\mathcal{F}), \|\cdot\|_{\mathcal{F}})$ is metricized by the Lévy-Prokhorov distance between $\alpha_n^{(N)}$ and $\mathbb{G}_n^{(N)}$ which is

$$\inf \left\{ \varepsilon > 0 : \mathbb{P}^{\alpha_n^{(N)}}(A) \leq \mathbb{P}^{\mathbb{G}_n^{(N)}}(A^\varepsilon) + \varepsilon, \mathbb{P}^{\mathbb{G}_n^{(N)}}(A) \leq \mathbb{P}^{\alpha_n^{(N)}}(A^\varepsilon) + \varepsilon \right\} \leq v_n. \quad (3.10)$$

To see this, recall that $v_n > 1/n^{\theta_0}$ and $v_n \rightarrow 0$ in Theorem 2.1, remind (1.6) then observe that

$$\begin{aligned} \mathbb{P}(\alpha_n^{(N)} \in A) &\leq \mathbb{P}\left(\left\{\alpha_n^{(N)} \in A\right\} \cap \left\{\left\|\alpha_n^{(N)} - \mathbb{G}_n^{(N)}\right\|_{\mathcal{F}} \leq v_n\right\} \cap B_{n,N_0}\right) \\ &\quad + \mathbb{P}\left(\left\{\left\|\alpha_n^{(N)} - \mathbb{G}_n^{(N)}\right\|_{\mathcal{F}} > v_n\right\} \cap B_{n,N_0}\right) + \mathbb{P}(B_{n,N_0}^c) \\ &\leq \mathbb{P}(\mathbb{G}_n^{(N)} \in A^{v_n}) + \frac{1}{n^{\theta_0}} + S_{N_0} (1 - \delta_{(N_0)})^n \end{aligned}$$

remains true by exchanging $\alpha_n^{(N)}$ and $\mathbb{G}_n^{(N)}$. Since v_n is the slowest sequence, if n_0 satisfies $v_{n_0} > 1/n_0^{\theta_0} + S_{N_0} (1 - \delta_{(N_0)})^{n_0}$ then $v_n > 1/n^{\theta_0} + S_{N_0} (1 - \delta_{(N_0)})^n$ for all $n > n_0$. Whence (3.10).

We next establish Theorem 2.1. Fix $N_0 \in \mathbb{N}$.

Step 1. Let introduce the transforms, for $f \in \mathcal{F}$, $N \geq 1$ and $1 \leq j \leq m_N$,

$$\begin{aligned} \phi_{(j,N)}f &= \left(f - \mathbb{E}\left(f \mid A_j^{(N)}\right)\right) 1_{A_j^{(N)}}, \\ \phi_{(N)}f &= \sum_{j=1}^{m_N} \phi_{(j,N)}f = f - \sum_{j=1}^{m_N} \mathbb{E}(f \mid A_j^{(N)}) 1_{A_j^{(N)}}. \end{aligned}$$

It holds $P(\phi_{(N)}f) = P(\phi_{(j,N)}f) = 0$ and, since $\mathcal{A}^{(N)}$ is a partition of \mathcal{X} ,

$$(\phi_{(j,N)}f)(\phi_{(j',N)}g) = 0, \quad 1 \leq j \neq j' \leq m_N. \quad (3.11)$$

Moreover, the $L_2(P)$ property of conditional expectations yields

$$\sigma_{\phi_{(j,N)}f}^2 = P(f_{(j,N)}^2) \leq \sigma_{\phi_{(N)}f}^2 = P(f_{(N)}^2) = \sum_{j=1}^{m_N} \sigma_{\phi_{(j,N)}f}^2 \leq \sigma_f^2. \quad (3.12)$$

Next consider the class of backward iterated transforms

$$\begin{aligned}\mathcal{F}_{(N)} &= \phi_{(1)} \circ \dots \circ \phi_{(N)}(\mathcal{F}), \\ \mathcal{H}_{(N)} &= \bigcup_{1 \leq k \leq N} \bigcup_{1 \leq j \leq m_k} \phi_{(j,k)} \circ \phi_{(k+1)} \circ \dots \circ \phi_{(N)}(\mathcal{F}),\end{aligned}$$

where $\phi_{(k+1)} \circ \dots \circ \phi_{(N)} = id$ if $k = N \geq 1$ and $\mathcal{F}_{(0)} = \mathcal{H}_{(0)} = \mathcal{F}$. Also write $\mathcal{F}_0 = \bigcup_{0 \leq N \leq N_0} \mathcal{F}_{(N)}$ and $\mathcal{H}_0 = \bigcup_{0 \leq N \leq N_0} \mathcal{H}_{(N)}$. By iterating (3.12) it comes $\sigma_{\mathcal{H}_0}^2 \leq \sigma_{\mathcal{F}_0}^2 \leq \sigma_{\mathcal{F}}^2$. We first show that properties of \mathcal{F} transfer to $\mathcal{F}_{(N)}$, $\mathcal{H}_{(N)}$ for $0 \leq N \leq N_0$ and thus to \mathcal{F}_0 and \mathcal{H}_0 . Remind the constants defined at (2.2).

Lemma 1. *Assume (1.2). If \mathcal{F} is pointwise measurable and bounded by M then $\mathcal{F}_{(N)}$ and $\mathcal{H}_{(N)}$ (resp. \mathcal{F}_0 and \mathcal{H}_0) are pointwise measurable and bounded by $(2M)^N/\Delta_N$ (resp. $(2M)^{N_0}/\Delta_{N_0}$). If (VC) (resp. (BR)) holds then \mathcal{F}_0 and \mathcal{H}_0 also satisfies (VC) (resp. (BR)) with the same power ν_0 (resp. r_0) as \mathcal{F} .*

Proof. If \mathcal{F} is uniformly bounded by M then for $N \leq N_0$ we have

$$\sup_{\mathcal{F}} \sup_{\mathcal{X}} |\phi_{(N)} f| = \sup_{\mathcal{F}} \max_{1 \leq j \leq m_N} \sup_{\mathcal{X}} |\phi_{(j,N)} f| \leq M \left(1 + \frac{1}{\delta_N}\right) \leq \frac{2M}{\delta_N}$$

thus, by backward induction from N_0 to 1, $\mathcal{F}_{(N_0)}$ and $\mathcal{H}_{(N_0)}$ are uniformly bounded by $(2M)^{N_0}/\Delta_{N_0}$. It readily follows that \mathcal{F}_0 and \mathcal{H}_0 are bounded by $(2M)^{N_0}/\Delta_{N_0}$. Assume that $f_k \in \mathcal{F}_*$ converges pointwise on \mathcal{X} to $f \in \mathcal{F}$. From

$$\lim_{k \rightarrow +\infty} 1_{A_j^{(N)}}(X) f_k(X) = 1_{A_j^{(N)}}(X) f(X) \quad \text{and} \quad P(1_{A_j^{(N)}} f_k) \leq P(f_k) \leq M$$

we deduce by dominated convergence that $\lim_{k \rightarrow +\infty} \mathbb{E}(f_k | A_j^{(N)}) = \mathbb{E}(f | A_j^{(N)})$. Thus $\phi_{(j,N)} f_k$ converges pointwise to $\phi_{(j,N)} f$ and $\phi_{(N)} f_k = \sum_{j=1}^{m_N} \phi_{(j,N)} f_k$ to $\phi_{(N)} f = \sum_{j=1}^{m_N} \phi_{(j,N)} f$. By iterating this reasoning backward from N to 1 we obtain that $\mathcal{F}_{(N)}$ and $\mathcal{H}_{(N)}$ are pointwise measurable, by using the countable classes $\phi_{(1)} \circ \dots \circ \phi_{(N)}(\mathcal{F}_*)$ and $\bigcup_{1 \leq k \leq N} \bigcup_{1 \leq j \leq m_k} \phi_{(j,k)} \circ \phi_{(k+1)} \circ \dots \circ \phi_{(N)}(\mathcal{F}_*)$ respectively. Assume next that \mathcal{F} satisfies (VC). By (3.11) we have

$$\begin{aligned}d_Q^2(\phi_{(N)} f, \phi_{(N)} g) &= \int_{\mathcal{X}} \left(\sum_{j=1}^{m_N} (\phi_{(j,N)} f - \phi_{(j,N)} g) \right)^2 dQ \\ &= \sum_{j=1}^{m_N} d_Q^2(\phi_{(j,N)} f, \phi_{(j,N)} g) \\ &= \sum_{j=1}^{m_N} \int_{A_j^{(N)}} \left(f - g - \mathbb{E}(f - g | A_j^{(N)}) \right)^2 dQ \\ &\leq \sum_{j=1}^{m_N} \int_{A_j^{(N)}} (f - g - (Qf - Qg))^2 dQ \\ &= d_Q^2(f, g) - (Qf - Qg)^2\end{aligned}$$

thus $d_Q(f, g) < \varepsilon$ implies $d_Q^2(\phi_{(N)} f, \phi_{(N)} g) \leq d_Q^2(f, g) < \varepsilon^2$. If \mathcal{F} can be covered by $N(\mathcal{F}, \varepsilon, d_Q)$ balls of d_Q -radius ε centered at some g then $\phi_{(N)}(\mathcal{F})$ can be

covered by the same number of balls, centered at the corresponding $\phi_{(N)}g$ and hence the same number of centers $\phi_{(1)} \circ \dots \circ \phi_{(N)}g$ suffices to cover $\mathcal{F}_{(N)}$. All the $\phi_{(j,k)} \circ \phi_{(k+1)} \circ \dots \circ \phi_{(N)}g$ are needed to cover $\mathcal{H}_{(N)}$, that is $S_N N(\mathcal{F}, \varepsilon, d_Q)$. This shows that \mathcal{F}_0 (resp. \mathcal{H}_0) obeys (VC) with the same power ν_0 and a constant $c_0(N_0 + 1)$ (resp. $c_0 \sum_{N=0}^{N_0} S_N$). Assume now that \mathcal{F} satisfies (BR). If $g_- \leq f \leq g_+$ then we have

$$\begin{aligned} h_{(j,N)}^- &= 1_{A_j^{(N)}} g_- - 1_{A_j^{(N)}} \mathbb{E}(g_+ | A_j^{(N)}) \\ &\leq \phi_{(j,N)} f \leq 1_{A_j^{(N)}} g_+ - 1_{A_j^{(N)}} \mathbb{E}(g_- | A_j^{(N)}) = h_{(j,N)}^+ \end{aligned}$$

and the $L_2(P)$ -size of the new bracket $[h_{(j,N)}^-, h_{(j,N)}^+]$ is

$$\begin{aligned} d_P^2(h_{(j,N)}^-, h_{(j,N)}^+) &= \int_{A_j^{(N)}} \left(g_+ - g_- + \mathbb{E}(g_+ - g_- | A_j^{(N)}) \right)^2 dP \\ &= P(1_{A_j^{(N)}}(g_+ - g_-)^2) + P(A_j^{(N)}) \mathbb{E}(g_+ - g_- | A_j^{(N)})^2 \\ &\quad + 2\mathbb{E}(g_+ - g_- | A_j^{(N)}) P(1_{A_j^{(N)}}(g_+ - g_-)). \end{aligned}$$

If $d_P(g_+, g_-) < \varepsilon$ the Hölder inequality yields $P(1_{A_j^{(N)}}(g_+ - g_-)) \leq \varepsilon \sqrt{P(A_j^{(N)})}$ and $\mathbb{E}(g_+ - g_- | A_j^{(N)}) \leq \varepsilon / \sqrt{P(A_j^{(N)})}$ hence

$$d_P^2(h_{(j,N)}^-, h_{(j,N)}^+) \leq P(1_{A_j^{(N)}}(g_+ - g_-)^2) + 3\varepsilon^2$$

so that $\phi_{(N)}f = \sum_{j=1}^{m_N} \phi_{(j,N)}f \in [h_{(N)}^-, h_{(N)}^+]$ where $h_{(N)}^\pm = \sum_{j=1}^{m_N} h_{(j,N)}^\pm$ satisfies

$$d_P^2(h_{(N)}^-, h_{(N)}^+) = \sum_{j=1}^{m_N} d_P^2(h_{(j,N)}^-, h_{(j,N)}^+) \leq d_P^2(g_-, g_+) + 3m_N \varepsilon^2 \leq 4m_N \varepsilon^2.$$

It ensues $N_{[\]}(\phi_{(N)}(\mathcal{F}), \varepsilon, d_P) \leq N_{[\]}(\mathcal{F}, \varepsilon/2\sqrt{m_N}, d_P)$ and $N_{[\]}(\mathcal{F}_{(N)}, \varepsilon, d_P) \leq N_{[\]}(\mathcal{F}, \varepsilon/2^N \sqrt{M_N}, d_P)$. To cover $\phi_{(j,k)} \circ \phi_{(k+1)} \circ \dots \circ \phi_{(N)}(\mathcal{F})$ one needs at most $m_k N_{[\]}(\mathcal{F}, \varepsilon/2^{N-k} \sqrt{m_{k+1} \dots m_N}, d_P)$ brackets. We have proved that

$$\begin{aligned} N_{[\]}(\mathcal{F}_0, \varepsilon, d_P) &\leq (N_0 + 1) N_{[\]}(\mathcal{F}, \varepsilon/2^{N_0} \sqrt{M_{N_0}}, d_P), \\ N_{[\]}(\mathcal{H}_0, \varepsilon, d_P) &\leq S_{N_0} N_{[\]}(\mathcal{F}, \varepsilon/2^{N_0} \sqrt{M_{N_0}}, d_P). \end{aligned}$$

Therefore $\mathcal{F}_0, \mathcal{H}_0$ satisfy (BR) with power r_0 and constant $2^{r_0 N_0} M_{N_0}^{r_0} b_0$. \square

Step 2. By (3.2) we have

$$\alpha_n^{(N)}(f) = \sum_{j=1}^{m_N} \frac{P(A_j^{(N)})}{\mathbb{P}_n^{(N-1)}(A_j^{(N)})} \alpha_n^{(N-1)}(\phi_{(j,N)}f) = \alpha_n^{(N-1)}(\phi_{(N)}f) + \Gamma_n^{(N)}(f), \quad (3.13)$$

$$\Gamma_n^{(N)}(f) = \sum_{j=1}^{m_N} q_n(j, N) \alpha_n^{(N-1)}(\phi_{(j,N)}f), \quad q_n(j, N) = \frac{P(A_j^{(N)})}{\mathbb{P}_n^{(N-1)}(A_j^{(N)})} - 1.$$

Under the convention that $\phi_{(N+1)} \circ \phi_{(N)} = id$, iterating (3.13) leads to

$$\begin{aligned}\alpha_n^{(N)}(f) &= \alpha_n^{(0)}(\phi_{(1)} \circ \dots \circ \phi_{(N)}f) + F_n^{(N)}(f), \\ F_n^{(N)}(f) &= \sum_{k=1}^N \Gamma_n^{(k)}(\phi_{(k+1)} \circ \dots \circ \phi_{(N)}f).\end{aligned}$$

Clearly the terms $\Gamma_n^{(k)}$ carry out some bias and variance distortion. However $\alpha_n^{(0)}(\mathcal{F}_{(N)})$ is the main contribution to $\alpha_n^{(N)}(\mathcal{F})$ and $F_n^{(N)}(\mathcal{F})$ is an error process.

Lemma 2. *Consider the sequence v_n defined at Theorem 2.1. If \mathcal{F} satisfies (VC) or (BR) then there exists $C_0 < +\infty$ such that we almost surely have, for all n large enough, $\max_{0 \leq N \leq N_0} \|F_n^{(N)}\|_{\mathcal{F}} \leq C_0 L \circ L(n) / \sqrt{n}$. Moreover, for any $\zeta > 0$ and $\theta > 0$ there exists $n_3(\zeta, \theta)$ such that we have, for all $n > n_3(\zeta, \theta)$,*

$$\mathbb{P} \left(\max_{0 \leq N \leq N_0} \|F_n^{(N)}\|_{\mathcal{F}} > \zeta v_n \right) \leq \frac{1}{2n^\theta}.$$

Proof. (i) Let us apply Proposition 2 to \mathcal{F} and, thanks to Lemma 1, to \mathcal{H}_0 and $\mathcal{H}_{(N)}$. The following statements are almost surely true, for all n large enough. On the one hand,

$$\max_{1 \leq N \leq N_0} \max_{1 \leq j \leq m_N} \left| \alpha_n^{(N-1)}(A_j^{(N)}) \right| \leq b_n = 2\sigma_{\mathcal{F}}\kappa_{N_0} \sqrt{L \circ L(n)}. \quad (3.14)$$

On the other hand, having $\sigma_{\mathcal{H}_0} \leq \sigma_{\mathcal{F}}$ by (3.12),

$$\max_{1 \leq N \leq N_0} \max_{1 \leq k \leq N} \max_{1 \leq j \leq m_k} \left| \alpha_n^{(k-1)}(\phi_{(j,k)} \circ \phi_{(k+1)} \circ \dots \circ \phi_{(N)}f) \right| \leq \max_{1 \leq k \leq N_0} \left\| \alpha_n^{(k-1)} \right\|_{\mathcal{H}_0} \leq b_n.$$

By (3.14), $q_n(j, N) = 1/(1 + \alpha_n^{(N-1)}(A_j^{(N)})/P(A_j^{(N)})\sqrt{n}) - 1$ satisfies

$$\max_{1 \leq N \leq N_0} \max_{1 \leq j \leq m_N} |q_n(j, N)| \frac{\sqrt{n}}{b_n} \delta_{(N_0)} \leq 2 \quad (3.15)$$

which implies

$$\left\| F_n^{(N)} \right\|_{\mathcal{F}} \leq \sum_{k=1}^N \max_{1 \leq j \leq m_k} |q_n(j, k)| \sum_{j=1}^{m_k} \left| \alpha_n^{(k-1)}(\phi_{(j,k)} \circ \phi_{(k+1)} \circ \dots \circ \phi_{(N)}f) \right| \quad (3.16)$$

$$\max_{1 \leq N \leq N_0} \left\| F_n^{(N)} \right\|_{\mathcal{F}} \leq \frac{2b_n}{\sqrt{n}\delta_{(N_0)}} S_{N_0} \max_{1 \leq k \leq N_0} \left\| \alpha_n^{(k-1)} \right\|_{\mathcal{H}_0} \leq \frac{2b_n^2 S_{N_0}}{\sqrt{n}\delta_{(N_0)}}.$$

The almost sure result then holds with $C_0 = 8\sigma_{\mathcal{F}}^2 \kappa_{N_0}^2 S_{N_0} / \delta_{(N_0)}$.

(ii) We now work on the event B_{n, N_0} of (3.1). There obviously exists n_1 such that if $n > n_1$ then $S_{N_0}(1 - \delta_{(N_0)})^n \leq 1/4n^\theta$. We can also find $\kappa > 0$ such that $n^{2\kappa}/\sqrt{n} = o(v_n)$ as $n \rightarrow +\infty$. Therefore, whatever $\zeta > 0$ there exists

$n_2(\kappa, S_{N_0}, \zeta, \mathcal{F}, P)$ such that $\zeta v_n > 2S_{N_0} n^{2\kappa} / \delta_{(N_0)} \sqrt{n}$ for any $n \geq n_2$. Choosing $n \geq \max(n_1, n_2)$ we deduce as for (3.15) and (3.16) that

$$\begin{aligned}
& \mathbb{P} \left(\max_{0 \leq N \leq N_0} \|F_n^{(N)}\|_{\mathcal{F}} > \zeta v_n \right) \\
& \leq \mathbb{P} \left(S_{N_0} \max_{1 \leq N \leq N_0} \left(\left\| \alpha_n^{(N-1)} \right\|_{\mathcal{H}_0} \max_{1 \leq j \leq m_N} |q_n(j, N)| \right) > \zeta v_n \right) \\
& \leq \mathbb{P} \left(\left(\max_{1 \leq N \leq N_0} \max_{1 \leq j \leq m_N} |q_n(j, N)| \right) > \frac{2n^\kappa}{\delta_{(N_0)} \sqrt{n}} \right) + \mathbb{P} \left(\max_{1 \leq N \leq N_0} \left\| \alpha_n^{(N-1)} \right\|_{\mathcal{H}_0} > n^\kappa \right) \\
& \leq 2\mathbb{P} \left(\max_{1 \leq N \leq N_0} \left\| \alpha_n^{(N-1)} \right\|_{\mathcal{H}_0} > n^\kappa \right).
\end{aligned}$$

By Proposition 3 we see that under (VC) or (BR) the latter probability can be made less than $1/8n^\theta$ for any $n > n_3(\zeta, \theta)$ and $n_3(\zeta, \theta)$ large enough. Clearly $n_3(\zeta, \theta)$ depends on ζ, θ, n_1, n_2 and on the entropy of \mathcal{H}_0 thus all constants in Lemma 1 and Proposition 3 are involved. \square

Step 3. Fix $\theta > 0$. By Lemma 1 we can apply Berthet and Mason to \mathcal{F}_0 , which ensures the following Gaussian approximation. For some constant $c_\theta(\mathcal{F}_0, P) > 0$ and $n_\theta(\mathcal{F}_0, P) > 0$ we can build on a probability space $(\Omega, \mathcal{T}, \mathbb{P})$ a version of the sequence $\{X_n\}$ of independent random variables with law P and a sequence $\{\mathbb{G}_n^{(0)}\}$ of coupling versions of $\mathbb{G}^{(0)}$ in such a way that, for all $n \geq n_\theta(\mathcal{F}_0, P)$,

$$\mathbb{P} \left(\left\| \alpha_n^{(0)} - \mathbb{G}_n^{(0)} \right\|_{\mathcal{F}_0} > c_\theta(\mathcal{F}_0, P) v_n \right) \leq \frac{1}{2n^\theta}. \quad (3.17)$$

Keep in mind that constants n_θ and c_θ only depend on the uniform boundedness and the entropy of \mathcal{F}_0 . By choosing $\theta > 1$, $d_\theta > c_\theta(\mathcal{F}_0, P)$ then applying Borel-Cantelli lemma to (3.17), it almost surely holds, for all n large enough,

$$\left\| \alpha_n^{(0)} - \mathbb{G}_n^{(0)} \right\|_{\mathcal{F}_0} < d_\theta v_n. \quad (3.18)$$

Step 4. Let $\theta_0 > 0$. We work under the event B_{n, N_0} of (3.1) with a probability at least $1 - 1/4n^{\theta_0}$ provided that $n > n_1$. The process $\mathbb{G}^{(0)}$ being linear on \mathcal{F} we see that the recursive definition (1.11) applied to the version $\mathbb{G}_n^{(0)}$ of $\mathbb{G}^{(0)}$ reads $\mathbb{G}_n^{(N)}(f) = \mathbb{G}_n^{(N-1)}(\phi_{(N)} f)$. This combined to (3.13) readily gives

$$\begin{aligned}
& \max_{1 \leq N \leq N_0} \left\| \alpha_n^{(N)} - \mathbb{G}_n^{(N)} \right\|_{\mathcal{F}} \\
& = \max_{1 \leq N \leq N_0} \left\| \alpha_n^{(N-1)}(\phi_{(N_0)} f) - \mathbb{G}_n^{(N-1)}(\phi_{(N_0)} f) + \Gamma_n^{(N)}(f) \right\|_{\mathcal{F}} \\
& = \max_{1 \leq N \leq N_0} \left\| \alpha_n^{(0)}(\phi_{(1)} \circ \dots \circ \phi_{(N)} f) - \mathbb{G}_n^{(0)}(\phi_{(1)} \circ \dots \circ \phi_{(N)} f) + F_n^{(N)}(f) \right\|_{\mathcal{F}} \\
& \leq \left\| \alpha_n^{(0)} - \mathbb{G}_n^{(0)} \right\|_{\mathcal{F}_0} + \max_{0 \leq N \leq N_0} \|F_n^{(N)}\|_{\mathcal{F}}. \quad (3.19)
\end{aligned}$$

Remind that $v_n > L(n)/\sqrt{n}$ and Lemma 2 holds. By (3.18) and (3.19) we almost surely have, for all n large enough and $d_0 = 2d_{\theta_0}$,

$$\max_{1 \leq N \leq N_0} \left\| \alpha_n^{(N)} - \mathbb{G}_n^{(N)} \right\|_{\mathcal{F}} \leq d_{\theta_0} v_n + C_0 \frac{L \circ L(n)}{\sqrt{n}} \leq d_0 v_n.$$

By Lemmas 1 and 2, (3.17) and (3.19), for $n_0 > \max(n_1, n_3(\zeta, \theta_0), n_{\theta_0}(\mathcal{F}_0, P))$ and $d_0 > c_{\theta_0}(\mathcal{F}_0, P) + \zeta$ we have, for all $n \geq n_0$,

$$\mathbb{P} \left(\max_{1 \leq N \leq N_0} \left\| \alpha_n^{(N)} - \mathbb{G}_n^{(N)} \right\|_{\mathcal{F}} > d_0 v_n \right) \leq \frac{1}{2n\theta_0} + \mathbb{P} \left(\max_{0 \leq N \leq N_0} \|F_n^{(N)}\|_{\mathcal{F}} > \zeta v_n \right) \leq \frac{1}{n\theta_0}.$$

To conclude observe that the parameters $N_0, M, M_{N_0}, S_{N_0}, \Delta_{N_0}, \delta_{(N_0)}, \theta_0, \nu_0, c_0, r_0, b_0$ have been used at one or several steps to finally define n_0 and d_0 .

3.6 Proof of Proposition 5

Theorem 2.1 implies, for $f \in \mathcal{F}$,

$$\mathbb{P}_n^{(N)}(f) - P(f) = \frac{1}{\sqrt{n}} \mathbb{G}_n^{(N)}(f) + \frac{1}{\sqrt{n}} \mathbb{R}_n^{(N)}(f) \quad (3.20)$$

where $\mathbb{G}_n^{(N)}$ is a sequence of versions of the centered Gaussian process $\mathbb{G}^{(N)}$ from (1.11) and the random sequence $r_n^{(N)} = \|\mathbb{R}_n^{(N)}\|_{\mathcal{F}}$ satisfies

$$r_n^{(N)} \leq \left\| \mathbb{G}_n^{(N)} \right\|_{\mathcal{F}} + \left\| \alpha_n^{(N)} \right\|_{\mathcal{F}} \leq \left\| \mathbb{G}_n^{(N)} \right\|_{\mathcal{F}} + 2M\sqrt{n}, \quad \lim_{n \rightarrow +\infty} \frac{r_n^{(N)}}{v_n} \leq d_0 \quad a.s.$$

We have to be a little careful with the expectation, variance and covariance of the coupling error process $\mathbb{R}_n^{(N)}$.

Step 1. Since $\mathbb{G}_n^{(N)}(f)$ is centered the bias is controlled by

$$\sup_{f \in \mathcal{F}} \frac{\sqrt{n}}{v_n} \left| \mathbb{E} \left(\mathbb{P}_n^{(N)}(f) \right) - P(f) \right| = \sup_{f \in \mathcal{F}} \left| \frac{1}{v_n} \mathbb{E} \left(\mathbb{R}_n^{(N)}(f) \right) \right| \leq \mathbb{E} \left(\frac{r_n^{(N)}}{v_n} \right). \quad (3.21)$$

Write $a_n = \sqrt{K \log n}$ where $K > 0$ and $\theta_0 > 1$ from Theorem 2.1 can be chosen as large as needed. Then, for $\theta > 1$, $\varepsilon > 0$ and $k \in \mathbb{N}^*$ consider the events

$$\begin{aligned} A_n &= \left\{ r_n^{(N)} \leq (d_0 + \varepsilon)v_n \right\}, \quad B_n = \left\{ \left\| \mathbb{G}_n^{(N)} \right\|_{\mathcal{F}} \leq a_n \right\}, \\ C_{n,k} &= \left\{ \theta^{k-1} a_n < \left\| \mathbb{G}_n^{(N)} \right\|_{\mathcal{F}} \leq \theta^k a_n \right\}. \end{aligned}$$

By Theorem 2.1, $\mathbb{P}(A_n^c) < 1/n^{\theta_0}$ and $v_n > a_n/\sqrt{n}$ for all n large enough, hence

$$\begin{aligned} \frac{1}{v_n} \mathbb{E} \left(r_n^{(N)} \right) &= \mathbb{E} \left(\frac{r_n^{(N)}}{v_n} 1_{A_n} \right) + \mathbb{E} \left(\frac{r_n^{(N)}}{v_n} 1_{A_n^c \cap B_n} \right) + \mathbb{E} \left(\frac{r_n^{(N)}}{v_n} 1_{A_n^c \cap B_n^c} \right) \\ &\leq d_0 + \varepsilon + \frac{a_n + 2M\sqrt{n}}{v_n} \mathbb{P}(A_n^c) + \mathbb{E} \left(\frac{r_n^{(N)}}{v_n} 1_{B_n^c} \right) \\ &\leq d_0 + 2\varepsilon + \sum_{k=1}^{+\infty} \mathbb{E} \left(\frac{r_n^{(N)}}{v_n} 1_{C_{n,k}} \right). \end{aligned}$$

By Theorems 7 and 9, $\mathbb{G}^{(N)}(f)$ is a centered Gaussian process indexed by \mathcal{F} such that, under (VC) or (BR),

$$\begin{aligned} \mathbb{E} \left(\left\| \mathbb{G}^{(N)} \right\|_{\mathcal{F}} \right) &< +\infty, \quad \sup_{f \in \mathcal{F}} \mathbb{V}(\mathbb{G}^{(N)}(f)) \leq \sigma_{\mathcal{F}}^2 < +\infty, \\ \mathbb{E} \left(\left\| \mathbb{G}^{(N)} \right\|_{\mathcal{F}}^2 \right) &\leq C_{\mathcal{F}}^2 = \sigma_{\mathcal{F}}^2 + \mathbb{E} \left(\left\| \mathbb{G}^{(N)} \right\|_{\mathcal{F}} \right)^2 < +\infty \end{aligned} \quad (3.22)$$

thus, by Borell's inequality, for any version $\mathbb{G}_n^{(N)}$ of $\mathbb{G}^{(N)}$,

$$\mathbb{P} \left(\left\| \mathbb{G}_n^{(N)} \right\|_{\mathcal{F}} > \lambda \right) \leq 2 \exp \left(-\frac{\lambda^2}{8C_{\mathcal{F}}^2} \right). \quad (3.23)$$

Therefore we have, since $\theta > 1$ and $v_n > 4M/\sqrt{n} > 2a_n/n$ for n large enough,

$$\begin{aligned} \mathbb{E} \left(\frac{r_n^{(N)}}{v_n} 1_{C_{n,k}} \right) &\leq \frac{\theta^k a_n + 2M\sqrt{n}}{v_n} \mathbb{P}(C_k) \leq \theta^k n \mathbb{P} \left(\left\| \mathbb{G}_n^{(N)} \right\|_{\mathcal{F}} > \theta^{k-1} a_n \right) \\ &\leq 2\theta^k n \exp \left(-\frac{(\theta^{k-1} a_n)^2}{8C_{\mathcal{F}}^2} \right) \end{aligned}$$

and the following series is converging to an arbitrarily small sum,

$$\begin{aligned} \sum_{k=1}^{+\infty} \mathbb{E} \left(\frac{r_n^{(N)}}{v_n} 1_{C_{n,k}} \right) &\leq 2n \exp \left(-\frac{a_n^2}{8C_{\mathcal{F}}^2} \right) \sum_{k=1}^{+\infty} \theta^k \exp \left(-\left(\frac{\theta^{2(k-1)} - 1}{8C_{\mathcal{F}}^2} \right) a_n^2 \right) \\ &\leq n \exp \left(-\frac{K \log n}{8C_{\mathcal{F}}^2} \right) \sum_{k=1}^{+\infty} 2e\theta^k \exp \left(-\theta^{2(k-1)} \right) \leq \frac{1}{n^\delta} \end{aligned}$$

where $\delta < K/8C_{\mathcal{F}}^2 - 1$. It follows that (3.21) is ultimately bounded by d_0 .

Step 2. Starting from (3.20) and the bias and variance decomposition, the quadratic risk is in turn controlled by

$$\begin{aligned} &\mathbb{E} \left(\left(\mathbb{P}_n^{(N)}(f) - P(f) \right)^2 \right) - \frac{1}{n} \mathbb{V} \left(\mathbb{G}^{(N)}(f) \right) \\ &= \left| \mathbb{E} \left(\mathbb{P}_n^{(N)}(f) \right) - P(f) \right|^2 + \frac{1}{n} \mathbb{V} \left(\mathbb{R}_n^{(N)}(f) \right) + \frac{2}{n} \text{Cov} \left(\mathbb{G}_n^{(N)}(f), \mathbb{R}_n^{(N)}(f) \right). \end{aligned} \quad (3.24)$$

(i) By step 1, the first right-hand term is the squared bias, of order $d_0^2 v_n^2/n$. Concerning the second right-hand term in (3.24), we bound $\mathbb{E}(\mathbb{R}_n^{(N)}(f)^2)$. Fix $\varepsilon > 0$ and assume that n is large enough for the following statements. By setting $s_n^{(N)} = (r_n^{(N)})^2$ then using $v_n > a_n/\sqrt{n}$, $a_n = K\sqrt{\log n} < \sqrt{n}$ we get, for $\theta_0 = 2$,

$$\begin{aligned} \frac{1}{v_n^2} \sup_{f \in \mathcal{F}} \mathbb{E} \left(\mathbb{R}_n^{(N)}(f)^2 \right) &\leq \mathbb{E} \left(\frac{s_n^{(N)}}{v_n^2} 1_{A_n} \right) + \mathbb{E} \left(\frac{s_n^{(N)}}{v_n^2} 1_{A_n^c \cap B_n} \right) + \mathbb{E} \left(\frac{s_n^{(N)}}{v_n^2} 1_{A_n^c \cap B_n^c} \right) \\ &\leq (d_0 + \varepsilon)^2 + \left(\frac{a_n + 2M\sqrt{n}}{v_n} \right)^2 \mathbb{P}(A_n^c) + \mathbb{E} \left(\frac{s_n^{(N)}}{v_n^2} 1_{B_n^c} \right) \\ &\leq (d_0 + 2\varepsilon)^2 + \left(\frac{3M\sqrt{n}}{\log n} \right)^2 \frac{1}{n^2} + \sum_{k=1}^{+\infty} \mathbb{E} \left(\frac{s_n^{(N)}}{v_n^2} 1_{C_{n,k}} \right) \\ &\leq (d_0 + 3\varepsilon)^2 + \sum_{k=1}^{+\infty} \theta^{2k} n^2 \mathbb{P} \left(\left\| \mathbb{G}_n^{(N)} \right\|_{\mathcal{F}} > \theta^{k-1} a_n \right) \leq (d_0 + 4\varepsilon)^2 \end{aligned}$$

where the series is equal to its first term $n^2 \exp(-a_n^2/8C_{\mathcal{F}}^2)$ times a convergent series, by using (3.23) as for step 1 with $K > 16C_{\mathcal{F}}^2$. We have shown that

$$\limsup_{n \rightarrow +\infty} \frac{1}{v_n^2} \sup_{f \in \mathcal{F}} \mathbb{V} \left(\mathbb{R}_n^{(N)}(f) \right) \leq \limsup_{n \rightarrow +\infty} \frac{1}{v_n^2} \mathbb{E} \left(s_n^{(N)} \right) \leq d_0^2. \quad (3.25)$$

(ii) Concerning the covariance term in (3.24) it holds

$$\begin{aligned} \frac{1}{v_n} \left| \text{Cov} \left(\mathbb{G}_n^{(N)}(f), \mathbb{R}_n^{(N)}(f) \right) \right| &= \frac{1}{v_n} \left| \mathbb{E} \left(\mathbb{G}_n^{(N)}(f) \mathbb{R}_n^{(N)}(f) \right) \right| \\ &\leq T_{A_n}(f) + T_{A_n^c \cap B_n}(f) + T_{A_n^c \cap B_n^c}(f) \end{aligned}$$

where

$$T_D(f) = \mathbb{E} \left(\left| \mathbb{G}_n^{(N)}(f) \right| \frac{r_n^{(N)}}{v_n} 1_D \right), \quad D \in \{A_n, A_n^c \cap B_n, A_n^c \cap B_n^c\}.$$

As a consequence of (3.22),

$$T_{A_n}(f) \leq \mathbb{E} \left(\left| \mathbb{G}_n^{(N)}(f) \right| (d_0 + \varepsilon) 1_{A_n} \right) \leq (d_0 + \varepsilon) \sigma_{\mathcal{F}} \mathbb{E}(|\mathcal{N}(0, 1)|).$$

By using again $\mathbb{P}(A_n^c) < 1/n^2$ we see that

$$T_{A_n^c \cap B_n}(f) \leq \mathbb{E} \left(a_n \left(\frac{2M\sqrt{n} + a_n}{v_n} \right) 1_{A_n^c \cap B_n} \right) \leq a_n \left(\frac{3M\sqrt{n}}{v_n} \right) \frac{1}{n^2} \leq \varepsilon.$$

Lastly, for $g_n^{(N)} = \left\| \mathbb{G}_n^{(N)} \right\|_{\mathcal{F}}$, K large and all n large enough it holds, by (3.23),

$$\begin{aligned} T_{A_n^c \cap B_n^c}(f) &\leq \mathbb{E} \left(g_n^{(N)} \left(\frac{2M\sqrt{n} + g_n^{(N)}}{v_n} \right) 1_{B_n^c} \right) \\ &= \sum_{k=1}^{+\infty} \mathbb{E} \left(g_n^{(N)} \left(\frac{2M\sqrt{n} + g_n^{(N)}}{v_n} \right) 1_{C_{n,k}} \right) \\ &\leq \sum_{k=1}^{+\infty} \theta^{2k} a_n^2 n \mathbb{P} \left(\left\| \mathbb{G}_n^{(N)} \right\|_{\mathcal{F}} > \theta^{k-1} a_n \right) \leq \varepsilon. \end{aligned}$$

The above upper bounds do not depend on f . By (3.24) and (3.25) this implies

$$\limsup_{n \rightarrow +\infty} \frac{n}{v_n} \sup_{f \in \mathcal{F}} \left| \mathbb{E} \left((\mathbb{P}_n^{(N)}(f) - P(f))^2 \right) - \frac{1}{n} \mathbb{V} \left(\mathbb{G}^{(N)}(f) \right) \right| \leq \sqrt{\frac{8}{\pi}} d_0 \sigma_{\mathcal{F}}.$$

Step 3. Let extend Step 2 to the covariance. By Step 1 we have, for all n large,

$$\begin{aligned} &\left| \text{Cov} \left(\mathbb{P}_n^{(N)}(f), \mathbb{P}_n^{(N)}(g) \right) - \mathbb{E} \left(\left(\mathbb{P}_n^{(N)}(f) - P(f) \right) \left(\mathbb{P}_n^{(N)}(g) - P(g) \right) \right) \right| \\ &= \left| \mathbb{E} \left(\mathbb{P}_n^{(N)}(f) \right) - P(f) + \mathbb{E} \left(\mathbb{P}_n^{(N)}(g) \right) - P(g) \right| < 2d_0 \frac{v_n^2}{n}. \end{aligned}$$

Now, by the upper bounds computed at (i) and (ii) of Step 2,

$$\begin{aligned} &\left| \mathbb{E} \left((\mathbb{P}_n^{(N)}(f) - P(f))(\mathbb{P}_n^{(N)}(g) - P(g)) \right) - \frac{1}{n} \text{Cov} \left(\mathbb{G}^{(N)}(f), \mathbb{G}^{(N)}(g) \right) \right| \\ &\leq \frac{1}{n} \mathbb{E} \left(\left| \mathbb{G}_n^{(N)}(f) \mathbb{R}_n^{(N)}(g) \right| \right) + \frac{1}{n} \mathbb{E} \left(\left| \mathbb{G}_n^{(N)}(g) \mathbb{R}_n^{(N)}(f) \right| \right) + \frac{1}{n} \mathbb{E} \left(\left| \mathbb{R}_n^{(N)}(f) \mathbb{R}_n^{(N)}(g) \right| \right) \\ &\leq \frac{2}{n} \sup_{f \in \mathcal{F}} \mathbb{E} \left(\left| \mathbb{G}_n^{(N)}(f) \right| r_n^{(N)} \right) + \frac{1}{n} \mathbb{E} \left(s_n^{(N)} \right) \\ &\leq \frac{2}{n} (d_0 + \varepsilon) \sigma_{\mathcal{F}} \sqrt{\frac{2}{\pi}} v_n + \frac{1}{n} (d_0 + \varepsilon)^2 v_n^2. \end{aligned}$$

3.7 Proof of Proposition 6.

Fix $N_0 \in \mathbb{N}$ and \mathcal{L} . Let apply Theorem 2.1 from which we also use n_0 and v_n . We have, for all $0 \leq N \leq N_0$, $\varphi \in \mathcal{L}$, $x \in \mathbb{R}$ and $n \geq n_0$

$$\begin{aligned} \mathbb{P} \left(\varphi(\alpha_n^{(N)}) \leq x \right) &\leq \frac{1}{n^2} + \mathbb{P} \left(\left\{ \varphi(\alpha_n^{(N)}) \leq x \right\} \cap \left\{ \left\| \alpha_n^{(N)} - \mathbb{G}_n^{(N)} \right\|_{\mathcal{F}} < c_0 v_n \right\} \right) \\ &\leq \frac{1}{n^2} + \mathbb{P} \left(\varphi(\mathbb{G}_n^{(N)}) \leq x + c_0 C_1 v_n \right) \\ &\leq \frac{1}{n^2} + \mathbb{P} \left(\varphi(\mathbb{G}_n^{(N)}) \leq x \right) + c_0 C_1 C_2 v_n \end{aligned}$$

and

$$\begin{aligned}
& \mathbb{P} \left(\varphi(\mathbb{G}_n^{(N)}) \leq x - c_0 C_1 v_n \right) \\
& \leq \frac{1}{n^2} + \mathbb{P} \left(\left\{ \varphi(\mathbb{G}_n^{(N)}) \leq x - c_0 C_1 v_n \right\} \cap \left\{ \left\| \alpha_n^{(N)} - \mathbb{G}_n^{(N)} \right\|_{\mathcal{F}} < c_0 v_n \right\} \right) \\
& \leq \frac{1}{n^2} + \mathbb{P} \left(\varphi(\alpha_n^{(N)}) \leq \varphi(\mathbb{G}_n^{(N)}) + c_0 C_1 v_n \leq x \right)
\end{aligned}$$

so that

$$\mathbb{P} \left(\varphi(\alpha_n^{(N)}) \leq x \right) \geq \mathbb{P} \left(\varphi(\mathbb{G}_n^{(N)}) \leq x \right) - c_0 C_1 C_2 v_n - \frac{1}{n^2}.$$

This establishes the second statement of Proposition 6 provided $c_1 > c_0$ and $n \geq n_1 \geq n_0$ where n_1 is large enough to have $(c_1 - c_0)v_n > n^{-2}$. The first statement coincides with the special case $\mathcal{L} = \{\varphi_f : f \in \mathcal{F}_0\}$ where $\varphi_f(g) = g(f)$ are pointwise projectors and we then have a Lipschitz constant $C_1 = 1$ whereas $\varphi_f(\mathbb{G}_n^{(N)}) = \mathbb{G}_n^{(N)}(f)$ has a Gaussian density bounded by

$$\frac{1}{\sqrt{2\pi\mathbb{V}(\mathbb{G}_n^{(N)}(f))}} \leq C_2 = \frac{1}{\sqrt{2\pi}\sigma_0} < +\infty.$$

4 Proofs concerning the limiting process

4.1 Proof of Proposition 7

Step 1. Let us first relate $\mathbb{G}^{(N)}(\mathcal{F})$ from (1.11) to $\mathbb{G}(\mathcal{F}) = \mathbb{G}^{(0)}(\mathcal{F})$ from (1.10) by means of the vectors $\Phi_k^{(N)}(f)$ introduced at (2.3) before Proposition 7.

Lemma 3. For all $N \in \mathbb{N}^*$ and $f \in \mathcal{F}$ it holds

$$\mathbb{G}^{(N)}(f) = \mathbb{G}(f) - \sum_{k=1}^N \Phi_k^{(N)}(f)^t \cdot \mathbb{G} \left[\mathcal{A}^{(k)} \right]$$

Proof. The formula is true for $N = 0$. Assume that it is the case for $N \geq 0$. Recall that sets $A \in \mathcal{A}_{\cup}^{(N_0)}$ from (1.9) are identified to $f = 1_A$. By (1.11),

$$\begin{aligned}
\mathbb{G}^{(N+1)}(f) &= \mathbb{G}^{(N)}(f) - \mathbb{E} \left[f \mid \mathcal{A}^{(N+1)} \right]^t \cdot \mathbb{G}^{(N)} \left[\mathcal{A}^{(N+1)} \right] \\
&= \mathbb{G}(f) - \sum_{k=1}^N \Phi_k^{(N)}(f)^t \cdot \mathbb{G} \left[\mathcal{A}^{(k)} \right] \\
&\quad - \sum_{j=1}^{m_{N+1}} \mathbb{E}(f \mid A_j^{(N+1)}) \left(\mathbb{G}(A_j^{(N+1)}) - \sum_{k=1}^N \Phi_k^{(N)}(A_j^{(N+1)})^t \cdot \mathbb{G} \left[\mathcal{A}^{(k)} \right] \right) \\
&= \mathbb{G}(f) - \mathbb{E} \left[f \mid \mathcal{A}^{(N+1)} \right]^t \cdot \mathbb{G} \left[\mathcal{A}^{(N+1)} \right] \\
&\quad - \sum_{k=1}^N \left[\Phi_k^{(N)}(f) - \Phi_k^{(N)}[\mathcal{A}^{(N+1)}] \cdot \mathbb{E}[f \mid \mathcal{A}^{(N+1)}] \right]^t \cdot \mathbb{G} \left[\mathcal{A}^{(k)} \right]
\end{aligned}$$

where the $m_k \times m_{N+1}$ matrix $\Phi_k^{(N)}[\mathcal{A}^{(N+1)}] = (\Phi_k^{(N)}(A_1^{(N+1)}), \dots, \Phi_k^{(N)}(A_{m_{N+1}}^{(N+1)}))$ satisfies

$$\sum_{j=1}^{m_{N+1}} \mathbb{E}(f | A_j^{(N+1)}) \Phi_k^{(N)}(A_j^{(N+1)})^t = \left[\Phi_k^{(N)}[\mathcal{A}^{(N+1)}] \cdot \mathbb{E}[f | \mathcal{A}^{(N+1)}] \right]^t.$$

Now observe that $\Phi_{N+1}^{(N+1)}(f) = \mathbb{E}[f | \mathcal{A}^{(N+1)}]$ since no index L matches in the sum (2.3) defining $\Phi_k^{(N)}$ when $N = k$. It remains to show that

$$\Phi_k^{(N)}(f) - \Phi_k^{(N)}[\mathcal{A}^{(N+1)}] \cdot \mathbb{E}[f | \mathcal{A}^{(N+1)}] = \Phi_k^{(N+1)}(f). \quad (4.1)$$

For $1 \leq k \leq N$ and $1 \leq j \leq m_{N+1}$ we have

$$\begin{aligned} \Phi_k^{(N)}(A_j^{(N+1)}) &= \mathbb{E} \left[A_j^{(N+1)} | \mathcal{A}^{(k)} \right] + \\ &\sum_{\substack{1 \leq L \leq N-k \\ k < l_1 < \dots < l_L \leq N}} (-1)^L \mathbf{P}_{\mathcal{A}^{(l_1)} | \mathcal{A}^{(k)}} \cdot \mathbf{P}_{\mathcal{A}^{(l_2)} | \mathcal{A}^{(l_1)}} \cdots \mathbf{P}_{\mathcal{A}^{(l_L)} | \mathcal{A}^{(l_{L-1})}} \cdot \mathbb{E} \left[A_j^{(N+1)} | \mathcal{A}^{(l_L)} \right] \end{aligned}$$

where, for $l = k, k+1, \dots, N$ the vector

$$\mathbb{E} \left[A_j^{(N+1)} | \mathcal{A}^{(l)} \right] = \left(\frac{P(A_j^{(N+1)} \cap A_1^{(l)})}{P(A_1^{(l)})}, \dots, \frac{P(A_j^{(N+1)} \cap A_{m_l}^{(l)})}{P(A_{m_l}^{(l)})} \right)^t$$

is also the j -th column of $\mathbf{P}_{\mathcal{A}^{(N+1)} | \mathcal{A}^{(l)}}$. Therefore, by turning L into $L' = L+1$,

$$\begin{aligned} &- \Phi_k^{(N)}[\mathcal{A}^{(N+1)}] \cdot \mathbb{E} \left[f | \mathcal{A}^{(N+1)} \right] \\ &= - \sum_{j=1}^{m_{N+1}} \mathbb{E}(f | A_j^{(N+1)}) \Phi_k^{(N)}(A_j^{(N+1)}) \\ &= - \sum_{j=1}^{m_{N+1}} \mathbb{E}(f | A_j^{(N+1)}) \mathbb{E} \left[A_j^{(N+1)} | \mathcal{A}^{(k)} \right] \\ &+ \sum_{j=1}^{m_{N+1}} \mathbb{E}(f | A_j^{(N+1)}) \sum_{\substack{1 \leq L \leq N-k \\ k < l_1 < \dots < l_L \leq N}} (-1)^{L+1} \mathbf{P}_{\mathcal{A}^{(l_1)} | \mathcal{A}^{(k)}} \cdots \mathbf{P}_{\mathcal{A}^{(l_L)} | \mathcal{A}^{(l_{L-1})}} \cdot \mathbb{E} \left[A_j^{(N+1)} | \mathcal{A}^{(l_L)} \right] \\ &= (-1)^1 \mathbf{P}_{\mathcal{A}^{(N+1)} | \mathcal{A}^{(k)}} \mathbb{E} \left[f | \mathcal{A}^{(N+1)} \right] \\ &+ \sum_{\substack{1 \leq L' \leq N+1-k \\ k < l_1 < \dots < l_{L'} = N+1}} (-1)^{L'} \mathbf{P}_{\mathcal{A}^{(l_1)} | \mathcal{A}^{(k)}} \cdots \mathbf{P}_{\mathcal{A}^{(l_{L'-1})} | \mathcal{A}^{(l_{L'-2})}} \cdot \mathbf{P}_{\mathcal{A}^{(N+1)} | \mathcal{A}^{(l_{L'-1})}} \cdot \mathbb{E} \left[f | \mathcal{A}^{(N+1)} \right] \end{aligned}$$

where all terms are different from those in

$$\Phi_k^{(N)}(f) = \mathbb{E} \left[f | \mathcal{A}^{(k)} \right] + \sum_{\substack{1 \leq L' \leq N+1-k \\ k < l_1 < \dots < l_{L'} < N+1}} (-1)^{L'} \mathbf{P}_{\mathcal{A}^{(l_1)} | \mathcal{A}^{(k)}} \cdots \mathbf{P}_{\mathcal{A}^{(l_{L'})} | \mathcal{A}^{(l_{L'-1})}} \cdot \mathbb{E} \left[f | \mathcal{A}^{(l_{L'})} \right].$$

Having collected all terms of $\Phi_k^{(N)}(f)$ in (2.3), this establishes (4.1). The proof is completed by induction. \square

The functions $\Phi_k^{(N)}$ and the process \mathbb{G} are linear, hence Lemma 3 implies that $\mathbb{G}^{(N)}$ is a linear process. Moreover $\mathbb{G}(f)$ and $\mathbb{G}[\mathcal{A}^{(k)}]$ being centered Gaussian, Lemma 3 proves that $\mathbb{G}^{(N)}(f)$ is a centered Gaussian random variable.

Step 2. To compute the covariance of $\mathbb{G}^{(N)}(\mathcal{F})$ we need the following properties. Recall that $\mathbf{P}_{\mathcal{A}^{(k)}|\mathcal{A}^{(k)}} = \text{Id}_{m_k}$ is the identity matrix of \mathbb{R}^{m_k} .

Lemma 4. For $1 \leq k, l \leq N$ and $f \in \mathcal{F}$ we have

$$\text{Cov}\left(\mathbb{G}[\mathcal{A}^{(k)}], \mathbb{G}(f)\right) = \mathbb{V}\left(\mathbb{G}[\mathcal{A}^{(k)}]\right) \cdot \mathbb{E}[f \mid \mathcal{A}^{(k)}], \quad (4.2)$$

$$\text{Cov}\left(\mathbb{G}[\mathcal{A}^{(k)}], \mathbb{G}[\mathcal{A}^{(l)}]\right) = \mathbb{V}\left(\mathbb{G}[\mathcal{A}^{(k)}]\right) \mathbf{P}_{\mathcal{A}^{(l)}|\mathcal{A}^{(k)}}, \quad (4.3)$$

$$\Phi_k^{(N)}(f) = \mathbb{E}\left[f \mid \mathcal{A}^{(k)}\right] - \sum_{k < l \leq N} \mathbf{P}_{\mathcal{A}^{(l)}|\mathcal{A}^{(k)}} \cdot \Phi_l^{(N)}(f). \quad (4.4)$$

Proof. The j -th coordinate of the vector $\mathbb{V}\left(\mathbb{G}[\mathcal{A}^{(k)}]\right) \cdot \mathbb{E}[f \mid \mathcal{A}^{(k)}]$ is

$$\begin{aligned} & P(A_j^{(k)})(1 - P(A_j^{(k)}))\mathbb{E}(f \mid A_j^{(k)}) - \sum_{j \neq i \leq m_k} P(A_i^{(k)})P(A_j^{(k)})\mathbb{E}(f \mid A_i^{(k)}) \\ &= \mathbb{E}(1_{A_j^{(k)}} f) - P(A_j^{(k)}) \sum_{1 \leq i \leq m_k} \mathbb{E}(1_{A_i^{(k)}} f) \\ &= \text{Cov}\left(\mathbb{G}(A_j^{(k)}), \mathbb{G}(f)\right). \end{aligned}$$

Likewise the (i, j) -th coordinate of the matrix $\mathbb{V}\left(\mathbb{G}[\mathcal{A}^{(k)}]\right) \cdot \mathbf{P}_{\mathcal{A}^{(l)}|\mathcal{A}^{(k)}}$ is

$$\begin{aligned} & P(A_i^{(k)})(1 - P(A_i^{(k)}))P(A_j^{(l)} \mid A_i^{(k)}) - \sum_{j \neq m \leq m_k} P(A_i^{(k)})P(A_m^{(k)})P(A_j^{(l)} \mid A_m^{(k)}) \\ &= P(A_j^{(l)} \cap A_i^{(k)}) - P(A_i^{(k)}) \sum_{1 \leq m \leq m_k} P(A_j^{(l)} \cap A_m^{(k)}) \\ &= \text{Cov}\left(\mathbb{G}(A_i^{(k)}), \mathbb{G}(A_j^{(l)})\right). \end{aligned}$$

By the definition (2.3) of the vectors $\Phi_l^{(N)}(f)$ we get

$$\begin{aligned}
& \sum_{k < l \leq N} \mathbf{P}_{\mathcal{A}^{(l)} | \mathcal{A}^{(k)}} \cdot \Phi_l^{(N)}(f) \\
&= \sum_{k < l \leq N} \mathbf{P}_{\mathcal{A}^{(l)} | \mathcal{A}^{(k)}} \cdot \mathbb{E} \left[f \mid \mathcal{A}^{(l)} \right] \\
&+ \sum_{\substack{k < l \leq N, 1 \leq L \leq N-l \\ l < l_1 < \dots < l_L \leq N}} (-1)^L \mathbf{P}_{\mathcal{A}^{(l)} | \mathcal{A}^{(k)}} \mathbf{P}_{\mathcal{A}^{(l_1)} | \mathcal{A}^{(l)}} \dots \mathbf{P}_{\mathcal{A}^{(l_L)} | \mathcal{A}^{(l_{L-1})}} \cdot \mathbb{E} \left[f \mid \mathcal{A}^{(l_L)} \right] \\
&= \sum_{\substack{1 \leq L \leq N-k \\ k < l_1 < \dots < l_L \leq N}} (-1)^{L+1} \mathbf{P}_{\mathcal{A}^{(l_1)} | \mathcal{A}^{(k)}} \dots \mathbf{P}_{\mathcal{A}^{(l_L)} | \mathcal{A}^{(l_{L-1})}} \cdot \mathbb{E} \left[f \mid \mathcal{A}^{(l_L)} \right] \\
&= \mathbb{E} \left[f \mid \mathcal{A}^{(k)} \right] - \Phi_k^{(N)}(f)
\end{aligned}$$

which yields (4.4). \square

Step 3. Let us first compute the variance of $\mathbb{G}^{(N)}(f)$. By Lemma 3 we have

$$\begin{aligned}
& \mathbb{V} \left(\mathbb{G}^{(N)}(f) \right) - \mathbb{V}(\mathbb{G}(f)) \\
&= \sum_{k=1}^N \Phi_k^{(N)}(f)^t \cdot \mathbb{V} \left(\mathbb{G}[\mathcal{A}^{(k)}] \right) \cdot \Phi_k^{(N)}(f) - 2 \sum_{k=1}^N \Phi_k^{(N)}(f)^t \cdot \text{Cov} \left(\mathbb{G}[\mathcal{A}^{(k)}], \mathbb{G}(f) \right) \\
&+ 2 \sum_{1 \leq k < l \leq N} \Phi_k^{(N)}(f)^t \cdot \text{Cov} \left(\mathbb{G}[\mathcal{A}^{(k)}], \mathbb{G}[\mathcal{A}^{(l)}] \right) \cdot \Phi_l^{(N)}(f)
\end{aligned}$$

hence Lemma 4 gives, through (4.2) and (4.3),

$$\mathbb{V} \left(\mathbb{G}^{(N)}(f) \right) - \mathbb{V}(\mathbb{G}(f)) = \sum_{k=1}^N \Phi_k^{(N)}(f)^t \cdot \mathbb{V} \left(\mathbb{G}[\mathcal{A}^{(k)}] \right) \cdot \Psi_k^{(N)}(f)$$

where, by (4.4),

$$\Psi_k^{(N)}(f) = \Phi_k^{(N)}(f) - 2 \mathbb{E} \left[f \mid \mathcal{A}^{(k)} \right] + 2 \sum_{k < l \leq N} \mathbf{P}_{\mathcal{A}^{(l)} | \mathcal{A}^{(k)}} \cdot \Phi_l^{(N)}(f) = -\Phi_k^{(N)}(f). \tag{4.5}$$

The formula (2.4) is proved. It extends to the covariance since, by Lemma 3,

$$\begin{aligned}
& \text{Cov}(\mathbb{G}^{(N)}(f), \mathbb{G}^{(N)}(g)) - \text{Cov}(\mathbb{G}(f), \mathbb{G}(g)) \\
&= \frac{1}{2} (\Upsilon_N(f, g) - 2\Gamma_N(f, g)) + \frac{1}{2} (\Upsilon_N(g, f) - 2\Gamma_N(g, f))
\end{aligned}$$

where, by (4.2) and (4.3) again,

$$\begin{aligned}
\Upsilon_N(f, g) &= \sum_{1 \leq k \leq l \leq N} \Phi_k^{(N)}(f)^t \cdot \mathbb{V} \left(\mathbb{G}[\mathcal{A}^{(k)}] \right) \mathbf{P}_{\mathcal{A}^{(l)} | \mathcal{A}^{(k)}} \cdot \Phi_l^{(N)}(g) \\
\Gamma_N(f, g) &= \sum_{l=1}^N \Phi_k^{(N)}(f)^t \cdot \mathbb{V} \left(\mathbb{G}[\mathcal{A}^{(k)}] \right) \cdot \mathbb{E} \left[g \mid \mathcal{A}^{(k)} \right].
\end{aligned}$$

By replacing $\Psi_k^{(N)}(g)$ with $-\Phi_k^{(N)}(g)$ according to (4.5), we obtain

$$\frac{1}{2} (\Upsilon_N(f, g) - 2\Gamma_N(f, g)) = -\frac{1}{2} \sum_{k=1}^N \Phi_k^{(N)}(f)^t \cdot \mathbb{V}(\mathbb{G}[\mathcal{A}^{(k)}]) \cdot \Phi_k^{(N)}(g)$$

which is symmetric in f and g . The covariance formula of Proposition 7 is proved.

4.2 Proof of Propositions 8 and 9

Since $\mathbb{V}(\mathbb{G}[\mathcal{A}^{(k)}])$ is semi-definite positive, for all $1 \leq k \leq N$ and $f \in \mathcal{F}$ we have

$$\Phi_k^{(N)}(f)^t \cdot \mathbb{V}(\mathbb{G}[\mathcal{A}^{(k)}]) \cdot \Phi_k^{(N)}(f) \geq 0$$

and the variance part (2.4) of Proposition 8 follows from Proposition 7. For any $m \in \mathbb{N}_*$, $(f_1, \dots, f_m) \in \mathcal{F}^m$ and $u \in \mathbb{R}^m$, it further holds, by Proposition 7 again,

$$\begin{aligned} & u^t \left(\Sigma_m^{(0)} - \Sigma_m^{(N)} \right) u \\ &= \sum_{1 \leq i, j \leq m} u_i u_j \left(\text{Cov}(\mathbb{G}(f_i), \mathbb{G}(f_j)) - \text{Cov}(\mathbb{G}^{(N)}(f_i), \mathbb{G}^{(N)}(f_j)) \right) \\ &= \sum_{k=1}^N \sum_{1 \leq i, j \leq m} \left(u_i \Phi_k^{(N)}(f_i) \right)^t \cdot \mathbb{V}(\mathbb{G}[\mathcal{A}^{(k)}]) \cdot \left(u_j \Phi_k^{(N)}(f_j) \right) \\ &= \sum_{k=1}^N \left(\sum_{1 \leq i \leq m} u_i \Phi_k^{(N)}(f_i) \right)^t \cdot \mathbb{V}(\mathbb{G}[\mathcal{A}^{(k)}]) \cdot \left(\sum_{1 \leq j \leq m} u_j \Phi_k^{(N)}(f_j) \right) \geq 0. \end{aligned}$$

Under the wrapping hypothesis of Proposition 9 we have

$$\Phi_{N_0-k}^{(N_0)} = \Phi_{N_1-k}^{(N_1)}, \quad 0 \leq k < N_0,$$

since the corresponding (2.3) only involves $\mathcal{A}^{(N_0-k)} = \mathcal{A}^{(N_1-k)}$ for $0 \leq k < N_0$. Assuming moreover $N_1 \geq 2N_0$ we get, by Proposition 7,

$$\begin{aligned} & \mathbb{V}(\mathbb{G}^{(N_0)}(f)) - \mathbb{V}(\mathbb{G}^{(N_1)}(f)) \\ &= \sum_{k=1}^{N_1} \Phi_k^{(N_1)}(f)^t \cdot \mathbb{V}(\mathbb{G}[\mathcal{A}^{(k)}]) \cdot \Phi_k^{(N_1)}(f) - \sum_{k=1}^{N_0} \Phi_k^{(N_0)}(f)^t \cdot \mathbb{V}(\mathbb{G}[\mathcal{A}^{(k)}]) \cdot \Phi_k^{(N_0)}(f) \\ &= \sum_{k=N_0+1}^{N_1-N_0} \Phi_k^{(N_1)}(f)^t \cdot \mathbb{V}(\mathbb{G}[\mathcal{A}^{(k)}]) \cdot \Phi_k^{(N_1)}(f) \geq 0 \end{aligned}$$

and, for $m \in \mathbb{N}_*$, $(f_1, \dots, f_m) \in \mathcal{F}^m$ and $u \in \mathbb{R}^m$,

$$\begin{aligned}
& u^t \left(\Sigma_m^{(N_0)} - \Sigma_m^{(N_1)} \right) u \\
&= u^t \left((\Sigma_m^{(0)} - \Sigma_m^{(N_1)}) - (\Sigma_m^{(0)} - \Sigma_m^{(N_0)}) \right) u \\
&= \sum_{k=N_0+1}^{N_1-N_0} \left(\sum_{1 \leq i \leq m} u_i \Phi_k^{(N)}(f_i) \right)^t \cdot \mathbb{V} \left(\mathbb{G}[\mathcal{A}^{(k)}] \right) \cdot \left(\sum_{1 \leq j \leq m} u_j \Phi_k^{(N)}(f_j) \right) \geq 0.
\end{aligned}$$

4.3 Proof of Proposition 10

We show the result by double induction. For $m = 0$ we have $\mathbb{G}^{(0)}(f) = \mathbb{G}(f)$ and, by (1.11), $\mathbb{G}^{(1)}(f) = \mathbb{G}(f) - \mathbb{E}[f|\mathcal{A}]^t \mathbb{G}[\mathcal{A}]$. Assume that (2.9) and (2.10) are true for $m \in \mathbb{N}$. For $m + 1$ we have, by the raking ratio transform (1.11),

$$\mathbb{G}^{(2m+2)}(f) = \mathbb{G}^{(2m+1)}(f) - \mathbb{E}[f|\mathcal{B}]^t \cdot \mathbb{G}^{(2m+1)}[\mathcal{B}], \quad (4.6)$$

$$\mathbb{G}^{(2m+3)}(f) = \mathbb{G}^{(2m+2)}(f) - \mathbb{E}[f|\mathcal{A}]^t \cdot \mathbb{G}^{(2m+2)}[\mathcal{A}]. \quad (4.7)$$

For $1 \leq j \leq m_2$ and $f = 1_{B_j}$ we get, by (2.10), (2.5), (2.6), (2.7) and (2.8),

$$\begin{aligned}
& \mathbb{G}^{(2m+1)}(B_j) \\
&= \mathbb{G}(B_j) - S_{1,\text{odd}}^{(m-1)}(1_{B_j})^t \cdot \mathbb{G}[\mathcal{A}] - S_{2,\text{odd}}^{(m-1)}(1_{B_j})^t \cdot \mathbb{G}[\mathcal{B}] \\
&= \mathbb{G}(B_j) - \left(\sum_{k=0}^{m-1} (\mathbf{P}_{\mathcal{A}|\mathcal{B}} \mathbf{P}_{\mathcal{B}|\mathcal{A}})^k \cdot (\mathbb{E}[1_{B_j}|\mathcal{B}] - \mathbf{P}_{\mathcal{A}|\mathcal{B}} \cdot \mathbb{E}[1_{B_j}|\mathcal{A}]) \right)^t \cdot \mathbb{G}[\mathcal{B}] \\
&\quad - \left(\sum_{k=0}^{m-1} (\mathbf{P}_{\mathcal{B}|\mathcal{A}} \mathbf{P}_{\mathcal{A}|\mathcal{B}})^k \cdot (\mathbb{E}[1_{B_j}|\mathcal{A}] - \mathbf{P}_{\mathcal{B}|\mathcal{A}} \cdot \mathbb{E}[1_{B_j}|\mathcal{B}]) + (\mathbf{P}_{\mathcal{B}|\mathcal{A}} \mathbf{P}_{\mathcal{A}|\mathcal{B}})^m \cdot \mathbb{E}[1_{B_j}|\mathcal{A}] \right)^t \cdot \mathbb{G}[\mathcal{A}]
\end{aligned}$$

where $\mathbb{E}[1_{B_j}|\mathcal{A}]$ is the j -th column of $\mathbf{P}_{\mathcal{B}|\mathcal{A}}$ and $\mathbb{E}[1_{B_j}|\mathcal{B}]$ is the j -th unit vector of \mathbb{R}^{m_2} . Therefore

$$\begin{aligned}
& \mathbb{G}^{(2m+1)}[\mathcal{B}] \\
&= \mathbb{G}[\mathcal{B}] - \left(\sum_{k=0}^{m-1} (\mathbf{P}_{\mathcal{A}|\mathcal{B}} \mathbf{P}_{\mathcal{B}|\mathcal{A}})^k (\text{Id}_{m_2} - \mathbf{P}_{\mathcal{A}|\mathcal{B}} \mathbf{P}_{\mathcal{B}|\mathcal{A}}) \right)^t \cdot \mathbb{G}[\mathcal{B}] \\
&\quad - \left(\sum_{k=0}^{m-1} (\mathbf{P}_{\mathcal{B}|\mathcal{A}} \mathbf{P}_{\mathcal{A}|\mathcal{B}})^k (\mathbf{P}_{\mathcal{B}|\mathcal{A}} - \mathbf{P}_{\mathcal{B}|\mathcal{A}} \text{Id}_{m_2}) + (\mathbf{P}_{\mathcal{B}|\mathcal{A}} \mathbf{P}_{\mathcal{A}|\mathcal{B}})^m \mathbf{P}_{\mathcal{B}|\mathcal{A}} \right)^t \cdot \mathbb{G}[\mathcal{A}] \\
&= ((\mathbf{P}_{\mathcal{A}|\mathcal{B}} \mathbf{P}_{\mathcal{B}|\mathcal{A}})^m)^t \cdot \mathbb{G}[\mathcal{B}] - ((\mathbf{P}_{\mathcal{B}|\mathcal{A}} \mathbf{P}_{\mathcal{A}|\mathcal{B}})^m \mathbf{P}_{\mathcal{B}|\mathcal{A}})^t \cdot \mathbb{G}[\mathcal{A}]
\end{aligned}$$

Finally (2.9) and (4.6) then again (2.5), (2.6), (2.7) and (2.8) together imply

$$\begin{aligned}
& \mathbb{G}^{(2m+2)}(f) \\
&= \mathbb{G}(f) - S_{1,\text{odd}}^{(m-1)}(f)^t \cdot \mathbb{G}[\mathcal{A}] - S_{2,\text{odd}}^{(m-1)}(f)^t \cdot \mathbb{G}[\mathcal{B}] - \mathbb{E}[f|\mathcal{B}]^t \cdot \mathbb{G}^{(2m+1)}[\mathcal{B}] \\
&= \mathbb{G}(f) - \left(S_{1,\text{odd}}^{(m-1)}(f) - (\mathbf{P}_{\mathcal{B}|\mathcal{A}}\mathbf{P}_{\mathcal{A}|\mathcal{B}})^m \mathbf{P}_{\mathcal{B}|\mathcal{A}} \cdot \mathbb{E}[f|\mathcal{B}] \right)^t \cdot \mathbb{G}[\mathcal{A}] \\
&\quad - \left(S_{2,\text{odd}}^{(m-1)}(f) + (\mathbf{P}_{\mathcal{A}|\mathcal{B}}\mathbf{P}_{\mathcal{B}|\mathcal{A}})^m \cdot \mathbb{E}[f|\mathcal{B}] \right)^t \cdot \mathbb{G}[\mathcal{B}] \\
&= \mathbb{G}(f) - S_{1,\text{even}}^{(m)}(f)^t \cdot \mathbb{G}[\mathcal{A}] - S_{2,\text{even}}^{(m-1)}(f)^t \cdot \mathbb{G}[\mathcal{B}]
\end{aligned}$$

and (2.9) is valid for $m+1$. If $1 \leq i \leq m_1$ then $\mathbb{E}[1_{A_i}|\mathcal{B}]$ is the i -th column of $\mathbf{P}_{\mathcal{A}|\mathcal{B}}$ and $\mathbb{E}[1_{A_i}|\mathcal{A}]$ is the i -th unit vector of \mathbb{R}^{m_1} thus (2.9) for $m+1$ and $f = 1_{B_i}$ in turn entails

$$\begin{aligned}
& \mathbb{G}^{(2m+2)}[\mathcal{A}] \\
&= \mathbb{G}[\mathcal{A}] - \left(\sum_{k=0}^m (\mathbf{P}_{\mathcal{B}|\mathcal{A}}\mathbf{P}_{\mathcal{A}|\mathcal{B}})^k (\text{Id}_{m_1} - \mathbf{P}_{\mathcal{B}|\mathcal{A}}\mathbf{P}_{\mathcal{A}|\mathcal{B}}) \right)^t \cdot \mathbb{G}[\mathcal{A}] \\
&\quad - \left(\sum_{k=0}^{m-1} (\mathbf{P}_{\mathcal{A}|\mathcal{B}}\mathbf{P}_{\mathcal{B}|\mathcal{A}})^k (\mathbf{P}_{\mathcal{A}|\mathcal{B}} - \mathbf{P}_{\mathcal{A}|\mathcal{B}}\text{Id}_{m_1}) + (\mathbf{P}_{\mathcal{A}|\mathcal{B}}\mathbf{P}_{\mathcal{B}|\mathcal{A}})^m \mathbf{P}_{\mathcal{A}|\mathcal{B}} \right)^t \cdot \mathbb{G}[\mathcal{B}] \\
&= ((\mathbf{P}_{\mathcal{B}|\mathcal{A}}\mathbf{P}_{\mathcal{A}|\mathcal{B}})^{m+1})^t \cdot \mathbb{G}[\mathcal{A}] - ((\mathbf{P}_{\mathcal{A}|\mathcal{B}}\mathbf{P}_{\mathcal{B}|\mathcal{A}})^m \mathbf{P}_{\mathcal{A}|\mathcal{B}})^t \cdot \mathbb{G}[\mathcal{B}]
\end{aligned}$$

and also, thanks to (2.10) and (4.7),

$$\begin{aligned}
& \mathbb{G}^{(2m+3)}(f) \\
&= \mathbb{G}(f) - \left(S_{1,\text{odd}}^{(m)}(f) - (\mathbf{P}_{\mathcal{B}|\mathcal{A}}\mathbf{P}_{\mathcal{A}|\mathcal{B}})^{m+1} \cdot \mathbb{E}[f|\mathcal{A}] \right)^t \cdot \mathbb{G}[\mathcal{A}] \\
&\quad - \left(S_{2,\text{odd}}^{(m-1)}(f) + (\mathbf{P}_{\mathcal{A}|\mathcal{B}}\mathbf{P}_{\mathcal{B}|\mathcal{A}})^m \cdot \mathbb{E}[f|\mathcal{B}] \right)^t \cdot \mathbb{G}[\mathcal{B}] - \mathbb{E}[f|\mathcal{A}]^t \cdot \mathbb{G}^{(2m+2)}[\mathcal{A}] \\
&= \mathbb{G}(f) - S_{1,\text{odd}}^{(m)}(f)^t \cdot \mathbb{G}[\mathcal{A}] - S_{2,\text{odd}}^{(m)}(f)^t \cdot \mathbb{G}[\mathcal{B}]
\end{aligned}$$

which is (2.10) for $m+1$.

4.4 Proof of Proposition 11

Step 1. For $m \geq 1$ let $\mathbf{0}_{m,m}$ be the $m \times m$ null matrix. Also recall the vectors $P(\mathcal{A}) = (P(A_1), \dots, P(A_{m_1}))$ and $P(\mathcal{B}) = (P(B_1), \dots, P(B_{m_2}))$.

Lemma 5. *Assume (ER). For $l = 1, 2$ there exists an invertible $m_l \times m_l$ matrix U_l and an upper triangular $(m_l - 1) \times (m_l - 1)$ matrix T_l such that*

$$\begin{aligned}
\mathbf{P}_{\mathcal{B}|\mathcal{A}}\mathbf{P}_{\mathcal{A}|\mathcal{B}} &= U_1 \begin{pmatrix} 1 & \mathbf{0}_{1,m_1-1} \\ \mathbf{0}_{m_1-1,1} & T_1 \end{pmatrix} U_1^{-1}, \quad \lim_{k \rightarrow +\infty} T_1^k = \mathbf{0}_{m_1-1,m_1-1}, \\
\mathbf{P}_{\mathcal{A}|\mathcal{B}}\mathbf{P}_{\mathcal{B}|\mathcal{A}} &= U_2 \begin{pmatrix} 1 & \mathbf{0}_{1,m_2-1} \\ \mathbf{0}_{m_2-1,1} & T_2 \end{pmatrix} U_2^{-1}, \quad \lim_{k \rightarrow +\infty} T_2^k = \mathbf{0}_{m_2-1,m_2-1}.
\end{aligned}$$

Proof. Since \mathcal{A} is a partition, for $1 \leq i \leq m_2$ the sum of the m_1 terms of row i of $\mathbf{P}_{\mathcal{A}|\mathcal{B}}$ is $\sum_{j=1}^{m_1} P(A_j | B_i) = 1$ hence $\mathbf{P}_{\mathcal{A}|\mathcal{B}}$ is stochastic. Likewise $\mathbf{P}_{\mathcal{B}|\mathcal{A}}$ is stochastic and, by stability, so are $\mathbf{P}_{\mathcal{A}|\mathcal{B}}\mathbf{P}_{\mathcal{B}|\mathcal{A}}$ and $\mathbf{P}_{\mathcal{B}|\mathcal{A}}\mathbf{P}_{\mathcal{A}|\mathcal{B}}$. Let the column of 1's associated to their eigenvalue 1 be in first position in their respective matrix U_1, U_2 of eigenvectors. The announced decomposition is always true with some upper triangular matrices T_l having Jordan decomposition $T_l = D_l + N_l$ where $D_l = P_l \Delta_l P_l^{-1}$, Δ_l is a diagonal $(m_l - 1) \times (m_l - 1)$ matrix, P_l is an invertible $(m_l - 1) \times (m_l - 1)$ matrix and N_l is a nilpotent $(m_l - 1) \times (m_l - 1)$ matrix of order $n_l \geq 1$ that commute with D_l . Next observe that

$$\begin{aligned} (P(\mathcal{A}) \cdot \mathbf{P}_{\mathcal{B}|\mathcal{A}}\mathbf{P}_{\mathcal{A}|\mathcal{B}})_k &= \sum_{i=1}^{m_1} P(A_i) (\mathbf{P}_{\mathcal{B}|\mathcal{A}}\mathbf{P}_{\mathcal{A}|\mathcal{B}})_{i,k} \\ &= \sum_{i=1}^{m_1} P(A_i) \sum_{j=1}^{m_2} (\mathbf{P}_{\mathcal{B}|\mathcal{A}})_{i,j} (\mathbf{P}_{\mathcal{A}|\mathcal{B}})_{j,k} \\ &= \sum_{j=1}^{m_2} \sum_{i=1}^{m_1} P(A_i) P(B_j | A_i) P(A_k | B_j) = P(A_k) \end{aligned}$$

which proves that $\mathbf{P}_{\mathcal{B}|\mathcal{A}}\mathbf{P}_{\mathcal{A}|\mathcal{B}}$ has invariant probability $P(\mathcal{A})$. Similarly, $P(\mathcal{B})$ is invariant for $\mathbf{P}_{\mathcal{A}|\mathcal{B}}\mathbf{P}_{\mathcal{B}|\mathcal{A}}$, and the first line of U_1^{-1} and U_2^{-1} is $P[\mathcal{A}]$ and $P[\mathcal{B}]$ respectively. Under (ER) these matrices are ergodic, which ensures that the eigenvalues of Δ_l have moduli strictly less than the dominant 1 since it is the case of eigenvalues of T_l hence D_l . It follows that

$$\lim_{k \rightarrow +\infty} \Delta_l^k = \mathbf{0}_{m_l-1, m_l-1}, \quad l = 1, 2.$$

Furthermore, since N_l and D_l commute it holds

$$T_l^k = \sum_{j=0}^{n_l-1} \binom{k}{j} N_l^j D_l^{k-j}, \quad l = 1, 2, \quad k \geq n_l.$$

We conclude that $\lim_{k \rightarrow +\infty} T_l^k = \mathbf{0}_{m_l-1, m_l-1}$. \square

Step 2. Let $V_1(f) = (\mathbb{E}[f|\mathcal{A}] - \mathbf{P}_{\mathcal{B}|\mathcal{A}}\mathbb{E}[f|\mathcal{B}])$ and $V_2(f) = (\mathbb{E}[f|\mathcal{B}] - \mathbf{P}_{\mathcal{A}|\mathcal{B}}\mathbb{E}[f|\mathcal{A}])$.

Lemma 6. *Under (ER) we have*

$$\lim_{k \rightarrow +\infty} (\mathbf{P}_{\mathcal{B}|\mathcal{A}}\mathbf{P}_{\mathcal{A}|\mathcal{B}})^k \cdot V_1(f) = \mathbf{0}_{m_1, 1}, \quad \lim_{k \rightarrow +\infty} (\mathbf{P}_{\mathcal{A}|\mathcal{B}}\mathbf{P}_{\mathcal{B}|\mathcal{A}})^k \cdot V_2(f) = \mathbf{0}_{m_2, 1}.$$

Proof. By Lemma 5 we have

$$\lim_{k \rightarrow +\infty} (\mathbf{P}_{\mathcal{B}|\mathcal{A}}\mathbf{P}_{\mathcal{A}|\mathcal{B}})^k = \begin{pmatrix} P(\mathcal{A}) \\ \vdots \\ P(\mathcal{A}) \end{pmatrix}, \quad \lim_{k \rightarrow +\infty} (\mathbf{P}_{\mathcal{A}|\mathcal{B}}\mathbf{P}_{\mathcal{B}|\mathcal{A}})^k = \begin{pmatrix} P(\mathcal{B}) \\ \vdots \\ P(\mathcal{B}) \end{pmatrix}. \quad (4.8)$$

The scalar product of $P(\mathcal{A})$ by $V_1(f)$ is null since $P(\mathcal{A}) \cdot \mathbb{E}[f|\mathcal{A}] = P(f)$ and

$$\begin{aligned} P(\mathcal{A}) \cdot \mathbf{P}_{\mathcal{B}|\mathcal{A}} \mathbb{E}[f|\mathcal{B}] &= \sum_{j=1}^{m_1} P(A_j) \sum_{k=1}^{m_2} P(B_k | A_j) \mathbb{E}(f|B_k) \\ &= \sum_{j=1}^{m_1} \sum_{k=1}^{m_2} P(A_j \cap B_k) \mathbb{E}[f|B_k] = P(f). \end{aligned}$$

Likewise we get $P(\mathcal{B}) \cdot V_2(f) = 0$. \square

The following convergences hold for any matrix norm. By Lemma 5 we have

$$\begin{aligned} \sum_{k=0}^N (\mathbf{P}_{\mathcal{B}|\mathcal{A}} \mathbf{P}_{\mathcal{A}|\mathcal{B}})^k &= U_1 \begin{pmatrix} N+1 & \mathbf{0}_{1,m_1-1} \\ \mathbf{0}_{m_1-1,1} & \sum_{k=0}^N T_1^k \end{pmatrix} U_1^{-1}, \\ \sum_{k=0}^N (\mathbf{P}_{\mathcal{A}|\mathcal{B}} \mathbf{P}_{\mathcal{B}|\mathcal{A}})^k &= U_2 \begin{pmatrix} N+1 & \mathbf{0}_{1,m_2-1} \\ \mathbf{0}_{m_2-1,1} & \sum_{k=0}^N T_2^k \end{pmatrix} U_2^{-1}. \end{aligned}$$

Now, the matrices $\text{Id}_{m_1-1} - T_1$ and $\text{Id}_{m_2-1} - T_2$ are nonsingular since 1 is a dominant eigenvalue of $\mathbf{P}_{\mathcal{B}|\mathcal{A}} \mathbf{P}_{\mathcal{A}|\mathcal{B}}$ and $\mathbf{P}_{\mathcal{A}|\mathcal{B}} \mathbf{P}_{\mathcal{B}|\mathcal{A}}$. It follows that

$$\begin{aligned} S_{1,\text{even}}^{(N)}(f) &= U_1 \begin{pmatrix} N+1 & \mathbf{0}_{1,m_1-1} \\ \mathbf{0}_{m_1-1,1} & (\text{Id}_{m_1-1} - T_1)^{-1} (\text{Id}_{m_1-1} - T_1^{N+1}) \end{pmatrix} U_1^{-1} \cdot V_1(f) \\ S_{2,\text{odd}}^{(N)}(f) &= U_2 \begin{pmatrix} N+1 & \mathbf{0}_{1,m_2-1} \\ \mathbf{0}_{m_2-1,1} & (\text{Id}_{m_2-1} - T_2)^{-1} (\text{Id}_{m_2-1} - T_2^{N+1}) \end{pmatrix} U_2^{-1} \cdot V_2(f) \end{aligned}$$

which, by Lemma 5, converge respectively to

$$\begin{aligned} S_{1,\text{even}}(f) &= U_1 \begin{pmatrix} N+1 & \mathbf{0}_{1,m_1-1} \\ \mathbf{0}_{m_1-1,1} & (\text{Id}_{m_1-1} - T_1)^{-1} \end{pmatrix} U_1^{-1} \cdot V_1(f) \\ S_{2,\text{odd}}(f) &= U_2 \begin{pmatrix} N+1 & \mathbf{0}_{1,m_2-1} \\ \mathbf{0}_{m_2-1,1} & (\text{Id}_{m_2-1} - T_2)^{-1} \end{pmatrix} U_2^{-1} \cdot V_2(f). \end{aligned}$$

Since we have already seen by using (4.8) that

$$\lim_{k \rightarrow +\infty} (\mathbf{P}_{\mathcal{B}|\mathcal{A}} \mathbf{P}_{\mathcal{A}|\mathcal{B}})^k \cdot \mathbb{E}[f|\mathcal{A}] = P_1[f], \quad \lim_{k \rightarrow +\infty} (\mathbf{P}_{\mathcal{A}|\mathcal{B}} \mathbf{P}_{\mathcal{B}|\mathcal{A}})^k \cdot \mathbb{E}[f|\mathcal{B}] = P_2[f],$$

we conclude by (2.7) and (2.8) that $S_{1,\text{odd}}^{(N)}(f)$, $S_{2,\text{even}}^{(N)}(f)$ converge to the vectors $S_{1,\text{odd}}(f) = S_{1,\text{even}}(f) + P_1[f]$, $S_{2,\text{even}}(f) = S_{1,\text{odd}}(f) + P_1[f]$ respectively.

Step 3. Given the spectral radius $\rho(T_l) < 1$ of T_l let $\lambda_l = \rho(T_l) + \varepsilon < 1$, $l = 1, 2$ for any $\varepsilon > 0$. Then there exists a vector norm $\|\cdot\|_l$ on \mathbb{C}^{m_l-1} such that its induced matrix norm $\|\cdot\|_l$ on matrices $(m_l - 1) \times (m_l - 1)$ satisfies $\|T_l\|_l \leq \lambda_l$. Introduce the vector norm $\|(x_1, \dots, x_{m_l})^t\|_l' = |x_1| + \|(x_2, \dots, x_{m_l})^t\|_l$ on \mathbb{C}^{m_l} and the induced operator norm $\|\cdot\|_l'$ for $m_l \times m_l$ matrices. Then we have

$$\left\| \begin{pmatrix} 0 & \mathbf{0}_{1,m_l-1} \\ \mathbf{0}_{m_l-1,1} & T \end{pmatrix} \right\|_l' = \sup \left\{ x \in \mathbb{C}^{m_l} : \frac{0 + \|T(x_2, \dots, x_{m_l})^t\|_l}{|x_1| + \|(x_2, \dots, x_{m_l})^t\|_l} \right\} = \|T\|_l$$

for any $m_l \times m_l$ matrix T . Let $K_l = \left\| (\text{Id}_{m_l-1} - T_l)^{-1} \right\|_l$, $\tilde{K}_l = \left\| U_l \right\|_l' \left\| U_l^{-1} \right\|_l'$ and $K_l' > 0$ be such that $\left\| \cdot \right\|_l' \leq K_l' \left\| \cdot \right\|_\infty$. By using Lemmas 5 and 6 we get

$$\begin{aligned} & \left\| S_{l,\text{even}}^{(N)}(f) - S_{l,\text{even}}(f) \right\|_l' \\ & \leq \left\| U_l \begin{pmatrix} 0 & \mathbf{0}_{1,m_l-1} \\ \mathbf{0}_{m_l-1,1} & -(\text{Id}_{m_l-1} - T_l)^{-1} T_l^{N+1} \end{pmatrix} U_l^{-1} \right\|_l' \left\| V_l(f) \right\|_l' \\ & \leq \tilde{K}_l \left\| (\text{Id}_{m_l-1} - T_l)^{-1} T_l^{N+1} \right\|_l \left\| V_l(f) \right\|_l' \\ & \leq \tilde{K}_l K_l \lambda_l^{N+1} K_l' \left\| V_l(f) \right\|_\infty \leq c_l \lambda_l^{N+1} \end{aligned}$$

where $c_l = \tilde{K}_l K_l K_l' M$. Similar constants show up for $\left\| S_{l,\text{odd}}^{(N)}(f) - S_{l,\text{odd}}(f) \right\|_l'$. The final constants c_1 and c_2 depend on $\lambda_1, \lambda_2, \varepsilon$, both matrices (ER) but also the two implicit constants relating the norms $\left\| \cdot \right\|_{m_1}, \left\| \cdot \right\|_{m_2}$ of Proposition 11 to the equivalent norms $\left\| \cdot \right\|_1, \left\| \cdot \right\|_2$.

4.5 Proof of Theorem 2.2

Write $Z_1 = \max_{1 \leq j \leq m_1} |\mathbb{G}(A_j)|$ and $Z_2 = \max_{1 \leq j \leq m_2} |\mathbb{G}(B_j)|$. According to Proposition 11 the following right hand side random variables, for $* \in \{\text{even}, \text{odd}\}$,

$$\begin{aligned} \sup_{f \in \mathcal{F}} \left| (S_{1,*}^{(N)}(f) - S_{1,*}(f))^t \cdot \mathbb{G}[\mathcal{A}] \right| & \leq c_1 \sup_{f \in \mathcal{F}} \left\| S_{1,*}^{(N)}(f) - S_{1,*}(f) \right\|_{m_1} Z_1 \\ \sup_{f \in \mathcal{F}} \left| (S_{2,*}^{(N)}(f) - S_{2,*}(f))^t \cdot \mathbb{G}[\mathcal{B}] \right| & \leq c_2 \sup_{f \in \mathcal{F}} \left\| S_{2,*}^{(N)}(f) - S_{2,*}(f) \right\|_{m_2} Z_2 \end{aligned}$$

almost surely converge to 0 since $\mathbb{P}(\|\mathbb{G}\|_{\mathcal{F}} < +\infty) = 1$. Hence the processes $\mathbb{G}^{(2m)}, \mathbb{G}^{(2m+1)}$ converge almost surely in $\ell^\infty(\mathcal{F})$ to $\mathbb{G}_{\text{even}}^{(\infty)}, \mathbb{G}_{\text{odd}}^{(\infty)}$ defined by

$$\begin{aligned} \mathbb{G}_{\text{even}}^{(\infty)}(f) & = \mathbb{G}(f) - S_{1,\text{even}}(f)^t \cdot \mathbb{G}[\mathcal{A}] - S_{2,\text{even}}(f)^t \cdot \mathbb{G}[\mathcal{B}] \\ & = \mathbb{G}^{(\infty)}(f) - P_2[f]^t \cdot \mathbb{G}[\mathcal{B}] \\ \mathbb{G}_{\text{odd}}^{(\infty)}(f) & = \mathbb{G}(f) - S_{1,\text{odd}}(f)^t \cdot \mathbb{G}[\mathcal{A}] - S_{2,\text{odd}}(f)^t \cdot \mathbb{G}[\mathcal{B}] \\ & = \mathbb{G}^{(\infty)}(f) - P_1[f]^t \cdot \mathbb{G}[\mathcal{A}] \end{aligned}$$

with $\mathbb{G}^{(\infty)}(f) = \mathbb{G}(f) - S_{1,\text{even}}(f)^t \cdot \mathbb{G}[\mathcal{A}] - S_{2,\text{odd}}(f)^t \cdot \mathbb{G}[\mathcal{B}]$ and using (2.8), (2.7). Since $P_1[f]^t \cdot \mathbb{G}[\mathcal{A}] = P(f) \sum_{j=1}^{m_1} \mathbb{G}(A_j) = P(f) \mathbb{G}(1) = 0$ and $P_2[f]^t \cdot \mathbb{G}[\mathcal{B}] = 0$ almost surely, we see that $\mathbb{G}_{\text{even}}^{(\infty)}(\mathcal{F}) = \mathbb{G}_{\text{odd}}^{(\infty)}(\mathcal{F}) = \mathbb{G}^{(\infty)}(\mathcal{F})$. Applying Proposition 11 with the supremum norms $\left\| \cdot \right\|_{m_1}$ and $\left\| \cdot \right\|_{m_2}$ further yields, for any $m \geq 0$ and $c_0 = m_1 c_1 + m_2 c_2$,

$$\begin{aligned} \left\| \mathbb{G}^{(2m)} - \mathbb{G}_{\text{even}}^{(\infty)} \right\|_{\mathcal{F}} & = \left\| (S_{1,\text{even}}^{(m-1)} - S_{1,\text{even}})^t \cdot \mathbb{G}[\mathcal{A}] + (S_{2,\text{even}}^{(m-2)} - S_{2,\text{even}})^t \cdot \mathbb{G}[\mathcal{B}] \right\|_{\mathcal{F}} \\ & \leq c_0 \max(\lambda_1, \lambda_2)^{m-2} Z \end{aligned} \quad (4.9)$$

where $Z = \max(Z_1, Z_2)$, and

$$\left\| \mathbb{G}^{(2m+1)} - \mathbb{G}_{\text{odd}}^{(\infty)} \right\|_{\mathcal{F}} \leq c_0 \max(\lambda_1, \lambda_2)^{m-1} Z. \quad (4.10)$$

Let $\varepsilon_N = q_N c_0 \max(\lambda_1, \lambda_2)^{N/2}$ and $q_N = F_Z^{-1}(c_0 \max(\lambda_1, \lambda_2)^{N/2})$, which is well defined for N large enough. From (4.9) and (4.10) we deduce that

$$\mathbb{P} \left(\left\| \mathbb{G}^{(N)} - \mathbb{G}^{(\infty)} \right\|_{\mathcal{F}} > \varepsilon_N \right) \leq \mathbb{P}(Z > q_N) \leq c_0 \max(\lambda_1, \lambda_2)^{N/2}$$

whence an upper bound for the Lévy-Prokhorov distance

$$d_{LP}(\mathbb{G}^{(N)}, \mathbb{G}^{(\infty)}) \leq \max \left(\mathbb{P} \left(\left\| \mathbb{G}^{(N)} - \mathbb{G}^{(\infty)} \right\|_{\mathcal{F}} > \varepsilon_N \right), \varepsilon_N \right) \leq c_0 q_N \max(\lambda_1, \lambda_2)^{N/2}.$$

Let Φ denote the standard Gaussian distribution function, $c_5 = m_1 + m_2$ and $c_4^2 = \max_{D \in \mathcal{A} \cup \mathcal{B}} \{P(D)(1 - P(D))\}$. The union bound

$$\mathbb{P}(Z > \lambda) \leq c_5 \left(1 - \Phi \left(\frac{\lambda}{c_4} \right) \right) \leq \frac{c_5 c_4}{\sqrt{2\pi} \lambda} \exp \left(-\frac{\lambda^2}{2c_4^2} \right)$$

shows that $q_N \leq c_6 c_4 \sqrt{N \log(1/c_0 \max(\lambda_1, \lambda_2))}$ for some $c_6 > 0$.

Appendix

.1 Elementary example

In many settings the auxiliary information concerns a few partitions and they are raked in a periodic order. A subcase is a multiple ways contingency table. The Raking-Ratio algorithm then changes the weights of cells of a contingency table in such a way that given margins are respected, just as if the sample should have respected the expected values of known probabilities. Let us start from the following basic table which is the most common subcase of Section 2.3.

$\mathbb{P}_n(1_{A_i^{(1)} \cap A_j^{(2)}})$	$A_1^{(2)}$	$A_2^{(2)}$	$A_3^{(2)}$	Total	Expected total
$A_1^{(1)}$	0.2	0.25	0.1	0.55	0.52
$A_2^{(1)}$	0.1	0.2	0.15	0.45	0.48
Total	0.3	0.45	0.25	1	
Excepted total	0.31	0.4	0.29		

The margins of this sample differ from the known margins, here called expected total. Firstly the weights of lines are corrected, hence each cell is multiplied by the ratio of the expected total and the actual one, this is step $N = 1$.

$\mathbb{P}_n^{(1)}(1_{A_i^{(1)} \cap A_j^{(2)}})$	$A_1^{(2)}$	$A_2^{(2)}$	$A_3^{(2)}$	Total	Expected total
$A_1^{(1)}$	0.189	0.236	0.095	0.52	0.52
$A_2^{(1)}$	0.11	0.21	0.16	0.48	0.48
Total	0.299	0.446	0.255	1	
Excepted total	0.31	0.4	0.29		

The totals for each column are similarly corrected at step $N = 2$. Typically the margins of the lines no longer match the expected frequencies. Here they move in the right direction. Some estimators based on $\mathbb{P}_n^{(2)}$ may be improved.

$\mathbb{P}_n^{(2)}(1_{A_i^{(1)} \cap A_j^{(2)}})$	$A_1^{(2)}$	$A_2^{(2)}$	$A_3^{(2)}$	Total	Excepted total
$A_1^{(1)}$	0.196	0.212	0.108	0.516	0.52
$A_2^{(1)}$	0.114	0.188	0.182	0.484	0.48
Total	0.31	0.4	0.29	1	
Excepted total	0.31	0.4	0.29		

The last two operations are repeated until stabilization. The algorithm converges to the Kullback projection of the initial joint law. The rate depends only on the initial table compared to the desired marginals. It takes only 7 iterations in our case to match the expected margins.

$\mathbb{P}_n^{(7)}(1_{A_i^{(1)} \cap A_j^{(2)}})$	$A_1^{(2)}$	$A_2^{(2)}$	$A_3^{(2)}$	Total	Excepted total
$A_1^{(1)}$	0.199	0.212	0.109	0.52	0.52
$A_2^{(1)}$	0.111	0.188	0.181	0.48	0.48
Total	0.31	0.4	0.29	1	
Excepted total	0.31	0.4	0.29		

The final raked frequencies are slightly moved away the initial ones, however this has to be compared with the natural sampling oscillation order $1/\sqrt{n}$ – insidiously n was not mentioned. Clearly, for very large samples such a correction is unlikely since the initial table was already close to the true one, which respects the margin constraint. For small samples such changes are likely to occur that may improve a large class of estimators, and worsen others. In both cases, a possible drawback is a slow rate of convergence of the empirical raking weights, thus enforcing a so large N that our results are no more relevant. This happens in particular when λ_1 and λ_2 of Proposition 11 are close to one.

.2 Counterexample of Remark J

Let assume that P satisfies the following probability values

$P(A_i \cap B_j)$	A_1	A_2	A_3	$P(B_j)$
B_1	0.2	0.25	0.1	0.55
B_2	0.25	0.1	0.1	0.45
$P(A_i)$	0.45	0.35	0.2	

and that f has the following conditional expectations

$\mathbb{E}(f A_i \cap B_j)$	A_1	A_2	A_3	$\mathbb{E}^{(2)}(f) \simeq$
B_1	0.75	-0.5	0.5	0.136
B_2	0.5	0.25	0.5	0.444
$\mathbb{E}^{(1)}(f) \simeq$	0.611	-0.286	0.5	

By supposing also that $\mathbb{V}(f|A_i \cap B_j) = 0.5$ for all $i = 1, 2, 3$ and $j = 1, 2$ we can compute the theoretical limiting variances from Proposition 7. We get $\mathbb{V}(\mathbb{G}^{(0)}(f)) \simeq 0.734$; $\mathbb{V}(\mathbb{G}^{(1)}(f)) \simeq 0.563$; $\mathbb{V}(\mathbb{G}^{(2)}(f)) \simeq 0.569$; $\mathbb{V}(\mathbb{G}^{(3)}(f)) \simeq 0.402$. The fact that $\mathbb{V}(\mathbb{G}^{(2)}(f)) > \mathbb{V}(\mathbb{G}^{(1)}(f))$ shows that the variance doesn't decrease necessarily at each step. As predicted by Propositions 8 and 9 we have $\mathbb{V}(\mathbb{G}^{(N)}(f)) > \mathbb{V}(\mathbb{G}^{(0)}(f))$ for $N = 1, 2, 3$ and $\mathbb{V}(\mathbb{G}^{(1)}(f)) > \mathbb{V}(\mathbb{G}^{(3)}(f))$.

.3 Raked empirical means over a class

General framework. Many specific settings in statistics may be modeled through \mathcal{F} . Typically \mathcal{X} is of very large or infinite dimension and each $f(X)$ is one variable with mean $P(f)$ in the population. To control correlations between such variables one needs to extend \mathcal{F} into $\mathcal{F}_\times = \{fg : f, g \in \mathcal{F}\}$ and consider the covariance process $\alpha_n^{(N)}(\mathcal{F}_\times)$. Random vectors $(Y_1, \dots, Y_k) = (f_1(X), \dots, f_k(X))$ can in turn be combined into real valued random variables $g_\theta(X) = \varphi_\theta(Y_1, \dots, Y_k)$ through parameters θ and functions g_θ that should be included in \mathcal{F} and so on. Consider for instance $g_\theta(X) = \theta_1 Y_1 + \dots + \theta_k Y_k + \varepsilon_\sigma(X)$ with a collection of possible residual functions ε_σ turning part of the randomness of X into a noise with variance σ^2 . The (VC) or (BR) entropy of \mathcal{F} rules the variety and complexity of models or statistics one can simultaneously deal with. We refer to Pollard [15], Shorack and Wellner [16] and Wellner [22] for classical statistical models where an empirical process indexed by functions is easily identified.

Direct applications. Since the limiting process $\mathbb{G}^{(N)}$ of $\alpha_n^{(N)}$ has less variance than $\mathbb{G}^{(0)}$, Theorem 2.1 can be applied to revisit the limiting behavior of classical estimators or tests by using $\mathbb{P}_n^{(N)}$ instead of \mathbb{P}_n and prove that the induced asymptotic variances or risk decrease. For instance, in the case of goodness of fit tests, the threshold decreases at any given test level while the power increases against any alternative distribution Q that do not satisfy the margin conditions. As a matter of fact, enforcing $\mathbb{P}_n^{(N)}$ to look like P instead of the true Q over all $\mathcal{A}^{(N)}$ makes $\mathbb{P}_n^{(N)}$ go very far from P on sets where Q was already far from P .

Example: two raked distribution functions. Let (X, Y) be a real centered Gaussian vector with covariance matrix $\begin{pmatrix} 3 & -1 \\ -1 & 1 \end{pmatrix}$. We consider the raked joint estimation of the two distribution functions F_X, F_Y . An auxiliary information provides their values at points -2 to 2 , every 0.5 . The class \mathcal{F} we need contains for all $t \in \mathbb{R}$ the functions $f_t^X(x, y) = \mathbf{1}_{]-\infty, t]}(x)$, $f_t^Y(x, y) = \mathbf{1}_{]-\infty, t]}(y)$ thus (VC) holds. For $Z = X, Y$ let $F_{Z,n}^{(N)}(t) = \sum_{Z_i \leq t} \mathbb{P}_n^{(N)}(\{Z_i\})$ be the N -th raked empirical distribution function and write $Z_{(1)} \leq \dots \leq Z_{(n)}$ the order statistics. To exploit at best the information we use $N = 2m$, $F_{X,n}^{(2m-1)}$ and $F_{Y,n}^{(2m)}$. Consider $d_{Z,n}^{(N)} = \sum_{i=1}^{n-1} (Z_{(i+1)} - Z_{(i)}) |F_{Z,n}^{(N)}(Z_{(i+1)}) - F_Z(Z_{(i+1)})|$ which approximates on $[Z_{(1)}, Z_{(n)}]$ the L^1 -distance between $F_{Z,n}^{(N)}$ and F_Z . Denote $\#_{Z,n}^{(N)}$ the random proportion of sample points where $F_{Z,n}^{(N)}$ is closer to F_Z than $F_{Z,n}^{(0)}$. The table below provides Monte-Carlo estimates of $D_{Z,n}^{(N)} = \mathbb{E}(d_{Z,n}^{(N)})$ and $p_{Z,n}^{(N)} = \mathbb{E}(\#_{Z,n}^{(N)})$ from 1000 simulations based on samples of size $n = 200$:

Z	$D_{Z,n}^{(0)}$	$D_{Z,n}^{(10)}$	$D_{Z,n}^{(\infty)}$	$p_{Z,n}^{(10)}$	$p_{Z,n}^{(\infty)}$
X	0.084	0.058	0.065	0.752	0.724
Y	0.085	0.043	0.053	0.731	0.681

This shows some improvement, especially for $N = 10$. For n rather small it seems not always relevant to wait for the stabilization of the algorithm – here denoted $N = \infty$. Our theoretical results provide guaranties only for small N and large n . We also observe on graphical representations that the way $F_{Z,n}^{(N)}$ leaves $F_{Z,n}^{(0)}$ to cross F_Z at the known points tends to accentuate the error at a few short intervals where $F_{Z,n}^{(0)}$ is far from F_Z . This is less and less the case as the auxiliary information partition is refined or the sample size increases.

Example: raked covariance matrices. Given $d \in \mathbb{N}_*$ and f_1, \dots, f_d let $\mathbb{V}(Y)$ denote the covariance matrix of the random vector $Y = (f_1(X), \dots, f_d(X))$ which we assume to be centered for simplicity. Instead of the empirical covariance $\mathbb{V}_n^{(0)}(Y) = n^{-1} \sum_{i=1}^n Y_i^t Y_i$ consider its raked version

$$\mathbb{V}_n^{(N)}(Y) = \left(\left(\mathbb{P}_n^{(N)}(f_i f_j) \right)_{i,j} \right).$$

Let $\|\cdot\|$ denote the Froebenius norm and define

$$\varphi_Y(\alpha_n^{(N)}) = \sqrt{n} \left\| \mathbb{V}_n^{(N)}(Y) - \mathbb{V}(Y) \right\|.$$

In other words,

$$\varphi_Y^2(\alpha_n^{(N)}) = \sum_{i=1}^d \sum_{j=1}^d \left(\alpha_n^{(N)}(f_i f_j) \right)^2, \quad \varphi_Y^2(\mathbb{G}^{(N)}) = \sum_{i=1}^d \sum_{j=1}^d \left(\mathbb{G}^{(N)}(f_i f_j) \right)^2.$$

In the context of Proposition 6 observe that φ_Y is $(\|\cdot\|_{\mathcal{F}}, \|\cdot\|)$ -Lipshitz with parameter $C_1 = d$. Clearly $\varphi_Y(\mathbb{G}^{(N)})$ has a bounded density since $\varphi_Y^2(\mathbb{G}^{(N)})$ is a quadratic form with Gaussian components and has a modified \mathcal{X}^2 distribution. Choosing a finite collection of such φ_Y ensures that $C_2 < +\infty$. More generally by letting (f_1, \dots, f_d) vary among a small entropy infinite subset \mathcal{L}_d of \mathcal{F}^d and imposing some regularity or localization constraints to the f_i one may have $C_2 < +\infty$ while $\{f_i f_j : f_i, f_j \in \mathcal{F}\}$ satisfies (BR). The largest C_2 still works for $\mathcal{L} = \bigcup_{d \leq d_0} \mathcal{L}_d$. Therefore Proposition 6 guaranties that

$$\max_{\substack{0 \leq N \leq N_0 \\ d \leq d_0}} \sup_{\substack{(f_1, \dots, f_d) \in \mathcal{L}_d \\ x > 0}} \left| \mathbb{P} \left(\varphi_Y(\alpha_n^{(N)}) \leq x \right) - \mathbb{P} \left(\varphi_Y(\mathbb{G}^{(N)}) \leq x \right) \right| \leq d_0 d_1 C_2 v_n$$

where it holds, for all $N \leq N_0$, $d \leq d_0$, $(f_1, \dots, f_d) \in \mathcal{L}_d$ and $x > 0$,

$$\mathbb{P} \left(\varphi_Y(\mathbb{G}^{(N)}) \leq x \right) \leq \mathbb{P} \left(\varphi_Y(\mathbb{G}^{(0)}) \leq x \right)$$

by the variance reduction property of Proposition 8. Hence we asymptotically have $\mathbb{P}(\varphi_Y(\alpha_n^{(N)}) \leq x) < \mathbb{P}(\varphi_Y(\alpha_n^{(0)}) \leq x) - \varepsilon$ uniformly among Y such that $\mathbb{P}(\varphi_Y(\mathbb{G}^{(N)}) \leq x) < \mathbb{P}(\varphi_Y(\mathbb{G}^{(0)}) \leq x) - 2\varepsilon$, for any fixed $\varepsilon > 0$.

References

- [1] ALEXANDER, K. S. (1984). Probability inequalities for empirical processes and a law of the iterated logarithm. *The Annals of Probability* **12** 1041–1067. MRMR0757769
- [2] BANKIER, M.D. (1986). Estimators based in several stratified samples with applications to multiple frame surveys. *Journal of the American Statistical Association* **81** 1074–1079.
- [3] BERTHET, P. and MASON, D. M. (2006). Revisiting two strong approximation results of Dudley and Philipp. *IMS Lecture Notes-Monograph Series High Dimensional Probability* **51** 155–172. MRMR2387767
- [4] BINDER, D. A. and THÉBERGE, A (1988). Estimating the variance of raking-ratio estimators. *The Canadian Journal of Statistics* **16** 47–55. MRMR0997121
- [5] BRACKSTONE, G. J. and RAO, J. N. K. (1979). An investigation of raking ratio estimators. *The Indian journal of Statistics* **41** 97–114.
- [6] BROWN, D. T. (1959). A note on approximations to discrete probability distributions. *Information and control* **2** 386–392. MRMR0110598
- [7] CHOUDHRY, G.H. and LEE, H. (1987). Variance estimation for the Canadian Labour Force Survey. *Survey Methodology* **13** 147–161.
- [8] DEMING, W. E. and STEPHAN, F. F. (1940). On a least squares adjustment of a sampled frequency table when the expected marginal totals are known. *The Annals of Mathematical Statistics* **11** 427–444. MRMR0003527
- [9] DEVILLE, J-C. and SÄRNDAL, C-E (1992). Calibration estimators in survey sampling. *Journal of the American Statistical Association* **87** 376–382. MRMR1173804
- [10] DEVILLE, J-C. and SÄRNDAL, C-E (1993). Generalized raking procedures in survey sampling. *Journal of the American Statistical Association* **88** 1013–1020.
- [11] FRANKLIN, J. and LORENZ, J. (1989). On the scaling of multidimensional matrices. *Linear Algebra and its Applications* **114-115** 717–735. MRMR0986904
- [12] IRELAND, C. T. and KULLBACK, S. (1968). Contingency tables with given marginals. *Biometrika* **55** 179–188. MRMR0229329
- [13] KONIJN, H. S. (1981). Biases, variances and covariances of raking ratio estimators for marginal and cell totals and averages of observed characteristics. *Metrika* **28** 109–121. MRMR0629356

- [14] LEWIS, P. M. (1959). Approximating probability distributions to reduce storage requirements. *Information and control* **2** 214–225. MRMR0110597
- [15] POLLARD, D. (1990). Empirical processes: theory and applications. *NSF-CBMS regional conference series in probability and statistics* MRMR1089429
- [16] SHORACK, G. R. and WELLNER, J. A. (1986). Empirical processes with applications to statistics. *Wiley Series in Probability and Mathematical Statistics: Probability and Mathematical Statistics* MRMR838963
- [17] SINKHORN, R. (1964). A relationship between arbitrary positive matrices and doubly stochastic matrices. *The Annals of Mathematical Statistics* **35** 876–879. MRMR0161868
- [18] SINKHORN, R. (1967). Diagonal equivalence to matrices with prescribed row and column sums. *The American Mathematical Monthly* **74** 402–405. MRMR0210730
- [19] STEPHAN, F. F. (1942). An iterative method of adjusting sample frequency tables when expected marginal totals are known. *The Annals of Mathematical Statistics* **13** 166–178. MRMR0006674
- [20] TALAGRAND, M. (1994). Sharper bounds for Gaussian and empirical processes. *The Annals of Probability* **22** 28–76. MR1258865
- [21] VAN DER VAART, A. W. and WELLNER, J. A.. Weak convergence and empirical processes with applications to statistics. *Springer series in statistics*
- [22] WELLNER, J. A. (1992). Empirical processes in action: a review. *International Statistical Review/Revue Internationale de Statistique* **60** 247–269.