



HAL
open science

A CONSERVATION LAW WITH SPATIALLY LOCALIZED SUBLINEAR DAMPING

Christophe Besse, Rémi Carles, Sylvain Ervedoza

► **To cite this version:**

Christophe Besse, Rémi Carles, Sylvain Ervedoza. A CONSERVATION LAW WITH SPATIALLY LOCALIZED SUBLINEAR DAMPING. 2018. hal-01741189v1

HAL Id: hal-01741189

<https://hal.science/hal-01741189v1>

Preprint submitted on 23 Mar 2018 (v1), last revised 1 Mar 2019 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A CONSERVATION LAW WITH SPATIALLY LOCALIZED SUBLINEAR DAMPING

CHRISTOPHE BESSE, RÉMI CARLES, AND SYLVAIN ERVEDOZA

ABSTRACT. We consider a general conservation law on the circle, in the presence of a sublinear damping. If the damping acts on the whole circle, then the solution becomes identically zero in finite time, following the same mechanism as the corresponding ordinary differential equation. When the damping acts only locally in space, we show a dichotomy: if the flux function is not zero at the origin, then the transport mechanism causes the extinction of the solution in finite time, as in the first case. On the other hand, if zero is a non-degenerate critical point of the flux function, then the solution becomes extinct in finite time only inside the damping zone, decays algebraically uniformly in space, and we exhibit a boundary layer, shrinking with time, around the damping zone. Numerical illustrations show how similar phenomena may be expected for other equations.

1. INTRODUCTION

We consider a general conservation law on the torus $\mathbb{T} = \mathbb{R}/\mathbb{Z}$, in the presence of a sublinear damping, possibly localized in space,

$$(1.1) \quad \partial_t u + \partial_x (f(u)) + a(x) \frac{u}{|u|^\alpha} = 0, \quad (t, x) \in \mathbb{R}_+ \times \mathbb{T},$$

with a smooth flux $f \in \mathcal{C}^\infty(\mathbb{R}, \mathbb{R})$, $0 < \alpha \leq 1$ and $a = a(x) \geq 0$. For the Cauchy problem, we prescribe the initial datum

$$(1.2) \quad u|_{t=0} = u_0, \quad x \in \mathbb{T}.$$

In the case where $a > 0$ is constant, the sublinear nonlinearity is motivated by the effect of friction forces that occur in almost every mechanism with moving parts, this process arising between all surfaces in contact. The first concepts go back to the work of Leonardo da Vinci on friction, rediscovered by Amontons [4] at the end of the 17th century, and then developed by Coulomb [16] in the 18th century. The main idea is that the friction is opposed to the movement and that the friction force is independent of the speed v and the contact surface. The friction force, known today as Coulomb friction, is therefore described as $F = F_c \operatorname{sgn}(v)$. Depending on how the sign function is defined, it can be zero or take any value in the interval $[-F_c, F_c]$. In the 19th century, the theory of hydrodynamics was developed leading to expressions for the frictional force caused by the viscosity of lubricants, and is usually modeled by $F = F_v v$. The linearity with respect to speed is not always correct and a more general relation is $F = F_v |v|^{\delta_v} \operatorname{sgn}(v)$ where δ_v depends on the geometry of the application (see e.g. [23, 1, 3] and references therein). The basic model for the motion of a body

Date: March 23, 2018.

S. E. was partially supported by IFSMACS ANR-15-CE40-0010 of the French National Research Agency (ANR).

lying on a surface is given by the Newton law. It reduces to the ordinary differential equation, for $\alpha \in (0, 1)$,

$$(1.3) \quad \dot{u} = -\frac{u}{|u|^\alpha}, \quad t \in \mathbb{R}, \quad u(0) = u_0 \in \mathbb{R}.$$

By separating the variables, explicit integration yields, in terms of $\rho = u^2$, since $\rho \geq 0$,

$$(1.4) \quad \dot{\rho} = -2\rho^{1-\alpha/2}, \quad \text{hence } \rho(t) = \begin{cases} (|u_0|^\alpha - \alpha t)^{2/\alpha} & \text{if } t \leq |u_0|^\alpha/\alpha, \\ 0 & \text{if } t > |u_0|^\alpha/\alpha. \end{cases}$$

Therefore, ρ becomes zero in finite time, and so does u . Note that for $\alpha = 1$, the equation (1.3) should be understood in the sense of Filippov (see [18, Chapter 2]): $u' \in -\text{Sign}(u)$, in which $\text{Sign}(u)$ is defined by

$$(1.5) \quad \text{Sign}(u) = \begin{cases} \{1\} & \text{if } u > 0, \\ \{-1\} & \text{if } u < 0, \\ [-1, 1] & \text{if } u = 0, \end{cases}$$

and the same argument as above still applies. Besides, note that solutions of the above ODE (1.3), whether $\alpha \in (0, 1)$ or $\alpha = 1$, are unique in positive time even if the source term is not \mathcal{C}^1 with respect to u , as a consequence of the one-sided Lipschitz condition satisfied by $h_\alpha(u) = -u/|u|^\alpha$, see [18, Chapter 2, Section 10, Theorem 1], which reads as follows: for all $(u, v) \in \mathbb{R}^2$,

$$(u - v)(h_\alpha(u) - h_\alpha(v)) \leq 0.$$

Such sublinear damping models have been considered for some partial differential equations: in the case of the wave equation [5, 24], in the case of various parabolic equations [2, 7, 8, 9, 10, 11], and in the case of the Schrödinger equation [12, 13]. The aspect that we now wish to investigate is the effect of such a damping when it is localized in space. Typically, the function a in (1.1) can be thought of as an indicating function.

Some of the results that we present can be adapted to the case where the space variable belongs to the whole line \mathbb{R} . The reason why we consider the periodic case is the following. On the whole line, the characteristics of the solution of (1.1) may cross the support of a without undergoing such a strong affect as in (1.4), that is, the sublinear damping occurs in too small a region to put u to zero. On the other hand, in a periodic box, and in the case where transport phenomenon is present, the solution will meet the support of a as long as it is not zero. These are typically the possibilities which we want to understand.

Assumption 1.1. *The function a is nonnegative, $a(x) \geq 0$ for all $x \in \mathbb{T}$, bounded, $a \in L^\infty(\mathbb{T})$, and satisfies*

$$\sup_{y>0} \frac{1}{y} \int_{\mathbb{T}} |a(x+y) - a(x)| \, dx < \infty.$$

Typically, this condition is satisfied for $a \in BV(\mathbb{T})$, see [17, Chapter 1 Theorem 1.7.1] (in fact, this is nearly equivalent of being in $BV(\mathbb{T})$). In particular, a may be an indicating function, $a(x) = \mathbf{1}_\omega(x)$ for some measurable set $\omega \subset \mathbb{T}$.

1.1. Cauchy problem. The notion of solution, as well as the vanishing viscosity method used to solve the Cauchy problem, follow from standard arguments (which we borrow from [17]). We shall see that the presence of the damping term in (1.1) requires only slight modifications of this approach. We emphasize however that the case $\alpha = 1$ is specific, and we shall treat it by adapting the approach of Filippov [18].

Definition 1.2 (Notion of solution, $0 < \alpha < 1$). *Let $\alpha \in (0, 1)$. A bounded measurable function u on $[0, T] \times \mathbb{T}$ is an admissible weak solution of (1.1)–(1.2), with $u_0 \in L^\infty(\mathbb{T})$, if the inequality*

$$(1.6) \quad \int_0^T \int_{\mathbb{T}} \left(\partial_t \psi \eta(u) + \partial_x \psi q(u) - \alpha \psi \eta'(u) \frac{u}{|u|^\alpha} \right) dx dt + \int_{\mathbb{T}} \psi(0, x) \eta(u_0(x)) dx \geq 0$$

holds for every convex function $\eta \in W^{1, \infty}$, with $q' = f'\eta'$, and all nonnegative Lipschitz continuous test function ψ on $[0, T] \times \mathbb{T}$.

Definition 1.3 (Notion of solution, $\alpha = 1$). *Let $\alpha = 1$. A bounded measurable function u on $[0, T] \times \mathbb{T}$ is an admissible weak solution of (1.1)–(1.2), with $u_0 \in L^\infty(\mathbb{T})$, if there exists $h \in L^\infty((0, T) \times \mathbb{T})$ such that*

$$\partial_t u + \partial_x(f(u)) + h = 0, \quad \text{in } \mathcal{D}'((0, T) \times \mathbb{T}),$$

with

$$h(t, x) \in a(x) \text{Sign}(u(t, x)), \quad \text{a.e. } (t, x) \in (0, T) \times \mathbb{T},$$

where Sign is defined in (1.5), and such that the inequality

$$(1.7) \quad \int_0^T \int_{\mathbb{T}} (\partial_t \psi \eta(u) + \partial_x \psi q(u) - \psi \eta'(u) h(t, x)) dx dt + \int_{\mathbb{T}} \psi(0, x) \eta(u_0(x)) dx \geq 0$$

holds for every convex function $\eta \in W^{1, \infty}$, with $q' = f'\eta'$, and all nonnegative Lipschitz continuous test function ψ on $[0, T] \times \mathbb{T}$.

In all that follows, the notion of solution refers either to Definition 1.2 (case $0 < \alpha < 1$), or to Definition 1.3 (case $\alpha = 1$). We show that the Cauchy problem is well-posed, regardless of the value of $\alpha \in (0, 1]$.

Proposition 1.4 (Cauchy problem). *Assume that a satisfies Assumption 1.1 and $\alpha \in (0, 1]$. Let $u_0 \in L^\infty(\mathbb{T})$. There exists a unique, global, admissible weak solution u of (1.1)–(1.2), $u \in \mathcal{C}^0(\mathbb{R}_+; L^1(\mathbb{T}))$.*

We will also need the following comparison result.

Proposition 1.5 (Comparison principles). *Let $\alpha \in (0, 1]$.*

1. Let a satisfying Assumption 1.1, u and v be solutions of (1.1) with respective initial data $u_0, v_0 \in L^\infty(\mathbb{T})$ such that $u_0 \leq v_0$. Then

$$u(t, x) \leq v(t, x), \quad \forall t \geq 0, \quad \text{a.e. } x \in \mathbb{T}.$$

Besides,

$$|u(t, x)| \leq \|u_0\|_{L^\infty(\mathbb{T})}, \quad \forall t \geq 0, \quad \text{a.e. } x \in \mathbb{T}.$$

2. Let a_1 and a_2 satisfying Assumption 1.1 such that for almost all $x \in \mathbb{T}$, $a_1(x) \leq a_2(x)$. Then, denoting by u_1 and u_2 the respective solutions to (1.1)–(1.2) with the same initial datum $u_0 \in L^\infty(\mathbb{T})$ with $u_0 \geq 0$, we have

$$u_1(t, x) \geq u_2(t, x) \geq 0, \quad \forall t \geq 0, \quad \text{a.e. } x \in \mathbb{T}.$$

Remark 1.6 (BV solutions). As a straightforward consequence of the proof of Proposition 1.5, given in Section 2, one can show that if $u_0 \in BV(\mathbb{T})$, then the solution remains in BV , $u \in L^\infty(\mathbb{R}_+; BV(\mathbb{T}))$. However, we shall not use this property in this paper.

1.2. Extinction results. We now focus on the core of this article, and give several results regarding the possible extinction of the solutions u of (1.1)–(1.2). The results depend on the flux f (its behavior near the origin), and the damping coefficient a , which is always assumed to satisfy Assumption 1.1.

The first case which we consider is the one corresponding to a damping coefficient acting everywhere.

Proposition 1.7 (Finite time extinction with damping everywhere). *Suppose that there exists $\delta > 0$ such that*

$$(1.8) \quad a(x) \geq \delta > 0, \quad \forall x \in \mathbb{T}.$$

Let $u_0 \in L^\infty(\mathbb{T})$. There exists $T > 0$ such that the solution to (1.1)–(1.2) satisfies

$$u(t, x) = 0, \quad \forall t \geq T, \text{ a.e. } x \in \mathbb{T}.$$

Besides, T can be chosen as

$$(1.9) \quad T = \frac{1}{\alpha\delta} \|u_0\|_{L^\infty(\mathbb{T})}^\alpha.$$

The proof of Proposition 1.7 is presented in Section 3 and is based on a Lyapunov approach. More precisely, we derive $L^p(\mathbb{T})$ estimates on the solutions of (1.1)–(1.2), and let then p go to infinity, so that we obtain a differential inequality for the $L^\infty(\mathbb{T})$ -norm of the solutions of (1.1)–(1.2), which in turn implies its extinction in finite time.

Next, as motivated above, we consider the case in which the damping coefficient acts only in some part of the domain:

$$(1.10) \quad \exists \text{ an open interval } \omega \subset \mathbb{T} \text{ and } \delta > 0 \text{ s.t. } a(x) \geq \delta, \quad \forall x \in \omega.$$

The extinction of the solution of (1.1)–(1.2) in this case will depend on the flux. Namely, we will treat two different cases, depending whether $f'(0)$ vanishes or not. The easier case corresponds to the presence of transport at the origin,

$$(1.11) \quad f'(0) \neq 0.$$

In this case, one expects that the transport phenomenon will steer the solution through the set ω an arbitrary number of times, so that the strong friction term will make the solution vanish after some finite time.

In agreement with these insights, in Section 4 we prove the following result:

Theorem 1.8 (Finite time extinction by transport). *Assume that the flux f is smooth and satisfies (1.11), and the damping profile a satisfies Assumption 1.1 and (1.10). Let K be such that*

$$(1.12) \quad \inf_{s \in [-K, K]} |f'(s)| > 0.$$

Then for any initial datum $u_0 \in L^\infty(\mathbb{T})$ satisfying

$$(1.13) \quad \|u_0\|_{L^\infty(\mathbb{T})} \leq K,$$

there exists $T > 0$ such that the solution u of (1.1)–(1.2) satisfies

$$(1.14) \quad u(t, x) = 0, \quad \forall t \geq T, \text{ a.e. } x \in \mathbb{T}.$$

To illustrate the typical behavior of such a solution, we plot on Figure 1.1 the evolution of the solution of the transport equation corresponding to $f(u) = 2u$ with initial datum $u_0(x) = 1.25$ for $(t, x) \in [0, 10] \times (0, 1)$, $a(x) = \mathbf{1}_{(0, 1/4)}$ and $\alpha = 1$. The solution is computed using the numerical procedure described in Section 6. The dashed line indicates the position of the support of a . Note that Theorem 1.8 is




FIGURE 1.1. Evolution of the solution for transport equation

a semi-global result, as it is valid for any initial datum u_0 whose $L^\infty(\mathbb{T})$ -norm is bounded by the constant K , which is chosen in (1.12) to guarantee that the transport phenomenon really occurs.

Our proof of Theorem 1.8 relies on Lyapunov functionals, in a similar spirit as the one developed in Section 3 to address the proof of Proposition 1.7. However, as the transport phenomenon is now essential to the decay process, we will introduce some weights in space in the functional. This approach is inspired by some recent works on the stabilization of hyperbolic systems of conservation laws, namely [15, 14] (see the recent book [6] for further references).

In the case

$$(1.15) \quad f'(0) = 0,$$

corresponding for instance to the celebrated example of Burgers equation

$$(1.16) \quad f(u) = \frac{u^2}{2},$$

the transport phenomenon competes with the dissipation of the solution, as the smaller the solution is, the slower the characteristics propagate. Our goal thus is to understand the interplay between these phenomena.

Theorem 1.9. *Let f be a smooth function such that*

$$(1.17) \quad f'(0) = 0 \quad \text{and} \quad \exists K > 0, \quad \inf_{s \in [-K, K]} |f''(s)| > 0.$$

Assume that the damping profile $a = a(x)$ satisfies Assumption 1.1 and (1.10). Then, for any initial datum $u_0 \in L^\infty(\mathbb{T})$ satisfying (1.13), the solution u of (1.1)–(1.2) satisfies the following property: There exist a time $t_ > 0$ and a constant $C > 0$ such that for all $t \geq t_*$, there exists an open subinterval $\omega(t) \subset \omega$ such that $u(t)|_{\omega(t)} = 0$, and, for all $t \geq t_*$,*

$$|\omega \setminus \omega(t)| \leq \frac{C}{t^{1+\alpha}}, \quad \|u(t)\|_{L^\infty(\mathbb{T})} \leq \frac{C}{t}.$$

The proof of Theorem 1.9 is based on a precise description of the solution corresponding to $u_0 = K$, where K is constant, \mathbb{T} is identified with $(0, 1)$ with periodic boundary conditions, and

$$(1.18) \quad a(x) = \delta \mathbf{1}_\omega, \quad \text{with } \omega = (0, A).$$

We can then deduce Theorem 1.9 from a simple comparison argument based on Proposition 1.5.

It is clear from these results that there still are cases which are not covered from our results, in particular cases in which the initial datum has an $L^\infty(\mathbb{T})$ -norm which is larger than the best K in (1.12) or (1.17). Some particular instance is numerically studied in Section 6, as well as other models for which the complete understanding of a (localized) strong friction on the dynamics of a system is still not well understood.

1.3. Outline of the paper. The Cauchy problem is addressed in Section 2, where Propositions 1.4 and 1.5 are established. The proof of Proposition 1.7 (finite time extinction with damping everywhere) is given in Section 3, thanks to suitable Lyapunov functionals. Theorem 1.8 (finite time extinction by transport) is proved in Section 4, by introducing refined Lyapunov functionals. For the case of the generalized Burgers equation, Theorem 1.9, a longer Section 5 is needed, where we first construct a rather explicit solution by following characteristics, which then turns out to be the solution provided by Proposition 1.4. Finally, Section 6 provides numerical illustrations in the case of (1.1) studied in this paper, as well as in the case of other equations for which the corresponding analysis turns out to be a challenging issue.

Acknowledgements. The authors wish to thank Jean-François Coulombel, Frédéric Lagoutière, Philippe Laurençot and Vincent Perrollaz for fruitful discussions.

2. CAUCHY PROBLEM AND COMPARISON PRINCIPLE

In this section, we prove Propositions 1.4 and 1.5.

We consider more generally the Cauchy problem

$$(2.1) \quad \partial_t u + \partial_x f(u) + h(x, u) = 0 \quad \text{in } \mathbb{R}_+ \times \mathbb{T}, \quad u|_{t=0} = u_0 \quad \text{in } \mathbb{T},$$

with a fairly general semilinear term h , possibly depending on x , in order to generalize the nonlinearity, typically of the form

$$(2.2) \quad h_\alpha(x, u) = a(x) \frac{u}{|u|^\alpha},$$

where $\alpha \leq 1$ corresponds to the case of (1.1), with the modification detailed in Definition 1.3 in the case $\alpha = 1$. We will use the following properties on the source term h , which encompass the framework of Definition 1.2 when $\alpha < 1$.

Assumption 2.1. *The map $h = h(x, u)$ satisfies:*

- $h \in L_{\text{loc}}^\infty(\mathbb{T} \times \mathbb{R})$.
- For all $u \in \mathbb{R}$, $h(x, u)u \geq 0$, for almost all $x \in \mathbb{T}$.
- For almost all fixed $x \in \mathbb{T}$, the map $u \mapsto h(x, u)$ is nondecreasing on \mathbb{R} .
- For every $R > 0$,

$$\sup_{|u| \leq R} \sup_{y > 0} \frac{1}{y} \int_{\mathbb{T}} |h(x+y, u) - h(x, u)| dx < \infty.$$

In the case (2.2), the first property corresponds to the assumption $a \in L^\infty(\mathbb{T})$, the second property to the fact that h is a damping term, the third property is straightforward (even in the case $\alpha = 1$ with the approach of Filippov), and the last property is a consequence of Assumption 1.1.

Definition 2.2 (Notion of solution). *Let h satisfy Assumption 2.1. A bounded measurable function u on $[0, T] \times \mathbb{T}$ is an admissible weak solution of (2.1), with $u_0 \in L^\infty(\mathbb{T})$, if the inequality*

$$(2.3) \quad \int_0^T \int_{\mathbb{T}} (\partial_t \psi \eta(u) + \partial_x \psi q(u) - \psi \eta'(u) h(x, u)) dx dt + \int_{\mathbb{T}} \psi(0, x) \eta(u_0(x)) dx \geq 0$$

holds for every convex function $\eta \in W^{1, \infty}$, with $q' = f'\eta'$, and all nonnegative Lipschitz continuous test function ψ on $[0, T] \times \mathbb{T}$.

Propositions 1.4 and 1.5 stem from the following result, which is slightly more general in view of Assumption 2.1.

Proposition 2.3. *Let Assumption 2.1 be satisfied.*

- (i) *Let $u_0 \in L^\infty(\mathbb{T})$. There exists a unique, global, admissible weak solution u of (2.1), $u \in \mathcal{C}^0(\mathbb{R}_+; L^1(\mathbb{T}))$.*
- (ii) *Let u and v be solutions of (2.1) with respective initial data $u_0, v_0 \in L^\infty(\mathbb{T})$, and $u_0(x) \leq v_0(x)$ for almost all $x \in \mathbb{T}$. Then*

$$(2.4) \quad u(t, x) \leq v(t, x), \quad \forall t \geq 0, \quad \text{a.e. } x \in \mathbb{T}.$$

Besides,

$$(2.5) \quad |u(t, x)| \leq \|u_0\|_{L^\infty(\mathbb{T})}, \quad \forall t \geq 0, \quad \text{a.e. } x \in \mathbb{T}.$$

- (iii) *If $h^{(1)}$ and $h^{(2)}$ satisfy Assumption 2.1, and in addition,*

$$(0 \leq) h^{(1)}(x, u) \leq h^{(2)}(x, u), \quad \text{a.e. } (x, u) \in \mathbb{T} \times (0, \infty),$$

then denoting by u_1 and u_2 the respective solutions to (2.1) with the same initial datum $u_0 \in L^\infty(\mathbb{T})$, $u_0 \geq 0$, we have

$$(2.6) \quad u_1(t, x) \geq u_2(t, x) \geq 0, \quad \forall t \geq 0, \quad \text{a.e. } x \in \mathbb{T}.$$

- (iv) *The same properties hold true for solutions of (1.1)–(1.2) for $\alpha = 1$, when considering solutions in the sense of Definition 1.3 and a satisfies Assumption 1.1.*

Proof. When Assumption 2.1 is satisfied, we follow very closely the approach of [17, Section 6.3], based on the method of vanishing viscosity.

• **Uniqueness.** For u and \underline{u} two solutions in the sense of Definition 2.2, introduce the entropy-entropy flux pair

$$\eta(u, \underline{u}) = |u - \underline{u}|, \quad q(u, \underline{u}) = \text{sign}(u - \underline{u}) (f(u) - f(\underline{u})).$$

This is an entropy-entropy flux pair for u when \underline{u} is fixed, and, conversely, for \underline{u} when u is fixed. For a nonnegative test function $\phi = \phi(t, x, \underline{t}, \underline{x})$, following successively the two points of view mentioned above, we have,

$$\begin{aligned} & \int_0^T \int_{\mathbb{T}} (\partial_t \phi(t, x, \underline{t}, \underline{x}) \eta(u(t, x), \underline{u}(\underline{t}, \underline{x})) + \partial_x \phi(t, x, \underline{t}, \underline{x}) q(u(t, x), \underline{u}(\underline{t}, \underline{x}))) dx dt \\ & \quad - \int_0^T \int_{\mathbb{T}} \phi(t, x, \underline{t}, \underline{x}) \text{sign}(u(t, x) - \underline{u}(\underline{t}, \underline{x})) h(x, u(t, x)) dx dt \\ & \quad + \int_{\mathbb{T}} \phi(0, x, \underline{t}, \underline{x}) \eta(u_0(x), \underline{u}(\underline{t}, \underline{x})) dx \geq 0, \quad (t, \underline{x}) \in \mathbb{R}_+ \times \mathbb{T}, \\ \\ & \int_0^T \int_{\mathbb{T}} (\partial_{\underline{t}} \phi(t, x, \underline{t}, \underline{x}) \eta(u(t, x), \underline{u}(\underline{t}, \underline{x})) + \partial_{\underline{x}} \phi(t, x, \underline{t}, \underline{x}) q(u(t, x), \underline{u}(\underline{t}, \underline{x}))) d\underline{x} d\underline{t} \\ & \quad - \int_0^T \int_{\mathbb{T}} \phi(t, x, \underline{t}, \underline{x}) \text{sign}(u(t, x) - \underline{u}(\underline{t}, \underline{x})) h(x, u(t, x)) d\underline{x} d\underline{t} \\ & \quad + \int_{\mathbb{T}} \phi(t, x, 0, \underline{x}) \eta(u(t, x), \underline{u}_0(\underline{x})) d\underline{x} \geq 0, \quad (t, x) \in \mathbb{R}_+ \times \mathbb{T}. \end{aligned}$$

Integrating each of these inequalities with respect to the last two variables and summing up the two resulting inequalities yields

$$\begin{aligned} & \int_0^T \int_0^T \iint_{\mathbb{T}^2} (\partial_t + \partial_{\underline{t}}) \phi(t, x, \underline{t}, \underline{x}) \eta(u(t, x), \underline{u}(\underline{t}, \underline{x})) dx d\underline{x} dt d\underline{t} \\ & + \int_0^T \int_0^T \iint_{\mathbb{T}^2} (\partial_x + \partial_{\underline{x}}) \phi(t, x, \underline{t}, \underline{x}) q(u(t, x), \underline{u}(\underline{t}, \underline{x})) dx d\underline{x} dt d\underline{t} \\ & - \int_0^T \int_0^T \iint_{\mathbb{T}^2} \phi(t, x, \underline{t}, \underline{x}) \text{sign}(u(t, x) - \underline{u}(\underline{t}, \underline{x})) (h(x, u(t, x)) - h(\underline{x}, \underline{u}(\underline{t}, \underline{x}))) dx d\underline{x} dt d\underline{t} \\ & + \int_0^T \iint_{\mathbb{T}^2} \phi(0, x, \underline{t}, \underline{x}) \eta(u_0(x), \underline{u}(\underline{t}, \underline{x})) dx d\underline{x} dt \\ & + \int_0^T \iint_{\mathbb{T}^2} \phi(t, x, 0, \underline{x}) \eta(u(t, x), \underline{u}_0(0, \underline{x})) dx d\underline{x} dt \geq 0. \end{aligned}$$

Pick ϕ of the form

$$\phi(t, x, \underline{t}, \underline{x}) = \frac{1}{\varepsilon^2} \psi \left(\frac{t + \underline{t}}{2} \right) \rho \left(\frac{t - \underline{t}}{2\varepsilon} \right) \rho \left(\frac{x - \underline{x}}{2\varepsilon} \right),$$

where the nonnegative function ψ depends only on time, and the nonnegative compactly supported function ρ is such that $\int_{\mathbb{R}} \rho = 1$, with an obvious abuse of notation for the last factor above. Letting $\varepsilon \rightarrow 0$, we find:

$$\begin{aligned} & \int_0^T \int_{\mathbb{T}} \psi'(t) \eta(u(t, x), \underline{u}(t, x)) dx dt \\ & \quad - \int_0^T \int_{\mathbb{T}} \psi(t) \text{sign}(u(t, x) - \underline{u}(t, x)) (h(x, u(t, x)) - h(x, \underline{u}(t, x))) dx dt \\ & \quad + \int_{\mathbb{T}} \psi(0) \eta(u_0(x), \underline{u}_0(x)) dx \geq 0. \end{aligned}$$

In view of the third point in Assumption 2.1, this implies

$$\int_0^T \int_{\mathbb{T}} \psi'(t) \eta(u(t, x), \underline{u}(t, x)) dx dt + \int_{\mathbb{T}} \psi(0) \eta(u_0(x), \underline{u}_0(x)) dx \geq 0.$$

For $0 < \tau < T$ and $k \in \mathbb{N}$, we define $\psi = \psi_k$ as

$$\psi_k(t) = \begin{cases} 1 & \text{if } 0 \leq t < \tau, \\ k(\tau - t) + 1 & \text{if } \tau \leq t < \tau + \frac{1}{k}, \\ 0 & \text{if } \tau + \frac{1}{k} \leq t < T. \end{cases}$$

Letting $k \rightarrow \infty$ now yields

$$\int_{\mathbb{T}} \eta(u_0(x), \underline{u}_0(x)) dx \geq \int_{\mathbb{T}} \eta(u(\tau, x), \underline{u}(\tau, x)) dx,$$

that is $\|u(\tau) - \underline{u}(\tau)\|_{L^1(\mathbb{T})} \leq \|u_0 - \underline{u}_0\|_{L^1(\mathbb{T})}$, hence uniqueness for solutions in the sense of Definition 2.2, since $\tau \in (0, T)$ is arbitrary.

• **Viscous approximation.** For $\mu > 0$, consider the equation

$$(2.7) \quad \partial_t u_\mu + \partial_x f(u_\mu) + h(x, u_\mu) = \mu \partial_x^2 u_\mu \text{ in } \mathbb{R}_+ \times \mathbb{T}, \quad u_\mu|_{t=0} = u_0 \text{ in } \mathbb{T}.$$

For a fixed $\mu > 0$, the solution to (2.7) is obtained by a fixed point argument applied to the associated Duhamel's formula,

$$(2.8) \quad u_\mu(t, x) = e^{\mu t \partial_x^2} u_0(x) - \int_0^t e^{\mu(t-s) \partial_x^2} (\partial_x f(u_\mu(s, x)) + h(x, u_\mu(s, x))) ds,$$

where we recall that the heat semigroup on \mathbb{T} acts on Fourier series as

$$e^{\mu t \partial_x^2} \left(\sum_{n \in \mathbb{Z}} a_n e^{i2\pi n x} \right) = \sum_{n \in \mathbb{Z}} a_n e^{i2\pi n x - 4\mu \pi^2 n^2 t}.$$

Up to considering the linear heat flow on \mathbb{R} and its uniqueness property to relate it to the heat flow on \mathbb{T} , we readily see that if $u_0 \in L^\infty(\mathbb{T})$, there exists $T > 0$ depending on $\|u_0\|_{L^\infty(\mathbb{T})}$ only and a unique solution $u_\mu \in \mathcal{C}^0([0, T]; L^\infty(\mathbb{T}))$ to (2.8).

• **A priori estimate.** The solution is global in time, $u_\mu \in \mathcal{C}^0(\mathbb{R}_+; L^\infty(\mathbb{T}))$, in view of the a priori estimate

$$(2.9) \quad \|u_\mu(t)\|_{L^\infty(\mathbb{T})} \leq \|u_0\|_{L^\infty(\mathbb{T})}, \quad \forall t \geq 0,$$

which we now establish. For $p > 2$, multiply (2.7) by $|u_\mu|^{p-2} u_\mu$, and integrate over \mathbb{T} . This yields

$$\begin{aligned} \frac{1}{p} \frac{d}{dt} \left(\int_{\mathbb{T}} |u_\mu|^p \right) + \int_{\mathbb{T}} f'(u_\mu) |u_\mu|^{p-2} u_\mu \partial_x u_\mu &= - \int_{\mathbb{T}} h(x, u_\mu) |u_\mu|^{p-2} u_\mu \\ &\quad - \mu(p-1) \int_{\mathbb{T}} |u_\mu|^{p-2} (\partial_x u_\mu)^2. \end{aligned}$$

If g is such that $g'(y) = p f'(y) |y|^{p-2} y$, the second term on the left hand side is

$$\int_{\mathbb{T}} g'(u_\mu) \partial_x u_\mu = \int_{\mathbb{T}} \partial_x g(u_\mu) = 0.$$

The first term on the right hand side is non-positive in view of Assumption 2.1, and the last term is obviously non-positive. We infer, for all $p > 2$,

$$\frac{d}{dt} \|u_\mu\|_{L^p(\mathbb{T})}^p \leq 0,$$

hence (2.9) by integrating in time and letting $p \rightarrow \infty$.

• **Entropy solution for (2.1).** The next step consists in showing that up to extracting a subsequence, u_μ converges to an entropy solution (Definition 2.2). The above uniqueness result shows that actually, no extraction is needed.

If η is a smooth convex entropy, with associated entropy flux q , multiplying (2.7) by $\eta'(u_\mu)$ yields

$$\partial_t \eta(u_\mu) + \partial_x q(u_\mu) + \eta'(u_\mu) h(x, u_\mu) = \mu \partial_x^2 \eta(u_\mu) - \mu \eta''(u_\mu) (\partial_x u_\mu)^2.$$

Multiply the above equation by a nonnegative test function ψ then yields, after an integration by parts and since $\eta'' \geq 0$,

$$(2.10) \quad \int_0^T \int_{\mathbb{T}} (\partial_t \psi \eta(u_\mu) + \partial_x \psi q(u_\mu) - \psi \eta'(u_\mu) h(x, u_\mu)) dx dt + \int_{\mathbb{T}} \psi(0, x) \eta(u_0(x)) dx \geq -\mu \int_0^T \int_{\mathbb{T}} \partial_x^2 \psi \eta(u_\mu) dx dt.$$

The same holds when we just have $\eta \in W^{1, \infty}$ by using an approximating entropy, as in the next paragraph.

Therefore, if for some sequence (μ_k) with $\mu_k \downarrow 0$, $(u_{\mu_k})_k$ converges to some function u , boundedly almost everywhere on $[0, \infty) \times \mathbb{T}$, then u is an admissible weak solution of (2.1) on $[0, \infty) \times \mathbb{T}$.

• **Comparison for the viscous solution.** Introduce as in the proof of [17, Theorem 6.3.2] the function η_ε defined for $\varepsilon > 0$ by

$$\eta_\varepsilon(w) = \begin{cases} 0 & \text{for } -\infty < w \leq 0, \\ \frac{w^2}{4\varepsilon} & \text{for } 0 < w \leq 2\varepsilon, \\ w - \varepsilon & \text{for } 2\varepsilon < w < \infty. \end{cases}$$

If u_μ and \bar{u}_μ solve (2.8), then by multiplying by $\eta'_\varepsilon(u_\mu - \bar{u}_\mu)$ the equation satisfied by $u_\mu - \bar{u}_\mu$, we compute

$$\begin{aligned} & \partial_t \eta_\varepsilon(u_\mu - \bar{u}_\mu) + \partial_x (\eta'_\varepsilon(u_\mu - \bar{u}_\mu) (f(u_\mu) - f(\bar{u}_\mu))) \\ & \quad - \eta''_\varepsilon(u_\mu - \bar{u}_\mu) (f(u_\mu) - f(\bar{u}_\mu)) \partial_x (u_\mu - \bar{u}_\mu) \\ & = -\eta'_\varepsilon(u_\mu - \bar{u}_\mu) (h(x, u_\mu) - h(x, \bar{u}_\mu)) + \mu \partial_x^2 \eta_\varepsilon(u_\mu - \bar{u}_\mu) - \mu \eta''_\varepsilon(u_\mu - \bar{u}_\mu) (\partial_x (u_\mu - \bar{u}_\mu))^2. \end{aligned}$$

The new term compared to the proof of [17, Theorem 6.3.2] is of course the first term of the right hand side (where h is present). We have more precisely, for $0 < s < t < \infty$, after integration on $(s, t) \times \mathbb{T}$,

$$(2.11) \quad \begin{aligned} & \int_{\mathbb{T}} \eta_\varepsilon(u_\mu(t) - \bar{u}_\mu(t)) - \int_{\mathbb{T}} \eta_\varepsilon(u_\mu(s) - \bar{u}_\mu(s)) \\ & \leq \int_s^t \int_{\mathbb{T}} \eta''_\varepsilon(u_\mu - \bar{u}_\mu) (f(u_\mu) - f(\bar{u}_\mu)) \partial_x (u_\mu - \bar{u}_\mu) \\ & \quad - \int_s^t \int_{\mathbb{T}} \eta'_\varepsilon(u_\mu - \bar{u}_\mu) (h(x, u_\mu) - h(x, \bar{u}_\mu)). \end{aligned}$$

In the limit $\varepsilon \rightarrow 0$, the first term on the right hand side goes to zero, while the second goes to

$$- \int_s^t \int_{\mathbb{T}} \mathbf{1}_{u_\mu > \bar{u}_\mu} (h(x, u_\mu) - h(x, \bar{u}_\mu)).$$

By Assumption 2.1 (third point), this term is non-positive, and we infer

$$\int_{\mathbb{T}} (u_\mu(t, x) - \bar{u}_\mu(t, x))_+ dx \leq \int_{\mathbb{T}} (u_\mu(s, x) - \bar{u}_\mu(s, x))_+ dx.$$

By letting $s \rightarrow 0$, this implies

$$(2.12) \quad \int_{\mathbb{T}} (u_\mu(t, x) - \bar{u}_\mu(t, x))_+ dx \leq \int_{\mathbb{T}} (u_0(x) - \bar{u}_0(x))_+ dx,$$

and by interchanging the roles of u_μ and \bar{u}_μ ,

$$\|u_\mu(t) - \bar{u}_\mu(t)\|_{L^1(\mathbb{T})} \leq \|u_0 - \bar{u}_0\|_{L^1(\mathbb{T})}.$$

Also, if

$$u_0(x) \leq \bar{u}_0(x), \quad \text{a.e. on } \mathbb{T},$$

then (2.12) yields

$$u_\mu(t, x) \leq \bar{u}_\mu(t, x), \quad \forall t \geq 0, \quad \text{a.e. } x \in \mathbb{T}.$$

This implies in particular uniqueness for (2.7).

• **Compactness.** To obtain compactness in space, as in [17], we consider $\bar{u}_\mu(t, x) = u_\mu(t, x + y)$. In the case where $h = 0$ (or more generally if h depends on u only), then \bar{u}_μ is a solution to (2.7), so (2.12) can be used directly. In our case, and precisely because we want to consider spatially localized damping, such \bar{u}_μ does not solve (2.7), and we have to resume the computations. Essentially, we go back to the previous computations, and replace $\bar{u}_\mu(t, x)$ with $u_\mu(t, x + y)$, noticing that $h(x, \bar{u}_\mu)$ has to be replaced by $h(x + y, u_\mu(t, x + y))$. We have

$$\begin{aligned} & \int_{\mathbb{T}} (u_\mu(t, x) - u_\mu(t, x + y))_+ dx - \int_{\mathbb{T}} (u_0(x) - u_0(x + y))_+ dx \leq \\ & \limsup_{\varepsilon \rightarrow 0} \left(- \int_0^t \int_{\mathbb{T}} \eta'_\varepsilon(u_\mu - \bar{u}_\mu) (h(x, u_\mu(\tau, x)) - h(x + y, u_\mu(\tau, x + y))) d\tau dx \right). \end{aligned}$$

In the above integral, insert $\pm h(x + y, u_\mu(\tau, x))$. By the same argument as above (third point in Assumption 2.1), we infer

$$\begin{aligned} & \int_{\mathbb{T}} (u_\mu(t, x) - u_\mu(t, x + y))_+ dx - \int_{\mathbb{T}} (u_0(x) - u_0(x + y))_+ dx \leq \\ & \limsup_{\varepsilon \rightarrow 0} \left(- \int_0^t \int_{\mathbb{T}} \eta'_\varepsilon(u_\mu - \bar{u}_\mu) (h(x, u_\mu(\tau, x)) - h(x + y, u_\mu(\tau, x))) d\tau dx \right), \end{aligned}$$

hence

$$\begin{aligned} & \int_{\mathbb{T}} (u_\mu(t, x) - u_\mu(t, x + y))_+ dx - \int_{\mathbb{T}} (u_0(x) - u_0(x + y))_+ dx \\ & \leq \int_0^t \int_{\mathbb{T}} |h(x, u_\mu(\tau, x)) - h(x + y, u_\mu(\tau, x))| d\tau dx. \end{aligned}$$

In view of (2.9) and of the last point in Assumption 2.1, we conclude

$$\int_{\mathbb{T}} (u_\mu(t, x) - u_\mu(t, x + y))_+ dx \leq \int_{\mathbb{T}} (u_0(x) - u_0(x + y))_+ dx + \mathcal{O}(y),$$

and

$$(2.13) \quad \int_{\mathbb{T}} |u_\mu(t, x) - u_\mu(t, x + y)| dx \leq \int_{\mathbb{T}} |u_0(x) - u_0(x + y)| dx + \mathcal{O}(y).$$

Equicontinuity in time is proved similarly by setting $\bar{u}_\mu(t, x) = u_\mu(t + \tau, x)$. Since h depends on x and u_μ only, the only extra term that we have to estimate is of the form

$$\left| \int_t^{t+\tau} \int_{\mathbb{T}} h(x, u_\mu(s, x)) \phi(x) ds d\tau \right| \leq \tau C (\|u_0\|_{L^\infty(\mathbb{T})}) \|\phi\|_{L^1(\mathbb{T})},$$

where we have used (2.9).

The above properties imply that the sequence $(u_\mu)_\mu$ is uniformly bounded and equicontinuous in $(0, \infty) \times \mathbb{T}$, so there is a subsequence of $(u_\mu)_\mu$, which converges boundedly almost everywhere on $(0, \infty) \times \mathbb{T}$ and strongly in $L^1_{\text{loc}}((0, \infty) \times \mathbb{T})$. We infer from (2.10) that the limit u is an entropy solution, hence the entropy solution by uniqueness of the entropy solution. We have thus proved the item (i), while item (ii) follows from (2.12) and (2.9), after passing to the limit $\mu \rightarrow 0$.

Remark 2.4. Dividing (2.13) by y yields the propagation of BV regularity mentioned in Remark 1.6.

• **Comparison when source terms are ordered.** It remains to prove (iii). Since we assume $u_0 \geq 0$, we know from (ii) that $u_1(t, x), u_2(t, x) \geq 0$ for $(t, x) \in (0, \infty) \times \mathbb{T}$, and so $h^{(2)}(x, u_j) \geq h^{(1)}(x, u_j)$ for $j = 1, 2$. We then consider the viscous approximations $u_{1,\mu}$ and $u_{2,\mu}$ of, respectively, u_1 and u_2 . The analogue of (2.11) reads

$$\begin{aligned} & \int_{\mathbb{T}} \eta_\varepsilon(u_{2,\mu}(t) - u_{1,\mu}(t)) - \int_{\mathbb{T}} \eta_\varepsilon(u_{2,\mu}(s) - u_{1,\mu}(s)) \leq \\ & \int_s^t \int_{\mathbb{T}} \eta''_\varepsilon(u_{2,\mu} - u_{1,\mu}) (f(u_{2,\mu}) - f(u_{1,\mu})) \partial_x(u_{2,\mu} - u_{1,\mu}) \\ & - \int_s^t \int_{\mathbb{T}} \eta'_\varepsilon(u_{2,\mu} - u_{1,\mu}) \left(h^{(2)}(x, u_{2,\mu}) - h^{(1)}(x, u_{1,\mu}) \right). \end{aligned}$$

Passing to the limits $\varepsilon \rightarrow 0$, $s \rightarrow 0$, and finally $\mu \rightarrow 0$, we infer, since u_1 and u_2 have the same initial datum:

$$\int_{\mathbb{T}} (u_2(t) - u_1(t))_+ \leq - \int_0^t \int_{\mathbb{T}} \mathbf{1}_{u_2 > u_1} \left(h^{(2)}(x, u_2) - h^{(1)}(x, u_1) \right).$$

Now since the integrand of the right hand side can be decomposed as

$$\underbrace{\mathbf{1}_{u_2 > u_1} \left(h^{(2)}(x, u_2) - h^{(2)}(x, u_1) \right)}_{\geq 0, \text{ by Assumption 2.1}} + \underbrace{\mathbf{1}_{u_2 > u_1} \left(h^{(2)}(x, u_1) - h^{(1)}(x, u_1) \right)}_{\geq 0, \text{ from above}},$$

we conclude that $\int_{\mathbb{T}} (u_2(t) - u_1(t))_+ = 0$, hence $u_1 \geq u_2 \geq 0$ as announced.

• **Item (iv): the case $h(x, u) = a(x)u/|u|$.**

In this case, the uniqueness of admissible weak solutions in the sense of Definition 1.3 holds without change. The difficulty then is to prove existence of admissible weak solutions. In order to do that, instead of approximating (1.1) by its viscous approximation (2.7), we also add an approximation of the function h . Namely, we consider the approximation given, for $\mu \in (0, 1)$, by

$$(2.14) \quad \partial_t u_\mu + \partial_x f(u_\mu) + h_{1-\mu}(x, u_\mu) = \mu \partial_x^2 u_\mu \text{ in } \mathbb{R}_+ \times \mathbb{T}, \quad u_\mu|_{t=0} = u_0 \text{ in } \mathbb{T},$$

where $h_{1-\mu}(x, u)$ is given by (2.2). For each $\mu > 0$, all the computations performed above can be repeated, so that the sequence of solutions $(u_\mu)_\mu$ is uniformly bounded and equicontinuous in $(0, \infty) \times \mathbb{T}$, so up to some subsequence, it converges boundedly almost everywhere on $(0, \infty) \times \mathbb{T}$ and strongly in $L^1_{\text{loc}}((0, \infty) \times \mathbb{T})$ to some u as $\mu \rightarrow 0$, so that a.e. $(t, x) \in (0, \infty) \times \mathbb{T}$,

$$h_{1-\mu}(x, u_\mu(t, x)) \xrightarrow{\mu \rightarrow 0} h(t, x),$$

where $h(t, x) = a(x) \frac{u(t, x)}{|u(t, x)|}$ if $u(t, x) \neq 0$, and $h(t, x) \in [-1, 1]$ if $u(t, x) = 0$.

It follows that u is an admissible weak solutions of (1.1)–(1.2) in the sense of Definition 1.3, hence the admissible weak solution in the sense of Definition 1.3 by

uniqueness. The comparison results can then be proved as before, by studying them for the solutions of (2.14) and passing to the limit $\mu \rightarrow 0$. \square

3. PROOF OF PROPOSITION 1.7: THE CASE OF A DAMPING ACTING EVERYWHERE

Proof of Proposition 1.7. Let $p > 2$. In view of (1.1), we formally have

$$\frac{d}{dt} \int_{\mathbb{T}} |u(t, x)|^p dx = p \int_{\mathbb{T}} |u|^{p-2} u \partial_t u dx = -p \int_{\mathbb{T}} |u|^{p-2} u \partial_x f(u) dx - p \int_{\mathbb{T}} a(x) |u|^{p-\alpha} dx.$$

Writing formally $p|u|^{p-2} u \partial_x f(u) = p|u|^{p-2} u f'(u) \partial_x u$, we have

$$p|u|^{p-2} u \partial_x f(u) = \partial_x g_p(u), \quad \text{with } g_p'(z) = p|z|^{p-2} z f'(z),$$

and so

$$\int_{\mathbb{T}} |u|^{p-2} u \partial_x f(u) dx = 0.$$

As a matter of fact, this reasoning is valid only for sufficiently smooth solutions. However, the conclusion remains true in our context, as can be seen by using Definitions 1.2 and 1.3 with $p > 2$, $\eta(u) = |u|^p u$, $\psi(t, x) = 1$:

$$\frac{d}{dt} \int_{\mathbb{T}} |u(t, x)|^p dx \leq -p \int_{\mathbb{T}} a(x) |u|^{p-\alpha} dx \leq -\frac{p\delta}{\|u(t)\|_{L^\infty(\mathbb{T})}^\alpha} \int_{\mathbb{T}} |u(t, x)|^p dx,$$

that is

$$\frac{d}{dt} \ln \|u(t)\|_{L^p(\mathbb{T})} \leq -\frac{\delta}{\|u(t)\|_{L^\infty(\mathbb{T})}^\alpha}.$$

By integration,

$$\|u(t)\|_{L^p(\mathbb{T})} \leq \|u_0\|_{L^p(\mathbb{T})} \exp\left(-\delta \int_0^t \frac{ds}{\|u(s)\|_{L^\infty(\mathbb{T})}^\alpha}\right),$$

and by letting $p \rightarrow \infty$,

$$(3.1) \quad \|u(t)\|_{L^\infty(\mathbb{T})} \leq \|u_0\|_{L^\infty(\mathbb{T})} \exp\left(-\delta \int_0^t \frac{ds}{\|u(s)\|_{L^\infty(\mathbb{T})}^\alpha}\right).$$

Set

$$\Phi(t) = \int_0^t \frac{ds}{\|u(s)\|_{L^\infty(\mathbb{T})}^\alpha}.$$

The above inequality reads

$$\Phi'(t) \geq \frac{1}{\|u_0\|_{L^\infty(\mathbb{T})}^\alpha} e^{\alpha\delta\Phi(t)}.$$

Since the solution to $\Psi' = C e^{\alpha\delta\Psi}$, $\Psi(0) = 0$, is given by

$$\Psi(t) = -\frac{1}{\alpha\delta} \ln(1 - C\alpha\delta t), \quad t \leq \frac{1}{C\alpha\delta},$$

we conclude by comparison that $\Phi(t) \rightarrow +\infty$ as $t \rightarrow \|u_0\|_{L^\infty(\mathbb{T})}^\alpha / (\alpha\delta)$, hence the result thanks to (3.1), with

$$T = \frac{\|u_0\|_{L^\infty(\mathbb{T})}^\alpha}{\alpha\delta},$$

which may not be the sharp extinction time, but an upper bound for it. \square

Remark 3.1 (Whole space). The above argument shows that the conclusion of Proposition 1.7 remains valid if (1.8) is set up on the whole line, $x \in \mathbb{R}$, provided that the solution which we consider goes to zero at $\pm\infty$.

Otherwise, a similar proof can be given, by considering estimates of $u(t)$ in $L^p(A_-(t), A_+(t))$ where

$$A_-(t) = A_-^0 + t \sup_{s \in [-K, K]} f'(s) \quad \text{and} \quad A_+(t) = A_+^0 + t \sup_{s \in [-K, K]} f'(s)$$

for any pair $(A_-^0, A_+^0) \in \mathbb{R}^2$; see e.g. the proof of Lemma 4.2 where the same kind of arguments are developed.

4. PROOF OF THEOREM 1.8: THE TRANSPORT CASE

The goal of this section is to prove Theorem 1.8. Thus, we consider the setting of Theorem 1.8, and we assume in particular that f is smooth, satisfies (1.11), K satisfies (1.12), and a satisfies Assumption 1.1 and (1.10).

4.1. Strategy. In order to ease the reading of the proof of Theorem 1.8, we decompose it into two lemmas, the first one stating that in the setting of Theorem 1.8 the solutions of (1.1)–(1.2) decay exponentially, while the second one will show that if the initial datum is small enough, then the corresponding solution of (1.1)–(1.2) vanishes in finite time.

Lemma 4.1. *Within the setting of Theorem 1.8, there exist $C > 0$ and $\mu > 0$ such that for any initial datum $u_0 \in L^\infty(\mathbb{T})$ satisfying (1.13), any solution u of (1.1)–(1.2) satisfies*

$$(4.1) \quad \|u(t)\|_{L^\infty(\mathbb{T})} \leq C e^{-\mu t} \|u_0\|_{L^\infty(\mathbb{T})}.$$

Lemma 4.2. *Within the setting of Theorem 1.8, there exist $\varepsilon_0 > 0$ and $T_0 > 0$ such that if*

$$(4.2) \quad \|u_0\|_{L^\infty(\mathbb{T})} \leq \varepsilon_0,$$

the solution u of (1.1)–(1.2) satisfies

$$(4.3) \quad u(t, x) = 0, \quad \forall t \geq T_0, \quad a.e. \quad x \in \mathbb{T}.$$

The proofs of Lemma 4.1 and Lemma 4.2 are given in Subsections 4.2 and 4.3, respectively. The proof of Theorem 1.8 is then given in Subsection 4.4.

4.2. Proof of Lemma 4.1.

Proof. We first choose a function $\varphi = \varphi(x)$ such that

$$\varphi \in \mathcal{C}^\infty(\mathbb{T}), \quad \text{with } \varphi(x) = x, \quad \forall x \in \mathbb{T} \setminus \omega.$$

Note that such a function φ satisfies in particular that

$$\forall x \in \mathbb{T} \setminus \omega, \quad \partial_x \varphi(x) = 1, \quad \text{and} \quad \partial_x \varphi \in \mathcal{C}^0(\mathbb{T}).$$

Now, let $u_0 \in L^\infty(\mathbb{T})$ and u the corresponding solution of (1.1)–(1.2). We consider the Lyapunov functionals, indexed by $p \geq 2$ and some parameter $\lambda \in \mathbb{R}$ chosen later,

$$(4.4) \quad E_{p, \lambda}(t) = \int_{\mathbb{T}} \left| e^{-\lambda \varphi(x)} u(t, x) \right|^p dx.$$

Formally, these functionals satisfy:

$$\begin{aligned} \frac{dE_{p,\lambda}(t)}{dt} &= p \int_{\mathbb{T}} e^{-p\lambda\varphi(x)} |u(t,x)|^{p-2} u(t,x) \partial_t u(t,x) dx \\ &= -p \int_{\mathbb{T}} e^{-p\lambda\varphi(x)} |u(t,x)|^{p-2} u(t,x) \partial_x (f(u(t,x))) dx \\ &\quad - p \int_{\mathbb{T}} a(x) e^{-p\lambda\varphi(x)} |u(t,x)|^{p-\alpha} dx. \end{aligned}$$

If u were smooth, we would write

$$\begin{aligned} p|u(t,x)|^{p-2} u(t,x) \partial_x (f(u(t,x))) &= p f'(u(t,x)) |u(t,x)|^{p-2} u(t,x) \partial_x u(t,x) \\ &= \partial_x (g_p(u(t,x))), \end{aligned}$$

where g_p is defined by

$$(4.5) \quad g_p(s) = p \int_0^s f'(\tau) |\tau|^{p-2} \tau d\tau,$$

so that we would write:

$$\begin{aligned} -p \int_{\mathbb{T}} e^{-p\lambda\varphi(x)} |u(t,x)|^{p-2} u(t,x) \partial_x (f(u(t,x))) dx \\ = -p\lambda \int_{\mathbb{T}} \partial_x \varphi(x) e^{-p\lambda\varphi(x)} g_p(u(t,x)) dx. \end{aligned}$$

Note in passing that g_p has the same sign as $f'(0)$. As solutions u may contain shocks, these estimates should be justified by using the definition of admissible weak solutions, i.e. inequalities (1.6) or (1.7), choosing $p > 2$, $\eta(u) = |u|^p u$ and $\psi(t,x) = e^{-p\lambda\varphi(x)}$. One obtains in that way:

$$\begin{aligned} \frac{dE_{p,\lambda}(t)}{dt} &\leq -p\lambda \int_{\mathbb{T}} \partial_x \varphi(x) e^{-p\lambda\varphi(x)} g_p(u(t,x)) dx \\ &\quad - p \int_{\mathbb{T}} a(x) e^{-p\lambda\varphi(x)} |u(t,x)|^{p-\alpha} dx \\ &\leq -p\lambda \int_{\mathbb{T}} e^{-p\lambda\varphi(x)} g_p(u(t,x)) dx \\ &\quad + p|\lambda| \|\partial_x \varphi - 1\|_{L^\infty(\mathbb{T})} \int_{\omega} e^{-p\lambda\varphi(x)} |g_p(u(t,x))| dx \\ &\quad - p\delta \int_{\omega} e^{-p\lambda\varphi(x)} |u(t,x)|^{p-\alpha} dx. \end{aligned}$$

It is thus natural to study the function g_p in (4.5). In order to do this, we first note that if $\|u_0\|_{L^\infty(\mathbb{T})} \leq K$, according to Proposition 1.5, for all time $t \geq 0$, the L^∞ -norm of $u(t)$ is bounded by K . Therefore, we introduce

$$\beta_- = \inf_{s \in [-K, K]} |f'(s)| \quad \text{and} \quad \beta_+ = \sup_{s \in [-K, K]} |f'(s)|,$$

so that for all $(t,x) \in [0, \infty) \times \mathbb{T}$,

$$(4.6) \quad \beta_- |u(t,x)|^p \leq |g_p(u(t,x))| \leq \beta_+ |u(t,x)|^p.$$

Therefore, we obtain

$$\begin{aligned} \frac{dE_{p,\lambda}(t)}{dt} &\leq -p\lambda \int_{\mathbb{T}} e^{-p\lambda\varphi(x)} g_p(u(t,x)) dx \\ &\quad + p \left(|\lambda| \|\partial_x \varphi - 1\|_{L^\infty(\mathbb{T})} \beta_+ - \delta \|u(t)\|_{L^\infty(\mathbb{T})}^{-\alpha} \right) \int_{\omega} e^{-p\lambda\varphi(x)} |u(t,x)|^p dx. \end{aligned}$$

In particular, for all $t \geq 0$, if

$$(4.7) \quad |\lambda| \|\partial_x \varphi - 1\|_{L^\infty(\mathbb{T})} \beta_+ \leq \delta \|u(t)\|_{L^\infty(\mathbb{T})}^{-\alpha},$$

we have

$$(4.8) \quad \frac{dE_{p,\lambda}(t)}{dt} \leq -p\lambda \int_{\mathbb{T}} e^{-p\lambda\varphi(x)} g_p(u(t,x)) dx.$$

As for all t , $\|u(t)\|_{L^\infty(\mathbb{T})} \leq K$, we therefore choose

$$\lambda_* = \text{sign}(f'(0)) \frac{\delta K^{-\alpha}}{\|\partial_x \varphi - 1\|_{L^\infty(\mathbb{T})} \beta_+},$$

so that (4.7) is satisfied and (4.8) becomes:

$$\frac{dE_{p,\lambda_*}(t)}{dt} \leq -p|\lambda_*| \beta_- E_{p,\lambda_*}(t).$$

Therefore, we get that

$$\|e^{-\lambda_* \varphi} u(t)\|_{L^p(\mathbb{T})} \leq e^{-|\lambda_*| \beta_- t} \|e^{-\lambda_* \varphi} u_0\|_{L^p(\mathbb{T})}, \quad \forall t \geq 0.$$

As λ_* does not depend on p , we can pass to the limit as $p \rightarrow \infty$, and obtain:

$$\|e^{-\lambda_* \varphi} u(t)\|_{L^\infty(\mathbb{T})} \leq e^{-|\lambda_*| \beta_- t} \|e^{-\lambda_* \varphi} u_0\|_{L^\infty(\mathbb{T})}, \quad \forall t \geq 0,$$

and thus,

$$\|u(t)\|_{L^\infty(\mathbb{T})} \leq e^{|\lambda_*|(\sup \varphi - \inf \varphi) - |\lambda_*| \beta_- t} \|u_0\|_{L^\infty(\mathbb{T})}, \quad \forall t \geq 0.$$

This concludes the proof of Lemma 4.1, as $|\lambda_*| > 0$. \square

Remark 4.3. Note that one can go further and show that the $L^\infty(\mathbb{T})$ norm of the solutions of (1.1)–(1.2) decays in fact faster than an exponential in time.

Indeed, if we set

$$T_1 = \frac{2}{\beta_-} (\sup \varphi - \inf \varphi),$$

and use the explicit choice

$$\lambda_* = \text{sign}(f'(0)) \frac{\delta \|u_0\|_{L^\infty(\mathbb{T})}^{-\alpha}}{\|\partial_x \varphi - 1\|_{L^\infty(\mathbb{T})} \beta_+},$$

which is admissible according to the above proof, one in fact gets

$$\begin{aligned} \|u(T_1)\|_{L^\infty(\mathbb{T})} &\leq e^{-|\lambda_*|(\sup \varphi - \inf \varphi)} \|u_0\|_{L^\infty(\mathbb{T})} \\ &\leq \exp\left(-c_0 \|u_0\|_{L^\infty(\mathbb{T})}^{-\alpha}\right) \|u_0\|_{L^\infty(\mathbb{T})}, \end{aligned}$$

where c_0 is given by

$$c_0 = \delta \frac{(\sup \varphi - \inf \varphi)}{\|\partial_x \varphi - 1\|_{L^\infty(\mathbb{T})} \beta_+}.$$

Starting from there and using the semi-group property, we introduce a sequence of time, indexed by $n \in \mathbb{N}$,

$$T_n = nT_1 \quad \left(= \frac{2n}{\beta_-} (\sup \varphi - \inf \varphi) \right),$$

for which one gets immediately, for all $n \in \mathbb{N}$,

$$\|u(T_{n+1})\|_{L^\infty(\mathbb{T})} \leq \exp\left(-c_0 \|u(T_n)\|_{L^\infty(\mathbb{T})}^{-\alpha}\right) \|u(T_n)\|_{L^\infty(\mathbb{T})}.$$

It is then easy to check that the sequence $(\|u(T_n)\|_{L^\infty(\mathbb{T})})_{n \in \mathbb{N}}$ goes to 0 faster than any (non-trivial) geometric sequence, which in turn implies that the map $t \mapsto \|u(t)\|_{L^\infty(\mathbb{T})}$ goes to 0 faster than any exponential.

It would be interesting to develop a direct proof of Theorem 1.8 based only on a suitable choice of Lyapunov functionals in the spirit of the one used above.

4.3. Proof of Lemma 4.2.

Proof. To simplify the presentation, in the proof of this lemma, \mathbb{T} is identified with an interval centered in 0, and ω is identified with an interval of the form $(-A, A)$.

The proof of Lemma 4.2 is divided in two steps. In the first step, we show that if $\varepsilon_0 > 0$ is chosen small enough, then necessarily, the solution u of (1.1)–(1.2) with $u_0 \in L^\infty(\mathbb{T})$ satisfying $\|u_0\|_{L^\infty(\mathbb{T})} \leq \varepsilon_0$ vanishes in some part of the domain ω after some (small) time. We then show that this implies that the solution u vanishes everywhere after some time.

• **Step 1.** We introduce the paths

$$A_-(t) = \sup_{[-K, K]} \{f'\}t - \frac{A}{2}, \quad A_+(t) = \inf_{[-K, K]} \{f'\}t + \frac{A}{2}.$$

We fix τ_* such for all $t \in [0, \tau_*)$,

$$-A < A_-(t) < A_+(t) < A,$$

that is

$$(4.9) \quad \tau_* = \frac{A}{\max\left\{2 \inf_{[-K, K]} |f'|, \sup_{[-K, K]} |f'| - \inf_{[-K, K]} |f'|\right\}}.$$

We then set, for $p \geq 2$ and $t \in [0, \tau_*]$,

$$E_{p, \text{loc}}(t) = \int_{A_-(t)}^{A_+(t)} |u(t, x)|^p dx.$$

Arguing as in the proof of Proposition 1.7, as $[A_-(t), A_+(t)] \subset \omega$ for all $t \in [0, \tau_*]$ and $\inf f' \leq f'(u(t, x)) \leq \sup f'$ for all $(t, x) \in \mathbb{R}_+ \times \mathbb{T}$, we get, for all $t \geq 0$,

$$(4.10) \quad \frac{d}{dt} E_{p, \text{loc}}(t) \leq - \frac{p\delta}{\|u(t)\|_{L^\infty(A_-(t), A_+(t))}^\alpha} E_{p, \text{loc}}(t).$$

In fact, to prove this estimate rigorously, we use the definition of admissible weak solutions, i.e. the inequalities (1.6) or (1.7), choosing $p > 2$, $\eta(u) = |u|^p u$ and $\psi(t, x) = \varphi_\epsilon(t, x)$, where

$$\begin{aligned} \varphi_\epsilon(t, x) = & \varphi_-^0 \left(\frac{x - A_-(t)}{\epsilon} \right) \mathbf{1}_{x \in (A_-(t) - \epsilon, A_-(t))} + \mathbf{1}_{x \in (A_-(t), A_+(t))} \\ & + \varphi_+^0 \left(\frac{x - A_+(t)}{\epsilon} \right) \mathbf{1}_{x \in (A_+(t), A_+(t) + \epsilon)}, \end{aligned}$$

with φ_-^0, φ_+^0 non-negative smooth cut-off functions, taking value 1 on \mathbb{R}_+ and vanishing on $(-\infty, -1)$ for φ_-^0 , taking value 1 on \mathbb{R}_- and vanishing on $(1, \infty)$ for φ_+^0 . We then pass to the limit $\epsilon \rightarrow 0$ in the inequalities (1.6) or (1.7) to show the estimate (4.10).

This yields, for all $t \in [0, \tau_*]$,

$$\frac{d}{dt} (\log(\|u\|_{L^p(A_-(t), A_+(t))})) \leq -\frac{\delta}{\|u(t)\|_{L^\infty(A_-(t), A_+(t))}^\alpha}.$$

Integrating this expression and letting $p \rightarrow \infty$, we obtain, similarly as in (3.1), that for all $t \in [0, \tau_*]$,

$$\|u(t)\|_{L^\infty(A_-(t), A_+(t))} \leq \|u_0\|_{L^\infty(-A/2, A/2)} \exp\left(-\delta \int_0^t \frac{ds}{\|u(s)\|_{L^\infty(A_-(s), A_+(s))}^\alpha}\right).$$

Arguing as in the proof of Proposition 1.7, this implies that $\|u(\tau_0)\|_{L^\infty(A_-(\tau_0), A_+(\tau_0))} = 0$ for

$$(4.11) \quad \tau_0 = \frac{\varepsilon_0^\alpha}{\alpha\delta},$$

which is smaller than τ_* for $\varepsilon_0 > 0$ small enough, i.e. for small enough initial datum (recall (4.2)).

Using the same argument on the solutions $u(\cdot + t_0)$ for all $t_0 \geq 0$, we see that in fact we have obtained

$$(4.12) \quad \forall t \geq \tau_0, \quad \|u(t)\|_{L^\infty(A_-(\tau_0), A_+(\tau_0))} = 0.$$

We end up this step by emphasizing that $A_-(\tau_0) < A_+(\tau_0)$, so that (4.12) really implies that $u(t, \cdot)$ vanishes on a constant interval for all time $t \geq \tau_0$.

• **Step 2.** In this step, to fix the ideas, we assume that $f'(0) > 0$, as a completely similar proof can be adapted to the case $f'(0) < 0$. We then look at the evolution of the L^2 -norm of $u(t)$ on the set $(A_-(\tau_0), B(t))$, where $B(t) = A_+(\tau_0) + \beta_-(t - \tau_0)_+$, with $\beta_- = \inf_{[-K, K]} f'$ (recall that we have assumed $f'(0) > 0$). Recall that $u(t, x) = 0$ for all $t \geq \tau_0$ and $x \in [A_-(\tau_0), A_+(\tau_0)]$ according to (4.12). Using the definition of admissible weak solutions, we infer:

$$\frac{d}{dt} \left(\int_{A_-(\tau_0)}^{B(t)} |u(t, x)|^2 dx \right) \leq 0.$$

Indeed, this comes from the definition of admissible weak solutions with $\eta(u) = |u|^2$ and $\psi(t, x) = \varphi_\epsilon(t, x)$, where

$$\begin{aligned} \varphi_\epsilon(t, x) &= \varphi_-^0 \left(\frac{x - A_+(\tau_0)}{\epsilon} \right) \mathbf{1}_{x \in (A_+(\tau_0) - \epsilon, A_+(\tau_0))} + \mathbf{1}_{x \in (A_+(\tau_0), B(t))} \\ &\quad + \varphi_+^0 \left(\frac{x - B(t)}{\epsilon} \right) \mathbf{1}_{x \in (B(t), B(t) + \epsilon)}, \end{aligned}$$

with φ_-^0, φ_+^0 non-negative smooth cut-off functions, taking value 1 on \mathbb{R}_+ and vanishing on $(-\infty, -1)$ for φ_-^0 , taking value 1 on \mathbb{R}_- and vanishing on $(1, \infty)$ for φ_+^0 .

Therefore, for all $t \geq \tau_0$,

$$\int_{A_-(\tau_0)}^{B(t)} |u(t, x)|^2 dx = 0.$$

In particular, waiting a time $T_0 > \tau_0$ such that $B(T_0) = A_-(\tau_0) + |\mathbb{T}|$, for all $t \geq T_0$, for almost all $x \in \mathbb{T}$, $u(t, x) = 0$. This concludes the proof of Lemma 4.2. \square

4.4. Proof of Theorem 1.8.

Proof. Theorem 1.8 easily follows from Lemma 4.1 and 4.2. Indeed, if one chooses an initial datum u_0 satisfying (1.13), the $L^\infty(\mathbb{T})$ -norm of the corresponding solution $u(t)$ of (1.1)–(1.2) decays exponentially. Thus, after some time, it becomes smaller than the parameter ε_0 in Lemma 4.2. It will therefore vanish after some time according to Lemma 4.2. \square

4.5. A control theoretic interpretation of Theorem 1.8. Let us mention that Theorem 1.8 is closely related to the following control problem: given ω a non-empty subinterval of \mathbb{T} , $T > 0$ and $u_0 \in L^2(\mathbb{T})$, find a control function $v \in L^2((0, T) \times \mathbb{T})$ such that the solution u of

$$(4.13) \quad \begin{cases} \partial_t u + \partial_x (f(u)) + \mathbf{1}_\omega v = 0, & (t, x) \in \mathbb{R}_+ \times \mathbb{T}, \\ u|_{t=0} = u_0, & x \in \mathbb{T}. \end{cases}$$

satisfies

$$(4.14) \quad u(T) = 0 \quad \text{in } \mathbb{T}.$$

Assuming (1.11) and defining K by (1.12), Theorem 1.8 implies that, for an initial datum $u_0 \in L^\infty(\mathbb{T})$ satisfying (1.13), choosing the control function v under the feedback form

$$(4.15) \quad v(t, x) = -\delta \frac{u(t, x)}{|u(t, x)|^\alpha}, \quad (t, x) \in \mathbb{R}_+ \times \mathbb{T},$$

for some $\delta > 0$, the controlled trajectory u solving (4.13) will satisfy the controllability requirement (4.14) in some time $T > 0$.

Looking more closely at the proof of Lemma 4.2, we can state the following result:

Proposition 4.4. *Let f be a smooth flux function satisfying (1.11) and define K by (1.12). Let ω a non-empty subinterval of \mathbb{T} .*

Given $\gamma > 0$, there exists a parameter δ in (4.15) such that, for any initial datum $u_0 \in L^\infty(\mathbb{T})$ satisfying (1.13), the corresponding solution u of (4.13)–(4.15) vanishes after the time

$$T = (1 + \gamma) \frac{|\mathbb{T}|}{\inf_{[-K, K]} |f'|}.$$

Proof. We do the same identifications as in the proof of Lemma 4.2. Indeed, choosing $\gamma > 0$ smaller if necessary, one can assume $\gamma|\mathbb{T}|/\inf |f'| < \tau_*$, with τ_* as in (4.9). Thus, taking

$$\delta = \frac{K^\alpha \inf |f'|}{\alpha \gamma |\mathbb{T}|},$$

for solutions $u_0 \in L^\infty(\mathbb{T})$ satisfying (1.13), the time τ_0 in (4.11) is smaller than τ_* in (4.9) and than $\gamma|\mathbb{T}|/\inf |f'|$, so that the proof of Lemma 4.2 easily yields that u vanishes after the time $T_0 = (1 + \gamma)|\mathbb{T}|/\inf |f'|$. \square

Note that the time of extinction given by Proposition 4.4 can be made arbitrarily close to the critical time expected to control (4.13), given by $|\mathbb{T}|/\inf_{[-K, K]} |f'|$. In this sense, we have produced a non-linear feedback operator which controls (4.13) in almost sharp time.

5. PROOF OF THEOREM 1.9: THE DEGENERATE CASE

The goal of this section is to discuss the case of a flux satisfying $f'(0) = 0$, with $f''(0) \neq 0$, and prove Theorem 1.9. As said in the introduction, we first prove Theorem 1.9 in the case of a constant initial datum and a strictly convex flux satisfying (1.17). We then deduce the other instances of Theorem 1.9 by using symmetry arguments and comparison arguments.

5.1. Computation and estimates for the solution u of (5.2) for a strictly convex flux f with $f'(0) = 0$. We first assume that f satisfies, for some $K > 0$:

$$(5.1) \quad f'(0) = 0 \quad \text{and} \quad \exists K > 0, \quad \inf_{s \in [-K, K]} f''(s) > 0,$$

and the damping is given by $a(x) = \delta \mathbf{1}_{(0, A)}$.

Let u be the solution of

$$(5.2) \quad \partial_t u + \partial_x f(u) + \delta \mathbf{1}_{(0, A)} \frac{u}{|u|^\alpha} = 0 \text{ in } \mathbb{R}_+ \times \mathbb{T}, \quad u|_{t=0} = K \text{ in } \mathbb{T},$$

where K is the positive constant in (5.1), $\delta > 0$ and $(0, A) \subset \mathbb{T}$.

Obviously, as $K > 0$, the solution u will stay non-negative for all times.

We develop a precise analysis of the characteristics curves of the solution, illustrated in Figure 5.1. We depict in Figure 5.1 both the evolution of the solution and its characteristic curves for the Burgers equation. The simulations are made following the process described in Section 6. The numerical parameters are $\alpha = 1$, $u_0(x) = K = 1.25$, $A = 0.25$, $\delta = 1$, $\delta x = \delta t = 5 \cdot 10^{-5}$ and the final time is $T_f = 10$. The dashed line indicates the location of the support of a . It appears that the solution becomes zero inside the support of a , with corresponding characteristics becoming vertical straight lines. Our goal in the following is to prove rigorously that this observed behavior indeed coincides with the theoretical one.

5.1.1. Formal computation of the solution u of (5.2) for a strictly convex flux f with $f'(0) = 0$. We shall compute u using characteristics as if the solution were regular. We will fully justify this assumption later (in Subsection 5.1.3, Lemma 5.1).

For $\alpha < 1$, when u is smooth, for $(t_0, x_0) \in \mathbb{R}_+ \times \mathbb{T}$, the characteristics are given by

$$(5.3) \quad \begin{cases} \frac{dX}{dt}(t, t_0, x_0) = f'(u(t, X(t, t_0, x_0))), & t \geq 0, \\ X(t_0, t_0, x_0) = x_0, \end{cases}$$

and the solution u along the characteristics satisfies:

$$(5.4) \quad \frac{d}{dt} (u(t, X(t, t_0, x_0))) = -a(X(t, t_0, x_0)) \frac{u(t, X(t, t_0, x_0))}{|u(t, X(t, t_0, x_0))|^\alpha}.$$

The solutions of (5.3)–(5.4) are then solutions of

$$(5.5) \quad \frac{d}{dt} \begin{pmatrix} X(t, t_0, x_0) \\ u(t, X(t, t_0, x_0)) \end{pmatrix} = F_\alpha(X(t, t_0, x_0), u(t, X(t, t_0, x_0))), \quad t \geq 0,$$

where

$$(5.6) \quad F_\alpha(X, u) = \begin{pmatrix} f'(u) \\ -a(X)u/|u|^\alpha \end{pmatrix}.$$

Of course, when $\alpha = 1$, similar computations can be performed as long as the solution u stays positive, but the corresponding definition of F_α for $\alpha = 1$ should be made




FIGURE 5.1. Evolution of the solution (left) and of the characteristic curves (right) for Burgers flux

more specific when u vanishes. We thus introduce

$$(5.7) \quad F_1(X, u) = \begin{pmatrix} f'(u) \\ -a(X) \text{Sign}(u) \end{pmatrix},$$

where Sign is defined in (1.5), and the corresponding counterpart of (5.5) should then read as

$$(5.8) \quad \frac{d}{dt} \begin{pmatrix} X(t, t_0, x_0) \\ u(t, X(t, t_0, x_0)) \end{pmatrix} \in F_1(X(t, t_0, x_0), u(t, X(t, t_0, x_0))), \quad t \geq 0.$$

In the computations given afterward, we will use the fact that as $u_0(x) = K \geq 0$, the solution u of (5.2) stays non-negative, so that we can in fact write $u/|u|^\alpha = u^{1-\alpha}$, which will make the various expressions slightly easier.

On the interval $(A, 1)$. As a vanishes on $(A, 1)$, (5.4) implies that the solution u stays constant along the characteristics in $(A, 1)$. Therefore, if we choose $x_0 = A$, we get

$$u(t, X(t, t_0, A)) = u(t_0, A), \quad \text{as long as } t \mapsto X(t, t_0, A) \text{ stays smaller than } 1,$$

and therefore

$$X(t, t_0, A) = A + (t - t_0)f'(u(t_0, A)) \quad \text{for } t \in \left[t_0, t_0 + \frac{1 - A}{f'(u(t_0, A))} \right].$$

Thus, we can write, for all $t \geq 0$,

$$(5.9) \quad u \left(t + \frac{1 - A}{f'(u(t, A))}, 0 \right) = u \left(t + \frac{1 - A}{f'(u(t, A))}, 1 \right) = u(t, A).$$

Let us finally note that easy computations show that

$$\forall t \in \left[0, \frac{1-A}{f'(K)}\right], \quad u(t, 0) = u(t, 1) = K.$$

On the interval $(0, A)$. We deal with this case as before. But now, as long as the characteristic $t \mapsto X(t, t_0, 0)$ defined for $t \geq t_0$ stays in $[0, A]$, we have

$$u(t, X(t, t_0, 0)) = (u(t_0, 0)^\alpha - \delta\alpha(t - t_0))_+^{1/\alpha}.$$

Therefore, the characteristic $X(t, t_0, 0)$ reaches $x = A$ for the first solution $t \geq t_0$ (if any) of

$$\int_0^{t-t_0} f' \left((u(t_0, 0)^\alpha - \delta\alpha\tau)_+^{1/\alpha} \right) d\tau = A,$$

for which we have

$$u(t, A) = (u(t_0, 0)^\alpha - \delta\alpha(t - t_0))_+^{1/\alpha}.$$

For $t \geq 0$, such that

$$\int_0^t f' \left((K^\alpha - \delta\alpha\tau)_+^{1/\alpha} \right) d\tau < A,$$

we get

$$u(t, A) = (K^\alpha - \delta\alpha t)_+^{1/\alpha}.$$

5.1.2. *Justification of the above formulae.* When the solution u is smooth and strictly positive, all the above computations are fully justified.

Besides, for $\alpha \in (0, 1]$, we only have *a priori* existence results for solutions of (5.5) or (5.8) due to Cauchy–Peano Theorem. Due to [18, Chapter 2 Section 10 Theorem 1], we also have forward uniqueness of the solutions of (5.5) as long as X stays in $(0, A)$ or as long as X stays in $(A, 1)$, as the function F_α satisfies the following one-sided Lipschitz condition: there exists $C > 0$ such that for all $(X_1, u_1) \in \mathbb{R}^2$, $(X_2, u_2) \in \mathbb{R}^2$ with $|u_1|, |u_2| \leq K$, and $(X_1, X_2) \in (0, A)^2 \cup (A, 1)^2$,

$$(5.10) \quad ((X_1, u_1) - (X_2, u_2)) \cdot (F_\alpha(X_1, u_1) - F_\alpha(X_2, u_2)) \leq C|(X_1, u_1) - (X_2, u_2)|^2.$$

The uniqueness across the set $\{X = 0\}$ in our setting will follow from the fact that the solution u of (5.2) with constant initial datum $u_0(x) = K > 0$ stays non-negative for all times, and strictly positive at $x = 0$ for all times (see Section 5.1.3), so the characteristics $t \mapsto X(t, t_0, x_0)$, when meeting $\{X = 0\}$, will simply follow the dynamics in $(0, A)$. This argument can be invoked similarly when characteristics meet the set $\{X = A\}$ while u stays positive. However, as we will see, there will be some time at which $u(t, A)$ vanishes. There, uniqueness should also hold across the set $\{X = A\}$ simply by continuity of $F_\alpha(x, 0)$ across $\{X = A\}$, at least for $\alpha \in (0, 1)$. This is however less clear to prove, especially when turning to the case $\alpha = 1$.

Thus, to properly justify the above computations, we construct explicitly the solution u of (5.2) using the characteristics formulae above. In turn, this will guarantee the characteristics formulae given above. Note that, strictly speaking, our arguments construct a solution of (5.2), but by uniqueness of the admissible solution of (5.2), see Proposition 1.4, this solution is *the* solution of (5.2) with initial datum $u_0(x) = K$.

5.1.3. *Regularity of the solution u .* The goal of this section is to prove the following regularity result on the solution u of (5.2) with $u_0(x) = K$.

Lemma 5.1. *Let f and K as in (5.1), $\delta > 0$ and $(0, A) \subset \mathbb{T}$.*

Then the solution u of (5.2) satisfies the following regularity properties: $u \in \mathcal{C}^0([0, \infty) \times \mathbb{T})$, u is piecewise $\mathcal{C}^1([0, \infty) \times \mathbb{T})$, and we have the more precise result:

- *The set $\mathcal{Z} = \{(t, x) \in [0, \infty) \times \mathbb{T}, \text{ s.t. } u(t, x) = 0\}$, if not empty, is a closed set of $[0, \infty) \times (0, A]$ whose boundary is globally Lipschitz and piecewise \mathcal{C}^1 .*
- *For all bounded open subset Ω such that $\bar{\Omega} \subset ([0, \infty) \times \mathbb{T} \setminus \mathcal{Z})$, there exists a finite number of smooth (\mathcal{C}^1) curves \mathcal{C}_i , which may intersect only transversally, such that u is \mathcal{C}^1 in the adherence of each of the connected component of $\Omega \setminus \mathcal{C}_i$.*
- *For all bounded open subset $\Omega \subset ([0, \infty) \times \mathbb{T})$, there exists a finite number of curves \mathcal{C}_i (globally Lipschitz and piecewise \mathcal{C}^1), which may intersect only transversally, such that u is \mathcal{C}^1 in each of the connected component of $\Omega \setminus \mathcal{C}_i$.*
- *For all $t \geq 0$, $x \mapsto u(t, x)$ is non-decreasing on $[A, 1]$.*

The proof of Lemma 5.1 relies on the explicit construction of u using characteristics formula. As we will see in the proof, some curve of strong \mathcal{C}^1 singularity (meaning that at each point of the curve, the function u on each side on the curve cannot be both extended as a \mathcal{C}^1 function up to the curve) may appear when the characteristic entering the zone in which the damping is active corresponds to a small value of u . In this case, characteristics may become vertical and merge after some time. This does not violate the forward uniqueness of the characteristics in the sense of Filippov. Still, we emphasize that when the characteristics merge, we cannot use them backward in time. This is in fact completely similar to the phenomenon which appears when solving the ODE (1.3).

To be more precise, we introduce $\varepsilon \in (0, K]$ as the solution, if it exists, of

$$(5.11) \quad \int_0^\infty f'((\varepsilon^\alpha - \delta\alpha\tau)_+^{1/\alpha}) d\tau = A,$$

and T_* as

$$(5.12) \quad T_* = \inf\{t \in [0, \infty), u(t, 0) \leq \varepsilon\}.$$

The role of ε will appear clearly in the proof below. Loosely speaking, when $u(t, 0) \leq \varepsilon$, the characteristic issued from $(t, 0)$ will never reach the set $\{x = A\}$: in other words, the characteristic issued from $(t, 0)$ is of too low energy to overpass the damping set.

The curves \mathcal{C} on which \mathcal{C}^1 singularities may appear will be constructed with the solution u itself. In fact, these curves will simply be

$$(5.13) \quad \begin{cases} \mathcal{C}_0 : t \mapsto (t, X(t, 0, 0)), \\ \mathcal{C}_1 : t \mapsto (t, X(t, 0, A)), \\ \mathcal{C}_2 : t \mapsto (t, 0), \\ \mathcal{C}_3 : t \mapsto (t, A), \end{cases}$$

to which, if $T_* < \infty$, one should add the boundary of the set \mathcal{Z} (if $\mathcal{Z} \neq \emptyset$), that will be shown to be delimited by

$$(5.14) \quad \mathcal{C} : t_0 \in (T_*, \infty) \mapsto (t(t_0), x(t_0)),$$

$$\text{where } t(t_0) = t_0 + \frac{(u(t_0, 0))^\alpha}{\delta\alpha}, \text{ and } x(t_0) = \int_0^\infty f'((u(t_0, 0)^\alpha - \delta\alpha\tau)_+^{1/\alpha}) d\tau,$$

and a part of \mathcal{C}_3 , namely $\{(t, A), \text{ for } t \geq t_*\}$ for

$$(5.15) \quad t_* = T_* + \frac{\varepsilon^\alpha}{\delta\alpha}.$$

The various discontinuity curves, \mathcal{Z} region and times T_* and t_* are displayed in Figure 5.2.




FIGURE 5.2. Discontinuity curves, \mathcal{Z} region and times T_* and t_* .

Proof.

• **Preliminary computations: Existence and uniqueness of ε in (5.11).** We first emphasize that condition (5.11) is satisfied by at most one parameter $\varepsilon > 0$ as the map

$$(5.16) \quad g : v \mapsto \int_0^\infty f'((v^\alpha - \delta\alpha\tau)_+^{1/\alpha}) d\tau$$

is well-defined and continuous on $[0, K]$, $g(0) = 0$, and g is strictly increasing. Indeed, if $0 \leq v_1 < v_2 \leq K$, using the fact that f' is strictly increasing on $[0, K]$ as f is

assumed to be strictly convex,

$$\begin{aligned} g(v_1) &= \int_0^\infty f'((v_1^\alpha - \delta\alpha\tau)_+^{1/\alpha}) d\tau = \int_0^{v_1^\alpha/(\delta\alpha)} f'((v_1^\alpha - \delta\alpha\tau)_+^{1/\alpha}) d\tau \\ &< \int_0^{v_2^\alpha/(\delta\alpha)} f'((v_2^\alpha - \delta\alpha\tau)_+^{1/\alpha}) d\tau \leq g(v_2). \end{aligned}$$

Thus, if $g(K) \geq A$, there exists a unique $\varepsilon \in [0, K]$ satisfying (5.11). (In case $g(K) < A$, there is no $\varepsilon \in [0, K]$ satisfying (5.11).) Note in particular that, when there exists $\varepsilon \in (0, K]$ satisfying (5.11), for all $v \in (\varepsilon, K]$, $g(v) \geq A$, so that

$$(5.17) \quad \forall v \in (\varepsilon, K], \quad A \leq f'(v) \frac{v^\alpha}{\delta\alpha} \leq f'(K) \frac{K^\alpha}{\delta\alpha}.$$

• **Construction of u .** In our construction below, we distinguish the cases $t < T_*$ and $t \geq T_*$. In particular, according to the definition (5.12) of T_* , for all $t \in [0, T_*)$, $u(t, 0) > \varepsilon$.

We restrict ourselves to the case $T_* > 0$, since the case $T_* = 0$ can be easily adapted from the case $T_* > 0$ and $t \geq T_*$. Hence we assume that $T_* > 0$, and deal separately with the cases $t \leq T_*$ and $t \geq T_*$.

We will not point out below, along the construction of u , that the curves delimiting the \mathcal{C}^1 singularities are exactly the ones in (5.13)–(5.14), but it will appear clearly from the construction of u .

As a matter of fact, we construct a sequence of time-space domains which eventually cover $[0, \infty) \times \mathbb{T}$, and a function u on each of these time-space domains, such that u is a globally \mathcal{C}^0 function there, and is a piecewise \mathcal{C}^1 function, where the curves of \mathcal{C}^1 discontinuities are given by (5.13)–(5.14). Besides, apart from these curves of singularities, the solution u is constructed to satisfy the characteristics equations (5.5) in the case $\alpha \in (0, 1)$, or (5.8) in the case $\alpha = 1$ away from the set \mathcal{X} . The regularity of the curves of \mathcal{C}^1 discontinuities then allows to check easily that the solution u constructed this way solves (5.2) in the sense of Definition 5.6 for $\alpha \in (0, 1)$ or of Definition 5.7 when $\alpha = 1$.

In the proof below, with a slight abuse of terminology, we call “smooth functions” functions which are \mathcal{C}^1 .

Case $t \leq T_*$. As the velocities involved for the solution u should belong to $[0, f'(K)]$, using the light cone of the equation, u should be fully determined in

$$\mathcal{T}_0 = \{(t, x) \text{ with } t \in [0, A/f'(K)] \text{ and } x \in [tf'(K), A]\}$$

by $u(0, \cdot)|_{[0, A]}$. For $(t, x) \in \mathcal{T}_0$, we shall therefore look for x_0 such that $X(t, 0, x_0) = x$, that is

$$x_0 + \int_0^t f'((K^\alpha - \delta\alpha\tau)_+^{1/\alpha}) d\tau = x.$$

For fixed $t \in [0, A/f'(K)]$, it is easily seen that the map

$$[0, A] \ni x_0 \mapsto k_t(x_0) = x_0 + \int_0^t f'((K^\alpha - \delta\alpha\tau)_+^{1/\alpha}) d\tau$$

is smooth, and strictly increasing, with image containing $[tf'(K), A]$. Therefore, for all $(t, x) \in \mathcal{T}_0$, there exists a unique $x_0 \in [0, A]$ such that $X(t, 0, x_0) = x$, so that we set $u(t, x) = (K^\alpha - \delta\alpha t)_+^{1/\alpha}$. As $t \leq A/f'(K)$, the bound (5.17) allows to guarantee that for all $(t, x) \in \mathcal{T}_0$, $K^\alpha - \delta\alpha t$ is strictly positive, so that $u \in \mathcal{C}^1(\mathcal{T}_0)$ and $t \mapsto u(t, A)$

is smooth on $[0, A/f'(K)]$, non-increasing, and strictly positive. We then consider the triangle

$$\mathcal{T}_1 = \{(t, x) \text{ with } t \in [0, 1/f'(K)] \text{ and } x \in [tf'(K), 1]\},$$

and construct u in this set. Of course, as we have already shown that u is smooth in the set \mathcal{T}_0 , we only focus on the set $\mathcal{T}_1 \setminus \mathcal{T}_0$. We determine u in $\mathcal{T}_1 \setminus \mathcal{T}_0$ from $u(\cdot, A)|_{[0, A/f'(K)]}$ and $u(0, \cdot)|_{[A, 1]}$. Introducing $X(t, 0, A) = A + tf'(u(0, A)) = A + tf'(K)$, for $(t, x) \in \mathcal{T}_1 \setminus \mathcal{T}_0$, we set $u(t, x) = K$ if $x \in [X(t, 0, A), 1]$, while if $x \in [\max\{tf'(K), A\}, X(t, 0, A)]$, we find a time $t_0 \in [0, A/f'(K)]$ such that $X(t, t_0, A) = x$, and set $u(t, x) = u(t_0, A)$. Indeed, this can be achieved since for $t \in [0, 1/f'(K)]$ and $x \in [\max\{tf'(K), A\}, X(t, 0, A)]$, finding $t_0 \in [0, \min\{t, A/f'(K)\}]$ such that $X(t, t_0, A) = x$ amounts to solving the equation

$$A + (t - t_0)f'(u(t_0, A)) = x.$$

But the map

$$(5.18) \quad h_t : t_0 \mapsto A + (t - t_0)f'(u(t_0, A))$$

satisfies:

- h_t is strictly decreasing on the interval $[0, \min\{t, A/f'(K)\}]$. This is a consequence of the fact that $t_0 \mapsto u(t_0, A)$ is non-increasing and strictly positive on $[0, A/f'(K)]$ and that f is strictly convex with $f'(0) = 0$. Besides, for all $t_0 \in [0, \min\{t, A/f'(K)\}]$,

$$h'_t(t_0) \leq -f'(u(A/f'(K), A)) < 0.$$

- $h_t(0) = A + tf'(K) = X(t, 0, A)$.
- If $t \leq A/f'(K)$, $h_t(t) = A$, and if $t \geq A/f'(K)$,

$$\begin{aligned} h_t\left(\frac{A}{f'(K)}\right) &= A + \left(t - \frac{A}{f'(K)}\right) f'\left(u\left(\frac{A}{f'(K)}, A\right)\right) \\ &\leq A + \left(t - \frac{A}{f'(K)}\right) f'(K) \leq tf'(K). \end{aligned}$$

- h_t is smooth.

It then follows that h_t is a diffeomorphism from the interval $[0, \min\{t, A/f'(K)\}]$ to its image, which contains $[\max\{tf'(K), A\}, X(t, 0, A)]$, so that we can write, for $t \in [0, 1/f'(K)]$ and x in the interval $[\max\{tf'(K), A\}, X(t, 0, A)]$, $u(t, x) = u(h_t^{-1}(x), A)$.

These formula easily show that the function u is smooth for $(t, x) \in \mathcal{T}_1$ with $x \in [\max\{tf'(K), A\}, X(t, 0, A)]$, and it is clear that u is smooth for $(t, x) \in \mathcal{T}_1$ with $x \in [X(t, 0, A), 1]$. However, though it is clear that u is continuous along the curve $t \mapsto (t, X(t, 0, A))$, u may not be locally \mathcal{C}^1 in a neighborhood of this curve, even if it can be extended as \mathcal{C}^1 functions up to this curve from each side. It follows that u is piecewise $\mathcal{C}^1(\mathcal{T}_1)$, and that $t \mapsto u(t, 1)$ is a piecewise \mathcal{C}^1 non-increasing function on $[0, 1/f'(K)]$, which remains strictly positive on the interval $[0, 1/f'(K)]$.

It is also easy to check that for all $t \in [0, A/f'(K)]$, $x \mapsto u(t, x)$ is non-decreasing on $[A, 1]$.

In fact, this is the starting point of an iterative argument showing that for all $n \in \mathbb{N}$ with $T_n = n/f'(K) < T_*$, one can construct a solution u of (5.2) in

$$\mathcal{T}_{2n+1} = \{(t, x) \in [0, (n+1)/f'(K)] \times [0, 1] \text{ with } x \in [\max\{tf'(K) - n, 0\}, 1]\},$$

such that:

- u is globally \mathcal{C}^0 and piecewise \mathcal{C}^1 on \mathcal{T}_{2n+1}

- $t \mapsto u(t, A)$ is strictly positive and non-increasing on $[0, (n + A)/f'(K)]$.
- $t \mapsto u(t, 1)$ is strictly positive non-increasing on $[0, (n + 1)/f'(K)]$.
- For all $t \in [0, T_n + A/f'(K)]$, $x \mapsto u(t, x)$ is non-decreasing on $[A, 1]$.

Indeed, these properties are already proved for $n = 0$. Let us show that if they hold for $n \in \mathbb{N}$, they also hold for $n + 1$ provided $T_{n+1} < T_*$.

Using the above properties for n , we first consider the equation in the trapezoid

$\mathcal{T}_{2n+2} = \{(t, x) \in [0, (n+1+A)/f'(K)] \times [0, A] \text{ with } x \in [\max\{tf'(K) - (n+1), 0\}, A]\}$, and use the boundary condition $u(\cdot, 0)|_{[0, (n+1)/f'(K)]}$ and $u(0, \cdot)|_{[0, A]}$ to construct u in \mathcal{T}_{2n+2} . In order to do this, we set, for $t \geq 0$,

$$x_0(t) = \min \left\{ \int_0^t f'((K^\alpha - \delta\alpha\tau)_+^{1/\alpha}) d\tau, A \right\}.$$

Now, let us fix $t \in [0, (n + 1 + A)/f'(K)]$ and $x \in [0, A]$. If $x > x_0(t)$, we define $u(t, x) = (K^\alpha - \delta\alpha t)_+^{1/\alpha}$. If $x < x_0(t)$, we look for $t_0 \in [0, t]$ such that $X(t, t_0, 0) = x$, i.e. such that

$$\int_0^{t-t_0} f'((u(t_0, 0)^\alpha - \delta\alpha\tau)_+^{1/\alpha}) d\tau = x.$$

We thus define, for $t_0 \in [0, t]$,

$$g_t(t_0) = \int_0^{t-t_0} f'((u(t_0, 0)^\alpha - \delta\alpha\tau)_+^{1/\alpha}) d\tau.$$

It is not difficult to check that, for $t \leq (n + 1 + A)/f'(K)$, g_t enjoys the following properties:

- g_t is strictly decreasing on $[0, \min\{t, T_n\}]$. This is a consequence of the fact that $t \mapsto u(t, 1)$ ($= u(t, 0)$) is non-increasing and strictly larger than ε on $[0, T_n]$. Besides, g_t is piecewise \mathcal{C}^1 on $[0, \min\{t, T_n\}]$ and for all t_0 such that $g_t(t_0) \leq A$ and for which g_t is differentiable,

$$\begin{aligned} g_t'(t_0) &\leq -f'((u(t_0, 0)^\alpha - \delta\alpha(t - t_0))_+^{1/\alpha}) \\ &\leq -f'((u(\min\{t, T_n\}, 0)^\alpha - \delta\alpha(\min\{t, T_n\} - t_0))_+^{1/\alpha}). \end{aligned}$$

Note that, if $f'((u(\min\{t, T_n\}, 0)^\alpha - \delta\alpha(\min\{t, T_n\} - t_0))_+^{1/\alpha}) = 0$ and $g_t(t_0) \leq A$, then $g(u(\min\{t, T_n\}, 0)) \leq A$, which is not compatible with $u(\min\{t, T_n\}, 0) > \varepsilon$. Thus there exists $\gamma > 0$ such that for all t_0 such that $g_t(t_0) \leq A$ and for which g_t is differentiable, $g_t'(t_0) \leq -\gamma$.

- $g_t(t) = 0$.
- $g_t(0) = x_0(t)$ if $\int_0^t f'((K^\alpha - \delta\alpha\tau)_+^{1/\alpha}) d\tau \leq A$, and is larger than A otherwise.

One then easily shows that for $t \leq (n + 1 + A)/f'(K)$, the map g_t is a piecewise \mathcal{C}^1 function from $[0, \min\{t, T_n\}]$ to its image, which contains $[0, x_0(t)]$. Besides, there exists a unique $t_0(t, A)$ such that g_t is a piecewise \mathcal{C}^1 diffeomorphism from $[t_0(t, A), \min\{t, T_n\}]$ to $[0, x_0(t)]$. We can then define u for $t \leq (n + 1 + A)/f'(K)$ and $[0, x_0(t)]$ by

$$u(t, x) = (u(g_t^{-1}(x), 0)^\alpha - \delta\alpha(t - g_t^{-1}(x)))_+^{1/\alpha},$$

and, for $x \in x_0(t)$, by $u(t, x) = (K^\alpha - \delta\alpha t)_+^{1/\alpha}$.

As g_t depends smoothly on the time parameter t , this defines u as a globally \mathcal{C}^0 and piecewise \mathcal{C}^1 function in \mathcal{T}_{2n+2} .

We then check that $t \mapsto u(t, A)$ is strictly positive, piecewise \mathcal{C}^1 and non-increasing on $[0, (n + 1 + A)/f'(K)]$. It is obviously \mathcal{C}^1 and decreasing in $\{t, x_0(t) < A\}$, as

$u(t, A) = (K^\alpha - \delta\alpha t)_+^{1/\alpha}$, the positivity coming from the fact that $K > \varepsilon$ and $t \leq T_{n+1} \leq T_*$. For $\{t, x_0(t) = A\}$, which is an interval of the form $[t_A, (n+1+A)/f'(K)]$, $u(t, A)$ is given by the formula

$$u(t, A) = (u(g_t^{-1}(A), 0)^\alpha - \delta\alpha(t - g_t^{-1}(A)))_+^{1/\alpha}.$$

Now, one can check that for $t^a, t^b \in [t_A, (n+1+A)/f'(K)]$ such that $t^a < t^b$, defining t_0^a and t_0^b by the formula

$$g_{t^a}(t_0^a) = A = g_{t^b}(t_0^b), \quad \text{i.e.} \quad t_0^a = g_{t^a}^{-1}(A), \quad t_0^b = g_{t^b}^{-1}(A),$$

the decay of $t_0 \mapsto u(t_0, 0)$ on $[0, T_n]$ implies that

$$t_0^a \leq t_0^b, \quad \text{and} \quad t^a - t_0^a \geq t^b - t_0^b.$$

Consequently, since $t_0 \mapsto u(t_0, 0)$ is non-increasing on $[0, T_{n+1}]$, we immediately have that $t \mapsto u(t, A)$ is non-increasing on $[0, (n+1+A)/f'(K)]$. It is also obviously piecewise \mathcal{C}^1 on $[0, (n+1+A)/f'(K)]$. The fact that $u(t, A)$ is strictly positive comes from the fact that $t \mapsto u(t, 0)$ stays strictly larger than ε for $t \leq T_n$.

We then construct the solution u in the set $\mathcal{T}_{2n+3} \setminus \mathcal{T}_{2n+2}$, using the information given by $u(\cdot, A)|_{[0, (n+1+A)/f'(K)]}$ and $u(0, \cdot)|_{[A, 1]}$. Again, as when working in \mathcal{T}_1 , we construct the solution u using characteristics and setting, for $(t, x) \in \mathcal{T}_{2n+3}$ with $x \in [A, \min\{A + tf'(K), 1\}]$,

$$u(t, x) = u(h_t^{-1}(x), A).$$

The other case, corresponding to $x \in [\min\{A + tf'(K), 1\}, 1]$, lies in fact in \mathcal{T}_1 , so regularity issues have been dealt with before. We only need to check that for all $t \in [0, T_{n+1} + A/f'(K)]$, $x \mapsto u(t, x)$ is non-decreasing on $[A, 1]$: this is obvious if $x \in [\min\{A + tf'(K), 1\}, 1]$ as $u(t, x) = K$ there; when $x \in [A, \min\{A + tf'(K), 1\}]$, the above formula and the facts that $t \mapsto u(t, A)$ is non-increasing and that h_t^{-1} is strictly decreasing imply that $x \mapsto u(t, x)$ is non-increasing on $[A, \min\{A + tf'(K), 1\}]$. Therefore, all the items in the above property also hold for $n+1$.

We can thus iterate these arguments while $T_n = n/f'(K) < T_*$. If $T_* = \infty$, this concludes Lemma 5.1. If $T_* < \infty$, we perform a similar iteration in the trapeze

$$\mathcal{T} = \{(t, x) \in [0, T_* + 1/f'(K)] \times [0, 1] \text{ with } x \in [\max\{(t - T_*)f'(K), 0\}, 1]\},$$

constructing a function u in \mathcal{T} such that:

- u is a solution of (5.2), is piecewise \mathcal{C}^1 on \mathcal{T} and globally \mathcal{C}^0 on \mathcal{T} .
- $t \mapsto u(t, A)$ is strictly positive and non-increasing on $[0, T_* + A/f'(K)]$.
- $t \mapsto u(t, 0)$ is strictly positive non-increasing on $[0, T_* + 1/f'(K)]$.
- $u(T_*, 0) = \varepsilon$.
- For all $t \in [0, T_* + A/f'(K)]$, $x \mapsto u(t, x)$ is non-decreasing on $[A, 1]$.

Case $t \geq T_*$. The difficulty when $t \geq T_*$ is to show that the solution u remains continuous in time-space. In order to do this, we introduce $(x_*(t), u_*(t))$ given by

$$(5.19) \quad \begin{cases} \frac{d}{dt} \begin{pmatrix} x_* \\ u_* \end{pmatrix} = F_\alpha \begin{pmatrix} x_* \\ u_* \end{pmatrix}, & t \geq T_*, \text{ if } \alpha \in (0, 1), \\ \text{or} \\ \frac{d}{dt} \begin{pmatrix} x_* \\ u_* \end{pmatrix} \in F_1 \begin{pmatrix} x_* \\ u_* \end{pmatrix}, & t \geq T_*, \text{ if } \alpha = 1, \end{cases} \quad \begin{pmatrix} x_*(T_*) \\ u_*(T_*) \end{pmatrix} = \begin{pmatrix} 0 \\ \varepsilon \end{pmatrix},$$

where F_α is defined in (5.6)–(5.7). Here, x_* corresponds to the characteristics $X(t, T_*, 0)$ $t \geq T_*$ and $x_*(t) \leq A$. Note that, due to the choice of ε in (5.11), there exists a time

t_* ($= T_* + \varepsilon^\alpha/(\delta\alpha)$), such that the solution (x_*, u_*) satisfies

$$(5.20) \quad \forall t \in [T_*, t_*), \quad x_*(t) \in [0, A], \quad \text{and} \quad \forall t \geq t_*, \quad (x_*(t), u_*(t)) = (A, 0),$$

thus guaranteeing the uniqueness of the solution of (5.19).

We then introduce the following sets:

$$\begin{aligned} \mathcal{R}_0 &= \{(t, x) \in [0, \infty) \times [0, A], \text{ such that, if } t \geq T_*, x \in [x_*(t), A]\}, \\ \mathcal{R}_1 &= \{(t, x) \in [0, \infty) \times [A, 1]\}, \\ \mathcal{R}_2 &= \{(t, x), \text{ with } t \geq T_*, x \in [0, x_*(t)]\}, \end{aligned}$$

and, similarly as before, for $n \geq 0$,

$$\begin{aligned} \mathcal{F}_{2n}^* &= \{(t, x) \in [0, T_* + (n + A)/f'(K)] \times [0, 1] \text{ with } x \in [((t - T_*)f'(K) - n)_+, A]\}, \\ \mathcal{F}_{2n+1}^* &= \{(t, x) \in [0, T_* + (n + 1)/f'(K)] \times [0, 1] \text{ with } x \in [((t - T_*)f'(K) - n)_+, 1]\}. \end{aligned}$$

As before, we construct iteratively a solution u of (5.2) in $\mathcal{F}_{2n+1}^* \setminus \mathcal{R}_2$ for all $n \in \mathbb{N}$ such that:

- u is piecewise \mathcal{C}^1 and globally \mathcal{C}^0 in $\mathcal{F}_{2n+1}^* \setminus \mathcal{R}_2$.
- $t \mapsto u(t, 1)$ is strictly positive non-increasing on $[0, T_* + (n + 1)/f'(K)]$ and piecewise \mathcal{C}^1 .
- for all $t \geq T_*$ $u(t, x_*(t)) = u_*(t)$, where (x_*, u_*) is the solution of (5.19).
- For all $t \in [0, T_* + (n + A)/f'(K)]$, $x \mapsto u(t, x)$ is non-decreasing on $[A, 1]$.

Of course, the previous paragraph shows that this is true for $n = 0$. Let us then assume that these properties are true for some $n \in \mathbb{N}$ and show that they are then true for $n + 1$.

Similarly as before, we work first on $\mathcal{F}_{2n+2}^* \cap \mathcal{R}_0$. The construction of the function u in the set $\mathcal{F}_{2n+2}^* \cap \mathcal{R}_0$ can then be handled as for \mathcal{F}_{2n+2} , and following the same arguments, we easily get that the function u there is piecewise \mathcal{C}^1 in $\mathcal{F}_{2n+2}^* \cap \mathcal{R}_0$, and that $t \mapsto u(t, A)$ is a non-increasing function on $[0, T_* + (n + 1 + A)/f'(K)]$, strictly positive while $t \leq t_*$, and vanishing for $t \geq t_*$.

The construction of the solution u in the set $\mathcal{F}_{2n+3}^* \cap \mathcal{R}_1$ can then be done similarly as the one corresponding to \mathcal{F}_{2n+3} , and following the same lines, we get that the solution u is piecewise \mathcal{C}^1 in $\mathcal{F}_{2n+3}^* \cap \mathcal{R}_1$. We nevertheless make the proof slightly more precise: for $(t, x) \in \mathcal{F}_{2n+3}^* \cap \mathcal{R}_1 \setminus \mathcal{F}_1$, $u(t, x)$ is constructed by

$$u(t, x) = u(h_t^{-1}(x), A),$$

where h_t is given by (5.18). In particular, to establish the strict positivity of $u(t, 1)$ for $t \leq T_* + (n + 2)/f'(K)$, we just remark that $t_0 = h_t^{-1}(x)$ is equivalent to $A + (t - t_0)f'(u(t_0, A)) = 1$, so that $u(t, x) = u(t_0, A)$ cannot be zero.

The continuity of the function u constructed above in $\mathcal{F}_{2n+3}^* \setminus \mathcal{R}_2$ follows easily from the continuity of u across the interfaces of $\mathcal{F}_{2n+2}^* \cap \mathcal{R}_0$, and $\mathcal{F}_{2n+3}^* \cap \mathcal{R}_1$.

The fact that for all $t \in [0, T_* + (n + 1 + A)/f'(K)]$, $x \mapsto u(t, x)$ is non-decreasing in $[A, 1]$ follows as before.

Our goal now is to construct u in \mathcal{R}_2 as a piecewise \mathcal{C}^1 function such that u is globally continuous on $[0, \infty) \times \mathbb{T}$ and solves (5.2) in $[0, \infty) \times \mathbb{T}$.

As $t \mapsto u(t, 0)$ is strictly positive and non-increasing on $[0, \infty)$, we can show, similarly as before that, for $(t, x) \in \mathcal{R}_2$, the map

$$g_t(t_0) = \int_0^{t-t_0} f' \left((u(t_0, 0)^\alpha - \delta\alpha\tau)_+^{1/\alpha} \right) d\tau$$

has the following properties:

- g_t is decreasing on $[T_*, t]$, takes values in $[0, x_*(t)]$, and is surjective on $[0, x_*(t)]$,
- For all $(t_0^a, t_0^b) \in [T_*, t]^2$, $g_t(t_0^a) = g_t(t_0^b)$ implies $t_0^a = t_0^b$ or $(u(t_0^a, 0)^\alpha - \delta\alpha(t - t_0^a))_+ = (u(t_0^b, 0)^\alpha - \delta\alpha(t - t_0^b))_+$.

It thus allows to set, for $(t, x) \in \mathcal{R}_2$,

$$(5.21) \quad u(t, x) = (u(g_t^{-1}(x), 0)^\alpha - \delta\alpha(t - g_t^{-1}(x)))_+^{1/\alpha},$$

where $g_t^{-1}(x)$ denotes any $t_0 \in [T_*, t]$ such that $g_t(t_0) = x$. Note that, according to the second item above, the definition above does not depend on the choice of t_0 such that $g_t(t_0) = x$. Besides, u defined this way is continuous on \mathcal{R}_2 as one can easily check, and if $t \geq T_*$, $u(t, x_*(t)) = u_*(t)$, where (x_*, u_*) is the solution of (5.19).

We then remark that, for $t \geq T_*$, and $T_* \leq t_0^a < t_0^b \leq t$,

$$\begin{aligned} g_t(t_0^a) &= \int_0^{t-t_0^a} f' \left((u(t_0^a, 0)^\alpha - \delta\alpha\tau)_+^{1/\alpha} \right) d\tau \\ &\geq \int_0^{t-t_0^b} f' \left((u(t_0^a, 0)^\alpha - \delta\alpha\tau)_+^{1/\alpha} \right) d\tau \\ &\geq \int_0^{t-t_0^b} f' \left((u(t_0^b, 0)^\alpha - \delta\alpha\tau)_+^{1/\alpha} \right) d\tau = g_t(t_0^b). \end{aligned}$$

In particular, analyzing the case of equality in the above estimates, we easily get that for $(t, x) \in \mathcal{R}_2$ such that $u(t, x) \neq 0$, $t_0(t, x) = g_t^{-1}(x)$ is uniquely defined and

$$(5.22) \quad t - t_0(t, x) \leq \frac{u(t_0(t, x), 0)^\alpha}{\delta\alpha} \leq \frac{\varepsilon^\alpha}{\delta\alpha}.$$

Besides, g_t is piecewise \mathcal{C}^1 locally around $t_0(t, x)$ and

$$\begin{aligned} g'_t(t_0) &= -f' \left((u(t_0, 0)^\alpha - \delta\alpha(t - t_0)^{1/\alpha}) \right) \\ &+ u(t_0, 0)^{\alpha-1} \partial_t u(t_0, 0) \int_0^{t-t_0} f' \left((u(t_0, 0)^\alpha - \delta\alpha\tau)^{1/\alpha} \right) (u(t_0, 0)^\alpha - \delta\alpha\tau)^{1/\alpha-1} d\tau. \end{aligned}$$

As $u(t, x) \neq 0$ and $t_0 \mapsto u(t_0, 0)$ decays, this implies that

$$g'_t(t_0(t, x)) \leq -f' \left((u(t_0, 0)^\alpha - \delta\alpha(t - t_0(t, x)))^{1/\alpha} \right) = -f'(u(t, x)) < 0,$$

and the bound is uniform in a neighborhood of t and x . As $t_0(t, x)$ is defined by $g_t(t_0(t, x)) = x$, we see that this implies that t_0 is piecewise \mathcal{C}^1 for $(t, x) \in \mathcal{R}_2$ such that $u(t, x) \neq 0$, so that the definition (5.21) shows that u is piecewise \mathcal{C}^1 for $(t, x) \in \mathcal{R}_2$ such that $u(t, x) \neq 0$.

It is thus also interesting to determine the area $\mathcal{Z} = \{(t, x), \text{ s.t. } u(t, x) = 0\} \cap \mathcal{R}_2$.

For $(t, x) \in \mathcal{R}_2$, it is clear, from the relation $g_t(t_0(t, x)) = x$ and from the fact that $t_0 \mapsto u(t_0, 0)$ decays, that for $t^a < t^b$ such that (t^a, x) and (t^b, x) are in \mathcal{R}_2 ,

$$t_0(t^a, x) \leq t_0(t^b, x), \quad \text{and} \quad t^a - t_0(t^a, x) \leq t^b - t_0(t^b, x),$$

so that we easily derive from the decay of $t_0 \mapsto u(t_0, 0)$ that

$$u(t^a, x) \leq u(t^b, x).$$

It follows that, for all $x \in [0, A]$, the map $t \mapsto u(t, x)$ decays while $(t, x) \in \mathcal{R}_2$.

We can in particular define, for $x \in (0, A]$, the time $t_*(x) = \inf\{t, \text{ s.t. } u(t, x) = 0\}$ (which may be infinite). If $t_*(x)$ is finite, for $t < t_*(x)$, $u(t, x) \neq 0$, so $t_0(t, x)$ is well-defined and unique, and is an increasing function of time which is bounded by

$t_*(x)$. Thus, the limit of $t_0(t, x)$ as $t \rightarrow t_*^-$ exists, and we call it $t_{0,*}(x)$. We then easily get that

$$u(t_{0,*}(x), 0)^\alpha - \delta\alpha(t_*(x) - t_{0,*}(x)) = 0,$$

so that

$$x = g_{t_*(x)}(t_{0,*}(x)) = g(u(t_{0,*}(x), 0)),$$

where g is defined in (5.16). Let us now note that there exists only one t_0 such that

$$(5.23) \quad u(t_0, 0)^\alpha - \delta\alpha(t_*(x) - t_0) = 0 \quad \text{and} \quad x = g(u(t_0, 0)).$$

Indeed, the second equation determines uniquely $u(t_0, 0)$, and the first equation then determines uniquely t_0 . Thus, if t_0 satisfies (5.23), $t_0 = t_{0,*}(x)$ and $t_*(x) = t_0 + u(t_0, 0)^\alpha/(\delta\alpha)$.

This suggests to study the parametric equation

$$(5.24) \quad \mathcal{C} : t_0 \in (T_*, \infty) \mapsto (t(t_0), x(t_0)),$$

$$\text{where } t(t_0) = t_0 + \frac{(u(t_0, 0))^\alpha}{\delta\alpha}, \text{ and } x(t_0) = g(u(t_0, 0)).$$

(This definition of course coincides with the definition of \mathcal{C} in (5.14), recall the definition of g in (5.16).) It is clear that by construction $u(t(t_0), x(t_0)) = 0$ for all $t_0 \geq T_*$. Besides, for $(t, x) \in \mathcal{R}_2$, if there exists $T_0 \geq T_*$ such that $x = g(u(t_0, 0))$, then

- if $t \geq \inf\{t(t_0), \text{ for } t_0 \text{ s.t. } x = g(u(t_0, 0))\}$, then $u(t, x) = 0$;
- if $t \geq \inf\{t(t_0), \text{ for } t_0 \text{ s.t. } x = g(u(t_0, 0))\}$, then $u(t, x) > 0$.

These statements follow immediately from the decay of $t \mapsto u(t, x)$ and the non-negativity of u . It follows that

$$(5.25) \quad \partial\mathcal{Z} = \mathcal{C} \cup \{(t, A), \text{ s.t. } t \geq t_*\}.$$

We remark that, as $t_0 \mapsto u(t_0, 0)$ is piecewise \mathcal{C}^1 and strictly positive, except at singularities (which are in finite number in any bounded interval), we have

$$\frac{d}{dt_0} \left(t_0 + \frac{(u(t_0, 0))^\alpha}{\delta\alpha}, g(u(t_0, 0)) \right) = \left(1 + \frac{u(t_0, 0)^{\alpha-1} \partial_t u(t_0, 0)}{\delta}, g'(u(t_0, 0)) \partial_t u(t_0, 0) \right),$$

so that

$$\left| \frac{d}{dt_0} \left(t_0 + \frac{(u(t_0, 0))^\alpha}{\delta\alpha}, g(u(t_0, 0)) \right) \right| \neq 0.$$

This proves that the tangent, hence the normal, of the curve \mathcal{C} is well-defined except at a locally finite number of points. This indicates that \mathcal{C} is a piecewise \mathcal{C}^1 parametric curve with finite limits at singularity points. It is thus a globally Lipschitz and piecewise \mathcal{C}^1 parametric curve.

We have thus proved the regularity properties stated in Lemma 5.1. In particular, in all connected component of $([0, \infty) \times \mathbb{T}) \setminus (\cup_{i=1}^4 \mathcal{C}_i \cup \mathcal{C})$, the function u is \mathcal{C}^1 and satisfies, by construction

$$\partial_t u + \partial_x(f(u)) + h(t, x) = 0,$$

where $h(t, x) = \delta \mathbf{1}_{(0, A)} u(t, x)/|u(t, x)|^\alpha$ if $u(t, x) > 0$ and $h(t, x) = 0$ if $u(t, x) = 0$. Since u is also continuous in the whole set $[0, \infty) \times \mathbb{T}$, a straightforward application of the integration by parts formula shows that the function u we constructed above is an admissible solution of (5.2) in $[0, \infty) \times \mathbb{T}$. \square

5.1.4. Dynamics of the solution u in the case of a strictly convex flux.

Lemma 5.2. *Let f and K as in (5.1), $\delta > 0$ and $\omega = (0, A) \subset \mathbb{T}$. Then the solution u of (5.2) satisfies:*

- (i) T_* defined in (5.12) is finite.
- (ii) For all $t \geq t_*$ (defined in (5.15)), there exists an open subinterval $\omega(t) \subset \omega$ such that $u(t)|_{\omega(t)} = 0$, and

$$|\omega \setminus \omega(t)| \leq \frac{C}{t^{1+\alpha}}, \quad \|u(t)\|_{L^\infty(\mathbb{T})} \leq \frac{C}{t}.$$

Proof of item (i) of Lemma 5.2. The proof of the first item of Lemma 5.2 follows from the analysis of the solution u of (5.2) along the characteristics $t \mapsto X(t, 0, 0)$, in particular when it crosses the set $\{x = 0\}$.

We thus introduce four sequences $(u_n)_{n \in \mathbb{N}}$, $(t_n)_{n \in \mathbb{N}}$, $(v_n)_{n \in \mathbb{N}}$, $(\tau_n)_{n \in \mathbb{N}}$, initialized by $u_0 = K$ and $t_0 = 0$, and defined iteratively for n such that $u_n > \varepsilon$ and $v_n > 0$ as follows:

- $\tau_n > t_n$ is the unique solution of

$$\int_0^{\tau_n - t_n} f'((u_n^\alpha - \delta\alpha\tau)_+^{1/\alpha}) d\tau = A.$$

- $v_n = (u_n^\alpha - \delta\alpha(\tau_n - t_n))_+^{1/\alpha}$.
- $t_{n+1} = \tau_n + (1 - A)/f'(v_n)$.
- $u_{n+1} = v_n$.

These choices are made so that for all $n \in \mathbb{N}$ such that $u_n > \varepsilon$, $u(t_n, 0) = u_n$, and $u(\tau_n, A) = v_n$.

We also set n_0 the first integer (if any) for which $u_n < \varepsilon$ or $v_n = 0$. This index n_0 , if any, is in fact such that $u_{n_0} \leq \varepsilon$ and $v_{n_0} = 0$. Indeed, if $u_n > \varepsilon$, one easily checks from the definition of ε that $v_n > 0$.

Note that we easily deduce from the formula of v_n that

$$\forall n \in \{0, \dots, n_0 - 1\}, \quad \delta\alpha(\tau_n - t_n) \leq u_n^\alpha.$$

In order to study these sequences, it will be convenient to have a good estimate on $\tau_n - t_n$ in terms of u_n only. We thus define

$$\beta_- = \inf_{[0, K]} \frac{f'(s)}{s}, \quad \beta_+ = \sup_{[0, K]} \frac{f'(s)}{s},$$

which are finite as $f'(0) = 0$ and which are both strictly positive as f is strictly convex.

We then recall that $1/\alpha \geq 1$. Thanks to the convexity of the function $s \mapsto s^{1/\alpha}$ at the point u^α , its graph on $[0, u^\alpha]$ is below the chord initiated from the origin, i.e. $s \mapsto su^{1-\alpha}$, and above its tangent $s \mapsto (1 - 1/\alpha)u + su^{1-\alpha}/\alpha$, which yields the following estimates: for all $u > 0$, $\tau \geq 0$,

$$(u - \delta\tau u^{1-\alpha})_+ \leq (u^\alpha - \delta\alpha\tau)_+^{1/\alpha} \leq (u - \delta\alpha\tau u^{1-\alpha})_+.$$

Combining the above two estimates, we infer

$$\beta_- \int_0^{\tau_n - t_n} (u_n - \delta\tau u_n^{1-\alpha})_+ d\tau \leq A \leq \beta_+ \int_0^{\tau_n - t_n} (u_n - \delta\alpha\tau u_n^{1-\alpha})_+ d\tau,$$

which implies in particular that

$$A \geq \begin{cases} \frac{\beta_-}{2\delta} u_n^{1+\alpha} & \text{if } u_n^\alpha \leq \delta(\tau_n - t_n), \\ \beta_- u_n(\tau_n - t_n) - \frac{\delta\beta_-}{2} u_n^{1-\alpha}(\tau_n - t_n)^2 & \text{if } u_n^\alpha \geq \delta(\tau_n - t_n), \end{cases}$$

and

$$A \leq \begin{cases} \frac{\beta_+}{2\alpha\delta} u_n^{1+\alpha} & \text{if } u_n^\alpha \leq \alpha\delta(\tau_n - t_n), \\ \beta_+ u_n(\tau_n - t_n) - \frac{\alpha\delta\beta_+}{2} u_n^{1-\alpha}(\tau_n - t_n)^2 & \text{if } u_n^\alpha \geq \alpha\delta(\tau_n - t_n). \end{cases}$$

We claim that we can deduce from this a lower bound on $\tau_n - t_n$ of the form

$$\forall n \in \{0, \dots, n_0 - 1\}, \alpha\delta(\tau_n - t_n) \geq u_n^\alpha \min \left\{ \alpha, 1 - \sqrt{\left(1 - \frac{2A\alpha\delta}{\beta_+ K^{1+\alpha}}\right)_+} \right\}.$$

Indeed, this is obvious when $\tau_n - t_n \geq u_n^\alpha/\delta$. Otherwise, when $\tau_n - t_n \leq u_n^\alpha/\delta$, we have $u_n^\alpha \geq \alpha\delta(\tau_n - t_n)$ and thus one should have

$$\frac{A\alpha\delta}{2\beta_+ u_n^{1+\alpha}} \leq \frac{\alpha\delta}{2} \frac{\tau_n - t_n}{u_n^\alpha} - \left(\frac{\alpha\delta}{2} \frac{\tau_n - t_n}{u_n^\alpha} \right)^2.$$

But $u_n \leq K$ as the sequence u_n is non-increasing, so that

$$\frac{A\alpha\delta}{2\beta_+ K^{1+\alpha}} \leq \left(\frac{\alpha\delta}{2} \frac{\tau_n - t_n}{u_n^\alpha} \right) - \left(\frac{\alpha\delta}{2} \frac{\tau_n - t_n}{u_n^\alpha} \right)^2.$$

Consequently, if $A\alpha\delta/(2\beta_+ K^{1+\alpha}) > 1/4$, this cannot happen. Besides, if

$$A\alpha\delta/(2\beta_+ K^{1+\alpha}) \leq 1/4,$$

we obtain immediately that

$$\frac{\tau_n - t_n}{u_n^\alpha} \geq \frac{1}{\alpha\delta} \left(1 - \sqrt{1 - \frac{2A\alpha\delta}{\beta_+ K^{1+\alpha}}} \right).$$

It follows from this estimate and the definition of v_n that there exists $c_0 < 1$ such that for all $n \in \{0, \dots, n_0 - 1\}$,

$$u_{n+1} = v_n \leq c_0 u_n.$$

Therefore,

$$\forall n \in \{0, \dots, n_0 - 1\}, \quad u_n \leq c_0^n K, \quad \text{and } v_n \leq c_0^{n-1} K,$$

and there indeed exists $n_0 \in \mathbb{N}$ such that $u_{n_0} \leq \varepsilon$. One can also estimate t_{n_0} and show that it is finite, and thus T_* is finite. \square

Proof of item (ii) of Lemma 5.2. In order to prove item (ii) of Lemma 5.2, we look at the solution u in $[t_*, \infty) \times [A, 1]$, where t_* is given by (5.15).

It follows from Lemma 5.1 that u is piecewise \mathcal{C}^1 and continuous in $[t_*, \infty) \times (A, 1)$, non-negative there, and $u(t, A) = 0$ for all $t \geq t_*$. Consequently, for all $t \geq t_*$,

$$u(t, 1) = u(t_*, x_0),$$

where x_0 is the unique solution of

$$x_0 + (t - t_*)f'(u(t_*, x_0)) = 1.$$

In particular, one should have

$$\beta_- u(t_*, x_0) \leq f'(u(t_*, x_0)) \leq \frac{1}{t - t_*}.$$

Therefore,

$$(5.26) \quad \forall t \geq t_*, \quad u(t, 1) \leq \frac{1}{\beta_-(t - t_*)}.$$

From Lemma 5.1 and the fact that for all $t \geq 0$, $x \mapsto u(t, x)$ is non-decreasing on $[A, 1]$ and non-negative, we get

$$\forall t \geq t_*, \quad \|u(t)\|_{L^\infty(A, 1)} \leq \frac{1}{\beta_-(t - t_*)}.$$

Now, the set \mathcal{Z} on which $u = 0$ is delimited by the curve $\{(t, A), \text{ for } t \geq t_*\}$ and the curve \mathcal{C} given by (5.14), that we now estimate: for $t_0 \geq T_*$,

$$t_0 + \frac{(u(t_0, 0))^\alpha}{\delta\alpha} \leq t_0 + \frac{1}{\beta_-^\alpha(t - t_*)^\alpha},$$

and

$$\begin{aligned} \int_0^\infty f'((u(t_0, 0)^\alpha - \delta\alpha\tau)_+^{1/\alpha}) d\tau &= \int_0^{u(t_0, 0)^\alpha/(\delta\alpha)} f'((u(t_0, 0)^\alpha - \delta\alpha\tau)_+^{1/\alpha}) d\tau \\ &\leq \beta_+ \int_0^{u(t_0, 0)^\alpha/(\delta\alpha)} (u(t_0, 0)^\alpha - \delta\alpha\tau u(t_0, 0)^{1-\alpha}) d\tau \\ &\leq \frac{\beta_+}{2\delta\alpha} (u(t_0, 0))^{1+\alpha} \leq \frac{\beta_+}{2\delta\alpha\beta_-^{1+\alpha}} \frac{1}{(t - t_*)^{1+\alpha}}. \end{aligned}$$

We then easily deduce that for $t \geq t_*$, there exists an open subinterval $\omega(t) \subset \omega$ such that $u(t)|_{\omega(t)} = 0$, and, for some constant C , for all time $t \geq t_*$,

$$|\omega \setminus \omega(t)| \leq \frac{C}{t^{1+\alpha}}.$$

Besides, for $t \geq t_*$, we easily get from (5.21) and (5.22) that

$$\forall x \in [0, A], \quad 0 \leq u(t, x) \leq \max_{t_0 \in [t - \varepsilon^\alpha/(\delta\alpha), t]} u(t_0, 0) \leq \frac{1}{\beta_-(t - t_* - \varepsilon^\alpha/(\delta\alpha))}.$$

The item (ii) of Lemma 5.2 easily follows. \square

5.2. Concave flux and positive constant initial datum. Let $K > 0$ and f be a strictly concave flux, and consider the solution u of

$$(5.27) \quad \partial_t u + \partial_x f(u) + a(x) \frac{u}{|u|^\alpha} = 0, \quad (t, x) \in \mathbb{R}_+ \times \mathbb{T}, \quad u|_{t=0} = K.$$

Setting

$$v(t, x) = u(t, -x), \quad (t, x) \in \mathbb{R}_+ \times \mathbb{T}, \quad \tilde{a}(x) = a(1 - x), \quad x \in \mathbb{T},$$

one easily checks that v formally solves

$$(5.28) \quad \partial_t v + \partial_x g(v) + \tilde{a}(x) \frac{v}{|v|^\alpha} = 0, \quad (t, x) \in \mathbb{R}_+ \times \mathbb{T}, \quad v|_{t=0} = K.$$

with $g = -f$, which satisfies $g'(0) = 0$ and $\inf_{[-K, K]} g'' > 0$. The fact that this transformation maps an admissible solution u of (5.27) to an admissible solution v of (5.28) is easy to check.

Therefore, the counterpart of Lemma 5.2 item (ii) also holds for solutions of (5.2) when f only satisfies (1.17):

Lemma 5.3. *Let f and K satisfy (1.17), $\delta > 0$ and $\omega = (0, A) \subset \mathbb{T}$. Then the solution u of (5.2) satisfies the following property: There exists $t_* \geq 0$ such that for all $t \geq t_*$, there exists an open subinterval $\omega(t) \subset \omega$ such that $u(t)|_{\omega(t)} = 0$, and*

$$|\omega \setminus \omega(t)| \leq \frac{C}{t^{1+\alpha}}, \quad \|u(t)\|_{L^\infty(\mathbb{T})} \leq \frac{C}{t}.$$

5.3. Negative constant initial datum. Let $K > 0$ and consider the solution u of

$$(5.29) \quad \begin{cases} \partial_t u + \partial_x f(u) + a(x) \frac{u}{|u|^\alpha} = 0, & (t, x) \in \mathbb{R}_+ \times \mathbb{T}, \\ u|_{t=0} = -K. \end{cases}$$

Then, setting

$$w(t, x) = -u(t, x), \quad (t, x) \in \mathbb{R}_+ \times \mathbb{T},$$

w formally solves

$$(5.30) \quad \begin{cases} \partial_t w + \partial_x h(w) + a(x) \frac{w}{|w|^\alpha} = 0, & (t, x) \in \mathbb{R}_+ \times \mathbb{T}, \\ w|_{t=0} = K, \end{cases}$$

where the flux h is given by

$$\forall s \in [-K, K], \quad h(s) = -f(-s).$$

It is easy to check that $h'(0) = 0$ and $\inf_{[-K, K]} |h''(s)| > 0$. Besides, this transformation also establishes the correspondence between the admissible solution u of (5.29) and the admissible solution w of (5.30).

Therefore, Lemma 5.2 item (ii) also holds when the initial datum is constant $= -K$, under the only condition that the flux f satisfies (1.17).

Lemma 5.4. *Let f and K satisfy (1.17), $\delta > 0$ and $\omega = (0, A) \subset \mathbb{T}$. Then the solutions u_\pm of*

$$(5.31) \quad \partial_t u_\pm + \partial_x f(u_\pm) + \delta 1_{(0, A)} \frac{u_\pm}{|u_\pm|^\alpha} = 0 \text{ in } \mathbb{R}_+ \times \mathbb{T}, \quad u_\pm|_{t=0} = \pm K \text{ in } \mathbb{T}.$$

satisfy the following property: There exists $t_ \geq 0$ such that for all $t \geq t_*$, there exists an open subinterval $\omega_\pm(t) \subset \omega$ such that $u_\pm(t)|_{\omega_\pm(t)} = 0$, and*

$$|\omega \setminus \omega_\pm(t)| \leq \frac{C}{t^{1+\alpha}}, \quad \|u_\pm(t)\|_{L^\infty(\mathbb{T})} \leq \frac{C}{t}.$$

5.4. Proof of Theorem 1.9. The proof of Theorem 1.9 follows by comparing the solution u of (1.1)–(1.2) with initial datum $u_0 \in L^\infty(\mathbb{T})$ with some reference solutions.

Namely, we assume that a satisfies (1.10) for some open interval $\omega \subset \mathbb{T}$. Up to a translation in space, we can assume that $\omega = (0, A)$. Therefore, the solution u of (1.1)–(1.2) with initial datum $u_0 \in L^\infty(\mathbb{T})$ with $\|u_0\|_{L^\infty} \leq K$, K as in (1.12), is sandwiched between the solutions u_\pm of (5.31).

We then immediately conclude Theorem 1.9 from Lemma 5.4.

6. NUMERICAL SIMULATIONS AND OPEN PROBLEMS

We present in this section some numerical experiments for various equations to which our theoretical results do not apply. The main numerical technique relies on the time-splitting scheme as time integrator. If one considers a general evolution equation

$$(6.1) \quad \begin{cases} \partial_t u = \mathcal{A}u + \mathcal{B}u, & (t, x) \in \mathbb{R}_+ \times \mathbb{T}, \\ u(0, x) = u_0(x), & x \in \mathbb{T}, \end{cases}$$

where \mathcal{A} and \mathcal{B} are (possibly non-linear) operators which need not commute. For a given time step $\delta t > 0$, set $t_n = n\delta t$, $n = 0, 1, 2, \dots$. Define the operators $S_{\mathcal{A}}$ and $S_{\mathcal{B}}$ associated respectively to the evolution equations

$$\partial_t u_{\mathcal{A}} = \mathcal{A}u_{\mathcal{A}}, \quad \partial_t u_{\mathcal{B}} = \mathcal{B}u_{\mathcal{B}}, \quad (t, x) \in \mathbb{R}_+ \times \mathbb{T},$$

The operators satisfy the following relations involving the exact solutions of the associated equations:

$$u_{\mathcal{A}}(t + \delta t) = S_{\mathcal{A}}(\delta t)u_{\mathcal{A}}(t) \quad \text{and} \quad u_{\mathcal{B}}(t + \delta t) = S_{\mathcal{B}}(\delta t)u_{\mathcal{B}}(t).$$

The splitting idea (see for example [19]) consists in approximating the continuous flow associated to (6.1) by a composition of operators $S_{\mathcal{A}}$ and $S_{\mathcal{B}}$ in the spirit of Trotter-Kato formula, the key for an efficient implementation being to solve efficiently these two reduced equations. We consider in this paper the second order Strang splitting scheme. Let $u^n(x)$ be the approximation of $u(t_n, x)$. The approximate solution to (6.1) at time t_{n+1} reads

$$(6.2) \quad u^{n+1} = S_{\mathcal{A}}(\delta t/2)S_{\mathcal{B}}(\delta t)S_{\mathcal{A}}(\delta t/2)u^n.$$

6.1. Scalar conservation laws. Let us now describe how it is applied to the equation (1.1). It involves the two reduced equation

$$(6.3) \quad \partial_t u + \partial_x f(u) = 0, \quad (t, x) \in \mathbb{R}_+ \times \mathbb{T},$$

and

$$(6.4) \quad \partial_t u = -a(x) \frac{u}{|u|^\alpha}, \quad (t, x) \in \mathbb{R}_+ \times \mathbb{T}.$$

The equation (6.3) is a standard nonlinear conservative hyperbolic equation. In the second equation (6.4), the space variable can be considered as a parameter and the equation reduced to an ordinary equation with solution

$$(6.5) \quad u(t, x) = \text{sign}(u_0(x)) (|u_0(x)|^\alpha - \alpha a(x)t)_+^{1/\alpha}.$$

If the flux f is linear, $f(u) = cu$, the solution to (6.3) is obviously

$$u(t, x) = u_0(x - ct).$$

For a general flux, we compute an approximate solution thanks to Rusanov scheme (see for example [22, p.233]). We identify the torus with $(0, 1)$ endowed with periodic boundary conditions and choose the spatial mesh size $\delta x > 0$ with $\delta x = 1/J$ with J denoting the number of nodes. The grid points are $x_j = j\delta x$, $j = 0, 1, \dots, J$. Let u_j^n be the full approximation to $u(t_n, x_j)$. The Rusanov scheme reads

$$u_j^{n+1} = u_j^n - \frac{\delta t}{\delta x} \left(F_{j+1/2}^n - F_{j-1/2}^n \right),$$

where the Rusanov numerical flux is given by

$$\begin{aligned} F_{j+1/2}^n &= F^{\text{Rus}}(u_j^n, u_{j+1}^n) \\ &= \frac{f(u_j^n) + f(u_{j+1}^n)}{2} - \frac{\max(|f'(u_j^n)|, |f'(u_{j+1}^n)|)}{2} (u_{j+1}^n - u_j^n). \end{aligned}$$

The theoretical results of previous sections apply to fluxes with assumption $f'(0) \neq 0$ or $f'(0) = 0$ with convexity hypothesis (convex or concave flux). Some fluxes do not satisfy such hypothesis. This is the case of the Buckley-Leverett flux which models

two phase fluid flow in a porous medium ([21]). In one space dimension the equation has the standard conservation law form ($k > 0$ is a parameter)

$$(6.6) \quad f_k^{BL}(u) = \frac{u^2}{u^2 + k(1-u)^2}.$$

We compute the evolution of the solution to (1.1) with $f_{1/4}^{BL}$ flux and the damping function $a(x)$ given by (1.18). The numerical parameters are $u_0(x) = K = 1.25$, $A = 1/4$, $\delta = 1$, $\delta t = 10^{-5}$ and $\delta x = 5 \cdot 10^{-5}$. The evolution of the solutions for $\alpha = 3/4$ and $\alpha = 1$ are plotted on Figure 6.1 and the evolution of their characteristic curves on Figure 6.2. The characteristic curves are computed as the evolution of a vector field with velocity given by the solution to (1.1). We see that contrary to convex (or concave) fluxes with $f'(0) = 0$, shock and rarefaction waves appear in finite time. Since the domain is a torus, the shock wave initiated from $x = A = 1/4$ propagates until the influence of the damping function a is enough important to annihilate the solution inside the support of a . We then recover a similar process as the one observed on Figure 5.1 where characteristic curves become vertical lines in finite time inside the support of the damping function a . The effect of decreasing α is to delay the extinction of the solution in $(0, A)$. The proof of this phenomenon is still missing.




FIGURE 6.1. Evolution of the solution for Buckley-Leverett flux $f_{1/4}^{BL}$ with $\alpha = 0.75$ (left) and $\alpha = 1$ (right)




FIGURE 6.2. Evolution of the characteristic curves for Buckley-Leverett flux $f_{1/4}^{BL}$ with $\alpha = 0.75$ (left) and $\alpha = 1$ (right)

6.2. Viscous Burgers equations. We consider here the convection diffusion equation given for $\mu > 0$ by

$$(6.7) \quad \partial_t u + \partial_x \left(\frac{u^2}{2} \right) = \mu \partial_x^2 u - a(x) \frac{u}{|u|^\alpha}, \quad (t, x) \in \mathbb{R}_+ \times \mathbb{T}.$$

We look at the solutions when $a(x) = \delta \mathbf{1}_\omega$, $\omega = (0, A)$, $A < 1$, the torus \mathbb{T} being the circle $(0, 1)$. This equation involves three different operators. We have to apply a second order three-operators splitting scheme which reads for the evolution equation $\partial_t u = (\mathcal{A} + \mathcal{B} + \mathcal{C})u$:

$$(6.8) \quad u^{n+1} = S_{\mathcal{A}}(\delta t/2) S_{\mathcal{B}}(\delta t/2) S_{\mathcal{C}}(\delta t) S_{\mathcal{B}}(\delta t/2) S_{\mathcal{A}}(\delta t/2) u^n,$$

where $S_{\mathcal{A}}$, $S_{\mathcal{B}}$ and $S_{\mathcal{C}}$ denote the flows associated to operators \mathcal{A} , \mathcal{B} and \mathcal{C} . Since we study the equation (6.7) on a torus, we benefit from the periodicity to use fast Fourier transform in order to make space approximation of the solutions of the heat equation

$$\partial_t w = \mu \partial_x^2 w.$$

The spatial mesh size is defined by $\delta x = 1/J$, $J = 2^P$, $P \in \mathbb{N}^*$. Since we discretize the heat equation by the Fourier spectral method, w_j^n and its Fourier transform satisfy the following relations:

$$w_j^n = \frac{1}{J} \sum_{m=-J/2}^{J/2-1} \hat{w}_m^n e^{i \xi_m (x_j - x_\ell)}, \quad j = 0, \dots, J-1,$$

and

$$\hat{w}_m^n = \sum_{j=0}^{J-1} w_j^n e^{-i\xi_m(x_j - x_\ell)}, \quad m = -\frac{J}{2}, \dots, \frac{J}{2} - 1,$$

where $\xi_m = 2\pi m$ for all $m = -\frac{J}{2}, \dots, \frac{J}{2} - 1$. The discrete Laplace operator Δ_P is therefore define by

$$(\widehat{\Delta_P v})_m = -\xi_m^2 \hat{v}_m, \quad v \in \mathbb{C}^M.$$

We present on Figure 6.3 the evolution of the logarithm of the solution. We choose the same numerical parameters used for Buckley-Leverett equation, the only difference relying on the mesh size which is $\delta x = 2^{-14}$. We present the logarithm to show that like in the hyperbolic case, the solution becomes zero on the support of the damping function a after a time T^* which depends on the parameter α . What is more surprising is the fact that after the time T^* , the solution on $(A, 1)$ behaves like the solution of the heat equation with homogeneous Dirichlet boundary conditions associated to the first eigenvalues of the Laplacian. We know that this solution on $(A, 1)$ is

$$v(t, x) = \exp(-\mu\lambda^2 t) \sin(\lambda(x - A)),$$

with $\lambda = \pi/(1 - A)$. We clearly identify this phenomenon by displaying the evolution of the L^∞ norm of the solution with respect to time on Figure 6.4. We plot both the L^∞ norm and a dashed line in log-scale with slope $-\mu\lambda^2$. The sin-like behavior of the solution on $[A, 1]$ for time $t = 10$ is also clearly present on Figure 6.4. A rigorous mathematical proof of the above observations is, to our knowledge, missing, despite the works [8, 11], where conditions for complete extinctions of the solutions are discussed (see also [7, 9, 10] for related results).




FIGURE 6.3. Evolution of the \log_{10} of the solution to (6.7) with $\alpha = 0.75$

We end up this paragraph by emphasizing that (6.7) is a viscous approximation of the Burgers equation, which is a conservation law fitting the assumptions of Theorem 1.9. It is thus completely natural to ask the behavior of (6.7) in large times, similarly to what has been done in Theorem 1.9. Though, as our numerical simulations underline, the large-time behavior of (6.7) is very different from the one of the Burgers equations predicted by Theorem 1.9. This is an evidence of the fact that the limit of large times and the limit of small viscosities do not commute, as observed in other contexts for instance in [20].




FIGURE 6.4. Evolution of the L^∞ norm of the solution to (6.7) in log-scale (left) and solution at time $t = 10$ (right) for $\alpha = 0.75$

6.3. Wave equation. We consider now the wave equation with homogeneous Dirichlet boundary conditions

$$(6.9) \quad \begin{cases} \partial_t^2 u - c^2 \partial_x^2 u = -a(x) \frac{\partial_t u}{|\partial_t u|^\alpha}, & (t, x) \in \mathbb{R}_+ \times (0, 1), \\ u(t, 0) = u(t, 1) = 0, \end{cases}$$

completed with initial conditions $u(0, x) = u_0(x)$ and $\partial_t u(0, x) = u_1(x)$.

To numerically simulate the solution to (6.9), we begin by transforming the equation as the first order hyperbolic system

$$\partial_t \begin{pmatrix} u \\ v \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ c^2 \partial_x^2 & 0 \end{pmatrix} \begin{pmatrix} u \\ v \end{pmatrix} + \begin{pmatrix} 0 \\ -a(x)v/|v|^\alpha \end{pmatrix}.$$

We can therefore apply the Strang splitting method (6.2). The solution to the ODE $\partial_t v = -a(x)v/|v|^\alpha$ is given by (6.5) and we approximate the free wave equation with the Newmark scheme ([25])

$$(6.10) \quad \begin{aligned} u_j^{n+1} &= u_j^n + \delta t v_j^n + \delta t^2 [\zeta c^2 w_j^{n+1} + (1/2 - \zeta) c^2 w_j^n], \\ v_j^{n+1} &= v_j^n + \delta t [(1 - \theta) c^2 w_j^n + \theta c^2 w_j^{n+1}], \end{aligned}$$

with $u_j^0 = u_0(x_j)$, $v_j^0 = u_1(x_j)$ and $w_j^k = (u_{j+1}^k - 2u_j^k + u_{j-1}^k)/(\delta x)^2$. We select for our numerical simulations $\theta = 1/2$ and $\zeta = 1/4$ for which the scheme is both second order in space and time and unconditionally stable.

We select the initial conditions

$$u_0(x) = K \begin{cases} 1 - e^{-0.1/(0.1-x)}, & \text{if } x < 0.1, \\ 1, & \text{if } 0.1 \leq x \leq 0.9, \\ 1 - e^{-0.1/(x-0.9)}, & \text{if } x > 0.9, \end{cases}$$

and $u_1(x) = 0$. The damping function is $a(x) = \delta \mathbf{1}_\omega$, $\omega = (3/8, 5/8)$. The numerical parameters are $\alpha = 1$, $c = 0.1$, $\delta t = 5 \cdot 10^{-4}$, $\delta x = 10^{-3}/3$, $\delta = 1$ and $K = 1.25$.

As can be expected (see Figure 6.5), the time derivative of the solution is annihilated on the support of a after a time T^* , the solution u becoming constant for $t > T^*$.

Let us underline that the linear wave equation is the prototype of a 2×2 system




FIGURE 6.5. Evolution of the solution to (6.9), u on the left and $\partial_t u$ on the right

of conservation laws, which can be easily seen with the use of characteristics. It is thus natural to consider such models as a generalization of (1.1)–(1.2). Note that the behavior of the solution of (6.9) when the damping acts everywhere in the domain has been studied in [5], or when the damping acts on the boundary [24], but the case of a localized damping term involving $\partial_t u$ still does not seem to be precisely described in the literature. In fact, the interested reader should also notice the close connection of this problem with the non-linear damped oscillator of the form

$$mx'' + \delta \frac{x'}{|x'|^\alpha} + \omega^2 x = 0, \quad t \geq 0,$$

with $m > 0$, $\alpha \in (0, 1]$, and $\omega > 0$, whose large time behavior is rather subtle, see e.g. [3, 26].

6.4. Schrödinger equation. The last equation we consider is the strongly damped cubic nonlinear Schrödinger (NLS) equation, motivated by the works [12, 13] in which the damping is effective everywhere. We thus wonder if the previous results for hyperbolic equations can be extended to the Schrödinger equation

$$(6.11) \quad i\partial_t u + \partial_x^2 u = -q|u|^2 u - ia(x) \frac{u}{|u|^\alpha}, \quad (t, x) \in \mathbb{R} \times \mathbb{T}.$$

with initial datum $u(0, x) = u_0(x)$. If the damping function a is zero, then the cubic NLS equation admits a special solution known as soliton. This solution is given by

$$(6.12) \quad u_{\text{sol}}(t, x) = \sqrt{\frac{2k}{q}} \operatorname{sech}(\sqrt{k}(x - ct)) \exp(i\frac{c}{2}(x - ct)) \exp(i(k + \frac{c^2}{4})t).$$

This solution evolves as the profile u_0 propagating at velocity c with time phase change. This solution for $c = 20$, $k = 0.81$ and the torus $x \in (-10, 10)$ is plotted on Figure 6.6. The numerical scheme again relies on the Strang splitting scheme for three operators (6.8). As for the Burgers heat equation, the space approximation is performed thanks to fast Fourier transform. The complex solution to ODE $\partial_t u = -a(x)u/|u|^\alpha$ is given by

$$u(t, x) = (|u_0|^\alpha - \alpha a(x)t)_+^{1/\alpha} \exp(i\operatorname{Arg}(u_0)),$$




FIGURE 6.6. Evolution of the modulus of the soliton (6.12) for $c = 20$ and $k = 0.81$.

whereas the solution to the ODE $i\partial_t u = -q|u|^2 u$ is given by

$$u(t, x) = \exp(itq|u_0(x)|^2)u_0(x).$$

We present the effects of the damping function $a(x) = \delta \mathbf{1}_\omega$, $\omega = (-10, -6) \cup (6, 10)$ on the soliton for $\mathbb{T} = (-10, 10)$ and $\alpha = 1$. The soliton initial datum overlaps the support of a . The numerical parameters are $\delta t = 5 \cdot 10^{-4}$ and $\delta x = 10 \cdot 2^{-12}$. As expected, the solution begins to propagate to the right direction and then vanishes on the support of a (see Figure 6.7). This is more clear on log scale. Again, to our knowledge, this behavior has not been proved rigorously in the literature.




FIGURE 6.7. Evolution of the solution to (6.11) in standard scale (left) and in log scale (right)

REFERENCES

- [1] S. ADLY, H. ATTOUCH, AND A. CABOT, *Finite time stabilization of nonlinear oscillators subject to dry friction*, in Nonsmooth mechanics and analysis, vol. 12 of Adv. Mech. Math., Springer, New York, 2006, pp. 289–304.
- [2] J. AGUIRRE AND M. ESCOBEDO, *A Cauchy problem for $u_t - \Delta u = u^p$ with $0 < p < 1$. Asymptotic behaviour of solutions*, Ann. Fac. Sci. Toulouse Math. (5), 8 (1986/87), pp. 175–203.

- [3] H. AMANN AND J. I. DÍAZ, *A note on the dynamics of an oscillator in the presence of strong friction*, *Nonlinear Anal.*, 55 (2003), pp. 209–216.
- [4] G. AMONTONS, *De la resistance causée dans les machines*, in *Mémoires de l'Académie Royale des Sciences Paris A*, 1699, pp. 257–282.
- [5] B. BAJI, A. CABOT, AND J. I. DÍAZ, *Asymptotics for some nonlinear damped wave equation: finite time convergence versus exponential decay results*, *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 24 (2007), pp. 1009–1028.
- [6] G. BASTIN AND J.-M. CORON, *Stability and boundary stabilization of 1-D hyperbolic systems*, vol. 88 of *Progress in Nonlinear Differential Equations and their Applications*, Birkhäuser/Springer, [Cham], 2016. Subseries in Control.
- [7] Y. BELAUD, *Time-vanishing properties of solutions of some degenerate parabolic equations with strong absorption*, *Adv. Nonlinear Stud.*, 1 (2001), pp. 117–152.
- [8] ———, *Extinction of solutions of some nonlinear parabolic equations*, *Sovrem. Mat. Fundam. Napravl.*, 36 (2010), pp. 5–11.
- [9] Y. BELAUD, B. HELFFER, AND L. VÉRON, *Long-time vanishing properties of solutions of some semilinear parabolic equations*, *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 18 (2001), pp. 43–68.
- [10] Y. BELAUD AND A. SHISHKOV, *Long-time extinction of solutions of some semilinear parabolic equations*, *J. Differential Equations*, 238 (2007), pp. 64–86.
- [11] Y. BELAUD AND A. SHISHKOV, *Extinction of solutions of semilinear higher order parabolic equations with degenerate absorption potential*, *J. Evol. Equ.*, 10 (2010), pp. 857–882.
- [12] R. CARLES AND C. GALLO, *Finite time extinction by nonlinear damping for the Schrödinger equation*, *Comm. Part. Diff. Eq.*, 36 (2011), pp. 961–975.
- [13] R. CARLES AND T. OZAWA, *Finite time extinction for nonlinear Schrödinger equation in 1D and 2D*, *Comm. in Partial Diff. Eq.*, 40 (2015), pp. 897–917.
- [14] J.-M. CORON, G. BASTIN, AND B. D'ANDRÉA NOVEL, *Dissipative boundary conditions for one-dimensional nonlinear hyperbolic systems*, *SIAM J. Control Optim.*, 47 (2008), pp. 1460–1498.
- [15] J.-M. CORON, B. D'ANDRÉA NOVEL, AND G. BASTIN, *A strict Lyapunov function for boundary control of hyperbolic systems of conservation laws*, *IEEE Trans. Automat. Control*, 52 (2007), pp. 2–11.
- [16] C. A. COULOMB, *Théorie des machines simples, en ayant égard au frottement de leurs parties, et la roideur des cordages*, *Mém. Math. Phys.*, X (1785), pp. 161–342.
- [17] C. M. DAFERMOS, *Hyperbolic conservation laws in continuum physics*, vol. 325 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*, Springer-Verlag, Berlin, 2000.
- [18] A. F. FILIPPOV, *Differential equations with discontinuous righthand sides*, vol. 18 of *Mathematics and its Applications (Soviet Series)*, Kluwer Academic Publishers Group, Dordrecht, 1988. Translated from the Russian.
- [19] E. HAIRER, C. LUBICH, AND G. WANNER, *Geometric Numerical Integration*, vol. 31 of *Springer Series in Computational Mathematics*, Springer, Heidelberg, 2010.
- [20] L. I. IGNAT, A. POZO, AND E. ZUAZUA, *Large-time asymptotics, vanishing viscosity and numerics for 1-D scalar conservation laws*, *Math. Comp.*, 84 (2015), pp. 1633–1662.
- [21] R. LEVEQUE, *Numerical Methods for Conservation Laws*, *Lectures in Mathematics*, ETH Zürich, Birkhäuser, second ed., 1992.
- [22] ———, *Finite volume methods for hyperbolic problems*, *Cambridge Texts in Applied Mathematics*, Cambridge University Press, Cambridge, 2002.
- [23] H. OLSSON, *Control systems with friction department of automatic control*, Lund Institute of Technology (LTH), (1996).
- [24] V. PERROLLAZ AND L. ROSIER, *Finite-time stabilization of 2×2 hyperbolic systems on tree-shaped networks*, *SIAM J. Control Optim.*, 52 (2014), pp. 143–163.
- [25] A. QUARTERONI AND A. VALLI, *Numerical Approximation of Partial Differential Equations*, *Springer Series in Computational Mathematics*, Springer, 2008.
- [26] J. L. VÁZQUEZ, *The nonlinearly damped oscillator*, *ESAIM Control Optim. Calc. Var.*, 9 (2003), pp. 231–246.

(C. Besse) INSTITUT DE MATHÉMATIQUES DE TOULOUSE, UMR5219, UNIVERSITÉ DE TOULOUSE,
CNRS, UPS IMT, F-31062 TOULOUSE CEDEX 9, FRANCE

E-mail address: `christophe.besse@math.univ-toulouse.fr`

(R. Carles) CNRS, IMAG, UMR 5149, MONTPELLIER, FRANCE

E-mail address: `Remi.Carles@math.cnrs.fr`

(S. Ervedoza) INSTITUT DE MATHÉMATIQUES DE TOULOUSE, UMR5219, UNIVERSITÉ DE TOULOUSE,
CNRS, UPS IMT, F-31062 TOULOUSE CEDEX 9, FRANCE

E-mail address: `sylvain.ervedoza@math.univ-toulouse.fr`