



HAL
open science

Quasi-isometric invariance of continuous group L^p -cohomology, and first applications to vanishings

Marc Bourdon, Bertrand Remy, Bertrand R ´ Emy

► **To cite this version:**

Marc Bourdon, Bertrand Remy, Bertrand R ´ Emy. Quasi-isometric invariance of continuous group
 L^p -cohomology, and first applications to vanishings. 2018. hal-01742591v1

HAL Id: hal-01742591

<https://hal.science/hal-01742591v1>

Preprint submitted on 9 Apr 2018 (v1), last revised 31 Dec 2020 (v3)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

QUASI-ISOMETRIC INVARIANCE OF CONTINUOUS GROUP L^p -COHOMOLOGY, AND FIRST APPLICATIONS TO VANISHINGS

MARC BOURDON AND BERTRAND RÉMY

ABSTRACT. We show that the continuous L^p -cohomology of locally compact second countable groups is a quasi-isometric invariant. As an application, we prove partial results supporting a positive answer to a question asked by M. Gromov, suggesting a classical behaviour of continuous L^p -cohomology of simple real Lie groups. In addition to quasi-isometric invariance, the ingredients are a spectral sequence argument and Pansu's vanishing results for real hyperbolic spaces. In the best adapted cases of simple Lie groups, we obtain nearly half of the relevant vanishings.

2010 Mathematics Subject Classification: 20J05, 20J06, 22E15, 22E41, 53C35, 55B35, 57T10, 57T15.

Keywords and Phrases: L^p -cohomology, topological group, Lie group, symmetric space, quasi-isometric invariance, spectral sequence, cohomology vanishing, root system.

CONTENTS

1. Introduction	2
2. Continuous cohomology	5
3. Continuous group L^p -cohomology	9
4. Proof of quasi-isometric invariance	12
5. Semi-direct products	15
6. The (possibly) non-vanishing strip	19
7. The admissible simple real Lie groups	22
8. Numerical efficiency of the method	25
9. Vanishing in degree one	31
References	35

1. INTRODUCTION

Let G be a locally compact second countable group. Then G is countable at infinity [Bou07, IX §2.9, Cor. to Prop. 16, p. 22] and it admits a compatible left invariant proper metric [CdlH16, Prop. 1.D.1]. In addition, it is also equipped with a left invariant Haar measure [Bou63, VII §1.2, Th. 1], which enables us to define the spaces $L^p(G)$ [Bou65, IV §3.4, Déf. 2].

In this paper we study the *group L^p -cohomology* of G for $p > 1$. By definition, this is the continuous cohomology of G [BW00, Chap. IX], with coefficients in the right-regular representation on $L^p(G)$. It shall be denoted by $H_{\text{ct}}^*(G, L^p(G))$. We also consider the associated *reduced cohomology*, denoted by $\overline{H}_{\text{ct}}^*(G, L^p(G))$ (they are the biggest Hausdorff quotients of the previous spaces – see Definition 2.2).

Our first result is the quasi-isometry invariance of these cohomology spaces (see Corollary 3.8 for details).

Theorem 1.1. *Let G_1 and G_2 be locally compact second countable groups, equipped with left-invariant proper metrics. If G_1 and G_2 are quasi-isometric, then $H_{\text{ct}}^*(G_1, L^p(G_1))$ and $H_{\text{ct}}^*(G_2, L^p(G_2))$ are isomorphic as graded topological vector spaces. The same holds for reduced cohomology.*

For example the group L^p -cohomology of a group G as above is isomorphic to the one of any of its cocompact lattices. When G is semi-simple, its group L^p -cohomology is isomorphic to the one of any of its parabolic subgroups; by Levi decomposition, this generalizes to (the points of) any connected real algebraic group.

Theorem 1.1 was already known for the finite $K(\pi, 1)$ -groups (see [Gro93, p. 219] and Remark 3.4), and then proved for finitely generated groups [Ele98]. In degree 1, the theorem is due to Cornuier-Tessera [CT11]. We prove it (in any degree) by relating the group L^p -cohomology with the asymptotic L^p -cohomology. The latter has been defined by Pansu, and it is known to be a quasi-isometric invariant [Pan95]. We recently learnt from R. Sauer and M. Schrödl that they had used the same comparison strategy with $p = 2$, in order to prove the quasi-isometric invariance of ℓ^2 -Betti numbers of unimodular locally compact second countable groups [SS17].

Our motivation for Theorem 1.1, which leads to the second main theorem of this paper, is understanding the L^p -cohomology of semi-simple groups of real rank ≥ 2 . In fact, we have in mind the following question, asked by M. Gromov [Gro93, p. 253].

Question 1.2. *Let G be a simple real Lie group. We assume that $l = \text{rk}_{\mathbf{R}}(G) \geq 2$. Let k be an integer $< l$ and p be a real number > 1 . Do we have: $H_{\text{ct}}^k(G, L^p(G)) = \{0\}$?*

The previous theorem allows one to reduce the study of the group L^p -cohomology of a Lie group G as in the question, to the one of any of its parabolic subgroups. Such a parabolic subgroup, say P , admits a Levi decomposition, which is a semi-direct product decomposition $P = M \ltimes AN$ where M is semi-simple and AN is the (solvable) radical of P . The class for which we can fruitfully use Theorem 1.1 consists of the simple Lie groups in which some maximal parabolic subgroup has suitable geometric properties.

Theorem 1.3. *Let G be a simple real Lie group admitting a parabolic subgroup whose radical is quasi-isometric to some real hyperbolic space whose dimension is denoted by d . Let D denote the dimension of the Riemannian symmetric space attached to G .*

- (1) *We have: $H_{\text{ct}}^0(G, L^p(G)) = \{0\}$, and also: $H_{\text{ct}}^k(G, L^p(G)) = \{0\}$ for $k \geq D$.*
- (2) *Suppose now that $k \in \{1, \dots, D - 1\}$. Then we have:*

$$H_{\text{ct}}^k(G, L^p(G)) = \{0\} \text{ for } k \leq \frac{d-1}{p} \text{ and for } k \geq \frac{d-1}{p} + D - d + 2.$$

Our results also deal with reduced cohomology – see Theorem 6.1 for details.

By lack of better terminology, the simple Lie groups admitting a parabolic subgroup whose radical is quasi-isometric to a real hyperbolic space are called *admissible* in the present paper. The prototype of simple Lie group, namely $\text{SL}_n(\mathbf{R})$, is admissible, and in this case we obtain:

Corollary 1.4. *For $G = \text{SL}_n(\mathbf{R})$, one has $H_{\text{ct}}^k(G, L^p(G)) = \{0\}$ for $k \leq \lfloor \frac{n^2}{4} \rfloor \cdot \frac{1}{p}$ and for $k \geq \lfloor \frac{n^2}{4} \rfloor \cdot \frac{1}{p} + \lfloor \frac{(n+1)^2}{4} \rfloor$.*

We also note that the same circle of ideas can be used to prove the following result, which is a particular case of a general result of Pansu [Pan07] and Cornuier-Tessera [CT11], saying that $H_{\text{ct}}^1(G, L^p(G)) =$

$\{0\}$ for every $p > 1$ and every connected Lie group G , unless G is Gromov hyperbolic or amenable unimodular.

Corollary 1.5. *For every admissible simple Lie group G such that $\text{rk}_{\mathbf{R}}(G) \geq 2$ and every $p > 1$, we have: $H_{\text{ct}}^1(G, L^p(G)) = \{0\}$.*

This result also follows from the fixed-point property for continuous affine isometric actions of semi-simple groups of real rank ≥ 2 on L^p -spaces, which is established in [BFGM07].

Let us finish this introduction by describing roughly our method and by discussing its efficiency and its validity.

The main idea is to combine tools from homological algebra, namely the Hochschild-Serre spectral sequence [BW00, IX.4], together with geometric results such as Pansu's description of the L^p -cohomology of real hyperbolic spaces [Pan99, Pan08]. We also use resolutions already considered in [Bla79], whose measurable nature makes them flexible enough to describe nicely intermediate spaces involved in the spectral sequences. The general outcome is a new description of the L^p -cohomology of G – see Theorem 6.1 and Remark 6.4.

In order to discuss the efficiency of the method, let us reformulate Theorem 1.3 as follows: for each admissible group G and each exponent $p > 1$, we exhibit an interval of degrees, whose explicit length does not depend on p and outside of which vanishing of L^p -cohomology is guaranteed (*i.e.*, this interval is a strip of potential non-vanishing of L^p -cohomology). First, by Lie-theoretic considerations, we can provide the full list of admissible simple Lie groups. It contains several infinite sequences of groups, say $\{G_l\}_{l \geq 2}$ where $l = \text{rk}_{\mathbf{R}}(G)$. In the best cases, the dimension D_l of the symmetric space of G_l and the dimension d_l of the radical of the suitable parabolic subgroup are quadratic polynomials in l , and we have $\lim_{l \rightarrow \infty} \frac{d_l}{D_l} = \frac{1}{2}$. Since the width of the strip of potential non-vanishing is $D_l - d_l + 2$, this makes us say that we obtain in the best cases of admissible groups nearly half of the relevant vanishings.

At last, our results can also be compared with Borel's result [Bor85] about the L^2 -cohomology of connected semi-simple Lie groups G , which makes heavy use of representation theory while we use a more geometric approach. It asserts that:

- $\overline{H_{\text{ct}}^k(G, L^2(G))} = 0$ unless $k = \frac{D}{2}$,
- $H_{\text{ct}}^k(G, L^2(G)) \neq 0$ at least for $k \in \left(\frac{D}{2} - \frac{l_0}{2}, \frac{D}{2} + \frac{l_0}{2}\right]$,

where D is the dimension of the Riemannian symmetric space G/K , and where l_0 is the difference between the complex rank of G and the complex rank of K . In the case $G = \mathrm{SL}_n(\mathbf{R})$, one has $D = \frac{n^2+n-2}{2}$ and $l_0 = \lfloor \frac{n-1}{2} \rfloor$. See also [BFS14] for related results about vanishing of the reduced L^2 -cohomology.

Organization of the paper. Section 2 is a brief presentation of some standard topics in cohomology of groups. Section 3 discusses group L^p -cohomology: we first consider the discrete group case and recall some classical connections with the simplicial L^p -cohomology; in the general case, we state the existence of an isomorphism between the group L^p -cohomology and the asymptotic L^p -cohomology (Theorem 3.7). This result, whose proof is given in Section 4, implies the first quasi-isometric invariance theorem (Theorem 1.1 of the introduction). Section 5 contains some preliminary results on the group L^p -cohomology of semi-direct products via the Hochschild-Serre spectral sequence. Section 6 focuses on applications of the previous sections to the group L^p -cohomology of admissible simple real Lie groups (an improved version of Theorem 1.3 of the introduction). Corollary 1.4 is proved in this section. Section 7 provides the list of admissible groups and Section 8 gives tables in which the numerical efficiency of the method can be discussed. Finally, we prove Corollary 1.5 on vanishing in degree 1 in Section 9.

Acknowledgements. We thank Nicolas Monod, whose paper [Mon10] was a valuable source of inspiration. We also thank Pierre Pansu: the present paper elaborates on several of his results. At last, we thank Y. Cornulier for useful remarks and R. Sauer and M. Schrödl for drawing our attention to their paper [SS17]. M.B. was partially supported by the Labex Cempi and B.R. by the GeoLie project (ANR-15-CE40-0012). Both authors were supported by the GDSous/GSG project (ANR-12-BS01-0003).

2. CONTINUOUS COHOMOLOGY

We review some of the basic notions of the continuous cohomology of groups. We refer to [BW00, Chapter IX] or to [Gui80] and their references for more details.

Let X, Y be topological spaces. The set of continuous mappings $f : X \rightarrow Y$ equipped with the compact open topology is denoted by $C(X, Y)$.

Let G be a locally compact second countable group, and let (π, V) be a *topological G -module* i.e. a Hausdorff locally convex vector space over \mathbf{R} on which G acts via a continuous representation π . We denote by $V^G \subset V$ the subspace of $\pi(G)$ -invariant vectors.

For $k \in \mathbf{N}$, let $C^k(G, V) := C(G^{k+1}, V)$ be the set of continuous maps from G^{k+1} to V equipped with the compact open topology. Then $C^k(G, V)$ is a topological G -module by means of the following action: for $g, x_0, \dots, x_k \in G$,

$$(g \cdot f)(x_0, \dots, x_k) = \pi(g)(f(g^{-1}x_0, \dots, g^{-1}x_k)).$$

Consider the following complex

$$C^0(G, V)^G \xrightarrow{d_0} C^1(G, V)^G \xrightarrow{d_1} C^2(G, V)^G \xrightarrow{d_2} \dots,$$

where, for $k \in \mathbf{N}$ and $x_0, \dots, x_{k+1} \in G$,

$$(2.1) \quad (d_k f)(x_0, \dots, x_{k+1}) = \sum_{i=0}^{k+1} (-1)^i f(x_0, \dots, \hat{x}_i, \dots, x_{k+1}).$$

Definition 2.2. (1) The *continuous cohomology of G with coefficients in (π, V)* is the cohomology of this complex. It is the collection of groups $H_{\text{ct}}^*(G, V) = \{H_{\text{ct}}^k(G, V)\}_{k \in \mathbf{N}}$, where $H_{\text{ct}}^0(G, V) = \text{Ker } d_0 \simeq V^G$ and $H_{\text{ct}}^k(G, V) = \text{Ker } d_k / \text{Im } d_{k-1}$ for $k \geq 1$.

- (2) The *reduced continuous cohomology* is the collection $\overline{H}_{\text{ct}}^*(G, V) = \{\overline{H}_{\text{ct}}^k(G, V)\}_{k \in \mathbf{N}}$, where $\overline{H}_{\text{ct}}^0(G, V) = \text{Ker } d_0 \simeq V^G$, and where $\overline{H}_{\text{ct}}^k(G, V) = \text{Ker } d_k / \overline{\text{Im } d_{k-1}}$ for $k \geq 1$ ($\overline{\text{Im } d_{k-1}}$ denotes the closure of $\text{Im } d_{k-1}$ in $C^k(G, V)$).
- (3) When G is a discrete group, we omit the subscript “ct” and simply write $H^k(G, V)$ and $\overline{H}^k(G, V)$.

We now recall some useful notions in homological algebra.

- A map $f : A \rightarrow B$ between topological G -modules is a *G -morphism* if it is a continuous G -equivariant linear map.
- A complex of topological G -modules and G -morphisms

$$0 \rightarrow V \xrightarrow{d_{-1}} A^0 \xrightarrow{d_0} A^1 \xrightarrow{d_1} A^2 \xrightarrow{d_2} \dots$$

is said to be a *strong G -resolution of V* if for every $k \in \mathbf{N}$, there exists a continuous linear map $h_k : A^k \rightarrow A^{k-1}$ (not a G -morphism) such that $h_0 \circ d_{-1} = \text{id}$ and for every $k \in \mathbf{N}$

$$d_{k-1} \circ h_k + h_{k+1} \circ d_k = \text{id}.$$

- A topological G -module U is *strong injective* if for every strong G -injection $0 \rightarrow A \rightarrow B$, every G -morphism $f : A \rightarrow U$ extends to a G -morphism $\bar{f} : B \rightarrow U$ (neither is required to be strong).
- A *strong injective G -resolution of V* is a strong G -resolution $0 \rightarrow V \xrightarrow{d_{-1}} A^0 \xrightarrow{d_0} A^1 \xrightarrow{d_1} A^2 \xrightarrow{d_2} \dots$ in which the A^k 's are strong injective.

Let $A^* = (A^0 \xrightarrow{d_0} A^1 \xrightarrow{d_1} A^2 \xrightarrow{d_2} \dots)$ and $B^* = (B^0 \xrightarrow{d_0} B^1 \xrightarrow{d_1} B^2 \xrightarrow{d_2} \dots)$ be complexes of Hausdorff locally convex vector spaces.

- They are *homotopy equivalent* if there exists continuous homomorphisms $\varphi_* : A^* \rightarrow B^*$, $\psi_* : B^* \rightarrow A^*$, $\alpha_* : A^* \rightarrow A^{*-1}$, $\beta_* : B^* \rightarrow B^{*-1}$, such that φ_* and ψ_* commute with d_* , and such that for every $k \in \mathbf{N}$:

$$\begin{aligned} d_{k-1} \circ \alpha_k + \alpha_{k+1} \circ d_k &= \text{id} - \psi_k \circ \varphi_k \\ d_{k-1} \circ \beta_k + \beta_{k+1} \circ d_k &= \text{id} - \varphi_k \circ \psi_k, \end{aligned}$$

with the convention $A^{-1} = B^{-1} = 0$ and $d_{-1} = \alpha_0 = \beta_0 = 0$.

- When B^* is a subcomplex of A^* , one says that A^* *retracts on B^** if they are homotopy equivalent as above, with ψ_* equal to the inclusion.

Example 2.3. The complex

$$0 \rightarrow V \xrightarrow{d_{-1}} C^0(G, V) \xrightarrow{d_0} C^1(G, V) \xrightarrow{d_1} C^2(G, V) \xrightarrow{d_2} \dots$$

is a strong injective G -resolution of V . Indeed, first it is a strong resolution; one defines the h_k 's by $(h_k f)(x_0, \dots, x_{n-1}) = f(1, x_0, \dots, x_{n-1})$. Secondly, $C^k(G, V)$ is isomorphic to $C^0(G, C^{k-1}(G, V))$; and for every topological G -module (π, W) the module $C^0(G, W)$ is strong injective.

Indeed given a G -injection $0 \rightarrow A \xrightarrow{i} B$ and a continuous linear map $h : B \rightarrow A$ such that $h \circ i = \text{id}$, every G -morphism $f : A \rightarrow C^0(G, W)$ extends to $\bar{f} : B \rightarrow C^0(G, W)$ by letting for $b \in B$ and $x \in G$:

$$\bar{f}(b)(x) = \pi(x) \left(f(h(x^{-1}b))(1) \right).$$

Example 2.4. Suppose G is a discrete group that acts properly discontinuously and freely on a contractible locally finite simplicial complex X by simplicial automorphisms. Let $X^{(k)}$ be the set of k -simplices of X and let $C^k(X, V)$ be the set of linear maps from the vector space $\mathbf{R}X^{(k+1)}$ to V . Let $\delta_k : C^k(X, V) \rightarrow C^{k+1}(X, V)$ be defined as follows; for $f \in C^k(X, V)$, and $\sigma \in X^{(k+1)}$,

$$(\delta_k f)(\sigma) = f(\partial\sigma).$$

The $C^k(X, V)$'s are G -modules for the following action: for $g \in G$, $f \in C^k(X, V)$ and $\sigma \in X^{(k)}$,

$$(g \cdot f)(\sigma) = \pi(g)(f(g^{-1}(\sigma))).$$

The δ_k 's are G -morphisms. The complex

$$0 \rightarrow V \xrightarrow{\delta_{-1}} C^0(X, V) \xrightarrow{\delta_0} C^1(X, V) \xrightarrow{\delta_1} C^2(X, V) \xrightarrow{\delta_2} \dots$$

is a strong injective G -resolution of V . Indeed one defines the h_k 's by induction on k by using a retraction of X to a point. To show that $C^k(X, V)$ is strong injective, one considers a fundamental domain D in X , the set $X_D^{(k)} := \{\sigma \in X^{(k)} \mid \text{the first vertex of } \sigma \text{ belongs to } D\}$ and the vector space $C_D^k(X, V)$ of linear maps from $\mathbf{R}X_D^{(k+1)}$ to V . Then one can show that $C^k(X, V) \simeq C^0(G, C_D^k(X, V))$. The latter is strong injective (see the discussion in Example 2.3 above).

Strong injective resolutions compute the continuous cohomology:

Proposition 2.5. *Suppose $0 \rightarrow V \rightarrow A^*$ and $0 \rightarrow V \rightarrow B^*$ are strong injective G -resolutions of V . Then the complexes $(A^*)^G$ and $(B^*)^G$ are homotopy equivalent.*

Since homotopy equivalent complexes have the same cohomology, one obtains

Corollary 2.6. *Suppose $0 \rightarrow V \rightarrow A^*$ is a strong injective G -resolution of V . Then, the cohomology and the reduced cohomology of the complex $(A^*)^G$ are topologically isomorphic to $H_{\text{ct}}^*(G, V)$ and $\overline{H}_{\text{ct}}^*(G, V)$, respectively.*

Proof. From Example 2.3 and Proposition 2.5 above, the complexes $(A^*)^G$ and $C^*(G, V)$ are homotopy equivalent. Therefore, by standard arguments, their cohomological spaces (equipped with the quotient topology) are topologically isomorphic. By definition, the cohomology of $C^*(G, V)$ is $H_{\text{ct}}^*(G, V)$. Thus the cohomology $H^*((A^*)^G)$ of $(A^*)^G$ is topologically isomorphic to $H_{\text{ct}}^*(G, V)$. The reduced cohomological spaces of $(A^*)^G$ and $C^*(G, V)$ are respectively topologically isomorphic to $H^k((A^*)^G)/\overline{\{0\}}$ and $H_{\text{ct}}^k(G, V)/\overline{\{0\}}$, where $\overline{\{0\}}$ denotes the closure of the null subspace. Since $H^k((A^*)^G)$ and $H_{\text{ct}}^k(G, V)$ are topologically isomorphic, so are $H^k((A^*)^G)/\overline{\{0\}}$ and $H_{\text{ct}}^k(G, V)/\overline{\{0\}}$. \square

3. CONTINUOUS GROUP L^p -COHOMOLOGY

The group L^p -cohomology is defined in this section (Definition 3.1). When the group is a finite $K(\pi, 1)$, we relate its group L^p -cohomology with the simplicial L^p -cohomology (Proposition 3.2). For locally compact second countable topological groups, the group L^p -cohomology is isomorphic to the asymptotic L^p -cohomology (Theorem 3.7). This implies Theorem 1.1 of the introduction.

Definition 3.1. Let G be a locally compact second countable topological group, and \mathcal{H} be a left-invariant Haar measure. Let $p \in (1, +\infty)$. The *group L^p -cohomology* of G is the continuous cohomology of G with coefficients in the right-regular representation of G on $L^p(G, \mathcal{H})$, *i.e.* the representation defined by

$$(\pi(g)u)(x) = u(xg) \quad \text{for } u \in L^p(G, \mathcal{H}) \quad \text{and } g, x \in G.$$

It will be denoted by $H_{\text{ct}}^*(G, L^p(G))$. Similarly is defined the *reduced group L^p -cohomology* of G denoted by $\overline{H}_{\text{ct}}^*(G, L^p(G))$. When G is discrete we omit the subscript “ct” and simply write $H^*(G, L^p(G))$ and $\overline{H}^*(G, L^p(G))$.

Observe that the right regular representation is isometric if and only if \mathcal{H} is bi-invariant.

The discrete group case. In this paragraph we relate the group L^p -cohomology of certain discrete groups with the simplicial L^p -cohomology of some simplicial complexes (Proposition 3.2). This is a standard material. We present it for completeness and also because its proof can be seen as a model for the general case.

Let X be a simplicial complex and let $X^{(k)}$ be the set of its k -simplices. We will always assume that X has *bounded geometry* *i.e.* that there is an $N \in \mathbf{N}$ such that each simplex intersects at most N simplices.

Let $C^{k,p}(X)$ be the Banach space of linear maps $u : \mathbf{R}X^{(k)} \rightarrow \mathbf{R}$ such that

$$\|u\|^p := \sum_{\sigma \in X^{(k)}} |u(\sigma)|^p < \infty.$$

The *simplicial L^p -cohomology* of X is the cohomology of the complex

$$C^{0,p}(X) \xrightarrow{\delta_0} C^{1,p}(X) \xrightarrow{\delta_1} C^{2,p}(X) \xrightarrow{\delta_2} \dots,$$

where the δ_k 's are defined as in Example 2.4. Similarly is defined its reduced simplicial L^p -cohomology. They are denoted by $L^p\mathbb{H}^*(X)$ and $L^p\overline{\mathbb{H}}^*(X)$ respectively.

Proposition 3.2. *Suppose that G acts by simplicial automorphisms, properly discontinuously, freely and cocompactly, on a locally finite contractible simplicial complex X . Then the complexes $C^*(G, L^p(G))^G$ and $C^{*,p}(X)$ are homotopy equivalent. In particular there are topological isomorphisms $\mathbb{H}^*(G, L^p(G)) \simeq L^p\mathbb{H}^*(X)$ and $\overline{\mathbb{H}}^*(G, L^p(G)) \simeq L^p\overline{\mathbb{H}}^*(X)$.*

Proof. According to Corollary 2.6 and Example 2.4, $C^*(G, L^p(G))^G$ is homotopy equivalent to the complex

$$(3.3) \quad C^0(X, L^p(G))^G \xrightarrow{\delta_0} C^1(X, L^p(G))^G \xrightarrow{\delta_1} C^2(X, L^p(G))^G \xrightarrow{\delta_2} \dots$$

In particular their cohomologies (reduced or not) are topologically isomorphic. To prove the proposition it is enough to show the latter complex is topologically isomorphic to the complex $C^{*,p}(X)$.

To this end, one considers the map $\Phi : C^{k,p}(X) \rightarrow C^k(X, L^p(G))$ defined by $\Phi(u) = f$ with $f(\sigma)(g) := u(g\sigma)$. With the norms expressions, one sees that it is a topological embedding. Its image is $C^k(X, L^p(G))^G$. Indeed $\Phi(u)$ is clearly G -invariant. Moreover, for $f \in C^k(X, L^p(G))^G$, one defines $u : X^{(k)} \rightarrow \mathbf{R}$ by $u(\sigma) = f(\sigma)(1)$. Since f is invariant, one has $u(g\sigma) = f(\sigma)(g)$. Because G acts properly discontinuously and cocompactly on X , by using the partition of $X^{(k)}$ into orbits, one gets that $u \in C^{k,p}(X)$. \square

Remark 3.4. Equip every simplicial complex X with the length metric obtained by identifying every simplex with the standard Euclidean one. A simplicial complex is called *uniformly contractible* if it is contractible and if there exists a function $\rho : (0, +\infty) \rightarrow (0, +\infty)$ such that every ball $B(x, r) \subset X$ can be contracted to a point in $B(x, \rho(r))$. Among bounded geometry uniformly contractible simplicial complexes, the simplicial L^p -cohomology (reduced or not) is invariant by quasi-isometry. Indeed if X and Y are quasi-isometric bounded geometry uniformly contractible simplicial complexes, then the complexes $C^{*,p}(X)$ and $C^{*,p}(Y)$ are homotopy equivalent (see [Gro93, p. 219], and [BP03] for a detailed proof). As a consequence, the above proposition implies that the group L^p -cohomology of fundamental groups of finite aspherical simplicial complexes is invariant by quasi-isometry. We will give below another proof of this fact that applies in a much larger generality.

The general case. Suppose that G is a locally compact second countable topological group. We will relate the group L^p -cohomology and the asymptotic L^p -cohomology of G . The latter has been considered by Pansu in [Pan95], see also [Gen14]. It is a quasi-isometric invariant (see Theorem 3.6 below).

Following [Pan95], we define the asymptotic L^p -cohomology of a metric measure space. Let $X = (X, d, \mu)$ be a metric measure space. Suppose it satisfies the following “bounded geometry” condition. There exists increasing functions $v, V : (0, +\infty) \rightarrow (0, +\infty)$ such for every ball $B(x, r) \subset X$ one has

$$v(r) \leq \mu(B(x, r)) \leq V(r).$$

For $s > 0$ and $k \in \mathbf{N}$, let $\Delta_s^{(k)} = \{(x_0, x_1, \dots, x_k) \in X^{k+1} \mid d(x_i, x_j) \leq s\}$. Let $AS^{k,p}(X)$ be the set of the (classes of) functions $u : X^{k+1} \rightarrow \mathbf{R}$ such that for every $s > 0$ one has

$$N_s(u)^p := \int_{\Delta_s^{(k)}} |u(x_0, \dots, x_k)|^p d\mu(x_0) \dots d\mu(x_k) < \infty.$$

We equip $AS^{k,p}(X)$ with the topology induced by the set of the seminorms N_s ($s > 0$).

Definition 3.5. The asymptotic L^p -cohomology of X is the cohomology of the complex $AS^{0,p}(X) \xrightarrow{d_0} AS^{1,p}(X) \xrightarrow{d_1} AS^{2,p}(X) \xrightarrow{d_2} \dots$, where the d_k 's are defined as in (2.1). Similarly is defined the reduced asymptotic L^p -cohomology. They are denoted by $L^pH_{AS}^*(X)$ and $L^p\overline{H}_{AS}^*(X)$ respectively.

Asymptotic L^p -cohomology (reduced or not) is invariant by quasi-isometry. In fact one has

Theorem 3.6 ([Pan95]). *Let X and Y be bounded geometry metric spaces and let $f : X \rightarrow Y$ be a quasi-isometry. Then f induces a homotopy equivalence between the complexes $AS^{*,p}(X)$ and $AS^{*,p}(Y)$. Moreover the associated isomorphism of graded topological vector spaces $f^* : L^pH_{AS}^*(Y) \rightarrow L^pH_{AS}^*(X)$ depends only the bounded perturbation class of f .*

See also [Gen14] for a more detailed proof. In the next section we will prove the

Theorem 3.7. *Suppose G is equipped with a left-invariant proper metric. Then $C^*(G, L^p(G))^G$ and $AS^{*,p}(G)$ are homotopy equivalent complexes. In particular there exist topological isomorphisms:*

$$H_{\text{ct}}^*(G, L^p(G)) \simeq L^p H_{AS}^*(G) \text{ and } \overline{H}_{\text{ct}}^*(G, L^p(G)) \simeq L^p \overline{H}_{AS}^*(G).$$

In combination with Theorem 3.6 one obtains the following result, which implies Theorem 1.1 of the introduction:

Corollary 3.8. *Let G_1 and G_2 be locally compact second countable topological groups, equipped with left-invariant proper metrics and let $f : G_1 \rightarrow G_2$ be a quasi-isometry. Then f induces a homotopy equivalence between the complexes $C^*(G_1, L^p(G_1))^{G_1}$ and $C^*(G_2, L^p(G_2))^{G_2}$. Moreover the associated isomorphism of graded topological vector spaces between the group L^p -cohomologies of G_1 and G_2 depends only the bounded perturbation class of f .*

Remark 3.9. Suppose X is a bounded geometry simplicial complex. In general its asymptotic and simplicial L^p -cohomologies are different (for example the asymptotic L^p -cohomology of a finite simplicial complex is trivial since it is quasi-isometric to a point). Pansu [Pan95] asked whether the asymptotic and the simplicial L^p -cohomologies coincide for uniformly contractile bounded geometry simplicial complexes. He proved that it is indeed the case for those which are in addition non-positively curved; for these spaces the complexes $C^{*,p}(X)$ and $AS^{*,p}(X)$ are homotopy equivalent.

4. PROOF OF QUASI-ISOMETRIC INVARIANCE

As explained in the previous section the quasi-isometric invariance of the group L^p -cohomology is a consequence of Theorem 3.7. We prove this theorem in this section. The proof is inspired by the one of Proposition 3.2. It will use a strong injective G -resolution (Lemma 4.1) that appears in Blanc [Bla79]. As already mentioned, a similar strategy was used by R. Sauer and M. Schrödl to prove the quasi-isometric invariance of ℓ^2 -Betti numbers of unimodular locally compact second countable groups [SS17].

Suppose that the topological vector space V is Fréchet (*i.e.* metrizable and complete). Let X be a locally compact second countable topological space endowed with a Borel measure μ . Let $p \in (1, +\infty)$. We denote by $L_{\text{loc}}^p(X, V)$ the set of (classes of) measurable functions

$f : X \rightarrow V$ such that for every compact $K \subset X$ and every continuous semi-norm N on V defining its topology, one has

$$\|f\|_{K,N} := \left(\int_K N(f(x))^p d\mu(x) \right)^{\frac{1}{p}} < \infty.$$

Suppose that G acts on X continuously by preserving μ . Equipped with the set of semi-norms $\|\cdot\|_{K,N}$, the vector space $L^p_{\text{loc}}(X, V)$ is a Fréchet G -module for the action $(g \cdot f)(x) = \pi(g)(f(g^{-1}x))$, see [Bla79]. Let \mathcal{H} be a left-invariant Haar measure on G . Denote by G^k the cartesian product of k copies of G equipped with the product measure.

Lemma 4.1 ([Bla79]). *The complex*

$$0 \rightarrow V \xrightarrow{d_{-1}} L^p_{\text{loc}}(G, V) \xrightarrow{d_0} L^p_{\text{loc}}(G^2, V) \xrightarrow{d_1} L^p_{\text{loc}}(G^3, V) \xrightarrow{d_2} \dots, \text{ with the } d_k \text{ 's defined as in (2.1), is a strong injective } G\text{-resolution of } V.$$

In the special case $V = L^p(G)$, one can express the spaces $L^p_{\text{loc}}(X, V)^G$ as follows.

Lemma 4.2. *Let X, μ, G, \mathcal{H} as above. Let E be the topological vector space that consists of the (classes of the) Borel functions $u : X \rightarrow \mathbf{R}$ such that for every compact subset $K \subset X$ one has*

$$\|u\|_K^p := \int_G \int_K |u(gx)|^p d\mu(x) d\mathcal{H}(g) < \infty.$$

Then E is topologically isomorphic to the Fréchet space $L^p_{\text{loc}}(X, L^p(G))^G$.

Proof. Let $u \in E$. Since u is Borel and since the map

$$\varphi : (g, x) \in G \times X \mapsto gx \in X$$

is continuous, the function $u \circ \varphi$ is Borel too, and its class depends only on the class of u . Moreover $u \circ \varphi = 0$ almost everywhere if and only if $u = 0$ almost everywhere. Indeed, given $A \subset X$, one has $\varphi^{-1}(A) = \cup_{g \in G} (\{g\} \times g^{-1}A)$, and thus $(\mathcal{H} \times \mu)(\varphi^{-1}(A)) = 0$ if and only if $\mu(A) = 0$.

Define $\Phi : E \rightarrow L^p_{\text{loc}}(X, L^p(G))$ by $\Phi(u) = f$ with

$$f(x)(g) := u(gx) = u \circ \varphi(g, x) \text{ for every } x \in X, g \in G.$$

An element $f \in L^p_{\text{loc}}(X, L^p(G))$ is null if and only if $f(x)(g) = 0$ for almost all $(g, x) \in G \times X$. Therefore the above discussion in combination with the semi-norms expressions and the Fubini-Tonelli theorem imply that Φ is a topological embedding. It remains to prove that its image is the subspace $L^p_{\text{loc}}(X, L^p(G))^G$.

Let $u \in E$, we show that $\Phi(u)$ is G -invariant. Write $f = \Phi(u)$ for simplicity. One has for $h \in G$ and for almost all $x \in X$ and $g \in G$:

$$(h \cdot f)(x)(g) = f(h^{-1}x)(gh) = u(gx) = f(x)(g).$$

Thus f is G -invariant.

Let $f \in L_{\text{loc}}^p(X, L^p(G))^G$, we are looking for $u \in E$ such that $\Phi(u) = f$. One would like to define u by $u(x) = f(x)(1)$; but this has no meaning in general since $f(x) \in L^p(G)$. Let $(U_n)_{n \in \mathbf{N}}$ be a decreasing sequence of relatively compact open neighborhoods of 1 in G such that $\bigcap_{n \in \mathbf{N}} U_n = \{1\}$. Define a measurable function $u : X \rightarrow \mathbf{R}$ by

$$u(x) = \limsup_{n \rightarrow \infty} \frac{1}{\mathcal{H}(U_n)} \int_{U_n} f(x)(h^{-1}) d\mathcal{H}(h).$$

Let $g \in G$. Since f is G -invariant and \mathcal{H} is left-invariant, one has for almost every $x \in X$

$$\int_{U_n} f(gx)(h^{-1}) d\mathcal{H}(h) = \int_{U_n} f(x)(h^{-1}g) d\mathcal{H}(h) = \int_{gU_n} f(x)(h^{-1}) d\mathcal{H}(h).$$

Therefore Lebesgue's theorem implies that $u(gx) = f(x)(g)$ for almost all $g \in G$ and $x \in X$. By modifying u to a Borel function in the same class if necessary, one obtains that $u \in E$ and that $\Phi(u) = f$. \square

According to Proposition 2.5 and Lemma 4.1, to finish the proof of Theorem 3.7, it is enough to establish:

Lemma 4.3. *The complexes $L_{\text{loc}}^p(G^{*+1}, L^p(G))^G$ and $AS^{*,p}(G)$ are topologically isomorphic.*

Proof. Denote by \mathcal{H}^n the product measure on G^n . According to Lemma 4.2, it is enough to show that the following two sets of semi-norms (on the space of measurable functions $u : G^{k+1} \rightarrow \mathbf{R}$) define the same topology. The first set of semi-norms is the one considered in Lemma 4.2 in the case $X = (G^{k+1}, \mathcal{H}^{k+1})$. These are the N_K 's, with $K \subset G^{k+1}$ compact, defined by

$$N_K(u)^p = \int_G \int_K |u(gx_0, \dots, gx_k)|^p d\mathcal{H}^{k+2}(g, x_0, \dots, x_k).$$

The second one is the set of semi-norms that appears in the definition of the asymptotic L^p -cohomology of (G, d, \mathcal{H}) . These are the N_s 's, with $s > 0$, defined by

$$N_s(u)^p = \int_{\Delta_s} |u(y_0, \dots, y_k)|^p d\mathcal{H}^{k+1}(y_0, \dots, y_k).$$

For a compact subset $U \subset G$, set

$$K_s^U := \{(x_0, x_1, \dots, x_k) \in \Delta_s \mid x_0 \in U\}.$$

Since the metric on G is proper, K_s^U is compact. Moreover every compact subset of G^{k+1} is contained in such a subset. Let Δ be the modular function on G . Since the metric and the measure are left-invariant, one has

$$\begin{aligned} N_{K_s^U}(u)^p &= \\ &= \int_{g \in G} \int_{x \in U} \int_{(y_1, \dots, y_k) \in \Delta_s} |u(gx, gxy_1, \dots, gxy_k)|^p d\mathcal{H}^{k+2}(g, x, y_1, \dots, y_k) \\ &= \int_{g \in G} \int_{x \in U} \int_{(y_1, \dots, y_k) \in \Delta_s} \Delta(x) |u(g, gy_1, \dots, gy_k)|^p d\mathcal{H}^{k+2}(g, x, y_1, \dots, y_k) \\ &= C(U) \cdot N_s(u)^p, \end{aligned}$$

where $C(U) := \int_U \Delta(x) d\mathcal{H}(x)$. Thus the two sets of semi-norms define the same topology. \square

5. SEMI-DIRECT PRODUCTS

We partially relate the L^p -cohomology of semi-direct products $P = R \rtimes Q$ with the L^p -cohomology of the normal subgroups R (see Corollaries 5.4 and 5.5). The next section will contain examples of application.

Some generalities. Let V be a Hausdorff locally convex topological vector space, and let X be a locally compact second countable topological space endowed with a Borel measure μ . Let $p \in (1, +\infty)$. We denote by $L^p(X, V)$ the set of (classes of) measurable functions $f : X \rightarrow V$ such that for every continuous semi-norm N on V defining the topology of V , one has

$$\|f\|_N^p := \int_X N(f(x))^p d\mu(x) < \infty.$$

Equipped with the set of semi-norms $\|\cdot\|_N$, the vector space $L^p(X, V)$ is a Hausdorff locally convex vector space. When V is a Fréchet space, or a Banach space, then so is $L^p(X, V)$.

Proposition 5.1. *Suppose X, Y are topological spaces as above endowed with Borel measures. Then $L^p(X, L_{\text{loc}}^p(Y, V))$ is topologically isomorphic to $L_{\text{loc}}^p(Y, L^p(X, V))$.*

Proof. Let $f : X \times Y \rightarrow V$ be a measurable function. For a compact subset $K \subset Y$ and a continuous semi-norm N on V , denote by $N_{K,N}$ and $\|\cdot\|_N$ the associated semi-norms on $L^p_{\text{loc}}(Y, V)$ and $L^p(X, V)$ respectively. One has by Fubini-Tonelli

$$\|f\|_{N_{K,N}}^p = \int_X \int_K N(f(x, y))^p d\mu_X(x) d\mu_Y(y) = \|f\|_{K, \|\cdot\|_N}^p.$$

The proposition follows. \square

Hochschild-Serre spectral sequence. Let R be a closed normal subgroup of a locally compact second countable group P and let (π, V) be a topological P -module. The P -action on $C^*(R, V)$, defined by

$$(g \cdot f)(x_0, \dots, x_k) = \pi(g)(f(g^{-1}x_0g, \dots, g^{-1}x_kg))$$

for every $g \in P$, $f \in C^k(R, V)$ and $x_0, \dots, x_k \in R$, induces a P/R -action on the topological vector spaces $H_{\text{ct}}^*(R, V)$ [Gui80, p. 50].

The Hochschild-Serre spectral sequence relates the continuous cohomology $H_{\text{ct}}^*(P, V)$ to the cohomologies $H_{\text{ct}}^*(P/R, H_{\text{ct}}^*(R, V))$ – see *e.g.* [BW00, Chapter IX, Th. 4.1 p. 178] or [Gui80, Prop. 5.1 p. 214].

In the special case (π, V) is the right regular representation on $L^p(P)$ and P is a semi-direct product $P = R \rtimes Q$ (where R and Q are closed subgroups) we give in this subsection a more comprehensive description of these cohomological spaces. In the following statement the measures \mathcal{H}_R and \mathcal{H}_Q are left-invariant Haar measures on R and Q respectively. We denote by Δ the modular function for the Q -action by conjugation on R , *i.e.* the function on Q defined by $q^{-1}\mathcal{H}_Rq = \Delta(q) \cdot \mathcal{H}_R$, for every $q \in Q$.

Proposition 5.2. *Let $k \in \mathbf{N}$. Assume that $H_{\text{ct}}^k(R, L^p(R))$ is Hausdorff and that the complex $C^*(R, L^p(R))^R$ is homotopically equivalent to a complex of Banach spaces. Then $H_{\text{ct}}^k(R, L^p(P))$ is Hausdorff and there is a canonical isomorphism of topological Q -modules:*

$$H_{\text{ct}}^k(R, L^p(P)) \simeq L^p(Q, H_{\text{ct}}^k(R, L^p(R))),$$

where, in the latter space, the group Q is equipped with the measure $\Delta^{-1} \cdot \mathcal{H}_Q$, and where the Q -action is induced by

$$(q \cdot f)(y)(x_0, \dots, x_k) = f(yq)(q^{-1}x_0q, \dots, q^{-1}x_kq),$$

for every $q, y \in Q$, $f : Q \rightarrow C^k(R, L^p(R))$, and $x_0, \dots, x_k \in R$.

Proof. As a consequence of Blanc's lemma (Lemma 4.1) the complex $L_{\text{loc}}^p(R^{*+1}, L^p(P))^R$ retracts to its subcomplex $C^*(R, L^p(P))^R$. Thus, it is not only that these two complexes have the same cohomology, but the Q -action on the cohomology can be expressed in the same manner.

We claim that $L^p(P) \simeq L^p(Q, L^p(R))$ as topological P -modules, where Q is equipped with the measure $\Delta^{-1} \cdot \mathcal{H}_Q$, and where the expression of the right-regular representation of P on the latter module is $(\pi(r, q)f)(x, y) = f(xryy^{-1}, yq)$.

Indeed this follows from Fubini, by observing that $\Delta^{-1} \cdot \mathcal{H}_R \times \mathcal{H}_Q$ is a left-invariant measure on $P = R \rtimes Q$.

Therefore, with Proposition 5.1, we obtain that $L_{\text{loc}}^p(R^{*+1}, L^p(P))^R$ is isomorphic to $L^p(Q, L_{\text{loc}}^p(R^{*+1}, L^p(R))^R)$. In this representation, one can check that the Q -action on the latter complex can be written as in the statement of the proposition.

It remains to prove that the k th-cohomology space of the complex $L^p(Q, L_{\text{loc}}^p(R^{*+1}, L^p(R))^R)$ is isomorphic to $L^p(Q, H_{\text{ct}}^k(R, L^p(R)))$. By assumption and from Lemma 4.1, the complex $L_{\text{loc}}^p(R^{*+1}, L^p(R))^R$ is homotopy equivalent to a complex of Banach spaces that we denote by B^* . Since every continuous linear map $\varphi : V_1 \rightarrow V_2$ between Hausdorff locally convex topological vector spaces, extends to a continuous linear map $\varphi_* : L^p(Q, V_1) \rightarrow L^p(Q, V_2)$ defined by $\varphi_*(f) = \varphi \circ f$, the above homotopy equivalence induces a homotopy equivalence between the complexes $L^p(Q, L_{\text{loc}}^p(R^{*+1}, L^p(R))^R)$ and $L^p(Q, B^*)$. Since $H^k(B^*) \simeq H_{\text{ct}}^k(R, L^p(R))$, the following lemma ends the proof of the proposition. \square

Lemma 5.3. *Suppose $k \in \mathbf{N}$ and that B^* is a complex of Banach spaces such that $H^k(B^*)$ is Hausdorff. Then the k th-cohomology space of the complex $L^p(Q, B^*)$ satisfies $H^k(L^p(Q, B^*)) \simeq L^p(Q, H^k(B^*))$ canonically and topologically.*

Proof of Lemma 5.3. Consider the complex $L^p(Q, B^*)$. One has clearly $\text{Ker } d|_{L^p(Q, B^k)} = L^p(Q, \text{Ker } d|_{B^k})$. We claim that

$$\text{Im}(d : L^p(Q, B^{k-1}) \rightarrow L^p(Q, B^k)) = L^p(Q, \text{Im}(d : B^{k-1} \rightarrow B^k)).$$

The direct inclusion is obvious. For the reverse one, we use the assumption that $\text{Im}(d : B^{k-1} \rightarrow B^k)$ is a Banach space in combination with the following Michael's theorem [Mic56, Prop. 7.2], that previously appears in [Mon01, p. 93] in the context of bounded cohomology:

Suppose $\varphi : B \rightarrow C$ is a continuous surjective linear map between Banach spaces. Then for every $\lambda > 1$ there exists a continuous (non linear) section $\sigma : C \rightarrow B$ such that for every $c \in C$ one has

$$\|\sigma(c)\| \leq \lambda \inf\{\|b\| ; \varphi(b) = c\}.$$

The natural map $L^p(Q, \text{Ker } d|_{B^k}) \rightarrow L^p(Q, \text{H}^k(B^*))$ and the above equalities induce an injective continuous map

$$\text{H}^k(L^p(Q, B^*)) \rightarrow L^p(Q, \text{H}^k(B^*)).$$

It is surjective, thanks again to Michael's theorem, since the projection map $\text{Ker } d|_{B^k} \rightarrow \text{H}^k(B^*)$ is a surjective continuous linear map between Banach spaces.

Since $\text{H}^k(B^*)$ is a Banach space, so is $L^p(Q, \text{H}^k(B^*))$. From the previous description of $\text{Ker } d|_{L^p(Q, B^k)}$ and $\text{Im } d|_{L^p(Q, B^{k-1})}$, the cohomological space $\text{H}^k(L^p(Q, B^*))$ is also a Banach space (for the quotient norm). Therefore Banach's theorem implies that the above isomorphism is a topological one. \square

As a consequence of Proposition 5.2, the Hochschild-Serre spectral sequence for L^p -cohomology takes the following form:

Corollary 5.4. *Suppose that $P = R \rtimes Q$ where R and Q are closed subgroups of P . Assume that $C^*(R, L^p(R))^R$ is homotopically equivalent to a complex of Banach spaces and that every cohomology space $\text{H}_{\text{ct}}^k(R, L^p(R))$ is Hausdorff. Then, there exists a spectral sequence (E_r) , abutting to $\text{H}_{\text{ct}}^*(P, L^p(P))$, in which*

$$E_2^{k,\ell} = \text{H}_{\text{ct}}^k\left(Q, L^p\left(Q, \text{H}_{\text{ct}}^\ell(R, L^p(R))\right)\right).$$

Corollary 5.5. *Suppose that $P = R \rtimes Q$ where R and Q are closed subgroups of P . Assume that $C^*(R, L^p(R))^R$ is homotopically equivalent to a complex of Banach spaces. Suppose also that there exists $n \in \mathbf{N}$, such that $\text{H}_{\text{ct}}^k(R, L^p(R)) = 0$ for $0 \leq k < n$ and such that $\text{H}_{\text{ct}}^n(R, L^p(R))$ is Hausdorff. Then $\text{H}_{\text{ct}}^k(P, L^p(P)) = 0$ for $0 \leq k < n$ and there is a linear isomorphism*

$$\text{H}_{\text{ct}}^n(P, L^p(P)) \simeq L^p\left(Q, \text{H}_{\text{ct}}^n(R, L^p(R))\right)^Q.$$

We notice that the latter isomorphism is just a linear one, in particular it does not imply that $\text{H}_{\text{ct}}^n(P, L^p(P))$ is Hausdorff.

6. THE (POSSIBLY) NON-VANISHING STRIP

Let G be a non-compact simple Lie group. As a consequence of Corollary 3.8, the group L^p -cohomology of G is isomorphic to the one of any of its parabolic subgroups. Indeed, every parabolic subgroup acts cocompactly on G , and thus is quasi-isometric to G .

Recall that a parabolic subgroup decomposes as $P = M \ltimes AN$ with M semi-simple, $A \simeq \mathbf{R}^r$, and N nilpotent. One can expect to use this decomposition to derive, from the Hochschild-Serre spectral sequence, some informations about the group L^p -cohomology of P . To do so, and according to Corollaries 5.4 and 5.5, one has to know a bit of the L^p -cohomology of the group AN . When the rank r of A is at least 2, very few is known about the L^p -cohomology of AN (apart from the vanishing of $H_{\text{ct}}^1(AN, L^p(AN))$ for all p – see [Pan07, CT11]).

In contrast, when the rank of A is one, *i.e.* when P is a maximal proper parabolic subgroup, the group AN admits an invariant negatively curved Riemannian metric, and the L^p -cohomology of negatively curved manifolds has been investigated by Pansu [Pan99, Pan08].

We focus on the class of simple Lie groups G that admit a maximal proper parabolic subgroup $P = M \ltimes AN$ with AN quasi-isometric to a real hyperbolic space \mathbb{H}^d of constant negative curvature. We call such groups G *admissible*. They will be classified in the next section. The elementary case $G = \text{SL}_n(\mathbf{R})$ is discussed in Example 6.5. We obtain the following partial description of their cohomology:

Theorem 6.1. *Let G be an admissible simple Lie group. Let $P = M \ltimes AN$ be a maximal proper parabolic subgroup with radical AN quasi-isometric to \mathbb{H}^d . Let $X = G/K$ be the associated symmetric space and let $D = \dim X$. One has:*

- (1) $H_{\text{ct}}^0(G, L^p(G)) = H_{\text{ct}}^k(G, L^p(G)) = 0$ for $k \geq D$.
- (2) *Suppose $k \in \{1, \dots, D-1\}$. Then:*
 - $H_{\text{ct}}^k(G, L^p(G)) = 0$ for $k \leq \frac{d-1}{p}$ and for $k \geq \frac{d-1}{p} + D - d + 2$,
 - $\overline{H}_{\text{ct}}^k(G, L^p(G)) = 0$ for $\frac{d-1}{p} + D - d + 1 \leq k < \frac{d-1}{p} + D - d + 2$,
 - *Suppose in addition that $\frac{d-1}{p} \notin \mathbf{N}$, and let $\ell = \lceil \frac{d-1}{p} \rceil + 1$. Then for $k > \frac{d-1}{p}$, there is a linear isomorphism:*

$$H_{\text{ct}}^k(G, L^p(G)) \simeq H_{\text{ct}}^{k-\ell} \left(M, L^p \left(M, H_{\text{ct}}^\ell(AN, L^p(AN)) \right) \right).$$

In particular one sees that the possible couples (p, k) for which $H_{\text{ct}}^k(G, L^p(G)) \neq 0$ lie in the strip $\frac{d-1}{p} < k < \frac{d-1}{p} + D - d + 2$ of

width $D - d + 2$. The numerical computation of the width will occupy Section 8.

A key ingredient in the proof of Theorem 6.1 is the following lemma which is a straightforward consequence of Pansu's results.

Lemma 6.2. *Suppose that AN is a Lie group quasi-isometric to a real hyperbolic space \mathbb{H}^d . Then for $p > 1$, its group L^p -cohomology satisfies*

- (1) $H_{\text{ct}}^0(AN, L^p(AN)) = H_{\text{ct}}^k(AN, L^p(AN)) = 0$ for $k \geq d$.
- (2) Suppose $k \in \{1, \dots, d - 1\}$. Then:
 - $H_{\text{ct}}^k(AN, L^p(AN))$ is Hausdorff if and only if $k \neq \frac{d-1}{p} + 1$,
 - $H_{\text{ct}}^k(AN, L^p(AN)) = 0$ if and only if either $k \leq \frac{d-1}{p}$ or $k > \frac{d-1}{p} + 1$.

Proof of Lemma 6.2. By Corollary 3.8, the group L^p -cohomology of AN is isomorphic to the one of any cocompact lattice in $\text{Isom}(\mathbb{H}^d)$; which in turn is the same as the simplicial L^p -cohomology of \mathbb{H}^d – see Proposition 3.2. By the simplicial L^p -cohomology of a complete Riemannian manifold X , we mean the simplicial L^p -cohomology of any bounded geometry quasi-isometric simplicial decomposition of X . Now, the simplicial L^p -cohomology of \mathbb{H}^d has been computed by Pansu [Pan99, Pan08] ⁽¹⁾. The result of this computation is precisely the statement of our lemma. \square

Parts of Theorem 6.1 rely on the following version of Poincaré duality established in [Pan08] Corollaire 14 ⁽²⁾.

Lemma 6.3. *Let X be a complete Riemannian manifold of bounded geometry and of dimension D . Denote by $L^pH^*(X)$ and $L^p\overline{H}^*(X)$ its simplicial L^p -cohomology and its reduced simplicial L^p -cohomology. Let $q = p/(p - 1)$ and $0 \leq k \leq D$. Then*

¹In [Pan99, Pan08], Pansu computes the de Rham L^p -cohomology of \mathbb{H}^d , i.e. the cohomology of the de Rham complex of L^p differential forms with differentials in L^p . In [Pan95], he proves an L^p -cohomology version of the de Rham theorem, namely the homotopy equivalence between the de Rham L^p -complex and the simplicial L^p -complex, for complete Riemannian manifolds of bounded geometry. The fact that the simplicial L^p -cohomology is Hausdorff for $k \neq \frac{d-1}{p} + 1$, and vanishes for $k < \frac{d-1}{p}$ and $k > \frac{d-1}{p} + 1$, can also be established without the de Rham theorem, as a consequence of Corollary B in [Bou16].

²Again, Pansu establishes this result for the de Rham L^p -cohomology. One obtains the result for the simplicial L^p -cohomology by using the L^p -cohomology version of the de Rham theorem. The lemma can also be established more directly without the de Rham theorem, as a consequence of Proposition 1.2 and Theorem 1.3 in [Bou16].

- (1) $L^p H^k(X)$ is Hausdorff if and only if $L^q \overline{H}^{D-k+1}(X)$ is Hausdorff.
 (2) $L^p \overline{H}^k(X) = 0$ if and only if $L^q \overline{H}^{D-k}(X) = 0$.

Proof of Theorem 6.1. (1). According to Corollary 3.8, the group L^p -cohomology of G is isomorphic to the one of any of its cocompact lattices; which in turn is isomorphic to the simplicial L^p -cohomology of X – see Proposition 3.2. The latter is trivially null in degree 0 and degrees $k > \dim X$. It is also null in degree $D = \dim X$. Indeed, the lemma above implies that $L^p \overline{H}^D(X) = 0$. Moreover, since X satisfies a linear isoperimetric inequality, $L^p H^1(X)$ is Hausdorff – see [Pan07] or [Gro93]. Hence the lemma implies that $L^p H^D(X)$ is Hausdorff, and so it is null.

(2). Since G and P are quasi-isometric, one has $H_{\text{ct}}^*(G, L^p(G)) \simeq H_{\text{ct}}^*(P, L^p(P))$ by Corollary 3.8. The vanishing of $H_{\text{ct}}^k(P, L^p(P))$ for every $k \leq \frac{d-1}{p}$, follows directly from the analogous result for AN in Lemma 6.2, and from Corollary 5.5 applied with $P = M \rtimes AN$, $Q = M$ and $R = AN$. We note that the assumptions of Corollary 5.5 are satisfied; indeed $C^*(AN, L^p(AN))^{AN}$ is homotopy equivalent to the simplicial L^p -complex of \mathbb{H}^d which is a complex of Banach spaces.

To show that $H_{\text{ct}}^k(G, L^p(G)) = 0$ for $k \geq \frac{d-1}{p} + D - d + 2$, we use again the isomorphism between the group L^p -cohomology of G and the simplicial L^p -cohomology of X . We know from above that the latter is null for $k \leq \frac{d-1}{p}$. Then Lemma 6.3 implies that $L^p \overline{H}^k(X)$ is null for $k \geq \frac{d-1}{p} + D - d + 1$ and that $L^p H^k(X)$ is Hausdorff for $k \geq \frac{d-1}{p} + D - d + 2$. The proof of the first two items is now complete.

Suppose that $\frac{d-1}{p} \notin \mathbf{N}$. Then, according to Lemma 6.2, the only degree ℓ such that $H_{\text{ct}}^\ell(AN, L^p(AN))$ is non-zero, is $\ell = \lfloor \frac{d-1}{p} \rfloor + 1$. Moreover $H_{\text{ct}}^\ell(AN, L^p(AN))$ is Hausdorff. By applying Corollary 5.4, one obtains the third item. \square

Remark 6.4. In view of the last item in Theorem 6.1, to analyse further the case where $\frac{d-1}{p} < k < \frac{d-1}{p} + D - d + 2$, in particular to decide whether $H_{\text{ct}}^k(G, L^p(G))$ vanishes or not, one would like to take advantage of a good description of $H_{\text{ct}}^\ell(AN, L^p(AN))$ and of the M -action on it. In particular one could try to exploit the description of $H_{\text{ct}}^\ell(AN, L^p(AN))$ as a functional space on $N \simeq \mathbf{R}^{d-1} \simeq \partial \mathbb{H}^d \setminus \{\infty\}$, that is established in Section 8.2 of [Pan99] (see also [Pan89] or [Rez08] or [BP03] for the case $\ell = 1$). We will follow this idea in Section 9 to prove the vanishing of the first group L^p -cohomology of admissible Lie groups of real rank ≥ 2 , see Corollary 1.5.

Example 6.5. Every maximal parabolic subgroup of $\mathrm{SL}_n(\mathbf{R})$ is conjugated to a subgroup of the form

$$P = \left\{ \begin{pmatrix} p_1 & p_3 \\ 0 & p_2 \end{pmatrix} \in \mathrm{SL}_n(\mathbf{R}) \mid p_1 \in M_{s,s}, p_2 \in M_{n-s,n-s}, p_3 \in M_{s,n-s} \right\},$$

with $0 < s < n$. Moreover $P = M \rtimes AN$ with

$$M^0 = \left\{ \begin{pmatrix} m_1 & 0 \\ 0 & m_2 \end{pmatrix} \mid m_1 \in \mathrm{SL}_s(\mathbf{R}), m_2 \in \mathrm{SL}_{n-s}(\mathbf{R}) \right\},$$

$$A = \left\{ \begin{pmatrix} \lambda_1 I & 0 \\ 0 & \lambda_2 I \end{pmatrix} \mid \lambda_1, \lambda_2 \in \mathbf{R}_+^*, \lambda_1^s \cdot \lambda_2^{n-s} = 1 \right\} \simeq \mathbf{R}_+^*,$$

and $N = \left\{ \begin{pmatrix} I & x \\ 0 & I \end{pmatrix} \mid x \in M_{s,n-s} \right\} \simeq (\mathbf{R}^{s(n-s)}, +)$. Since one has

$$\begin{pmatrix} \lambda_1 I & 0 \\ 0 & \lambda_2 I \end{pmatrix} \begin{pmatrix} I & x \\ 0 & I \end{pmatrix} \begin{pmatrix} \lambda_1 I & 0 \\ 0 & \lambda_2 I \end{pmatrix}^{-1} = \begin{pmatrix} I & \lambda_1 \lambda_2^{-1} x \\ 0 & I \end{pmatrix} = \begin{pmatrix} I & \lambda_1^{\frac{n}{n-s}} x \\ 0 & I \end{pmatrix},$$

we get that $AN \simeq \mathbf{R} \rtimes_{\varphi} \mathbf{R}^{s(n-s)}$, with $\varphi(t)(x) = e^{\frac{tn}{n-s}} x$. Therefore AN is isometric to $\mathbb{H}^{s(n-s)+1}$ and $\mathrm{SL}_n(\mathbf{R})$ is admissible. Take $s = \lfloor \frac{n}{2} \rfloor$. Then one has $d-1 = \dim N = s(n-s) = \frac{n^2}{4}$ if n is even, and $\frac{n^2-1}{4}$ otherwise. Since $D = \dim X = \frac{(n-1)(n+2)}{2}$, Theorem 6.1 gives the following possibly non-vanishing strip for the L^p -cohomology of $\mathrm{SL}_n(\mathbf{R})$:

- $\frac{n^2}{4p} < k < \frac{n^2}{4p} + \frac{n(n+2)}{4}$ if n is even, and
- $\frac{n^2-1}{4p} < k < \frac{n^2-1}{4p} + \frac{(n+1)^2}{4}$ if n is odd,

which can be summarized by: $\lfloor \frac{n^2}{4} \rfloor \cdot \frac{1}{p} < k < \lfloor \frac{n^2}{4} \rfloor \cdot \frac{1}{p} + \lfloor \frac{(n+1)^2}{4} \rfloor$.

7. THE ADMISSIBLE SIMPLE REAL LIE GROUPS

We are looking for the simple Lie groups that we called admissible, *i.e.* that contain a proper parabolic subgroup, say P with Levi decomposition $M_P \rtimes A^P N^P$ where A^P is a split torus and N^P is a unipotent group normalized by A^P , for which the following metric condition holds: (*) *the underlying Riemannian variety of the radical $A^P N^P$ is quasi-isometric to a real hyperbolic space.*

The list of admissible simple Lie groups is given by the following.

Proposition 7.1. *The admissible simple real Lie groups are those whose relative root system is of type A_l , B_l , C_l , D_l , E_6 and E_7 . In particular, for classical types the only excluded groups are those with non-reduced root system BC_l .*

Proof. Let P_\emptyset be a minimal parabolic subgroup in G containing A , a maximal split torus. The subgroup A defines a set Φ of roots with respect to A and the inclusion $A \subset P_\emptyset$ defines, inside Φ , a subset Φ^+ of positive roots and a subset Δ of simple roots. We have: $\Delta \subset \Phi^+ \subset \Phi$. Any parabolic subgroup is conjugated to exactly one of the parabolic subgroups containing P_\emptyset , and the latter subgroups are parametrized by subsets of Δ . More precisely, they are of the form $P_{\Delta'} = M_{\Delta'} \ltimes A^{\Delta'} N^{\Delta'}$, where $M_{\Delta'}$ is the Levi subgroup generated by the roots of Φ in the linear span $\mathbf{Z}\Delta'$ of $\Delta' \subset \Delta$, the subtorus $A^{\Delta'}$ of A lies in the kernels of the roots in Δ' and the Lie algebra of the unipotent group $N^{\Delta'}$ is linearly spanned by the weight spaces indexed by the roots in $\Phi^+ \setminus \Phi(\Delta')$ where $\Phi(\Delta') = \Phi \cap \mathbf{Z}\Delta'$. This description is consistent with the notation P_\emptyset for a minimal parabolic subgroup; note that we also have $G = P_\Delta$.

Let us go back to condition (*). By homogeneity, it implies that $A^P N^P = A^P \ltimes N^P$ is actually homothetic to a real hyperbolic space, and in terms of roots it implies that A^P must act on N^P via a single character. The first consequence is that P must be maximal for inclusion among proper parabolic subgroups. Therefore, since we work up to conjugacy, this implies that we may – and shall – assume that there is a simple root $\gamma \in \Delta$ such that $P = P_{\Delta \setminus \{\gamma\}}$. To simplify notation, we set: $A_\gamma = A^{\Delta \setminus \{\gamma\}}$ and $N_\gamma = N^{\Delta \setminus \{\gamma\}}$, so that $P = P_{\Delta \setminus \{\gamma\}} = M_{\Delta \setminus \{\gamma\}} \ltimes A_\gamma N_\gamma$.

The torus A_γ is 1-dimensional and it acts on N_γ (more precisely, on the root subspaces of the Lie algebra of N_γ) via powers of γ since it lies in the kernel of any other simple root of the basis Δ . Note that the simple root γ itself always appears as a weight in the root space decomposition of $\text{Lie}(N_\gamma)$. Therefore, for $A_\gamma N_\gamma$ to be homothetic to a real hyperbolic space, it is necessary and sufficient that A_γ act on N_γ via the single character γ (*i.e.* we must avoid powers γ^k with $k \geq 2$).

We have obtained this way a combinatorial interpretation of condition (*). Indeed, let δ be a root occurring as a weight in the root space decomposition of $\text{Lie}(N_\gamma)$: it is a root in Φ^+ for which the coordinate along γ in the basis Δ has a coefficient $m \geq 1$ (since vanishing amounts to being in the root system $\Phi(\Delta \setminus \{\gamma\})$ of $M_{\Delta \setminus \{\gamma\}}$). In this case, any $a \in A_\gamma$ acts on the root space of weight δ via the scalar $\delta(a)^m$.

The remaining task now is to investigate the descriptions of root systems, and to check for which ones there is a simple root $\gamma \in \Delta$ with respect to which the γ -coordinate in the basis Δ of each positive root is equal to 0 or 1: the first case says that the positive root belongs to $\Phi(\Delta \setminus \{\gamma\})$ and the second one says that it belongs to Φ^+ hence occurs as a weight for the A_γ -action on N_γ . We refer now to [Bou68,

Planches] and its notation. Let us investigate the root systems by a case-by-case analysis of the formulas given for the linear decompositions of the positive roots according to the given bases (formulas (II) in [loc. cit.]):

- for type A_l , any simple root can be chosen for γ since no coefficient ≥ 2 occurs in formula (II) of [loc. cit., Planche I];
- for type B_l , the root γ can be chosen to be α_1 in the notation of [loc. cit., Planche II];
- for type C_l , the root γ can be chosen to be α_l which occurs only for two kinds of positive roots, each time with coefficient 1, in formula (II) of [loc. cit., Planche III];
- for type D_l , we use the formulas in [loc. cit., p.208] instead of Planche IV (where the formulas are not correct), to see that the root γ can be chosen to be α_1 or α_l ;
- for type E_6 , the root γ can be chosen to be α_1 or α_6 , as inspection of the positive roots with some coefficient ≥ 2 shows in [loc. cit., Planche V];
- for type E_7 , the root γ can be chosen to be α_7 , as inspection of the positive roots whose support contains α_7 and with some coefficient ≥ 2 shows in [loc. cit., Planche VI].

To exclude the remaining case, it is enough to note that:

- for type E_8 , the last positive root described on the first page of [loc. cit., Planche VII] has all its coefficients ≥ 2 ;
- for type F_4 (resp. G_2), the last root in (II) of [loc. cit., Planche VIII (resp. IX)] has the same property;
- for the only non-reduced irreducible root system of rank l , namely BC_l , the positive root ε_1 has all its coefficients ≥ 2 [loc. cit., p. 222].

This finishes the determinations of the admissible simple Lie groups by means of their relative root systems. \square

As a complement, we can say that the groups that are not admissible can be listed thanks to É. Cartan's classification as stated for instance in [Hel01, Table VI, pp. 532-534]. The classical groups with non-reduced relative root system correspond to the cases where the value of $m_{2\lambda}$ is ≥ 1 in this table. To sum up, we can reformulate the previous proposition in more concrete terms:

Proposition 7.2. *The admissible simple real Lie groups are the split simple real Lie groups of classical types and of types E_6 et E_7 , the complex simple groups of classical types and of types E_6 et E_7 (all seen as real Lie groups), all orthogonal groups $SO_{p,q}(\mathbf{R})$, all special linear groups $SL_n(\mathbf{H})$ over the quaternions \mathbf{H} , the special unitary groups $SU_{n,n}(\mathbf{R})$ with $n \geq 2$, the groups $Sp_{2n,2n}(\mathbf{R}) = SU_{n,n}(\mathbf{H})$ with $n \geq 2$, the groups $SO_{4n}^*(\mathbf{R})$ with $n \geq 2$, the exceptional groups of absolute type E_6 and real rank 2 and of absolute type E_7 and real rank 3.*

8. NUMERICAL EFFICIENCY OF THE METHOD

Let us turn now to the numerical efficiency of the method, namely the computation of the actual width of the (possibly) non-vanishing strips, *i.e.* the intervals outside of which our method proves vanishing of continuous group L^p -cohomology (Theorem 6.1).

Recall that for a given admissible group G with maximal compact subgroup K , this width is equal to $D-d+2$ where D is the dimension of the symmetric space $X = G/K$ and d is the dimension of the solvable radical $A_\gamma N_\gamma$. Since $d = 1 + \dim N_\gamma$, we have:

$$d - 1 = \dim N_\gamma = \sum_{\alpha \in \Phi^+ \setminus \Phi(\Delta \setminus \{\gamma\})} m_\alpha,$$

where m_α is the multiplicity $\dim N(\alpha)$ the root space $N(\alpha)$ attached to α . Now, we have just seen that the root system of an admissible group is reduced, so we can freely use [Hum72] (in which root systems are assumed to be reduced by definition). By [loc. cit., Lemma C p. 53] there are at most two root lengths in Φ and the Weyl group acts transitively on the roots of given length, therefore since this Weyl group action lifts to the action of $N_G(A)$ on the root spaces $N(\alpha)$, we deduce that there are at most two multiplicities, say m_1 and m_2 , and with the notation $\Psi = \Phi^+ \setminus \Phi(\Delta \setminus \{\gamma\})$ the previous formula becomes:

$$d - 1 = m_1 \cdot |\{\alpha \in \Psi : m_\alpha = m_1\}| + m_2 \cdot |\{\alpha \in \Psi : m_\alpha = m_2\}|.$$

The case of split groups is easy since all multiplicities m_α are equal to 1 (hence in this case $d = 1 + |\Psi|$), and the case of Weil restrictions (*i.e.* complex groups seen as real ones) is easy too: all multiplicities are equal to 2, hence in this case $d = 1 + 2 \cdot |\Psi|$.

For the other cases, where two multiplicities may occur, we proceed as follows. For each admissible group, thanks to the previous section we make the choice (often, but not systematically, unique) of a *good* root, that is a simple root γ such that $A_\gamma N_\gamma$ is quasi-isometric to a real

hyperbolic space. Thanks to [Bou68, Planches], this gives easily $|\Psi|$ since the root system of $\Phi(\Delta \setminus \{\gamma\})$ is given by the Dynkin diagram of Φ minus the vertex of type γ and the edges emanating from it. We must then sort the roots of Ψ according to their length, which is done again thanks to the concrete descriptions of [loc. cit.]. It remains then to apply the last formula for $d - 1$ and to use the multiplicities given in [Hel01, Table VI, pp. 532-534].

The rest of this section is dedicated to computing the width $d = D - d + 2$ of the (possibly) non-vanishing strip, and to determine the minimal one when the choice of a good root γ is not unique for the group G under consideration.

The case of split groups.— Investigating the class of admissible split simple real Lie groups is the opportunity to provide a description of the roots sets Ψ of nilpotent radicals, that will be useful in the more complicated cases; we use the notation of [Bou68, Planches].

- When $G = \mathrm{SL}_{n+1}(\mathbf{R})$, the root system has type A_n , any simple root α_i ($1 \leq i \leq n$) is a good root, and we have $|\Psi| = i \times (n - i)$; all roots have the same length.
- When $G = \mathrm{SO}_{n+1,n}(\mathbf{R})$, the root system has type B_n , the only good root is α_1 and we have: $|\Psi| = |\Phi^+(B_n)| - |\Phi^+(B_{n-1})| = n^2 - (n - 1)^2 = 2n - 1$, with one root of length 1 and $(n - 1) + (n - 1) = 2n - 2$ roots of length $\sqrt{2}$.
- When $G = \mathrm{Sp}_{2n}(\mathbf{R})$, the root system has type C_n , the only good root is α_n and we have: $|\Psi| = |\Phi^+(C_n)| - |\Phi^+(A_{n-1})| = n^2 - \frac{n(n-1)}{2} = \frac{n(n+1)}{2}$, with $\frac{n(n-1)}{2}$ roots of length $\sqrt{2}$ and n roots of length 2.
- When $G = \mathrm{SO}_{n,n}(\mathbf{R})$, the root system has type D_n , the good roots are α_1 , α_{n-1} and α_n ; all roots have the same length. For the choice $\gamma = \alpha_1$, we have: $|\Psi| = |\Phi^+(D_n)| - |\Phi^+(D_{n-1})| = n(n - 1) - (n - 1)(n - 2) = 2(n - 1)$ and for $\gamma = \alpha_{n-1}$ or α_n , we have: $|\Psi| = |\Phi^+(D_n)| - |\Phi^+(A_{n-1})| = n(n - 1) - \frac{n(n-1)}{2} = \frac{n(n-1)}{2}$.
- When $G = \mathrm{E}_6^6(\mathbf{R})$, the root system has type E_6 , the good roots are α_1 and α_6 ; all roots have the same length. Whatever the choice, we have: $|\Psi| = |\Phi^+(E_6)| - |\Phi^+(D_5)| = 36 - 20 = 16$.
- When $G = \mathrm{E}_7^7(\mathbf{R})$, the root system has type E_7 , the only good root is α_7 ; all roots have the same length and we have $|\Psi| = |\Phi^+(E_7)| - |\Phi^+(E_6)| = 63 - 36 = 27$.

As already mentioned, in this case we can then compute $d = 1 + |\Psi|$ and take $w = D - d + 2$ to be minimal when several good roots are available.

The case of complex groups seen as real ones.— Then the multiplicities are then 2; in this case we have $d = 1 + 2|\Psi|$ and again we can take $w = D - d + 2$ to be minimal when several good roots are available.

The remaining admissible simple real Lie groups.— We use here the formula $d - 1 = m_1 \cdot |\{\alpha \in \Psi : m_\alpha = m_1\}| + m_2 \cdot |\{\alpha \in \Psi : m_\alpha = m_2\}|$ and the above partition of Ψ into short and long roots when the root system Ψ is not simply laced, combined with the multiplicities given in [Hel01, Table VI, pp. 532-534].

- When $G = \mathrm{SL}_{n+1}(\mathbf{H})$, the root system has type A_n , any simple root α_i ($1 \leq i \leq n$) is good, we have $|\Psi| = i \times (n - i)$ and all roots have multiplicity 4, so that $d - 1 = 4i(n - i)$ for $\gamma = \alpha_i$.
- When $G = \mathrm{SU}_{n,n}(\mathbf{R})$, the relative root system has type C_n , the good root is α_n , the $\frac{n(n-1)}{2}$ roots of length $\sqrt{2}$ have multiplicity 2 and the n roots of length 2 have multiplicity 1, so that $d - 1 = n(n - 1) + n = n^2$.
- When $G = \mathrm{Sp}_{2n,2n}(\mathbf{R})$, the relative root system has type C_n , the good root is α_n , the $\frac{n(n-1)}{2}$ roots of length $\sqrt{2}$ have multiplicity 4 and the n roots of length 2 have multiplicity 3, so that $d - 1 = 2n(n - 1) + 3n = 2n^2 + n$.
- When $G = \mathrm{SO}_{4n}^*(\mathbf{R})$, the relative root system has type C_n , the good root is α_n , the $\frac{n(n-1)}{2}$ roots of length $\sqrt{2}$ have multiplicity 4 and the n roots of length 2 have multiplicity 1, so that $d - 1 = 2n(n - 1) + n = 2n^2 - n$.
- When $G = \mathrm{E}_6^2(\mathbf{R})$, the relative root system has type A_2 , any of two simple roots is good and all roots have multiplicity 8, so that $d - 1 = 16$.
- When $G = \mathrm{E}_7^3(\mathbf{R})$, the relative root system has type C_3 , the good root is α_3 , there are 3 roots of each length $\sqrt{2}$ and 2, and since the multiplicities are 8 and 1, this gives $d - 1 = 27$.

At last, in the case of non-split orthogonal groups $\mathrm{SO}_{p,q}(\mathbf{R})$ (*i.e.*, where $q - 2 \geq p \geq 2$), we always obtain a relative root system of type B_p , with $2p - 2$ roots of length $\sqrt{2}$ and multiplicity 1 and one root of length 1 and multiplicity $q - p$, so that $d - 1 = p + q - 2$.

The next two pages contain tables which summarize the vanishing results obtained by our method. In conclusion, apart from the general family of non-split orthogonal groups (which is special since it depends

on two parameters), the infinite families of classical admissible groups provide a vanishing proportion of $\frac{1}{2}$ asymptotically in the rank. For non-split orthogonal groups, fixing the rank (*i.e.* the smallest number in the signature) and letting the bigger parameter go to infinity provide a vanishing proportion equal to the inverse of the rank.

Admissible group	Cartan type	Relative root system Φ	Good root γ	$ \Psi $	Multiplicities	$d - 1 = \dim N_\gamma$	$D = \dim X$	Asymptotic vanishing proportion $\lim_{l \rightarrow +\infty} \frac{a(l)}{D(l)}$
$SL_{l+1}(\mathbf{R})$	A I	A_l	any simple root α_i	$i(l+1-i)$	1	best choice: $\frac{i(l+1-i)}{(l+1)^2}$ (odd l), $\frac{l(l+2)}{4}$ (even l)	$\frac{l(l+3)}{2}$	$\frac{1}{2}$
$SL_{l+1}(\mathbf{H})$	A II	A_l	any simple root α_i	$i(l+1-i)$	4	$4i(l+1-i)$ best choice: $\frac{i(l+1-i)}{(l+1)^2}$ (odd l), $\frac{l(l+2)}{4}$ (even l)	$l(2l+3)$	$\frac{1}{2}$
$SU_{l,l}(\mathbf{R})$	A III	C_l	α_l	$\frac{l(l+1)}{2}$	2 and 1	l^2	$2l^2$	$\frac{1}{2}$
$Sp_{2l}(\mathbf{R})$	C I	C_l	α_l	$\frac{l(l+1)}{2}$	1	$\frac{l(l+1)}{2}$	$l(l+1)$	$\frac{1}{2}$
$Sp_{2l,2l}(\mathbf{R})$	C II	C_l	α_l	$\frac{l(l+1)}{2}$	4 and 3	$2l^2 + l$	$4l^2$	$\frac{1}{2}$
$SO_{l,l}(\mathbf{R})$	D I	C_l	α_1 , α_{l-1} and α_l	$2(l-1)$ for $\gamma = \alpha_1$, $\frac{l(l-1)}{2}$ otherwise	1	best choice: $\frac{l(l-1)}{2}$	l^2	$\frac{1}{2}$
$SO_{4l}^*(\mathbf{R})$	D III	C_l	α_l	$\frac{l(l+1)}{2}$	4 and 1	$2l^2 - l$	$2l(2l-1)$	$\frac{1}{2}$
$E_6^6(\mathbf{R})$	E I	E_6	α_1 and α_6	16	1	16	42	$\frac{5}{14}$
$E_6^2(\mathbf{R})$	E IV	A_2	α_1 and α_2	2	8	16	26	$\frac{15}{26}$
$E_7^7(\mathbf{R})$	E V	E_7	α_7	27	1	27	70	$\frac{13}{35}$
$E_7^3(\mathbf{R})$	E VII	C_3	α_3	6	8 and 1	27	54	$\frac{13}{27}$

Admissible group	Cartan type	Relative root system	Good root γ	$ \Psi $	Multiplicities	$d - 1 = \dim N_\gamma$	$D = \dim X$	Asymptotic vanishing proportion $\lim_{l \rightarrow +\infty} \frac{d(l)}{D(l)}$
$SL_{l+1}(\mathbf{C})$	–	A_l	any simple root α_i	$i(l+1-i)$	2	$2i(l+1-i)$ best choice: $\frac{(l+1)^2}{2}$ (odd l), $\frac{l(l+2)}{2}$ (even l)	$l(l+2)$	$\frac{1}{2}$
$SO_{l+1}(\mathbf{C})$	–	B_l	α_1	$2l-1$	2	$4l-2$	$l(2l+1)$	0
$Sp_{2l}(\mathbf{C})$	–	C_l	α_l	$\frac{l(l+1)}{2}$	2	$l(l+1)$	$l(2l+1)$	$\frac{1}{2}$
$SO_{l,l}(\mathbf{C})$	–	D_l	α_1 , α_{l-1} and α_l	$2(l-1)$ for $\gamma = \alpha_1$, $\frac{l(l-1)}{2}$ otherwise	2	best choice: $l(l-1)$	$l(2l-1)$	$\frac{1}{2}$
$E_6(\mathbf{C})$	–	E_6	α_1 and α_6	16	2	32	78	$\frac{31}{78}$
$E_7(\mathbf{C})$	–	E_7	α_7	27	2	54	133	$\frac{53}{133}$

9. VANISHING IN DEGREE ONE

As a further application of Theorem 6.1, we give in this section a proof of Corollary 1.5, namely vanishing of $H_{\text{ct}}^1(G, L^p(G))$ for every $p > 1$ and every admissible simple Lie group G of real rank ≥ 2 .

Let G be an admissible simple Lie group, and let $P = M \ltimes AN$ be a maximal parabolic subgroup with AN quasi-isometric to \mathbb{H}^d . One has $H_{\text{ct}}^1(G, L^p(G)) = \{0\}$ for $p \leq d - 1$, thanks to Theorem 6.1. Suppose that $p > d - 1$. Then Theorem 6.1 again implies that $H_{\text{ct}}^1(G, L^p(G))$ is linearly isomorphic to

$$H_{\text{ct}}^0\left(M, L^p\left(M, H_{\text{ct}}^1(AN, L^p(AN))\right)\right) \simeq L^p\left(M, H_{\text{ct}}^1(AN, L^p(AN))\right)^M.$$

By [Pan89] (see also [Rez08] or [BP03]) the topological vector space $H_{\text{ct}}^1(AN, L^p(AN))$ is canonically isomorphic to the following Besov space on $\mathbf{R}^{d-1} \simeq N \simeq \partial\mathbb{H}^d \setminus \{\infty\}$:

$$B_{p,p}^{(d-1)/p}(\mathbf{R}^{d-1}) := \{u : \mathbf{R}^{d-1} \rightarrow \mathbf{R} \text{ measurable ; } \|u\|_B < +\infty\} / \mathbf{R}$$

$$\text{where } \|u\|_B^p = \int_{\mathbf{R}^{d-1} \times \mathbf{R}^{d-1}} \frac{|u(x) - u(x')|^p}{\|x - x'\|^{2(d-1)}} dx dx'.$$

The M -action on $L^p\left(M, H_{\text{ct}}^1(AN, L^p(AN))\right)$ described in Proposition 5.2, takes now the following form. Denote by $(m, x) \mapsto m(x)$ the linear action of M by conjugation on N . Then for $m, y \in M$, $f : M \rightarrow B_{p,p}^{(d-1)/p}(\mathbf{R}^{d-1})$ and $x \in N \simeq \mathbf{R}^{d-1}$, one has

$$(m \cdot f)(y)(x) = f(y m)(m^{-1}(x)).$$

Therefore $H_{\text{ct}}^1(G, L^p(G))$ is linearly isomorphic to the fixed point space $L^p\left(M, B_{p,p}^{(d-1)/p}(\mathbf{R}^{d-1})\right)^M$, which in turn is isomorphic to

$$\left\{ u \in B_{p,p}^{(d-1)/p}(\mathbf{R}^{d-1}) \mid \int_M \|u \circ m\|_B^p d\mathcal{H}(m) < \infty \right\}.$$

Since M acts on \mathbf{R}^{d-1} linearly and preserves the volume, a change of variable and Fubini-Tonelli yield

$$(9.1) \quad \int_M \|u \circ m\|_B^p d\mathcal{H}(m) = \int_{\mathbf{R}^{d-1} \times \mathbf{R}^{d-1}} \frac{|u(x) - u(x')|^p}{\|x - x'\|^{2(d-1)}} \varphi(x - x') dx dx',$$

where φ is the following positive function on $\mathbf{R}^{d-1} \setminus \{0\}$:

$$(9.2) \quad \varphi(v) = \int_M \frac{d\mathcal{H}(m)}{\|m^{-1}\left(\frac{v}{\|v\|}\right)\|^{2(d-1)}}.$$

We claim that $\int_M \|u \circ m\|_B^p d\mathcal{H}(m) = +\infty$ unless u is constant *a.e.*; the corollary will follow. The proof relies on three lemmata. The first one gives a sufficient condition for u to be constant.

Lemma 9.3. *Let $u \in B_{p,p}^{(d-1)/p}(\mathbf{R}^{d-1})$ that is invariant by a 1-parameter group of translations. Then u is constant *a.e.**

Proof. Express the Besov norm of u as $\|u\|_B^p = \int_{\mathbf{R}^{d-1} \times \mathbf{R}^{d-1}} \psi \, dx dx'$ with

$$\psi(x, x') = \frac{|u(x) - u(x')|^p}{\|x - x'\|^{2(d-1)}}.$$

Suppose that u is invariant by a 1-parameter group of translations along some vector line $\mathbf{R}v_0$. Then ψ is invariant by the 1-parameter group of translations along $\mathbf{R}(v_0, v_0)$. If in addition one has $\int_{\mathbf{R}^{d-1} \times \mathbf{R}^{d-1}} \psi \, dx dx' < +\infty$, then Fubini-Tonelli implies that ψ is null *a.e.* This in turn implies that u is constant *a.e.* \square

The second lemma collects some simple properties of the function φ defined in (9.2).

Lemma 9.4. *The function φ admits the following properties:*

- (1) $\varphi(\lambda v) = \varphi(v)$ for every $\lambda \in \mathbf{R}^*$ and every $v \in \mathbf{R}^{d-1} \setminus \{0\}$.
- (2) There exists a positive constant c_0 such that $\varphi \geq c_0$.
- (3) The function $v \mapsto \frac{\varphi(v)}{\|v\|^{2(d-1)}}$ is M -invariant on $\mathbf{R}^{d-1} \setminus \{0\}$.
- (4) If $v \in \mathbf{R}^{d-1} \setminus \{0\}$, $g \in M$ and $\lambda \in \mathbf{R}$, are such that $g(v) = \lambda v$ with $|\lambda| \neq 1$, then $\varphi(v) = +\infty$.

Proof. The third item follows from the fact that \mathcal{H} is a left-invariant measure. To obtain the last item, one observes that the first and third items imply that

$$\varphi(v) = \varphi(\lambda v) = \varphi(g(v)) = \frac{\|g(v)\|^{2(d-1)}}{\|v\|^{2(d-1)}} \varphi(v) = |\lambda|^{2(d-1)} \varphi(v).$$

Thus $\varphi(v) = +\infty$. \square

We now describe the behaviour of φ in a neighborhood of an eigenvector.

Lemma 9.5. *Let $(g^t)_{t \in \mathbf{R}} \subset M$ be a non-trivial 1-parameter subgroup which acts diagonally with positive eigenvalues on \mathbf{R}^{d-1} . Let $\lambda_{\max} > 1$ and $\lambda_{\min} < 1$ be the maximum and minimum eigenvalues of g . Let $v_0 \in \mathbf{R}^{n-1}$ be an eigenvector of g for the eigenvalue λ_{\max} . There exist a*

constant $c_2 > 0$ and a neighborhood U of v_0 such that for every $v \in U$ one has

$$\varphi(v) \geq \frac{c_2}{\|v - v_0\|^\alpha}$$

with $\alpha = 2(d-1) \frac{\log \lambda_{\max}}{\log \lambda_{\max} - \log \lambda_{\min}}$.

Proof. If v is an eigenvector then $\varphi(v) = +\infty$ thanks to Lemma 9.4. Suppose now that v is not an eigenvector. Let $W \subset \mathbf{R}^{d-1}$ be a subspace such that $\mathbf{R}^{d-1} = \mathbf{R}v_0 \oplus W$ is a $\{g^t\}_{t \in \mathbf{R}}$ -invariant decomposition. We write every vector $v \in \mathbf{R}^{d-1}$ as $v = v_1 + v'$ according to this decomposition. For simplicity we will abusively denote by the same symbol $\|\cdot\|$ the standard norm on \mathbf{R}^{d-1} and the norm $\max(\|v_1\|, \|v'\|)$.

Suppose that $\|v'\| \leq \|v_1\|$. Then there exists $t \geq 0$ such that $\|g^{-t}(v')\| = \|g^{-t}(v_1)\|$. Write $w = g^{-t}(v) = g^{-t}(v_1) + g^{-t}(v') = w_1 + w'$. It follows from the previous lemma that

$$\begin{aligned} \varphi(v) = \varphi(g^t(w)) &= \frac{\|g^t(w)\|^{2(d-1)}}{\|w\|^{2(d-1)}} \varphi(w) = \frac{\|\lambda_{\max}^t w_1 + g^t(w')\|^{2(d-1)}}{\|w_1 + w'\|^{2(d-1)}} \varphi(w) \\ &\geq \frac{(\lambda_{\max}^t \|w_1\|)^{2(d-1)}}{(2\|w_1\|)^{2(d-1)}} \varphi(w) \geq c_1 \lambda_{\max}^{2t(d-1)}, \end{aligned}$$

with $c_1 = 2^{-2(d-1)} c_0$. On the other hand, we have

$$\frac{\|v_1\|}{\|v'\|} = \frac{\|\lambda_{\max}^t w_1\|}{\|g^t(w')\|} \leq \frac{\lambda_{\max}^t \|w_1\|}{\lambda_{\min}^t \|w'\|} = \frac{\lambda_{\max}^t}{\lambda_{\min}^t}.$$

Since $(\lambda_{\max})^{2(d-1)} = (\lambda_{\max}/\lambda_{\min})^\alpha$ we obtain by combining the above inequalities

$$\varphi(v) \geq c_1 \lambda_{\max}^{2t(d-1)} = c_1 \frac{\lambda_{\max}^{t\alpha}}{\lambda_{\min}^{t\alpha}} \geq c_1 \frac{\|v_1\|^\alpha}{\|v'\|^\alpha}.$$

Now there is a neighborhood of v_0 in which the ratio $\|v_1\|^\alpha/\|v'\|^\alpha$ is comparable to $1/\|v'\|^\alpha$, which in turn is larger than $1/\|v - v_0\|^\alpha$; the lemma follows. \square

Finally we complete the

Proof of Corollary 1.5. Let $u \in B_{p,p}^{(d-1)/p}(\mathbf{R}^{d-1})$ be a function such that $\int_M \|u \circ m\|_B^p d\mathcal{H}(m) < \infty$. We want to prove that u is constant *a.e.* According to Lemma 9.3, it is enough to show that there is a non-zero vector $v_0 \in \mathbf{R}^{d-1}$ such that u is constant almost everywhere along almost every line directed by v_0 . Since M is a semi-simple non-compact Lie group, it contains a 1-parameter subgroup $(g^t)_{t \in \mathbf{R}}$ which acts on \mathbf{R}^{d-1} as in Lemma 9.5. Let λ_{\max} , λ_{\min} , v_0 , c_2 , U and α be as in Lemma

9.5. By changing g to its inverse, if necessary, we can assume that $\lambda_{\max} \geq \lambda_{\min}^{-1}$; so that we have $\alpha \geq d - 1$. Since u is measurable, it is approximately continuous *a.e.* (see [Fed69, 2.9.13]). Let a, b be distinct points on a line directed by v_0 , such that u is approximately continuous at a and b . We want to show that $u(a) = u(b)$. By multiplying v_0 by a real number if necessary, we can assume that $v_0 = b - a$. For every $\varepsilon > 0$, there exists $r_\varepsilon > 0$ such that for every $0 < r \leq r_\varepsilon$ the set

$$E_r(a) := \{x \in B(a, r) \mid |u(x) - u(a)| < \varepsilon\}$$

contains at least 99% of the measure of $B(a, r)$. Similarly is defined $E_r(b)$.

Suppose by contradiction that $u(a) \neq u(b)$. Let $\varepsilon = |u(a) - u(b)|/4$ and $0 < r \leq r_\varepsilon$. Then for every $(x, x') \in E_r(a) \times E_r(b)$ one has $|u(x) - u(x')| \geq |u(a) - u(b)|/2$. Therefore by making the change of variable $v = x' - x$, we obtain with relation 9.1:

$$\begin{aligned} \int_M \|u \circ m\|_B^p d\mathcal{H}(m) &\geq \int_{E_r(a) \times E_r(b)} \frac{|u(x) - u(x')|^p}{\|x - x'\|^{2(d-1)}} \varphi(x - x') dx dx' \\ &\geq c_3 |u(a) - u(b)|^p \int_{E_r(a) \times E_r(b)} \varphi(x - x') dx dx' \\ &= c_3 |u(a) - u(b)|^p \int_{v \in \mathbf{R}^{d-1}} \text{meas}\{E_r(a) \cap (E_r(b) - v)\} \varphi(v) dv, \end{aligned}$$

where c_3 is a positive constant that depends only on $\|a - b\|$. Since $b - a = v_0$, for v close enough to v_0 , one has

$$\text{meas}\{E_r(a) \cap (E_r(b) - v)\} \geq (3/10) \text{meas}\{B(a, r)\}.$$

With Lemma 9.5 and the inequality $\alpha \geq d - 1$ one gets that for every small enough neighbourhood U of v_0 :

$$\int_M \|u \circ m\|_B^p d\mathcal{H}(m) \geq c_4 |u(a) - u(b)|^p \int_U \frac{dv}{\|v - v_0\|^{d-1}},$$

where c_4 depends only on c_2, c_3, r . Since the last integral is equal to $+\infty$, we obtain a contradiction. \square

REFERENCES

- [BFGM07] Y. Bader, A. Furman, T. Gelander, and N. Monod: Property (T) and rigidity for actions on Banach spaces. *Acta Math.*, 198(1):57–105, 2007.
- [BFS14] Y. Bader, A. Furman, and R. Sauer: Weak notions of normality and vanishing up to rank in L^2 -cohomology. *Int. Math. Res. Not. IMRN*, 12:3177–3189, 2014.
- [Bla79] Ph. Blanc: Sur la cohomologie continue des groupes localement compacts. *Ann. Sci. Ecole Norm. Sup.*, 12(2):137–168, 1979.
- [Bor85] A. Borel: The L^2 -cohomology of negatively curved Riemannian symmetric spaces. *Ann. Acad. Sci. Fenn. Ser. A I Math.*, 10:95–105, 1985.
- [Bou63] N. Bourbaki: *Intégration. Chapitres 7 à 8*. Actualités Scientifiques et Industrielles, No. 1306. Hermann, Paris, 1963.
- [Bou65] ———: *Intégration. Chapitres 1 à 4*. Actualités Scientifiques et Industrielles, No. 1175. Hermann, Paris, 1965.
- [Bou68] ———: *Groupes et algèbres de Lie. Chapitres 4 à 6*. Actualités Scientifiques et Industrielles, No. 1337. Hermann, Paris, 1968.
- [Bou07] ———: *Topologie générale. Chapitres 5 à 10*. Springer, 2007.
- [Bou16] M. Bourdon: Cohomologie ℓ_p en degrés supérieurs et dimension conforme. *Ann. Inst. Fourier*, 66:1013–1043, 2016.
- [BP03] M. Bourdon and H. Pajot: Cohomologie ℓ_p et espaces de Besov. *J. Reine Angew. Math.*, 558:85–108, 2003.
- [BW00] A. Borel and N. Wallach: *Continuous cohomology, discrete subgroups, and representations of reductive groups. Second edition*. American Mathematical Society, Providence, RI, 2000. Mathematical Surveys and Monographs, 67.
- [CdIH16] Y. Cornuier and P. de la Harpe: *Metric geometry of locally compact groups*, volume 25 of *EMS Tracts in Mathematics*. European Mathematical Society (EMS), Zürich, 2016.
- [CT11] Y. Cornuier and R. Tessera: Contracting automorphisms and L^p -cohomology in degree one. *Ark. Mat.*, 49(2):295–324, 2011.
- [Ele98] G. Elek: Coarse cohomology and ℓ^p -cohomology. *K-Theory*, 13: 1-22, 1998.
- [Fed69] H. Federer: *Geometric measure theory*, volume 153 of *Die Grundlehren der mathematischen Wissenschaften*. Springer-Verlag, New-York, 1969.
- [Gen14] L. Genton: *Scaled Alexander-Spanier Cohomology and $L^{q,p}$ Cohomology for Metric Spaces*. These no 6330. EPFL, Lausanne, 2014.
- [Gro93] M. Gromov: Asymptotic invariants of infinite groups. In *Geometric group theory, Vol. 2 (Sussex, 1991)*, volume 182 of *London Math. Soc. Lecture Note Ser.*, pages 1–295. Cambridge Univ. Press, Cambridge, 1993.
- [Gui80] A. Guichardet: *Cohomologie des groupes topologiques et des algèbres de Lie*. Textes Mathématiques. CEDIC/Fernand Nathan, Paris, 1980.
- [Hel01] S. Helgason: *Differential geometry, Lie groups, and symmetric spaces*, volume 34 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2001. Corrected reprint of the 1978 original.
- [Hum72] J. E. Humphreys: *Introduction to Lie algebras and representation theory*. Springer, 1972. Graduate Texts in Mathematics, Vol. 9.
- [Mic56] E. Michael: Continuous selections I. *Ann. of Maths.*, 63:361–382, 1956.

- [Mon01] N. Monod: *Continuous bounded cohomology of locally compact groups*, volume 1758 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 2001.
- [Mon10] ———: On the bounded cohomology of semi-simple groups, S -arithmetic groups and products. *J. Reine Angew. Math.*, 640(1):97–128, 2010.
- [Pan89] P. Pansu: Cohomologie L^p des variétés à courbure négative, cas du degré 1. *Rend. Sem. Mat. Univ. Politec. Torino*, (Special Issue):95–120, 1989. Conference on Partial Differential Equations and Geometry (Torino, 1988).
- [Pan95] ———: Cohomologie L^p : invariance sous quasiisométrie. Preprint 1995.
- [Pan99] ———: Cohomologie L^p , espaces homogènes et pincement. Preprint 1999.
- [Pan07] ———: Cohomologie L^p en degré 1 des espaces homogènes. *Potential Anal.*, 27:151–165, 2007.
- [Pan08] ———: Cohomologie L^p et pincement. *Comment. Math. Helv.*, 83(2):327–357, 2008.
- [Rez08] A. Reznikov: Analytic topology of groups, actions, strings and varieties. In *Geometry and dynamics of groups and spaces*, volume 265 of *Progr. Math.*, pages 3–93. Birkhäuser, Basel, 2008.
- [SS17] R. Sauer and M. Schrödl: Vanishing of ℓ^2 -Betti numbers of locally compact groups as an invariant of coarse equivalence. Preprint arXiv:1702.01685, available at <https://arxiv.org/abs/1702.01685>, 2017.

Laboratoire Paul Painlevé, UMR 8524 de CNRS, Université de Lille, Cité Scientifique, Bat. M2, 59655 Villeneuve d’Ascq, France.

E-mail: bourdon@math.univ-lille1.fr.

Centre de Mathématiques Laurent Schwartz, UMR 7640 de CNRS, École polytechnique, 91128 Palaiseau, France.

E-mail: bertrand.remy@polytechnique.edu.