



HAL
open science

Field Theories for Loop-Erased Random Walks

Kay Joerg Wiese, Andrei A. Fedorenko

► **To cite this version:**

Kay Joerg Wiese, Andrei A. Fedorenko. Field Theories for Loop-Erased Random Walks. Nuclear Physics B, 2019, 946, pp.114696. 10.1016/j.nuclphysb.2019.114696 . hal-01744049

HAL Id: hal-01744049

<https://hal.science/hal-01744049v1>

Submitted on 25 Oct 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

Field Theories for Loop-Erased Random Walks

Kay Jörg Wiese^a, Andrei A. Fedorenko^b

^a*Laboratoire de Physique de l'École normale supérieure, ENS, Université PSL, CNRS, Sorbonne Université, Université Paris-Diderot, Sorbonne Paris Cité, 24 rue Lhomond, 75005 Paris, France.*

^b*Univ. Lyon, ENS de Lyon, Univ. Claude Bernard, CNRS, Laboratoire de Physique, F-69342 Lyon, France*

Abstract

Self-avoiding walks (SAWs) and loop-erased random walks (LERWs) are two ensembles of random paths with numerous applications in mathematics, statistical physics and quantum field theory. While SAWs are described by the $n \rightarrow 0$ limit of ϕ^4 -theory with $O(n)$ -symmetry, LERWs have no obvious field-theoretic description. We analyse two candidates for a field theory of LERWs, and discover a connection between the corresponding and a priori unrelated theories. The first such candidate is the $O(n)$ -symmetric ϕ^4 theory at $n = -2$ whose link to LERWs was known in two dimensions due to conformal field theory. Here it is established in arbitrary dimension via a perturbation expansion in the coupling constant. The second candidate is a field theory for charge-density waves pinned by quenched disorder, whose relation to LERWs had been conjectured earlier using analogies with Abelian sandpiles. We explicitly show that both theories yield identical results to 4-loop order and give both a perturbative and a non-perturbative proof of their equivalence. This allows us to compute the fractal dimension of LERWs to order ε^5 where $\varepsilon = 4 - d$. In particular, in $d = 3$ our theory gives $z_{\text{LERW}}(d = 3) = 1.6243 \pm 0.001$, in excellent agreement with the estimate $z = 1.62400 \pm 0.00005$ of numerical simulations.

1. Introduction

Random walks (RWs) which are not allowed to self-intersect play an important role in combinatorics, statistical physics and quantum field theory. The two most prominent models are *self-avoiding walks* (SAWs) and *loop-erased random walks* (LERWs). The SAW was first introduced in polymer physics to model long polymer chains with self-repulsion due to excluded-volume effects. It is defined as the uniform measure on RW paths of a given length conditioned on having no self-intersection. Though this model is difficult to analyze rigorously, it was discovered by de Gennes [1] that its scaling behavior in d dimensions is given by the $O(n)$ symmetric ϕ^4 theory in the unusual limit of $n \rightarrow 0$. A loop-erased random walk (LERW) is defined as the trajectory of a random walk (RW) in which any loop is erased as soon as it is formed [2]. An example is shown on figure 1, where the underlying RW is drawn in red, and the LERW remaining after erasure in blue. Similar to a self-avoiding walk it has a scaling limit in all dimensions, e.g. the end-to-end distance R scales with the intrinsic length ℓ as $R \sim \ell^{1/z}$, where z is the fractal dimension [3]. It is crucial to note that while both LERWs and SAWs are non-self-intersecting, their fractal dimensions do not agree since they have a different statistics on the same set of allowed trajectories. LERWs appear in many combinatorial problems, e.g. the shortest path on a uniform spanning tree

Preprint submitted to Nucl. Phys. B

March 12, 2020




Figure 1: Trace of a LERW in blue, with the erased loops in red, on a 2-dimensional Honeycomb lattice.

is a LERW. While LERWs are non-Markovian RWs, their traces are equivalent to those of the *Laplacian Random Walk* [4, 5], which is Markovian, if one considers the whole trace as state variable. It is constructed on the lattice by solving the Laplace equation $\nabla^2\Phi(x) = 0$ with boundary conditions $\Phi(x) = 0$ on the already constructed curve, while $\Phi(x) = 1$ at the destination of the walk, either a chosen point, or infinity. The walk then advances from its tip x to a neighbouring point y , with probability proportional to $\Phi(y)$. In a variant of this model growth is allowed not only from the tip, but from any point on the already constructed object. This is known as the *dielectric breakdown model* [6], the simplest model for lightning. The same construction pertains to diffusion-limited aggregation [7]. In contrast to SAWs, LERWs have no obvious field-theoretic description. In three dimensions LERWs have been studied numerically [8, 9, 10, 11], while in two dimensions they are described by SLE with $\kappa = 2$ [12, 13], predicting a fractal dimension $z_{\text{LERW}}(d = 2) = \frac{5}{4}$. Coulomb-gas techniques link this to the 2d $O(n)$ -model at $n = -2$ [14, 15]. Below, we give perturbative arguments that this construction can be done in any dimension d via the $O(n)$ -symmetric ϕ^4 theory at $n = -2$.

Coming from a different direction, it was conjectured in [16] that the field theory of the depinning transition of charge-density waves (CDWs) pinned by disorder is a field theory for LERWs. This equivalence is based on the conjecture of Narayan and Middleton [17] that pinned CDWs can be mapped onto the Abelian sandpile model. The connection of Abelian sandpiles with uniform spanning trees, and thus with LERWs, was earlier established by Majumdar [18, 19]. Despite the lack of a proof of this equivalence, the corresponding 2-loop predictions agree with rigorous




Figure 2: The different field theories for LERWs.

bounds [16] and have been tested against numerical simulations at the upper critical dimension $d = 4$ in [10] where they correctly reproduce the leading and subleading logarithmic corrections of [16]. The depinning transition of CDWs is described by the functional RG (FRG) fixed point for random periodic systems, proposed by Narayan and Fisher [20, 21, 22], and developed in [23, 24, 25, 26, 27, 28, 29].

If true, both ϕ^4 -theory at $n = -2$ and the FRG for CDWs must agree, at least for observables related to LERWs. We show below that both the β -function and the fractal dimension of LERWs coincide. This is done using (i) graph-theoretical arguments valid at all orders in perturbation theory, (ii) non-perturbative supersymmetry techniques, and (iii) an explicit 4-loop calculation. This does not mean that the theories are identical: for example, at depinning CDWs exhibit avalanches which are seemingly absent in the ϕ^4 model. Our statement is that in the *sector* in which we can compare the two theories, they agree. This is illustrated on figure 2. It does not exclude that one of the theories can handle observables the other cannot. A classical example for this behavior is the solution of the 2d Ising model via the conformal bootstrap, as proposed by Belavin, Polyakov and Zamolodchikov (BPZ) [30]. Here a theory with three operators, the energy ε , the spin σ , and the identity $\mathbb{1}$, closes under OPE. However, other observables can be constructed from the Ising model on a lattice. Examples in case are domain walls, at criticality described by SLE. On the other hand, SLE does not (at least obviously) describe the operators of the original BPZ construction. Thus, SLE and BPZ describe different *sectors* of the same theory.

2. Mapping of LERWs onto $O(n)$ ϕ^4 -theory at $n = -2$

We now map LERWs onto the n -component ϕ^4 theory at $n = -2$. The latter is defined by the action

$$\mathcal{S}[\vec{\phi}] := \int_x \frac{1}{2} [\nabla \vec{\phi}(x)]^2 + \frac{m^2}{2} \vec{\phi}(x)^2 + \frac{g}{8} [\vec{\phi}(x)^2]^2. \quad (1)$$

One checks that, for $n = -2$, the full 2-point correlation function is given in Fourier space by the free-theory result

$$\langle \phi_i(k) \phi_j(k) \rangle = \langle \phi_i(k) \phi_j(k) \rangle_0 = \frac{\delta_{ij}}{k^2 + m^2}, \quad (2)$$

independent of g . This fact is well known perturbatively [31, 32, 33, 34]. A non-perturbative derivation is given below, by mapping onto complex fermions. Eq. (2) is the Laplace transform of


the k -dependent Green function for a random walk (RW),

$$\langle \phi_i(k) \phi_j(k) \rangle = \int_0^\infty dt e^{-m^2 t} \times e^{-k^2 t}. \quad (3)$$

Here $t \geq \ell$ is the time of the RW used to construct the LERW of length ℓ , which scales as $\ell \sim t^{z/2} \sim m^{-z}$, and z is the fractal dimension of the LERW. Let us convene that we draw the trajectory of a random walk in blue, and when it hits itself we do not erase the emerging loop, but color it in red; see figure 1 for an example. We claim that we can reconstruct these *colored* trajectories from ϕ^4 theory. To this aim, we first reformulate the theory (1) in terms of $N = n/2$ complex bosons Φ and Φ^* , with $\Phi_i(x) := \frac{1}{\sqrt{2}}[\phi_{2i-1}(x) + i\phi_{2i}(x)]$, $i = 1, \dots, N = n/2$. Its action reads

$$\mathcal{S}[\vec{\Phi}] := \int_x \nabla \vec{\Phi}^*(x) \nabla \vec{\Phi}(x) + m^2 \vec{\Phi}^*(x) \vec{\Phi}(x) + \frac{g}{2} [\vec{\Phi}^*(x) \vec{\Phi}(x)]^2. \quad (4)$$


This reformulation gives propagators a direction, making it easier to interpret them as RWs. Consider a specific path with s intersections in the path-integral representation. This conditioning introduces a small-length cutoff a [35, 36]: One may think of putting a grid of box-size a , and introduce for each box a Θ -function, which is one if the path passes through twice, and zero otherwise. It is crucial that for the self-intersections encoded in the quartic term we use the same prescription, that it squares to one, and is symmetric under exchange of the momenta. The contributions to paths with $s = 1$ self-intersections are



$$\text{LERW with self-intersection} \longrightarrow -g \text{ (loop on blue)} - gN \text{ (loop on red)} \text{ (loop on blue)} \quad (5)$$

The drawing on the l.h.s. of equation (5) is a LERW starting at x , ending in z , and passing through the segments numbered 1 to 3. Due to the crossing at y , the loop labeled 2 is erased; we draw it in red. The r.h.s of equation (5) gives all diagrams of ϕ^4 theory up to order g^s . The first term is the free-theory result, proportional to g^0 . The second term $\sim g$ cancels the first term, if one puts $g \rightarrow 1$. Here it is crucial to have the same regularization for the interaction as for the conditioning. The third term is proportional to N , due to the loop, indicated in red. Setting $N \rightarrow -1$ compensates for the subtracted second term. Thus setting $g \rightarrow 1$ and $N \rightarrow -1$, the probability to go from x to z remains unchanged as compared to the free theory. This is a necessary condition to be satisfied. Since the first two terms cancel, what remains is the last diagram, corresponding to the drawing for the trajectory of the LERW we started with.


Let us consider how this continues for $s = 2$ intersections. Once a first loop has been formed, there are two possibilities: The walk can either hit a blue or a red part of its own trace. Let us first assume it hits a blue part. Then a second loop is formed, and the mapping at $g = 1$, and $N = -1$ reads



$$\text{LERW with two self-intersections} \longrightarrow \text{diagram 1} - \text{diagram 2} - \text{diagram 3} \quad (6)$$

This is a result of multiple cancelations, which we can analyze vertex by vertex. The tricky part is what happens at point y : We can either not use any interaction, use the interaction following the lines of the original walk, or reconnect the lines of the walk to form a loop. Here we used cancelation of the first two terms, retaining the last one, which resulted in the given drawing. (Note that the drawn red trace has the statistics of Brownian motion, as the two possible interaction terms mutually cancel.)

The other possibility is to hit a red part of the trace, say at point y



Here nothing should happen, since the walk does not see the erased part of its trace. The appropriate cancelation is between “no interaction” and “reconnecting”, since the latter would result in the erased loop appearing again in blue in perturbation theory. Thus also in this case we map onto the appropriate diagram of ϕ^4 -theory. Continuing these arguments inductively allows us to prove this for any number of intersections s .

We have thus established a one-to-one correspondence between traces of LERWs and diagrams in perturbation theory. We still need an observable which is 1 when inserted into a blue part of the trace, and 0 within a red part. This can be achieved by modifying the probability to diffuse from x to z , given by the expectation of $\Phi_1^*(x)\Phi_1(z)$, to

$$\mathcal{O}(x, y, z) := \Phi_1^*(x)\Phi_1(z) [\Phi_1^*(y)\Phi_1(y) - \Phi_2^*(y)\Phi_2(y)] . \quad (8)$$

As shown in appendix [Appendix A](#) this observable can be simplified to

$$\mathcal{O}(x, y, z) = \Phi_1^*(x)\Phi_1(y)\Phi_2^*(y)\Phi_2(z) . \quad (9)$$

The fractal dimension z of a LERW is extracted from the length of the walk after erasure (blue part)

$$\frac{\left\langle \int_{y,z} \mathcal{O}(x, y, z) \right\rangle}{\left\langle \int_z \Phi_1^*(x)\Phi_1(z) \right\rangle} \equiv m^2 \left\langle \int_{y,z} \mathcal{O}(x, y, z) \right\rangle \sim m^{-z} . \quad (10)$$

We recall that the fractal dimension of SAWs is given by $z_{\text{SAW}} = 1/\nu$ where the critical exponent ν is taken in the limit of $n \rightarrow 0$. Remarkably, we can deform the measure on the RWs in a continuous way by tuning the parameter n . This interpolates between SAWs and LERWs, and more generally gives critical curves in the n -component vector model. Indeed, the operator in square brackets of (8) is related to the so-called *crossover exponent* [37, 38, 39]. It is defined as (see e.g. [38], Eq. (8))

$$\phi_c = \frac{2 + \gamma_{\phi\phi} - \eta}{2 + \gamma_{\phi^2} - \eta} \equiv (2 + \gamma_{\phi\phi} - \eta) \nu . \quad (11)$$

Here $\gamma_{\phi\phi}$ is the RG function for the operator $\Phi_1\Phi_2^*$, and η is the anomalous exponent for the dimension of the fields. For the LERWs considered here, $\eta = 0$ and $\nu = 1/2$. More generally [37, 38, 40],

$$z = d_f = 2 + \gamma_{\phi\phi} - \eta \quad (12)$$

is the fractal dimension of a curve in the critical $O(n)$ symmetric ϕ^4 theory. Let us discuss what this means for LERWs and SAWs. First, for $n = -2$, the 2-point function between two points x and z in the ϕ^4 theory is the sum over all LERWs from x to z , weighted by the chemical potential m^2 conjugate to its construction time t . It equals the free propagator from x to z , since coloring loops in red or erasing them does not change the propagator. On the other hand, if we apply our picture to SAWs, red loops carry a factor of $N = 0$, and only configurations without self-intersections survive. In this case the ratio (10) equals 1, and the fractal dimension can be inferred from the 2-point function alone, i.e. $\phi_c(n = 0) = 1$ and $d_f = 1/\nu(n = 0) = z_{\text{SAW}}$. Thus, the 2-point function of ϕ^4 theory with $n = 0$ equals the sum over all SAWs connecting the two points.

Let us now turn our attention to $d = 2$: There it is known that SLE_κ for $0 \leq \kappa \leq 4$ is a simple curve. RWs generated by the 2-dimensional $O(n)$ -symmetric ϕ^4 theory with $n = -2 \cos(4\pi/\kappa)$ are non-self-intersecting for $-2 \leq n \leq 2$, corresponding to $2 \leq \kappa \leq 4$. For $\kappa > 4$ the curves are self-intersecting. For $|n| > 2$ the field theory becomes massive, and the connection breaks down. As we now know that $n = -2$ and $n = 0$ correspond to LERWs and SAWs in arbitrary dimension d , we conjecture that there is a family of non-selfintersecting random walks in arbitrary dimension d , parameterized by n , such that their measure can be continuously deformed from that of LERWs at $n = -2$ to that of SAWs at $n = 0$, and beyond to $n > 0$. Interestingly, the barrier of $|n| = 2$ in two dimensions is seemingly absent in higher dimensions. This may be understood from the fact that the return probability of a random walk to the origin decreases with increasing dimension.

3. ϕ^4 -theory at $N = -1$, and fermions

Up to now, we worked with $N = -1$ families of complex bosons. We show below that instead one can consider the limit of $N_f \rightarrow 1$ in a theory with N_f complex fermions, or more generally with N_b bosons and N_f fermions, where $N = N_b - N_f$. Among others, this provides a non-perturbative proof that the propagator at $N = -1$ is independent of g .

The correspondence is based on the observation that N_b -component bosons $\vec{\Phi}$ carry a factor of N_b per loop, while N_f -component fermions $\vec{\Psi}$ yield a factor of $-N_f$. On a more formal level, this can be inferred from the path integrals for both theories. Setting $\mathbb{H}_0 := -\nabla^2 + \tau(x)$ then

$$\mathcal{Z}_0^b = \int \mathcal{D}[\vec{\Phi}^*] \mathcal{D}[\vec{\Phi}] e^{-\int_x \vec{\Phi}^*(x) \mathbb{H}_0 \vec{\Phi}(x)} = e^{-N_b \text{tr} \ln(\mathbb{H}_0)}, \quad (13)$$

$$\mathcal{Z}_0^f = \int \mathcal{D}[\vec{\Psi}^*] \mathcal{D}[\vec{\Psi}] e^{-\int_x \vec{\Psi}^*(x) \mathbb{H}_0 \vec{\Psi}(x)} = e^{N_f \text{tr} \ln(\mathbb{H}_0)}. \quad (14)$$

The bosonic correlation function is given by

$$\langle \Phi_i(x) \Phi_j^*(y) \rangle = \delta_{ij} (\mathbb{H}_0^{-1})_{x,y} = \frac{1}{\mathcal{Z}_0^b} \int \mathcal{D}[\vec{\Phi}^*] \mathcal{D}[\vec{\Phi}] \Phi_i(x) \Phi_j^*(y) e^{-\int_x \vec{\Phi}^*(x) \mathbb{H}_0 \vec{\Phi}(x)}. \quad (15)$$

For fermions an equivalent expression holds, and $\langle \Psi_i(x) \Psi_j^*(y) \rangle = \langle \Phi_i(x) \Phi_j^*(y) \rangle$. Setting $\tau(x) = m^2 + i\rho(x)$, a Hubbard Stratonovich transformation allows us to decouple the quartic interaction in a theory of N_b bosons Φ_i and N_f fermions Ψ_i ,

$$e^{-\frac{g}{2} \int_x [\vec{\Phi}^*(x) \vec{\Phi}(x) + \vec{\Psi}^*(x) \vec{\Psi}(x)]^2} = \int \mathcal{D}[\rho] e^{-\frac{1}{2g} \int_x \rho(x)^2 - i\rho(x) [\vec{\Phi}^*(x) \vec{\Phi}(x) + \vec{\Psi}^*(x) \vec{\Psi}(x)]}. \quad (16)$$

As a consequence, a system of N_b bosons and N_f fermions with the interaction (16) has partition function

$$\begin{aligned} \mathcal{Z}^{b+f} &= \int \mathcal{D}[\vec{\Phi}^*] \mathcal{D}[\vec{\Phi}] \mathcal{D}[\vec{\Psi}^*] \mathcal{D}[\vec{\Psi}] e^{-\int_x \vec{\Phi}^*(x)(-\nabla^2+m^2)\vec{\Phi}(x)+\vec{\Psi}^*(x)(-\nabla^2+m^2)\vec{\Psi}(x)} \\ &\times e^{-\frac{g}{2} \int_x [\vec{\Phi}^*(x)\vec{\Phi}(x)+\vec{\Psi}^*(x)\vec{\Psi}(x)]^2} = \int \mathcal{D}[\rho] e^{(N_f-N_b) \text{tr} \ln(\mathbb{H}_0) - \frac{1}{2g} \int_x \rho(x)^2}. \end{aligned} \quad (17)$$

More generally, the correlations $\langle \Phi_1(y)\Phi_1^*(x) \rangle$ in complex N -component Φ^4 theory can be calculated from a theory with N_b bosons and N_f fermions, where $N = N_b - N_f$. For $N_b = 0$, $N_f = 1$ one gets $N = -1$, and the interaction is $[\vec{\Psi}^*(x)\vec{\Psi}(x)]^2$. It vanishes due to the squares of Grassmann variables: This theory of *complex fermions*,

$$\mathcal{S}_{N_b=0}^{N_f=1}[\Psi^*, \Psi] = \int_x \nabla \Psi^*(x) \nabla \Psi(x) + m^2 \Psi^*(x) \Psi(x), \quad (18)$$

is a free theory. It provides a non-perturbative proof that correlation functions of complex Φ^4 -theory at $N = -1$ ($n = -2$ real fields) are equivalent to those of the free theory. In $d = 2$ this is also known from lattice models [14]. However, it does not yield the renormalization of the coupling g at $N = -1$. To obtain the latter, one has to study $N_f \neq 1$, and take the limit of $N_f \rightarrow 1$ at the end. Or one uses one family of complex bosons $N_b = 1$ and two families of complex fermions $N_f = 2$, a formulation onto which we will map CDWs at depinning later.

Finally, care has to be taken in identifying observables in both theories: While the 2-point functions of bosonic fields are symmetric under their exchange, those of the fermionic theory are antisymmetric. As a consequence $\langle \phi_1 \phi_1 \rangle \neq 0$, whereas $\langle \psi_1 \psi_1 \rangle = 0$.

4. Equivalence between ϕ^4 -theory at $N = -1$ and CDWs at depinning

Charge-density waves are ground states of solids, where the charge density is varying spatially, with a period set to 1. Coupling these elastic objects to quenched disorder leads after averaging over disorder to the dynamic field theory [42, 43, 20, 21, 22, 44]

$$\mathcal{S}^{\text{CDW}} = \int_{x,t} \tilde{u}(x,t) (\partial_t - \nabla^2 + m^2) u(x,t) - \frac{1}{2} \int_{x,t,t'} \tilde{u}(x,t) \tilde{u}(x,t') \Delta(u(x,t) - u(x,t')). \quad (19)$$

The function $\Delta(u)$ is an even function with period 1. Its renormalization can be studied using functional RG (FRG) [20, 22, 21, 23, 24, 25, 26, 27]. The analysis of the FRG flow for the function $\Delta(u)$ shows that the fixed point has the form


$$\Delta(u) = \Delta(0) - \frac{g}{2} u(1-u). \quad (20)$$

In the absence of higher-order terms in u , the RG flow closes in the space of polynomials of degree 2, and for the quadratic term one is left with the renormalization of a single coupling constant g .

They both vanish, since the first is an acausal loop, thus zero, and the second has only one connected component, thus does not contribute to the renormalization of g . Now consider


(28)

This diagram contains an acausal loop, thus does not contribute to the SAW theory (22) where it vanishes due to a factor of $N = 0$. The diagram however has *children*; together they are


(29)

The modified lines are in red. We first remark that none of them restricts the time-difference between the left and right-most vertices, and they all contribute to the effective coupling. Their coefficients are $0 \times 1 - 1 - 1 + 1 = -1$. Integrating over times, the result is the same as in ϕ^4 -theory at $N = -1$, graphically represented as


(30)

Noting the pairwise cancelations in loops of the form (29), this can be interpreted as the *missing* contribution of the first (acausal) diagram. To summarize: We showed that at 1-loop order the action (21) has the same effective coupling as the action (4), diagram by diagram (after time integration). These considerations can be formalized to higher orders, and we checked them explicitly up to 4 loops.

The theory (21) has a second renormalization, namely of friction, or time, which shows up in a renormalization of $\int_{x,t} \tilde{u}(x,t) \dot{u}(x,t)$. The standard route to study this is to write down all diagrams constructed from (21), in which one field \tilde{u} and one field u remain [26]. Due to the structure of the action, the latter has the form $u(x,t) - u(x,t')$ and can be expanded as $\dot{u}(x,t)(t - t')$. The additional time difference, when appearing together with a response function, acts by inserting an additional point into the latter, as can be seen from the definition (23), and the relation

$$tR(k, t) = \int_0^t dt' R(k, t') R(k, t - t'). \quad (31)$$

Following this strategy, we checked that up to 4-loop order all diagrams appearing after time-integration are equivalent to those encountered in expectations of \mathcal{O} , defined in Eq. (8). Graphically, this is proven by first realizing that one can alternatively study the renormalization of friction by considering insertions of $\int_{x,t} \tilde{u}(x,t) \dot{u}(x,t)$. Doing this, the time derivative on \dot{u} can be passed through a closed string of response functions to either the earliest time in the diagram, and will then act on the remaining uncontracted field u , or it will end on a vertex with no further u field, and thus

vanish. The same argument can be done by moving the time derivative to the field \tilde{u} . These operations restrict the class of diagrams. Graphically, inserting $\int_{x,t} \tilde{u}(x,t) \dot{u}(x,t)$ corresponds again to inserting a point into diagrams correcting expectations of $\tilde{u}(x,t)u(x',t')$. The final step of the proof is to realize that this is equivalent to insertions of the crossover operator $\Phi_1(y)\Phi_2^*(y)$ in the theory (4).

Finally note that the absence of a renormalization of $-\nabla^2 + m^2$ in Eqs. (19) and (21) implied by the statistical tilt symmetry is equivalent to the absence of a renormalization of the theory (18).

5. A non-perturbative proof for the equivalence of ϕ^4 -theory at $N = -1$ and CDWs

The method introduced in [45, 46] allows one to write the disorder average of any observable $\mathcal{O}[u_i]$ as

$$\overline{\mathcal{O}[u_i]} = \int \prod_{a=1}^r \mathcal{D}[\tilde{u}_a] \mathcal{D}[u_a] \mathcal{D}[\bar{\psi}_a] \mathcal{D}[\psi_a] \mathcal{O}[u_i] \times \overline{\exp \left[- \int_x \tilde{u}_a(x) \frac{\delta \mathcal{H}[u_a]}{\delta u_a(x)} + \bar{\psi}_a(x) \frac{\delta^2 \mathcal{H}[u_a]}{\delta u_a(x) \delta u_a(y)} \psi_a(y) \right]}. \quad (32)$$

The integral over \tilde{u}_a ensures that u_a is at a minimum. $\bar{\psi}_a$ and ψ_a are fermionic degrees of freedom (Grassmann variables), which compensate for the functional determinant appearing in the integration over u , yielding a partition function $\mathcal{Z} = 1$. Averaging over disorder gives an effective action [47]

$$\begin{aligned} \mathcal{S}[\tilde{u}_a, u_a, \bar{\psi}_a, \psi_a] &= \sum_a \int_x \tilde{u}_a(x) (-\nabla^2 + m^2) u_a(x) + \bar{\psi}_a(x) (-\nabla^2 + m^2) \psi_a(x) \\ &- \sum_{a,b} \int_x \left[\frac{1}{2} \tilde{u}_a(x) \Delta(u_a(x) - u_b(x)) \tilde{u}_b(x) - \tilde{u}_a(x) \Delta'(u_a(x) - u_b(x)) \bar{\psi}_b(x) \psi_b(x) \right. \\ &\quad \left. - \frac{1}{2} \bar{\psi}_a(x) \psi_a(x) \Delta''(u_a(x) - u_b(x)) \bar{\psi}_b(x) \psi_b(x) \right]. \end{aligned} \quad (33)$$

The function $\Delta(u)$ is the same as in Eq. (19). Note that we allow for an arbitrary number of replicas r . In the work [45] the focus was on $r = 1$, which does not allow to extract the second cumulant of the disorder, i.e. its correlations. To do the latter, one needs at least $r = 2$ copies, to which we specify now. We introduce center-of-mass coordinates,

$$u_1(x) = u(x) + \frac{1}{2} \phi(x), \quad u_2(x) = u(x) - \frac{1}{2} \phi(x), \quad (34)$$

$$\tilde{u}_1(x) = \frac{1}{2} \tilde{u}(x) + \tilde{\phi}(x), \quad \tilde{u}_2(x) = \frac{1}{2} \tilde{u}(x) - \tilde{\phi}(x). \quad (35)$$

The action (33) can then be written as

$$\begin{aligned}
\mathcal{S} = & \int_x \tilde{\phi}(x)(-\nabla^2 + m^2)\phi(x) + \tilde{u}(x)(-\nabla^2 + m^2)u(x) + \sum_{a=1}^2 \bar{\psi}_a(x)(-\nabla^2 + m^2)\psi_a(x) \\
& + \tilde{\phi}(x)^2 \left[\Delta(\phi(x)) - \Delta(0) \right] - \frac{1}{4} \tilde{u}(x)^2 \left[\Delta(\phi(x)) + \Delta(0) \right] \\
& + \frac{1}{2} \tilde{u}(x) \Delta'(\phi(x)) \left[\bar{\psi}_2(x)\psi_2(x) - \bar{\psi}_1(x)\psi_1(x) \right] \\
& + \tilde{\phi}(x) \Delta'(\phi(x)) \left[\bar{\psi}_2(x)\psi_2(x) + \bar{\psi}_1(x)\psi_1(x) \right] \\
& + \bar{\psi}_2(x)\psi_2(x)\bar{\psi}_1(x)\psi_1(x)\Delta''(\phi(x)) .
\end{aligned} \tag{36}$$

As in the derivation of the action (21) replacing $\Delta(u) \rightarrow \frac{g}{2}u^2$, the action (36) takes the form

$$\begin{aligned}
\mathcal{S} = & \int_x \tilde{\phi}(x)(-\nabla^2 + m^2)\phi(x) + \tilde{u}(x)(-\nabla^2 + m^2)u(x) + \sum_{a=1}^2 \bar{\psi}_a(x)(-\nabla^2 + m^2)\psi_a(x) \\
& + \frac{g}{2} \tilde{u}(x) \phi(x) \left[\bar{\psi}_2(x)\psi_2(x) - \bar{\psi}_1(x)\psi_1(x) \right] - \frac{g}{8} \tilde{u}(x)^2 \phi(x)^2 \\
& + \frac{g}{2} \left[\tilde{\phi}(x)\phi(x) + \bar{\psi}_1(x)\psi_1(x) + \bar{\psi}_2(x)\psi_2(x) \right]^2 .
\end{aligned} \tag{37}$$

Note that only $\tilde{u}(x)$, but not the center-of-mass position $u(x)$ appear in the interaction. While $u(x)$ may have non-trivial expectations, it does not contribute to the renormalization of g , and the latter can be obtained by dropping the third line of Eq. (37). What remains is a ϕ^4 -type theory as in Eq. (17), with one complex boson, and two complex fermions. It can equivalently be viewed as complex ϕ^4 -theory at $N = -1$, or real ϕ^4 -theory at $n = -2$.

What is yet missing is information about the exponent z . One can use the operator \mathcal{O} defined in Eqs. (8) or (9), replacing Φ_i by ψ_i , and Φ_i^* by $\bar{\psi}_i$. Another possibility is to introduce time, adding a time argument to all fields, and replacing $-\nabla^2 + m^2$ by $\partial_t - \nabla^2 + m^2$. The interaction part, i.e. the last line of Eq. (37), then becomes bilocal in time, i.e. the time integral appears inside the square bracket. The tricky part is to ensure that time-integrated vacuum bubbles retain their static expectations. This can be done by specifying an initial condition, once again adding the action (37) where all fields are evaluated at some initial time t_0 . This implies that

$$R(k; t', t) = \left\langle \phi(-k, t') \tilde{\phi}(k, t) \right\rangle = \left\langle \psi_i(-k, t') \bar{\psi}_i(k, t) \right\rangle = \Theta(t' - t) e^{-(k^2 + m^2)(t' - t)} + \frac{\delta_{t, t_0} \delta_{t', t_0}}{k^2 + m^2} . \tag{38}$$

The δ -functions are to be understood s.t.

$$\int_t R(k_1, t, t) \dots R(k_n, t, t) = \frac{1}{(k_1^2 + m^2) \dots (k_n^2 + m^2)} . \tag{39}$$

We explicitly checked to 3-loop order that terms proportional to ∂_t receive the same renormalization as at depinning. Furthermore we can analyze the renormalization of $\tilde{\phi}(x, t) \partial_t \phi(x, t)$ as an insertion. Contributing diagrams carry two external fields, one $\tilde{\phi}$, and one ϕ . Passing the time

derivative $\partial_t \phi(x, t)$ of the insertion to this external field, what remains is the insertion of a single point in the line connecting the external fields $\tilde{\phi}$ and ϕ , but no contribution from insertions into loops. After integration over times, this is equivalent to the insertion of \mathcal{O} defined above in Eqs. (8) or (9).

6. Fractal dimension of LERWs at 5-loop order

We generated all diagrams entering into $\mathcal{O}(x, y, z)$ at 5-loop order, and into the renormalization of the coupling constant at 4-loop order, supplemented by the diagrams of [34] at 5-loop order. At 3-loop order we obtain the fractal dimension z of LERWs using the massive diagrams of Ref. [28]. To 4- and 5-loop order, we use diagrams in a massless minimal subtraction scheme obtained in [33, 34]. The result reproduces at 4-loop order the one given for the crossover exponent in Ref. [38], setting there $n = -2$. This yields for the fractal dimension z of LERWs in dimension $d = 4 - \varepsilon$, equivalent to the dynamical exponent of CDWs at depinning,

$$\begin{aligned} z &= 2 - \frac{\varepsilon}{3} - \frac{\varepsilon^2}{9} + \left[\frac{2\zeta(3)}{9} - \frac{1}{18} \right] \varepsilon^3 - \left[\frac{70\zeta(5)}{81} - \frac{\zeta(4)}{6} - \frac{17\zeta(3)}{162} + \frac{7}{324} \right] \varepsilon^4 \\ &\quad + \left[\frac{121\zeta(3)}{972} - \frac{8\zeta(3)^2}{81} + \frac{17\zeta(4)}{216} - \frac{103\zeta(5)}{243} - \frac{175\zeta(6)}{162} + \frac{833\zeta(7)}{216} - \frac{17}{1944} \right] \varepsilon^5 + \mathcal{O}(\varepsilon^6) \\ &= 2 - \frac{\varepsilon}{3} - \frac{\varepsilon^2}{9} + 0.211568\varepsilon^3 - 0.611186\varepsilon^4 + 2.43354\varepsilon^5 + \mathcal{O}(\varepsilon^6). \end{aligned} \quad (40)$$

Using Borel resummation of (40), where to improve the precision we have included the 6-loop term yields [41]

$$z(d=2) = 1.244 \pm 0.01, \quad (41)$$

$$z(d=3) = 1.6243 \pm 0.001. \quad (42)$$

This can be compared to the exact value $z(d=2) = 5/4$ [12, 13] and the most precise numerical simulations to date by David Wilson [11],

$$z(d=3) = 1.62400 \pm 0.00005. \quad (43)$$

7. Summary and Perspectives

We presented evidence that both ϕ^4 theory with $O(n)$ -symmetry at $n = -2$, as well as the field theory for CDWs at depinning describe loop-erased random walks. We sketched a proof of this equivalence, based on a diagrammatic expansion, and gave an algebraic proof for the latter. All claims were explicitly checked to 4-loop order. This equivalence gives a strong support for the Narayan-Middleton conjecture [17] that CDWs pinned by disorder can be mapped onto the Abelian sandpile model, and thus for the conjecture of [16]. Remarkably, while CDWs at depinning map onto Abelian sandpiles, disordered elastic interfaces at depinning map onto Manna

sandpiles [48, 49]. Thus each main universality class at depinning corresponds to a specific sandpile model.

The result is surprising, since a simple ϕ^4 -type theory contains all necessary information to obtain the FRG fixed point of CDWs, a disordered system. It does not directly yield the renormalized 2-point function, or the physics of avalanches. As sketched on Fig. 2, our understanding is that the different field theories are not equivalent, but when restricted to the same *physical sector* make the same predictions. This opens a path to eventually tackle other systems which necessitate FRG via a simpler scalar field theory.

Finally, the mapping of ϕ^4 -theory at $n = -2$ onto LERWs was done at a microscopic coupling $g = 1$. Changing the latter to $p < 1$ can be interpreted as a random walk where loops are erased with probability p . Since the RG fixed point is reached for any $0 < p < 1$, we conjecture that these *partially* loop-erased random walks are in the same universality class as LERWs. We propose taking p close to 1 to measure the correction-to-scaling exponent ω precisely. While its ε expansion is known to 6-loop order [34], it is only slowly converging, and we estimate $\omega = 0.83 \pm 0.01$.

8. Acknowledgements

It is a pleasure to thank E. Brézin, J. Cardy, F. David, K. Gawedzki, P. Grassberger, J. Jacobsen, M.V. Kompaniets, A. Nahum, S. Rychkov, D. Wilson and J. Zinn-Justin for valuable discussions.

Appendix A. Rewriting the operator \mathcal{O}

We had defined in Eq. (8)

$$\mathcal{O}(x, y, z) := \phi_1^*(x) \left[\phi_1^*(y) \phi_1(y) - \phi_2^*(y) \phi_2(y) \right] \phi_1(z). \quad (\text{A.1})$$

Due to the symmetry of the action under the mapping $\phi_2 \rightarrow -\phi_2$, (ibid. for ϕ_2^*) with all other components fixed, it follows that

$$0 = \langle \phi_1^*(x) \phi_2(z) \rangle = \left\langle \phi_1^*(x) \left[\phi_1^*(y) \phi_1(y) - \phi_2^*(y) \phi_2(y) \right] \phi_2(z) \right\rangle. \quad (\text{A.2})$$

Symmetry upon exchange of components 1 and 2 further yields

$$\left\langle \phi_1^*(x) \left[\phi_1^*(y) \phi_1(y) - \phi_2^*(y) \phi_2(y) \right] \phi_1(z) \right\rangle = - \left\langle \phi_2^*(x) \left[\phi_1^*(y) \phi_1(y) - \phi_2^*(y) \phi_2(y) \right] \phi_2(z) \right\rangle. \quad (\text{A.3})$$

Using Eqs. (A.2) and (A.3) we deduce that

$$\begin{aligned} & \left\langle \phi_1^*(x) \left[\phi_1^*(y) \phi_1(y) - \phi_2^*(y) \phi_2(y) \right] \phi_1(z) \right\rangle \\ &= \frac{1}{2} \left\langle [\phi_1^*(x) - \phi_2^*(x)] \left[\phi_1^*(y) \phi_1(y) - \phi_2^*(y) \phi_2(y) \right] [\phi_1(z) + \phi_2(z)] \right\rangle. \end{aligned} \quad (\text{A.4})$$

Define (ibid. for the complex conjugate fields)

$$\phi_+(x) = \frac{1}{\sqrt{2}} [\phi_1(x) + \phi_2(x)], \quad \phi_-(x) = \frac{1}{\sqrt{2}} [\phi_1(x) - \phi_2(x)]. \quad (\text{A.5})$$

Then expectations of the observable $\mathcal{O}(x, y, z)$ can be rewritten as

$$\langle \mathcal{O}(x, y, z) \rangle = \langle \phi_-^*(x) \phi_+(y) \phi_-^*(y) \phi_+(z) \rangle . \quad (\text{A.6})$$

Eq. (A.5) is a rotation. Relabeling $\phi_- \rightarrow \phi_1$ and $\phi_+ \rightarrow \phi_2$, this proves the equivalence of the definitions (8) and (9).

Appendix B. Renormalization group functions and critical exponents

The β -function is defined as

$$\beta(g) := -m \partial_m g(g_0) \Big|_{g_0=g_0(g)} . \quad (\text{B.1})$$

It has a non-trivial fixed point at g^* , s.t. $\beta(g^*) = 0$. The slope of the β -function at $g = g^*$ yields the correction-to-scaling exponent ω ,

$$\begin{aligned} \omega := -\beta'(g^*) &= \varepsilon - \frac{2\varepsilon^2}{3} + \left(\frac{4\zeta(3)}{3} + \frac{5}{9} \right) \varepsilon^3 + \left(\frac{\pi^4}{90} - \frac{4\zeta(3)}{3} - \frac{70\zeta(5)}{9} - \frac{31}{54} \right) \varepsilon^4 \\ &+ \left(\frac{19\zeta(3)^2}{27} + \frac{163\zeta(3)}{54} + \frac{479\zeta(5)}{81} + \frac{833\zeta(7)}{18} - \frac{5\pi^6}{486} - \frac{\pi^4}{90} + \frac{23}{36} \right) \varepsilon^5 \\ &+ \mathcal{O}(\varepsilon^6) \end{aligned} \quad (\text{B.2})$$

This result agrees with the literature on the $O(n)$ -model [33, 34] and for CDWs [29] (where $\omega := \beta'(g^*)$). To obtain the exponent z , we use that

$$\left\langle \int_z \phi_1^*(x) \phi_1(z) \right\rangle = \frac{1}{m^2} , \quad \frac{\left\langle \int_{y,z} \mathcal{O}(x, y, z) \right\rangle}{\left\langle \int_z \phi_1^*(x) \phi_1(z) \right\rangle} \equiv m^2 \left\langle \int_x \mathcal{O}(x, y, z) \right\rangle \sim m^{-z} . \quad (\text{B.3})$$

With the help of Eqs. (C.1) and (D.1), this yields

$$z = 2 + \frac{\partial \gamma_{\phi\phi}}{\partial g} \beta(g) \Big|_{g=g^*} . \quad (\text{B.4})$$

The result is given in Eq. (40) in the main text. Note that when using the diagrams of Ref. [34], the mass m gets replaced by a momentum scale. Due to universality, the critical exponents are independent of the scheme.

Appendix C. The effective coupling at 5-loop order

$$\begin{aligned}
g = & g_0 - 3g_0^2 \left[\text{diagram} \right] + 3g_0^3 \left[\text{diagram} + 4 \text{diagram} \right] \\
& - 3g_0^4 \left[\text{diagram} + 4 \text{diagram} + 2 \text{diagram} + 16 \text{diagram} + 4 \text{diagram} + 4 \text{diagram} + 4 \text{diagram} \right] \\
& + 3g_0^5 \left[1 \text{diagram} + 4 \text{diagram} + 2 \text{diagram} + 8 \text{diagram} + 4 \text{diagram} + 16 \text{diagram} + 4 \text{diagram} + 8 \text{diagram} \right. \\
& \quad + 8 \text{diagram} + 16 \text{diagram} + 8 \text{diagram} + 24 \text{diagram} + 32 \text{diagram} + 32 \text{diagram} + 16 \text{diagram} \\
& \quad \left. + 48 \text{diagram} + 8 \text{diagram} + 32 \text{diagram} + 4 \text{diagram} + 8 \text{diagram} + 4 \text{diagram} + 28 \text{diagram} \right] \\
& - 3g_0^6 \left[8 \text{diagram} + 32 \text{diagram} + 16 \text{diagram} + 112 \text{diagram} + 96 \text{diagram} + 4 \text{diagram} + 4 \text{diagram} \right. \\
& \quad + 16 \text{diagram} + 24 \text{diagram} + 32 \text{diagram} + 8 \text{diagram} + 32 \text{diagram} + 4 \text{diagram} + 8 \text{diagram} \\
& \quad + 1 \text{diagram} + 2 \text{diagram} + 4 \text{diagram} + 8 \text{diagram} + 16 \text{diagram} + 32 \text{diagram} + 32 \text{diagram} \\
& \quad + 16 \text{diagram} + 32 \text{diagram} + 36 \text{diagram} + 56 \text{diagram} + 96 \text{diagram} + 2 \text{diagram} + 4 \text{diagram} \\
& \quad + 8 \text{diagram} + 16 \text{diagram} + 32 \text{diagram} + 4 \text{diagram} + 8 \text{diagram} + 32 \text{diagram} + 32 \text{diagram} \\
& \quad + 32 \text{diagram} + 16 \text{diagram} + 16 \text{diagram} + 16 \text{diagram} + 96 \text{diagram} + 32 \text{diagram} + 16 \text{diagram} \\
& \quad + 32 \text{diagram} + 48 \text{diagram} + 64 \text{diagram} + 32 \text{diagram} + 32 \text{diagram} + 32 \text{diagram} + 32 \text{diagram} \\
& \quad + 16 \text{diagram} + 4 \text{diagram} + 4 \text{diagram} + 4 \text{diagram} + 4 \text{diagram} + 8 \text{diagram} + 32 \text{diagram} \\
& \quad + 16 \text{diagram} + 4 \text{diagram} + 8 \text{diagram} + 64 \text{diagram} + 32 \text{diagram} + 48 \text{diagram} + 48 \text{diagram} \\
& \quad + 96 \text{diagram} + 224 \text{diagram} + 32 \text{diagram} + 48 \text{diagram} + 16 \text{diagram} + 32 \text{diagram} + 272 \text{diagram} \\
& \quad + 48 \text{diagram} + 8 \text{diagram} + 48 \text{diagram} + 32 \text{diagram} + 24 \text{diagram} + 112 \text{diagram} + 96 \text{diagram} \\
& \quad + 64 \text{diagram} + 12 \text{diagram} + 8 \text{diagram} + 32 \text{diagram} + 64 \text{diagram} + 32 \text{diagram} + 32 \text{diagram} \\
& \quad + 16 \text{diagram} + 8 \text{diagram} + 64 \text{diagram} + 32 \text{diagram} + 8 \text{diagram} + 16 \text{diagram} + 48 \text{diagram} \\
& \quad \left. + 32 \text{diagram} + 96 \text{diagram} + 64 \text{diagram} + 8 \text{diagram} + 28 \text{diagram} + 56 \text{diagram} + 4 \text{diagram} \right] \\
& + \dots
\end{aligned} \tag{C.1}$$

Note that at 5-loop order the vertices are sitting on a circle, and crossings inside this circle are not vertices. There are $(2n + 1)!!$ terms at n -loop order.

Appendix D. The observable \mathcal{O} at 5-loop order

Amputating the external points of the observable \mathcal{O} , and integrating over all but one point, we find

$$\begin{aligned}
e^{\gamma_{\phi\phi}(g_0)} &= m^4 \left\langle \int_{x,z} \mathcal{O}(x, y, z) \right\rangle \\
&= 1 - g_0 \text{ (circle) } + g_0^2 \left[\text{circle with 2 vertices} + 2 \text{ (circle with 3 vertices)} \right] \\
&\quad - g_0^3 \left[\text{circle with 4 vertices} + 2 \text{ (circle with 5 vertices)} + 2 \text{ (circle with 6 vertices)} + 8 \text{ (circle with 7 vertices)} + 2 \text{ (circle with 8 vertices)} \right] \\
&\quad + g_0^4 \left[\text{circle with 8 vertices} + 2 \text{ (circle with 9 vertices)} + 4 \text{ (circle with 10 vertices)} + 8 \text{ (circle with 11 vertices)} + 2 \text{ (circle with 12 vertices)} + 4 \text{ (circle with 13 vertices)} + 8 \text{ (circle with 14 vertices)} + 16 \text{ (circle with 15 vertices)} \right. \\
&\quad + 16 \text{ (circle with 16 vertices)} + 8 \text{ (circle with 17 vertices)} + 2 \text{ (circle with 18 vertices)} + 8 \text{ (circle with 19 vertices)} + 2 \text{ (circle with 20 vertices)} + 4 \text{ (circle with 21 vertices)} + 4 \text{ (circle with 22 vertices)} \\
&\quad \left. + 4 \text{ (circle with 23 vertices)} + 12 \text{ (circle with 24 vertices)} \right] \\
&\quad - g_0^5 \left[\text{circle with 24 vertices} + 2 \text{ (circle with 25 vertices)} + 4 \text{ (circle with 26 vertices)} + 8 \text{ (circle with 27 vertices)} + 2 \text{ (circle with 28 vertices)} + 4 \text{ (circle with 29 vertices)} + 4 \text{ (circle with 30 vertices)} + 16 \text{ (circle with 31 vertices)} \right. \\
&\quad + 4 \text{ (circle with 32 vertices)} + 2 \text{ (circle with 33 vertices)} + 8 \text{ (circle with 34 vertices)} + 4 \text{ (circle with 35 vertices)} + 16 \text{ (circle with 36 vertices)} + 16 \text{ (circle with 37 vertices)} + 16 \text{ (circle with 38 vertices)} \\
&\quad + 8 \text{ (circle with 39 vertices)} + 8 \text{ (circle with 40 vertices)} + 2 \text{ (circle with 41 vertices)} + 16 \text{ (circle with 42 vertices)} + 4 \text{ (circle with 43 vertices)} + 16 \text{ (circle with 44 vertices)} + 4 \text{ (circle with 45 vertices)} \\
&\quad + 8 \text{ (circle with 46 vertices)} + 16 \text{ (circle with 47 vertices)} + 16 \text{ (circle with 48 vertices)} + 8 \text{ (circle with 49 vertices)} + 4 \text{ (circle with 50 vertices)} + 8 \text{ (circle with 51 vertices)} + 8 \text{ (circle with 52 vertices)} \\
&\quad + 4 \text{ (circle with 53 vertices)} + 12 \text{ (circle with 54 vertices)} + 48 \text{ (circle with 55 vertices)} + 12 \text{ (circle with 56 vertices)} + 32 \text{ (circle with 57 vertices)} + 32 \text{ (circle with 58 vertices)} + 32 \text{ (circle with 59 vertices)} \\
&\quad + 16 \text{ (circle with 60 vertices)} + 32 \text{ (circle with 61 vertices)} + 16 \text{ (circle with 62 vertices)} + 32 \text{ (circle with 63 vertices)} + 16 \text{ (circle with 64 vertices)} + 8 \text{ (circle with 65 vertices)} + 8 \text{ (circle with 66 vertices)} \\
&\quad + 16 \text{ (circle with 67 vertices)} + 48 \text{ (circle with 68 vertices)} + 24 \text{ (circle with 69 vertices)} + 24 \text{ (circle with 70 vertices)} + 48 \text{ (circle with 71 vertices)} + 16 \text{ (circle with 72 vertices)} + 4 \text{ (circle with 73 vertices)} \\
&\quad + 4 \text{ (circle with 74 vertices)} + 16 \text{ (circle with 75 vertices)} + 16 \text{ (circle with 76 vertices)} + 16 \text{ (circle with 77 vertices)} + 16 \text{ (circle with 78 vertices)} + 16 \text{ (circle with 79 vertices)} + 16 \text{ (circle with 80 vertices)} + 16 \text{ (circle with 81 vertices)} \\
&\quad + 8 \text{ (circle with 82 vertices)} + 2 \text{ (circle with 83 vertices)} + 2 \text{ (circle with 84 vertices)} + 4 \text{ (circle with 85 vertices)} + 16 \text{ (circle with 86 vertices)} + 4 \text{ (circle with 87 vertices)} + 8 \text{ (circle with 88 vertices)} \\
&\quad \left. + 4 \text{ (circle with 89 vertices)} + 28 \text{ (circle with 90 vertices)} + 56 \text{ (circle with 91 vertices)} \right] + \dots \tag{D.1}
\end{aligned}$$

There are $(2n - 1)!!$ diagrams at n -loop order. At 5-loop order the vertices are sitting on a circle, and crossings inside this circle are not vertices.

References

- [1] P.-G. De Gennes, *Exponents for the excluded volume problem as derived by the Wilson method*, [Phys. Lett. A **38** \(1972\) 339–340](#).
- [2] G.F. Lawler, *A self-avoiding random walk*, [Duke Math. J. **47** \(1980\) 655–693](#).
- [3] G. Kozma, *The scaling limit of loop-erased random walk in three dimensions*, [Acta Mathematica **199** \(2007\) 29–152](#).
- [4] J. W. Lyklema, C. Evertsz and L. Pietronero, *The Laplacian random walk*, [EPL **2** \(1986\) 77](#).
- [5] Gregory F. Lawler, *The Laplacian-b random walk and the Schramm-Loewner evolution*, [Illinois J. Math. **50** \(2006\) 701–746](#).
- [6] L. Niemeyer, L. Pietronero and H. J. Wiesmann, *Fractal dimension of dielectric breakdown*, [Phys. Rev. Lett. **52** \(1984\) 1033–1036](#).
- [7] T. A. Witten and L. M. Sander, *Diffusion-limited aggregation, a kinetic critical phenomenon*, [Phys. Rev. Lett. **47** \(1981\) 1400–1403](#).
- [8] A.J. Guttmann and R.J. Bursill, *Critical exponent for the loop erased self-avoiding walk by monte carlo methods*, [J. Stat. Phys. **59** \(1990\) 1–9](#).
- [9] H. Agrawal and D. Dhar, *Distribution of sizes of erased loops of loop-erased random walks in two and three dimensions*, [Phys. Rev. E **63** \(2001\) 056115](#).
- [10] P. Grassberger, *Scaling of loop-erased walks in 2 to 4 dimensions*, [J. Stat. Phys. **136** \(2009\) 399–404](#), [arXiv:0905.3440](#).
- [11] D.B. Wilson, *Dimension of the loop-erased random walk in three dimensions*, [Phys. Rev. E **82** \(2010\) 062102](#), [arXiv:1008.1147](#).
- [12] O. Schramm, *Scaling limits of loop-erased random walks and uniform spanning trees*, [Israel J. Math. **118** \(2000\) 221–288](#), [arXiv:math/9904022](#).
- [13] G.F. Lawler, O. Schramm and W. Werner, *Conformal invariance of planar loop-erased random walks and uniform spanning trees*, [Ann. Probab. **32** \(2004\) 939–995](#), [arXiv:math/0112234](#).
- [14] B. Nienhuis, *Exact critical point and critical exponents of $O(n)$ models in two dimensions*, [Phys. Rev. Lett. **49** \(1982\) 1062–1065](#).
- [15] B. Duplantier, *Loop-erased self-avoiding walks in two dimensions: exact critical exponents and winding numbers*, [Physica A **191** \(1992\) 516–522](#).
- [16] A.A. Fedorenko, P. Le Doussal and K.J. Wiese, *Field theory conjecture for loop-erased random walks*, [J. Stat. Phys. **133** \(2008\) 805–812](#), [arXiv:0803.2357](#).
- [17] O. Narayan and A.A. Middleton, *Avalanches and the renormalization-group for pinned charge-density waves*, [Phys. Rev. B **49** \(1994\) 244–256](#).
- [18] S.N. Majumdar, *Exact fractal dimension of the loop-erased self-avoiding walk in two dimensions*, [Phys. Rev. Lett. **68** \(1992\) 2329–2331](#).
- [19] D. Dhar, *Theoretical studies of self-organized criticality*, [Physica A **369** \(2006\) 29–70](#).
- [20] D.S. Fisher, *Sliding charge-density waves as a dynamical critical phenomena*, [Phys. Rev. B **31** \(1985\) 1396–1427](#).
- [21] O. Narayan and D.S. Fisher, *Critical behavior of sliding charge-density waves in 4-epsilon dimensions*, [Phys. Rev. B **46** \(1992\) 11520–49](#).
- [22] O. Narayan and D.S. Fisher, *Dynamics of sliding charge-density waves in 4-epsilon dimensions*, [Phys. Rev. Lett. **68** \(1992\) 3615–18](#).
- [23] H. Leschhorn, T. Nattermann, S. Stepanow and L.-H. Tang, *Driven interface depinning in a disordered medium*, [Annalen der Physik **509** \(1997\) 1–34](#), [arXiv:cond-mat/9603114](#).
- [24] T. Nattermann, S. Stepanow, L.-H. Tang and H. Leschhorn, *Dynamics of interface depinning in a disordered medium*, [J. Phys. II \(France\) **2** \(1992\) 1483–8](#).
- [25] P. Le Doussal, K.J. Wiese and P. Chauve, *Functional renormalization group and the field theory of disordered elastic systems*, [Phys. Rev. E **69** \(2004\) 026112](#), [cond-mat/0304614](#).
- [26] P. Le Doussal, K.J. Wiese and P. Chauve, *2-loop functional renormalization group analysis of the depinning transition*, [Phys. Rev. B **66** \(2002\) 174201](#), [cond-mat/0205108](#).

- [27] P. Chauve, P. Le Doussal and K.J. Wiese, *Renormalization of pinned elastic systems: How does it work beyond one loop?*, *Phys. Rev. Lett.* **86** (2001) 1785–1788, [cond-mat/0006056](#).
- [28] K.J. Wiese, C. Husemann and P. Le Doussal, *Field theory of disordered elastic interfaces at 3-loop order: The β -function*, *Nucl. Phys. B* (2018) in press, [arXiv:1801.08483](#).
- [29] C. Husemann and K.J. Wiese, *Field theory of disordered elastic interfaces to 3-loop order: Results*, *Nucl. Phys. B* (2017) in press, [arXiv:1707.09802](#).
- [30] A.A. Belavin, A.M. Polyakov and A.B. Zamolodchikov, *Infinite conformal symmetry in two-dimensional quantum field theory*, *Nucl. Phys. B* **241** (1984) 333–380.
- [31] J. Zinn-Justin, *Phase Transitions and Renormalization Group*, Oxford University Press, Oxford, 2007.
- [32] H. Kleinert and V. Schulte-Frohlinde, *Critical Properties of ϕ^4 -Theories*, World Scientific Publishing, 2001.
- [33] H. Kleinert, J. Neu, N. Schulte-Frohlinde and S.A. Larin, *Five-loop renormalization group functions of $O(N)$ -symmetric ϕ^4 -theory and ε -expansion of critical exponents up to ε^5* , *Phys. Lett. B* **272** (1991) 39–44, [arXiv:hep-th/9503230](#).
- [34] M.V. Kompaniets and E. Panzer, *Minimally subtracted six-loop renormalization of $O(n)$ -symmetric ϕ^4 theory and critical exponents*, *Phys. Rev. D* **96** (2017) 036016, [arXiv:1705.06483](#).
- [35] J. des Cloizeaux and G. Jannink, *Polymers in Solution, Their Modelling and Structure*, Clarendon Press, Oxford, 1990.
- [36] J. des Cloizeaux, *Polymers in solutions: Principles and applications of a direct renormalization method*, *J. de Physique* **42** (1981) 635–652.
- [37] D.J. Amit and V. Martin-Mayor, *Field Theory, the Renormalization Group, and Critical Phenomena*, World Scientific, Singapore, 3rd edition, 1984.
- [38] J.E. Kirkham, *Calculation of crossover exponent from Heisenberg to Ising behaviour using the fourth-order ε expansion*, *J. Phys. A* **14** (1981) L437–L442.
- [39] H. Shimada and S. Hikami, *Fractal dimensions of self-avoiding walks and Ising high-temperature graphs in 3d conformal bootstrap*, *J. Stat. Phys.* **165** (2016) 1006–1035, [arXiv:1509.04039](#).
- [40] J. Kiskis, R. Narayanan, and P. Vranas, *The Hausdorff dimension of random walks and the correlation length critical exponent in Euclidean field theory*, *J. Stat. Phys.* **73** (1993) 765–774, [arXiv:hep-th/9303055](#).
- [41] M.V. Kompaniets and K.J. Wiese, *Crossover exponent for $O(n)$ -symmetric ϕ^4 -theory, and fractal dimension of loop-erased random walks at 6-loop order*, unpublished.
- [42] H. Fukuyama and P.A. Lee, *Dynamics of the charge-density wave. I. Impurity pinning in a single chain*, *Phys. Rev. B* **17** (1978) 535.
- [43] P.A. Lee and T.M. Rice, *Electric-field depinning of charge-density waves*, *Phys. Rev. B* **19** (1979) 3970–3980.
- [44] T. Nattermann and S. Scheidl, *Vortex-glass phases in type-II superconductors*, *Advances in Physics* **49** (2000) 607–704, [cond-mat/0003052](#).
- [45] G. Parisi and N. Surlas, *Random magnetic fields, supersymmetry, and negative dimensions*, *Phys. Rev. Lett.* **43** (1979) 744–5.
- [46] G. Parisi and N. Surlas, *Supersymmetric field theories and stochastic differential equations*, *Nucl. Phys. B* **B206** (1982) 321–32.
- [47] K.J. Wiese, *Supersymmetry breaking in disordered systems and relation to functional renormalization and replica-symmetry breaking*, *J. Phys.: Condens. Matter.* **17** (2005) S1889–S1898, [cond-mat/0411656](#).
- [48] P. Le Doussal and K.J. Wiese, *An exact mapping of the stochastic field theory for Manna sandpiles to interfaces in random media*, *Phys. Rev. Lett.* **114** (2014) 110601, [arXiv:1410.1930](#).
- [49] K.J. Wiese, *Coherent-state path integral versus coarse-grained effective stochastic equation of motion: From reaction diffusion to stochastic sandpiles*, *Phys. Rev. E* **93** (2016) 042117, [arXiv:1501.06514](#).