



HAL
open science

Scaled lattices of closed p -adic semi-algebraic sets

Luck Darnière

► **To cite this version:**

| Luck Darnière. Scaled lattices of closed p -adic semi-algebraic sets. 2018. hal-01756160v1

HAL Id: hal-01756160

<https://hal.science/hal-01756160v1>

Preprint submitted on 31 Mar 2018 (v1), last revised 28 Oct 2018 (v3)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Scaled lattices of closed p -adic semi-algebraic sets*

Luck Darnière[†]

March 31, 2018

Abstract

Let p be prime number, K be a p -adically closed field, $X \subseteq K^m$ a semi-algebraic set defined over K and $L(X)$ the lattice of semi-algebraic subsets of X which are closed in X . We prove that the complete theory of $L(X)$ eliminates the quantifiers in a certain language \mathcal{L}_{ASC} , the \mathcal{L}_{ASC} -structure on $L(X)$ being an extension by definition of the lattice structure. Moreover it is decidable, contrary to what happens over a real closed field. We classify these \mathcal{L}_{ASC} -structures up to elementary equivalence, and get in particular that the complete theory of $L(K^m)$ only depends on m , not on K nor even on p . As an application we obtain a classification of semi-algebraic sets over countable p -adically closed fields up to so-called “pre-algebraic” homeomorphisms.

1 Introduction

Our knowledge of the geometry of semi-algebraic sets over p -adic fields, that is the field \mathbf{Q}_p of p -adic numbers and its finite extensions, has grown intensively in the past decades. After Macintyre’s quantifier elimination theorem [Mac76], Denef’s cell decomposition theorem [Den84] lead to develop a good dimension theory for p -adic semi-algebraic sets [SvdD88], and to classify them up to semi-algebraic bijection [Clu01]. There has been recently a new step forward with the triangulation of p -adic semi-algebraic sets [Dar17]. All this has been generalised to p -adically closed fields, that is fields which are elementarily equivalent to a p -adic one. In the present paper we use a consequence of the p -adic triangulation to prove the next result (Theorem 10.2).

Theorem A *Let K be a p -adically closed field, $X \subseteq K^m$ a semi-algebraic set and $L(X)$ the lattice of all semi-algebraic subsets of X which are closed in X . Then the complete theory of $L(X)$ is decidable, and eliminates the quantifier in \mathcal{L}_{ASC} (a certain expansion by definition of the lattice language, see below).*

Remark 1.1 In particular $L(\mathbf{Q}_p^m)$ is decidable for every positive integer m . This contrasts with the known results on archimedean local fields: $L(\mathbf{R}^m)$ and $L(\mathbf{C}^m)$ are undecidable for every $m \geq 2$ by [Grz51] (see [Tre17] for more on this topic). Note that the decomposition of semi-algebraic sets over \mathbf{R} in finitely many semi-algebraic connected components is crucial in the proof of Grzegorzczuk’s undecidability result, a property which disappears in the p -adic case.

The above result follows from a fine-grain study of the model-theory of various classes of geometric lattices such as the following ones.

*This paper is a major revision of [Dar06], with extended results.

Keywords: model-theory, p -adic, scaled lattice, Heyting algebra, quantifier elimination, decidability.

MSC classes: 03C10, 06D20, 06D99.

[†]Département de mathématiques, Université d’Angers, 2 Boulevard Lavoisier, 49045 Angers cedex 01 (France)

Example 1.2 Let \mathcal{K} be an o -minimal or P -minimal expansion¹ of a field K . For every definable sets $A, B \subseteq X \subseteq K^m$ let $A - B = \overline{A \setminus B} \cap A$ where the overline stands for the topological closure w.r.t. the topology induced on K^m by the order or the p -valuation of K . For every $a \in A$, the **local dimension** of A at a is the maximum of the dimensions of the definable neighborhoods of a in A , and A is called **pure dimensional** if it has the same local dimension at every point. For every non-negative integer i let

$$C^i(A) = \overline{\{a \in A \mid \dim(A, a) = i\}} \cap A.$$

This is a definable subset of A called the **i -pure component** of A . We call $L_{\text{def}}(X)$ the lattice of all the definable subsets of X which are closed in X , enriched with the above functions “ $-$ ” and C^i for every i .

Let $\mathcal{L}_{\text{lat}} = \{\mathbf{0}, \mathbf{1}, \vee, \wedge\}$ be the language of lattices, and $\mathcal{L}_{\text{SC}} = \mathcal{L}_{\text{lat}} \cup \{-, (C^i)_{i \geq 0}\}$ be its expansion the above function symbols. Finally let $\text{SC}_{\text{def}}(\mathcal{K}, d)$ be the class of these \mathcal{L}_{SC} -structures $L_{\text{def}}(X)$ for all the sets X of dimension at most d definable over \mathcal{K} . A similar construction can be done over a pure field K , with the Zariski topology on K^m . We let $\text{SC}_{\text{Zar}}(K, d)$ denote the corresponding class of \mathcal{L}_{SC} -structures.

Theorem B *Given any non-negative integer d , the universal theories of $\text{SC}_{\text{def}}(\mathcal{K}, d)$ (resp. $\text{SC}_{\text{Zar}}(K, d)$) in the language \mathcal{L}_{SC} are the same for every o -minimal or P -minimal expansion \mathcal{K} of a field K (resp. for every infinite field K).*

In order to prove this we give in Section 2 an explicit list of universal axioms for a theory T_d in \mathcal{L}_{SC} , the models of which we call d -subscaled lattices. All the examples given above are d -scaled lattices, a natural subclass of d -subscaled lattices (the class of d -scaled lattices is elementary but not universal). After some technical preliminaries in Section 3 we prove in Section 4 that every finitely generated d -subscaled lattice is finite. Combining this with a linear representation for finite d -subscaled lattices (Proposition 5.3) and with the model-theoretic compactness theorem, we then prove in Section 5 the theory of d -subscaled lattices is exactly the universal theory of $\text{SC}_{\text{def}}(\mathcal{K}, d)$ and of $\text{SC}_{\text{Zar}}(K, d)$ (Theorem 5.3).

A detailed study of the minimal finitely generated extensions of finite d -subscaled lattices, achieved in Section 6, leads us in Section 7 to our third main result (Theorem 7.3 and Corollary 7.5).

Theorem C *For every non-negative integer d , the theory of d -subscaled lattices admits a model-completion \bar{T}_d which is finitely axiomatizable and \aleph_0 -categorical. Moreover \bar{T}_d has finitely many prime models, hence it is decidable as well as all its completions.*

Remark 1.3 Since 0-subscaled lattices are exactly non-trivial boolean algebras (in which the \mathcal{L}_{SC} -structure and the boolean structure are quantifier-free bi-definable) the above model-completion result for subscaled lattices is a generalisation to arbitrary finite dimension d of the well known theorem on the model-completion of boolean algebras.

Remark 1.4 Our d -subscaled lattices form (after language reduction) an elementary class of co-Heyting algebras, closely related to the “slices” studied in [Hos67], [Ono71] and others (see [DJ11] for a more precise account of this relation). So this paper might also be considered as a contribution to the model-theory of co-Heyting algebras with at most d slices.

¹The reader unfamiliar with o -minimality, and P -minimality may consult [vdD98], [HM97], or restrict to the special case where $K = \mathbf{R}$ or K is a finite extension of \mathbf{Q}_p , and consider that “definable” is a synonym for “semi-algebraic”.

The axiomatization of \bar{T}_d given in Section 7 consists in a pair of axioms expressing a ‘‘Catenarity’’ and a ‘‘Splitting’’ property which both have a natural topological and geometric meaning. In particular the Splitting Property expresses a very strong form of disconnectedness, which implies that the models of \bar{T}_d are atomless. So we develop in Sections 8 and 9 a variant of this quantifier elimination result in a language $\mathcal{L}_{\text{ASC}} = \mathcal{L}_{\text{SC}} \cup \{\text{At}_k\}_{k \geq 1}$, where each At_k is a unary predicate symbol, to be interpreted as the set of elements which are the join of exactly k atoms. The model-completion \bar{T}_d^{At} that we obtain is axiomatized by the Catenarity property and a little restriction of the Splitting Property which preserves the atoms. This theory \bar{T}_d^{At} has \aleph_0 prime models which can easily be classified in terms of the prime models of \bar{T}_d , from which it follows that it is decidable as well as all its completions (Theorem 9.4).

In the initial version of this paper [Dar04] we conjectured that $L_{\text{def}}(\mathbf{Q}_p^d)$ might be a natural model of \bar{T}_d^{At} . This intuition proved to be crucial in the proof of the triangulation of semi-algebraic sets over a p -adically closed field [Dar17]. Conversely it follows from this triangulation that $L_{\text{def}}(X)$ is indeed a model of \bar{T}_d^{At} , for every semi-algebraic² set $X \subseteq K^m$ of dimension $\leq d$, from which Theorem A follows in the last Section 10.

The prime \mathcal{L}_{ASC} -substructure of $L_{\text{def}}(X)$ (which is generated by the empty set) is finite. We expect it to be an essential invariant for the classification of semi-algebraic sets over p -adically closed fields up to semi-algebraic homeomorphisms. Such a classification is far from being achieved, but the next result exemplifies the role that this invariant can play in a weaker classification, up to ‘‘pre-algebraic’’ homeomorphisms (this is defined in Section 10).

Theorem D *Let K, F be countable p -adically closed fields, and $X \subseteq K^m, Y \subseteq F^n$ be two semi-algebraic sets. Let $L^0(X)$ and $L^0(Y)$ be the prime \mathcal{L}_{ASC} -substructures of $L(X)$ and $L(Y)$ respectively. Then X and Y are pre-algebraically homeomorphic if and only if $L^0(X)$ and $L^0(Y)$ are \mathcal{L}_{ASC} -isomorphic. In particular, any two semi-algebraic sets over K and F with the same pure dimension $d \geq 1$ are pre-algebraically homeomorphic.*

2 Notation and definitions

\mathbf{N} denotes the set of non-negative integers, and $\mathbf{N}^* = \mathbf{N} \setminus \{0\}$. If \mathcal{N} is an unbounded non-empty subset of \mathbf{N} (resp. the empty subset) we set $\max \mathcal{N} = +\infty$ (resp. $\max \mathcal{N} = -\infty$). The symbols \subseteq and \subset denote respectively the inclusion and the strict inclusion. The logical connectives ‘or’, ‘and’ and their iterated forms will be denoted by \vee, \wedge, \bigvee and \bigwedge respectively.

2.1 Lattices and dimension

In this paper a **lattice** is a partially ordered set in which every finite subset has a greatest lower element and a least greater element. This applies in particular to the empty subset, hence our lattices must have a least and a greatest element. $\mathcal{L}_{\text{lat}} = \{\mathbf{0}, \mathbf{1}, \vee, \wedge\}$ is the language of lattices, each symbol having its obvious meaning. As usually $b \leq a$ is an abbreviation for $a \vee b = a$ and similarly for $b < a, b \geq a$ and $b > a$. Iterated \vee and \wedge operations are denoted by $\bigvee_{i \in I} a_i$ and $\bigwedge_{i \in I} a_i$ respectively. If the index set I is empty then $\bigvee_{i \in I} a_i = \mathbf{0}$ and $\bigwedge_{i \in I} a_i = \mathbf{1}$. Given a subset S of a lattice L , the **upper semi-lattice** generated in L by S is the set of finite joins of elements of L . We will make intensive use of the following definable relation:

$$b \ll a \iff \forall c (c < a \Rightarrow b \vee c < a)$$

This is a strict order on $L \setminus \{\mathbf{0}\}$ (but not on L because $\mathbf{0} \ll a$ for every a including $a = \mathbf{0}$).

²A generalization to definable sets over more general P -minimal fields, if possible, has still to be done.

The **spectrum** of a lattice L is the set $\text{Spec}(L)$ of all prime filters of L , endowed with the so-called Zariski topology, defined by taking as a basis of closed sets all the sets

$$P(a) = \{\mathfrak{p} \in \text{Spec}(L) \mid a \in \mathfrak{p}\}$$

for a ranging over L . Stone-Priestley's duality asserts that if L is distributive (which is always the case in this paper) the map $a \mapsto P(a)$ is an isomorphism between L and the lattice of closed subsets S of $\text{Spec}(L)$ such that the complement of S in $\text{Spec}(L)$ is compact.

We call a lattice **noetherian** if it is isomorphic to the lattice of closed sets of a noetherian topological space. By Stone-Priestley's duality a lattice L is noetherian if and only if its spectrum is a noetherian topological space. In such a lattice every filter is principal and every element a writes uniquely as the join of its (finitely many) **\vee -irreducible components**, which are the maximal elements in the set of \vee -irreducible³ elements of L smaller than a . We denote by $\mathcal{I}(L)$ the set of all \vee -irreducible elements of L .

We define the **dimension of an element a in a lattice L** , denoted $\dim_L a$, as the least upper bound (in $\mathbf{N} \cup \{-\infty, +\infty\}$) of the set of non-negative integers n such that

$$\exists \mathfrak{p}_0 \subset \cdots \subset \mathfrak{p}_n \in P(a).$$

This is nothing but the ordinary topological or Krull dimension (defined by chains of irreducible closed subsets) of the spectral space $P(a)$. By construction $\dim_L \mathbf{0} = -\infty$ and $\dim_L a \vee b = \max(\dim_L a, \dim_L b)$. The index L is necessary since $\dim_L a$ is not preserved by \mathcal{L}_{lat} -embeddings, but we will omit it whenever the ambient lattice is clear from the context. We let the **dimension of L** be the dimension of $\mathbf{1}_L$ in L .

In all the geometric examples given the introduction, a set A is pure dimensional if and only if $\dim U = \dim A$ for every non-empty definable subset U of A which is open in A . This motivates the next definition: given an integer k we say that an element a of a distributive lattice L is **k -pure in L** if and only if

$$\forall b \in L (a - b \neq \mathbf{0} \Rightarrow \dim_L a - b = k).$$

Then either $a = \mathbf{0}$ or $\dim_L a = k$. In the latter case we say that a **has pure dimension k in L** .

Fact 2.2 *If $L = L(X)$ is any of the lattices of the introduction, then for any $A \in L(X)$, $\dim_{L(X)} A$ is exactly the usual dimension of A as a semi-algebraic set over K , and A is pure dimensional in $L(X)$ if and only if it so in the geometric sense.*

The second assertion in Fact 2.2 follows from the first, which itself follows from Fact 2.4 below.

2.3 Co-Heyting algebras

We let $\mathcal{L}_{\text{TC}} = \mathcal{L}_{\text{lat}} \cup \{-\}$ with $'-\'$ a binary function symbol. An \mathcal{L}_{TC} -structure L is a **co-Heyting algebra** if its \mathcal{L}_{lat} -reduct is a lattice and if every element b has in L a **topological complement** relatively to every element, denoted $a - b$. By definition $a - b$ is the least element c such that $a \leq b \vee c$. Equivalently $P(a - b)$ is the topological closure of the relative complement $P(a) \setminus P(b)$, so the name of $a - b$. Reversing the order of a co-Heyting Heyting algebra L gives a Heyting algebra L^* , with $b \rightarrow a$ in L^* corresponding to $a - b$ in L , and every co-Heyting algebra is of this form. From the theory of Heyting algebras (see for example [Joh82]) we know that every

³An element c of a lattice L is \vee -irreducible if it is *non-zero* and if $a \vee b = c$ implies $a = c$ or $b = c$.

co-Heyting algebra is distributive and that the class of all co-Heyting algebras is a variety (in the sense of universal algebra). Observe that in co-Heyting algebras the \ll relation is quantifier-free definable since

$$b \ll a \iff b \leq a - b.$$

So it will be preserved by \mathcal{L}_{TC} -embeddings. On the other hand the dimension will not be preserved in general by \mathcal{L}_{TC} -embeddings.

We will use the following rules, the proof of which are elementary exercises (using either Stone-Priestley's duality or corresponding properties of Heyting algebras).

$$\mathbf{TC}_1 : a = (a \wedge b) \vee (a - b).$$

In particular if a is \vee -irreducible then $b < a \implies b \ll a$.

$$\mathbf{TC}_2 : (a_1 \vee a_2) - b = (a_1 - b) \vee (a_2 - b).$$

$$\mathbf{TC}_3 : (a - b) - b = a - b.$$

In particular $(a - b) \wedge b \ll a - b \leq a$.

$$\mathbf{TC}_4 : \text{More generally } a - (b_1 \vee b_2) = (a - b_1) - b_2.$$

So if $a - b_1 = a$ then $a - (b_1 \vee b_2) = a - b_2$.

Fact 2.4 (Theorem 3.8 in [DJ11]) *For every element a in a co-Heyting algebra L , $\dim_L a$ is the least upper bound of the set of positive integers n such that there exists $a_0, \dots, a_n \in L$ such that*

$$\mathbf{0} \neq a_0 \ll a_1 \ll \dots \ll a_n \leq a.$$

There is a well-established duality between finite (co-)Heyting algebras and finite posets, from which we will pick up Fact 2.5 below. We first need a notation. Given an element x in a poset I and a subset X of I let

$$x^\downarrow = \{y \in I \mid y \leq x\} \qquad X^\downarrow = \bigcup_{x \in I} x^\downarrow.$$

The dual notation x^\uparrow and X^\uparrow is defined accordingly. The family $\mathcal{D}^\downarrow(I)$ of decreasing subsets of I (that is the sets $X \subseteq I$ such that $X^\downarrow = X$) are the closed sets of a topology on I , hence a co-Heyting algebra with respect to the following operations.

$$X \vee Y = X \cup Y \qquad X \wedge Y = X \cap Y \qquad X - Y = (X \setminus Y)^\downarrow$$

The \vee -irreducible elements of $\mathcal{D}^\downarrow(I)$ are precisely the sets x^\downarrow for $x \in I$.

Fact 2.5 (Proposition 2.7 in [DJ]) *Let L be a finite co-Heyting algebra and \mathcal{I} an ordered set. Assume that there is a surjective increasing map $\pi : \mathcal{I} \rightarrow \mathcal{I}(L)$ such that $\pi(x^\uparrow) \subseteq \pi(x)^\uparrow$ for every $x \in \mathcal{I}(L)$. Then there exists an \mathcal{L}_{TC} -embedding φ of L into $\mathcal{D}^\downarrow(\mathcal{I})$ such that⁴ $\pi(\varphi(a)) = a^\downarrow \cap \mathcal{I}(L)$ for every $a \in L$.*

⁴Note that the compositum $\pi \circ \varphi$ is not defined. In this proposition $\varphi(a)$ is a decreasing subset of \mathcal{I} and $\pi(\varphi(a)) = \{\pi(\xi) \mid \xi \in \varphi(a)\}$.

2.6 (Sub)scaled lattices.

Recall that $\mathcal{L}_{\text{SC}} = \mathcal{L}_{\text{lat}} \cup \{-, C^i\}_{i \in \mathbf{N}} = \mathcal{L}_{\text{TC}} \cup \{C^i\}_{i \in \mathbf{N}}$ where $\{C^i\}_{i \in \mathbf{N}}$ is a family of new unary function symbols. With the examples of the introduction in mind, define the **sc-dimension** of a non-zero element a of an \mathcal{L}_{SC} -structure L as

$$\text{sc-dim } a = \min\{k \in \mathbf{N} / a = \bigvee_{0 \leq i \leq k} C^i(a)\}.$$

Of course this is defined only if $\text{sc-dim } a = \bigvee_{0 \leq i \leq k} C^i(a)$ for some k . If it is not defined we let $\text{sc-dim } a = +\infty$, and by convention $\text{sc-dim } \mathbf{0} = -\infty$. The **sc-dimension of L** , denoted $\text{sc-dim}(L)$, is the sc-dimension of $\mathbf{1}_L$. In general the dimension of an element in a co-Heyting algebra is not preserved by \mathcal{L}_{TC} -embeddings. At the contrary the sc-dimension of an element is obviously preserved by \mathcal{L}_{SC} -embeddings, and this is the “raison d’être” of this structure.

A **d -subscaled lattice** is an \mathcal{L}_{SC} -structure whose \mathcal{L}_{TC} -reduct is a co-Heyting algebra and which satisfies the following list of axioms:

$$\mathbf{SC}_1^d : \bigvee_{0 \leq i \leq d} C^i(a) = a \quad \text{and} \quad \forall i > d, C^i(a) = \mathbf{0}.$$

$$\mathbf{SC}_2 : \forall I \subseteq \{0, \dots, d\}, \forall k:$$

$$C^k\left(\bigvee_{i \in I} C^i(a)\right) = \begin{cases} \mathbf{0} & \text{if } k \notin I \\ C^k(a) & \text{if } k \in I \end{cases}$$

$$\mathbf{SC}_3 : \forall k \geq \max(\text{sc-dim}(a), \text{sc-dim}(b)), \quad C^k(a \vee b) = C^k(a) \vee C^k(b)$$

$$\mathbf{SC}_4 : \forall i \neq j, \quad \text{sc-dim}(C^i(a) \wedge C^j(a)) < \min(i, j)$$

$$\mathbf{SC}_5 : \forall k \geq \text{sc-dim}(b), \quad C^k(a) - b = C^k(a) - C^k(b).$$

$$\mathbf{SC}_6 : \text{If } b \ll a \text{ then } \text{sc-dim } b < \text{sc-dim } a.$$

It is a **d -scaled lattice** if it satisfies in addition the following property:

$$\mathbf{SC}_0 : \text{sc-dim } a = \dim a$$

All the geometric \mathcal{L}_{SC} -structures in $\text{SC}_{\text{def}}(\mathcal{K}, d)$ or $\text{SC}_{\text{Zar}}(K, d)$ (defined after Example 1.2) are d -scaled lattices (see Fact 2.2). However SC_0 does not follow from the other axioms as the following example shows.

Example 2.7 Let L be an arbitrary noetherian lattice, and $D: \mathcal{I}(L) \rightarrow \{0, \dots, d\}$ be a strictly increasing map. For every $a, b \in L$, if $\mathcal{C}(a)$ denotes the set of all \vee -irreducible components of a , let

$$\begin{aligned} a - b &= \bigvee\{c \in \mathcal{C}(a) / c \not\leq b\}, \\ (\forall k) \quad C_D^k(a) &= \bigvee\{c \in \mathcal{C}(a) / D(c) = k\}. \end{aligned}$$

This is a typical example of a d -subscaled lattice in which the sc-dimension does not coincide with the dimension, except of course if $D(a) = \dim_L a$ for every $a \in \mathcal{I}(L)$. Conversely, every noetherian (in particular every finite) d -subscaled lattice is of that kind.

We call **(sub)scaled lattices** the \mathcal{L}_{SC} -structures whose \mathcal{L}_{SC} -reduct is a d -(sub)scaled for some $d \in \mathbf{N}$. Of course this is not an elementary class. At the contrary, for any fixed $d \in \mathbf{N}$, SC_1^d to SC_5 are expressible by equations, SC_6 by a universal formula and SC_0 by a first order formula in \mathcal{L}_{SC} , hence d -scaled (resp. d -subscaled) lattices form elementary class. As the terminology

suggests, we will see that d -subscaled lattices are precisely the \mathcal{L}_{SC} -substructures of d -scaled lattices.

By analogy with our guiding geometric examples, we say that an element a in a d -subscaled lattice is **k -sc-pure** if

$$\forall b \in L \ (a - b \neq \mathbf{0} \Rightarrow \text{sc-dim}(a - b) = k).$$

We will see that a is k -sc-pure if and only if $a = C^k(a)$ (this is SC_{13} in Section 3). Then either $a = \mathbf{0}$ or $\text{sc-dim } a = k$. In the latter case we say that a **has pure sc-dimension k** . For any a , the element $C^k(a)$ is called the **k -sc-pure component** of a , or simply its **k -pure component** if L is a scaled lattice. By construction these notions coincide with their geometric counterparts in $\text{SC}_{\text{def}}(\mathcal{K}, d)$ and $\text{SC}_{\text{Zar}}(K, d)$.

The following notation will be convenient in induction arguments. If \mathcal{L} is any of our languages \mathcal{L}_{lat} , \mathcal{L}_{TC} or \mathcal{L}_{SC} we let $\mathcal{L}^* = \mathcal{L} \setminus \{\mathbf{1}\}$. Given an \mathcal{L} -structure L whose reduct to \mathcal{L}_{lat} is a lattice, for any $a \in L$ we let

$$L(a) = \{b \in L \mid b \leq a\}.$$

$L(a)$ is a typical example of \mathcal{L}^* -substructure of L .

3 Basic properties and embeddings

The next properties follow easily from the axioms of d -subscaled lattices.

SC₇ $\text{sc-dim } a = \max\{k \mid C^k(a) \neq \mathbf{0}\}$.

In particular $\forall k$, $\text{sc-dim } C^k(a) = k \iff C^k(a) \neq \mathbf{0}$.

SC₈ $\forall k \geq \text{sc-dim}(a)$, $\text{sc-dim}(b \wedge a) < k \implies C^k(a) - b = C^k(a)$.

SC₉ $\text{sc-dim } a \vee b = \max(\text{sc-dim } a, \text{sc-dim } b)$.

In particular $b \leq a \implies \text{sc-dim } b \leq \text{sc-dim } a$.

SC₁₀ $\forall k \geq \text{sc-dim}(a)$, $C^k(a)$ is the largest k -sc-pure element lower than a .

SC₁₁ $\dim a \leq \text{sc-dim } a$.

SC₁₂ $\forall I \subseteq \{0, \dots, d\}$, $a - \bigvee_{i \in I} C^i(a) = \bigvee_{i \notin I} C^i(a)$.

In particular $\text{sc-dim} \left(a - \bigvee_{i \geq k} C^i(a) \right) < k$.

SC₁₃ $\forall k$, $C^k(a) = a \iff \forall b \ (a - b \neq \mathbf{0} \implies \text{sc-dim } a - b = k)$.

That is a is k -sc-pure if and only if $a = C^k(a)$.

In particular if a is \vee -irreducible then a is sc-pure by SC_1^d .

Proof: (Sketch) SC_7 follows from SC_1^d and SC_2 ; SC_8 from SC_5 and SC_7 ; SC_9 from SC_2 , SC_3 and SC_7 ; SC_{10} from SC_2 and SC_3 ; SC_{11} from SC_6 by Fact 2.4. Only the two last properties require a little effort.

SC₁₂: For every $l \in I$, $C^l(a) \leq \bigvee_{i \in I} C^i(a)$ hence $C^l(a) - \bigvee_{i \in I} C^i(a) = \mathbf{0}$. On the other hand for every $l \notin I$ and every $i \in I$, $C^l(a) - C^i(a) = C^l(a)$ by SC_4 and SC_5 . So $C^l(a) - \bigvee_{i \in I} C^i(a) = C^l(a)$ by TC_4 . Finally by SC_1^d and TC_2 ,

$$a - \bigvee_{i \in I} C^i(a) = \bigvee_{l \leq d} \left(C^l(a) - \bigvee_{i \in I} C^i(a) \right) = \bigvee_{l \notin I} C^l(a).$$

SC₁₃: Assume that $a = C^k(a)$ and $a - b \neq \mathbf{0}$ for some b . Then $\text{sc-dim } a = k$ so $\text{sc-dim}(a - b) \leq k$ and $\text{sc-dim}(a \wedge b) \leq k$ by SC₉. Since $a = (a - b) \vee (a \wedge b)$ it follows by SC₃ that

$$C^k(a) = C^k((a - b) \vee (a \wedge b)) = C^k(a - b) \vee C^k(a \wedge b).$$

By assumption $C^k(a) = a$, and $C^k(a \wedge b) \leq a \wedge b$ by SC₁^d. So $a \leq C^k(a - b) \vee (a \wedge b) \leq C^k(a - b) \vee b$ which implies that $a - b \leq C^k(a - b)$. In particular $C^k(a - b) \neq \mathbf{0}$. Since $\text{sc-dim}(a - b) \leq k$ it follows that $\text{sc-dim}(a - b) = k$ by SC₇.

Conversely assume that $a \neq C^k(a)$ (hence $a \neq \mathbf{0}$). For $b = C^k(a)$ we then have $a - b \neq \mathbf{0}$ on one hand and $\text{sc-dim}(a - b) \neq k$ by SC₇ on the other hand, because $C^k(a - b) = \mathbf{0}$ by SC₁₂ and SC₂.

■

Proposition 3.1 *The \mathcal{L}_{SC} -structure of a d -scaled lattice L is uniformly definable in the \mathcal{L}_{lat} -structure of L . In particular it is uniquely determined by this \mathcal{L}_{lat} -structure.*

Proof: Clearly the \mathcal{L}_{TC} -structure is an extension by definition of the lattice structure of L . For every positive integer k the class of k -pure elements is uniformly definable, using the definability of \ll and Fact 2.4. Then so is the function C^k for every k , by decreasing induction on k . Indeed by SC₁₀ and SC₁₂, $C^k(a)$ is the largest k -pure element c such that $c \leq a - \mathbb{W}_{i>k} C^i(a)$.

■

We need a reasonably easy criterion for an \mathcal{L}_{lat} -embedding of subscaled lattices to be an \mathcal{L}_{SC} -embedding. In the special case of a noetherian⁵ embedded lattice, it is given by Proposition 3.3 below, whose proof will use the next characterisation of sc-pure components.

Proposition 3.2 *Let L be a subscaled lattice and $a, a_0, \dots, a_d \in L$ be such that $a = \mathbb{W}_{i \leq d} a_i$, each a_i is i -sc-pure and $\text{sc-dim}(a_i \wedge a_j) < \min(i, j)$ for every $i \neq j$. Then $C^i(a) = a_i$ for every i .*

Proof: Note first that $C^k(a_k) = a_k$ for every $k \leq d$ by SC₁₃. Hence for every $k \leq i \leq d$ we have by SC₃ and SC₂

$$C^i\left(\mathbb{W}_{k \leq i} a_k\right) = \mathbb{W}_{k \leq i} C^i(a_k) = \mathbb{W}_{k \leq i} C^i(C^k(a_k)) = C^i(a_i) = a_i. \quad (1)$$

In particular $C^d(a) = a_d$. Now assume that for some $i < d$ we have proved that $C^j(a) = a_j$ for $i < j \leq d$. By SC₁₂ and SC₂ we then have

$$C^i\left(a - \mathbb{W}_{j>i} a_j\right) = C^i\left(a - \mathbb{W}_{j>i} C^j(a)\right) = C^i\left(\mathbb{W}_{j \leq i} C^j(a)\right) = C^i(a). \quad (2)$$

On the other hand $a - \mathbb{W}_{j>i} a_j = \mathbb{W}_{k \leq d} (a_k - \mathbb{W}_{j>i} a_j)$ by TC₂. For $k > i$ obviously $a_k - \mathbb{W}_{j>i} a_j = \mathbf{0}$. For $k < i$, $a_k = C^k(a)$ and $a_j = C^j(a_j)$ imply that $\text{sc-dim}(a_k \wedge a_j) < k$ for $j > i$ by SC₄. Hence $a_k - a_j = a_k$ by SC₈ and finally $a_k - \mathbb{W}_{j>i} a_j = a_k$ by TC₄, so

$$a - \mathbb{W}_{j>i} a_j = \mathbb{W}_{k \leq i} a_k. \quad (3)$$

By (1), (2), (3) we conclude that $C^i(a) = a_i$. The result follows for every i by decreasing induction.

■

⁵Although we won't use it, let us mention that in the general case of an \mathcal{L}_{lat} -embedding $\varphi : L \rightarrow L'$ between arbitrary subscaled lattices, one may easily derive from Proposition 3.3, by means of the Local Finiteness Theorem 4.1 and the model-theoretic compactness theorem, that φ is an \mathcal{L}_{SC} -embedding if and only if it preserves sc-dimension and sc-purity, that is for every $a \in L$ and every $k \in \mathbf{N}$, $C^k(a) = a \Rightarrow C^k(\varphi(a)) = \varphi(a)$.

Proposition 3.3 *Let L_0 be a noetherian subscaled lattice, L a subscaled lattice, and $\varphi : L_0 \rightarrow L$ an \mathcal{L}_{lat} -embedding such that for every $a \in \mathcal{I}(L_0)$, $\varphi(a)$ is sc-pure and has the same sc-dimension as a . Then φ is an \mathcal{L}_{SC} -embedding.*

Remark 3.4 Clearly the same statement remains true with \mathcal{L}_{lat} and \mathcal{L}_{SC} replaced respectively by $\mathcal{L}_{\text{lat}}^*$ and $\mathcal{L}_{\text{SC}}^*$. We will freely use these variants.

Proof: L_0 and L are d -subscaled lattices for some $d \in \mathbf{N}$. Given $a \in L_0$ and k a non-negative integer, we first check that $\varphi(C^k(a))$ is k -sc-pure. Note that every \vee -irreducible component c of $C^k(a)$ in L_0 has sc-pure dimension k . Indeed $C^k(a)$ is k -sc-pure by SC_{13} , and $c = C^k(a) - b \neq \mathbf{0}$ where b is the join of all the other \vee -irreducible components of $C^k(a)$, hence $\text{sc-dim}(c) = \text{sc-dim}(C^k(a) - b) = k$. Moreover c is sc-pure because it is \vee -irreducible, hence c is k -pure. By our assumption on φ it follows that $\varphi(c)$ is k -sc-pure. Every finite union of k -sc-pure elements being k -sc-pure by SC_3 , it follows that

$$\varphi(C^k(a)) \text{ is } k\text{-sc-pure.} \quad (4)$$

Now for every $l \neq k$ we have $\text{sc-dim}(C^k(a) \wedge C^l(a)) < \min(k, l)$ by SC_4 . It follows that each \vee -irreducible component c of $C^k(a) \wedge C^l(a)$ has sc-dimension strictly less than $\min(k, l)$, hence so does $\varphi(c)$ by assumption. By SC_9 we conclude that

$$\text{sc-dim}(C^k(a) \wedge C^l(a)) < \min(k, l) \quad (\forall l \neq k). \quad (5)$$

$\varphi(a) = \mathbb{W}_{k \leq d} \varphi(C^k(a))$ by SC_1^d and because φ is an \mathcal{L}_{lat} -embedding. By (4), (5) and Proposition 3.2 it follows that $C^k(\varphi(a)) = \varphi(C^k(a))$ for every $k \leq d$. Since φ is injective, this implies by SC_7 that for every $a \in L_0$

$$\text{sc-dim } a = \text{sc-dim } \varphi(a). \quad (6)$$

It only remains to check that $\varphi(a - b) = \varphi(a) - \varphi(b)$ for every $a, b \in L_0$. By TC_2 , replacing if necessary a by its \vee -irreducible components, we may assume w.l.o.g. that a itself is \vee -irreducible in L_0 . This implies that $a = C^k(a)$ for some k , hence $\varphi(a)$ is k -sc-pure by assumption on φ . It then remains two possibilities for $a - b$:

- If $b \geq a$ then $\varphi(b) \geq \varphi(a)$, hence $a - b = \mathbf{0}$ and $\varphi(a) - \varphi(b) = \mathbf{0}$, so $\varphi(a - b) = \varphi(\mathbf{0}) = \mathbf{0} = \varphi(a) - \varphi(b)$.
- Otherwise $b \wedge a < a$ hence $b \wedge a \ll a$ by TC_1 , that is $a - b = a$. So we have to prove that $\varphi(a) - \varphi(b) = \varphi(a)$. By SC_6 $\text{sc-dim } b \wedge a < \text{sc-dim } a$, hence $\text{sc-dim}(\varphi(b) \wedge \varphi(a)) < \text{sc-dim}(\varphi(a)) = k$ by (6). Since $\varphi(a)$ is k -sc-pure it follows that $\varphi(a) - \varphi(b) = \varphi(a)$ by SC_5 .

■

Corollary 3.5 *Let L_0 be a noetherian lattice embedded in a subscaled lattice L . Assume that every $b' < b \in \mathcal{I}(L_0)$ are sc-pure in L and $\text{sc-dim } b' < \text{sc-dim } b$ in L . Then the restrictions to L_0 of the \mathcal{L}_{SC} -operations “ $-$ ” and “ C_i ” of L turn L_0 into an \mathcal{L}_{SC} -substructure which is a subscaled lattice.*

Proof: The assumptions imply that the map $D : a \mapsto \text{sc-dim } a$ is a strictly increasing map from $\mathcal{I}(L_0)$ to \mathbf{N} . Endow L_0 with the structure of subscaled lattice determined by D as in Example 2.7. By construction the inclusion map from L_0 to L is an \mathcal{L}_{lat} -embedding which preserves the sc-purity and sc-dimension of every $b \in \mathcal{I}(L_0)$, hence is an \mathcal{L}_{SC} -embedding by Proposition 3.3.

■

4 Local finiteness

We prove in this section that every finitely generated subscaled lattice is finite. This result is far non-obvious, due to the lack of any known normal form for terms in \mathcal{L}_{SC} . It contrasts with the general situation in co-Heyting algebras, which can be both infinite and generated by a single element. Our main ingredient, which explains this difference, is the uniform bound given *a priori* for the sc-dimension of any element in a given d -subscaled lattice.

Theorem 4.1 *Any d -subscaled lattice L generated by n elements is finite. More precisely, the cardinality of $\mathcal{I}(L)$ is bounded by the function $\mu(n, d)$ defined by*

$$\mu(n, d) = 2^n + \mu(2^{n+1}, d - 1).$$

for $d \geq 0$, and $\mu(n, d) = 0$ for $d < 0$.

Proof: The only subscaled lattice of sc-dimension $d < 0$ is the one-element lattice $\{\mathbf{0}\}$, so the result is trivial in this case. Assume that $d \geq 0$ and that the result is proved for every $d' < d$ and every non-negative integer n .

Let L be a subscaled lattice of sc-dimension d generated by elements x_1, \dots, x_n . Let Ω_n be the family of all subsets of $\{1, \dots, n\}$ (so $\Omega_0 = \{\emptyset\}$). For every $I \in \Omega_n$ let $I^c = \Omega_n \setminus I$ and

$$y_I = \left(\bigwedge_{i \in I} x_i \right) - \left(\bigvee_{i \in I^c} x_i \right), \quad z_I = C^d(y_I).$$

The family of all $\mathcal{Y}_I = \bigcap_{i \in I} P(x_i) \cap \bigcap_{i \in I^c} P(x_i)^c$ is a partition of $\text{Spec}(L)$. Indeed the \mathcal{Y}_i 's are the atoms of the boolean algebra generated in the power set $\mathcal{P}(\text{Spec}(L))$ by the $P(x_i)$'s. Moreover each $P(y_I)$ is the topological closure $\bar{\mathcal{Y}}_I$ of \mathcal{Y}_I in $\text{Spec}(L)$ hence for every $x \in L$

$$P(x) = \bigcup_{I \in \Omega_n} P(x) \cap \mathcal{Y}_I \subseteq \bigcup_{I \in \Omega_n} P(x) \cap \bar{\mathcal{Y}}_I = P\left(\bigvee_{I \in \Omega_n} x \wedge y_I\right).$$

So $x \leq \bigvee_{I \in \Omega_n} (x \wedge y_I)$ by Stone-Priestley's duality. The reverse inequality being obvious we have proved that

$$\forall x \in L, \quad x = \bigvee_{I \in \Omega_n} (x \wedge y_I). \quad (7)$$

In particular SC_3 also gives

$$C^d(\mathbf{1}) = C^d\left(\bigvee_{I \in \Omega_n} y_I\right) = \bigvee_{I \in \Omega_n} z_I. \quad (8)$$

For every $I \neq J \in \Omega_n$, if for example $I \not\subseteq J$ choose any $i \in I \setminus J$ and observe that $y_I \leq x_i$ and $y_J \leq \mathbf{1} - x_i$ so $y_I \wedge y_J \ll \mathbf{1} - x_i$ by TC_3 . By SC_6 and the d -sc-purity of the z_I 's it follows that

$$\text{sc-dim } z_I \wedge z_J < d \quad \text{hence} \quad z_I - z_J = z_I. \quad (9)$$

It follows from SC_9 , SC_{12} and (9) above, that the element

$$a = (\mathbf{1} - C^d(\mathbf{1})) \vee \left(\bigvee_{I \in \Omega_n} (y_I - z_I) \right) \vee \left(\bigvee_{I \neq J \in \Omega_n} (z_I \wedge z_J) \right)$$

has sc-dimension strictly smaller than d . So the induction hypothesis applies to the \mathcal{L}_{SC} -substructure L_0^- of $L(a)$ generated by the $(y_I - z_I)$'s and the $(z_I \wedge z_J)$'s: L_0^- is finite, with at most $\mu(2|\Omega_n|, d-1)$ \vee -irreducible elements. Note that L_0^- is an $\mathcal{L}_{\text{SC}}^*$ -substructure of L (recall that

$\mathcal{L}_{\text{SC}}^* = \mathcal{L}_{\text{SC}} \setminus \{\mathbf{1}\}$. Finally let L_1 be the upper semi-lattice generated in L by $L_0^- \cup \{z_I\}_{I \in \Omega_n}$. By construction L_1 is finite and $\mathcal{I}(L_1) \subseteq \mathcal{I}(L_0^-) \cup \{z_I\}_{I \in \Omega_n}$, so $|\mathcal{I}(L_1)| \leq 2^n + \mu(2^{n+1}, d-1) = \mu(n, d)$. It is then sufficient to show that $L_1 = L$.

We first prove that L_1 is a lattice. By (8), $C^d(\mathbf{1}) \vee a = \mathbb{W}_{I \in \Omega_n} z_I \vee a \in L_1$ hence $\mathbf{1} = C^d(\mathbf{1}) \vee a \in L_1$. For every $I \in \Omega_n$ and every $b \in L_0^-$, $z_I \wedge b = (z_I \wedge a) \wedge b \in L_0^-$. For every $I \neq J \in \Omega_n$, $z_I \wedge z_J = (z_I \wedge a) \wedge (z_J \wedge a) \in L_0^-$. So by the distributivity law, L_1 is a sublattice of L .

In order to conclude that L_1 is an \mathcal{L}_{SC} -substructure of L , by Corollary 3.5 it only remains to check that for every $b' < b \in \mathcal{I}(L_1)$, b is sc-pure in L and $\text{sc-dim } b' < \text{sc-dim } b$ in L . Since $\mathcal{I}(L_1) \subseteq \mathcal{I}(L_0^-) \cup \{z_I\}_{I \in \Omega_n}$ we can distinguish two cases.

Case 1: $b \in \mathcal{I}(L_0^-)$. Then b is sc-pure in L_0^- by SC_{13} hence also in L since L_0^- is an $\mathcal{L}_{\text{SC}}^*$ -substructure of L . Similarly $b' \ll b$ in L_0^- by TC_1 that is $b - b' = b$ in L_0^- hence also in L . Thus $b' \ll b$ in L which implies that $\text{sc-dim } b' < \text{sc-dim } b$ in L by SC_6 .

Case 2: $b = z_I$ for some $I \in \Omega_n$. Then $b = C^d(y_I)$ is sc-pure in L and $\text{sc-dim } b = d$. If $b' = z_J$ for some other $J \in \Omega_n$ then on one hand $\text{sc-dim}(b') = d$ and on the other hand $I \neq J$ hence $b' = b' \wedge b = z_I \wedge z_J$ has sc-dimension $< d$ by (9), a contradiction. So necessarily $b' \in \mathcal{I}(L_0^-)$, in particular $b' \leq a$ hence $\text{sc-dim}(b') \leq \text{sc-dim}(a) < d$.

So L_1 is indeed an \mathcal{L}_{SC} -substructure of L . Finally every $y_I = (y_I - z_I) \vee z_I \in L_1$ and (7) gives, for every $i \leq n$,

$$x_i = \mathbb{W}_{I \in \Omega_n} x_i \wedge y_I \leq \mathbb{W}_{\substack{I \in \Omega_n \\ i \in I}} y_I \leq x_i.$$

So equality holds, hence each $x_i \in L_1$, which finally proves that $L_1 = L$.

■

Corollary 4.2 *For every n, d there are finitely many non-isomorphic subscaled lattices of sc-dimension d generated by n elements.*

Proof: Any such subscaled lattice L is finite, with $|\mathcal{I}(L)| \leq \mu(n, d)$ by Theorem 4.1. Clearly there are finitely many non-isomorphic lattices such that $|\mathcal{I}(L)| \leq \mu(n, d)$ and each of them admits finitely many non-isomorphic \mathcal{L}_{SC} -structures of d -subscaled lattices.

■

5 Linear representation

In this section we prove that the theory of d -subscaled lattices is the universal theory of various natural classes of \mathcal{L}_{SC} -structures, including $\text{SC}_{\text{Zar}}(K, d)$. The argument is based on an elementary representation theorem for d -subscaled lattices, combined with the local finiteness result of Section 4.

Given an arbitrary field K , a non-empty linear variety $X \subseteq K^m$ is determined by the data of an arbitrary point $P \in X$ and the vector subspace \vec{X} of K^m , *via* the relation $X = P + \vec{X}$ (the orbit of P under the action of \vec{X} by translation). We call X a **special linear variety** (resp. a **special linear set**) if X is a linear variety such that \vec{X} is generated by a subset of the canonical basis of K^m (resp. if X is a finite union of special linear varieties). The family $L_{\text{lin}}(X)$ of all special linear subsets of X is the family of closed sets of a noetherian topology on X , hence a noetherian lattice. For every $A \in L_{\text{lin}}(X)$ we let $D(A)$ be the dimension of A in the sense of linear algebra. This endows $L_{\text{lin}}(X)$ with a natural structure of scaled lattice as in Example 2.7.

Remark 5.1 For every $A \in \mathbf{L}_{\text{lin}}(X)$, if K is infinite then $\text{sc-dim } A = \dim_{\mathbf{L}_{\text{lin}}(X)} A =$ the dimension of A as defined in linear algebra. It coincides with the Krull dimension as well. If moreover A is \vee -irreducible in $\mathbf{L}_{\text{lin}}(X)$ then it is pure dimensional, hence it is sc-pure both in $\mathbf{L}_{\text{lin}}(X)$ and $\mathbf{L}_{\text{Zar}}(X)$. By Proposition 3.3 it follows that $\mathbf{L}_{\text{lin}}(X)$ is an \mathcal{L}_{SC} -substructure of $\mathbf{L}_{\text{Zar}}(X)$. Similarly if K is a real closed field or p -adically closed field then $\mathbf{L}_{\text{lin}}(X)$ is an \mathcal{L}_{SC} -substructure of $\mathbf{L}_{\text{def}}(X)$.

In what follows K^m is identified to $K^m \times \{0\}^r \subseteq K^{m+r}$. The very easy result below prepares the proof of Proposition 5.3.

Proposition 5.2 *For every two special linear sets $C \subseteq B \subseteq K^m$ and every non-negative integer $n \geq \dim C$ there exists a special linear set $A \subseteq K^{m+n}$ of pure dimension n such that $A \cap B = C$.*

Proof: The result being rather trivial if C is empty, we can assume w.l.o.g. that $C \neq \emptyset$. Let (e_1, \dots, e_{m+n}) be the canonical basis of K^{m+n} . If I is a subset of $\{1, \dots, m+n\}$ we let $\vec{E}(I)$ denote the vector space generated in K^n by $(e_i)_{i \in I}$. Decompose C as a union of special linear varieties C_1, \dots, C_p , and write each $C_i = P_i + \vec{E}(J_i)$ with $|J_i| = \dim C_i \leq n$. Let $I_i = J_i \cup \{m+1, \dots, m+n-|J_i|\}$ and $A_i = P_i + \vec{E}(I_i)$ for every $i \leq p$. Finally let $A = A_1 \cup \dots \cup A_p$. By construction each A_i has pure dimension $|I_i| = n$, hence A has pure dimension n . Clearly each $A_i \cap K^m = C_i$, hence $A \cap K^m = C$ and *a fortiori* $A \cap B = C$. ■

Proposition 5.3 (Linear representation) *Let K be an infinite field, $d \geq 0$ an integer and L a finite d -subscaled lattice. Then there exists a special linear set X over K of dimension $\leq d$ and an \mathcal{L}_{SC} -embedding $\varphi : L \rightarrow \mathbf{L}_{\text{lin}}(X)$.*

Proof: By induction on the number r of \vee -irreducible elements of an arbitrary d -subscaled lattice L , we prove that there exists an $\mathcal{L}_{\text{SC}}^*$ -embedding φ of L into $\mathbf{L}_{\text{lin}}(K^m)$ for some m depending on L . Taking $X = \varphi(\mathbf{1}_L)$ then gives the conclusion. Indeed X is a special linear set over K , $\dim X = \text{sc-dim}(\mathbf{1}_L) \leq d$ because φ preserves the sc-dimension, and φ is obviously an \mathcal{L}_{SC} -embedding of L into $\mathbf{L}_{\text{lin}}(X)$.

If $r = 0$ then L is the one-element lattice $\{\mathbf{0}\}$, hence an $\mathcal{L}_{\text{SC}}^*$ -substructure of $\mathbf{L}_{\text{lin}}(K)$. So, given a fixed $r \geq 1$, we can assume by induction that the result is proved for $r - 1$. Let L be a d -subscaled lattice with \vee -irreducible elements a_1, \dots, a_r .

Renumbering if necessary we may assume that a_r is maximal among the a_i 's. Let $a = a_r$, $b = \bigvee_{1 \leq i < r} a_i$, $c = a \wedge b$ and φ an $\mathcal{L}_{\text{SC}}^*$ -embedding of $L(b)$ into some $\mathbf{L}_{\text{lin}}(K^m)$ given by the induction hypothesis. Since a is \vee -irreducible in L it is sc-pure. Moreover $c \leq a$ by TC_1 , hence a has pure sc-dimension n for some $n \geq \text{sc-dim}(c)$ by SC_6 . Let B, C be the respective images of b, c by φ . Proposition 5.2 gives a special linear set $A \subseteq K^{m+n}$ of pure dimension n such that $A \cap B = C$. Identifying K^m with $K^m \times \{0\}^n \subseteq K^{m+n}$ turns φ into an $\mathcal{L}_{\text{SC}}^*$ -embedding of $L(b)$ into $\mathbf{L}_{\text{lin}}(K^{m+n})$.

Every $x \in L$ writes uniquely $x_a \vee x_b$ with $x_a \in \{\mathbf{0}, a\}$ and $x_b \in L(b)$ by grouping appropriately the \vee -irreducible components of x , using the maximality of a_r . So we can let

$$\bar{\varphi}(x) = \begin{cases} \varphi(x_b) & \text{if } x_a = \mathbf{0}, \\ A \cup \varphi(x_b) & \text{if } x_a = a. \end{cases}$$

This is a well-defined $\mathcal{L}_{\text{lat}}^*$ -embedding of L into $\mathbf{L}_{\text{lin}}(K^{m+n})$. Moreover $\bar{\varphi}$ is an $\mathcal{L}_{\text{SC}}^*$ -embedding by Proposition 3.3. This finishes the induction. ■

For any infinite field K and positive integer d let $\text{SC}_{\text{lin}}(K, d)$ be the class of d -scaled lattices $\mathbf{L}_{\text{lin}}(X)$ with X ranging over the special linear sets over K of dimension at most d .

Theorem 5.4 *Given any non-negative integer d , the universal theories of $\text{SC}_{\text{def}}(K, d)$ (resp. $\text{SC}_{\text{zar}}(K, d)$, $\text{SC}_{\text{lin}}(K, d)$) in the language \mathcal{L}_{SC} are the same for every o -minimal or P -minimal expansion \mathcal{K} of a field K (resp. for every infinite field K). This is the theory of d -subscaled lattices.*

Proof: The universal theory of any of these classes obviously contains the theory of d -subscaled lattices. For the converse, thanks to Remark 5.1 it suffices to prove that every d -subscaled lattice L embeds into a model of the theory of $\text{SC}_{\text{lin}}(K, d)$. If L is finite this is Proposition 5.3. The general case then follows from the model-theoretic compactness theorem, because L is locally finite by Theorem 4.1.

■

6 Minimal extensions

Minimal proper extensions of any finite subscaled lattices are entirely determined by so-called “SC-signatures” (see below). Since this is a special case of minimal extensions of finite co-Heyting algebras, we first recall what the main results of [DJ] on this subject, and try to reduce to them as much as possible.

We need some specific notation and definitions. Given a finite lattice L_0 , an \mathcal{L}_{lat} -extension L , elements $a \in L_0$ and $x \in L$ we write:

- $a^- = \mathbb{W}\{b \in L_0 \mid b < a\}$.
- $g(x, L_0) = \mathbb{M}\{a \in L_0 \mid x \leq a\}$.

Clearly $a \in \mathcal{I}(L_0)$ if and only if a^- is the unique predecessor of a in L_0 (otherwise $a^- = a$).

Assume that L_0 and L are co-Heyting algebras (or topologically complemented lattices, or TC-lattice for short). A **TC-signature** in L_0 is a triple (g, H, r) where $g \in \mathcal{I}(L_0)$, H is a set of one or two elements $h_1, h_2 \in L_0$ and $r \in \{1, 2\}$ are such that:

- either $r = 1$ and $h_1 = h_2 < g$;
- or $r = 2$ and $h_1 \vee h_2 < g$.

A couple (x_1, x_2) of elements of L is **TC-primitive over** L_0 if there is $g \in \mathcal{I}(L_0)$ such that

P1 $g^- \wedge x_1$ and $g^- \wedge x_2$ belong to L_0 .

P2 One of the following happens:

1. $x_1 = x_2$ and $g^- \wedge x_1 \ll x_1 \ll g$.
2. $x_1 \neq x_2$, $x_1 \wedge x_2 \in L_0$ and $g - x_1 = x_2$, $g - x_2 = x_1$.

This implies that each $x_i \notin L_0$, that $g = g(x_1, L_0) = g(x_2, L_0)$ and that the triple $\sigma_{\text{TC}}(x_1, x_2) = (g, H, r)$ defined as follows is a signature in L_0 , called the **signature of** (x_1, x_2) **in** L_0 .

$$g = g(x_1, L_0) \qquad H = \{g^- \wedge x_1, g^- \wedge x_2\} \qquad r = \text{Card}\{x_1, x_2\}$$

Finally we say that L is a **TC-primitive extension** of L_0 if it is \mathcal{L}_{TC} -generated over L_0 by a primitive tuple. For the convenience of the reader we collect here all the properties of TC-signatures and TC-primitive extensions that we are going to use. All the references are taken from [DJ].

Proposition 6.1 *Let L_0 be a finite co-Heyting algebra and L an \mathcal{L}_{TC} -extension.*

1. [Theorem 3.3] *If L is \mathcal{L}_{TC} -generated over L_0 by a TC-primitive tuple (x_1, x_2) , then L is exactly the upper semi-lattice generated over L_0 by x_1 and x_2 . It is a finite co-Heyting algebra and one of the following happens :*

- (a) $x_1 = x_2$ and $\mathcal{I}(L) = \mathcal{I}(L_0) \cup \{x_1\}$.
- (b) $x_1 \neq x_2$ and $\mathcal{I}(L) = (\mathcal{I}(L_0) \setminus \{g\}) \cup \{x_1, x_2\}$.

*In particular (x_1, x_2) is (up to permutation) the only TC-primitive tuple over L_0 in L . We call $\sigma_{\text{TC}}(x_1, x_2)$ the **TC-signature of L in L_0** and denote it $\sigma_{\text{TC}}(L)$.*

2. [Remark 3.6] *TC-signatures in L_0 and TC-primitive extensions of L_0 are in one-to-one correspondence: every TC-signature in L_0 is the TC-signature of a TC-primitive extension, and two TC-primitive extensions of L_0 are \mathcal{L}_{TC} -isomorphic over L_0 if and only if they have the same TC-signature in L_0 .*

3. [Corollary 3.4] *If L is finite, the following are equivalent.*

- (a) L is a minimal proper extension of L_0 .
- (b) L is a TC-primitive extension of L_0 .
- (c) $\text{Card}(\mathcal{I}(L)) = \text{Card}(\mathcal{I}(L_0)) + 1$.

As a consequence every finite \mathcal{L}_{TC} -extension L' of L_0 is the union of a tower of TC-primitive extensions $L_0 \subset L_1 \subset \dots \subset L_n = L'$ with $n = \text{Card}(\mathcal{I}(L')) - \text{Card}(\mathcal{I}(L_0))$.

We will refer to the k -th item of the above proposition as to Proposition 6.1. k .

Now let L_0 be a finite subscaled lattice and L and \mathcal{L}_{SC} -extension. An **SC-signature** in L_0 is a triple $\sigma = (g, H, q)$ where $g \in \mathcal{I}(L_0)$, H is a set of one or two elements $h_1, h_2 \in L_0$ and $q \in \mathbf{N}$ are such that:

- either $\text{sc-dim } h_1 < q < \text{sc-dim } g$ and $h_1 = h_2 < g$;
- or $q = \text{sc-dim } g$ and $h_1 \vee h_2 = g^-$.

Let $r_\sigma = 1$ if $q < \text{sc-dim } g$, $r_\sigma = 2$ if $q = \text{sc-dim } g$, and $\sigma^{\text{TC}} = (g, H, r_\sigma)$. By construction this is a TC-signature in L_0 . Given an \mathcal{L}_{SC} -extension L of L_0 , a tuple (x_1, x_2) of elements of L is **SC-primitive over L_0** if it is TC-primitive over L_0^{TC} and if in addition

P3 x_1, x_2 are sc-pure of the same sc-dimension.

Such an SC-primitive tuple determines the so-called **SC-signature of (x_1, x_2) in L_0** , denoted $\sigma_{\text{SC}}(x_1, x_2) = (g, H, q)$ and defined as follows.

$$g = g(x_1, L_0) \qquad H = \{g^- \wedge x_1, g^- \wedge x_2\} \qquad q = \text{sc-dim } x_1$$

Note that, by condition **P2** of the definition of TC-signatures, $x_1 = x_2$ if and only if $x_1 \ll g$, and otherwise $\text{sc-dim } x_1 = \text{sc-dim } x_2 = \text{sc-dim } g$. This ensures that $\sigma_{\text{TC}}(x_1, x_2) = (\sigma_{\text{SC}}(x_1, x_2))^{\text{TC}}$.

Let L_0^{TC} and L^{TC} denote the respective \mathcal{L}_{TC} -reducts of L_0 and L . For every subset X_0 of L we let:

- $L_0 \langle X_0 \rangle$ = the \mathcal{L}_{SC} -structure generated by $L_0 \cup X_0$ in L ;

- $L_0^{\text{TC}}\langle X_0 \rangle =$ the \mathcal{L}_{TC} -structure generated by $L_0^{\text{TC}} \cup X_0$ in L^{TC} .

We say that L is an **SC-primitive extension** of L_0 , if there exists a tuple (x_1, x_2) SC-primitive over L_0 such that $L = L_0\langle x_1, x_2 \rangle$ (then clearly $L = L_0\langle x_1 \rangle = L_0\langle x_2 \rangle$). By Lemma 6.2 below and Proposition 6.1.1 such a tuple is necessarily unique.

Lemma 6.2 *Let L_0 be finite subscaled lattice, and L an \mathcal{L}_{SC} -extension generated over L_0 by an SC-primitive tuple (x_1, x_2) . Then $L^{\text{TC}} = L_0\langle x_1, x_2 \rangle$, (x_1, x_2) is TC-primitive over L_0^{TC} and $\sigma_{\text{TC}}(x_1, x_2) = (\sigma_{\text{SC}}(x_1, x_2))^{\text{TC}}$.*

Proof: That (x_1, x_2) is TC-primitive over L_0^{TC} and $\sigma_{\text{TC}}(x_1, x_2) = (\sigma_{\text{SC}}(x_1, x_2))^{\text{TC}}$ is only a reminder: it follows directly from the definitions. Let $L_1 = L_0\langle x_1, x_2 \rangle$, in order to conclude that $L_1 = L$ it only remains to prove that L_1 is an \mathcal{L}_{SC} -substructure of L . By Corollary 3.5 it suffices to check that for every $b' < b \in \mathcal{I}(L_1)$, b is sc-pure in L and $\text{sc-dim } b' < \text{sc-dim } b$ in L .

If $b \in \mathcal{I}(L_0)$, then b is sc-pure in L_0 by SC_{13} , hence also in L because L_0 is an \mathcal{L}_{SC} -substructure of L . Otherwise $b = x_i$ for some $i \in \{1, 2\}$. Then b is sc-pure in L by definition of SC-primitive tuples over L_0 .

In both cases $b' \ll b$ in L_1 by TC_1 , that is $b - b' = b$ in L_1 , hence also in L because L_1 is an \mathcal{L}_{TC} -substructure of L . So $b' \ll b$ in L hence $\text{sc-dim}(b') < \text{sc-dim}(b)$ in L by SC_6 .

■

Lemma 6.3 *Let L_0 be finite subscaled lattice, L_1 an \mathcal{L}_{TC} -extension generated over L_0^{TC} by a TC-primitive tuple (x_1, x_2) , and $\tau = (g, \{h_1, h_2\}, q)$ an SC-signature in L_0 such that $\tau^{\text{TC}} = \sigma_{\text{TC}}(x_1, x_2)$. Then there exists a unique structure of subscaled lattice expanding L_1 which makes it an \mathcal{L}_{SC} -extension of L_0 such that (x_1, x_2) is SC-primitive over L_0 and $\sigma_{\text{SC}}(x_1, x_2) = \tau$.*

Proof: By Proposition 6.1.1, $\mathcal{I}(L_1) \subseteq \mathcal{I}(L_0) \cup \{x_1, x_2\}$. For every $x \in \mathcal{I}(L_0)$ let $D(x) = \text{sc-dim } x$, and let $D(x_1) = D(x_2) = q$. This defines by restriction a function from $\mathcal{I}(L_1)$ to \mathbf{N} . Assume that D is strictly increasing. Then it determines as in Example 2.7 an \mathcal{L}_{SC} -structure on L_1 expanding its \mathcal{L}_{TC} -structure. Let us denote it L , so that $L^{\text{TC}} = L_1$. Every \vee -irreducible element of L_0 remains sc-pure in L with the same sc-dimension, hence by Proposition 3.3 the inclusion of L_0 into L is an \mathcal{L}_{SC} -embedding. This is clearly the only possible \mathcal{L}_{SC} -structure on L_1 which makes it an \mathcal{L}_{SC} -extension of L_0 such that $\text{sc-dim } x_1 = \text{sc-dim } x_2 = q$. So it only remains to prove that D is strictly increasing.

Let $b < a$ in $\mathcal{I}(L)$, if $a, b \in \mathcal{I}(L_0)$ then $D(b) < D(a)$ by 6. So we can assume that a or b does not belong to $\mathcal{I}(L_0)$. By Proposition 6.1.1 one of them must belong to $\{x_1, x_2\}$ and the other one to $\mathcal{I}(L_0)$. Note that our assumption $\sigma_{\text{TC}}(x_1, x_2) = \sigma^{\text{TC}}$ implies that (for $i = 1, 2$) $g = g(x_i, L_0)$, and $h_i = x_i \wedge g^-$ (up to re-numbering) and: either $x_1 = x_2$, $h_1 = h_2 < g$ and $\text{sc-dim } h_1 < q < \text{sc-dim } g$; or $x_1 \neq x_2$, $h_1 \vee h_2 = g^-$ and $q = \text{sc-dim } g$.

Case 1: $b = x_1$ or $b = x_2$, hence $D(b) = q$. Then $a \in \mathcal{I}(L_0)$, in particular $a \in L_0$, hence $g \leq a$ and so $\text{sc-dim } g \leq \text{sc-dim } a$. If $x_1 = x_2$ then $q < \text{sc-dim } g \leq \text{sc-dim } a$ hence $D(b) < D(a)$. If $x_1 \neq x_2$ then $q = \text{sc-dim } g$, and g is not \vee -irreducible in L hence $g \neq a$. So $g \ll a$ (because $g < a$ and a is \vee -irreducible) hence $\text{sc-dim } g < \text{sc-dim } a$ by 6, that is $D(b) < D(a)$.

Case 2: $a = x_1$ or $a = x_2$, hence $D(a) = q$. Then again $b \in L_0$, and $b < a \leq g$ hence $b \leq g^-$. If $x_1 = x_2$, since $b \leq a \wedge g^- = h_1$ we get $\text{sc-dim } b \leq \text{sc-dim } h_1 < q$, hence $D(b) < D(a)$. If $x_1 \neq x_2$ then $\text{sc-dim } g = q$. Since $b < g$ we have $b \ll g$ (because g is \vee -irreducible) hence $\text{sc-dim } b < \text{sc-dim } g$ by 6. So $\text{sc-dim } b < q$, that is $D(b) < D(a)$.

■

We can now pack all this together.

Proposition 6.4 *Let L_0 be a finite subscaled lattice and L an \mathcal{L}_{SC} -extension.*

1. If L is \mathcal{L}_{SC} -generated over L_0 by an SC-primitive tuple (x_1, x_2) , then L is exactly the upper semi-lattice generated over L_0 by x_1 and x_2 . It is a finite subscaled lattice and one of the following happens :

- (a) $x_1 = x_2$ and $\mathcal{I}(L) = \mathcal{I}(L_0) \cup \{x_1\}$.
- (b) $x_1 \neq x_2$ and $\mathcal{I}(L) = (\mathcal{I}(L_0) \setminus \{g\}) \cup \{x_1, x_2\}$.

In particular (x_1, x_2) is (up to permutation) the only SC-primitive tuple over L_0 in L . We call $\sigma_{\text{SC}}(x_1, x_2)$ the **SC-signature of L in L_0** and denote it $\sigma_{\text{SC}}(L)$.

2. SC-signatures in L_0 and SC-primitive extensions of L_0 are in one-to-one correspondence: every SC-signature in L_0 is the SC-signature of an SC-primitive extension, and two SC-primitive extensions of L_0 are \mathcal{L}_{SC} -isomorphic over L_0 if and only if they have the same SC-signature in L_0 .

3. If L is finite, the following are equivalent.

- (a) L is a minimal proper \mathcal{L}_{SC} -extension of L_0 .
- (b) L is an SC-primitive extension of L_0 .
- (c) $\text{Card}(\mathcal{I}(L)) = \text{Card}(\mathcal{I}(L_0)) + 1$.

As a consequence every finite \mathcal{L}_{SC} -extension L' of L_0 is the union of a tower of SC-primitive extensions $L_0 \subset L_1 \subset \dots \subset L_n = L'$ with $n = \text{Card}(\mathcal{I}(L')) - \text{Card}(\mathcal{I}(L_0))$.

We will refer to the k -th item of the above proposition as to Proposition 6.4. k .

Proof: If L is \mathcal{L}_{SC} -generated over L_0 by an SC-primitive tuple (x_1, x_2) , then by Lemma 6.2 L^{TC} is also \mathcal{L}_{TC} -generated over L_0^{TC} by (x_1, x_2) , which is TC-primitive. The first item follows from Proposition 6.1.1.

Let σ be an SC-signature in L_0 . Then σ^{TC} is a TC-signature in L_0 . Proposition 6.1.2 gives a TC-primitive \mathcal{L}_{TC} -extension L_1 of L_0^{TC} with TC-signature σ^{TC} in L_0 . Lemma 6.3 then gives a unique structure of subscaled lattice expanding L_1 which makes it an SC-primitive extension of L_0 with signature σ in L_0 . Let us denote it L , so that $L^{\text{TC}} = L_1$. Now if L' is another SC-primitive extension with signature σ in L_0 , by Proposition 6.1.2 L'^{TC} is \mathcal{L}_{TC} -isomorphic to L^{TC} over L_0 . The image of L' via this endomorphism defines an \mathcal{L}_{SC} -structure expanding L^{TC} , which makes it an SC-primitive extension of L_0 with the same signature as L . By the uniqueness of such a structure, given by Lemma 6.3, it follows that this \mathcal{L}_{TC} -isomorphism from L'^{TC} to L^{TC} is actually an \mathcal{L}_{SC} -isomorphism, which proves the result.

We finely prove item (3) by cyclic implication. Note that (3b) \Rightarrow (3c) follows from item 1) above.

(3c) \Rightarrow (3a). Let L' be a proper \mathcal{L}_{SC} -extension of L_0 contained in L . Then L'^{TC} is a proper \mathcal{L}_{TC} -extension of L_0^{TC} contained in L^{TC} . By Proposition 6.1.3, (3c) implies that L'^{TC} is a minimal proper \mathcal{L}_{TC} -extension of L_0^{TC} . So $L'^{\text{TC}} = L^{\text{TC}}$, thus necessarily $L' = L$, which proves that L is minimal.

(3a) \Rightarrow (3b). Let x_1 be a minimal element in $\mathcal{I}(L) \setminus \mathcal{I}(L_0)$. Let $g = g(x_1, L_0)$; if $x_1 \ll g$ let $x_2 = x_1$, otherwise let $x_2 = g - x_1$. The proof of Corollary 3.4 in [DJ] shows that (x_1, x_2) is TC-primitive over L_0^{TC} . In particular $x_1, x_2 \in \mathcal{I}(L)$ so they are sc-pure by SC₁₃. The same holds true for g , hence if $x_1 \neq x_2$ then $x_1 = g - x_2$ and $x_2 = g - x_1$ have the same dimension (the dimension of g , by definition of the sc-purity of g). So (x_1, x_2) is actually SC-primitive. Since $L_0^{\text{TC}}\langle x_1, x_2 \rangle = L^{\text{TC}}$, a fortiori $L_0\langle x_1, x_2 \rangle = L$ hence L is SC-primitive over L_0 .
■

7 Model-completion of scaled lattices

We call **super scaled lattice** every subscaled lattice L which satisfy the following additional properties, both of which are clearly axiomatizable by $\forall\exists$ -formulas in \mathcal{L}_{SC} . If $\text{sc-dim } L \leq d$ we call it a **super d -scaled lattice**.

Catenarity For every non-negative integers $r \leq q \leq p$ and every elements $c \leq a \neq \mathbf{0}$, if c is r -sc-pure and a is p -sc-pure then there exists a non-zero q -sc-pure element b such that $c \leq b \leq a$.

If $\text{Spec}L$ is noetherian this property is equivalent to the usual notion of catenarity, namely that any two maximal chains in $\text{Spec}L$ having the same first and last elements have the same length. In particular every d -scaled lattice L of type $L_{\text{Zar}}(X)$ or $L_{\text{lin}}(X)$ satisfies this property. At the contrary none of them satisfy the next property, as it implies that L is atomless.

Splitting: For every elements b_1, b_2, a , if $b_1 \vee b_2 \ll a \neq \mathbf{0}$ then there exists non-zero elements $a_1 \geq b_1$ and $a_2 \geq b_2$ such that:

$$\begin{cases} a_1 = a - a_2 \\ a_2 = a - a_1 \\ a_1 \wedge a_2 = b_1 \wedge b_2 \end{cases}$$

We will then say a_1, a_2 **split a along b_1, b_2** .

Remark 7.1 If $r < p \leq q$ in the Catenarity axiom, the conclusion can be strengthened to $c \ll b \leq a$. Indeed b has pure sc-dimension q and $c \wedge b = c$ has sc-dimension $< q$ hence $b - c = b$ by SC_8 . In particular *every subscaled lattice satisfying the Catenarity axiom is a scaled lattice*. Indeed, given any element a of sc-dimension $d \geq 1$, repeated applications of the Catenarity axiom to $C^d(a)$, $c = \mathbf{0}$ and each integer p from 0 to d , gives a chain of sc-pure elements a_0, \dots, a_d such that

$$\mathbf{0} \neq a_0 \ll a_1 \ll \dots \ll a_d \leq a.$$

By Fact 2.2 it follows that $\dim a \geq d$, and by SC_{11} that $\dim a = d$.

Lemma 7.2 *Let a, b_1, b_2 be elements of a finite subscaled lattice L . If $b_1 \vee b_2 \ll a \neq \mathbf{0}$ then L embeds in a finite subscaled lattice L' containing non-zero elements a_1, a_2 which split a along b_1, b_2 . Moreover, if $C^0(a) \neq \mathbf{0}$ we can require⁶ that all the atoms of L' belong to L_0 .*

Proof: We are going to prove by induction on $d = \text{sc-dim } a$ a slightly more precise result, namely that in addition $x \leq a$ for every $x \in \mathcal{I}(L') \setminus \mathcal{I}(L)$. Let g_1, \dots, g_n be the \vee -irreducible components of a in L . ($n \geq 1$ because $a \neq \mathbf{0}$). If $d = 0$ our assumption that $b_1 \vee b_2 \ll a$ implies by SC_6 that $b_1 = b_2 = \mathbf{0}$. If $n = 1$, that is $a = g_1$ is \vee -irreducible, then $\sigma = (g, \{\mathbf{0}\}, 0)$ is a signature in L . Proposition 6.4.2 gives an SC-primitive tuple (a_1, a_2) generating an \mathcal{L}_{SC} -extension L_1 over L with signature σ . This signature ensures that (a_1, a_2) splits a along $(\mathbf{0}, \mathbf{0})$. If $n \geq 2$, $a_1 = g_1$ and $a_2 = a - a_1$ will do the job. So the result is proved for $d = 0$.

Now assume that $d \geq 1$ and the result is valid until $d - 1$. Note that $g_1^- \vee \dots \vee g_n^-$ is the greatest element $c \in L$ such that $c \ll a$, in particular

$$b_1 \vee b_2 \leq g_1^- \vee \dots \vee g_n^-. \tag{10}$$

⁶This additional condition if $C^0(a) \neq \mathbf{0}$ will be used only later, in Section 9.

Let $u = (\bigvee_{i \leq n} g_i^-) - (b_1 \vee b_2)$ and $u^* = u - C^0(u)$. Since $u \ll a$ we have $\text{sc-dim } u < d$ by SC₆. We are claiming that L embeds in a finite subscaled lattice L without new atoms, in which all the g_i 's are still \vee -irreducible with the same predecessor as in L_0 , and in which there are elements u_1^*, u_2^* which satisfy all the conditions to split u^* along $b_1 \wedge u^*, b_2 \wedge u^*$, except that u_1^*, u_2^* might be zero elements.

By TC₃, $(b_1 \vee b_2) \wedge u^* \ll u^*$ so if $\text{sc-dim } u \leq 0$ we can simply take $u_1^* = u_2^* = \mathbf{0}$ and $L_0 = L$. On the other hand, if $\text{sc-dim } u > 0$ the induction hypothesis applies to $u^*, b_1 \wedge u^*, b_2 \wedge u^*$. It gives a finite subscaled lattice L_0 containing L and elements $u_1^*, u_2^* \in L_0$ which split u^* along $b_1 \wedge u^*, b_2 \wedge u^*$. Moreover we can require that L_0 do not contain any new atom because $C^0(u^*) = \mathbf{0}$, and that $x \leq u^*$ for every $x \in \mathcal{I}(L) \setminus \mathcal{I}(L_0)$. For every $x \in \mathcal{I}(L)$ such that $x < g_i$ for some $i \leq n$, if $x \in L_0$ then $x \leq g_i^-$ (where g_i^- still denotes the predecessor of g_i in L_0). If $x \notin L_0$ then $x \leq u^*$ by construction hence $x \leq g_i \wedge u^*$. The latter belongs to L_0 and is strictly lower than g_i , hence lower than g_i^- , so $x < g_i$. It follows that g_i has a unique predecessor in L which is g_i^- , in particular g_i remains \vee -irreducible in L . This proves our claim in both cases.

Now let $u_1 = C^0(u) \vee u_1^*$ and $u_2 = u_2^*$. We have in particular

$$u_1 \vee u_2 = \bigvee_{i \leq n} g_i^- - (b_1 \vee b_2). \quad (11)$$

Since $u - b_2 = u$ by TC₃ necessarily $b_2 \wedge c \ll c$ for every \vee -irreducible component c of u , hence $b_2 \wedge C^0(u) \ll C^0(u)$. By SC₆ it follows that $b_2 \wedge C^0(u) = \mathbf{0}$ hence $b_2 \wedge u_1 = b_2 \wedge u_1^*$. Similarly $u^* \wedge C^0(u) = \mathbf{0}$ because $u^* - C^0(u) = u^*$ by SC₁₂ and TC₃. *A fortiori* $u_2^* \wedge C^0(u) = \mathbf{0}$ hence $u_2^* \wedge u_1 = u_2^* \wedge u_1^*$. Note also that $b_1 \wedge u_2^* = b_1 \wedge u^* \wedge u_2^* \leq u_1^* \wedge u_2^*$, and symmetrically $b_2 \wedge u_1^* \leq u_1^* \wedge u_2^*$. Altogether, since $u_2 = u_2^*$ and $u_1^* \wedge u_2^* \leq b_1 \wedge b_2$ by construction, this gives

$$(b_1 \wedge u_2) \vee (b_2 \wedge u_1) \vee (u_1 \wedge u_2) \leq (b_1 \wedge b_2)$$

hence

$$(b_1 \vee u_1) \wedge (b_2 \vee u_2) = (b_1 \wedge b_2) \vee (b_1 \wedge u_2) \vee (b_2 \wedge u_1) \vee (u_1 \wedge u_2) = (b_1 \wedge b_2). \quad (12)$$

After this preparation, for each i let

$$h_{i,1} = g_i^- \wedge (b_1 \vee u_1), \quad h_{i,2} = g_i^- \wedge (b_2 \vee u_2), \quad \sigma_i = (g_i, \{h_{i,1}, h_{i,2}\}, \text{sc-dim } g_i)$$

Using (11) we get

$$\begin{aligned} h_{i,1} \vee h_{i,2} &= g_i^- \wedge (b_1 \vee u_1 \vee b_2 \vee u_2) \\ &= g_i^- \wedge \left[b_1 \vee b_2 \vee \left(\bigvee_{j \leq n} g_j^- - (b_1 \vee b_2) \right) \right] \\ &= g_i^- \wedge \bigvee_{j \leq n} g_j^- = g_i^-. \end{aligned}$$

So each σ_i is an SC-signature in L_0 . In particular Proposition 6.4.2 gives an SC-primitive extension $L_1 = L_0 \langle a_{1,1}, a_{1,2} \rangle$ with SC-signature σ_1 in L_0 . By Proposition 6.4.1, $\mathcal{I}(L_1) = (\mathcal{I}(L_0) \setminus \{a_1\}) \cup \{a_{1,1}, a_{1,2}\}$. In particular $g_2 \in \mathcal{I}(L_1)$, hence σ_2 is still an SC-signature in L_1 . Repeating the construction n times (note that $a \neq \mathbf{0}$ ensures that $n \geq 1$) gives a chain of \mathcal{L}_{SC} -extensions $(L_i)_{i \leq n}$ and for each $i > 0$, an SC-primitive tuple $(a_{i,1}, a_{i,2})$ generating L_i over L_{i-1} with signature σ_i in L_{i-1} . Each $a_i = a_{i,1} \vee a_{i,2}$ and by Proposition 6.4.1

$$\mathcal{I}(L_n) = (\mathcal{I}(L_0) \setminus \{a_1, \dots, a_n\}) \cup \{a_{1,1}, a_{1,2}, \dots, a_{n,1}, a_{n,2}\} \quad (13)$$

so $a_{1,1}, a_{1,2}, \dots, a_{n,1}, a_{n,2}$ are the \vee -irreducible components of a in L_n . Moreover every $c \in \mathcal{I}(L)$ such that $c < a_{i,k}$ for some i, k must belong to L_0 , hence the predecessor of $a_{i,k}$ is the same in

every L_j and belongs to L_0 . We can then denote it $a_{i,k}^-$ without ambiguity, and by construction we have

$$a_{i,k}^- = a_{i,k} \wedge g(a_i, L_{i-1}) = a_{i,k} \wedge g_i = h_{i,k}. \quad (14)$$

Let $a_1 = \mathbb{W}_{i \leq n} a_{i,1}$, $a_2 = \mathbb{W}_{i \leq n} a_{i,2}$, $h_1 = \mathbb{W}_{i \leq n} h_{i,1}$ and $h_2 = \mathbb{W}_{i \leq n} h_{i,2}$. We are going to check that a_1, a_2 split a along b_1, b_2 . Both of them are non-zero and since the $a_{i,k}$'s are the \vee -irreducible components of a we have $a - a_1 = a_2$, $a - a_2 = a_1$. Each $a_{i,1} \geq h_{i,1}$ by construction, hence $a_1 \geq h_1$ and symmetrically $a_2 \geq h_2$. Moreover for $k \in \{1, 2\}$

$$h_k = \mathbb{W}_{i \leq n} h_{i,k} \geq \mathbb{W}_{i \leq n} g_i^- \wedge b_k = b_k$$

where the last equality comes from (10), so $a_k \geq b_k$. It remains to check that $a_1 \wedge a_2 = b_1 \wedge b_2$.

For $i \neq j$, $a_{i,1}$ and $a_{j,2}$ are mutually incomparable hence by (14)

$$a_{i,1} \wedge a_{j,2} = a_{i,1}^- \wedge a_{j,2}^- = h_{i,1} \wedge h_{j,2}.$$

On the other hand $a_{i,1} \wedge a_{i,2} = h_{i,1} \wedge h_{i,2}$ by construction. The conclusion follows, using (12).

$$\begin{aligned} a_1 \wedge a_2 &= \mathbb{W}_{i,j} a_{i,1} \wedge a_{j,2} = \mathbb{W}_{i,j} h_{i,1} \wedge h_{j,2} \\ &= \mathbb{W}_{i,j} [g_i^- \wedge (b_1 \vee u_1)] \wedge [g_j^- \wedge (b_2 \vee u_2)] \\ &= \mathbb{W}_{i,j} (g_i^- \wedge g_j^-) \wedge [(b_1 \vee u_1) \wedge (b_2 \vee u_2)] \\ &= \left(\mathbb{W}_i g_i^- \right) \wedge \left(\mathbb{W}_j g_j^- \right) \wedge [(b_1 \vee u_1) \wedge (b_2 \vee u_2)] \\ &= (b_1 \vee u_1) \wedge (b_2 \vee u_2) = b_1 \wedge b_2. \end{aligned}$$

■

Theorem 7.3 *The theory of super d -scaled lattices is the model-completion of the theory of d -subscaled lattices*

Proof: By standard model-theoretic arguments it suffices to prove that every existentially closed d -subscaled lattice is super d -scaled, and that for every super d -scaled lattice \hat{L} , every finitely generated d -subscaled lattice L and every common \mathcal{L}_{SC} -substructure L_0 , there is an embedding of L into \hat{L} over L_0 .

Let L be an existentially closed d -subscaled lattice, and L_0 a finitely generated substructure. By Theorem 4.1, L_0 is finite. By Proposition 5.3, L_0 \mathcal{L}_{SC} -embeds the d -scaled lattice $L_{\text{def}}(X)$ of some special linear set X , which is in particular a Catenary lattice. By the model-theoretic compactness Theorem it follows that L is catenary. Similarly Theorem 4.1, Lemma 7.2 and the model-theoretic compactness Theorem prove that L has the Splitting property, hence L is super d -scaled.

Conversely assume that L is a super d -scaled lattice, L is a finitely generated d -subscaled lattice and L_0 common \mathcal{L}_{SC} -substructure. By Theorem 4.1 and Proposition 6.4.3a, we are reduced to the case where L is a primitive extension of L_0 . Let $\sigma = (g, \{h_1, h_2\}, q)$ be its SC-signature. By Proposition 6.4.2 it suffices to find a $x_1, x_2 \in \hat{L}$ such that (x_1, x_2) is SC-primitive over L_0 and $\sigma_{\text{SC}}(x_1, x_2) = \sigma$. We distinguish two cases, and let g^- denotes the predecessor of g in L_0 .

Case 1: $\text{sc-dim } h_1 < q < \text{sc-dim } g$ and $h_1 = h_2 < g$. Let $p = \text{sc-dim } g$ and $r = \text{sc-dim } h_1$. Let $y_1, y_2 \in \hat{L}$ which split g along h_1, g^- . For $0 \leq i \leq d$, either $i < q$ or $C^i(h_1) = \mathbf{0}$ (because $\text{sc-dim } h_1 = r < q$), hence $\text{sc-dim } C^i(h_1) < q < \text{sc-dim } g$. The Catenarity property then applies to $C^i(h_1) \leq y_1$ and gives $x_i \in \hat{L}$ such that $C^i(h_1) \leq x_i \leq y_1$ and x_i has pure dimension q . Let

$x = \bigvee_{0 \leq i \leq d} x_i$, by construction $h_1 = \bigvee_{i \leq d} C^i(h_1) \leq x \leq y_1$ and x has pure dimension q . In particular

$$h_1 \leq x \wedge g^- \leq y_1 \wedge y_2 = h_1 \wedge g^- = h_1$$

hence $x \wedge g^- = h_1 \in L_0$. Moreover $x \wedge g^- = h_1 \ll x$ because $\dim h_1 < q$ and x has pure dimension q . Finally $x \ll g$ because $\dim x_i = q < p$ and g has pure dimension p . Altogether this proves that (x, x) is an SC-primitive tuple over L_0 with SC-signature σ .

Case 2: $q = \text{sc-dim } g$ and $h_1 \vee h_2 = g^-$. Let $y_1, y_2 \in \hat{L}$ which split g along h_1, h_2 . By construction $y_1 \vee y_2 = g$, and since g has pure SC-dimension q so does each y_i . In addition $y_1 \wedge y_2 = h_1 \wedge h_2 \in L_0$. Moreover

$$y_1 \wedge h_2 \leq y_1 \wedge y_2 = h_1 \wedge h_2$$

hence $y_1 \wedge (h_1 \vee h_2) = h_1 \vee (y_1 \wedge h_2) = h_1$. Since $h_1 \vee h_2 = g^-$ it follows that $y_1 \wedge g^- = h_1 \in L_0$, and symmetrically $y_2 \wedge g^- = h_2 \in L_0$. So (y_1, y_2) is an SC-primitive tuple over L_0 with SC-signature σ .

■

Remark 7.4 The proof of Theorem 7.3 shows that if L_0 be a finite \mathcal{L}_{SC} -substructure of a super scaled lattice \hat{L} , then every signature σ in L_0 is the signature of an SC-primitive extension of L_0 in \hat{L} .

The completions of the theory of super d -scaled lattices are easy to classify. Let us say that a d -subscaled lattice is **prime** if it does not contain any proper d -subscaled lattice, or equivalently if it is generated by the empty set. Every prime d -subscaled lattice is finite. By Corollary 4.2 there exists finitely many prime d -subscaled lattices up to isomorphism.

Corollary 7.5 *The theory of super d -scaled lattices containing (a copy of) a given prime d -subscaled lattice is \aleph_0 -categorical, hence complete. It is also recursively axiomatizable, hence decidable. Since every completion of the theory of super d -scaled lattices is of that kind, the theory of super d -scaled lattices is decidable.*

Proof: Let L, L' be any two countable super d -scaled lattices containing isomorphic prime d -subscaled lattices L_0 and L'_0 . By Remark 7.4 any partial isomorphism between L and L' , extending the given isomorphism between L_0 and L'_0 , can be extended by va-et-vient. This proves the first statement. The other ones are immediate consequences.

■

8 Atomic scaled lattices

Every super scaled lattice is atomless because of the Splitting Property, hence none of the geometric scaled lattice in $\text{SC}_{\text{def}}(\mathcal{K}, d)$, $\text{SC}_{\text{Zar}}(K, d)$, $\text{SC}_{\text{lin}}(K, d)$ can be super scaled. In order to apply our study to some of them we introduce now a variant of subscaled lattices intended to protect atoms against splitting.

Let $\mathcal{L}_{\text{ASC}} = \mathcal{L}_{\text{SC}} \cup \{\text{At}_k\}_{k \in \mathbb{N}^*}$ with each At_k a new unary predicate symbol. For any \mathcal{L}_{ASC} -structure L we denote by $\text{At}_k(L)$ the set of elements a in L such that $L \models \text{At}_k(a)$, and we let $\text{At}_0(L) = L \setminus \bigcup_{k > 0} \text{At}_k(L)$. We call L an **ASC-lattice** if its \mathcal{L}_{SC} -reduct is a scaled lattice and if satisfies the following condition.

ASC₀ : $(\forall k > 0), a \in \text{At}_k(L)$ if and only if a is the join of exactly k atoms in L .

Remark 8.1 This condition can be expressed by $\forall\exists$ formulas in \mathcal{L}_{ASC} by saying first that $\text{At}_1(L)$ is the set of atoms of L , and then that $\text{At}_k(L)$ is the set of elements of L which are the join of exactly k elements of $\text{At}_1(L)$.

Every ASC-lattice obviously satisfies also the following schemes (for $k, l > 0$) of universal axioms:

$$\text{ASC}_1 : (\forall k \neq l), \quad \forall a, \text{At}_k(a) \rightarrow \neg \text{At}_l(a)$$

$$\text{ASC}_2 : (\forall k), \quad \forall a, a_0, \dots, a_{2^k}, \quad \text{At}_k(a) \longrightarrow \left[\bigwedge_{0 \leq i \leq 2^k} (a_i \leq a) \longrightarrow \bigvee_{0 \leq i < j \leq 2^k} (a_i = a_j) \right] \bigwedge \text{sc-dim } a = 0$$

$$\text{ASC}_3 : (\forall k), \quad \forall a, a_1, a_2, \quad \left[(a = a_1 \vee a_2) \bigwedge (a_1 \wedge a_2 = \mathbf{0}) \bigwedge (a_1 \neq \mathbf{0}) \bigwedge (a_2 \neq \mathbf{0}) \right] \longrightarrow \left[\text{At}_k(a) \longleftrightarrow \bigvee_{0 < l < k} (\text{At}_l(a_1) \bigwedge \text{At}_{k-l}(a_2)) \right]$$

We call **sub-ASC-lattices** the \mathcal{L}_{ASC} -structures L whose \mathcal{L}_{SC} -reduct is a subscaled lattice and which satisfy ASC_1 to ASC_3 (but not necessarily ASC_0).

ASC_1 obviously means that $(\text{At}_k(L))_{k \in \mathbf{N}}$ is a partition of L . For any $a \in L$ we then define $\text{asc}(a)$ as the unique $k \in \mathbf{N}$ such that $a \in \text{At}_k(L)$.

ASC_2 says that if $\text{asc}(a) = k > 0$ then $L(a)$ has at most 2^k element and $\text{sc-dim}(a) = 0$. Then $\text{dim}(a) = 0$ by SC_{11} hence $L(a)$ is a co-Heyting algebra with dimension 0, that is a Boolean algebra. So ASC_2 actually says that $\text{sc-dim } a = 0$ and $L(a)$ is a Boolean algebra with n atoms for some non zero $n \leq k$. In particular every $a \in \text{At}_1(L)$ is an atom of L .

ASC_3 says that if a is the join of two non-zero disjoint elements a_1, a_2 then $\text{asc}(a)$ is non-zero if and only if $\text{asc}(a_1)$ and $\text{asc}(a_2)$ are non-zero, in which case $\text{asc}(a) = \text{asc}(a_1) + \text{asc}(a_2)$. By a straightforward induction this extends to any decomposition of a as the join of finitely many pairwise disjoint elements. In view of ASC_2 it then says that $\text{asc}(a) > 0$ if and only if a is the join of finitely many atoms a_1, \dots, a_n of L such that each $\text{asc}(a_i) > 0$, in which case $\text{asc}(a) = \sum_{1 \leq i \leq n} \text{asc}(a_i)$.

Remark 8.2 It follows immediately that an \mathcal{L}_{SC} -embedding of sub-ASC-lattices $\varphi: L \rightarrow L'$ is an \mathcal{L}_{ASC} -embedding if and only if $\text{asc}(a) = \text{asc}(\varphi(a))$ for every atom $a \in L$.

Remark 8.3 Obviously every finitely generated substructure of a sub-ASC-lattices is finite by the Local Finiteness Theorem 4.1 because \mathcal{L}_{ASC} expands \mathcal{L}_{SC} only by relational symbols.

Every scaled lattice L admits a unique structure of ASC-lattice which is an expansion by definition of its lattice structure. We denote by L^{At} this expansion of L .

Proposition 8.4 (Linear representation) *Let K be an infinite field and L_0 be a finite sub-ASC-lattice. For every integer $N \geq 0$ there exists a special linear set X_N over K and an $\mathcal{L}_{\text{ASC}}^*$ -embedding $\varphi_N: L_0 \rightarrow L_{\text{lin}}^{\text{At}}(K^m)$ such that for every atom a of L_0 :*

- If $\text{asc}(a) > 0$ then $\text{asc}(\varphi_N(a)) = \text{asc}(a)$.
- If $\text{asc}(a) = 0$ then $\text{asc}(\varphi_N(a)) \geq N$.

Proof: By induction on lexicographically ordered tuples of integers (r, s) we prove that the result is true for every finite sub-ASC-lattice L_0 having r \vee -irreducible elements, s of which have the same sc-dimension as L_0 .

If $r = 0$ then $s = 0$ and the unique embedding of $L_0 = \{\mathbf{0}\}$ into $L_{\text{lin}}^{\text{At}}(P)$, for an arbitrary point P of K , has the required property. So let us assume that $r \geq 1$ and that the result is proved for every $(r', s') < (r, s)$. Let $d = \text{sc-dim } L$ and a_1, \dots, a_r be the elements of $\mathcal{I}(L_0)$ ordered by increasing sc-dimension, so that $\text{sc-dim } a_r = d \geq 0$.

Case 1: $d = 0$. Then L is a boolean algebra and a_1, \dots, a_r are its atoms. Let A_1, \dots, A_r be pairwise disjoint subsets of K such that:

- If $\text{asc}(a_i) > 0$ then A_i has $\text{asc}(a_i)$ elements, so $\text{asc}(A_i) = \text{asc}(a_i)$.
- If $\text{asc}(a_i) = 0$ then A_i has N elements, so $\text{asc}(A_i) = N$.

Let X be the union of all these A_i 's. Clearly the map φ which maps each a_i to A_i extends uniquely to an \mathcal{L}_{SC} -embedding of L_0 into $L_{\text{lin}}^{\text{At}}(X)$ which has the required properties.

Case 2: $d > 0$. The upper semi-lattice L_0^- generated by a_1, \dots, a_{r-1} is an $\mathcal{L}_{\text{ASC}}^*$ -substructure of L_0 to which the induction hypothesis applies. This gives for some integer m a special linear set $B \subseteq K^m$ over K and an \mathcal{L}_{SC} -embedding $\psi: L_0^- \rightarrow L_{\text{lin}}^{\text{At}}(B)$ having the required properties. Let $C = \varphi(\mathbf{1}_{L_0^-} \wedge a_r)$ and $n = \text{sc-dim } a_r$. Proposition 5.2 gives a special linear set $A \subseteq K^{m+n}$ such that $A \cap B = C$. One can extend ψ to an \mathcal{L}_{SC} -embedding φ of L_0 into $L_{\text{lin}}^{\text{At}}(A \cup B)$ exactly like in the proof of Proposition 5.3. Then φ inherits from ψ the required properties because all the atoms of L already belong to L_0^- .

■

Let $\text{ASC}_{\text{Zar}}(K, d)$, $\text{ASC}_{\text{lin}}(K, d)$, $\text{ASC}_{\text{def}}(K, d)$ denote the class of all ASC-lattices L^{At} for L ranging over $\text{SC}_{\text{Zar}}(K, d)$, $\text{SC}_{\text{lin}}(K, d)$, $\text{SC}_{\text{def}}(K, d)$ respectively.

Corollary 8.5 *For every integer $d \geq 0$, the universal theories of $\text{ASC}_{\text{def}}(K, d)$ (resp. of $\text{ASC}_{\text{Zar}}(K, d)$ or $\text{ASC}_{\text{lin}}(K, d)$) is the same for every o -minimal or P -minimal expansion of a field K (resp. every infinite field K). This is the theory of sub-ASC-lattices.*

Proof: Since $\text{ASC}_{\text{lin}}(K, d)$ is contained in the other classes, all of which are contained in the class of ASC-lattices, it suffices to prove that conversely every sub-ASC-lattice \mathcal{L}_{ASC} -embeds into an ultraproduct of elements of $\text{ASC}_{\text{lin}}(K, d)$. By the model-theoretic compactness theorem, it suffices to prove it for any finitely generated sub-ASC-lattice L_0 .

By Theorem 4.1, L_0 is finite. For any integer $N \geq 0$ let $\varphi_N : L_0 \rightarrow L_{\text{lin}}^{\text{At}}(X_N)$ be an \mathcal{L}_{SC} -embedding given by Proposition 8.4. Let \mathcal{U} be a non principal ultrafilter in the Boolean algebra of subsets of \mathbf{N} , and consider the ultraproduct $L = \prod_{N \in \mathbf{N}} L_{\text{lin}}^{\text{At}}(X_N) / \mathcal{U}$. Then $\varphi = \prod_{N \in \mathbf{N}} \varphi_N / \mathcal{U}$ is an \mathcal{L}_{SC} -embedding of L_0 into the L . In order to prove that it is an \mathcal{L}_{ASC} -embedding, by Remark 8.2 it remains check that for every atom a of L_0 , $\text{asc}(\varphi(a)) = \text{asc}(a)$. So let a be an atom of L_0 and $k = \text{asc}(a)$.

If $k > 0$ then for every $N \geq k$, $L_{\text{lin}}^{\text{At}}(X_N) \models \text{At}_k(\varphi_N(a))$ by construction. So $L \models \text{At}_k(\varphi(a))$, that is $\text{asc}(\varphi(a)) = k$.

If $k = 0$, let l be any strictly positive integer. For every $N \geq l$, $L_{\text{lin}}^{\text{At}}(X_N) \models \text{At}_N(\varphi_N(a))$ by construction, hence $L_{\text{lin}}^{\text{At}}(X_N) \not\models \text{At}_l(\varphi_N(a))$. So $L \not\models \text{At}_l(\varphi(a))$, and this being true for every $l > 0$ it follows that $\text{asc}(\varphi(a)) = 0$.

■

9 Model-completion of atomic scaled lattices

Let us call **super ASC-lattices** those ASC-lattices which satisfy the following axioms, all of which are axiomatizable by $\forall\exists$ -formulas in \mathcal{L}_{ASC} . We are going to show that its theory is the model-completion of the theory of sub-ASC-lattices of dimension at most d (resp. exactly d).

Atomicity: Every element x is the least upper bound of the set of atoms smaller than x .

Catenarity: For every non-negative integers $r \leq q \leq p$ and every elements $c \leq a \neq \mathbf{0}$, if c is r -sc-pure and a is p -sc-pure then there exists a non-zero q -sc-pure element b such that $c \leq b \leq a$.

ASC-Splitting: For every b_1, b_2, a , if $b_1 \vee b_2 \ll a \neq \mathbf{0}$ and $C^0(a) = \mathbf{0}$ there exists non-zero elements $a_1 \geq b_1$ and $a_2 \geq b_2$ such that:

$$\begin{cases} a_1 = a - a_2 \\ a_2 = a - a_1 \\ a_1 \wedge a_2 = b_1 \wedge b_2 \end{cases}$$

Remark 9.1 An immediate consequence of the atomicity axiom is that for every elements x, y in a super ASC-lattice L such that $y < x$ and $\text{sc-dim}(x - y) \geq 1$, there are infinitely many atoms $a \in L$ such that $a \leq x$ and $a \wedge y = \mathbf{0}$. Indeed let A be the set of atoms $a \in L$ such that $a \leq x - y$, and B the subset of those a such that $a \wedge y \neq \mathbf{0}$. Assume for a contradiction that B is finite and let $b = \bigvee_{a \in B} a$. Note that $b \leq y$ and $\text{sc-dim } b = \dim b = 0$. Then by the Atomicity axiom

$$x - y = \bigvee_{a \in A} a \leq y \vee b, \text{ hence } x - y \leq (y \vee b) - y = b - y \leq b.$$

This implies that $\text{sc-dim}(x - y) \leq \text{sc-dim } b = 0$, a contradiction.

Primitive tuples and sc-primitive extensions are defined for sub-ASC-lattices exactly like for subscaled lattices.

We define **ASC-signatures** in a finite sub-ASC-lattice L_0 as triples (g, H, q) with H a set of non-necessarily distinct pairs $(h_1, k_1), (h_2, k_2)$ in $L_0 \times \mathbf{N}$ such that $(g, \{h_1, h_2\}, q)$ is an SC-signature in the \mathcal{L}_{SC} -reduct of L_0 and (see Example 9.2):

1. If $q < \text{sc-dim } g$ then $k_1 = k_2$.
Note that $h_1 = h_2$ in this case because $(g, \{h_1, h_2\}, q)$ is a signature.
2. If $q \neq 0$ then $k_1 = k_2 = 0$.
3. If $k_1 = 0$ or $k_2 = 0$ then $\text{asc}(g) = 0$.
4. If $k_1 \neq 0, k_2 \neq 0$ and $\text{sc-dim } g = 0$ then $\text{asc}(g) = k_1 + k_2$.

Example 9.2 Let L_0 be a finite sub-ASC-lattice, and L an \mathcal{L}_{SC} -extension of L_0 generated by a (necessarily unique) SC-primitive tuple (x_1, x_2) . Let $(g, \{h_1, h_2\}, q)$ be the SC-signature of L in L_0 and $k_i = \text{asc}(x_i)$. Then $(g, \{(h_1, k_1), (h_2, k_2)\}, q)$ is easily an ASC-signature in L_0 . We call it the **ASC-signature** of L and of (x_1, x_2) in L_0 .

The same argument as in Proposition 6.4.2 shows (using Remark 8.2) that two SC-primitive extensions of a finite sub-ASC-lattice L_0 are \mathcal{L}_{ASC} -isomorphic over L_0 if and only if they have the same ASC-signature in L_0 .

Lemma 9.3 *Let L_0 be a finite \mathcal{L}_{ASC} -substructure of a super ASC-lattice \hat{L} . Let $\sigma_{\text{At}} = (g, q, \{(h_1, k_1), (h_2, k_2)\})$ be an ASC-signature in L_0 . Assume that $q \neq 0$ or $k_1 k_2 \neq 0$. Otherwise assume that \hat{L} is \aleph_0 -saturated. Then there exists a primitive tuple $(x_1, x_2) \in \hat{L}$ over L_0 whose signature is σ_{At} .*

Proof: Let $\sigma = (g, \{h_1, h_2\}, q)$. This is an SC-signature in L_0 (more precisely in its \mathcal{L}_{SC} -reduct).

Case 1: $\text{sc-dim } g \geq 1$ and $q \geq 1$. Then $C^0(g) = 0$ and by definition of ASC-signatures $k_1 = k_2 = 0$. By Remark 7.4 there is an SC-primitive tuple (x_1, x_2) in \hat{L} with signature σ in L_0 . Moreover each $\text{asc } x_i = 0$ (because $\text{sc-dim } x_i = p \geq 1$) and each $k_i = 0$, so the ASC-signature of (x_1, x_2) is σ_{At} .

Case 2: $\text{sc-dim } g \geq 1$ and $q = 0$. Then $C^0(g) = 0$ again and since $\text{sc-dim}(h_1 \vee h_2) < q = 0$ by definition of SC-signatures we get that $h_1 = h_2 = \mathbf{0}$. Finally $k_1 = k_2$ by definition of ASC-signatures since $q = 0 < \text{sc-dim } g$. By Remark 9.1 there are infinitely many atoms z in \hat{L} such that $z \leq g$ and $z \wedge g^- = \mathbf{0}$. If $k_1 > 0$ let x be the join of k_1 such atoms of \hat{L} . Otherwise \hat{L} is \aleph_0 -saturated by assumption hence it contains an element $x \leq g$ of dimension 0 such that $x \wedge g^- = \mathbf{0}$ and $\hat{L}(x)$ has infinitely many atoms. By the Atomicity Property $\text{asc}(x) = 0$. So in both cases (x, x) is an SC-primitive tuple over L_0 with ASC-signature σ_{At} .

Case 3: $\text{sc-dim } g = 0$. Then $q = 0$, g is an atom of L_0 and $h_1 = h_2 = \mathbf{0}$. In each of the two remaining sub-cases, we build a tuple (x_1, x_2) and leave as an exercise to check that (x_1, x_2) is SC-primitive over L_0 with ASC-signature σ_{At} .

If k_1 and k_2 are non-zero then $\text{asc}(g) = k_1 + k_2$ hence $\hat{L}(g)$ contains $k_1 + k_2$ atoms. Let x_1 be the join of k_1 of them and x_2 be the join of the others.

Otherwise, by symmetry we can assume that $k_1 = 0$. Then $\text{asc}(g) = 0$ by definition of ASC-signatures so $\hat{L}(g)$ contains infinitely many atoms. By \aleph_0 -saturation it follows that \hat{L} contains an element x lower than g such that both $L(x)$ and $L(g - x)$ contain infinitely many atoms, hence $\text{asc}(x) = \text{asc}(g - x) = 0$. If $k_2 = 0$ let $(x_1, x_2) = (x, g - x)$. Otherwise let x_2 be the join of k_2 atoms in $\hat{L}(g)$ and let $x_1 = g - x_2$.

■

Theorem 9.4 *The theory of super ASC-lattices of sc-dimension at most d (resp. exactly d) is the model-completion of the theory of ASC-lattices of dimension at most d (resp. exactly d). It admits \aleph_0 completions, each of which is decidable, and it is decidable.*

Proof: We only sketch the proof of the first statement as it essentially the same as for Theorem 7.3.

On one hand, given a finite sub-ASC-lattice L_0 , we can embed it in an extension satisfying the Atomicity and Catenary Property by Proposition 8.4, and the ASC-Splitting Property by means of Lemma 7.2 applied to any $a, b_1, b_2 \in L_0$ such that $b_1 \vee b_2 \ll a \neq \mathbf{0}$ and $C^0(a) = \mathbf{0}$ (note that this last assumption ensures that the extension built in Lemma 7.2 is an \mathcal{L}_{ASC} -extension). That every existentially closed sub-ASC-lattice is a super ASC-lattice then follows, by the model-theoretic compactness theorem.

On the other hand, given an \aleph_0 -saturated super ASC-lattice \hat{L} , a finite \mathcal{L}_{ASC} -substructure L_0 and a finite extension L of L_0 , we reduce to the case where L is SC-primitive and let σ be its ASC-signature in L_0 . Lemma 9.3 gives an SC-primitive extension L_1 of L_0 in \hat{L} with the same signature in L_0 , hence an embedding of L_1 into \hat{L} over L_0 (which maps L to L_1).

This proves the first statement. The last one follows because there are finitely many subscaled lattices of dimension at most d (resp. exactly d) and on each of them \aleph_0 different structures of

sub-ASC-lattices. So the completions of the theory of super ASC-lattices, which are determined by their prime model, can be recursively enumerated.

■

We say that a sub-ASC-lattice L is **standard** if every element of sc-dimension 0 belongs to some $\text{At}_k(L)$ for some $k > 0$. The existence of standard super ASC-lattices (see Section 10) and non-standard super ASC-lattices (by the model theoretic compactness theorem) implies that the theory of super ASC-lattices containing a given prime sub-ASC-lattice is not \aleph_0 -categorical, contrary to what happens for super scaled lattice. However we can recover \aleph_0 -categoricity by restricting to standard models.

Proposition 9.5 *Let L_1, L_2 be two standard countable super ASC-lattices. Then every \mathcal{L}_{ASC} -isomorphism from a finite sub-ASC-lattice $L_{1,0} \subset L_1$ to a sub-ASC-lattice $L_{2,0} \subset L_2$ extends to an \mathcal{L}_{ASC} -isomorphism from L_1 to L_2 . In particular L_1 and L_2 are isomorphic if and only if their prime \mathcal{L}_{ASC} -substructures (those generated by the empty set) are isomorphic.*

Proof: Let φ be an \mathcal{L}_{ASC} -isomorphism from $L_{1,0}$ to $L_{2,0}$. Pick any element $x \in L_1 \setminus L_{1,0}$. The subscaled lattice generated in L_1 by $L_{1,0} \cup \{x\}$ (more precisely their \mathcal{L}_{SC} -reducts) is finite hence by Proposition 6.4.3 there is a chain $L_{1,0} \subset L_{1,1} \subset \dots \subset L_{1,r}$ of SC-primitive extensions of subscaled lattices such that $L_{1,0} \cup \{x\} \subseteq L_{1,r}$. Endow each $L_{1,i}$ with the \mathcal{L}_{ASC} -structure induced by L_1 . It suffices to prove that φ extends to an \mathcal{L}_{ASC} -embedding $\varphi_1 : L_{1,1} \rightarrow L_2$. Indeed, repeating the argument will give an \mathcal{L}_{ASC} -embedding $\varphi_r : L_{1,r} \rightarrow L_2$ extending φ , and by symmetry the conclusion will then follow by Fraïssé's va-et-vient.

Identifying $L_{1,0}$ with its image by φ we can replace $L_{1,0}$ and $L_{2,0}$ by a common \mathcal{L}_{ASC} -structure L_0 of L_1 and L_2 . Now $L_{1,1}$ is generated over L_0 by an SC-primitive tuple (x_1, x_2) with signature $\sigma_{\text{At}} = (g, \{(h_1, k_1), (h_2, k_2)\}, q)$. In particular $q = \text{sc-dim } x_i$ and $k_i = \text{asc}(x_i)$ for $i = 1, 2$. If $q = 0$ then for each i , $\text{sc-dim } x_i = 0$ hence $k_i > 0$ because L_1 is standard. With other words $q \neq 0$ or $k_1 k_2 \neq 0$ hence Lemma 9.3 gives an sc-primitive tuple (y_1, y_2) in L_2 with signature σ_{At} . Let $L_{2,1}$ be the asc-substructure of L_2 generated by $L_{1,1} \cup \{y_1, y_2\}$. By Proposition 6.4.2 φ extends to an \mathcal{L}_{SC} -isomorphism φ_1 from $L_{1,1}$ to $L_{2,1}$ which maps each x_i to y_i . By construction $\text{asc}(x_i) = \text{asc}(y_i)$, and by Proposition 6.4.1 φ_1 is the identity map on L_0 , so $\text{asc}(\varphi_1(z)) = \text{asc}(z)$ for every $z \in \mathcal{I}(L_{1,1})$. Hence φ_1 is an \mathcal{L}_{ASC} -isomorphism by Remark 8.2, which proves the result.

■

10 Applications to lattices of p -adic semi-algebraic sets

In all this section K denotes a fixed p -adically closed field. For every semi-algebraic set X contained in K^m we let $L(X)$ denote the lattice of semi-algebraic subsets of X closed in X , endowed with its natural structure of ASC-lattice. Note that every $A \in L(X)$ of dimension 0 is finite, hence $L(X)$ is *standard*.

As already mentioned in the introduction, the results of the previous section lead us to conjecture in [Dar04] and finally to prove in [Dar17] the following result.

Theorem 10.1 (Theorem 3.4 in [Dar17]) *Let X be a non-empty semi-algebraic subset of K^m without isolated points. Assume that X is open in its topological closure \overline{X} and let Y_1, \dots, Y_s be a collection of closed semi-algebraic subsets of $\partial X = \overline{X} \setminus X$ such that $Y_1 \cup \dots \cup Y_s = \partial X$. Then there is a partition of X in non-empty semi-algebraic sets X_1, \dots, X_s such that $\partial X_i = Y_i$ for $1 \leq i \leq s$.*

We can now combine this theorem with the results of Section 10 in order to get the following applications.

Theorem 10.2 *Let X be any semi-algebraic subset of K^m . Then $L(X)$ is a super ASC-lattice. In particular its complete theory is decidable and eliminates the quantifiers in \mathcal{L}_{ASC} .*

Proof: By construction $L(X)$ is an ASC-lattice satisfying the Atomicity property. The Catenarity Property is well known (it follows for example from Denef's Cell Decomposition) hence we only have to check the Splitting Property. So let $A, B_1, B_2 \in L(X)$ such that $B_1 \cup B_2 \ll A$ and A has no isolated point.

The same holds true for their closures in K^m , denoted $\bar{A}, \bar{B}_1, \bar{B}_2$. Apply Theorem 10.1 to $X = \bar{A} \setminus (\bar{B}_1 \cup \bar{B}_2)$, $Y_1 = \bar{B}_1$ and $Y_2 = \bar{B}_2$. It gives a partition of X in non-empty semi-algebraic sets X_1, X_2 whose frontiers are respectively \bar{B}_1, \bar{B}_2 . Then $\bar{X}_1 \cup \bar{X}_2 = \bar{A}$, $\bar{X}_1 \cap \bar{X}_2 = \bar{B}_1 \cap \bar{B}_2$ and each $\bar{X}_i = X_i \cup \bar{B}_i$. Let $A_1 = \bar{X}_1 \cap A = (X_1 \cap A) \cup B_1$ and define A_2 accordingly. We have to check that A_1, A_2 split A along B_1, B_2 .

A is dense in $\bar{A} = \bar{X}$ and $X_1 = \bar{X} \setminus (X_2 \cup \bar{B}_1 \cup \bar{B}_2) = \bar{X} \setminus (\bar{X}_2 \cup \bar{B}_1)$ is open in \bar{X} , hence $A \cap X_1$ is dense in X_1 . In particular $A \cap X_1 \neq \emptyset$, and symmetrically $A \cap X_2 \neq \emptyset$. Clearly $A_1 \cup A_2 = A$, $A_1 \cap A_2 = B_1 \cap B_2$ and each $A_i \supseteq B_i$ by construction. So it only remains to check that $A - A_1 = A_2$ in $L(X)$, that is that the closure of $A \setminus A_1$ in X (hence in A) is A_2 . Note that $A \setminus A_1 = A \setminus \bar{X}_1$ and

$$\bar{A} \setminus \bar{X}_1 = (X_1 \cup X_2 \cup \bar{B}_1 \cup \bar{B}_2) \setminus (X_1 \cup \bar{B}_1) = X_2 \cup (\bar{B}_2 \setminus \bar{B}_1).$$

In particular $A \setminus A_1 = A \setminus \bar{X}_1 = (\bar{A} \setminus \bar{X}_1) \cap A$ contains $X_2 \cap A$ and is contained in $(X_2 \cup \bar{B}_2) \cap A = \bar{X}_2 \cap A = A_2$. The conclusion will follow, if we can prove that $X_2 \cap A$ is dense in A_2 . Since $A_2 = (X_2 \cap A) \cup B_2$ it suffices to check that $B_2 \subseteq \bar{X}_2 \cap \bar{A}$. But this is clear since $X_2 \cap A$ is dense in X_2 , hence in $\bar{X}_2 = X_2 \cup \bar{B}_2$.

■

Corollary 10.3 *Let F be a q -adically closed field (for some prime q not necessarily equal to p). Let $X \subseteq K^m$ and $Y \subseteq F^n$ be two semi-algebraic sets.*

1. *If $m = n$, $K \preceq F$ and $X = Y \cap K^n$ then $L(X) \preceq L(Y)$.*
2. *$L(X) \equiv L(Y) \iff$ their prime \mathcal{L}_{ASC} -substructures are isomorphic.
In particular $L(K^m) \equiv L(F^n)$ if and only if $m = n$.*
3. *If K and F are countable then $L(X) \equiv L(Y) \iff L(X) \simeq L(Y)$.*

Proof: The two first points follow immediately from Theorem 10.2. Note that $L(K^m) \equiv L(F^m)$ is a special case because their prime sublattice is just the two-element lattice with the same \mathcal{L}_{ASC} -structure, because K^m and F^m both have pure dimension m . The last point follows from Proposition 9.5 since both $L(X)$ and $L(Y)$ are standard and countable.

■

Given a pair of semi-algebraic sets $X \subseteq K^m$ and $Y \subseteq F^n$, we say that a homeomorphism $\psi : X \rightarrow Y$ is **pre-algebraic** if for every semi-algebraic sets $A \subseteq X$ and $B \subseteq Y$ defined over K and F respectively, $\varphi(A)$ and $\varphi^{-1}(B)$ are still semi-algebraic sets defined over K and F . It is obviously sufficient to check this for semi-algebraic sets A, B closed in X, Y respectively. With other words, a bijection $\psi : X \rightarrow Y$ is a pre-algebraic if and only if taking direct images by ψ defines an \mathcal{L}_{ASC} -isomorphism from $L(X)$ to $L(Y)$ (which also ensures that ψ is a homeomorphism).

Remark 10.4 When $K = F$, semi-algebraic homeomorphisms are obviously pre-algebraic but the converse is false. Indeed, applying the next proposition to $X = K$ and $Y = R$, the p -valuation of K , gives a pre-algebraic homeomorphism $\varphi : K \rightarrow R$. If the value group of K is \mathbf{Z} , the p -valuation defines a metric on K and its completion K' is known to be an elementary extension of K . Since its valuation ring R' is compact and K' is not, they cannot be homeomorphic. Hence φ do not extend to a homeomorphism from K' to R' , in particular φ is not semi-algebraic.

Theorem 10.5 *Let K, F be countable p -adically closed fields, and $X \subseteq K^m, Y \subseteq F^n$ be two semi-algebraic sets. Let $L^0(X)$ and $L^0(Y)$ be the prime \mathcal{L}_{ASC} -substructures of $L(X)$ and $L(Y)$ respectively. Then X and Y are pre-algebraically homeomorphic if and only if $L^0(X)$ and $L^0(Y)$ are \mathcal{L}_{ASC} -isomorphic. In particular, any two semi-algebraic sets over K and F with the same pure dimension $d \geq 1$ are pre-algebraically homeomorphic.*

Proof: One direction is obvious: every pre-algebraic homeomorphism $\psi : X \rightarrow Y$ induces an \mathcal{L}_{ASC} -isomorphism from $L(X)$ to $L(Y)$, which maps their respective prime \mathcal{L}_{ASC} -substructures one to each other. Conversely, assume that an \mathcal{L}_{ASC} -isomorphism is given from $L^0(X)$ to $L^0(Y)$. By Proposition 9.5 it extends to an \mathcal{L}_{ASC} -isomorphism $\varphi : L(X) \rightarrow L(Y)$. For every $t \in X$, φ maps $\{t\}$ to an atom $\{t'\}$ of $L(Y)$. Let $\psi(t) = t'$, this defines a bijection $\psi : X \rightarrow Y$ such that $\psi(A) = \varphi(A)$ for every $A \in L(X)$, hence ψ is a pre-algebraic homeomorphism. The last statement follows.

■

References

- [Clu01] Raf Cluckers. Classification of semi-algebraic p -adic sets up to semi-algebraic bijection. *J. Reine Angew. Math.*, 540:105–114, 2001.
- [Dar04] Luck Darnière. Model-completion of scaled lattices. *Prépublications mathématiques d'Angers* 191, Département de mathématiques de l'université d'Angers (France), may 2004. See also arXiv:math/0606792v1.
- [Dar06] Luck Darnière. Model-completion of scaled lattices. arXiv:math/0606792, 2006.
- [Dar17] Luck Darnière. Semi-algebraic triangulation over p -adically closed fields. arXiv:1702.05030, 2017.
- [Den84] Jan Denef. The rationality of the Poincaré series associated to the p -adic points on a variety. *Invent. Math.*, 77(1):1–23, 1984.
- [DJ] Luck Darnière and Markus Junker. Model-completion of varieties of co-Heyting algebras. To appear in the Houston Journal of Mathematics.
- [DJ11] Luck Darnière and Markus Junker. Codimension and pseudometric in co-Heyting algebras. *Algebra Universalis*, 64(3):251–282, 2011.
- [Grz51] Andrzej Grzegorzczak. Undecidability of some topological theories. *Fund. Math.*, 38:137–152, 1951.
- [HM97] Deirdre Haskell and Dugald Macpherson. A version of o-minimality for the p -adics. *J. Symbolic Logic*, 62(4):1075–1092, 1997.
- [Hos67] Tsutomu Hosoi. On intermediate logics. I. *J. Fac. Sci. Univ. Tokyo Sect. I*, 14:293–312, 1967.
- [Joh82] Peter T. Johnstone. *Stone spaces*, volume 3 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 1982.
- [Mac76] Angus Macintyre. On definable subsets of p -adic fields. *J. Symbolic Logic*, 41(3):605–610, 1976.

- [Ono71] Hiroakira Ono. Kripke models and intermediate logics. *Publ. Res. Inst. Math. Sci.*, 6:461–476, 1970/71.
- [SvdD88] Philip Scowcroft and Lou van den Dries. On the structure of semialgebraic sets over p -adic fields. *J. Symbolic Logic*, 53(4):1138–1164, 1988.
- [Tre17] Marcus Tressl. On the strength of some topological lattices. *Contemp. Math.*, 2017.
- [vdD98] Lou van den Dries. *Tame topology and o-minimal structures*, volume 248 of *London Mathematical Society Lecture Note Series*. Cambridge University Press, Cambridge, 1998.