



Post-1980 shifts in the sensitivity of boreal tree growth to North Atlantic Ocean dynamics and seasonal climate

Clémentine Ols, Valerie Trouet, Martin Girardin, Annika Hofgaard, Yves Bergeron, Igor Drobyshev

► To cite this version:

Clémentine Ols, Valerie Trouet, Martin Girardin, Annika Hofgaard, Yves Bergeron, et al.. Post-1980 shifts in the sensitivity of boreal tree growth to North Atlantic Ocean dynamics and seasonal climate. Global and Planetary Change, 2018, 165, pp.1 - 12. 10.1016/j.gloplacha.2018.03.006 . hal-01756849

HAL Id: hal-01756849

<https://hal.science/hal-01756849>

Submitted on 3 Apr 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Post-1980 shifts in the sensitivity of boreal tree growth to North Atlantic Ocean dynamics and seasonal climate

Tree growth responses to North Atlantic Ocean dynamics

Clémentine Ols^{a,*}, Valérie Trouet^b, Martin P. Girardin^c, Annika Hofgaard^d, Yves Bergeron^a & Igor Drobyshev^{a,c}

^a Chaire Industrielle en Aménagement Forestier Durable UQAM-UQAT, Institut de recherche sur les forêts, Université du Québec en Abitibi-Témiscamingue, 445 boul. de l'Université, Rouyn-Noranda, QC J9X 5E4, Canada

^b Laboratory of Tree-Ring Research, University of Arizona, 1215 E. Lowell Street, Tucson, AZ 85721, USA

^c Natural Resources Canada, Canadian Forest Service, Laurentian Forestry Centre, 1055 du P.E.P.S. P.O. Box 10380, Stn. Sainte-Foy, Quebec, QC G1V 4C7, Canada

^d Norwegian Institute for Nature Research, P.O. Box 5685 Torgarden, NO-7485 Trondheim, Norway

^e Southern Swedish Forest Research Centre, Swedish University of Agricultural Sciences, P.O. Box 49, SE-230 53 Alnarp, Sweden; * Corresponding author.

*Corresponding author

clementine.ols88@gmail.com

¹ Present address : Institut National de l'Information Géographique et Forestière, Laboratoire de l'Inventaire Forestier, 14 rue Girardet, 54000 Nancy, France

Highlights

- A significant boreal tree growth response to oceanic and atmospheric indices emerged during the 1980s.
- This was particularly observed in western and central boreal Quebec and in central and northern boreal Sweden.
- The post-1980 sensitivity to large-scale indices synchronized with changes in tree growth responses to local climate.
- Future large-scale dynamics may impact forest growth and carbon sequestration to a greater extent than previously thought.

Abstract

The mid-20th century changes in North Atlantic Ocean dynamics, e.g. slow-down of the Atlantic meridional overturning thermohaline circulation (AMOC), have been considered as early signs of tipping points in the Earth climate system. We hypothesized that these changes have significantly altered boreal forest growth dynamics in northeastern North America (NA) and northern Europe (NE), two areas geographically adjacent to the North Atlantic Ocean. To test our hypothesis, we investigated tree growth responses to seasonal large-scale oceanic and atmospheric indices (the AMOC, North Atlantic Oscillation (NAO), and Arctic Oscillation (AO)) and climate (temperature and precipitation) from 1950 onwards, both at the regional and local levels. We developed a network of 6,876 black spruce (NA) and 14,437 Norway spruce (NE) tree-ring width series, extracted from forest inventory databases. Analyses revealed post-1980 shifts from insignificant to significant tree growth responses to summer oceanic and atmospheric dynamics both in NA (negative responses to NAO and AO indices) and NE (positive response to NAO and AMOC indices). The strength and sign of these responses varied, however, through space with stronger responses in western and central boreal Quebec and in central and northern boreal Sweden, and across scales with stronger responses at the regional level than at the local level. Emerging post-1980 associations with North Atlantic Ocean dynamics synchronized with stronger tree growth responses to local seasonal climate, particularly to winter temperatures. Our results suggest that ongoing and future anomalies in oceanic and atmospheric dynamics may impact forest growth and carbon sequestration to a greater extent than previously thought. Cross-scale differences in responses to North Atlantic Ocean dynamics highlight complex interplays in the effects of local climate and ocean-atmosphere dynamics on tree growth processes and advocate for the use of different spatial scales in climate-growth research to better understand factors controlling tree growth.

Keywords

Climate change, Dendrochronology, Climate-growth interactions, Response functions, Teleconnections, Arctic amplification.

1 Introduction

Terrestrial biomes on both sides of the North Atlantic Ocean are strongly influenced by Arctic and Atlantic oceanic and atmospheric dynamics (D'Arrigo *et al.*, 1993; Ottersen *et al.*, 2001; Girardin *et al.*, 2014). Some mid-20th century changes in the dynamics of the North Atlantic Ocean have been considered as early signs of tipping points in the Earth climate system (Lenton *et al.*, 2008; Lenton, 2011). The Atlantic Meridional Overturning Circulation (AMOC) exhibited an exceptional slow-down in the 1970s (Rahmstorf *et al.*, 2015). The cause of this slow-down is still under debate, but possible explanations include the weakening of the vertical structure of surface waters through the discharge of low-salinity fresh water into the North Atlantic Ocean, due to the disintegration of the Greenland ice sheet and the melting of Canadian Arctic glaciers. A further weakening of the AMOC may possibly lead to a wide-spread cooling and decrease in precipitation in the North Atlantic region (Sgubin *et al.*, 2017), subsequently lowering the productivity of land vegetation both over northeastern North America and northern Europe (Zickfeld *et al.*, 2008; Jackson *et al.*, 2015). Despite increasing research efforts in monitoring climate-change impacts on ecosystems, effects of late 20th century changes in North Atlantic Ocean dynamics on mid- to high-latitude terrestrial ecosystems remain poorly understood.

The dynamics of North Atlantic oceanic and atmospheric circulation, as measured through the AMOC, North Atlantic Oscillation (NAO) and Arctic Oscillation (AO) indices, strongly influence climate variability in northeastern North America (NA) and northern Europe (NE) (Hurrell, 1995; Baldwin & Dunkerton, 1999; Wettstein & Mearns, 2002). NAO and AO indices integrate differences in sea-level pressure between the Iceland Low and the

Azores High (Walker, 1924), with high indices representative of increased west-east air circulation over the North Atlantic. Variability in AMOC, NAO and AO indices affects climate dynamics, both in terms of temperatures and precipitation regimes. Periods of high winter NAO and AO indices are associated with below-average temperatures and more sea ice in NA and a warmer- and wetter-than-average climate in NE. Periods of low winter NAO and AO indices are, in turn, associated with above-average temperatures and less sea ice in NA and a colder- and dryer-than-average climate in NE (Wallace & Gutzler, 1981; Chen & Hellström, 1999). Low AMOC indices induce a wide-spread cooling and decrease of precipitation across the high latitudes of the North Atlantic region (Jackson *et al.*, 2015).

Boreal forests cover most of mid- and high-latitude terrestrial regions of NA and NE and play an important role in terrestrial carbon sequestration and land-atmosphere energy exchange (Betts, 2000; Bala *et al.*, 2007; de Wit *et al.*, 2014). Boreal forests are sensitive to climate change (Gauthier *et al.*, 2015). Despite general warming and lengthening of the growing season at mid- and high-latitudes (Karlsen *et al.*, 2009; IPCC, 2014), tree growth in many boreal regions lost its positive response to rising temperatures during the late-20th century (Briffa *et al.*, 1998). An increasing dependence on soil moisture in the face of the rapid rise in summer temperatures may counterbalance potential positive effects on boreal forest growth of increased atmospheric CO₂ concentrations (Girardin *et al.*, 2016). During the late 20th century, large-scale growth declines (Girardin *et al.*, 2014) and more frequent low growth anomalies (Ols *et al.*, 2016)- in comparison with the early 20th century- have been reported for pristine boreal spruce forests of NA. In coastal NE, climatic changes over the 20th century have triggered shifts from negative significant to non-significant spruce responses to winter precipitation (Solberg *et al.*, 2002). Annual variability in boreal forest tree growth patterns have shown sensitivity to sea ice conditions (Girardin *et al.*, 2014; Drobyshev *et al.*, 2016) and variability in SSTs (Lindholm *et al.*, 2001). All changes in boreal tree growth

patterns and climate-growth interactions listed above may be driven by the dynamics of the North Atlantic Ocean. Understanding current and projected future impacts of North Atlantic Ocean dynamics on boreal forest ecosystems and their carbon sequestration capacity calls for a deeper spatiotemporal analysis of tree growth sensitivity to large-scale oceanic and atmospheric dynamics.

The present study investigates tree growth responses to changes in North Atlantic Ocean dynamics of two widely distributed tree species in the boreal forests of northeastern North America (black spruce) and northern Europe (Norway spruce). We investigated tree-growth sensitivity to seasonal large-scale indices (AMOC, NAO; AO) and seasonal climate (temperature and precipitation) over the second half of the 20th century. We hypothesize that shifts in tree growth sensitivity to large-scale indices and local climate are linked to major changes in North Atlantic Ocean dynamics. This study aims to answer two questions: (i) has boreal tree growth shown sensitivity to North-Atlantic Ocean dynamics? and (ii) does tree growth sensitivity to such dynamics vary through space and time, both within and across NA and NE?

2 Material and methods

2.1 Study areas

We studied two boreal forest dominated areas under the influence of large-scale atmospheric circulation patterns originating in the North Atlantic: the northern boreal biome of the Canadian province of Quebec (50°N-52°N, 58°W-82°W) in NA and the boreal biome of Sweden (59°N-68°N, 12°E-24°E) in NE (Fig. 1a). The selection of the study areas was based on the availability of accurate annually-resolved tree growth measurements acquired from forest inventories.

In northern boreal Quebec, mean annual temperature increases from north to south (-5

to 0.8 °C) and total annual precipitation increases from west to east (550 to 1300 mm), mainly due to winter moisture advection from the North Atlantic Ocean (Gerardin & McKenney, 2001). In boreal Sweden, annual mean temperature increases from north to south (-2 to 6 °C) and annual total precipitation decreases from west to east (900 to 500 mm), mostly because of winter moisture advection from the North Atlantic Ocean that condenses and precipitates over the Scandinavian mountains in the west (Sveriges meteorologiska och hydrologiska institut (SMHI), 2016).

The topography in northern boreal Quebec reveals a gradient from low plains in the west (200-350 m above sea level [a.s.l.]) to hills in the east (400-800 m a.s.l.). In boreal Sweden, the topography varies from high mountains (1500-2000 m a.s.l.) in the west to low lands (50-200 m a.s.l.) in the east along the Baltic Sea. However, mountainous coniferous forests are only found up to ca. 400 m a.s.l. in the north (68°N) and ca. 800 m a.s.l. in the south (61°N).

2.2 Tree growth data

We studied tree growth patterns of the most common and widely distributed spruce species in each study area: black spruce (*Picea mariana* (Mill.) Britton) in Quebec and Norway spruce (*P. abies* (L.) H. Karst) in Sweden. A total of 6,876 and 14,438 tree-ring width series were retrieved from the Quebec (Ministère des Ressources naturelles du Québec, 2014) and Swedish forest inventory database (Riksskogstaxeringen, 2016), respectively. We adapted data selection procedures to each database to provide as high local coherence in growth patterns as possible.

For Quebec, core series were collected from dominant trees on permanent plots (three trees per plot, four cores per tree) between 2007 and 2014. Permanent plots were situated in unmanaged old-growth black spruce forests north of the northern limit for timber exploitation.

Core series were aggregated into individual tree series using a robust bi-weighted mean (robust average unaffected by outliers, Affymetrix 2002). To enhance growth coherence at the local level, we further selected tree series presenting strong correlation ($r > 0.4$) with their respective local landscape unit master chronology. This master chronology corresponds to the average of all other tree series within the same landscape unit (landscape units are 6341 km² on average and delimit a territory characterized by specific bioclimatic and physiographic factors (Robitaille & Saucier, 1998)). This resulted in the selection of 790 tree series that were averaged at the plot level using a robust bi-weighted mean. The obtained 444 plot chronologies had a common period of 1885-2006 (Table 1). Plot chronologies were detrended using a log transformation and a 32-year spline de-trending, and pre-whitened using autocorrelation removal (Cook & Peters, 1981). Detrending aims at removing the low-frequency age-linked variability in tree-ring series (decreasing tree-ring width with increasing age) while keeping most of the high-frequency variability (mainly linked to climate). Pre-whitening removes all but the high frequency variation in the series by fitting an autoregressive model to the detrended series. The order of the auto-regressive model was selected by Akaike Information Criterion (Akaike 1974).

For Sweden, core series were collected within the boreal zone of the country (59°N-68°N) on temporary plots between 1983 and 2010. Temporary plots were situated in productive forests, i.e. those with an annual timber production of at least 1m³/ha. These forests encompass protected, semi-natural and managed forests. In each plot, one to three trees were sampled, with two cores per tree. Swedish inventory procedures do not include any visual and statistical cross-dating of core series at the plot level. To filter out misdated series, we aggregated core series into 4067 plot chronologies using a robust bi-weighted mean, and compared them to Norway spruce reference chronologies from the International Tree-Ring Data Base (International Tree Ring Data Bank (ITRDB), 2016). In total, seven ITRDB

reference chronologies were selected (Fig. 1b), all representative of tree growth at mesic sites in boreal Sweden. Plot and reference chronologies were detrended and pre-whitened using the same standard procedures used for the Quebec data. Each plot chronology was then compared with its geographically nearest reference chronology - determined based on Euclidean distance - using Student's t-test analysis (Student 1908). Plot chronologies with a t-test value lower than 2.5 with their respective nearest reference chronology were removed from further analyses (the t-test value threshold was set up according to the mean length of plot chronologies (Table 1)). A total of 1256 plot chronologies (with a common period of 1936-1995) passed this quality test (Table 1).

Table 1. Characteristics of tree-ring width chronologies*.

	Quebec	Sweden198
Plot chronologies		
Number	444	1256
Mean length (SD) [yrs.]	191 (59)	80 (3)
Grid cell chronologies		
Number	36	56
Plot chronologies per grid cell (SD)	12 (8)	23 (13)
Mean length (SD) [yrs.]	230 (47)	81 (13)
Common period	1885-2006	1936-1995
Regional chronologies		
Number	3	3
Grid cell chronologies per cluster	7/10/19*	14/19/23**
Length [yrs.]	212/196/263*	81/81/79**
Common period	1812-2008	1929-2008

*Data for Q_W, Q_C and Q_E chronologies respectively.

**Data for S_S, S_C and S_N chronologies respectively.

2.3 Spatial aggregation of plot chronologies into regional chronologies in each study area

Quality checked chronologies at the plot level were aggregated into 1° x 1° latitude-longitude grid cell chronologies within each study area (Fig. 1b). Grid cell chronologies were calculated as the robust bi-weighted mean of all plot chronologies within each grid cell. Grid cells

containing less than three plot chronologies were removed from further analyses. This resulted in a total of 36 and 56 grid cell chronologies in Quebec and Sweden, respectively (Fig. 1b, Table 1). Grid cells contained on average 12 and 23 plot chronologies in Quebec and Sweden, respectively (Table 1).

To investigate the influence of spatial scale in climate-growth sensitivity analyses, we performed an ordination of grid cell chronologies within each study area over their common period (Fig. 1c). The common period between grid cell chronologies was 1885-2006 and 1936-1995 in Quebec and Sweden, respectively. Ordination analyses were performed in R using Euclidean dissimilarities matrices (*dist* function) and Ward agglomeration (*hclust* function) methods. Three main clusters were identified in each study area (Fig. 1c). Spatial extents of all clusters were consistent with well-defined bioclimatic regions, providing support to data selection procedures. In Quebec, clusters identified in the West (Q_W) and the East (Q_E) corresponded well to the drier and wetter northern boreal region, respectively (Fig. 1b & c). In Sweden, the cluster identified in the South (S_S) corresponded to a combination of the nemo-boreal and southern boreal zones (Moen, 1999). The Swedish central (S_C) and northern (S_N) clusters corresponded to the mid-boreal and northern boreal zones, respectively (Fig. 1b & c) (Moen, 1999). Regional chronologies were built as the average of all grid cell chronologies within a cluster. In Sweden, inter-cluster correlations were all significant and ranged from 0.77 (S_S vs S_N) to 0.94 (S_C vs S_N). In Quebec, inter-cluster correlations were all significant and ranged from 0.44 (Q_W vs Q_E) to 0.52 (Q_C vs Q_E) (see Appendix S1-S3 in Supporting Information). Henceforward, the terms ‘local level’ and ‘regional level’ refer to analyses focusing on the grid cell chronologies and the six regional chronologies, respectively.




Fig. 1 a: Location of the two study areas (black frame); b & c: Clusters identified in each study area by ordination of $1^\circ \times 1^\circ$ latitude-longitude grid cell chronologies. Ordination analyses were performed over the common period between grid cell chronologies in each study area using Euclidean dissimilarities matrices and Ward agglomeration methods. The common period was 1885-2006 for Quebec and 1936-1995 for Sweden. Ordinations included 36 and 56 grid cell chronologies in Quebec and Sweden, respectively. A western (Q_W), central (Q_C) and eastern (Q_E) cluster were identified in Quebec and a southern (S_S), central (S_C) and northern (S_N) cluster were identified in Sweden. Reference chronologies

from the ITRDB used for the cross-dating of plot chronologies in Sweden are indicated with a * (swed011, swed012, swed013, swed014, swed015, swed017 and swed312). The grey shading indicates the boreal zone delimitation according to Brant *et al.* (2013).

2.4 Climate data

For each grid cell, we extracted local seasonal mean temperature and total precipitation data (1950-2008) from the CRU TS 3.24 1° x 1° (Harris *et al.*, 2014), with seasons spanning from the previous (pJJA) through the current summer (JJA). Climate data were further aggregated at the regional level as the robust bi-weighted mean of climate data of all grid cells contained in each regional cluster (Fig. 1b & c). Seasonal AMOC indices (1961-2005, first AMOC measurements in 1961) were extracted from the European Center for Medium-Range Weather Forecast (Ocean Reanalysis System ORA-S3). Seasonal AO and NAO indices (1950-2008) were extracted from the Climate Prediction Center database (NOAA, 2016). Seasonal AMOC, NAO, and AO indices included previous summer, winter (DJF), and current summer. All seasonal climate data were downloaded using the KNMI Climate Explorer (Trouet & Van Oldenborgh, 2013) and were detrended using linear regression and thereafter pre-whitened (autocorrelation of order 1 removed from time series).

2.5 Links between seasonal climate and growth patterns

Analyses were run over the 1950-2008 period (the longest common period between tree growth and climate data), except with AMOC indices which were only available for 1961-2005. Tree growth patterns were correlated with seasonal climate variables (previous-to-current summer temperature averages and precipitation sums) and seasonal indices (previous summer, winter, and current summer AMOC, NAO, and AO) at the regional and local levels. To minimize type I errors, each correlation analysis was tested for 95% confidence intervals using 1000 bootstrap samples. In addition, moving correlation analyses (21-yr windows

moved one year at a time) were performed at the regional level using the same procedures as above. All calculations were performed using the R package *treeclim* (Zang & Biondi, 2015). For more details regarding bootstrapping procedures please see the description of the “*dcc*” function of this package.

3 Results

3.1 Tree growth responses to seasonal climate

Some significant climate-growth associations were observed at the regional level (Fig. 2). Significant associations at the local level displayed strong spatial patterns and revealed heterogeneous within-region growth responses (Figs. 3 and 4). Moving correlations revealed numerous shifts in the significance of climate-growth associations around 1980 (Fig. 5).

3.1.1 Quebec

No significant climate-growth associations were observed at the regional level in western boreal Quebec over the entire study period (Fig. 2). Some significant positive responses to previous winter and current spring temperatures were observed at the local level, but these concerned a minority of cells (Fig. 3). Moving correlations revealed that Q_W significantly correlated with previous summer precipitation (negatively) before the 1970s, with previous winter temperatures (positively) from the 1970s and with current spring temperatures (positively) from 1980 (Fig. 5).

Tree growth in central boreal Quebec significantly and positively correlated with current summer temperatures at the regional and local levels (Figs. 2 and 3). Numerous negative correlations between tree growth and spring precipitation were observed at the local level (Fig. 3). Moving correlations revealed an emerging correlation between Q_C and previous winter temperatures in the early 1970s (significant during most intervals up to most recent years) (Fig. 5).

No significant climate-growth associations were observed in eastern boreal Quebec at the regional level (Fig. 2). At the local level, some positive significant correlations with current summer temperatures were observed (Fig. 3). Moving correlations revealed that Q_E correlated significantly and positively with current summer temperatures up to the early 1970s (Fig. 5).




Fig. 2. Tree growth responses to seasonal temperature averages (a) and precipitation sums (b) at the regional level over the 1950-2008 period, as revealed by correlation analyses. Analyses were computed between the six regional chronologies (Q_W, Q_C, and Q_E in NA; and S_S, S_C and S_N in NE) and seasonal climate data. Climate data were first extracted from the CRU TS 3.24 1° x 1° (Harris *et al.*, 2014) for each grid cell and then aggregated at the regional level by a robust bi-weighted mean. Seasons included previous summer (pJJA), previous autumn (SON), winter (DJF); current spring (MAM) and current summer (JJA). Significant correlations ($P < 0.05$) are marked with a star.




Fig. 3. Tree growth responses to seasonal temperature averages (a) and precipitation sums (b) at the local level over the 1950-2008 period in Quebec, as revealed by correlation analyses. Analyses were computed between grid cell chronologies and local seasonal climate data extracted for each grid cell from the CRU TS 3.24 1° x 1° (Harris *et al.*, 2014). Seasons included previous summer (pJJA), previous autumn (SON), winter (DJF); current spring (MAM) and current summer (JJA). To visualize separation between regional clusters (Q_W, Q_C, and Q_E, cf. Fig. 1) correlation values at Q_C grid cells are plotted with circles. Significant correlations ($P < 0.05$) are marked with a black dot.

3.1.2 Sweden

Tree growth in southern boreal Sweden correlated significantly and negatively with previous summer and winter temperatures at the regional and local levels, the correlation with winter temperatures concerning however only a minority of cells (Figs. 2 and 4). Moving

correlations indicated that the negative association with previous summer temperatures remained significant up to the early 1990s and that the negative association with winter temperatures emerged after 1980 (Fig. 5).

In central boreal Sweden, tree growth significantly and negatively correlated with previous summer temperatures both at the regional and local levels (Figs. 2 and 4). Some additional significant correlations with winter temperatures (negative) and with current summer temperatures (positive) were observed at the local level (Fig. 4). Moving correlation analyses revealed a significant positive correlation between S_C and current summer temperatures that dropped and became non-significant at the end of the study period (Fig. 5). In addition, the correlation between S_C and previous summer precipitation shifted from significantly negative to significantly positive during the 1980s (Fig. 5). S_C became significantly and negatively correlated with previous summer temperatures after the 1980s and stopped being significantly and negatively correlated with previous autumn precipitation and with winter temperatures at the end of the 1970s (Fig. 5).

Tree growth in northern boreal Sweden correlated significantly with previous summer (negatively) and current summer temperatures (positively) both at the regional and local levels (Figs. 2 and 4). At the local level, tree growth in some cells significantly and negatively correlated with winter temperatures (Fig. 4). Significant and negative responses to current summer precipitation were observed at northernmost cells (Fig. 4). Moving correlations revealed that the positive association with current summer temperatures was only significant at the beginning and at the end of the study period (Fig. 5). After the 1980s, significant positive associations with previous autumn temperatures emerged (Fig. 5) and the significant negative association with winter temperatures disappeared.




Fig. 4. Tree growth responses to seasonal temperature averages (a) and precipitation sums (b) at the local level over the 1950-2008 period in Sweden, as revealed by correlation analyses. Analyses were computed between grid cell chronologies and local seasonal climate data extracted for each grid cell from the CRU TS 3.24 1° x 1° (Harris *et al.*, 2014). Seasons included previous summer (pJJA), previous autumn (SON), winter (DJF); current spring (MAM) and current summer (JJA). To visualize the separation between regional clusters (S_S, S_C, and S_N, cf. Fig. 1) correlation values at S_C grid cells are plotted with circles. Significant correlations ($P < 0.05$) are marked with a black dot.




Fig. 5. Moving correlations between regional seasonal temperature averages (red lines) and precipitation sums (blue lines), and the six regional chronologies (Q_W, Q_C, and Q_E in NA; and S_S, S_C and S_N in NE) over the 1950-2008 period. Climate data were first extracted for each grid cell from the CRU TS 3.24 1° x 1° (Harris *et al.*, 2014) and then aggregated at the regional level by robust bi-weighted mean. Seasons included previous summer (pJJA), previous autumn (SON), winter (DJF); current spring (MAM) and current summer (JJA). Moving correlations were calculated using 21-yr windows moved one year at a time and are plotted using the central year of each window. Windows of significant correlations ($P < 0.05$) are marked with a dot.

3.2 Links between tree growth patterns and large-scale indices

Some significant associations were found between tree growth and large-scale indices (Figs. 6, 7, and 8). Moving correlation analyses revealed some shifts from pre-1980 insignificant to

post-1980 significant correlations (Fig. 9). The seasonal indices involved in these shifts varied across regional chronologies.

3.2.1 Quebec

Tree growth in western boreal Quebec was significantly and negatively associated with the winter AMOC and the winter AO indices at the regional level (Fig. 6). At the local level, these associations concerned, however, a minority of cells (Fig. 7). Moving correlations revealed that the regional negative association with winter AMOC was only significant in the most recent part of the study period (Fig. 9). Significant negative correlations between Q_W and current summer NAO and AO indices were observed from the 1980s up to the most recent years, at which point they show a steep increase and become non-significant (Fig. 9).

In central boreal Quebec, no significant associations between tree growth and seasonal indices were identified at the regional or local level (Figs. 6 and 7). Moving correlations indicated significant negative correlations with previous summer NAO and AO indices of during the 1970s, with winter NAO and AO indices during the 1980s and with current summer NAO and AO indices from the 1980s up to the most recent years (Fig. 9).

No significant association was identified between large-scale indices and tree growth in eastern boreal Quebec (Figs. 6, 7, and 9).





Fig. 6. Correlation between seasonal AMOC (a), NAO (b), and AO (c) indices and the six regional chronologies (Q_W, Q_C, and Q_E in NA; and S_S, S_C and S_N in NE). Seasonal indices include previous summer (pJJA), winter (DJF), and current summer (JJA), and were calculated as mean of monthly indices. Correlations were calculated over the 1961-2005 period for AMOC, and over the 1950-2008 period for NAO and AO. Significant correlations ($P < 0.05$) are marked with a star.



391

392 **Fig. 7.** Correlation between seasonal AMOC (a), NAO (b), and AO (c) indices, and growth patterns at the local level in Quebec. Seasonal indices
 393 include previous summer (left-hand panels), winter (middle panels), and current summer (right-hand panels), and were calculated as mean of
 394 monthly indices. Correlations were calculated over the 1961-2005 period for AMOC, and over the 1950-2008 period for NAO and AO. To
 395 visualize the separation between regional clusters, correlation values at Q_C grid cells are plotted with circles. Significant correlations ($P < 0.05$)
 396 are marked with a black dot.

3.2.2 Sweden

No significant association between tree growth in southern boreal Sweden and seasonal large-scale indices was identified at the regional or local level (Figs. 6 and 8). Moving correlations revealed, however, significant negative associations between S_S and the winter AMOC index before the 1980s (Fig. 9).

In central boreal Sweden, tree growth significantly and positively correlated with the current summer NAO index at the regional level (Fig. 6). At the local level, this correlation concerned, however, a minority of cells (Fig. 8). Moving correlations revealed that the significant positive association with the current summer NAO index emerged in the early 1980s (Fig. 9) and that S_C significantly correlated with the current summer AMOC index during the 1980s (Fig. 9).

In northern boreal Sweden, tree growth significantly correlated with the current summer NAO index (positively) and with the winter AO index (negatively) at the regional level (Fig. 6). At the local level, the positive association with summer NAO concerned a large majority of cells and the negative association with the winter AO index concerned only very few cells (Fig. 8). Moving correlation analyses indicated that the positive association between S_N and the current summer NAO index was only significant after the 1980s and that S_N significantly correlated with current summer AMOC during most of the 1980s (Fig. 9).




Fig. 8. Correlation between seasonal AMOC (a), NAO (b), and AO (c) indices, and growth patterns at the local level in Sweden. Seasonal indices were calculated as mean of monthly indices and include previous summer (left-hand panels), winter (middle panels), and current summer (right-hand panels). Correlations were calculated over the 1961-2005 period for AMOC, and over the 1950-2008 period for NAO and AO. To visualize the separation between regional clusters, correlation values at S_C grid cells are plotted with circles. Significant correlations ($P < 0.05$) are marked with a black dot.




Fig. 9. Moving correlations between previous summer (pJJA; left-hand panels), winter (DJF; middle panels) and current summer (JJA; right-hand panels) large-scale indices, and the six regional chronologies (Q_W, Q_C, and Q_E in NA; and S_S, S_C and S_N in NE). Large-scale indices include AMOC (black), NAO (red), and AO (blue). Moving correlations were calculated using 21-yr windows moved one year at a time and are plotted using the central year of each window. Correlations were calculated over the 1961-2005 period for AMOC, and over the 1950-2008 period for NAO and AO. Windows of significant correlations ($P < 0.05$) are marked with a dot.

4 DISCUSSION

4.1 Spatial aggregation of tree growth data

The high correlation between the regional chronologies in NE (Appendix S1), especially between the central and northern chronologies, could have supported the construction of one single boreal Sweden-wide regional chronology. Climate-growth analyses at the regional and local level revealed, nevertheless, clear differences across space in tree growth sensitivity to climate (Fig. 4) and to large-scale indices (Fig. 8), with a higher sensitivity in northernmost forests. The aggregation of tree growth data across space, even if based on objective similarity statistics (Appendix S1), may, therefore, mask important local differences in climate-growth interactions (Macias *et al.* 2004). Our results demonstrate that spatial aggregation should not be performed without accounting for bioclimatic domains especially when studying climate-growth interactions. In practice, one should at least check that a spatial similarity in tree growth patterns is associated with spatial similarity in seasonal climate. The use of both the regional and local scales regarding climate-growth interactions, as in the present study, is, therefore, recommended to exhaustively and more precisely capture cross-scale diverging and emerging tree growth patterns and sensitivity to climate.

4.2 Post-1980 shifts towards significant influence of large-scale indices on boreal tree growth

The emergence of a post-1980 significant positive tree growth response to the current summer NAO index in central and northern boreal Sweden (Fig. 9) appears to be linked to spatial variability in the NAO influence on seasonal climate (Fig. 10). Summer NAO has had little to no influence on summer climate variability over the entire period 1950-2008 in boreal Quebec or Sweden (Appendix S4). However, the partitioning of the period into two sub-periods of similar length (1950-1980 and 1981-2008) revealed a northeastward migration of the

significant-correspondence field between the summer NAO index and local climate, particularly in NE (Fig. 10). Over the 1981-2008 period, the summer NAO index was significantly and positively associated with temperature and negatively with precipitation in boreal Sweden (Fig. 10). Higher growing-season temperatures, induced by a higher summer NAO, might have promoted the growth of temperature-limited Swedish boreal forest ecosystems, explaining recent positive response of tree growth to this large-scale index in the central and northern regions (Fig. 9). The northeastward migration of the NAO-climate spatial field may be an early sign of a northward migration of the North Atlantic Gulf stream (Taylor & Stephens, 1998) or a spatial reorganization of the Icelandic-low and Azores-high pressure NAO's nodes (Portis *et al.*, 2001; Wassenburg *et al.*, 2016). The August Northern Hemisphere Jet over NE reached its northernmost position in 1976 but thereafter moved southward, despite increasing variability in its position (Trouet *et al* 2018). This southward migration of the jet may weaken the strength of the observed post-1980 positive association between boreal tree growth and the summer NAO index in NE in the coming decades.




Fig. 10. Correspondence between summer NAO (a) and AO (b) indices and local summer climate (mean temperature and total precipitation) between 1950 and 1980 (left-hand panels) and between 1981 and 2008 (right-hand panels). NAO and AO indices over the 1950-2008 period were extracted from NOAA's climate prediction center. Summer mean temperature and total precipitation are those of CRU TS 3.24 1° x 1° (Harris *et al.*, 2014). All correlations were computed in the KNMI Climate Explorer (<https://climexp.knmi.nl> (Trouet & Van Oldenborgh, 2013)). Indices and climate variables were normalized (linear regression) prior to analyses. Only correlations significant at $P < 0.05$ are plotted.

The post-1980 significant negative associations between tree growth and summer NAO and AO indices in boreal Quebec are more challenging to interpret. There was no evident significant tree growth response to summer temperature in these regions when analyzed over the full 1950-2008 period (Fig. 4). Yet, some significant positive associations between tree growth and temperatures were observed with winter temperatures from the 1970s (in central Quebec) and with spring temperatures from the 1980s (in western Quebec only) (Fig. 5). These associations indicate that tree growth in boreal Quebec has been limited by winter and spring climate since the 1970s and 1980s, respectively. Below-average summer temperatures induced by high summer NAO and AO may exacerbate the sensitivity of tree growth to low temperatures. Noting that no significant post-1980 association was observed between temperature and summer NAO and AO indices in Quebec (Fig. 10), the emerging negative tree growth response to summer NAO and AO indices may indicate a complex interplay between large-scale indices and air mass dynamics and lagged effects over several seasons (Boucher *et al.*, 2017).

In western Quebec, tree growth was negatively influenced by the winter AMOC index at the regional level (Fig. 6). This relationship appears to be linked to a significant positive association between tree growth and spring temperature (Figs. 5 and 9). Positive winter AMOC indices are generally associated with cold temperatures in Quebec, and particularly so in the West (Appendix S4). Positive winter AMOC indices are associated with the dominance

of dry winter air masses of Arctic origin over Quebec, and may thereby delay the start of the growing season and reduce tree-growth potential.

Forest dynamics in NA have been reported to correlate with Pacific Ocean indices such as the Pacific Decadal Oscillation (PDO) or the El-Nino Southern Oscillation (ENSO), particularly through their control upon fire activities (Macias Fauria & Johnson 2006, Le Goff *et al.*, 2007). These indices have not been investigated in the present study but might present some additional interesting features.

4.3 Contrasting climate-growth associations among boreal regions

Post-1980 shifts in tree growth sensitivity to seasonal climate differed among boreal regions. In NA, we observed the emergence of significantly positive growth responses to winter and spring temperatures. In NE, observed post-1980 shifts mainly concerned the significance of negative growth responses to previous summer and winter temperatures. Warmer temperatures at boreal latitudes have been reported to trigger contrasting growth responses to climate (Wilmking *et al.*, 2004) and to enhance the control of site factors upon growth (Nicklen *et al.*, 2016). This is particularly true with site factors influencing soil water retention, such as soil type, micro-topography, and vegetation cover (Düthorn *et al.*, 2013). Despite a generalized warming at high latitudes (Serreze *et al.*, 2009), no increased sensitivity of boreal tree growth to precipitation was identified in the present study, except in central Sweden where tree growth became positively and significantly correlated to previous summer precipitation (Fig. 5). This result underlines that temperature remains the major-growth limiting factor in our study regions.

The observed differences in tree growth response to winter temperature highlight diverging non-growing season temperature constraints on boreal forest growth. While warmer winters appear to promote boreal tree growth in NA, they appear to constrain tree growth in

boreal NE. Such opposite responses to winter climate from two boreal tree species of the same genus might be linked to different winter conditions between Quebec and Sweden. In NA, winters conditions are more continental and harsher than in NE (Appendix S5). Warmer winters may therefore stimulate an earlier start of the growing season and increase growth potential (Rossi *et al.*, 2014). However, warmer winters, combined with shallower snow-pack, have been shown to induce a delay in the spring tree growth onset, through lower thermal inertia and a slower transition from winter to spring (Contosta *et al.*, 2017). This phenomenon might explain the negative association between tree growth and winter temperatures observed in NE.

The post-1970s growth-promoting effects of winter and spring temperature in NA (Fig. 5) suggest, as earlier reported by Charney *et al.* (2016) and Girardin *et al.* (2016), that, under sufficient soil water availability and limited heat stress conditions, tree growth at mid- to high-latitudes can increase in the future. However, warmer winters may also negatively affect growth by triggering an earlier bud break and increasing risks of frost damages to developing buds (Cannell and Smith, 1986) or by postponing the start of the growing season (see above, Contosta *et al.*, 2017). This might provide an argument against a sustained growth-promoting effect of higher seasonal temperatures (Girardin *et al.*, 2014).

4.4 Gradients in the sensitivity of tree growth to North Atlantic Ocean dynamics across boreal Quebec and Sweden

Trees in western and central boreal Quebec, despite being furthest away from the North Atlantic Ocean in comparison to trees in eastern boreal Quebec, were the most sensitive to oceanic and atmospheric dynamics, and particularly to current summer NAO and AO indices after the 1970s. In these two boreal regions, tree growth responses to large-scale indices were stronger and more spatially homogeneous than tree growth responses to regional climate. This

suggests that growth dynamics in western and central boreal Quebec, despite being mainly temperature-limited, can be strongly governed by large-scale oceanic and atmospheric dynamics (Boucher *et al.*, 2017). The tree growth sensitivity to the winter AMOC index observed at regional level in western boreal Quebec might directly emerge from the correspondence between AMOC and winter snow fall. Western boreal Quebec is the driest and most fire-prone of the Quebec regions studied here. Soil water availability in this region strongly depends on winter precipitation. High winter AMOC indices are associated with the dominance of Arctic air masses over NA and leads to decreased snowfall (Appendix S4). Large-scale indices, through their correlation with regional fire activity, can also possibly override the direct effects of climate on boreal forest dynamics (Drobyshev *et al.*, 2014; Zhang *et al.*, 2015). Fire activity in NA strongly correlates with variability in atmospheric circulation, with summer high-pressure anomalies promoting the drying of forest fuels and increasing fire hazard (Skinner *et al.*, 1999, Macias Fauria & Johnson 2006) and low-pressure anomalies bringing precipitation and decreasing fire activity.

In Sweden, the northernmost forests were the most sensitive to North Atlantic Ocean dynamics, particularly to the summer NAO (Fig. 8). These high-latitude forests, considered to be 'Europe's last wilderness' (Kuuluvainen *et al.*, 2017), are experiencing the fastest climate changes (Hansen *et al.*, 2010). Numerous studies have highlighted a correspondence between tree growth and NAO (both winter and summer) across Sweden (D'Arrigo *et al.*, 1993; Cullen *et al.*, 2001; Linderholm *et al.*, 2010), with possible shifts in the sign of this correspondence along north-south (Linderholm *et al.*, 2001) and west-east gradients (Linderholm *et al.*, 2003). Our results identified a post-1980 positive correspondence between tree growth and summer NAO, spatially restricted to the northernmost regions (Figs. 8 and 9). This emerging correspondence appears linked to the combination of a growth-promoting effect of higher temperature at these latitudes (Fig. 5) and a northeastward migration of the spatial

correspondence between NAO and local climate (Fig. 10). Boreal forests of Quebec (western and central) and Sweden (central and northern) emerged as regions sensitive to large-scale climate dynamics. We, therefore, consider them as suitable for a long-term survey of impacts of ocean-atmosphere dynamics on boreal forest ecosystems.

Acknowledgements

This study was financed by the Natural Sciences and Engineering Research Council of Canada (NSERC) through the project ‘Natural disturbances, forest resilience and forest management: the study case of the northern limit for timber allocation in Quebec in a climate change context’ (STPGP 41344-11). We acknowledged financial support from the Nordic Forest Research Cooperation Committee (SNS) through the network project entitled ‘Understanding the impacts of future climate change on boreal forests of northern Europe and eastern Canada’, from the EU Belmont Forum (project PREREAL), NINA’s strategic institute program portfolio funded by the Research Council of Norway (grant no. 160022/F40), the Forest Complexity Modelling (FCM), an NSERC funded program in Canada and a US National Science Foundation CAREER grant (AGS-1349942). We are thankful to the Ministry of Forests, Wildlife and Parks (MFFP) in Quebec and to the Swedish National Forest Inventory (Bertil Westerlund, Riksskogstaxeringen, SLU) in Sweden for providing tree-growth data. ID thanks the Swedish Institute for support of this study done within the framework of CLIMECO project.

Appendix A – Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1902/j.gloplacha.2018.03.006>

References

Affymetrix (2002) Appendix I- Statistical Algorithms Description Document

- Akaike H. (1974). A new look at the statistical model identification. *IEEE Trans. Autom. Control* **16**, 716-723.
- Bala G., Caldeira K., Wickett M., Phillips T.J., Lobell D.B., Delire C. & Mirin A. (2007) Combined climate and carbon-cycle effects of large-scale deforestation. *Proc. Natl. Acad. Sci. U. S. A.*, **104**, 6550–6555.
- Baldwin M.P. & Dunkerton T.M. (1999) Propagation of the Arctic Oscillation from the stratosphere to the troposphere. *J. Geophys. Res.* **104**, 30,937-30,946.
- Betts R.A. (2000) Offset of the potential carbon sink from boreal forestation by decreases in surface albedo. *Nature* **408**, 187-190.
- Boucher E., Nicault A., Arseneault D., Bégin Y. & Pasha M. (2017) Decadal variations in eastern Canada's taiga wood biomass production forced by ocean-atmosphere interactions. *Sci. Rep.* **7**, 2457.
- Brant J.P., Flannigan M.D., Maynard D.G., Thompson I.D. & Volney W.J.A. (2013) An introduction to Canada's boreal zone: ecosystem processes, health, sustainability, and environmental issues. *Environ. Rev.* **21**, 207–226.
- Briffa K.R., Schweingruber F.H., Jones P.D., Osborn T.J., Shiyatov S.G. & Vaganov E.A. (1998) Reduced sensitivity of recent tree-growth to temperature at high northern latitudes. *Nature*, **391**, 678-682.
- Cannell M.G.R. & Smith R.I. (1986) Climatic warming, spring budburst, and frost damage on trees. *J Appl Ecol*, **23**, 177–191.
- Charney N.D., Babst F., Poulter B., Record S., Trouet V.M., Frank D., Enquist B.J. & Evans M.E. (2016) Observed forest sensitivity to climate implies large changes in 21st century North American forest growth. *Ecology Letters*, **19**, 1119-1128
- Chen D. & Hellström C. (1999) The influence of the North Atlantic Oscillation on the regional temperature variability in Sweden: spatial and temporal variations. *Tellus A*, **51**, 505-516.
- Contosta A.R., Adolph A., Burchsted D., Burakowski E., Green M., Guerra D., Albert M., Dibb J., Martin M., McDowell W.H., Routhier M., Wake C., Whitaker R. & Wollheim W. (2017) A longer vernal window: the role of winter coldness and snowpack in driving spring transitions and lags. *Glob Change Biol*, **23**, 1610-1625.
- Cook E.R. & Peters K. (1981) The smoothing spline: an approach to standardizing forest interior tree-ring width series for dendroclimatic studies *Tree-Ring Bull*, **41**, 45-53.
- Cullen H.M., D'Arrigo R.D. & Cook E.R. (2001) Multiproxy reconstructions of the North Atlantic Oscillation. *Paleoceanography*, **16**, 27-39.
- D'Arrigo R.D., Cook E.R., Jacoby G.C. & Briffa K.R. (1993) NAO and sea surface temperature signatures in tree-ring records from the North Atlantic sector. *Quat Sci Rev*, **12**, 431-440.
- de Wit H.A., Bryn A., Hofgaard A., Karstensen J., Kvilevag M.M. & Peters G.P. (2014) Climate warming feedback from mountain birch forest expansion: Reduced albedo dominates carbon uptake. *Glob Change Biol*, **20**, 2344-55.
- Drobyshev I., Guitard M.-A., Asselin H., Genries A. & Bergeron Y. (2014) Environmental controls of the northern distribution limit of yellow birch in eastern Canada. *Can J For Res*, **44**, 720-731.
- Drobyshev I., Bergeron Y., Vernal A., Moberg A., Ali A.A. & Niklasson M. (2016) Atlantic SSTs control regime shifts in forest fire activity of Northern Scandinavia. *Sci Rep*, **6**, 22532.
- Düthorn E, Holzkämper S., Timonen M. & Esper J. (2013) Influence of micro-site conditions on tree-ring climate signals and trends in central and northern Sweden. *Trees*, **27**, 1395-1404.

- Gauthier S., Bernier P., Kuuluvainen T., Shvidenko A.Z. & Schepaschenko D.G. (2015) Boreal forest health and global change. *Science*, **349**, 819-822.
- Gerardin V. & McKenney D. (2001) Une classification climatique du Québec à partir de modèles de distribution spatiale de données climatiques mensuelles : vers une définition des bioclimats du Québec. *Direction du patrimoine écologique et du développement durable, ministère de l'Environnement, Québec*,
- Girardin M.P., Guo X.J., De Jong R., Kinnard C., Bernier P. & Raulier F. (2014) Unusual forest growth decline in boreal North America covaries with the retreat of Arctic sea ice. *Glob Change Biol*, **20**, 851-66.
- Girardin M.P., Bouriaud O., Hogg E.H., Kurz W., Zimmermann N.E., Metsaranta J.M., de Jong R., Frank D.C., Esper J., Buntgen U., Guo X.J. & Bhatti J. (2016) No growth stimulation of Canada's boreal forest under half-century of combined warming and CO₂ fertilization. *Proc Natl Acad Sci U S A*, **113**, E8406-E8414.
- Hansen J., Ruedy R., Sato M. & Lo K. (2010) Global surface temperature change. *Rev Geophys*, **48**
- Harris I., Jones P.D., Osborn T.J. & Lister D.H. (2014) Updated high-resolution grids of monthly climatic observations --- the CRU TS3.10 dataset. *Int J Climatol*, **34**, 623-642.
- Hurrell J.W. (1995) Decadal trends in the north atlantic oscillation: regional temperatures and precipitation. *Science*, **269**, 676-9.
- International Tree Ring Data Bank (ITRDB) (2016) Available at: <http://www.ncdc.noaa.gov/data-access/paleoclimatology-data/datasets/tree-ring> (accessed 15 October 2014).
- IPCC (2014) Intergovernmental Panel on Climate Change. Fifth assessment report climate change 2014. synthesis report---summary for policymakers. 1-35.
- Jackson L.C., Kahana R., Graham T., Ringer M.A., Woollings T., Mecking J.V. & Wood R.A. (2015) Global and European climate impacts of a slowdown of the AMOC in a high resolution GCM. *Clim. Dyn.*, **45**, 3299-3316.
- Karlsen S.R., Høgda K.A., Wielgolaski F.E., Tolvanen A., Tømmervik H., Poikolainen J. & Kubin E. (2009) Growing-season trends in Fennoscandia 1982–2006, determined from satellite and phenology data. *Clim Res*, **39**, 275-286.
- Kuuluvainen T., Hofgaard A., Aakala T. & Jonsson B.G. (2017) North Fennoscandian mountain forests: History, composition, disturbance dynamics and the unpredictable future. *For Ecol Manage*, **385**, 140-149.
- Le Goff H., Flannigan M.D., Bergeron Y. & Girardin M.P. (2007) Historical fire regime shifts related to climate teleconnections in the Waswanipi area, central Quebec, Canada. *Int. J. Wild Fire*. **16**, 607-618
- Lenton T.M. (2011) Early warning of climate tipping points. *Nat Clim Chan*, **1**, 201-209.
- Lenton T.M., Held H., Kriegler E., Hall J.W., Lucht W., Rahmstorf S. & Schellnhuber H.J. (2008) Tipping elements in the Earth's climate system. *P Natl Acad Sci USA*, **105**, 1786-93.
- Linderholm H.W., Solberg B. Ø. & Lindholm M. (2003) Tree-ring records from central Fennoscandia: the relationship between tree growth and climate along a west–east transect. *The Holocene*, **13**, 887-895
- Linderholm H.W., Björklund J.A., Seftigen K., Gunnarson B.E., Grudd H., Jeong J.-H., Drobyshev I. & Liu Y. (2010) Dendroclimatology in Fennoscandia – from past accomplishments to future potential. *Clim Past*, **9**, 93-114.
- Lindholm M., Eggertsson O., Lovelius N., Raspopov O., Shumilov O. & Läänelaid A. (2001) Growth indices of North European Scots pine record the seasonal North Atlantic Oscillation. *Boreal Environ Res*, **6**, 275-284.

- Macias M., Timonen M., Kirchhefer A.J., Lindholm M., Eronen M. & Gutiérrez E. (2004). Growth variability of Scots pine (*Pinus sylvestris*) along a west-east gradient across northern Fennoscandia: A dendroclimatic approach. *Arctic, Antarctic, and Alpine Research*, **36**, 565-574.
- Macias Faurias M. & Johnson E.A. (2006) Large-scale climatic patterns control large lightning fire occurrence in Canada and Alaska forest regions. *J Geophys Res*, **111**, G04008, doi:10.1029/2006JG000181
- Ministère des Ressources naturelles du Québec (2014) Norme d'inventaire écodendrométrie nordique. In: (ed. S.D. Forêts.), p. 156 p. + 10 annexes.
- Moen A. (1999) National Atlas of Norway. Vegetation. Norway Mapping Authority, Hønefoss.
- Nicklen E.F., Roland C.A., Ruess R.W., Schmidt J.H. & Lloyd A.H. (2016) Local site conditions drive climate–growth responses of *Picea mariana* and *Picea glauca* in interior Alaska. *Ecosphere*, **7**
- NOAA (2016) *National Oceanic and Atmospheric Administration*. Available at: <http://www.cpc.ncep.noaa.gov/data/> (accessed 01-09 2016).
- Ols C., Hofgaard A., Bergeron Y. & Drobyshev I. (2016) Previous growing season climate controls the occurrence of black spruce growth anomalies in boreal forests of Eastern Canada. *Can J For Res*, **46**, 696-705.
- Ottersen G., Planque B., Belgrano A., Post E., Reid P.C. & Stenseth N.C. (2001) Ecological effects of the North Atlantic Oscillation. *Oecologia*, **128**, 1-14.
- Portis D.H., Walsh J.E., El Hamly M. & Lamb P.J. (2001) Seasonality of the North Atlantic Oscillation. *J Clim*, **14**, 2069-2078.
- Rahmstorf S., Box J.E., Feulner G., Mann M.E., Robinson A., Rutherford S. & Schaffernicht E.J. (2015) Exceptional twentieth-century slowdown in Atlantic Ocean overturning circulation. *Nat Clim Chan*, **5**, 475-480
- Riksskogstaxeringen (2016) *Statistik om skog*. Available at: <http://www.slu.se/riksskogstaxeringen> (accessed 10 09 2014).
- Robitaille A. & Saucier J.-P. (1998) *Paysages régionaux du Québec méridional*. Les Publications du Québec, Québec.
- Rossi S., Girard M.J. & Morin H. (2014) Lengthening of the duration of xylogenesis engenders disproportionate increases in xylem production. *Glob Chang Bio*, **20**, 2261-2271
- Serreze M.C., Barrett A.P., Stroeve J.C., Kindig D.N. and Holland M.M. (2009) The emergence of surface-based Arctic amplification, *The Cryosphere*, **3**, 11-19, <https://doi.org/10.5194/tc-3-11-2009>.
- Sgubin G., Swingedouw D., Drijfhout S., Mary Y. & Bennadi A. (2017) Abrupt cooling over the North Atlantic in modern climate models. *Nat Commun*, **8**, 1-12.
- Skinner W.R., Stocks B.J., Martell D.L., Bonsal B. & Shabbar A. (1999) The Association Between Circulation Anomalies in the Mid-Troposphere and Area Burned by Wildland Fire in Canada. *Theor Appl Climatol*, **63**, 89-105.
- Solberg B.O., Hofgaard A. & Hytteborn H. (2002) Shifts in radial growth responses of coastal *Picea abies* induced by climatic change during the 20th century, central Norway. *Ecoscience* **9**, 79-88.
- Student (1908). The probable error of the mean. *Biometrika*, **6**, 1-25.
- Sveriges meteorologiska och hydrologiska institut (SMHI) (2016) *Sveriges Meteorologi: Temperatur och Nederbörd över 1961-1990*. Available at: <http://www.smhi.se/klimatdata/meteorologi/> (accessed 15 June 2016).
- Taylor A.H. & Stephens J.A. (1998) The North Atlantic Oscillation and the latitude of the Gulf Stream. *Tellus*, **50A**, 134-141.

- Trouet V. & Van Oldenborgh G.J. (2013) KNMI Climate Explorer: A web-based research tool for high-resolution paleoclimatology. *Tree-Ring Res*, **69**, 3-13.
- Trouet V., Babst F. & Meko M. (2018) Recent enhanced high-summer North Atlantic Jet variability emerges from three-century context. *Nat Commun*, **9**: 180
- Walker G.T. (1924) Correlation in seasonal variation of weather. IX. A further study of world weather. *Memoirs of the India Meteorological Department*, **24**, 275–333.
- Wallace J.M. & Gutzler D.S. (1981) Teleconnections in the geopotential height field during the Northern Hemisphere Winter. *Mon Weather Rev*, **109**, 784-812.
- Wassenburg J.A., Dietrich S., Fietzke J., Fohlmeister J., Jochum K.P., Scholz D., Richter D.K., Sabaoui A., Spotl C., Lohmann G., Andreae M.O. & Immenhauser A. (2016) Reorganization of the North Atlantic Oscillation during early Holocene deglaciation. *Nature Geosci*, **9**, 602-605.
- Wettstein J.J. & Mearns L.O. (2002) The influence of the North Atlantic–Arctic Oscillation on mean, variance, and extremes of temperature in the Northeastern United States and Canada. *J Clim*, **15**, 3582-3600.
- Wilmking M., Juday G.P., Barber V.A. & Zald H.S.J. (2004) Recent climate warming forces contrasting growth responses of white spruce at treeline in Alaska through temperature thresholds. *Glob Chang Biol*, **10**, 1724-1736.
- Zang C. & Biondi F. (2015) treeclim: an R package for the numerical calibration of proxy-climate relationships. *Ecography*, **38**, 431-436.
- Zhang Y., Bergeron Y., Zhao X.-H. & Drobyshev I. (2015) Stand history is more important than climate in controlling red maple (*Acer rubrum* L.) growth at its northern distribution limit in western Quebec, Canada. *J Plant Ecol*, **8**, 368-379.
- Zickfeld K., Eby M. & Weaver A.J. (2008) Carbon-cycle feedbacks of changes in the Atlantic meridional overturning circulation under future atmospheric CO₂. *Glob Biogeochem Cycles*, **22**.