



HAL
open science

Notes on two kinds of special values for the Bell polynomials of the second kind

Feng Qi, Dongkyu Lim, Yong-Hong Yao

► **To cite this version:**

Feng Qi, Dongkyu Lim, Yong-Hong Yao. Notes on two kinds of special values for the Bell polynomials of the second kind: Notes on special values of the Bell polynomials. *Miskolc Mathematical Notes*, 2019, 20 (1), pp.465–474. 10.18514/MMN.2019.2635 . hal-01757740v2

HAL Id: hal-01757740

<https://hal.science/hal-01757740v2>

Submitted on 18 Jan 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

NOTES ON TWO KINDS OF SPECIAL VALUES FOR THE BELL POLYNOMIALS OF THE SECOND KIND

FENG QI, DONGKYU LIM, AND YONG-HONG YAO

ABSTRACT. In the paper, by methods and techniques in combinatorial analysis and the theory of special functions, the authors discuss two kinds of special values for the Bell polynomials of the second kind for two special sequences, find a relation between these two kinds of special values for the Bell polynomials of the second kind, and derive an identity involving the combinatorial numbers.

1. MOTIVATION

In [1, Definition 11.2] and [2, p. 134, Theorem A], the Bell polynomials of the second kind, denoted by $B_{n,k}(x_1, x_2, \dots, x_{n-k+1})$ for $n \geq k \geq 0$, are defined by

$$B_{n,k}(x_1, x_2, \dots, x_{n-k+1}) = \sum_{\substack{1 \leq i \leq n-k+1 \\ \ell_i \in \{0\} \cup \mathbb{N} \\ \sum_{i=1}^{n-k+1} i \ell_i = n \\ \sum_{i=1}^{n-k+1} \ell_i = k}} \frac{n!}{\prod_{i=1}^{n-k+1} \ell_i!} \prod_{i=1}^{n-k+1} \left(\frac{x_i}{i!}\right)^{\ell_i}.$$

For more information on the Bell polynomials of the second kind $B_{n,k}$, please refer to the monographs and handbooks [1–3] and closely related references therein.

In [1, p. 451], the formulas

$$B_{2r,k}(0, 2!, \dots, 0, (2r)!) = \frac{(2r)!}{k!} \binom{r-1}{k-1},$$

$$B_{2r-1,k}(0, 2!, \dots, (2r-2)!, 0) = 0,$$

2010 *Mathematics Subject Classification*. Primary 11B83; Secondary 11C08, 12E10, 26C05.

Key words and phrases. special value; Bell polynomial of the second kind; factorial; relation; Faà di Bruno formula; identity; beta function; combinatorial analysis; Vandermonde convolution formula.

Please cite this article as “Feng Qi, Dongkyu Lim, and Yong-Hong Yao, *Notes on two kinds of special values for the Bell polynomials of the second kind*, Miskolc Mathematical Notes **20** (2019), no. 1, 465–474; available online at <https://doi.org/10.18514/MMN.2019.2635>.”

$$\begin{aligned}
B_{2r,2s}(1!, 0, \dots, (2r-1)!, 0) &= \frac{(2r)!}{(2s)!} \binom{r+s-1}{2s-1}, \\
B_{2r,2s-1}(1!, 0, \dots, (2r-1)!, 0) &= 0, \\
B_{2r-1,2s-1}(1!, 0, \dots, (2r-1)!, 0) &= \frac{(2r-1)!}{(2s-1)!} \binom{r+s-2}{2s-2}, \\
B_{2r-1,2s}(1!, 0, \dots, 0, (2r-1)!) &= 0
\end{aligned}$$

were stated, but no proof was supplied for them there.

For simplicity, we denote

$$\lambda(n, k) = B_{n,k} \left(0, 2!, 0, 4!, \dots, (n-k+1)! \frac{1 + (-1)^{n-k+1}}{2} \right)$$

and

$$\mu(n, k) = B_{n,k} \left(1!, 0, 3!, 0, \dots, (n-k+1)! \frac{1 - (-1)^{n-k+1}}{2} \right).$$

In terms of these notations, the above claims in [1, p. 451] can be restated as

$$\begin{aligned}
\lambda(2r, k) &= \frac{(2r)!}{k!} \binom{r-1}{k-1}, & \lambda(2r-1, k) &= 0, \\
\mu(2r, 2s) &= \frac{(2r)!}{(2s)!} \binom{r+s-1}{2s-1}, & \mu(2r, 2s-1) &= 0, \\
\mu(2r-1, 2s-1) &= \frac{(2r-1)!}{(2s-1)!} \binom{r+s-2}{2s-2}, & \mu(2r-1, 2s) &= 0.
\end{aligned}$$

In this paper, by methods and techniques in combinatorial analysis and the theory of special functions, we will provide alternative proofs for the above six formulas, find a relation between them, and derive an identity involving combinatorial numbers.

2. MAIN RESULTS

We first derive an identity involving combinatorial numbers, which will be useful in next proofs of our main results.

Theorem 1. *For $k \geq 1$ and $n \geq 0$, we have*

$$\sum_{q=0}^n \frac{(-1)^q}{k+q} \binom{n}{q} = \frac{1}{k \binom{k+n}{k}}. \quad (2.1)$$

Proof. Let

$$f(x) = \sum_{q=0}^n \frac{(-1)^q}{k+q} \binom{n}{q} x^{k+q}, \quad x \in [-1, 1].$$

Then $f(0) = 0$ and

$$f'(x) = \sum_{q=0}^n (-1)^q \binom{n}{q} x^{k+q-1} = x^{k-1} \sum_{q=0}^n (-1)^q \binom{n}{q} x^q = x^{k-1} (1-x)^n.$$

Integrating from 0 to $t \in (0, 1]$ on both sides of the above equality yields

$$f(t) = \int_0^t x^{k-1} (1-x)^n dx = B(t; k, n+1),$$

where $B(z; a, b)$ denotes the incomplete beta function, see [3, p. 183]. Therefore, we obtain

$$\begin{aligned} f(1) &= B(1; k, n+1) = B(k, n+1) \\ &= \frac{\Gamma(k)\Gamma(n+1)}{\Gamma(k+n+1)} = \frac{(k-1)!n!}{(k+n)!} = \frac{1}{k} \frac{k!n!}{(k+n)!} = \frac{1}{k \binom{n+k}{k}}, \end{aligned}$$

where $B(a, b)$ and $\Gamma(z)$ denote the beta function and the classical Euler gamma function respectively, see [3, p. 142] and [3, Chapter 5]. The formula (2.1) is thus proved. The proof of Theorem 1 is complete. \square

We are now in a position to state and prove our main results.

Theorem 2. *For $n \geq k \geq 0$, we have the relation*

$$\frac{\mu(n, k)}{n!} = \frac{\lambda(n+k, k)}{(n+k)!} \quad (2.2)$$

and two explicit formulas

$$\lambda(n, k) = \frac{1 + (-1)^n}{2} \frac{n!}{k!} \binom{\frac{n}{2} - 1}{k-1} \quad (2.3)$$

and

$$\mu(n, k) = \frac{1 + (-1)^{n+k}}{2} \frac{n!}{k!} \binom{\frac{n+k}{2} - 1}{k-1}. \quad (2.4)$$

Proof. It is known [2, 13] that the quantities

$$\langle x \rangle_n = \prod_{\ell=0}^{n-1} (x - \ell) = \begin{cases} x(x-1) \cdots (x-n+1), & n \geq 1 \\ 1, & n = 0 \end{cases}$$

and

$$(x)_n = \prod_{\ell=0}^{n-1} (x + \ell) = \begin{cases} x(x+1) \cdots (x+n-1), & n \geq 1 \\ 1, & n = 0 \end{cases}$$

are called the falling and rising factorials respectively. In [2, p. 133], it was listed that

$$\frac{1}{k!} \left(\sum_{m=1}^{\infty} x_m \frac{t^m}{m!} \right)^k = \sum_{n=k}^{\infty} B_{n,k}(x_1, x_2, \dots, x_{n-k+1}) \frac{t^n}{n!} \quad (2.5)$$

for $k \geq 0$. Hence, we have

$$\sum_{n=k}^{\infty} \lambda(n, k) \frac{t^n}{n!} = \frac{1}{k!} \left[\sum_{m=1}^{\infty} m! \frac{1 + (-1)^m t^m}{2} \frac{t^m}{m!} \right]^k = \frac{1}{k!} \left(\frac{t^2}{1-t^2} \right)^k$$

which is equivalent to

$$\sum_{n=0}^{\infty} \lambda(n+k, k) \frac{1}{\binom{n+k}{n}} \frac{t^n}{n!} = \left(\frac{t}{1-t^2} \right)^k.$$

Since the function $\frac{t}{1-t^2}$ is odd, we derive $\lambda(2r-1, k) = 0$. Further computation yields

$$\begin{aligned} \lambda(n+k, k) &= \binom{n+k}{n} \lim_{t \rightarrow 0} \frac{d^n}{dt^n} \left(\frac{t}{1-t^2} \right)^k \\ &= \binom{n+k}{n} \frac{1}{2^k} \lim_{t \rightarrow 0} \left[\left(\frac{1}{1-t} - \frac{1}{1+t} \right)^k \right]^{(n)} \\ &= \binom{n+k}{n} \frac{1}{2^k} \lim_{t \rightarrow 0} \left[\sum_{\ell=0}^k \binom{k}{\ell} \left(\frac{1}{1-t} \right)^\ell (-1)^{k-\ell} \left(\frac{1}{1+t} \right)^{k-\ell} \right]^{(n)} \\ &= \binom{n+k}{n} \frac{1}{2^k} \sum_{\ell=0}^k (-1)^{k-\ell} \binom{k}{\ell} \lim_{t \rightarrow 0} \sum_{q=0}^n \binom{n}{q} \left[\left(\frac{1}{1-t} \right)^\ell \right]^{(q)} \left[\left(\frac{1}{1+t} \right)^{k-\ell} \right]^{(n-q)} \\ &= \binom{n+k}{n} \frac{1}{2^k} \sum_{\ell=0}^k (-1)^{k-\ell} \binom{k}{\ell} \lim_{t \rightarrow 0} \sum_{q=0}^n \binom{n}{q} \\ &\quad \times \langle -\ell \rangle_q (-1)^q \left(\frac{1}{1-t} \right)^{\ell+q} \langle -(k-\ell) \rangle_{n-q} \left(\frac{1}{1+t} \right)^{k-\ell+(n-q)} \\ &= \frac{(-1)^k}{2^k} \binom{n+k}{n} \sum_{\ell=0}^k (-1)^\ell \binom{k}{\ell} \sum_{q=0}^n (-1)^q \binom{n}{q} \langle -\ell \rangle_q \langle \ell-k \rangle_{n-q} \\ &= \frac{(-1)^{n+k}}{2^k} \binom{n+k}{n} \sum_{\ell=0}^k (-1)^\ell \binom{k}{\ell} \sum_{q=0}^n (-1)^q \binom{n}{q} (\ell)_q (k-\ell)_{n-q}. \end{aligned}$$

In summary, we obtain

$$\begin{aligned} \lambda(n, k) &= \frac{(-1)^k}{2^k} \binom{n}{k} \sum_{\ell=0}^k (-1)^\ell \binom{k}{\ell} \sum_{q=0}^{n-k} (-1)^q \binom{n-k}{q} \langle -\ell \rangle_q \langle \ell-k \rangle_{n-k-q} \\ &= \frac{(-1)^n}{2^k} \binom{n}{k} \sum_{\ell=0}^k (-1)^\ell \binom{k}{\ell} \sum_{q=0}^{n-k} (-1)^q \binom{n-k}{q} (\ell)_q (k-\ell)_{n-k-q}. \quad (2.6) \end{aligned}$$

By similar argument as above, it follows that

$$\sum_{n=k}^{\infty} \mu(n, k) \frac{t^n}{n!} = \frac{1}{k!} \left[\sum_{m=1}^{\infty} m! \frac{1 - (-1)^m t^m}{2} \frac{1}{m!} \right]^k = \frac{(-1)^k}{k! 2^k} \left(\frac{1}{t-1} + \frac{1}{t+1} \right)^k$$

which is equivalent to

$$\sum_{n=0}^{\infty} \mu(n+k, k) \frac{1}{\binom{n+k}{n}} \frac{t^n}{n!} = \frac{(-1)^k}{2^k} \frac{1}{t^k} \left(\frac{1}{t-1} + \frac{1}{t+1} \right)^k = \left(\frac{1}{1-t^2} \right)^k. \quad (2.7)$$

Since $\frac{1}{1-t^2}$ is even, we deduce immediately that $\mu(2r-1, 2s) = 0$ and $\mu(2r, 2s-1) = 0$. By the L'Hôpital rule and the identity (2.1) in Theorem 1, it follows that

$$\begin{aligned} \mu(n+k, k) &= \binom{n+k}{n} \frac{(-1)^k}{2^k} \lim_{t \rightarrow 0} \frac{d^n}{d t^n} \left[\frac{1}{t^k} \left(\frac{1}{t-1} + \frac{1}{t+1} \right)^k \right] \\ &= \binom{n+k}{n} \frac{(-1)^k}{2^k} \lim_{t \rightarrow 0} \sum_{q=0}^n \binom{n}{q} \frac{\langle -k \rangle_q}{t^{k+q}} \left[\left(\frac{1}{t-1} + \frac{1}{t+1} \right)^k \right]^{(n-q)} \\ &= \binom{n+k}{n} \frac{(-1)^k}{2^k} \sum_{q=0}^n \binom{n}{q} \frac{\langle -k \rangle_q}{(k+q)!} \lim_{t \rightarrow 0} \left[\left(\frac{1}{t-1} + \frac{1}{t+1} \right)^k \right]^{(n+k)} \\ &= \binom{n+k}{n} \frac{(-1)^k}{2^k} \sum_{q=0}^n \binom{n}{q} \frac{(-1)^q (k+q-1)!}{(k-1)! (k+q)!} \\ &\quad \times \lim_{t \rightarrow 0} \left[\sum_{\ell=0}^k \binom{k}{\ell} \left(\frac{1}{t-1} \right)^\ell \left(\frac{1}{t+1} \right)^{k-\ell} \right]^{(n+k)} \\ &= \binom{n+k}{n} \frac{(-1)^k}{2^k (k-1)!} \sum_{q=0}^n \frac{(-1)^q}{k+q} \binom{n}{q} \sum_{\ell=0}^k \binom{k}{\ell} \\ &\quad \times \lim_{t \rightarrow 0} \sum_{m=0}^{n+k} \binom{n+k}{m} \left[\left(\frac{1}{t-1} \right)^\ell \right]^{(m)} \left[\left(\frac{1}{t+1} \right)^{k-\ell} \right]^{(n+k-m)} \\ &= \frac{(-1)^k}{2^k k!} \sum_{\ell=0}^k \binom{k}{\ell} \lim_{t \rightarrow 0} \sum_{m=0}^{n+k} \binom{n+k}{m} \\ &\quad \times \langle -\ell \rangle_m \left(\frac{1}{t-1} \right)^{\ell+m} \langle \ell - k \rangle_{n+k-m} \left(\frac{1}{t+1} \right)^{(k-\ell)+(n+k-m)} \\ &= \frac{(-1)^k}{2^k k!} \sum_{\ell=0}^k (-1)^\ell \binom{k}{\ell} \sum_{m=0}^{n+k} (-1)^m \binom{n+k}{m} \langle -\ell \rangle_m \langle \ell - k \rangle_{n+k-m}. \end{aligned}$$

In short, we obtain

$$\begin{aligned}
\mu(n, k) &= \frac{(-1)^k}{2^k k!} \sum_{\ell=0}^k (-1)^\ell \binom{k}{\ell} \sum_{m=0}^n (-1)^m \binom{n}{m} \langle -\ell \rangle_m \langle \ell - k \rangle_{n-m} \\
&= \frac{(-1)^{n+k}}{2^k k!} \sum_{\ell=0}^k (-1)^\ell \binom{k}{\ell} \sum_{m=0}^n (-1)^m \binom{n}{m} (\ell)_m (k - \ell)_{n-m}. \quad (2.8)
\end{aligned}$$

Comparing between (2.6) and (2.8) reveals

$$\mu(n, k) = \frac{(-1)^{n+k}}{2^k k!} \frac{2^k}{(-1)^{n+k}} \frac{1}{\binom{n+k}{k}} \lambda(n+k, k)$$

which can be rearranged as (2.2).

The Faà di Bruno formula, see [1, Theorem 11.4] and [2, p. 139, Theorem C], can be described in terms of $B_{n,k}(x_1, x_2, \dots, x_{n-k+1})$ by

$$\frac{d^n}{dx^n} f \circ h(x) = \sum_{k=0}^n f^{(k)}(h(x)) B_{n,k}(h'(x), h''(x), \dots, h^{(n-k+1)}(x)). \quad (2.9)$$

In [1, p. 412] and [2, p. 135], one can find the identity

$$\begin{aligned}
B_{n,k}(abx_1, ab^2x_2, \dots, ab^{n-k+1}x_{n-k+1}) \\
= a^k b^n B_{n,k}(x_1, x_2, \dots, x_{n-k+1}) \quad (2.10)
\end{aligned}$$

for $n \geq k \geq 0$ and $a, b \in \mathbb{C}$. In [8, Theorem 5.1] and [15, Section 3], it was set up little by little that

$$B_{n,k}(x, 1, 0, \dots, 0) = \frac{1}{2^{n-k}} \frac{n!}{k!} \binom{k}{n-k} x^{2k-n}, \quad (2.11)$$

where $\binom{0}{0} = 1$ and $\binom{p}{q} = 0$ for $q > p \geq 0$. For detailed information on applications of the formula (2.11), please refer to the papers [4–8, 12, 14–17] and closely related references therein. Then it follows from (2.7), (2.9), (2.10), and (2.11) that, when denoting $u = u(t) = 1 - t^2$,

$$\begin{aligned}
\mu(n+k, k) &= \binom{n+k}{n} \lim_{t \rightarrow 0} \frac{d^n}{dt^n} \left(\frac{1}{1-t^2} \right)^k \\
&= \binom{n+k}{n} \lim_{t \rightarrow 0} \sum_{\ell=0}^n \left(\frac{1}{u^k} \right)^{(\ell)} B_{n,\ell}(-2t, -2, 0, \dots, 0) \\
&= \binom{n+k}{n} \lim_{t \rightarrow 0} \sum_{\ell=0}^n \langle -k \rangle_\ell \frac{1}{u^{k+\ell}} (-2)^\ell B_{n,\ell}(t, 1, 0, \dots, 0) \\
&= \binom{n+k}{n} \lim_{t \rightarrow 0} \sum_{\ell=0}^n \langle -k \rangle_\ell \frac{1}{(1-t^2)^{k+\ell}} (-2)^\ell \frac{1}{2^{n-\ell}} \frac{n!}{\ell!} \binom{\ell}{n-\ell} t^{2\ell-n}
\end{aligned}$$

$$\begin{aligned}
 &= \binom{n+k}{n} \lim_{t \rightarrow 0} \sum_{\ell=0}^n \binom{k}{\ell} \frac{n!}{\ell!} \binom{\ell}{n-\ell} \frac{(2t)^{2\ell-n}}{(1-t^2)^{k+\ell}} \\
 &= \begin{cases} 0, & n = 2m-1 \\ \binom{2m+k}{2m} \binom{k}{m} \frac{(2m)!}{m!} \binom{m}{2m-m}, & n = 2m \end{cases} \\
 &= \begin{cases} 0, & n = 2m-1 \\ \frac{(2m+k)!}{k!} \binom{m+k-1}{k-1}, & n = 2m \end{cases} \\
 &= \begin{cases} 0, & n = 2m-1 \\ \frac{(n+k)!}{k!} \binom{\frac{n+2k}{2}-1}{k-1}, & n = 2m \end{cases}
 \end{aligned}$$

for $m \in \mathbb{N}$. The formula in (2.4) is thus proved.

Substituting the formula in (2.4) into (2.2) and simplifying lead readily to the formula in (2.3). The proof of Theorem 2 is complete. \square

3. REMARKS

In this section, we state several remarks on something related.

Remark 1. In [8, Theorem 2.1], it was proved that

$$B_{n,k} \left(1, 0, 1, \dots, \frac{1 - (-1)^{n-k+1}}{2} \right) = \frac{1}{2^k k!} \sum_{\ell=0}^k (-1)^\ell \binom{k}{\ell} (k-2\ell)^n$$

and

$$B_{n,k} \left(0, 1, 0, \dots, \frac{1 + (-1)^{n-k+1}}{2} \right) = \frac{1}{2^k k!} \sum_{\ell=0}^{2k} (-1)^\ell \binom{2k}{\ell} (k-\ell)^n$$

for $n \geq k \geq 0$, where 0^0 is regarded as 1. In [8, Section 3], basing on numerical calculation, the authors guessed that

$$B_{2\ell-1,k} \left(0, 1, 0, \dots, \frac{1 + (-1)^{(2\ell-1)-k+1}}{2} \right) = 0, \quad 2\ell-1 \geq k \geq 0, \quad (3.1)$$

$$B_{2\ell,k} \left(0, 1, 0, \dots, \frac{1 - (-1)^{2\ell-k+1}}{2} \right) = 0, \quad 2\ell > k > \ell, \quad (3.2)$$

$$B_{2\ell,k} \left(0, 1, 0, \dots, \frac{1 - (-1)^{2\ell-k+1}}{2} \right) \neq 0, \quad \ell \geq k \geq 1, \quad (3.3)$$

$$B_{k+2\ell,k}(1, 0, 1, \dots, 1, 0, 1) \neq 0, \quad k, \ell \in \mathbb{N}, \quad (3.4)$$

and

$$B_{k+2\ell-1,k}(1, 0, 1, \dots, 0, 1, 0) = 0, \quad k, \ell \in \mathbb{N}. \quad (3.5)$$

In [8, Theorem 3.1], the identity (3.5) was proved to be true. In fact, the identities (3.1) and (3.5) can be concluded readily from the proof of [8, Theorem 2.1] as follows. From the formula (2.5), it followed that

$$\sum_{n=k}^{\infty} B_{n,k} \left(0, 1, 0, \dots, \frac{1 + (-1)^{n-k+1}}{2} \right) \frac{t^n}{n!} = \frac{(\cosh t - 1)^k}{k!} \quad (3.6)$$

and

$$\sum_{n=k}^{\infty} B_{n,k} \left(1, 0, 1, \dots, \frac{1 - (-1)^{n-k+1}}{2} \right) \frac{t^n}{n!} = \frac{\sinh^k t}{k!}. \quad (3.7)$$

In (3.6), the function $\cosh t - 1$ is even, so the identity (3.1) is clearly valid. Since the function $\sinh t$ in (3.7) is odd, then

$$B_{2n,2k-1}(1, 0, 1, \dots, 1, 0) = 0 \quad \text{and} \quad B_{2n-1,2k}(1, 0, 1, \dots, 1, 0) = 0$$

which are equivalent to the identity (3.5). However, the validity of the identities (3.2), (3.3), and (3.4) has not been verified yet.

Remark 2. The first formula in (2.3) of Theorem 2 was also discussed in [9, Lemma 2.5], [11] and closely related references therein.

Remark 3. The formula (2.8) can be rewritten as

$$\begin{aligned} \mu(n, k) &= \frac{(-1)^{n+k} n!}{2^k k!} \\ &\times \sum_{\ell=0}^k \sum_{m=0}^n (-1)^{\ell+m} \binom{k}{\ell} \binom{\ell+m-1}{\ell-1} \binom{(k-\ell)+(n-m)-1}{k-\ell-1}. \end{aligned} \quad (3.8)$$

Then combining the formulas (2.8) and (3.8) with the formula in (2.4) and rearranging arrive at

$$\begin{aligned} \sum_{\ell=0}^k (-1)^{\ell} \binom{k}{\ell} \sum_{m=0}^n (-1)^m \binom{n}{m} (\ell)_m (k-\ell)_{n-m} \\ = [1 + (-1)^{n+k}] 2^{k-1} n! \binom{\frac{n+k}{2} - 1}{k-1} \end{aligned}$$

and

$$\begin{aligned} \sum_{m=0}^n \sum_{\ell=0}^k (-1)^{\ell+m} \binom{k}{\ell} \binom{\ell+m-1}{\ell-1} \binom{(k-\ell)+(n-m)-1}{k-\ell-1} \\ = [1 + (-1)^{n+k}] 2^{k-1} \binom{\frac{n+k}{2} - 1}{k-1}. \end{aligned}$$

Comparing these identities with the Vandermonde convolution formula

$$\sum_{k=0}^n \binom{n}{k} \langle x \rangle_k \langle y \rangle_{n-k} = \langle x + y \rangle_n$$

in [1, Theorem 3.1] and [2, p. 44] motivates us to ask a question: is the quantity

$$\sum_{k=0}^n (-1)^k \binom{n}{k} \langle x \rangle_k \langle y \rangle_{n-k}$$

summable?

Remark 4. This paper is a slightly revised version of the preprint [10].

Acknowledgements. The second author was supported by the National Research Foundation of Korea (Grant No. 2018R1D1A1B07041846).

REFERENCES

- [1] C. A. Charalambides, *Enumerative Combinatorics*, CRC Press Series on Discrete Mathematics and its Applications. Chapman & Hall/CRC, Boca Raton, FL, 2002.
- [2] L. Comtet, *Advanced Combinatorics: The Art of Finite and Infinite Expansions*, Revised and Enlarged Edition, D. Reidel Publishing Co., Dordrecht and Boston, 1974; available online at <https://doi.org/10.1007/978-94-010-2196-8>.
- [3] F. W. J. Olver, D. W. Lozier, R. F. Boisvert, and C. W. Clark (eds.), *NIST Handbook of Mathematical Functions*, Cambridge University Press, New York, 2010; available online at <http://dlmf.nist.gov/>.
- [4] F. Qi, *Integral representations for multivariate logarithmic polynomials*, J. Comput. Appl. Math. **336** (2018), 54–62; available online at <https://doi.org/10.1016/j.cam.2017.11.047>.
- [5] F. Qi and V. Čerňanová, *Some discussions on a kind of improper integrals*, Int. J. Anal. Appl. **11** (2016), no. 2, 101–109.
- [6] F. Qi, V. Čerňanová, X.-T. Shi, and B.-N. Guo, *Some properties of central Delannoy numbers*, J. Comput. Appl. Math. **328** (2018), 101–115; available online at <https://doi.org/10.1016/j.cam.2017.07.013>.
- [7] F. Qi and B.-N. Guo, *Explicit and recursive formulas, integral representations, and properties of the large Schröder numbers*, Kragujevac J. Math. **41** (2017), no. 1, 121–141; available online at <https://doi.org/10.5937/KgJMath1701121F>.
- [8] F. Qi and B.-N. Guo, *Explicit formulas for special values of the Bell polynomials of the second kind and for the Euler numbers and polynomials*, Mediterr. J. Math. **14** (2017), no. 3, Article 140, 14 pages; available online at <https://doi.org/10.1007/s00009-017-0939-1>.
- [9] F. Qi and B.-N. Guo, *Some properties of the Hermite polynomials*, Georgian Math. J. **27** (2020), in press; available online at <https://doi.org/10.20944/preprints201611.0145.v1>.
- [10] F. Qi, D. Lim, and Y.-H. Yao, *Notes on two kinds of special values for the Bell polynomials of the second kind*, HAL preprint (2018), available online at <https://hal.archives-ouvertes.fr/hal-01757740>.
- [11] F. Qi, D.-W. Niu, D. Lim, and Y.-H. Yao, *Special values of the Bell polynomials of the second kind for some sequences and functions*, HAL preprint (2018), available online at <https://hal.archives-ouvertes.fr/hal-01766566>.

- [12] F. Qi, X.-T. Shi, and B.-N. Guo, *Two explicit formulas of the Schröder numbers*, *Integers* **16** (2016), Paper No. A23, 15 pages.
- [13] F. Qi, X.-T. Shi, and F.-F. Liu, *Several identities involving the falling and rising factorials and the Cauchy, Lah, and Stirling numbers*, *Acta Univ. Sapientiae Math.* **8** (2016), no. 2, 282–297; available online at <https://doi.org/10.1515/ausm-2016-0019>.
- [14] F. Qi, X.-T. Shi, F.-F. Liu, and D. V. Kruchinin, *Several formulas for special values of the Bell polynomials of the second kind and applications*, *J. Appl. Anal. Comput.* **7** (2017), no. 3, 857–871; available online at <https://doi.org/10.11948/2017054>.
- [15] F. Qi and M.-M. Zheng, *Explicit expressions for a family of the Bell polynomials and applications*, *Appl. Math. Comput.* **258** (2015), 597–607; available online at <https://doi.org/10.1016/j.amc.2015.02.027>.
- [16] C.-F. Wei and F. Qi, *Several closed expressions for the Euler numbers*, *J. Inequal. Appl.* **2015**, 2015:219, 8 pages; available online at <https://doi.org/10.1186/s13660-015-0738-9>.
- [17] J.-L. Zhao and F. Qi, *Two explicit formulas for the generalized Motzkin numbers*, *J. Inequal. Appl.* **2017**, 2017:44, 8 pages; available online at <https://doi.org/10.1186/s13660-017-1313-3>.

Authors' addresses

Feng Qi

INSTITUTE OF MATHEMATICS, HENAN POLYTECHNIC UNIVERSITY, JIAOZUO 454010, HENAN, CHINA; COLLEGE OF MATHEMATICS, INNER MONGOLIA UNIVERSITY FOR NATIONALITIES, TONGLIAO 028043, INNER MONGOLIA, CHINA; SCHOOL OF MATHEMATICAL SCIENCES, TIANJIN POLYTECHNIC UNIVERSITY, TIANJIN 300387, CHINA

Email address: qifeng618@gmail.com, qifeng618@hotmail.com, qifeng618@qq.com

URL: <https://qifeng618.wordpress.com>, <http://orcid.org/0000-0001-6239-2968>

Dongkyu Lim

DEPARTMENT OF MATHEMATICS EDUCATION, ANDONG NATIONAL UNIVERSITY, ANDONG 36729, REPUBLIC OF KOREA

Email address: dgrim84@gmail.com, dklim@andong.ac.kr

URL: <http://orcid.org/0000-0002-0928-8480>

Yong-Hong Yao

SCHOOL OF MATHEMATICAL SCIENCES, TIANJIN POLYTECHNIC UNIVERSITY, TIANJIN 300387, CHINA

Email address: yaoyonghong@aliyun.com

URL: <https://orcid.org/0000-0002-0452-785X>