This file is the postprint of the article accepted in the Journal of Computational Physics accessible throught the following DOI: https://doi.org/10.1016/j.jcp.2018.03.027. To access freely the final version until May 24, 2018: https://authors.elsevier.com/a/1WqZA508HeRgq

Cite as: M. Pigou, J. Morchain, P. Fede, M. I. Penet, G. Laronze, New developments of the Extended Quadrature Method of Moments to solve Population Balance Equations, J. Comput. Phys. 365 (2018) 243-268, ISSN 0021-9991, doi:10.1016/ j.jcp.2018.03.027

New developments of the Extended Quadrature Method of Moments to solve Population Balance Equations

Maxime PIGOU^{a,b,*}, Jérôme MORCHAIN^a, Pascal FEDE^b, Marie-Isabelle PENET^c, Geoffrey LARONZE^c

^aLISBP, Université de Toulouse, CNRS, INRA, INSA, Toulouse, France

^bInstitut de Mécanique des Fluides de Toulouse - Université de Toulouse, CNRS-INPT-UPS, Toulouse,

France

^cSanofi Chimie - C&BD Biochemistry Vitry - 9 quai Jules Guesde 94400 Vitry-sur-Seine, France

Abstract

Population Balance Models have a wide range of applications in many industrial fields as they allow accounting for heterogeneity among properties which are crucial for some system modelling. They actually describe the evolution of a Number Density Function (NDF) in a Population Balance Equation (PBE). For instance, they are applied to gas-liquid columns or stirred reactors, aerosol technology, crystallisation processes, fine particles or biological systems. There is a significant interest for fast, stable and accurate numerical methods in order to solve for PBEs, a class of such methods actually does not solve directly the NDF but resolves their moments. These methods of moments, and in particular quadraturebased methods of moments, have been successfully applied to a variety of systems. Pointwise values of the NDF are sometimes required but are not directly accessible from the moments. To address these issues, the Extended Quadrature Method of Moments (EQMOM) has been developed in the past few years and approximates the NDF, from its moments, as a convex mixture of Kernel Density Functions (KDFs) of the same parametric family. In the present work EQMOM is further developed on two aspects. The main one is a significant improvement of the core iterative procedure of that method, the corresponding reduction of its computational cost is estimated to range from 60% up to 95%. The second aspect is an extension of EQMOM to two new KDFs used for the approximation, the Weibull and the Laplace kernels. All MATLAB source codes used for this article are provided with this article.

Keywords: Extended Quadrature Method of Moments (EQMOM), Quadrature Based Method of Moments (QBMM), Population Balance, Mathematical modelling, Gauss quadrature

1 1. Introduction

Population Balance Equations (PBEs) are particular formalisms that allows describing
 the evolution of properties among heterogeneous populations. They are used to track the

Preprint submitted to Journal of Computational Physics

©2018. This manuscript version is made available under the CC-BY-NC-ND 4.0 license http://creativecommons.org/licenses/by-nc-nd/4.0/

^{*}Corresponding author, Phone: +33561559275

Email address: maxime.pigou@insa-toulouse.fr (Maxime PIGOU)

Nomenclature						
Greek symbols			orthogonal polynomials recurrence			
ε	relative tolerance		coefficient			
λ_j	j-th nested quadrature node	${\cal H}$	Hankel determinant			
μ	positive measure	$oldsymbol{J}_n$	n order Jacobi matrix			
ω_j	j-th nested quadrature weight	m_k	moment of order k			
Ω_{ξ}	NDF support	\mathcal{M}	realisable moment space			
π_k	k order orthogonal polynomial	n	number density function			
σ	shape parameter	\widetilde{a}	approximation of a			
ξ	random variable	n	approximation of n			
ξ_i	i-th main quadrature node	N	order of moment set			
ζ	realisability criteria on $]0,+\infty[$	\mathcal{N}	order of realisability			
Roman		p_k	canonical moment of order k			
a	orthogonal polynomials recurrence	P	number of main quadrature nodes			
	coefficient	0	number of nested quadrature nodes			
\boldsymbol{A}	transition matrix to degenerated mo-	~ 111.	<i>i</i> -th main quadrature weight			
	monto	ω_i	e-un main quadrature weight			

size distribution of fine particles [1]; the bubble size distribution in gas-liquid stirred-tank 4 reactors or bubble columns [2, 3]; the crystal-size distribution in crystallizers; the distribution 5 of biological cell properties in bioreactors [4, 5]; the volume and/or surface distribution of 6 soot particles in flames [6, 7] or the formation of nano-particles [8], among other examples. 7 A PBE describes the evolution and transport of a Number Density Function (NDF), 8 under the influence of multiple processes which modify the tracked property distribution 9 (e.q. erosion, dissolution, aggregation, breakage, coalescence, nucleation, adaptation, etc.). 10 One often requires low-cost numerical methods to solve PBEs, for instance when coupling 11 with a flow solver (e.g. Computational Fluid Dynamics software). Monte-Carlo methods 12 constitute a stochastic resolution of the population balance and can be applied to such 13 PBE-CFD simulations [9]. Similarly, sectional methods allow direct numerical resolutions of 14 the PBE through the discretisation of the property space [10, 11]. They respectively require 15 a high number of parcels or sections in order to reach high accuracy and are thus often 16 discarded for large-scale simulations. 17

An interesting alternative approach lies in the field of methods of moments. A PBE, which describes the evolution of a NDF, is transformed in a set of equations which describes the evolution of the moments of that distribution. Moments are integral properties of NDFs, the first low order integer moments are related to the mean, variance, skewness and flatness of the statistical distributions described by NDFs. This approach then reduces the number of resolved variables to a finite set of NDF moments. It also comes with some difficulties when one must compute non-moment integral properties, or point-wise evaluations, of the

²⁵ distribution [12].

To tackle these issues, one can try to recover a NDF from a finite set of its moments. In most cases, this reverse problem has an infinite number of solutions and different approaches exist to identify one or an other out of them. The simplest is probably to assume that the NDF is a standard distribution (Gaussian, Log-normal, ...) whose parameters will be deduced from its first few moments. Other methods that lead to continuous approximations, and which preserve a higher number of moments, are the Spline method [13], the Maximum-Entropy approach [12, 14, 15] or the Kernel Density Element Method (KDEM) [16].

³³ More recently, the Extended Quadrature Method of Moments (EQMOM) was proposed ³⁴ as a new approach which is more stable than the previous ones, and yields either continuous ³⁵ or discrete NDFs depending on the moments [1, 17, 18]. EQMOM has been implemented in ³⁶ OpenFOAM [19] for the purpose of PBE-CFD coupling. The core of this method relies on

an iterative procedure that is a computational bottleneck.

The current work focuses on EQMOM and develops a new core procedure whose computational cost is significantly lower than previous implementations by reducing both (i) the cost of each iteration and (ii) the total number of required iterations.

The previous core procedure [1] will be recalled before describing how it can be shifted toward the new –cheaper– approach. Both implementations will be compared in terms of computational cost (number of required floating-point operations) and run-time.

Multiple variations of EQMOM exist, the Gauss EQMOM [17, 20], Log-normal EQMOM
[21] as well as Gamma and Beta EQMOM [18]. Two new variations, namely Laplace EQMOM and Weibull EQMOM, are proposed along with a unified formalism among all six
variations.

The whole source code used to write this article (figures and data generation) is provided as supplementary data, as well as our implementations of EQMOM in the form of a MATLAB functions library [22].

⁵¹ 2. Quadrature Based Methods of Moments: QMOM and EQMOM

52 2.1. Definitions

Let $d\mu(\xi)$ be a positive measure, induced by a non-decreasing function $\mu(\xi)$ defined on a support Ω_{ξ} . This measure is associated to a Number Density Function $n(\xi)$ such that $d\mu(\xi) = n(\xi)d\xi$. Let \boldsymbol{m}_N be the vector of the first N + 1 integer moments of this measure:

$$\boldsymbol{m}_{N} = \begin{bmatrix} m_{0} \\ m_{1} \\ \vdots \\ m_{N} \end{bmatrix}, \qquad m_{k} = \int_{\Omega_{\xi}} \xi^{k} n(\xi) \mathrm{d}\xi \qquad (1)$$

Three actual supports will be considered: (i) $\Omega_{\xi} =]-\infty, +\infty[$, (ii) $\Omega_{\xi} =]0, +\infty[$ and (iii) $\Omega_{\xi} =]0, 1[$. For each support, one can define the associated realisable moment space, $\mathcal{M}_N(\Omega_{\xi})$, as the set of all vectors of finite moments \boldsymbol{m}_N induced by all possible positive measures defined on Ω_{ξ} . ⁶⁰ A moment set is said to be "weakly realisable" if located on the boundary of the realisable ⁶¹ moment space ($\mathbf{m}_N \in \partial \mathcal{M}_N(\Omega_{\xi})$). Otherwise, if located within the realisable moment space, ⁶² \mathbf{m}_N is said to be "strictly realisable".

63 2.2. Quadrature Method of Moments

EQMOM is based on the Quadrature Method of Moments (QMOM) that was first introduced by McGraw [23]. It is used to approximate integral properties of a distribution where only a finite number of its moments is known. By making use of an even number of moments 2P, one can compute a Gauss quadrature rule characterised by its weights $\boldsymbol{w}_P = [w_1, \ldots, w_P]^{\mathsf{T}}$ and nodes $\boldsymbol{\xi}_P = [\xi_1, \ldots, \xi_P]^{\mathsf{T}}$ such that:

$$\int_{\Omega_{\xi}} f(\xi) \mathrm{d}\mu(\xi) = \sum_{i=1}^{P} w_i f(\xi_i)$$
(2)

⁶⁹ holds true if $f(\xi) = \xi^k$, $\forall k \in \{0, ..., 2P-1\}$. Otherwise, this quadrature rule will produce ⁷⁰ an approximation of the integral property. The computation of the quadrature rule (*i.e.* ⁷¹ the vectors \boldsymbol{w}_P and $\boldsymbol{\xi}_P$) is of special interest for us, which is why its two main steps will be ⁷² detailed.

Any positive measure $d\mu(\xi)$ is associated with a sequence of monic polynomials (*i.e.* polynomial whose leading coefficient equals 1) denoted π_k –with k the order of the polynomial– such that:

$$\int_{\Omega_{\xi}} \pi_i(\xi) \pi_j(\xi) d\mu(\xi) = 0, \quad \text{for } i \neq j$$
(3)

These polynomials are said orthogonal with respect to the measure $d\mu(\xi)$ and are defined to by:

$$\pi_{k}(\xi) = \frac{1}{c_{k}} \begin{vmatrix} m_{0} & m_{1} & \cdots & m_{k-1} & m_{k} \\ m_{1} & m_{2} & \cdots & m_{k} & m_{k+1} \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ m_{k-1} & m_{k} & \cdots & m_{2k-2} & m_{2k-1} \\ 1 & \xi & \cdots & \xi^{k-1} & \xi^{k} \end{vmatrix}$$
(4)

⁷⁸ with c_k a constant chosen so that the leading coefficient (of order k) of π_k equals 1, hence ⁷⁹ making π_k a monic polynomial.

⁸⁰ It is known that monic orthogonal polynomials satisfy a three-term recurrence relation ⁸¹ [24]:

$$\pi_{k+1}(\xi) = (\xi - a_k)\pi_k(\xi) - b_k\pi_{k-1}(\xi)$$
(5)

with a_k and b_k being the recurrence coefficients specific to the measure $d\mu(\xi)$, $\pi_{-1}(\xi) = 0$ and $\pi_0(\xi) = 1$.

Let $J_n(d\mu)$ be the $n \times n$ Jacobi matrix associated to the measure $d\mu$. This is a tridiagonal symmetric matrix defined as:

$$\boldsymbol{J}_{n}(\mathrm{d}\boldsymbol{\mu}) = \begin{pmatrix} a_{0} & \sqrt{b_{1}} & 0 \\ \sqrt{b_{1}} & a_{1} & \ddots & \\ & \ddots & \ddots & \\ 0 & & \sqrt{b_{n-1}} & a_{n-1} \end{pmatrix}$$
(6)

The weights and nodes of the quadrature rule from Eq. (2) are given by spectral properties of $J_P(d\mu)$. The nodes ξ_P of the rule are the eigenvalues of $J_P(d\mu)$. The weights of the rule are given by:

$$w_i = m_0 v_{1,i}^2 \tag{7}$$

where $v_{1,i}$ is the first component of the normalised eigenvector belonging to the eigenvalue ξ_i . The computation of the quadrature rule (Eq. (2)) then relies on two steps:

⁹¹ 1. The computation of the recurrence coefficients $\boldsymbol{a}_{P-1} = [a_0, \ldots, a_{P-1}]^\mathsf{T}$ and $\boldsymbol{b}_{P-1} = \begin{bmatrix} b_1, \ldots, b_{P-1} \end{bmatrix}^\mathsf{T}$.

⁹³ 2. The computation of the eigenvalues and the normalised eigenvectors of $J_P(d\mu)$.

⁹⁴ Multiple algorithms are available in the literature to compute the recurrence coefficients:

- The Quotient-Difference algorithm [25, 26]
- The Product-Difference algorithm [27]
- The Chebyshev algorithm [28]

The Chebyshev algorithm was found to be the stablest one of the three [1, 28], its description
is given in Appendix A.

100 2.3. Extended Quadrature Method of Moments

The QMOM method is well suited for the approximation of integral properties of the NDF, which is actually the main purpose of Gauss quadratures. However, in many applications such as evaporation [12] or dissolution [29] processes, point-wise values of the NDF $n(\xi)$ are required but not directly accessible from the moments. For that purpose, a method is needed to produce an approximation $\tilde{n}(\xi)$ of the original distribution $n(\xi)$, by knowing only a finite set of its moments.

In a sense, one can consider that the Gaussian quadrature computed with QMOM approximates $n(\xi)$ as a weighted sum of Dirac distributions:

$$\widetilde{n}(\xi) = \sum_{i=1}^{P} w_i \delta(\xi, \xi_i) \tag{8}$$

with the Dirac δ distribution defined by its sifting property

$$\int_{-\infty}^{+\infty} f(\xi) \,\delta(\xi,\xi_m) \mathrm{d}\xi = f(\xi_m) \tag{9}$$

For most applications, $n(\xi)$ is expected to be a continuous distribution whilst QMOM yields monodisperse or discrete polydisperse reconstructions of $n(\xi)$, with $\tilde{n}(\xi) = 0$ for all values of ξ except some finite number of these values.

Many methods were suggested to tackle this problem and to propose a continuous recon-113 struction $\widetilde{n}(\xi)$ from a finite number of moments \boldsymbol{m}_N . Some of them are the Spline method 114 [13], the Maximum-Entropy approach [14, 15, 12] or the Kernel Density Element Method 115 [16]. Their properties will not be discussed here but one only underlines that they tend to 116 be unstable, ill-conditionned, or have a high sensitivity to numerical parameters [13, 29, 30]. 117 In particular, none of them can handle the case of a weakly realisable moment set. Such 118 a moment set is associated to a discrete (or degenerated) distribution and, in this specific 119 case, the distribution provided by QMOM is the only possible reconstruction (see Eq. 8). 120

¹²¹ Note that a failure –or instabilities– in a numerical method can compromise the integrity ¹²² of large-scale simulations. For this reason, Chalons et al. [17], Yuan et al. [18] and Nguyen ¹²³ et al. [1] proposed a robust and stable method to tackle this reconstruction problem by han-¹²⁴ dling both continuous approximations and discrete solutions. Their approach, the Extended ¹²⁵ Quadrature Method of Moments, approximates $n(\xi)$ as a convex mixture of Kernel Density ¹²⁶ Functions (KDFs) of the same parametric family:

$$\widetilde{n}(\mu) = \sum_{i=1}^{P} w_i \delta_\sigma(\xi, \xi_i) \tag{10}$$

127 with

• w_i : the weight of the i-th node, $w_i \ge 0, \forall i \in \{1, \dots, P\}$

• ξ_i : the location parameter of the i-th node, $\xi_i \in \Omega_{\xi}, \forall i \in \{1, \dots, P\}$

• δ_{σ} : a KDF chosen to perform the approximation, referred later to as the reconstruction kernel. σ is the shape parameter of the approximation.

The computation of the weights $\boldsymbol{w}_P = [w_1, \ldots, w_P]^\mathsf{T}$, the nodes $\boldsymbol{\xi}_P = [\xi_1, \ldots, \xi_P]^\mathsf{T}$ and the shape parameter σ from the moment set \boldsymbol{m}_{2P} is performed by the EQMOM momentinversion procedure. The improvement of this procedure constitutes the core of this article and is detailed in section 3.

¹³⁶ Multiple standard normalized distribution functions can be used as the reconstruction ¹³⁷ kernel δ_{σ} (*e.g.* Gaussian, Log-normal, etc.). A list of them is given in Appendix B. All ¹³⁸ of these kernels degenerate into Dirac distribution if their shape parameters are sufficiently ¹³⁹ small:

$$\lim_{\sigma \to 0} \delta_{\sigma}(\xi, \xi_m) = \delta(\xi, \xi_m) \tag{11}$$

This allows EQMOM to be numerically stable in the case of a moment set m_{2P} being on the boundary of the realisable moment space $\partial \mathcal{M}_{2P}(\Omega_{\xi})$. Indeed, in such cases, the EQMOM approximation simply degenerates in a weighted sum of Dirac distribution and the definition given in Eq. 10 still holds true, with $\sigma = 0$. EQMOM can also be used to compute integral properties of the NDF with high accuracy. This comes with the introduction of nested quadratures. The main quadrature proposes the following approximation of integral terms:

$$\int_{\Omega_{\xi}} f(\xi) n(\xi) d\xi \approx \sum_{i=1}^{P} w_i \left[\int_{\Omega_{\xi}} f(\xi) \delta_{\sigma}(\xi, \xi_i) d\xi \right]$$
(12)

¹⁴⁷ Moreover, a quadrature rule can be used to approximate the bracketed integral in Eq. ¹⁴⁸ (12). This will be the nested quadrature that actually depends on the kernel $\delta_{\sigma}(\xi, \xi_m)$. For ¹⁴⁹ instance, Gauss-Hermite quadratures can be used to approximate integrals over a Gaussian ¹⁵⁰ kernel (see Appendix B.1). Nested quadratures then give the following approximation:

$$\int_{\Omega_{\xi}} f(\xi) n(\xi) d\xi \approx \sum_{i=1}^{P} w_i \sum_{j=1}^{Q} \omega_j f\left(g(\sigma, \xi_i, \lambda_j)\right)$$
(13)

with Q the order, $\boldsymbol{\omega}_Q = [\omega_1, \dots, \omega_Q]^\mathsf{T}$ the weights and $\boldsymbol{\lambda}_Q = [\lambda_1, \dots, \lambda_Q]^\mathsf{T}$ the nodes of the sub-quadrature. g defines the nodes of the nested quadrature from σ , ξ_i and λ_j . These nested quadratures are detailed for all KDFs in Appendix B and Appendix C.

¹⁵⁴ 3. Moment inversion procedure

The EQMOM moment-inversion procedure comes with analytical solutions for some kernels in the case of low-order quadratures. The one-node analytical solutions are detailed for all kernels in Appendix B. When they exist, the two-nodes analytical solutions are implemented in MATLAB code (see supplementary data) but are not detailed in this article. The current section is focusing on the numerical procedure used to compute the reconstruction parameters in absence of an analytical solution.

The procedure proposed by Yuan et al. [18] and Nguyen et al. [1] is first recalled in section 3.1. The section 3.2 details how their approach can be shifted toward a new convergence criteria that will be applied to the specific cases of

- the Hamburger moment problem (section 3.3): NDF defined on the whole phase space $\Omega_{\xi} =]-\infty, +\infty[$
- the *Stieltjes* moment problem (section 3.4): NDF defined on the positive phase space $\Omega_{\xi} =]0, +\infty[$
- the Hausdorff moment problem (section 3.5): NDF defined on the closed support 169 $\Omega_{\xi} =]0,1[$

Some moment sets lead to ill-conditioned situations that need to be specifically handled by EQMOM implementations. These are addressed in section 3.6.

172 3.1. Standard procedure

Let \boldsymbol{m}_N be the vector of the first N+1 integer moments of the measure $d\mu(\xi) = n(\xi)d\xi$, with N = 2P an even integer:

$$\boldsymbol{m}_{N} = \begin{bmatrix} m_{0} \\ m_{1} \\ \vdots \\ m_{N} \end{bmatrix}, \qquad m_{k} = \int_{\Omega_{\xi}} \xi^{k} n(\xi) \mathrm{d}\xi \qquad (14)$$

The EQMOM moment-inversion procedure aims to identify the parameters σ , $\boldsymbol{w}_P = \begin{bmatrix} w_1, \ldots, w_P \end{bmatrix}^{\mathsf{T}}$ and $\boldsymbol{\xi}_P = \begin{bmatrix} \xi_1, \ldots, \xi_P \end{bmatrix}^{\mathsf{T}}$ such that $\boldsymbol{m}_N = \widetilde{\boldsymbol{m}}_N$ with:

$$\widetilde{\boldsymbol{m}}_{N} = \begin{bmatrix} \widetilde{m}_{0} \\ \widetilde{m}_{1} \\ \vdots \\ \widetilde{m}_{N} \end{bmatrix}, \qquad \widetilde{m}_{k} = \int_{\Omega_{\xi}} \xi^{k} \widetilde{n}(\xi) \mathrm{d}\xi, \qquad \widetilde{n}(\xi) = \sum_{i=1}^{P} w_{i} \delta_{\sigma}(\xi, \xi_{i})$$
(15)

For any value of σ , Yuan et al. [18] identified a procedure which leads to the parameters \boldsymbol{w}_P and $\boldsymbol{\xi}_P$ such that $\boldsymbol{m}_{N-1} = \widetilde{\boldsymbol{m}}_{N-1}$. The EQMOM moment-inversion problem has then been reduced to solving a scalar non-linear equation by looking for a root of the function $D_N(\sigma) = m_N - \widetilde{m}_N(\sigma)$.

The approach developed by Yuan et al. [18] and then improved by Nguyen et al. [1] is based on the fact that, for the KDFs used in EQMOM, it is possible to write the following linear system:

$$\widetilde{\boldsymbol{m}}_n = \boldsymbol{A}_n(\sigma) \cdot \boldsymbol{m}_n^* \tag{16}$$

where $A_n(\sigma)$ is a lower-triangular $(n+1) \times (n+1)$ matrix whose elements depend only on the chosen KDF and on the value σ , whereas m_n^* is defined as:

$$\boldsymbol{m}_{n}^{*} = \begin{bmatrix} m_{0}^{*} \\ m_{1}^{*} \\ \vdots \\ m_{n}^{*} \end{bmatrix}, \qquad m_{k}^{*} = \sum_{i=1}^{P} w_{i} \xi_{i}^{k}$$
(17)

By their definition, the moments m_n^* correspond to the moments of a degenerated distribution (*i.e.* a finite sum of Dirac distributions), hence these moments will be referred as the *degenerated moments of the approximation*. Degenerated moments are defined in such a way that the vectors w_P and ξ_P can be computed from m_{2P-1}^* using a Gauss Quadrature (see 2.2).

At this point, one has the basis required to compute the objective function $D_N(\sigma)$ and to search for its root. The computation of $D_N(\sigma)$ from a vector \boldsymbol{m}_N is as follow (see also Fig. 1a):

- 194 1. Compute $\boldsymbol{m}_{N-1}^*(\sigma) = \boldsymbol{A}_{N-1}^{-1}(\sigma) \cdot \boldsymbol{m}_{N-1}$.
- ¹⁹⁵ 2. Compute the recurrence coefficients $\boldsymbol{a}_{P-1}^*(\sigma)$ and $\boldsymbol{b}_{P-1}^*(\sigma)$ by applying the Chebyshev ¹⁹⁶ algorithm to $\boldsymbol{m}_{N-1}^*(\sigma)$.
- ¹⁹⁷ 3. Use the recurrence coefficients to compute the Gaussian quadrature rule $\boldsymbol{w}_P(\sigma)$ and ¹⁹⁸ $\boldsymbol{\xi}_P(\sigma)$.

4. Knowing the parameters σ , $\boldsymbol{w}_P(\sigma)$ and $\boldsymbol{\xi}_P(\sigma)$ of the reconstruction, compute $\widetilde{m}_N(\sigma)$, this can be done easily by:

201

• Computing the N-th order degenerated moment of the approximated NDF: $\overline{m}_N^*(\sigma) = \sum_{i=1}^P w_i(\sigma)\xi_i(\sigma)^N$.

202

203 204 • Multiplying the last line of $\mathbf{A}_N(\sigma)$ and the vector of degenerated moments: $\widetilde{m}_N(\sigma) = [0, 0, \dots, 1] \cdot \mathbf{A}_N(\sigma) \cdot [m_0^*(\sigma), \dots, m_{N-1}^*(\sigma), \overline{m}_N^*(\sigma)]^{\mathsf{T}}$.

5. Compute
$$D_N(\sigma) = m_N - \widetilde{m}_N(\sigma)$$

For each compatible KDF, it is possible to use the low order moments to compute an upper bound σ_{max} so that the search of a root of D_N is restricted to the interval $\sigma \in [0, \sigma_{max}]$. Then a bounded non-linear equation solver such as Ridder's method can be applied to actually find the root of the function.

Two specific cases were discarded in the previous description of the method. First, it happens that the function D_N does not admit any root, in such a case the procedure is switched toward the minimisation of this function in order to reduce the error on the last moment of the approximation.

Second, during the computation of $D_N(\sigma)$, one must compute degenerated moments 214 from which the weights and nodes are extracted. If the degenerated moments $m_{N-1}^*(\sigma)$ turn 215 out not to be realisable on the support Ω_{ξ} of the NDF, the quadrature performed on this 216 vector will lead to nodes outside Ω_{ξ} , or even to negative/complex weights. Nguyen et al. [1] 217 then suggest to check for the realisability of the degenerated moments, and if these are not 218 realisable, to set $\widetilde{m}_N(\sigma)$ to a arbitrarily high value such as 10^{100} . This will force the non-219 linear equation solver to test a lower value of σ in order to bring back the vector $\boldsymbol{m}_{N-1}^{*}(\sigma)$ 220 within the realisable moment space. However note that this is only a numerical trick to 221 converge toward the actual root, but $D_N(\sigma)$ is actually undefined as soon as $\boldsymbol{m}_{N-1}^*(\sigma)$ is not 222 realisable. 223

224 3.2. A new procedure based on moment realisability

The reversible linear system linking the raw moments of the approximation \widetilde{m}_N to its degenerated moments m_N^* is such that a new objective function $D_N^*(\sigma)$ –whose root is the same as that of $D_N(\sigma)$ – can be formulated. Its computation is as follow (see also Fig. 1b):

1. Compute
$$\boldsymbol{m}_N^*(\sigma) = \boldsymbol{A}_N^{-1}(\sigma) \cdot \boldsymbol{m}_N$$

229 2. Compute a quadrature on the vector $\boldsymbol{m}_{N-1}^*(\sigma)$ to obtain the vectors $\boldsymbol{w}_P(\sigma)$ and $\boldsymbol{\xi}_P(\sigma)$.

230 3. Compute
$$\overline{m}_N^*(\sigma) = \sum_{i=1}^P w_i(\sigma)\xi_i(\sigma)^N$$
.

231 4. Compute
$$D_N^*(\sigma) = m_N^*(\sigma) - \overline{m}_N^*(\sigma)$$
.

Note that $D_N(\sigma) = D_N^*(\sigma) \times A_{N,N}(\sigma)$. As shown in Appendix B for all kernels, diagonal elements of $A_n(\sigma)$ are always strictly positive so the two objective functions do share the same roots.

The benefit of this new objective function is that it only requires the matrix $A_N^{-1}(\sigma)$ instead of both the matrix $A_{N-1}^{-1}(\sigma)$ and the last line of $A_N(\sigma)$. This only increases the clarity of the method, but has hardly no effect on its numerical cost. The point of this alternative approach is however to underline a crucial element for the new EQMOM implementation: we actually look for a value of σ for which $m_{2P}^*(\sigma) = \overline{m}_{2P}^*(\sigma)$. This implies that, for this specific searched σ value, the vector $\boldsymbol{m}_{2P}^*(\sigma)$ reads

$$\boldsymbol{m}_{2P}^{*}(\sigma) = \begin{bmatrix} \sum_{i=1}^{P} w_i \xi_i^{0} \\ \sum_{i=1}^{P} w_i \xi_i^{1} \\ \vdots \\ \sum_{i=1}^{P} w_i \xi_i^{2P} \end{bmatrix}$$
(18)

which is, by construction, the vector of the first 2P + 1 moments of the sum of P Dirac distributions.

Under the condition that a *P*-node EQMOM reconstruction exists for the moment set m_{2P} with $\sigma > 0$, $w_i > 0$, $\xi_i \neq 0, i \in \{1, \ldots, P\}$, the vector $m_{2P}^*(\sigma)$ will have the following specific properties:

1. The vector $\boldsymbol{m}_{2P-1}^*(\sigma)$ must be strictly within the realisable moment space $\mathcal{M}_{N-1}(\Omega_{\xi})$. 247 2. The vector $\boldsymbol{m}_{2P}^*(\sigma)$ must be on the boundary of the realisable moment space $\mathcal{M}_N(\Omega_{\xi})$.

EQMOM procedure will then rely on the realisability of the vector $m_{2P}^*(\sigma)$ instead of the computation of the error on the last moment, this will be a cheaper approach.

Situations were the EQMOM reconstruction exist but with $\sigma = 0$, or $\exists i \in \{1, \dots, P\}, w_i = 0$ or $\xi_i = 0$ are tackled in section 3.6 but are always based on checking the realisability of $m_{2P}^*(\sigma)$.

The actual definition of the realisable moment space of order n, \mathcal{M}_n , depends on the support Ω_{ξ} of the NDF. The three classical supports, corresponding to the *Hamburger*, *Stieltjes* and *Hausdorff* moment problems, come with different constraints on a moment set to ensure its realisability. The realisability criteria for each of these supports will then be detailed.

Fig. 1 sums up the "standard approach" based on $D_N(\sigma)$, the shifted approach, based on $D_N^*(\sigma)$, as well as the new approach based on the realisability criteria of $m_{2P}^*(\sigma)$ for all three supports.

²⁶¹ 3.3. Application to the Hamburger problem

As stated in 2.2, it is known that the monic polynomials which are orthogonal to a measure $d\mu(\xi) = n(\xi)d\xi$ satisfy a three-term recurrence relation (Eq. (5)) with a_k and $b_k, k \in \mathbb{N}$, the recurrence coefficients specific to the measure $d\mu(\xi)$. The Favard's theorem [31] and its converse [32] imply that the measure $d\mu(\xi)$ is realisable on $\Omega_{\xi} =]-\infty, +\infty[$ if and only if $a_k \in \mathbb{R}$ and $b_k > 0, \forall k \in \mathbb{N}$.

One looks for a value of σ such that the associated degenerated moments $\boldsymbol{m}_{2P-1}^*(\sigma)$ are strictly realisable (*i.e.* within the moment space), and the moments $\boldsymbol{m}_{2P}^*(\sigma)$ are weakly realisable (*i.e.* on the frontier of realisability). Then, if the Chebyshev algorithm is used to compute the recurrence coefficients $\boldsymbol{a}_{P-1}^*(\sigma) = [\boldsymbol{a}_0^*(\sigma), \ldots, \boldsymbol{a}_{P-1}^*(\sigma)]^{\mathsf{T}}$ and $\boldsymbol{b}_P^*(\sigma) = [\boldsymbol{b}_1^*(\sigma), \ldots, \boldsymbol{b}_P^*(\sigma)]^{\mathsf{T}}$ from the vector $\boldsymbol{m}_{2P}^*(\sigma)$, the condition of realisability can be written in terms of values of $\boldsymbol{b}_P^*(\sigma)$: looking for the EQMOM reconstruction parameters with the Gaussian and Laplace

²⁷³ kernels is equivalent to looking for a value of σ such as:

Figure 1: Comparison of the computation of convergence criteria based on (a) $D_N(\sigma)$, (b) $D_N^*(\sigma)$ and (c) the realisability criteria of the support Ω_{ξ} . CA: Chebyshev Algorithm. QC: Quadrature Computation. The convergence criteria are highlighted in light blue. Inspired by Fig. 1 from Nguyen et al. [1].

•
$$b_k^*(\sigma) > 0, \quad \forall k \in \{1, \dots, P-1\}$$

•
$$b_P^*(\sigma) = 0$$

Fig. 2 makes use of the developments from Appendix B.1 and Appendix B.2, about the Gaussian and Laplace kernels respectively, to show the evolution of $D_6(\sigma)$, $D_6^*(\sigma)$ and $b_k^*(\sigma)$, $k \in \{1, 2, 3\}$ for two sets of 7 moments (P = 3). This figure illustrates the fact that indeed the approaches based on $D_N(\sigma)$, $D_N^*(\sigma)$ and $b_P^*(\sigma)$ are equivalent as they share the same circled root.

Let denote σ_k the root of $b_k(\sigma)$. One can notice that the root σ_k lies within the interval [0, σ_{k-1}]. We actually observed the existence of all roots σ_k , $k \in \{1, \ldots, P\}$ on numerous (about 10⁶) randomly selected moment sets of N + 1 = 13 moments, and never observed an undefined root. The generality of this observation has not been mathematically proved, but it seems that indeed σ_k is always defined and always lies in $\sigma_k \in [0, \sigma_{k-1}], k \in \{2, \ldots, P\}$. σ_1 is defined analytically.

The previous observations were used to design a simple algorithm which allows identifying the root σ_P . This algorithm is based on the fact that it is possible to check whether a value σ_t is higher or lower than σ_P at low cost and with no prior knowledge of σ_P value:

- If $b_k^*(\sigma_t) > 0$, $\forall k \in \{1, \ldots, P\}$, then $\sigma_t < \sigma_P$.
- Otherwise, that is if $\exists k \in \{1, \ldots, P\}$, $b_k^*(\sigma_t) < 0$, then $\sigma_t > \sigma_P$.
- ²⁹² One can then use an iterative approach that will



Figure 2: Evolution of the different convergence criteria for both Gaussian (a and b) and Laplace (c and d) kernels depending on σ value. The two initial moment sets are $\boldsymbol{m}_6^{(1)} = [1 \ 1 \ 2 \ 5 \ 12 \ 42 \ 133]^{\mathsf{T}}$ and $\boldsymbol{m}_6^{(2)} = [1 \ 2 \ 7 \ 17 \ 58 \ 149 \ 493]^{\mathsf{T}}$.

- 1. Check the realisability of the raw moments $\boldsymbol{m}_{2P} = \boldsymbol{m}_{2P}^*(0)$ by computing $\boldsymbol{b}_P^*(0)$ and checking the positivity of all elements.
- 295 2. Initialise an interval $\left[\sigma_l^{(0)}, \sigma_r^{(0)}\right]$ such that $\sigma_l^{(0)} < \sigma_P$ and $\sigma_r^{(0)} > \sigma_P$, and then update 296 these bounds to shrink the search interval. These initial values will be $\sigma_l^{(0)} = 0$ and 297 $\sigma_r^{(0)} = \sigma_1$ with σ_1 the analytical solution of $b_1^*(\sigma) = 0$.
- $_{298}$ 3. Iterate over k

29

302

(a) Choose
$$\sigma_t \in \left[\sigma_l^{(k-1)}, \sigma_r^{(k-1)}\right]$$

300 (b) Compute $\boldsymbol{b}_P^*(\boldsymbol{\sigma}_t)$.

(c) If all elements of
$$\boldsymbol{b}_P^*(\sigma_t)$$
 are positive, set $\sigma_l^{(k)} = \sigma_t$ and $\sigma_r^{(k)} = \sigma_r^{(k-1)}$.

(d) Otherwise, set
$$\sigma_l^{(k)} = \sigma_l^{(k-1)}$$
 and $\sigma_r^{(k)} = \sigma_t$.

The choice of σ_t at step 3a will be made by trying to locate the root σ_j of $b_j^*(\sigma)$ with j the index of the first negative element of $\boldsymbol{b}_P^*(\sigma_r^{(k)})$. Following Nguyen et al. [1] developments, the use of Ridder's method is advised to select σ_t . This method actually tests two σ values per iteration. Consequently, the step 3 of the previous algorithm becomes:

307 3. Iterate over k

(a) Identify
$$j$$
 the index of the first negative element of $\boldsymbol{b}_P^*\left(\sigma_r^{(k-1)}\right)$
(b) Compute $\sigma_{t_1} = \frac{1}{2}\left(\sigma_l^{(k-1)} + \sigma_r^{(k-1)}\right)$ and $\boldsymbol{b}_P^*(\sigma_{t_1})$.

(c) Compute
$$\sigma_{t_2} = \sigma_{t_1} + \left(\sigma_{t_1} - \sigma_l^{(k-1)}\right) \frac{b_j^*(\sigma_{t_1})}{\sqrt{b_j^*(\sigma_{t_1})^2 - b_j^*(\sigma_l^{(k-1)}) * b_j^*(\sigma_r^{(k-1)})}}$$
 and $\boldsymbol{b}_P^*(\sigma_{t_2})$.

- (d) Set $\sigma_l^{(k)}$ as the highest value between $\sigma_l^{(k-1)}$, σ_{t_1} and σ_{t_2} such that the corresponding vector \boldsymbol{b}_P^* contains only positive values.
- 313 314

(e) Set $\sigma_r^{(k)}$ as the lowest value between $\sigma_r^{(k-1)}$, σ_{t_1} and σ_{t_2} such that the corresponding vector \boldsymbol{b}_P^* contains at least one negative value.

Stop the computation if $\sigma_r^{(k)} - \sigma_l^{(k)} < \varepsilon \sigma_1$ or if $b_P^* \left(\sigma_l^{(k)} \right) < \varepsilon b_P^*(0)$, with ε a relative tolerance (e.g. $\varepsilon = 10^{-10}$). Then compute the weights \boldsymbol{w}_P and nodes $\boldsymbol{\xi}_P$ of the EQMOM reconstruction by computing a Gauss quadrature based on the recurrence coefficients $\boldsymbol{a}_{P-1}^* \left(\sigma_l^{(k)} \right)$ and $\boldsymbol{b}_{P-1}^* \left(\sigma_l^{(k)} \right)$.

Actual implementations of this algorithm for both kernels are provided as supplementary data.

321 3.4. Application to the Stieltjes problem

It is well known that the realisability of a moment set \mathbf{m}_N on the support $\Omega_{\xi} =]0, +\infty[$ is strictly equivalent to the positivity of the Hankel determinants $\underline{\mathcal{H}}_{2n+d}$ [33] defined as:

$$\underline{\mathcal{H}}_{2n+d} = \begin{vmatrix} m_d & \cdots & m_{n+d} \\ \vdots & \ddots & \vdots \\ m_{n+d} & \cdots & m_{2n+d} \end{vmatrix}$$
(19)

with $d \in \{0, 1\}$ and $n \in \mathbb{N}$, $2n + d \leq N$.

This condition on the positivity of Hankel determinants can be translated into a condition on the positivity of the numbers ζ_k [32] defined by :

$$\zeta_k = \frac{\underline{\mathcal{H}}_{k-3}\underline{\mathcal{H}}_k}{\underline{\mathcal{H}}_{k-2}\underline{\mathcal{H}}_{k-1}}, \qquad \underline{\mathcal{H}}_j = 1 \text{ if } j < 0$$
(20)

These numbers can be directly computed from the recurrence coefficients a_P and b_P defined in 2.2 through the following relations:

$$\zeta_{2k} = \frac{b_k}{\zeta_{2k-1}}, \qquad \zeta_{2k+1} = a_k - \zeta_{2k} \tag{21}$$

329 with $\zeta_1 = a_0 = m_1/m_0$.

The goal here is to use these realisability criteria to compute the parameters of EQMOM quadrature with either the Log-normal, the Gamma or the Weibull kernel (see Appendix B.3, Appendix B.4 and Appendix B.5 respectively). In these cases, one must

- 1. Compute $\boldsymbol{m}_N^*(\sigma) = \boldsymbol{A}_N^{-1}(\sigma) \cdot \boldsymbol{m}_N$ with $\boldsymbol{A}_N(\sigma)$ the matrix associated to the chosen kernel (see Appendix B.3, Appendix B.4, Appendix B.5).
- 2. Apply the Chebyshev algorithm to $\boldsymbol{m}_N^*(\sigma)$ to access the recurrence coefficients $\boldsymbol{a}_P^*(\sigma)$ and $\boldsymbol{b}_P^*(\sigma)$.

337 3. Compute
$$\boldsymbol{\zeta}_N^*(\sigma) = [\boldsymbol{\zeta}_1^*(\sigma), \dots, \boldsymbol{\zeta}_N^*(\sigma)]^\mathsf{T}$$
 using the relations in Eq. (21).

338 One actually looks for σ such that



Figure 3: Evolution of the different convergence criteria for the Weibull kernel depending on σ value. The initial moment sets are $\boldsymbol{m}_{6}^{(a)} = [1\ 1.5\ 12\ 131\ 15200\ 18033\ 2.16e5]^{\mathsf{T}}$, $\boldsymbol{m}_{6}^{(b)} = [1\ 5.5\ 78\ 1285\ 22225\ 4.05e5\ 7.88e6]^{\mathsf{T}}$ and $\boldsymbol{m}_{6}^{(c)} = [1\ 1\ 2\ 5\ 14\ 42\ 133]^{\mathsf{T}}$.

339 • $\zeta_k^*(\sigma) > 0, \quad \forall k \in \{1, \dots, N-1\}$

• $\zeta_N^*(\sigma) = 0$

Let σ_k be the root of $\zeta_k^*(\sigma)$. In all cases, the root σ_2 is defined, analytically for the Log-normal and Gamma kernels, and numerically for the Weibull kernel. Fig. 3 shows the evolution of $D_6(\sigma)$, $D_6^*(\sigma)$ and $\zeta_6^*(\sigma)$ for three moment sets when the developments relative to the Weibull (see Appendix B.5) kernel are used. Three situations can be observed on that figure:

1. All roots σ_k , $k \in \{2, \ldots, N\}$ are defined (Fig. 3a).

2. Some intermediary roots σ_k , $k \in \{3, \ldots, N-1\}$, are not defined but the root σ_N still exists (Fig. 3b).

349 3. The root σ_N is not defined (Fig. 3c).

³⁵⁰ These three cases can be observed for the Gamma and Log-normal kernels too.

In the first two cases, when σ_N exists, the EQMOM approximation is well defined. The last case –where $\zeta_N^*(\sigma)$ admits no root in $[0, \sigma_{N-1}]$ – actually corresponds to the case described by Nguyen et al. [1] where $D_N(\sigma)$ did not admit any root either. In this case, it was suggested to minimise $D_N(\sigma)$ in order to reduce the difference between m_N and $\tilde{m}_N(\sigma)$ as much as possible.

 $D_N(\sigma)$ tends to be a decreasing function, but is undefined as soon as any element of $\zeta_{N-1}^*(\sigma)$ is negative. The minimum of $D_N(\sigma)$ is then usually located at the highest order defined root. For instance, in the case shown in Fig. 3c, the minimum of $D_6(\sigma)$ is located at the root σ_5 of $\zeta_5^*(\sigma)$.

The moment-inversion procedure for reconstruction kernels defined on $\Omega_{\xi} =]0, +\infty[$ is then reduced to the identification of the defined root $\sigma_k, \ k \in \{2, \ldots, N\}$, of highest index. The algorithm proposed in section 3.3 already converges toward this root and only requires little adjustments:

1. Check the realisability of the raw moments $\boldsymbol{m}_{2P} = \boldsymbol{m}_{2P}^*(0)$ by computing $\boldsymbol{\zeta}_N^*(0)$ and checking the positivity of all elements.

26. Initialise an interval $\left[\sigma_l^{(0)}, \sigma_r^{(0)}\right]$ with $\sigma_l^{(0)} = 0$ and $\sigma_r^{(0)} = \sigma_2$ with σ_2 the solution of $\zeta_2^*(\sigma) = 0$.

368 3. Iterate over k

370

37

(a) Identify j the index of the first negative element of $\zeta_N^*\left(\sigma_r^{(k-1)}\right)$.

(b) Compute
$$\sigma_{t_1} = \frac{1}{2} \left(\sigma_l^{(k-1)} + \sigma_r^{(k-1)} \right)$$
 and $\boldsymbol{\zeta}_N^*(\sigma_{t_1})$.

(c) Compute
$$\sigma_{t_2} = \sigma_{t_1} + \left(\sigma_{t_1} - \sigma_l^{(k-1)}\right) \frac{\zeta_j^*(\sigma_{t_1})}{\sqrt{\zeta_j^*(\sigma_{t_1})^2 - \zeta_j^*(\sigma_l^{(k-1)}) * \zeta_j^*(\sigma_r^{(k-1)})}}$$
 and $\boldsymbol{\zeta}_N^*(\sigma_{t_2})$.

(d) Set $\sigma_l^{(k)}$ as the highest value between $\sigma_l^{(k-1)}$, σ_{t_1} and σ_{t_2} such that the corresponding vector $\boldsymbol{\zeta}_N^*$ contains only positive values.

(e) Set $\sigma_r^{(k)}$ as the lowest value between $\sigma_r^{(k-1)}$, σ_{t_1} and σ_{t_2} such that the corresponding vector $\boldsymbol{\zeta}_N^*$ contains at least one negative value.

Stop the computation if $\sigma_r^{(k)} - \sigma_l^{(k)} < \varepsilon \sigma_1$ or if $\zeta_N^* \left(\sigma_l^{(k)} \right) < \varepsilon \zeta_N^*(0)$, with ε a relative tolerance (*e.g.* $\varepsilon = 10^{-10}$). Then compute the weights \boldsymbol{w}_P and nodes $\boldsymbol{\xi}_P$ of the EQMOM reconstruction by computing a Gaussian-quadrature based on the recurrence coefficients $\boldsymbol{a}_{P-1}^* \left(\sigma_l^{(k)} \right)$ and $\boldsymbol{b}_{P-1}^* \left(\sigma_l^{(k)} \right)$.

380 3.5. Application to the Hausdorff problem

The moments of a distribution defined on the closed support $\Omega_{\xi} =]0, 1[$ must obey two sets of conditions in order to be within the realisable moment space [15, 26]. The moment set \boldsymbol{m}_N is interior to the realisable moment space associated to the support $\Omega_{\xi} =]0, 1[$ if and only if:

385 • $\underline{\mathcal{H}}_k > 0, \qquad \forall k \in \{0, \dots, N\}$

 $\bullet \ \overline{\mathcal{H}}_k > 0, \qquad \forall k \in \{1, \dots, N\}$

³⁸⁷ with $\underline{\mathcal{H}}_k$ defined in Eq. (19) and $\overline{\mathcal{H}}_k$ defined by

$$\overline{\mathcal{H}}_{2n+d} = \begin{vmatrix} m_{d-1} - m_d & \cdots & m_{n+d-1} - m_{n+d} \\ \vdots & \ddots & \vdots \\ m_{n+d-1} - m_{n+d} & \cdots & m_{2n+d-1} - m_{2n+d} \end{vmatrix}$$
(22)

Leaving aside the obvious condition $\underline{\mathcal{H}}_0 = m_0 > 0$, the conditions $\overline{\mathcal{H}}_k > 0$ and $\underline{\mathcal{H}}_k > 0$ induce a lower bound m_k^- and an upper bound m_k^+ for the values of m_k , $k \in \{1, \ldots, N\}$. Consequently, one can define the canonical moments of the distribution $\boldsymbol{p}_N = [p_1, \ldots, p_N]^{\mathsf{T}}$ as

$$p_k = \frac{m_k - m_k^-}{m_k^+ - m_k^-} \tag{23}$$

³⁹² A moment set \boldsymbol{m}_N is realisable if and only if the associated canonical moment set \boldsymbol{p}_N lies ³⁹³ in the hypercube $]0,1[^N$. The canonical moments can be computed through the recurrence ³⁹⁴ relation [34]:

$$p_k = \frac{\zeta_k}{1 - p_{k-1}} \tag{24}$$

with ζ_k defined in Eq. (20) and $p_1 = m_1$.

In the case of the Beta kernel (see Appendix B.6), one is looking for a value of σ such that the vector $\mathbf{p}_N^*(\sigma)$ has the following properties:

398 •
$$p_k^*(\sigma) \in [0, 1[, \forall k \in \{1, \dots, N-1\}]$$

$$\bullet \ p_N^*(\sigma) = 0$$

 $\boldsymbol{p}_{N}^{*}(\sigma) \text{ is computed from the vector } \boldsymbol{\zeta}_{N}^{*}(\sigma) \text{ which is deduced from the recurrence coef-}$ $\boldsymbol{p}_{N}^{*}(\sigma) \text{ and } \boldsymbol{b}_{P}^{*}(\sigma). \text{ These are computed -like previously- through the Chebyshev }$ $\boldsymbol{p}_{N}^{*}(\sigma) = \boldsymbol{h}_{N}^{-1}(\sigma) \cdot \boldsymbol{m}_{N}.$

Fig. 4 shows the evolution of the canonical moments and the convergence criteria $D_6(\sigma)$ and $D_6^*(\sigma)$ for four different sets of 7 moments with the developments relative to the Beta kernel (see Appendix B.6). Each of these sets corresponds to one of the four situations encountered when dealing with Beta EQMOM:

- Fig. 4a: the root σ_N of $D_N(\sigma)$, $D_N^*(\sigma)$ and $p_N^*(\sigma)$ exists and can be identified through a similar procedure than that described in sections 3.3 and 3.4.
- Fig. 4b: the root σ_N is not defined but the minimum of $D_N(\sigma)$ is located at the σ value for which $\boldsymbol{p}_{N-1}^*(\sigma)$ is on the boundary of the hypercube $]0,1[^{N-1}]$.
- Fig. 4c: $D_N(\sigma)$, $D_N^*(\sigma)$ and $p_N^*(\sigma)$ admit multiple roots.

• Fig. 4d: the root σ_N is defined, but there is a range $]\sigma_{v_1}, \sigma_{v_2}[$ with $\sigma_{v_2} < \sigma_N$, highlighted in light grey, such that in this interval the convergence criteria are undefined because $\forall \sigma \in]\sigma_{v_1}, \sigma_{v_2}[, \mathbf{p}_{N-1}^*(\sigma) \notin]0, 1[^{N-1}]$.

The algorithm proposed in sections 3.3 and 3.4 can still be applied here by replacing the convergence criteria by the canonical moments, and by checking that the values of $\boldsymbol{p}_N^*(\sigma)$ all lie in the interval]0, 1[instead of checking only for positivity:

⁴¹⁸ 1. Check the realisability of the raw moments $\boldsymbol{m}_{2P} = \boldsymbol{m}_{2P}^{*}(0)$ by computing $\boldsymbol{p}_{N}^{*}(0)$ and ⁴¹⁹ checking that all elements lie in]0, 1[.

420 2. Initialise an interval $\left[\sigma_l^{(0)}, \sigma_r^{(0)}\right]$ with $\sigma_l^{(0)} = 0$ and $\sigma_r^{(0)} = \sigma_2$ with σ_2 the analytical 421 solution of $p_2^*(\sigma) = 0$.

422 3. Iterate over k



Figure 4: Evolution of the different convergence criteria for the Beta reconstruction kernel and four initial moment sets. These sets can be found in the figure source code provided as supplementary data.

(a) Identify
$$j$$
 the index of the first element of $\boldsymbol{p}_{N}^{*}\left(\sigma_{r}^{(k-1)}\right)$ that is either negative or
higher than 1.
(b) Compute $\sigma_{t_{1}} = \frac{1}{2}\left(\sigma_{l}^{(k-1)} + \sigma_{r}^{(k-1)}\right)$ and $\boldsymbol{p}_{N}^{*}(\sigma_{t_{1}})$.
(c) If $j < N$ and $p_{j}^{*}\left(\sigma_{r}^{(k-1)}\right) > 1$
(c) Compute $\sigma_{t_{2}} = \sigma_{t_{1}} + \left(\sigma_{t_{1}} - \sigma_{l}^{(k-1)}\right) \frac{q_{j}^{*}(\sigma_{t_{1}})^{2}}{\sqrt{q_{j}^{*}(\sigma_{t_{1}})^{2} - q_{j}^{*}(\sigma_{l}^{(k-1)})}}$ and $\boldsymbol{p}_{N}^{*}(\sigma_{t_{2}})$,
with $q_{j}^{*}(\sigma) = 1 - p_{j}^{*}(\sigma)$.
(d) Else, that is if $j = N$ or $p_{j}^{*}\left(\sigma_{r}^{(k-1)}\right) < 0$
(e) Compute $\sigma_{t_{2}} = \sigma_{t_{1}} + \left(\sigma_{t_{1}} - \sigma_{l}^{(k-1)}\right) \frac{p_{j}^{*}(\sigma_{t_{1}})^{2} - p_{j}^{*}(\sigma_{l}^{(k-1)})}{\sqrt{p_{j}^{*}(\sigma_{t_{1}})^{2} - p_{j}^{*}(\sigma_{l}^{(k-1)})}}$ and $\boldsymbol{p}_{N}^{*}(\sigma_{t_{2}})$.
(e) Set $\sigma_{l}^{(k)}$ as the highest value between $\sigma_{l}^{(k-1)}$, $\sigma_{t_{1}}$ and $\sigma_{t_{2}}$ such that the correspond-
ing vector \boldsymbol{p}_{N}^{*} lies in $]0, 1[^{N}$.

⁴³⁵ Stop the computation if $\sigma_r^{(k)} - \sigma_l^{(k)} < \varepsilon \ \sigma_2$ or if $p_N^* \left(\sigma_l^{(k)} \right) < \varepsilon \ p_N^*(0)$, with ε a relative ⁴³⁶ tolerance (*e.g.* $\varepsilon = 10^{-10}$). As previously, once convergence is achieved, the weights \boldsymbol{w}_P and ⁴³⁷ nodes $\boldsymbol{\xi}_P$ of the reconstruction can be obtained by computing a Gaussian quadrature rule ⁴³⁸ based on the recurrence coefficients $\boldsymbol{a}_{P-1}^* \left(\sigma_l^{(k)} \right)$ and $\boldsymbol{b}_{P-1}^* \left(\sigma_l^{(k)} \right)$. This algorithm will converge to the root σ_N for cases similar to Fig. 4a; to the minimum of $D_N(\sigma)$ for cases similar to Fig. 4b; to one of the multiple roots for cases similar to Fig. 441 4c. In the case illustrated in Fig. 4d, the algorithm may or may not identify the existing 442 root, depending on whether one of the intermediate tested σ values lies in the greyed area.

One could try to develop a more robust algorithm, that will always find the root if it 443 is defined, even in the case shown in Fig. 4d. An other improvement would be to ensure 444 a consistent result when multiple roots exist, for instance by converging toward the lowest 445 root, so that a small perturbation in the raw moments will only cause a small change on the 446 resulting σ value. Nothing prevents the current algorithm from converging toward one root 447 for a moment set and toward another one after a small perturbation of this set which could 448 induce instabilities in large-scale simulations. Note that these limitations already existed in 449 previous EQMOM implementations and do not result from the new approach developed in 450 this article. 451

452 3.6. Handling weakly realisable and ill-conditioned moment sets

⁴⁵³ The EQMOM moment-inversion procedure attempts to identify a NDF defined by

$$\widetilde{n}(\xi) = \sum_{i=1}^{P} w_i \delta_{\sigma}(\xi, \xi_i)$$
(25)

454 whose first 2P + 1 integer moments are given by m_{2P} .

This approximation is not always possible as shown in sections 3.4 and 3.5. When the EQMOM approximation exists, it may be ill-conditioned if at least one of the followings holds true:

- 458 $\sigma = 0$
- $\exists i, w_i = 0$
- $\bullet \exists i, \ \xi_i = 0$

The first situation is that of m_{2P} being weakly realisable. The second situation occurs if m_{2P} is the moment set of a convex mixture of the reconstruction kernel with less than Pnodes. These situations are not mutually exclusive, a vector m_6 could be the vector of the 7 first moments of a bi-Dirac distribution, one of which could be located in $\xi = 0$.

Accounting for these situations requires introducing the order of realisability of a moment set, $\mathcal{N}(\boldsymbol{m}_N)$. This notation was introduced by Nguyen et al. [1] but was only defined on $\Omega_{\xi} =]0, +\infty[$ in terms of Hankel determinants. The following definition is broader as it encompasses theirs but extends it to other supports. $\mathcal{N}(\boldsymbol{m}_N)$ is the number of moments in the largest strictly realisable subset of \boldsymbol{m}_N . For each support, the order of realisability is defined in terms of the realisability criterion:

- For $\Omega_{\xi} =]-\infty, +\infty[$, compute \boldsymbol{b}_P from \boldsymbol{m}_{2P} ;
- if all elements are positive, $\mathcal{N}(\boldsymbol{m}_{2P}) = 2P + 1;$
- 473 else, if there is n such that $b_n = 0$, $\mathcal{N}(\boldsymbol{m}_{2P}) = 2n$;

474	- else, if there is n such that $b_n < 0$, $\mathcal{N}(\boldsymbol{m}_{2P}) = 2n - 1$.
475	• For $\Omega_{\xi} =]0, +\infty[$, compute $\boldsymbol{\zeta}_{2P}$ from $\boldsymbol{m}_{2P};$
476	- if all elements are positive, $\mathcal{N}(\boldsymbol{m}_{2P}) = 2P + 1;$
477	- else identify n such that $\zeta_n \leq 0$, $\mathcal{N}(\boldsymbol{m}_{2P}) = n$.
478	• For $\Omega_{\xi} =]0, 1[$, compute \boldsymbol{p}_{2P} from \boldsymbol{m}_{2P} ;
479	- if all elements are included on $]0, 1[, \mathcal{N}(\boldsymbol{m}_{2P}) = 2P + 1;$

480 - else identify n such that $p_n \notin]0, 1[, \mathcal{N}(\boldsymbol{m}_{2P}) = n.$

Detecting the situations where $\sigma = 0$ requires to check the order of realisability of raw moments. If $\mathcal{N}(\boldsymbol{m}_{2P})$ is even, set $\sigma = 0$; otherwise apply the iterative procedure to $\boldsymbol{m}_{2P'}$ with $\mathcal{N}(\boldsymbol{m}_{2P}) = 2P' - 1$ to identify σ [1].

The actual number of nodes required by the EQMOM approximation, *i.e.* the number of non-zero weights P'', is determined from $\mathcal{N}(\boldsymbol{m}_{2P}^*(\sigma))$. If it is even, $P'' = \mathcal{N}(\boldsymbol{m}_{2P}^*(\sigma))/2$; otherwise, $P'' = (\mathcal{N}(\boldsymbol{m}_{2P}^*(\sigma)) + 1)/2$ but one node will be located in $\xi = 0$ which might be an issue for KDFs defined on $\Omega_{\xi} =]0, +\infty[$ or $\Omega_{\xi} =]0, 1[$. The weights and nodes will be computed from the recurrence coefficients $\boldsymbol{a}_{P''-1}^*(\sigma)$ and $\boldsymbol{b}_{P''-1}^*(\sigma)$. If P'' < P, let $w_k = 0, \xi_k = 1/2, \ \forall k \in \{P'' + 1, \dots, P\}.$

These adjustments of the first and last steps of algorithms described in sections 3.3, 3.4 and 3.5 give great stability to the moment-inversion procedure at low cost.

In the situation where $\mathcal{N}(\boldsymbol{m}_{2P}^*(\sigma)) = 2P$, the EQMOM approximation is guaranteed to preserve the whole moment set \boldsymbol{m}_{2P} . However, if $\mathcal{N}(\boldsymbol{m}_{2P}^*(\sigma)) < 2P$, the approximation may, or may not, preserve all moments with no simple method to check for this. One should compute the moments of the EQMOM approximation and measure the relative error from original moments.

497 4. Comparison of EQMOM approaches

498 4.1. Method

The new EQMOM moment-inversion procedure only requires computation of the realisability criteria of the vector of degenerated moments $m_{2P}^*(\sigma)$ in order to identify σ . These computations were already performed in the original approach [1] to ensure the realisability of the vector $m_{2P-1}^*(\sigma)$ prior to the quadrature computation and ulterior steps.

It is therefore obvious that the new approach will always require a lower number of floating point operations (FLOP). In order to quantify this reduction on FLOP number, and the actual performance gain, different implementations of EQMOM are compared, they are based either on the realizability criteria, or on a quadrature-based objective function.

507 4.1.1. Tested EQMOM implementations

⁵⁰⁸ Comparison are performed for kernels defined on $\Omega_{\xi} =]-\infty, +\infty[$ (*i.e.* Gauss and Laplace ⁵⁰⁹ kernels), and on $\Omega_{\xi} =]0, +\infty[$ (*i.e.* Log-Normal, Gamma and Weibull kernels), using MAT-⁵¹⁰ LAB [22] implementations. Implementations that are based on the realizability criteria of $m_{2P}^*(\sigma)$ use algorithms that were fully described in sections 3.3 and 3.4 and adjustments from section 3.6.

For quadrature-based moment-inversion implementations, we optimized codes from Marchisio and Fox [20] and the OpenQBMM project [19] by implementing optimizations suggested by Nguyen et al. [1] and adjustments from section 3.6. Instead of searching for the root of $D_{2P}(\sigma)$ (see Fig. 1a), these implementations directly search the root of $D_{2P}^*(\sigma)$ (Fig. 1b). Doing so, all compared implementations only require the matrix $A_{2P}^{-1}(\sigma)$ and can benefit from the same code optimization when computing $m_{2P}^*(\sigma) = A_{2P}^{-1}(\sigma) \cdot m_{2P}$.

For kernels defined on $\Omega_{\xi} =]0, +\infty[$, if Ridder's method fails to identify a root of $D_{2P}^*(\sigma)$, the golden-ratio method is used to minimize $D_{2P}(\sigma)^2 = (D_{2P}^*(\sigma) \cdot A_{2P,2P}(\sigma))^2$. The goldenratio minimization method was already used in OpenQBMM [19].

522 4.1.2. Performance measurements

The main element of comparison is the number of floating-point operations required for the whole moment-inversion procedure. The MATLAB implementations embed a simple FLOP counter that distinguishes each operation $(+, -, *, /, \exp, \sqrt{\cdot}, \Gamma(\cdot), ...)$ and counts them for each step of the moment-inversion procedure (linear system, Chebyshev algorithm, quadrature computation and others).

In order to evaluate the number of operations used in the computation of the eigenvalues and eigenvectors of the Jacobi matrix (Eq. (6)), the Jacobi and the Francis algorithms which are suited for symmetric matrices [35] are used in place of the MATLAB built-in "eig" function [22]. The Jacobi algorithm is used for matrices of size up to 3×3 and the Francis algorithm for larger matrices in order to always use the fastest method.

Two others metrics are measured for each call to the moment-inversion procedure: the number of tested σ values and the wall-time of function calls.

535 4.1.3. Tested moment sets

Each comparison was performed on 10^4 randomly generated moment sets. These have varying size $2P + 1 \in \{5, 7, 9, 11\}$ and were either far from, or close to, the boundary of the realisable moment space.

⁵³⁹ Moments sets for kernels defined on $\Omega_{\xi} =]-\infty, +\infty[$ were computed from random vectors ⁵⁴⁰ \boldsymbol{a}_{P-1} and \boldsymbol{b}_P using a reversed Chebyshev algorithm. Distribution laws for the elements of ⁵⁴¹ these vectors are

- $a_k \sim \mathcal{N}(0, 25), \quad k \in \{0, \dots, P-1\}.$
- $b_k \sim 1 + \operatorname{Exp}(4), \quad k \in \{1, \dots, P\}.$
- $b_P \sim \text{Exp}(0.5)$ for moment sets close from the frontier of realisability.

Similarly, moments sets for kernels defined on $\Omega_{\xi} =]0, +\infty[$ were computed from random vectors ζ_{2P} using a reversed ζ -Chebyshev algorithm [1]. Elements of these vectors are generated using following distribution laws:

- $\zeta_k \sim 1 + \operatorname{Exp}(4), \quad k \in \{1, \dots, 2P\}.$
- $\zeta_{2P} \sim \text{Exp}(0.5)$ for moment sets close from the frontier of realisability.

550 4.1.4. Reproducibility

To allow reproducibility of results described hereafter, every source codes previously described, and randomly generated data, are available as supplementary data.

553 4.2. Results

Results of the comparison performed on Gauss-EQMOM for moment sets far from the boundary of the realisable moment space are given in Table 1. Similar tables are available as supplementary data for all kernels and moment sets.

Table 1: Comparison of Gauss EQMOM implementations corresponding to Fig. 1b and 1c for moment sets far from the frontier of realisability. The count of FLOP details the operations related to (i) the matrix-vector product $A_{2P}^{-1}(\sigma) \cdot m_{2P}$, (ii) the Chebyshev Algorithm (CA), (iii) the Quadrature Computation (QC) and (iv) a miscellaneous category. Results are given as mean±standard-deviation among 10⁴ moment sets.

	0 1					0
			P = 2	P = 3	P = 4	P = 5
New		$oldsymbol{A}_{2P}^{-1}(\sigma)$	$237_{\pm 59}$	$767_{\pm 141}$	$1709_{\pm 253}$	$3201_{\pm 476}$
approach	CA FLOP QC Misc.		$177_{\pm 40}$	$477_{\pm 83}$	$979_{\pm 139}$	$1751_{\pm 251}$
			$52_{\pm 0}$	$474_{\pm 42}$	$995_{\pm 120}$	$1746_{\pm 188}$
			$54_{\pm 12}$	$65_{\pm 11}$	$75_{\pm 11}$	$86_{\pm 12}$
		Total	$519_{\pm 112}$	$1783_{\pm 242}$	$3759_{\pm 441}$	$6784_{\pm 830}$
	Evaluat	tions	$12_{\pm 3}$	$14_{\pm 2}$	$17_{\pm 2}$	$19_{\pm 3}$
	Run-tir	ne (ms)	$1_{\pm 0}$	$2_{\pm 0}$	$3_{\pm 0}$	$4_{\pm 1}$
Former		$oldsymbol{A}_{2P}^{-1}(\sigma)$	$295_{\pm 161}$	$1433_{\pm 423}$	$4060_{\pm 869}$	$8516_{\pm 1870}$
approach		CA	$202_{\pm 102}$	$853_{\pm 241}$	$2246_{\pm 467}$	$4509_{\pm 967}$
	FLOP	QC	$742_{\pm 377}$	$9171_{\pm 2910}$	$24997_{\pm 9966}$	$52312_{\pm 14096}$
		Misc.	$191_{\pm 99}$	$430_{\pm 129}$	$804_{\pm 156}$	$1298_{\pm 251}$
		Total	$1429_{\pm 739}$	$11887_{\pm 3603}$	$32108_{\pm 10645}$	$66635_{\pm 16085}$
	Evaluations		$14_{\pm 7}$	$26_{\pm 7}$	$39_{\pm 8}$	$50_{\pm 11}$
	Run-time (ms)		$1_{\pm 1}$	$9_{\pm 3}$	$17_{\pm 5}$	$31_{\pm 7}$
Gain in	FLOP		$59.1\%_{\pm 12.3\%}$	$84.2\%_{\pm 3.5\%}$	$87.9\%_{\pm 2.5\%}$	$88.0\%_{\pm 13.1\%}$
	Evaluations		$8.6\%_{\pm 27.7\%}$	$40.9\%_{\pm 17.7\%}$	$54.2\%_{\pm 12.8\%}$	$53.0\%_{\pm 55.2\%}$
	Run-time		$53.2\%_{\pm 13.2\%}$	$81.9\%_{\pm 4.2\%}$	$84.0\%_{\pm 3.6\%}$	$83.3\%_{\pm 18.0\%}$

Table 2 underlines a decrease in the number of tested σ values, in particular for high order reconstructions. This decrease is mainly due to the fact that in the former approach, if $m_{N-1}^*(\sigma)$ turns out not to be realisable, the objective function is set to a arbitrarily high negative value. The use of such an arbitrary value slows down the convergence of the nonlinear equation solver. Meanwhile, the new approach never makes use of arbitrary values, all the elements of the vectors of realisability criteria $(\boldsymbol{b}_P^*(\sigma), \boldsymbol{\zeta}_{2P}^*(\sigma) \text{ or } \boldsymbol{p}_{2P}^*(\sigma))$ are used one after the other which yields a better choice of the next tested σ value.

⁵⁶⁴ Moreover, for kernels defined on $\Omega_{\xi} =]0, +\infty[$ and in situations illustrated in Fig. 3c, ⁵⁶⁵ the former approach may switch from a root search to a minimization process if no root is ⁵⁶⁶ found. This induces numerous supplementary tested σ values before convergence is reached ⁵⁶⁷ while this situation never occurs in the new approach.

A significant drop in the total number of FLOP can be observed in Table 3. This was expected and is mainly justified by the fact that the quadrature computation is only called

		P = 2	P = 3	P = 4	P = 5
Course	Strict	$8.6\%_{\pm 27.7\%}$	$40.9\%_{\pm 17.7\%}$	$54.2\%_{\pm 12.8\%}$	$53.0\%_{\pm 55.2\%}$
Gauss	Weak	$10.2\%_{\pm 25.5\%}$	$43.0\%_{\pm 17.4\%}$	$54.7\%_{\pm 20.0\%}$	$42.4\%_{\pm 86.3\%}$
Laplaco	Strict	$8.6\%_{\pm 28.3\%}$	$41.1\%_{\pm 17.6\%}$	$54.3\%_{\pm 12.7\%}$	$53.2\%_{\pm 54.4\%}$
царіасе	Weak	$9.9\%_{\pm 23.7\%}$	$43.3\%_{\pm 17.4\%}$	$54.8\%_{\pm 19.9\%}$	$42.5\%_{\pm 85.7\%}$
I am nameal	Strict	$8.8\%_{\pm 45.9\%}$	$21.9\%_{\pm 32.8\%}$	$30.5\%_{\pm 27.0\%}$	$49.5\%_{\pm 23.8\%}$
Log-normai	Weak	$4.0\%_{\pm 38.1\%}$	$17.8\%_{\pm 26.4\%}$	$40.0\%_{\pm 27.0\%}$	$59.1\%_{\pm 16.1\%}$
Carrana	Strict	$15.5\%_{\pm 38.2\%}$	$24.7\%_{\pm 31.1\%}$	$34.9\%_{\pm 30.1\%}$	$57.2\%_{\pm 24.9\%}$
Gaiiiiia	Weak	$7.5\%_{\pm 28.3\%}$	$20.7\%_{\pm 26.0\%}$	$47.2\%_{\pm 29.6\%}$	$65.4\%_{\pm 16.0\%}$
Waibull	Strict	$26.3\%_{\pm 35.0\%}$	$27.2\%_{\pm 30.2\%}$	$32.4\%_{\pm 28.1\%}$	$54.0\%_{\pm 25.8\%}$
	Weak	$9.4\%_{\pm 15.7\%}$	$19.4\%_{\pm 17.7\%}$	$41.8\%_{\pm 25.3\%}$	$63.3\%_{\pm 16.4\%}$

Table 2: Gain in number of tested σ values.

once in the new approach whilst it is called for most tested σ values in the former momentinversion procedure. This quadrature, which consists in the computation of the eigenvalues and eigenvectors of a tridiagonal symmetric matrix, is the most expensive operation used in the EQMOM moment-inversion procedure.

Concerning the impact of whether the moment sets are close or far from the boundary of the moment space, no significant difference appears in Tables 2 and 3. This implies that there are no preferential situations where the former approach could be more interesting. The new EQMOM core procedure should be used against all moment sets.

La	Table 5. Gain in FLOT for an tested kernels and moment sets.					
		P=2	P = 3	P = 4	P = 5	
Cause	Strict	$59.1\%_{\pm 12.3\%}$	$84.2\%_{\pm 3.5\%}$	$87.9\%_{\pm 2.5\%}$	$88.0\%_{\pm 13.1\%}$	
Gauss	Weak	$59.3\%_{\pm 11.3\%}$	$84.4\%_{\pm 3.7\%}$	$87.8\%_{\pm 4.8\%}$	$85.3\%_{\pm 20.4\%}$	
Laplaco	Strict	$64.2\%_{\pm 10.7\%}$	$87.5\%_{\pm 2.7\%}$	$91.0\%_{\pm 1.7\%}$	$91.6\%_{\pm 8.9\%}$	
Laplace	Weak	$64.1\%_{\pm 9.1\%}$	$87.5\%_{\pm 2.9\%}$	$90.8\%_{\pm 3.5\%}$	$89.6\%_{\pm 14.1\%}$	
I or normal	Strict	$58.9\%_{\pm 20.0\%}$	$85.6\%_{\pm 5.8\%}$	$89.1\%_{\pm 4.3\%}$	$93.3\%_{\pm 3.5\%}$	
Log-normai	Weak	$56.7\%_{\pm 16.7\%}$	$84.7\%_{\pm 4.6\%}$	$90.8\%_{\pm 4.4\%}$	$94.7\%_{\pm 2.4\%}$	
Commo	Strict	$58.2\%_{\pm 18.8\%}$	$82.1\%_{\pm 7.3\%}$	$85.7\%_{\pm 6.8\%}$	$91.6\%_{\pm 5.4\%}$	
Gaiiiiia	Weak	$54.2\%_{\pm 14.1\%}$	$81.1\%_{\pm 6.1\%}$	$88.7\%_{\pm 6.9\%}$	$93.4\%_{\pm 3.5\%}$	
Waibull	Strict	$67.7\%_{\pm 15.1\%}$	$87.4\%_{\pm 5.1\%}$	$90.0\%_{\pm 4.2\%}$	$94.2\%_{\pm 3.6\%}$	
WEIDUII	Weak	$61.0\%_{\pm 6.5\%}$	$86.7\%_{\pm 3.0\%}$	$91.8\%_{\pm 3.9\%}$	$95.7\%_{\pm 2.3\%}$	

Table 3: Gain in FLOP for all tested kernels and moment sets.

Overall, one observes a net decrease in the number of floating-point operations and in the computation run-times of 60% up to 95% for these implementations of EQMOM and the tested moment sets.

One final interesting observation is the evolution of variability in the computational cost of each EQMOM reconstruction. This is illustrated by the ratios standard-deviation/mean shown in Table 4. The high ratios occurring for the former approach, ranging from 24% up to 794%, show that the distributions of required FLOP by EQMOM reconstruction are highly skewed. This leads to high probabilities of significantly different computational costs between different moment sets. On the other hand, that ratio never goes higher than 26% for the new approach if P = 2, and 16% for higher order reconstructions. It implies that the distribution of numerical cost is more narrow and that this new approach will induce more consistent numerical costs among different moment sets. This is a salient feature of this new moment-inversion algorithm as it allows better load-balancing in high performance computing, in particular in highly parallelized CFD softwares.

•	-	P=2	P=3	P=4	P = 5
Cauga	New approach	0.22	0.14	0.12	0.12
Gauss	Former approach	0.52	0.30	0.33	0.24
Laplaco	New approach	0.21	0.12	0.11	0.11
Laplace	Former approach	0.67	0.30	0.43	0.43
Log pormal	New approach	0.26	0.15	0.12	0.11
Log-normai	Former approach	1.10	0.97	7.94	6.48
Commo	New approach	0.25	0.16	0.15	0.15
Gamma	Former approach	1.26	1.14	0.87	3.23
Weibul	New approach	0.21	0.13	0.12	0.11
weibul	Former approach	1.05	1.46	7.21	6.41

Table 4: Ratio standard-deviation/mean of the distribution of total number of FLOP required by EQMOM reconstructions. Only moment sets generated far from the frontier of realisability are considered.

593 5. Conclusion

The first developments relative to the Extended Quadrature Method of Moments are 594 quite recent [17]. Most of these developments were dedicated to widening the use of this 595 method to new application cases, in particular by adding new reconstruction kernels to 596 the EQMOM formalism, and to demonstrate its stability and accuracy compared to other 597 methods. This article summarised all of these developments, relative to the Gaussian kernel 598 [17], to the Log-normal kernel [21] and to the Gamma and Beta kernels [18]. It was also 599 shown that at least two other kernels are perfectly compatible with the EQMOM formalism: 600 the Laplace and Weibull kernels. 601

The youth of EQMOM explains that there is still room left for improvements. The core of this method –the moment-inversion procedure– is an iterative process which is its computational bottleneck. Nguyen et al. [1] proposed some modifications, compared to previous implementations, in order to stabilise the method and to speed-up its resolution, namely the use of Ridder's method instead of bounded-secant or dichotomic methods to solve the non-linear problem, and the realisability checks performed prior to the quadrature computation.

Further improvements were proposed by shifting the resolution toward a new paradigm. This results in a significant decrease in computational cost of about 60% - 95% in terms of required floating-point operations. This resulted in our MATLAB implementations in a similar gain in terms of computation wall-time. Moreover, the new approach offers more consistent numerical costs which will be beneficial to load-balancing in parallelized software. In multiple works [1, 18, 30], EQMOM has been compared to other methods (Maximum Entropy approach or sectional methods) and exhibited (i) similar accuracy even with a lower number of resolved variables, and (ii) faster or comparable computation times. The new improvements of EQMOM will make it even more competitive as its stability and accuracy are kept while reducing the gap in terms of numerical cost between EQMOM and other cheaper methods such as Gauss or Gauss-Radau quadratures.

We strongly believe that transparency about these developments will help further refine-620 ments of EQMOM. For that reason, all sources used to generate figures and data in this article 621 are provided as supplementary data. We also release all our EQMOM source codes both with 622 this article and in an open-access GIT repository (url: https://gitlab.com/open-eqmom). 623 It will be updated as well as supplemented with implementations of EQMOM in languages 624 other than MATLAB. In the case of the Beta reconstruction kernel, some suggestions for 625 further improvements in terms of accuracy and stability were listed in section 3.5. These 626 will be tackled in ulterior work. 627

628 Acknowledgments

The authors thank Sanofi Chimie - C&BD Biochemistry Vitry for its financial support. The authors would also like to thank Prof. Frédérique Laurent and Prof. Rodney O. Fox for the precious discussions and insights about EQMOM and the developments presented in this paper. The authors declare no conflict of interest.

633 References

- [1] T. Nguyen, F. Laurent, R. Fox, M. Massot, Solution of population balance equations
 in applications with fine particles: Mathematical modeling and numerical schemes, J.
 Comput. Phys. 325 (2016) 129–156, ISSN 0021-9991, doi:10.1016/j.jcp.2016.08.017.
- [2] P. Moilanen, M. Laakkonen, O. Visuri, V. Alopaeus, J. Aittamaa, Modelling mass transfer in an aerated 0.2 m3 vessel agitated by Rushton, Phasejet and Combijet impellers,
 Chem. Eng. J. 142 (1) (2008) 95–108, ISSN 1385-8947, doi:10.1016/j.cej.2008.01.033.
- [3] A. Buffo, V. Alopaeus, A novel simplified multivariate PBE solution method for mass
 transfer problems, Chem. Eng. Sci. 172 (Supplement C) (2017) 463–475, ISSN 00092509, doi:10.1016/j.ces.2017.06.036.
- [4] J. Morchain, J. Gabelle, A. Cockx, A coupled population balance model and CFD approach for the simulation of mixing issues in lab-scale and industrial bioreactors, AIChE J. 60 (1) (2014) 27–40, ISSN 1547-5905, doi:10.1002/aic.14238.
- [5] A. Heins, R. L. Fernandes, K. V. Gernaey, A. E. Lantz, Experimental and in silico investigation of population heterogeneity in continuous Sachharomyces cerevisiae scale-down fermentation in a two-compartment setup, J. Chem. Technol. Biotechnol. 90 (2) (2015) 324–340, ISSN 1097-4660, doi:10.1002/jctb.4532.

- [6] S. Salenbauch, A. Cuoci, A. Frassoldati, C. Saggese, T. Faravelli, C. Hasse, Modeling
 soot formation in premixed flames using an Extended Conditional Quadrature Method
 of Moments, Combust. Flame 162 (6) (2015) 2529–2543, ISSN 0010-2180, doi:10.1016/
 j.combustflame.2015.03.002.
- [7] A. Wick, T.-T. Nguyen, F. Laurent, R. O. Fox, H. Pitsch, Modeling soot oxidation
 with the Extended Quadrature Method of Moments, Proc. Combust. Inst. 36 (1) (2017)
 789–797, ISSN 1540-7489, doi:10.1016/j.proci.2016.08.004.
- [8] J. Akroyd, A. J. Smith, R. Shirley, L. R. McGlashan, M. Kraft, A coupled CFD-population balance approach for nanoparticle synthesis in turbulent reacting flows, Chem. Eng. Sci. 66 (17) (2011) 3792–3805, ISSN 0009-2509, doi:10.1016/j.ces.2011.05.
 006.
- [9] P. Fede, O. Simonin, P. Villedieu, Monte-Carlo simulation of colliding particles or coalescing droplets transported by a turbulent flow in the framework of a joint fluidparticle pdf approach, Int. J. Multiph. Flow 74 (Supplement C) (2015) 165–183, ISSN 0301-9322, doi:10.1016/j.ijmultiphaseflow.2015.04.006.
- [11] S. Kumar, D. Ramkrishna, On the solution of population balance equations by dis cretization II. A moving pivot technique, Chem. Eng. Sci. 51 (8) (1996) 1333–1342,
 ISSN 0009-2509, doi:10.1016/0009-2509(95)00355-X.
- [12] M. Massot, F. Laurent, D. K. S. de Chaisemartin, A robust moment method for evaluation of the disappearance rate of evaporating sprays, SIAM J. Appl. Math. 70 (8)
 (2010) 3203–3234, doi:10.1137/080740027.
- [13] V. John, I. Angelov, A. Öncül, D. Thévenin, Techniques for the reconstruction of a distribution from a finite number of its moments, Chem. Eng. Sci. 62 (11) (2007) 2890–2904, ISSN 0009-2509, doi:10.1016/j.ces.2007.02.041.
- [14] L. Mead, N. Papanicolaou, Maximum entropy in the problem of moments, J. Math.
 Phys. 25 (8) (1984) 2404–2417.
- [15] A. Tagliani, Hausdorff moment problem and maximum entropy: A unified approach, Appl. Math. Comput. 105 (2-3) (1999) 291–305, ISSN 0096-3003, doi:10.1016/
 S0096-3003(98)10084-X.
- [16] G. Athanassoulis, P. Gavriliadis, The truncated Hausdorff moment problem solved by
 using kernel density functions, Probab. Eng. Mech. 17 (3) (2002) 273–291, ISSN 02668920, doi:10.1016/S0266-8920(02)00012-7.
- [17] C. Chalons, R. Fox, M. Massot, A multi-Gaussian quadrature method of moments for
 gas-particle flows in a LES framework, in: Proceedings of the 2010 Summer Program,
 Center for turbulence Research, Stanford University, 347–358, 2010.

- [18] C. Yuan, F. Laurent, R. Fox, An extended quadrature method of moments for population balance equations, J. Aerosol Sci. 51 (0) (2012) 1–23, ISSN 0021-8502, doi: 10.1016/j.jaerosci.2012.04.003.
- [19] A. Passalacqua, F. Laurent, E. Madadi-Kandjani, J. Heylmun, R. Fox, An open-source
 quadrature-based population balance solver for OpenFOAM, Chem. Eng. Sci. ISSN
 0009-2509, doi:10.1016/j.ces.2017.10.043.
- [20] D. Marchisio, R. Fox, Computational Models for Polydisperse Particulate and Multi phase Systems, Cambridge Series in Chemical Engineering, Cambridge University Press,
 ISBN 9781107328174, 2013.
- E. Madadi-Kandjani, A. Passalacqua, An extended quadrature-based moment method
 with log-normal kernel density functions, Chem. Eng. Sci. 131 (0) (2015) 323–339, ISSN 0009-2509, doi:10.1016/j.ces.2015.04.005.
- [22] MATLAB, version 9.0 (R2016a), The MathWorks, Inc., Natick, Massachusetts, United
 States, 2016.
- [23] R. McGraw, Description of Aerosol Dynamics by the Quadrature Method of Moments,
 Aerosol Sci. Technol. 27 (2) (1997) 255–265, doi:10.1080/02786829708965471.
- [24] W. Gautschi, Orthogonal Polynomials: Computation and Approximation, Numerical
 mathematics and scientific computation, Oxford University Press, ISBN 9780198506720,
 2004.
- [25] P. Henrici, The Quotient-Difference algorithm, Natl. Bur. Stand., Appl. Math. Ser. 49 (1958) 23–46.
- [26] H. Dette, W. Studden, The theory of canonical moments with applications in statistics,
 probability, and analysis, John Wiley & Sons, New York; Chichester, 1997.
- [27] R. Gordon, Error Bounds in Equilibrium Statistical Mechanics, J. Math. Phys. 9 (1968)
 655, doi:10.1063/1.1664624.
- [28] J. Wheeler, Modified moments and Gaussian quadratures, Rocky Mountain J. Math.
 4 (2) (1974) 287–296, doi:RMJ-1974-4-2-287.
- [29] N. Lebaz, A. Cockx, M. Spérandio, J. Morchain, Reconstruction of a distribution
 from a finite number of its moments: A comparative study in the case of depolymerization process, Comput. Chem. Eng. 84 (2016) 326–337, ISSN 0098-1354, doi:
 10.1016/j.compchemeng.2015.09.008.
- [30] M. Pigou, J. Morchain, P. Fede, M. Penet, G. Laronze, An assessment of methods of moments for the simulation of population dynamics in large-scale bioreactors, Chem.
 Eng. Sci. 171 (2017) 218–232, ISSN 0009-2509, doi:10.1016/j.ces.2017.05.026.
- [31] J. Favard, Sur les polynomes de Tchebicheff, C. r. hebd. séances Acad. sci. 200 (1935)
 2052–2053.

- [32] T. Chihara, An introduction to orthogonal polynomials, no. vol. 13 in Mathematics and its applications, Gordon and Breach, ISBN 9780677041506, 1978.
- [33] J. Shohat, J. Tamarkin, The Problem of Moments, Mathematical Surveys and Mono graphs, American Mathematical Society, 4 edn., 1943.
- [34] H. Wall, Analytic Theory of Continued Fractions, The University series in higher math ematics, Van Nostrand, 1948.
- [35] W. Ford, Chapter 19 The Symmetric Eigenvalue Problem, in: Numerical Linear Al gebra with Applications, Academic Press, Boston, ISBN 978-0-12-394435-1, 439-468,
 doi:10.1016/B978-0-12-394435-1.00019-3, 2015.
- [36] M. Wilck, A general approximation method for solving integrals containing a lognormal
 weighting function, J. Aerosol Sci. 32 (9) (2001) 1111–1116, ISSN 0021-8502, doi:10.
 1016/S0021-8502(01)00044-1.
- [37] J. Shen, T. Tang, L.-L. Wang, Orthogonal Polynomials and Related Approximation
 Results, Springer Berlin Heidelberg, Berlin, Heidelberg, ISBN 978-3-540-71041-7, 47–
 140, doi:10.1007/978-3-540-71041-7_3, 2011.

Appendix A. Chebyshev algorithm 739

The Chebyshev algorithm allows to compute the three-term recurrence coefficients of the 740 monic polynomials orthogonal to a measure $d\mu(\xi)$ whose moments are given by the vector 741 $\boldsymbol{m}_N = [m_0, \ldots, m_N]$. This version of the algorithm fills column-wise a $N + 1 \times \lfloor \frac{N+1}{2} \rfloor$ matrix 742 denoted \boldsymbol{S} . 743

First, fill the first column with the moments $S_{i,0} = m_i$, compute $a_0 = m_1/m_0$ and fill the 744 second column with $S_{i,1} = S_{i+1,0} - a_0 S_{i,0}, \quad \forall i \in \{1, \dots, N-1\}.$ 745

Then iterate for $j \in \{2, \dots, \lceil \frac{N-1}{2} \rceil\}$: 746

$$a_{j-1} = \frac{S_{j,j-1}}{S_{j-1,j-1}} - \frac{S_{j-1,j-2}}{S_{j-2,j-2}}$$

$$b_{j-1} = \frac{S_{j-1,j-1}}{S_{j-2,j-2}}$$

$$S_{i,j} = S_{i+1,j-1} - a_{j-1}S_{i,j-1} - b_{j-1}S_{i,j-2}, \qquad i \in \{j, \dots, N-j\}$$

Appendix B. Kernels for EQMOM 747

There exists multiple variations of the EQMOM method depending on the Kernel Density 748 Function that is used for the reconstruction in Eq. (15). This section details the specificities 749 of multiple KDF that were found to be compatible with the EQMOM procedure. It details 750 for each kernel 751

1. the actual expression of that kernel $\delta_{\sigma}(\xi, \xi_m)$; 752

- 2. the expression of its moments; 753
- 3. the matrix $A_n(\sigma)$ that allows the transfer between the raw moments of the reconstruc-754 tion $\widetilde{\boldsymbol{m}}_n$ and its degenerated moments \boldsymbol{m}_n^* ; 755
- 4. the nested quadrature rules suiting this kernel; 756
- 5. the analytical solutions available for one-node EQMOM (P = 1). 757

Two-nodes analytical solutions exist for the Gaussian, Gamma, Laplace and Log-normal 758 kernels and are accessible using the same methodology than that used by Chalons et al. [17] 759 for the Gaussian kernel. These solutions are not detailed here but are implemented in the 760 MATLAB code given in supplementary data. 761

All definitions of matrices $A_n(\sigma)$ are given using zero-offset. The element of the first line 762 and column of this matrix then reads $A_{0,0}(\sigma)$. 763

- Appendix B.1. Gaussian kernel 764
- Appendix B.1.1. Definition 765

The Gaussian kernel $\delta_{\sigma}^{(G)}(\xi,\xi_m)$ was first used in EQMOM by Chalons et al. [17]. It is 766 defined on $\Omega_{\xi} = \left] -\infty, +\infty \right[$ by 767

$$\delta_{\sigma}^{(G)}(\xi,\xi_m) = \frac{1}{\sigma\sqrt{2\pi}} \exp\left(-\frac{(\xi-\xi_m)^2}{2\sigma^2}\right) \tag{B.1}$$

768 Appendix B.1.2. Moments and linear system

⁷⁶⁹ Moments of the Gaussian kernel are given by:

$$\int_{-\infty}^{+\infty} \xi^k \delta_{\sigma}^{(G)}(\xi, \xi_m) \mathrm{d}\xi = \sum_{j=0}^{\lfloor k/2 \rfloor} \frac{k!}{j!(k-2j)!} \left(\frac{\sigma^2}{2}\right)^j \xi_m^{k-2j}$$
(B.2)

Moments of the distribution $\widetilde{n}(\xi) = \sum_{i=1}^{P} w_i \delta_{\sigma}^{(G)}(\xi, \xi_i)$ are given by the linear system

$$\widetilde{\boldsymbol{m}}_n = \boldsymbol{A}_n^{(G)}(\sigma) \cdot \boldsymbol{m}_n^* \tag{B.3}$$

771 with

$$A_{i,j}^{(G)}(\sigma) = \begin{cases} 0 & \text{if } j > i \text{ or } (i-j \mod 2) = 1\\ \frac{i!}{\left(\frac{i-j}{2}\right)!j!} \left(\frac{\sigma^2}{2}\right)^{\frac{i-j}{2}} & \text{otherwise} \end{cases}$$
(B.4)

The inverse of this matrix is given by:

$$A_{i,j}^{(G)-1}(\sigma) = \begin{cases} 0 & \text{if } j > i \text{ or } (i-j \mod 2) = 1\\ \frac{i!}{\left(\frac{i-j}{2}\right)!j!} \left(-\frac{\sigma^2}{2}\right)^{\frac{i-j}{2}} & \text{otherwise} \end{cases}$$
(B.5)

which translates, for the case n = 4, into:

$$\begin{bmatrix} \tilde{m}_{0} \\ \tilde{m}_{1} \\ \tilde{m}_{2} \\ \tilde{m}_{3} \\ \tilde{m}_{4} \end{bmatrix} = \begin{pmatrix} 1 & & & 0 \\ 0 & 1 & & & \\ \sigma^{2} & 0 & 1 & & \\ 0 & 3\sigma^{2} & 0 & 1 & \\ 3\sigma^{4} & 0 & 6\sigma^{2} & 0 & 1 \end{pmatrix} \cdot \begin{bmatrix} m_{0}^{*} \\ m_{1}^{*} \\ m_{2}^{*} \\ m_{3}^{*} \\ m_{4}^{*} \end{bmatrix} = \begin{pmatrix} 1 & & & 0 \\ 0 & 1 & & & \\ -\sigma^{2} & 0 & 1 & & \\ 0 & -3\sigma^{2} & 0 & 1 & \\ 3\sigma^{4} & 0 & -6\sigma^{2} & 0 & 1 \end{pmatrix} \cdot \begin{bmatrix} \tilde{m}_{0} \\ \tilde{m}_{1} \\ \tilde{m}_{2} \\ \tilde{m}_{3} \\ \tilde{m}_{4} \end{bmatrix}$$
(B.6)

774 Appendix B.1.3. Moment preserving nested quadrature

The approximation of integral properties using Gauss EQMOM is performed through the following nested quadrature:

$$\int_{-\infty}^{+\infty} f(\xi) n(\xi) d\xi \approx \frac{1}{\sqrt{\pi}} \sum_{i=1}^{P} w_i \sum_{j=1}^{Q} \omega_j f\left(\xi_i + \sigma \lambda_j \sqrt{2}\right)$$
(B.8)

with \boldsymbol{w}_P , $\boldsymbol{\xi}_P$ and σ the EQMOM reconstruction parameters computed from \boldsymbol{m}_{2P} ; $\boldsymbol{\omega}_Q$ and $\boldsymbol{\lambda}_Q$ are the weights and nodes of a Q-nodes Gauss-Hermite quadrature rule (see Appendix C).

- 780 Appendix B.1.4. Single node analytical solution
- The case P = 1 has the following analytical solution:

$$w_{1} = m_{0}$$

$$\xi_{1} = \frac{m_{1}}{m_{0}}$$

$$\sigma = \frac{\sqrt{m_{2}m_{0} - m_{1}^{2}}}{m_{0}}$$

- 782 Appendix B.2. Laplace kernel
- 783 Appendix B.2.1. Definition
- The Laplace kernel $\delta_{\sigma}^{(\lambda)}(\xi,\xi_m)$ is defined on $\Omega_{\xi} =]-\infty, +\infty[$ by

$$\delta_{\sigma}^{(\lambda)}(\xi,\xi_m) = \frac{1}{2\sigma} \exp\left(-\frac{|\xi-\xi_m|}{\sigma}\right) \tag{B.9}$$

- 785 Appendix B.2.2. Moments and linear system
- ⁷⁸⁶ Moments of the Laplace kernel are given by

$$\int_{-\infty}^{+\infty} \xi^k \delta_{\sigma}^{(\lambda)}(\xi, \xi_m) \mathrm{d}\xi = \sum_{j=0}^k \frac{k!}{(k-j)!} \frac{1 + (-1)^j}{2} \xi_m^{k-j} \sigma^j \tag{B.10}$$

Moments of the distribution $\widetilde{n}(\xi) = \sum_{i=1}^{P} w_i \delta_{\sigma}^{(\lambda)}(\xi, \xi_i)$ are given by the linear system

$$\widetilde{\boldsymbol{m}}_n = \boldsymbol{A}_n^{(\lambda)}(\sigma) \cdot \boldsymbol{m}_n^* \tag{B.11}$$

788 with

$$A_{i,j}^{(\lambda)}(\sigma) = \begin{cases} 0 & \text{if } j > i \text{ or } (i-j \mod 2) = 1\\ \frac{i!}{j!}\sigma^{i-j} & \text{otherwise} \end{cases}$$
(B.12)

⁷⁸⁹ The inverse matrix is defined by

$$A_{i,j}^{(\lambda) -1}(\sigma) = \begin{cases} 1 & \text{if } i = j \\ -(j+1)(j+2)\sigma^2 & \text{if } i = j+2 \\ 0 & \text{otherwise} \end{cases}$$
(B.13)

⁷⁹⁰ which translates for n = 6 into

⁷⁹¹ Appendix B.2.3. Moment preserving nested quadrature

The approximation of integral properties using Laplace EQMOM is performed through
 the following nested quadrature:

$$\int_{-\infty}^{+\infty} f(\xi) n(\xi) d\xi \approx \sum_{i=1}^{P} w_i \sum_{j=1}^{Q} \omega_j f\left(\xi_i + \sigma \lambda_j\right)$$
(B.16)

with \boldsymbol{w}_P , $\boldsymbol{\xi}_P$ and σ the EQMOM reconstruction parameters computed from \boldsymbol{m}_{2P} ; $\boldsymbol{\omega}_Q$ and $\boldsymbol{\lambda}_Q$ are the weights and nodes of a Q-nodes "Gauss-Laplace" quadrature rule (see Appendix C).

⁷⁹⁷ Appendix B.2.4. Single node analytical solution

The case P = 1 has the following analytical solution:

$$w_{1} = m_{0}$$

$$\xi_{1} = \frac{m_{1}}{m_{0}}$$

$$\sigma = \sqrt{\frac{m_{2}m_{0} - m_{1}^{2}}{2m_{0}^{2}}}$$

799 Appendix B.3. Log-normal kernel

800 Appendix B.3.1. Definition

The Log-normal kernel $\delta_{\sigma}^{(L)}(\xi, \xi_m)$ was first used in EQMOM by Madadi-Kandjani and Passalacqua [21]. It is defined on $\Omega_{\xi} =]0, +\infty[$ by

$$\delta_{\sigma}^{(L)}(\xi,\xi_m) = \frac{1}{\sigma\xi\sqrt{2\pi}} \exp\left(-\frac{\left(\log(\xi) - \log(\xi_m)\right)^2}{2\sigma^2}\right) \tag{B.17}$$

⁸⁰³ Appendix B.3.2. Moments and linear system

⁸⁰⁴ Moments of the Log-normal kernel are given by

$$\int_0^{+\infty} \xi^k \delta_\sigma^{(L)}(\xi, \xi_m) \mathrm{d}\xi = \xi_m^k z^{k^2} \qquad \text{with } z = e^{\sigma^2/2} \tag{B.18}$$

Moments of the distribution $\widetilde{n}(\xi) = \sum_{i=1}^{P} w_i \delta_{\sigma}^{(L)}(\xi, \xi_i)$ are given by

$$\widetilde{m}_k = m_k^* z^{k^2} \tag{B.19}$$

⁸⁰⁶ This can be translated into a linear system

$$\widetilde{\boldsymbol{m}}_n = \boldsymbol{A}_n^{(L)}(\sigma) \cdot \boldsymbol{m}_n^* \tag{B.20}$$

with $A_n^{(L)}(\sigma)$ a diagonal matrix:

$$A_{i,j}^{(L)}(\sigma) = \begin{cases} z^{i^2} & \text{if } i = j \\ 0 & \text{otherwise} \end{cases}$$
(B.21)

⁸⁰⁸ whose inverse matrix is directly given by

$$A_{i,j}^{(L)-1}(\sigma) = \begin{cases} z^{-i^2} & \text{if } i = j\\ 0 & \text{otherwise} \end{cases}$$
(B.22)

Appendix B.3.3. Low cost nested quadrature

A variable change allows approximating integral properties over a LogN EQMOM reconstruction using Gauss-Hermite quadratures [21]:

$$\int_{0}^{+\infty} f(\xi) n(\xi) d\xi \approx \frac{1}{\sqrt{\pi}} \sum_{i=1}^{P} w_i \sum_{j=1}^{Q} \omega_j f\left(\xi_i \exp\left(\sigma \lambda_j \sqrt{2}\right)\right)$$
(B.23)

with \boldsymbol{w}_P , $\boldsymbol{\xi}_P$ and σ the EQMOM reconstruction parameters computed from \boldsymbol{m}_{2P} ; $\boldsymbol{\omega}_Q$ and $\boldsymbol{\lambda}_Q$ are the weights and nodes of a Q-nodes Gauss-Hermite quadrature rule (see Appendix C).

Parameters of this nested quadrature do not depend on σ of the main quadrature nodes $\boldsymbol{\xi}_P$. Consequently, $\boldsymbol{\omega}_Q$ and $\boldsymbol{\lambda}_Q$ only need to be computed once. It is worth noting that this quadrature does not preserve the moments of the distribution and only yields exact results for $f(\xi) = \log(\xi)^k$, $k \in \{0, \ldots, 2\min(P, Q) - 1\}$.

⁸¹⁹ Appendix B.3.4. Moment preserving nested quadrature

Passalacqua et al. [19] suggested the use of Gauss-Wigert quadratures [36] to preserve the moments of a LogN EQMOM reconstruction:

$$\int_{0}^{+\infty} f(\xi) n(\xi) \mathrm{d}\xi \approx \sum_{i=1}^{P} w_i \sum_{j=1}^{Q} \omega_j^{(\sigma)} f\left(\xi_i \lambda_j^{(\sigma)}\right) \tag{B.24}$$

with \boldsymbol{w}_P , $\boldsymbol{\xi}_P$ and σ the EQMOM reconstruction parameters computed from \boldsymbol{m}_{2P} ; $\boldsymbol{\omega}_Q^{(\sigma)}$ and $\boldsymbol{\lambda}_Q^{(\sigma)}$ are the weights and nodes of a Q-nodes Gauss-Wigert quadrature rule of parameter σ (see Appendix C). This quadrature rule must be computed for each value of σ , *i.e.* for each LogN EQMOM reconstruction.

- ⁸²⁶ Appendix B.3.5. Single node analytical solution
- The case P = 1 has the following analytical solution:

$$w_1 = m_0$$

$$\xi_1 = \sqrt{\frac{m_1^4}{m_2 m_0^3}}$$

$$\sigma = \sqrt{\log\left(\frac{m_2 m_0}{m_1^2}\right)}$$

- 828 Appendix B.4. Gamma kernel
- 829 Appendix B.4.1. Definition

The Gamma kernel $\delta_{\sigma}^{(\Gamma)}(\xi, \xi_m)$ was first used in EQMOM by Yuan et al. [18]. It is defined on $\Omega_{\xi} =]0, +\infty[$ by

$$\delta_{\sigma}^{(\Gamma)}(\xi,\xi_m) = \frac{\xi^{(l-1)}\exp(-\xi/\sigma)}{\Gamma(l)\sigma^l} \quad \text{with } l = \frac{\xi_m}{\sigma} \text{ and } \Gamma(x) = \int_0^{+\infty} t^{x-1} e^{-t} dt \quad (B.25)$$

- ⁸³² Appendix B.4.2. Moments and linear system
- ⁸³³ Moments of the Gamma kernel are given by

$$\int_{0}^{+\infty} \xi^{k} \delta_{\sigma}^{(\Gamma)}(\xi, \xi_{m}) \mathrm{d}\xi = G_{k}(\xi_{m}, \sigma) = \begin{cases} 1 & \text{if } k = 0\\ \prod_{j=0}^{k-1} (\xi_{m} + j\sigma) & \text{otherwise} \end{cases}$$
(B.26)

⁸³⁴ Moments of the distribution $\tilde{n}(\xi) = \sum_{i=1}^{P} w_i \delta_{\sigma}^{(\Gamma)}(\xi, \xi_i)$ are given by the linear system

$$\widetilde{\boldsymbol{m}}_n = \boldsymbol{A}_n^{(\Gamma)}(\sigma) \cdot \boldsymbol{m}_n^* \tag{B.27}$$

835 with

$$A_{i,j}^{(\Gamma)}(\sigma) = \begin{cases} 0 & \text{if } j > i \text{ or } i = 0 \text{ or } j = 0\\ 1 & \text{if } i = 0 \text{ and } j = 0\\ A_{i-1,j-1}^{(\Gamma)}(\sigma) + (i-1)\sigma A_{i-1,j}^{(\Gamma)}(\sigma) & \text{otherwise} \end{cases}$$
(B.28)

⁸³⁶ The inverse of this matrix is given by

$$A_{i,j}^{(\Gamma)-1}(\sigma) = \begin{cases} 0 & \text{if } j > i \text{ or } i = 0 \text{ or } j = 0\\ 1 & \text{if } i = 0 \text{ and } j = 0\\ A_{i-1,j-1}^{(\Gamma)-1}(\sigma) - j\sigma A_{i-1,j}^{(\Gamma)-1}(\sigma) & \text{otherwise} \end{cases}$$
(B.29)

which translates, for n = 6 into

$$\begin{bmatrix} \tilde{m}_{0} \\ \tilde{m}_{1} \\ \tilde{m}_{2} \\ \tilde{m}_{3} \\ \tilde{m}_{4} \\ \tilde{m}_{5} \\ \tilde{m}_{6} \end{bmatrix} = \begin{pmatrix} 1 & & & & & & & & & \\ 0 & 1\sigma & 1 & & & & & & \\ 0 & 2\sigma^{2} & 3\sigma & 1 & & & & & \\ 0 & 2\sigma^{2} & 3\sigma & 1 & & & & & \\ 0 & 2\sigma^{3} & 11\sigma^{2} & 6\sigma & 1 & & & \\ 0 & 24\sigma^{4} & 50\sigma^{3} & 35\sigma^{2} & 10\sigma & 1 & & \\ 0 & 24\sigma^{4} & 50\sigma^{3} & 35\sigma^{2} & 10\sigma & 1 & & \\ 0 & 24\sigma^{4} & 50\sigma^{3} & 35\sigma^{2} & 15\sigma & 1 & \end{pmatrix} \cdot \begin{bmatrix} m_{0}^{*} \\ m_{1}^{*} \\ m_{5}^{*} \\ m_{6}^{*} \end{bmatrix} = \begin{pmatrix} 1 & & & & & & \\ 0 & 1 & & & & & \\ 0 & -\sigma & 1 & & & & \\ 0 & -\sigma & 1 & & & & \\ 0 & -\sigma^{3} & 7\sigma^{2} & -6\sigma & 1 & & \\ 0 & -\sigma^{5} & 31\sigma^{4} & -90\sigma^{3} & 65\sigma^{2} & -15\sigma & 1 \end{pmatrix} \cdot \begin{bmatrix} \tilde{m}_{0} \\ \tilde{m}_{1} \\ \tilde{m}_{2} \\ \tilde{m}_{3} \\ \tilde{m}_{4} \\ \tilde{m}_{5} \\ \tilde{m}_{6} \end{bmatrix}$$
(B.31)

⁸³⁸ Appendix B.4.3. Low cost nested quadrature

A Gauss-Laguerre quadrature can be used to approximate integral properties over a Gamma EQMOM reconstruction:

$$\int_{0}^{+\infty} f(\xi) n(\xi) d\xi \approx \sum_{j=1}^{Q} \omega_j f(\sigma \lambda_j) \sum_{i=1}^{P} \frac{w_i}{\Gamma\left(\frac{\xi_i}{\sigma}\right)} \lambda_j^{\frac{\xi_i}{\sigma} - 1}$$
(B.32)

with \boldsymbol{w}_P , $\boldsymbol{\xi}_P$ and σ the EQMOM reconstruction parameters computed from \boldsymbol{m}_{2P} ; $\boldsymbol{\omega}_Q$ and $\boldsymbol{\lambda}_Q$ are the weights and nodes of a Q-nodes Gauss-Laguerre quadrature rule of parameter $\boldsymbol{\alpha} = 0$ (see Appendix C). The advantage of this quadrature is that it only requires $\boldsymbol{\omega}_Q$ and $\boldsymbol{\lambda}_Q$ to be computed once. However, this quadrature will not preserve the moments of the distribution.

846 Appendix B.4.4. Moment preserving nested quadrature

A generalized Gauss-Laguerre quadrature preserves the moments of a Gamma EQMOM reconstruction:

$$\int_{0}^{+\infty} f(\xi) n(\xi) d\xi \approx \sum_{i=1}^{P} \frac{w_i}{\Gamma\left(\frac{\xi_i}{\sigma}\right)} \sum_{j=1}^{Q} \omega_j^{(\alpha_i)} f\left(\sigma \lambda_j^{(\alpha_i)}\right)$$
(B.33)

with $\boldsymbol{w}_P, \boldsymbol{\xi}_P$ and σ the EQMOM reconstruction parameters computed from $\boldsymbol{m}_{2P}; \boldsymbol{\omega}_Q^{(\alpha_i)}$ and $\boldsymbol{\lambda}_Q^{(\alpha_i)}$ are the weights and nodes of a *Q*-nodes Gauss-Laguerre quadrature rule of parameter ⁸⁵¹ $\alpha_i = \frac{\xi_i}{\sigma} - 1$ (see Appendix C).

The accuracy of this quadrature comes with a cost related to the computation of $\boldsymbol{\omega}_Q^{(\alpha_i)}$ and $\boldsymbol{\lambda}_Q^{(\alpha_i)}$ for each value of α_i .

Appendix B.4.5. Single node analytical solution

The case P = 1 has the following analytical solution:

$$w_1 = m_0$$

$$\xi_1 = \frac{m_1}{m_0}$$

$$\sigma = \frac{m_2}{m_1} - \frac{m_1}{m_0}$$

- 856 Appendix B.5. Weibull kernel
- 857 Appendix B.5.1. Definition
- The Weibull kernel $\delta_{\sigma}^{(W)}(\xi,\xi_m)$ is defined on $\Omega_{\xi} =]0, +\infty[$ by

$$\delta_{\sigma}^{(W)}(\xi,\xi_m) = \frac{1}{\sigma\xi_m} \left(\frac{\xi}{\xi_m}\right)^{\frac{1-\sigma}{\sigma}} \exp\left(-\left(\frac{\xi}{\xi_m}\right)^{1/\sigma}\right) \tag{B.34}$$

- Appendix B.5.2. Moments and linear system
- Moments of the Weibull kernel are given by

$$\int_{0}^{+\infty} \xi^{k} \delta_{\sigma}^{(W)}(\xi, \xi_{m}) \mathrm{d}\xi = \xi_{m}^{k} \Gamma(1 + k\sigma)$$
(B.35)

Moments of the distribution $\widetilde{n}(\xi) = \sum_{i=1}^{P} w_i \delta_{\sigma}^{(W)}(\xi, \xi_i)$ are given by

$$\widetilde{m}_k = m_k^* \Gamma(1 + k\sigma) \tag{B.36}$$

⁸⁶² This can be translated into a linear system

$$\widetilde{\boldsymbol{m}}_n = \boldsymbol{A}_n^{(W)}(\sigma) \cdot \boldsymbol{m}_n^* \tag{B.37}$$

with $A_n^{(W)}(\sigma)$ a diagonal matrix:

$$A_{i,j}^{(W)}(\sigma) = \begin{cases} \Gamma(1+i\sigma) & \text{if } i=j\\ 0 & \text{otherwise} \end{cases}$$
(B.38)

⁸⁶⁴ whose inverse matrix is directly given by

$$A_{i,j}^{(W)-1}(\sigma) = \begin{cases} \frac{1}{\Gamma(1+i\sigma)} & \text{if } i = j\\ 0 & \text{otherwise} \end{cases}$$
(B.39)

⁸⁶⁵ Appendix B.5.3. Low cost nested quadrature

A Gauss-Laguerre quadrature can be used to approximate integral properties over a Weibull EQMOM reconstruction:

$$\int_{0}^{+\infty} f(\xi) n(\xi) d\xi \approx \sum_{i=1}^{P} w_i \sum_{j=1}^{Q} \omega_j f\left(\xi_i \lambda_j^{\sigma}\right)$$
(B.40)

with \boldsymbol{w}_P , $\boldsymbol{\xi}_P$ and σ the EQMOM reconstruction parameters computed from \boldsymbol{m}_{2P} ; $\boldsymbol{\omega}_Q$ and $\boldsymbol{\lambda}_Q$ are the weights and nodes of a Q-nodes Gauss-Laguerre quadrature rule of parameter $\alpha = 0$ (see Appendix C). The advantage of this quadrature is that it only requires $\boldsymbol{\omega}_Q$ and $\boldsymbol{\lambda}_Q$ to be computed once. However, this quadrature will not preserve the moments of the distribution and only yields exact results for $f(\xi) = \xi^{k/\sigma}, \ k \in \{0, \dots, 2\min(P, Q) - 1\}$

⁸⁷³ Appendix B.5.4. Moment preserving nested quadrature

One can produce a Gauss quadrature that preserves the moments of Weibull EQMOM approximations:

$$\int_{0}^{+\infty} f(\xi) n(\xi) \mathrm{d}\xi \approx \sum_{i=1}^{P} w_i \sum_{j=1}^{Q} \omega_j^{(\sigma)} f\left(\xi_i \lambda_j^{(\sigma)}\right) \tag{B.41}$$

with \boldsymbol{w}_P , $\boldsymbol{\xi}_P$ and σ the EQMOM reconstruction parameters computed from \boldsymbol{m}_{2P} ; $\boldsymbol{\omega}_Q^{(\sigma)}$ and $\boldsymbol{\lambda}_Q^{(\sigma)}$ are the weights and nodes of a *Q*-nodes "Gauss-Weibull" quadrature rule of parameter σ (see Appendix C). The weights and nodes of the nested quadrature need to be computed for each value of σ , *i.e.* for each Weibull EQMOM approximation of the NDF.

Appendix B.5.5. Single node numerical solution

The parameters w_1 , ξ_1 and σ of the one-node Weibull EQMOM must be solution of the following system:

$$m_0 = w_1$$
$$\frac{m_1}{\Gamma(1+\sigma)} = w_1 \xi_1$$
$$\frac{m_2}{\Gamma(1+2\sigma)} = w_1 \xi_1^2$$

The first equation gives $w_1 = m_0$ but no explicit solution exists for the two other equations.

⁸⁸⁴ One can however notice that $s = \frac{\sigma}{1+\sigma}$ must be a root of

$$G(s) = \frac{m_2 m_0}{m_1^2} - \frac{\Gamma(\frac{1+s}{1-s})}{\Gamma(\frac{1}{1-s})^2}$$
(B.42)

which is monotonous, defined on $s \in [0, 1]$ and has the following limits

$$G(0) = \frac{m_2 m_0}{m_1^2} > 0$$
$$\lim_{s \to 1^-} G(s) < 0$$

G(s) then admits a single root that can be computed numerically with the Ridder's method. 886 One can also narrow down, at a very low cost, the search interval [0, 1] by using the property 887

$$g_n = G\left(\frac{n}{n+1}\right) = \frac{m_2 m_0}{m_1^2} - \frac{(2n)!}{(n!)^2}$$
(B.43)

with n an integer, which induces the following recurrence relation: 888

s-

$$g_n = c - h_n \tag{B.44}$$

$$h_{n+1} = \left(4 - \frac{2}{n+1}\right)h_n \tag{B.45}$$

with $c = \frac{m_2 m_0}{m_1^2}$ and $h_1 = 2$. 889 The proposed algorithm to identify the root of G(s) is 890 1. Compute $c = \frac{m_2 m_0}{m_1^2}$ 891 • if c < 1, cancel the operation as the moments are not realisable; 892 • if c = 1, s = 0 is the root of G(s); 893 • if c < 2, set $s_l = 0$, $v_l = c - 1$, $s_r = \frac{1}{2}$ and $v_r = c - 2$ and go to step 3. 894 • otherwise, set $s_l = 0$, $v_l = c - 1$ and go to step 2. 895 2. Initialise i = 1, h = 2 and iterate 896 (a) increment i by 1; 897 (b) compute $h = h * (4 - \frac{2}{i})$ 898 • if h = c, then $s = \frac{i}{i+1}$ is a root of G(s); 899 • if h < c, set $s_l = \frac{i}{i+1}$ and $v_l = c - h$; 900 • if h > c, set $s_r = \frac{i}{i+1}$, $v_r = c - h$ and go to step 3. 901 3. Apply the Ridder's method to G(s) on the interval $[s_l, s_r]$ 902 (a) compute $s_{t_1} = \frac{1}{2}(s_l + s_r)$ and $v_{t_1} = G(s_{t_1})$; (b) compute $s_{t_2} = s_{t_1} + (s_{t_1} - s_l) \frac{v_{t_1}}{\sqrt{v_{t_1}^2 - v_l v_r}}$ and $v_{t_2} = G(s_{t_2})$; 903 904 (c) set s_l the highest value between s_l , s_{t_1} and s_{t_2} whose image by G is positive; 905 (d) set s_r the lowest value between s_r , s_{t_1} and s_{t_2} whose image by G is negative; 906 (e) stop the computation if $v_l < \varepsilon(c-1)$ with ε a relative tolerance (e.g. $\varepsilon = 10^{-10}$) 907 and consider s_l as a root of G(s). 908

Once the root of G(s) is identified, compute 909

$$\sigma = \frac{s}{1-s}$$

$$\xi_1 = \frac{m_1}{m_0 \Gamma(1+\sigma)}$$

Note that each iteration of the Ridder's method requires two computations of G(s), that implies four computations of the Gamma function –which is quite expensive– by iteration. This explains the interest of the second step which allows to narrow down the research interval at hardly no cost.

- 914 Appendix B.6. Beta kernel
- 915 Appendix B.6.1. Definition

The Beta kernel $\delta_{\sigma}^{(\beta)}(\xi, \xi_m)$ was first used in EQMOM by Yuan et al. [18]. It is defined on $\Omega_{\xi} =]0, 1[$ by

$$\delta_{\sigma}^{(\beta)}(\xi,\xi_m) = \frac{\xi^{(l-1)}(1-\xi)^{(m-1)}}{B(l,m)} \quad \text{with } l = \frac{\xi_m}{\sigma} \text{ and } m = \frac{1-\xi_m}{\sigma}$$
(B.46)

with $B(l,m) = \int_0^1 x^{(l-1)} (1-x)^{(m-1)} dx$ the beta function.

919 Appendix B.6.2. Moments and linear system

⁹²⁰ Moments of the Beta kernel are given by

$$\int_{0}^{1} \xi^{k} \delta_{\sigma}^{(\beta)}(\xi, \xi_{m}) \mathrm{d}\xi = H_{k}(\xi_{m}, \sigma) = \begin{cases} 1 & \text{if } k = 0\\ \prod_{j=0}^{k-1} \left(\frac{\xi_{m}+j\sigma}{1+j\sigma}\right) & \text{otherwise} \end{cases}$$
(B.47)

Moments of the distribution $\tilde{n}(\xi) = \sum_{i=1}^{P} w_i \delta_{\sigma}^{(\beta)}(\xi, \xi_i)$ are given by the linear system

$$\widetilde{\boldsymbol{m}}_n = \boldsymbol{A}_n^{(\beta)}(\sigma) \cdot \boldsymbol{m}_n^* \tag{B.48}$$

with the elements of $A_n^{(\beta)}(\sigma)$ being computed from the elements of the matrix relative to Gamma EQMOM, $A_n^{(\Gamma)}(\sigma)$:

$$A_{i,j}^{(\beta)}(\sigma) = \frac{A_{i,j}^{(\Gamma)}(\sigma)}{F_i(\sigma)}$$
(B.49)

$$F_i(\sigma) = \begin{cases} 1 & \text{if } i \le 1\\ (1 + (i - 1)\sigma)F_{i-1}(\sigma) & \text{otherwise} \end{cases}$$
(B.50)

The inverse of this matrix is also easily defined from $A_n^{(\Gamma)-1}(\sigma)$:

$$A_{i,j}^{(\beta)-1}(\sigma) = A_{i,j}^{(\Gamma)-1}(\sigma)F_j(\sigma)$$
(B.51)

925 Appendix B.6.3. Low cost nested quadrature

A Gauss-Legendre quadrature can be used to approximate integral properties over a Beta EQMOM reconstruction:

$$\int_0^1 f(\xi) n(\xi) \mathrm{d}\xi \approx \frac{1}{2} \sum_{i=1}^P \frac{w_i}{B\left(\alpha_i + 1, \beta_i + 1\right)} \sum_{j=1}^Q \omega_j f\left(\frac{1-\lambda_j}{2}\right) \left(\frac{1-\lambda_j}{2}\right)^{\alpha_i} \left(\frac{1+\lambda_j}{2}\right)^{\beta_i} \quad (B.52)$$

with \boldsymbol{w}_P , $\boldsymbol{\xi}_P$ and σ the EQMOM reconstruction parameters computed from \boldsymbol{m}_{2P} ; $\boldsymbol{\omega}_Q$ and $\boldsymbol{\lambda}_Q$ are the weights and nodes of a Q-nodes Gauss-Legendre quadrature rule (see Appendix C); $\alpha_i = \frac{\xi_i - \sigma}{\sigma}$ and $\beta_i = \frac{1 - \xi_i - \sigma}{\sigma}$. This nested quadrature only requires $\boldsymbol{\omega}_Q$ and $\boldsymbol{\lambda}_Q$ to be computed once, but will not preserve the moments of the distribution.

932 Appendix B.6.4. Moment preserving nested quadrature

A Gauss-Jacobi quadrature will preserve the moments of the distribution:

$$\int_{0}^{1} f(\xi) n(\xi) d\xi \approx 2^{\frac{\sigma-1}{\sigma}} \sum_{i=1}^{P} \frac{w_i}{B(\alpha_i + 1, \beta_i + 1)} \sum_{j=1}^{Q} \omega_j^{(\alpha_i, \beta_i)} f\left(\frac{1 - \lambda_j^{(\alpha_i, \beta_i)}}{2}\right)$$
(B.53)

with \boldsymbol{w}_P , $\boldsymbol{\xi}_P$ and σ the EQMOM reconstruction parameters computed from \boldsymbol{m}_{2P} ; $\boldsymbol{\omega}_Q^{(\alpha_i,\beta_i)}$ and $\boldsymbol{\lambda}_Q^{(\alpha_i,\beta_i)}$ are the weights and nodes of a Q-nodes Gauss-Jacobi quadrature rule of parameters $\boldsymbol{\alpha}_i = \frac{\xi_i - \sigma}{\sigma}$ and $\beta_i = \frac{1 - \xi_i - \sigma}{\sigma}$ (see Appendix C). The moment-preserving property of this quadrature comes with the need to compute $\boldsymbol{\omega}_Q^{(\alpha_i,\beta_i)}$ and $\boldsymbol{\lambda}_Q^{(\alpha_i,\beta_i)}$ for each node of the main Beta EQMOM quadrature.

⁹³⁹ Appendix B.6.5. Single node analytical solution ⁹⁴⁰ The case P = 1 has the following analytical solution:

$$w_{1} = m_{0}$$

$$\xi_{1} = \frac{m_{1}}{m_{0}}$$

$$\sigma = \frac{m_{1}^{2} - m_{0}m_{2}}{m_{0}(m_{2} - m_{1})}$$

941 Appendix C. Gaussian quadratures

A Q-node Gaussian quadrature allows to approximate a function integral as a weighted sum of point wise values of this function over an interval I:

$$\int_{I} f(x)p(x)dx \approx \sum_{j=1}^{Q} \omega_{j}f(\lambda_{j})$$
(C.1)

p(x) is a weight function, and the quadrature rule yields accurate integral evaluations if $f(x) = x^k, k \in \{0, ..., 2Q - 1\}$. The computation of the weights ω_Q and nodes λ_Q is performed as detailed in 2.2 by considering polynomials that are orthogonal with respect to the weight function p(x).

Table C.1 details for each Gauss quadrature:

- the weight function p(x);
- the integration support I;
- the computation of recurrence coefficients a_{Q-1} and b_{Q-1} ;
- the zero-th order moment P_0 of p(x).

The recurrence coefficients are used to construct the Jacobi matrix J_Q associated with p_{54} p(x) on I (see Eq. 6). The nodes λ_Q are the eigenvalues of J_Q , and the weights ω_Q are given by $\omega_j = P_0 v_{1,j}^2$ with $v_{1,j}$ the first component of the normalised eigenvector belonging to the eigenvalue λ_j .

Gauss-	Ι	p(x)	$oldsymbol{a}_Q ext{and} oldsymbol{b}_Q$	P_0
Hermite	$\mathbb R$	$\exp\left(-x^2\right)$	$a_k = 0$	$\sqrt{\pi}$
			$b_k = k/2$	
			Apply Chebyshev algorithm to	
Laplace ^c	\mathbb{R}	$\exp\left(-\left x\right \right)/2$	$\mathbf{p} \text{if } k \text{ odd}$	1
			P_{2Q-1} with $P_k = \begin{cases} k! & \text{if } k \text{ even} \end{cases}$	
Laguerre ^f	\mathbb{R}^+	$x^{\alpha}\exp\left(-x\right)$	$a_0 = 1 + \alpha$	$\Gamma (1 + \alpha)^{\mathrm{d}}$
			$a_k = 2 + a_{k-1}$	
			$b_k = k(k + \alpha)$	
Wigert ^{a,f}	\mathbb{R}^+	$\frac{1}{\gamma x \sqrt{2\pi}} \exp\left(\frac{\log^2(x)}{2\gamma^2}\right)$	$a_k = \left(\left(z^2 + 1 \right) z^{2k} - 1 \right) z^{2k-1}$	1
			$b_k = (z^{2k} - 1) z^{6k-4}$	
			$z = \exp(\gamma^2/2)$	
Weibull ^{c,f}	T₽+	$\gamma x^{\gamma-1} \exp\left(-x^{\gamma}\right)$	Apply Chebyshev algorithm to	1
weibuli	11/2	$fx = \exp(-x)$	P_{2Q-1} with $P_k = \Gamma \left(1 + k/\gamma\right)$	1
Legendre ^b]-1,1[1	$a_k = 0$	2
			$h_{\rm r} = \frac{k^2}{2}$	
			$\frac{0k}{2} - \frac{4k^2 - 1}{4k^2 - 1}$	201811
Jacobi ^{b,f}]-1,1[$(1-x)^{\alpha}(1+x)^{\beta}$	$a_k = \frac{\beta^2 - \alpha^2}{\beta_k (\beta_k - \alpha_k)}$	$2^{\alpha+\rho+1}\times$
	J , L		$\delta_k(\delta_k+2)$	$B(\alpha+1,\beta+1)^{e}$
			$b_{k} = \frac{4k(k+\alpha)(k+\beta)(k+\alpha+\beta)}{4k(k+\alpha)(k+\alpha+\beta)}$	
			$\delta_k^2(\delta_k^2-1)$	
			$\delta_k = 2k + \alpha + \beta$	

Table C.1: Specifics of Gauss quadratures used for EQMOM nested quadratures.

^aWilck [36]. ^bShen et al. [37]. ^c Not standard Gauss-quadrature. ^d $\Gamma(x) = \int_0^{+\infty} t^{x-1} e^{-t} dt$. ^e $B(x,y) = \frac{\Gamma(x)\Gamma(y)}{\Gamma(x+y)}$. ^f $\alpha > -1$, $\beta > -1$, $\gamma > 0$.