



HAL
open science

Does pore scale biogeography exist in different soil types ?

Claire Chenu, Valerie Pouteau, Naoise Nunan

► **To cite this version:**

Claire Chenu, Valerie Pouteau, Naoise Nunan. Does pore scale biogeography exist in different soil types?. EGU 2018, European Geosciences Union General Assembly 2018, European Geosciences Union, Apr 2018, Vienne, Austria. hal-01765428

HAL Id: hal-01765428

<https://hal.science/hal-01765428>

Submitted on 4 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



Does pore scale biogeography exist in different soil types ?

Claire Chenu (1), Valérie Pouteau (1), and Naoise Nunan (2)

(1) INRA, AgroParisTech, Université Paris Saclay, UMR Ecosys, Grignon, France (claire.chenu@inra.fr), (2) CNRS, UPMC, IRD, UMR IEES, Paris, France

Microbial activity is largely controlled by the abiotic conditions prevailing in their habitats, which are very heterogeneous at the microscopic scale. A few studies have demonstrated a microbial biogeography at the pore scale. Different regions in the soil pore network can be considered to be different microbial habitats and these different regions have been shown to be well correlated with organic carbon mineralization rates. We aimed to test whether such a functional biogeography exists for different soil types and whether it is consistent across soils.

We selected six topsoils with contrasted texture, soil organic matter content and pH (2 cambisols, 3 luvisols and 1 podzol under different managements). We added a ^{13}C labelled, easily mineralisable organic substrate, pyruvate, to soil samples previously equilibrated at different matric potentials, in order to place the substrate preferentially in soil pores with neck diameters of 3 to 10 μm or 30 to 100 μm , according to the Jurin-Laplace law. The soil samples were then incubated at pF 1.5 for 3 weeks and CO_2 and $^{13}\text{C}\text{-CO}_2$ were monitored. At the end of incubation, total and $^{13}\text{C}\text{-PLFA}$ were extracted and analysed.

Basal mineralisation, expressed as % total organic C was affected by soil type, mainly related soil pH and the quality of the organic matter. The 6 soils exhibited contrasted microbial community composition, as shown by their PLFA profiles. The addition of pyruvate did not induce any priming effect in soils, except in the long term bare fallow soil, where the mineralization of SOM was presumably limited by energy. In the long term bare fallow soil, pyruvate mineralization was the same whatever the region it was placed in, suggesting other controls of its mineralization than the characteristics of pore scale habitats. In four soils out of six, the mineralization of pyruvate was more rapid when it was initially placed in large pores (30 to 100 μm) than in small pores (3 to 10 μm), suggesting that pore scale biogeography may be a general feature in soils and that coarser pores are more favourable habitats for soil organic matter mineralization.