



HAL
open science

Asymptotic expansions for the conductivity problem with nearly touching inclusions with corner

Virginie Bonnaillie-Noël, Clair Poignard, Grégory Vial

► **To cite this version:**

Virginie Bonnaillie-Noël, Clair Poignard, Grégory Vial. Asymptotic expansions for the conductivity problem with nearly touching inclusions with corner. 2018. hal-01766052v1

HAL Id: hal-01766052

<https://hal.science/hal-01766052v1>

Preprint submitted on 13 Apr 2018 (v1), last revised 28 Jun 2019 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Asymptotic expansions for the conductivity problem with nearly touching inclusions with corner

V. Bonnaillie-Noël*, C. Poinard† and G. Vial‡

April 13, 2018

Abstract

We investigate the case of a medium with two inclusions or inhomogeneities with nearly touching corner singularities. We present two different asymptotic models to describe the phenomenon under specific geometrical assumptions. These asymptotic expansions are analysed and compared in a common framework. We conclude by a representation formula to characterise the detachment of the corners and we provide the possible extensions of the geometrical hypotheses.

Keywords Asymptotic expansion, corner singularity

2010 MSC 35B25, 35C20

1 Introduction

The aim of the paper is to investigate the asymptotic behavior of the solution to the conductivity (also called thermal) problem in a domain with two nearly touching inclusions with corner singularity. It is well-known that behavior of the electrostatic field near a corner of an inclusion depends on the angle of the corner, see [8] for instance. For acute angles – from the domain of computation – the electric field remains bounded but as soon as the angle is larger than π , the electric field blows-up.

Problems with inclusions have given rise to an important literature, in various contexts, mainly motivated by mechanical and electrical engineering, or image processing. In the special case of smooth inclusions close to each other, or close to the boundary of the domain, we can mention for example the works [1, 3, 4, 12]. For nonsmooth inclusions, we can refer to [11, 13].

The configuration we are looking at here is quite trickier since we consider a domain with two inclusions with corner singularity at distance $\delta \ll 1$ from each other. Depending on the geometrical configuration, the corner asymptotics of the solution to the conductivity problem at $\delta \neq 0$ may be dramatically different from the asymptotics of the solution at $\delta = 0$. In this paper, we provide a common framework for two different methods to derive the asymptotic expansions of the solution at $\delta \neq 0$. These expansions are mainly based upon the solution to the limit problem $\delta = 0$ and the appropriate so-called profile terms. These profiles are solution to the conductivity

*Département de mathématiques et applications, Ecole normale supérieure, CNRS, PSL Research University, 45 rue d’Ulm, 75005 Paris, France, BONNAILLIE@math.cnrs.fr

†Institut de Mathématiques de Bordeaux, INRIA, CNRS, Université de Bordeaux, 351, cours de la Libération, 33405 Talence, France, clair.poinard@inria.fr

‡Univ Lyon, Ecole Centrale de Lyon, CNRS UMR 5208, Institut Camille Jordan, F-69134 Ecully, France, gregory.vial@ec-lyon.fr

problem in a sectorial domain with appropriate source terms. They appear naturally from the expansion, in the same vein as in [5].

Precisely, we are interested in the model problem

$$\begin{cases} -\Delta u_\delta = 0 & \text{in } \Pi_\delta, \\ \partial_\nu u_\delta = g & \text{on } \partial\Pi_\delta, \\ u_\delta \rightarrow 0 & \text{at infinity,} \end{cases} \quad (1)$$

where g is a given datum (trace of a smooth function), and the unbounded domain Π_δ is defined by

$$\Pi_\delta = \mathbb{R}^2 \setminus \overline{\Omega_\delta^L \cup \Omega_\delta^R},$$

the domains Ω_δ^L and Ω_δ^R standing for the inclusions. The distance between Ω_δ^L and Ω_δ^R is 2δ , see Figure 1.




Figure 1: The domain Π_δ with the inclusions $\Omega_\delta^L, \Omega_\delta^R$, and the corresponding limit domain Π_0 .

Naturally, the formal limit case where the inclusions touch each other corresponds to

$$\begin{cases} -\Delta u_0 = 0 & \text{in } \Pi_0, \\ \partial_\nu u_0 = g & \text{on } \partial\Pi_0, \\ u_0 \rightarrow 0 & \text{at infinity.} \end{cases} \quad (2)$$

Nevertheless, the asymptotic model is not unique since the dependence of the domains $\Omega_\delta^L, \Omega_\delta^R$ with respect to the parameter δ is not explicitly given by the physical situation. We will investigate the following two choices:

- **Translation method.** The size of the inclusions Ω_δ^L and Ω_δ^R is supposed to be independent of δ . Changing the value of the parameter δ merely corresponds to a translation of the inclusions along a given unit vector \mathbf{d} . The subsequent asymptotic expansion is detailed in Section 3.1 where the domain Π_δ will be denoted by $\dot{\Pi}_\delta$ for notation consistency.
- **Contraction method.** The inclusions do not have a constant size with respect to δ , but are contractions of the limit case (i.e. $\delta = 0$). This case is precisely addressed in Section 3.2. In particular, the geometrical setting will be made clear in Subsection 3.2.1, where, here again, the domain Π_δ will be denoted $\dot{\Pi}_\delta^*$.

If the two presented asymptotic models coincide for $\delta = 0$, both have pros and cons since they can handle different geometrical frameworks of inclusions. For comparison purposes, which yield one of the main goal of this paper, we restrict the geometrical configurations to a situation where both models might be considered. The precise framework is detailed in the following assumptions, but the reader can keep in mind the situation of Figure 1.

Assumption 1.1 *The geometrical hypotheses we make are the following*

(H1) *For $\delta = 0$, the inclusions touch each other at one point, where we put the origin $\mathbf{0}$.*

(H2) *The inclusions Ω_0^L and Ω_0^R are smooth domains, except at point $\mathbf{0}$, where they coincide with plane sectors.*

(H3) *There exists a direction \mathbf{d} such that*

$$\exists \rho_0 > 0 \quad \{\mathbf{0} + \rho \mathbf{d} ; 0 < |\rho| < \rho_0\} \subset \Omega_0^L \cup \Omega_0^R.$$

By convention, we fix the horizontal axis along this direction. Moreover, Ω_0^L and Ω_0^R are supposed to be included into $\mathbb{R}_- \times \mathbb{R}$, and $\mathbb{R}_+ \times \mathbb{R}$, respectively.

Remark 1.2 *Replacing in Assumption (H1) one point by a finite number of points generates only technical difficulties. We do not address the case of inclusions without corners (for example two disks), which generates a cuspidal limit domain. Hypothesis (H2) prevents from this situation. Besides, the regularity of the boundaries far from the origin is not necessary and is only assumed here for simplicity. Likewise, hypothesis (H3) is technical and might be weakened. The relaxation of these hypotheses will be discussed in Section 5.*

Section 2 recalls the decomposition in regular and singular parts of the Laplacian equation (2) with adding a source term. Then we consider the two asymptotic models. First in Section 3, we present a formal asymptotic expansion where we do not take care of the corner singularities. Even though formal, this construction is useful for the complete analysis and enables to deal with difficulties step by step. We present asymptotic model for the translation and contraction case and propose a unified formulation in Section 3.3. In Section 4, we take the singularities into account, we construct the corner profiles and we give the complete asymptotic expansions obtained by the two methods.

As far as we know, the comparison of these two asymptotic methods have not been addressed before. In this paper the two methods are studied in parallel and presented in a unified formulation which allows us to deduce a generic full asymptotic expansion of the solution of (1).

For $\delta > 0$, the solution u_δ of problem (1) and the limit solution u_0 have both singularities at the origin, but with different singular exponents because they are associated with different angles. In the geometry of Figure 1, u_δ has singularities due to the re-entrant corners whereas u_0 has only singularities of the convex corners (of angle α^\pm , see Figure 1). The asymptotic procedures that will be described in the paper will allow to compare u_δ and u_0 . Precisely, the first terms in the asymptotic expansion read

$$u_\delta(\mathbf{x}) = \psi\left(\frac{\mathbf{x}}{\delta}\right)u_0(\mathbf{x}) - \chi(\mathbf{x}) \left(g(\mathbf{0})\mathfrak{M}\left(\frac{\mathbf{x}}{\delta}\right) + \sum_{\pm} c_{0,0}^\pm \mathfrak{R}_0^\pm \frac{\mathbf{x}}{\delta} \right), \quad (3)$$

where

- The functions ψ and χ are smooth radial cutoff functions satisfying

$$\psi(\mathbf{x}) = 0 \text{ and } \chi(\mathbf{x}) = 1, \text{ near } \mathbf{0}.$$

- The coefficients $c_{0,0}^\pm$ are the first singular coefficients of u_0 , solution to (2).
- The profiles \mathfrak{R}_0^\pm and \mathfrak{M} are defined in the infinite domain \mathfrak{F} – see Figure 4 – as the respective solutions to Problems (30)–(31).

As we can see in (3), the singularities of u_0 are canceled through the cutoff function ψ (at distance $\mathcal{O}(\delta)$ from the origin), and replaced by adapted counterparts through the profiles terms, rescaled to fit the δ -depending domain Π_δ . It is worth noting whatever the method, the zeroth-order terms of the expansion of u_δ are given by (3). The differences in the expansions will appear at the next order terms. The goal of the paper is to push forward the expansion for each method to provide different ways to approximate u_δ .

2 Corner asymptotics for the limit problem

Let us consider now the solution u to the generic problem:

$$\begin{cases} -\Delta u = f & \text{in } \Pi_0, \\ \partial_\nu u = h & \text{on } \partial\Pi_0, \\ u \rightarrow 0 & \text{at infinity,} \end{cases} \quad (4)$$

where f and h are smooth given data (for $f = 0$ and $h = g$ the solution u is nothing but u_0).

Since corners appear in the limit domain Π_0 , singularities arise for the solution u near the origin, preventing a full elliptic regularity. Precisely, if α^\pm denote the absolute value of the sector angles involved in Π_0 – see Figure 1 one has:

$$\alpha^\pm = \theta_2^\pm - \theta_1^\pm.$$

Notice that according to (H3), $\alpha^\pm \in (0, \pi)$.

Assumption 2.1 *Let us add a geometrical hypothesis on α^\pm*

(H4) $\alpha^\pm \notin \pi\mathbb{Q}$.

With Assumption (H4), the following decomposition (5) is valid. Otherwise, logarithmic singularities may appear in the corner asymptotics. We do not consider this case here for simplicity.

The potential u admits a splitting into regular and singular parts for any integer p :

$$u(\mathbf{x}) = u_{\text{flat}}^p(\mathbf{x}) + \chi(\mathbf{x}) \sum_{\pm} \sum_{\lambda \in \Lambda_p^\pm} c_\lambda^\pm \mathfrak{s}_\lambda^\pm(\mathbf{x}) + \chi(\mathbf{x}) \sum_{\pm} \sum_{\sigma=1,2} \sum_{1 \leq n \leq p-1} a_n^{\pm,\sigma} \mathfrak{a}_n^{\pm,\sigma}(\mathbf{x}), \quad (5)$$

where, using polar coordinates (r, θ) ,

- $u_{\text{flat}}^p \in H_{\text{loc}}^p(\Pi_0)$, and for any multi-index β with $|\beta| \leq p$, we have

$$\partial^\beta u_{\text{flat}}^p(\mathbf{x}) = \mathcal{O}(r^{p-|\beta|}), \quad \text{near the corner } \mathbf{0}, \quad (6)$$

- $\chi : \mathbb{R}^2 \rightarrow \mathbb{R}_+$ is a smooth radial cutoff function, with compact support and equal to 1 near $\mathbf{0}$,
- we denote by

$$\Lambda_p^\pm = \left\{ \lambda = \frac{q\pi}{\alpha^\pm} ; 0 \leq \frac{q\pi}{\alpha^\pm} \leq p \text{ with } q \in \mathbb{N} \right\},$$

- the coefficients $c_\lambda^\pm \in \mathbb{R}$ depend on f and h and are called the *singular coefficients* of order λ of u in the upper or lower part of Π_0 . They can be numerically computed from the solution u via extraction formulae (see [8, Chap. 8]),

- the *singular functions* \mathfrak{s}_λ^\pm are defined for any $\lambda \in \Lambda_p^\pm$ by

$$\mathfrak{s}_\lambda^\pm(\mathbf{x}) = \begin{cases} r^\lambda \cos(\lambda(\theta - \theta_1^\pm)) & \text{if } \theta_1^\pm < \theta < \theta_2^\pm, \\ 0 & \text{otherwise,} \end{cases}$$

see Figure 1 for the definition of θ_1^\pm and θ_2^\pm . Note that for $\lambda = 0$, the functions \mathfrak{s}_0^\pm equal 1 for $\theta_1^\pm < \theta < \theta_2^\pm$.

- The last series in (5) comes from the data f and h , which do not necessarily vanish near the corner. The coefficients $a_n^{\pm, \sigma}$ are explicitly determined by the Taylor expansions of f and h . The functions $\mathfrak{a}_n^{\pm, \sigma}$ are defined for any positive integer n and any $\sigma \in \{1, 2\}$ by

$$\mathfrak{a}_n^{\pm, \sigma}(\mathbf{x}) = \begin{cases} r^n \cos(n(\theta - \theta_\sigma^\pm)) & \text{if } \theta_1^\pm < \theta < \theta_2^\pm, \\ 0 & \text{otherwise.} \end{cases}$$

Such a decomposition is standard, and described in more details in the theory of corner domains, see [6, 8, 9] for example.

Before dealing with singularities, it is important to derive the asymptotic expansion in the ideal case where no singularity appears at any order. Actually, understanding how such a derivation occurs will facilitate the accurate asymptotic expansion with the singularities.

3 Asymptotic models without singularities

The goal of the section is to present the formal derivation of the asymptotic expansions for both translation and contraction cases, without accounting for the influence of corner singularities. We emphasize that throughout this section, we assume that no singularity appears in the derivation of the asymptotics, meaning that all the coefficients c_λ^\pm and $a_n^{\pm, \sigma}$ of any expansion of type (5) equal zero. Therefore the following expansion only involves coefficients with entire powers of δ .

This ideal case has no chance to hold in general. The singularities will make appear non entire powers of δ , and they will lead to a modification of the source terms of the following elementary problems, but the procedure will remain unchanged. Therefore even though very specific, this ideal framework is interesting for understanding the way the cascade of elementary problems is derived at any order to get the asymptotics. We derive such “ideal” asymptotics, in the two different geometrical frameworks. In addition, we unify both approach but introducing appropriate notations, making thus possible a comparison of the two methods. The difficulties arising from the corner singularities are postponed to Section 4 where the corner profiles are introduced.

3.1 Asymptotic model with translation of inclusions

This section is devoted to the construction of the asymptotic expansion for Problem (1) in the case where the inclusions are constant patterns translated from each other by a distance of 2δ . As mentioned previously, to distinguish the two geometrical frameworks, the notation will be enriched by a \bullet for the translation case.

3.1.1 Geometrical setting

We assume here that the domains $\dot{\Omega}_\delta^L$ and $\dot{\Omega}_\delta^R$ are translations of constant size inclusions :

$$\dot{\Omega}_\delta^L = \dot{\Omega}_0^L - \delta \mathbf{d} \quad \text{and} \quad \dot{\Omega}_\delta^R = \dot{\Omega}_0^R + \delta \mathbf{d}.$$

We recall that

$$\dot{\Pi}_\delta = \Pi_\delta = \mathbb{R}^2 \setminus \overline{\dot{\Omega}_\delta^L \cup \dot{\Omega}_\delta^R}.$$

It is classical to transform the problem into a domain independent of δ in order to make the parameter δ appear in the equations. In the present case, the ‘gap’ between the inclusions can be seen as a thin layer. It is then natural to set the change of variables

$$\dot{\Phi}_\delta : \mathbf{x} = (x_1, x_2) \mapsto \boldsymbol{\xi} = (\eta, \tau),$$

defined as

$$\boldsymbol{\xi} = \dot{\Phi}_\delta(\mathbf{x}) = \begin{cases} (x_1 + \delta - 1, x_2) & \text{if } x_1 \leq -\delta, \\ (\eta, \tau) = \left(\frac{x_1}{\delta}, x_2\right) & \text{if } |x_1| \leq \delta, \\ (x_1 - \delta + 1, x_2) & \text{if } x_1 \geq \delta. \end{cases} \quad (7)$$

Through this change of variables, the domain $\dot{\Pi}_\delta$ is transformed into $\dot{\Pi}$, see Figure 2. Here, we obviously have $\dot{\Pi} = \dot{\Pi}_1$. We set $\dot{\Pi}_\delta^{\text{lay}} = \{|x_1| < \delta\}$, and $\dot{\Pi}_\delta^{\text{ext}} = \dot{\Pi}_\delta \cap \{|x_1| > \delta\}$.




Figure 2: The domain $\dot{\Pi}_\delta$ and the domain $\dot{\Pi}$ obtained after change of variables $\dot{\Phi}_\delta$ given by (7).

For any function $v \in L_{\text{loc}}^2(\dot{\Pi}_\delta)$, we denote by \dot{v}^{ext} and \dot{v}^{lay} the restrictions of the function $v \circ \dot{\Phi}_\delta^{-1}$ to $\dot{\Pi}^{\text{ext}} = \dot{\Pi}_1^{\text{ext}}$ (the exterior part), and $\dot{\Pi}^{\text{lay}} = \dot{\Pi}_1^{\text{lay}}$ (the layer), respectively.

3.1.2 Elementary problems for the translation case

We are now able to set the transformed problem in a domain independent of the parameter δ . Expressed in terms of the unknowns $\dot{\mathbf{u}}^{\text{ext},\delta}$ and $\dot{\mathbf{u}}^{\text{lay},\delta}$, it reads

$$\left\{ \begin{array}{ll} \partial_\eta^2 \dot{\mathbf{u}}^{\text{lay},\delta} = -\delta^2 \partial_\tau^2 \dot{\mathbf{u}}^{\text{lay},\delta} & \text{in } \dot{\Pi}^{\text{lay}}, \\ \dot{\mathbf{u}}^{\text{lay},\delta} = \dot{\mathbf{u}}^{\text{ext},\delta} & \text{on } \dot{\Gamma}^{\text{trans}}, \\ -\Delta \dot{\mathbf{u}}^{\text{ext},\delta} = 0 & \text{in } \dot{\Pi}^{\text{ext}}, \\ \partial_\nu \dot{\mathbf{u}}^{\text{ext},\delta} = \frac{1}{\delta} \partial_\eta \dot{\mathbf{u}}^{\text{lay},\delta} & \text{on } \dot{\Gamma}^{\text{trans}}, \\ \partial_\nu \dot{\mathbf{u}}^{\text{ext},\delta} = g \circ \dot{\Phi}_\delta^{-1}(\boldsymbol{\xi}) & \text{on } \partial \dot{\Pi}^{\text{ext}} \setminus \dot{\Gamma}^{\text{trans}}, \\ \dot{\mathbf{u}}^{\text{ext},\delta} \rightarrow 0 & \text{at infinity.} \end{array} \right. \quad (8)$$

We now replace $g \circ \dot{\Phi}_\delta^{-1}(\boldsymbol{\xi})$ by its Taylor expansion:

$$g \circ \dot{\Phi}_\delta^{-1}(\boldsymbol{\xi}) = \sum_{n=0}^N \dot{\mathbf{g}}_n(\boldsymbol{\xi}) \delta^n + o(\delta^N). \quad (9)$$

Considering Problem (8), we postulate the two following Ansätze

$$\dot{\mathbf{u}}^{\text{ext},\delta} = \sum_{n \geq 0} \delta^n \dot{\mathbf{u}}_n^{\text{ext}}, \quad \dot{\mathbf{u}}^{\text{lay},\delta} = \sum_{n \geq 0} \delta^n \dot{\mathbf{u}}_n^{\text{lay}}.$$

Note that the series are written in the sense of asymptotic expansions (*i.e.* truncate the series and make $\delta \rightarrow 0$, see [10] for example), we do not expect – nor need – convergence as $n \rightarrow \infty$. Inserting these *Ansätze* into Problem (8), we get the following cascades of elementary problems:

$$\left\{ \begin{array}{ll} \partial_\eta^2 \dot{\mathbf{u}}_{n+1}^{\text{lay}} = -\partial_\tau^2 \dot{\mathbf{u}}_{n-1}^{\text{lay}} & \text{in } \dot{\Pi}^{\text{lay}}, \\ \dot{\mathbf{u}}_{n+1}^{\text{lay}} = \dot{\mathbf{u}}_{n+1}^{\text{ext}} & \text{on } \dot{\Gamma}^{\text{trans}}, \\ -\Delta \dot{\mathbf{u}}_n^{\text{ext}} = 0 & \text{in } \dot{\Pi}^{\text{ext}}, \\ \partial_\nu \dot{\mathbf{u}}_n^{\text{ext}} = \partial_\eta \dot{\mathbf{u}}_{n+1}^{\text{lay}} & \text{on } \dot{\Gamma}^{\text{trans}}, \\ \partial_\nu \dot{\mathbf{u}}_n^{\text{ext}} = \dot{\mathbf{g}}_n & \text{on } \partial \dot{\Pi}^{\text{ext}} \setminus \dot{\Gamma}^{\text{trans}}, \\ \dot{\mathbf{u}}_n^{\text{ext}} \rightarrow 0 & \text{at infinity,} \end{array} \right. \quad (10)$$

with the convention $\dot{\mathbf{u}}_\ell^{\text{ext}} = 0$ and $\dot{\mathbf{u}}_\ell^{\text{lay}} = 0$ for $\ell < 0$. In the first two equations of the problem, solved by $\dot{\mathbf{u}}_n^{\text{lay}}$, the variable τ is actually only a parameter. The question of the solvability of this sequence of problems will be discussed in section 3.3.

3.2 Asymptotic model with contraction of inclusions

As mentioned previously, the way each asymptotic expansion is derived is somehow arbitrary, and it depends mostly of the point of view of the modeling. In Section 3.1, we considered the case of two inclusions with δ -independent size, whose corners were distant from 2δ .

In this section we adopt another point of view, leading to the same limit geometry: we assume that the inclusions are contractions of the limit inclusion $\Omega_0^L \cup \Omega_0^R$ given by Figure 1. Let us first make precise the geometrical assumptions. As mentioned previously, to distinguish the two geometrical frameworks, the notation is enriched by a \star for the contraction case.

3.2.1 Geometrical setting

Identification of the thin layer For δ small enough, we define here $\dot{\Omega}_\delta^R$ and $\dot{\Omega}_\delta^L$ as contractions in the normal direction ν of the respective limit inclusions $\dot{\Omega}_0^L$ and $\dot{\Omega}_0^R$. A thin layer $\dot{\Pi}_\delta^{\text{lay}}$ appears naturally, when passing from the domain $\dot{\Pi}_\delta$ to the limit domain $\dot{\Pi}_0 := \Pi_0$ (see Fig. 3).

We denote respectively by $\dot{\Pi}^{\text{ext}}$, $\dot{\Pi}_\delta^{\text{lay}}$ and $\dot{\Pi}_\delta$ the 3 domains (see Figure 3)

$$\dot{\Pi}^{\text{ext}} = \Pi_0, \quad \dot{\Pi}_\delta^{\text{lay}} = \left(\dot{\Omega}_0^L \cup \dot{\Omega}_0^R \right) \setminus \overline{\left(\dot{\Omega}_\delta^L \cup \dot{\Omega}_\delta^R \right)}, \quad \dot{\Pi}_\delta = \overline{\dot{\Pi}^{\text{ext}} \cup \dot{\Pi}_\delta^{\text{lay}}}.$$




Figure 3: The domain Π_δ^* with the inclusions Ω_δ^{LR} (non symmetric and symmetric cases).

Let us introduce a suitable parameterization of $\partial\Pi_\delta^*$: Inside the ball $\mathcal{B}_{\rho_*} = \{|\mathbf{x}| < \rho_*\}$, the boundary $\partial\Pi_\delta^*$ is deduced by translation by $\pm\delta\mathbf{d}$ from the right and left parts of $\partial\Pi_0^*$.

$$\left(\partial\Pi_\delta^*\right) \cap \mathcal{B}_{\rho_*} = \left(\left\{\mathbf{x} - \delta\mathbf{d} ; \mathbf{x} \in \partial\Omega_0^{L*}\right\} \cup \left\{\mathbf{x} + \delta\mathbf{d} ; \mathbf{x} \in \partial\Omega_0^{R*}\right\}\right) \cap \mathcal{B}_{\rho_*}.$$

Outside the ball $\mathcal{B}_{\frac{\rho_*}{2}} = \{|\mathbf{x}| < \frac{\rho_*}{2}\}$, we choose a smooth, positive and δ -independent function f to parameterize $\partial\Pi_\delta^*$:

$$\partial\Pi_\delta^* \setminus \mathcal{B}_{\frac{\rho_*}{2}} = \left\{\mathbf{x} + \delta f(\mathbf{x})\nu(\mathbf{x}) ; \mathbf{x} \in \partial\Pi_0^*\right\} \setminus \mathcal{B}_{\frac{\rho_*}{2}},$$

designed in such a way the two definitions are consistent in the intersection $\{\frac{\rho_*}{2} \leq |\mathbf{x}| \leq \rho_*\}$.

Remark 3.1 (Arbitrary choice for f near $\mathbf{0}$) *The choice of ρ_* is somehow arbitrary, but it does not change dramatically the derivation of the expansion, since the function u_0 is supposed to be flat near the origin $\mathbf{0}$, as stated by (5)–(6).*

Remark 3.2 (The case of constant thickness) *In the simple geometrical framework where the plane sectors are equally distributed from both side of the horizontal axis $(\mathbf{0}, \mathbf{d})$, the simplest choice for f is the constant function defined by*

$$\forall \mathbf{x} \in \partial\Pi_\delta^*, \quad f(\mathbf{x}) = \sin\left(\frac{\omega}{2}\right), \quad \text{where } \omega = \theta_1^+ \text{ (and } \theta_2^+ = \pi - \omega, \theta_1^- = \pi + \omega, \theta_2^- = 2\pi - \omega).$$

Laplace operator in local coordinates Once the identification and the parameterization of Π_δ^{lay} is performed, it is natural to introduce new variables (η, τ) , which respectively correspond to the normal and tangential variables along $\partial\Pi_0$, in order to rewrite the Laplace operator in these new coordinates. We parameterize $\partial\Pi_0$ by the vector field Ψ defined on the torus \mathbb{T} :

$$\partial\Pi_0 = \{\Psi(\tau), \quad \tau \in \mathbb{T}\}.$$

By abuse of notation, we still denote by f and ν the functions $f \circ \Psi$ and $\nu \circ \Psi$ defined on \mathbb{T} . We then define the mapping Φ_δ^* from $(0, 1) \times \mathbb{T}$ into \mathbb{R}^2 as

$$\xi = \Phi_\delta^*(\eta, \tau) = \Psi(\tau) + \delta f(\tau)\eta\nu(\tau), \quad \forall (\eta, \tau) \in (0, 1) \times \mathbb{T}, \quad (11)$$

such that

$$\Pi_\delta^{\text{lay}} \setminus \mathcal{B}_{\frac{\rho_*}{2}} = \Phi_\delta^*((0, 1) \times \mathbb{T}) \setminus \mathcal{B}_{\frac{\rho_*}{2}}.$$

Note that $\dot{\Phi}_\delta^*$ applies only to the layer part (contrary to $\dot{\Phi}_\delta$ which is defined in \mathbb{R}^2). In local coordinates (η, τ) , the Euclidean metric $(G_{ij})_{i,j \in \{1, \dots, 2\}}$ is given by

$$G_{11} = |\partial_\eta \dot{\Phi}_\delta^*|^2, \quad G_{22} = |\partial_\tau \dot{\Phi}_\delta^*|^2, \quad G_{12} = G_{21} = \langle \partial_\eta \dot{\Phi}_\delta^*, \partial_\tau \dot{\Phi}_\delta^* \rangle.$$

Denoting by $|G|$ its determinant:

$$|G| = \det(G_{ij}) = G_{11}G_{22} - G_{12}^2,$$

Laplace's operator reads then

$$\Delta = \frac{1}{\sqrt{|G|}} \partial_\eta \left(\frac{1}{\sqrt{|G|}} (G_{22} \partial_\eta - G_{12} \partial_\tau) \right) + \frac{1}{\sqrt{|G|}} \partial_\tau \left(\frac{1}{\sqrt{|G|}} (-G_{12} \partial_\eta + G_{11} \partial_\tau) \right)$$

and thus it can be expanded as

$$\begin{aligned} \Delta &= \frac{1}{\delta^2 f^2} \left\{ \partial_\eta^2 + \delta f \kappa \partial_\eta \right. \\ &\quad \left. + \delta^2 \left(\eta^2 (f')^2 \partial_\eta^2 + \left(2 (f')^2 - f f'' - \eta \kappa^2 f^2 \right) \partial_\eta - 2 \eta f f' \partial_\eta \partial_\tau + f \partial_\tau^2 \right) + \dots \right\} \\ &= \frac{1}{\delta^2 f^2} \left(\partial_\eta^2 + \sum_{k \geq 1} \delta^k \dot{\mathbb{T}}_k \right), \end{aligned} \quad (12)$$

where $\dot{\mathbb{T}}_k$ are differential operators independent of δ and κ denotes the curvature (extended by 0 at the origin). On the other hand, the normal derivative is rescaled by a factor $1/(\delta f)$:

$$\partial_\nu|_{\eta=0} = \frac{1}{\delta f} \partial_\eta, \quad (13a)$$

$$\partial_\nu|_{\eta=1} = \frac{1}{\delta f} \left(\partial_\eta + \sum_{k \geq 1} \delta^k \dot{\mathbb{B}}_k \right), \quad (13b)$$

where $\dot{\mathbb{B}}_k$ are differential operators independent of δ .

3.2.2 Elementary problems for the contraction case

By analogy with the translation case, we denote by $\dot{\Gamma}^{\text{trans}}$ the interface across which transmissions occur. In the contraction case, $\dot{\Gamma}^{\text{trans}}$ is nothing but $\partial \Pi_0$:

$$\dot{\Gamma}^{\text{trans}} = \partial \Pi_0 = \partial \dot{\Pi}_{\text{ext}}.$$

We denote respectively by $\dot{u}^{\text{ext}, \delta}$ and $\dot{u}^{\text{lay}, \delta}$ the function $u \circ \dot{\Phi}_\delta^{-1}$ in $\dot{\Pi}^{\text{ext}}$ and $\dot{\Pi}^{\text{lay}}$. After this change of variables, the elementary problems read

$$\left\{ \begin{array}{ll} \partial_\eta^2 \dot{u}^{\text{lay}, \delta} = - \sum_{k \geq 1} \delta^k \dot{\mathbb{T}}_k(\dot{u}^{\text{lay}, \delta}) & \text{in } \dot{\Pi}^{\text{lay}}, \\ \dot{u}^{\text{lay}, \delta} = \dot{u}^{\text{ext}, \delta} & \text{on } \dot{\Gamma}^{\text{trans}}, \\ \frac{1}{\delta f} \left(\partial_\eta + \sum_{k \geq 1} \delta^k \dot{\mathbb{B}}_k \right) \dot{u}^{\text{lay}, \delta} = \sum_{n \geq 0} \delta^n \dot{\mathbb{G}}_n & \eta = 1, \\ -\Delta \dot{u}^{\text{ext}, \delta} = 0 & \text{in } \dot{\Pi}^{\text{ext}}, \\ \partial_\nu \dot{u}^{\text{ext}, \delta} = \frac{1}{\delta f} \partial_\eta \dot{u}^{\text{lay}, \delta} & \text{on } \dot{\Gamma}^{\text{trans}}, \\ \dot{u}^{\text{ext}, \delta} \rightarrow 0 & \text{at infinity,} \end{array} \right. \quad (14)$$

where $\xi \mapsto \mathring{g}_n(\xi)$ are obtained from the Taylor expansion of $g \circ \Phi_\delta^{-1}(\xi)$. Polynomial Ansätze similar to the translation case,

$$\mathring{u}^{\text{ext},\delta} = \sum_{n \geq 0} \delta^n \mathring{u}_n^{\text{ext}}, \quad \mathring{u}^{\text{lay},\delta} = \sum_{n \geq 0} \delta^n \mathring{u}_n^{\text{lay}},$$

lead to the following sequence of elementary problems:

$$\left\{ \begin{array}{ll} \partial_\eta^2 \mathring{u}_{n+1}^{\text{lay}} = - \sum_{k+\ell=n+1} \mathring{T}_k \mathring{u}_\ell^{\text{lay}} & \text{in } \mathring{\Pi}^{\text{lay}}, \\ \mathring{u}_{n+1}^{\text{lay}} = \mathring{u}_{n+1}^{\text{ext}} & \text{for } \eta = 0, \\ \partial_\eta \mathring{u}_{n+1}^{\text{lay}} = \mathring{f} \mathring{g}_n - \sum_{k=1}^n \mathring{B}_k \mathring{u}_{n-k}^{\text{lay}} & \text{for } \eta = 1, \\ -\Delta \mathring{u}_n^{\text{ext}} = 0 & \text{in } \mathring{\Pi}^{\text{ext}}, \\ \partial_\nu \mathring{u}_n^{\text{ext}} = \frac{1}{f} \partial_\eta \mathring{u}_{n+1}^{\text{lay}} & \text{on } \eta = 0, \\ \mathring{u}_n^{\text{ext}} \rightarrow 0 & \text{at infinity,} \end{array} \right. \quad (15)$$

Again, we postpone the discussion of the well-posedness of this sequence of equations to Section 3.3.

3.3 A unified formulation

In Sections 3.1 and 3.2 we have derived an asymptotic expansion for two different asymptotic families converging to the same limit problem. Actually, it is possible to describe a general framework containing the two procedures. Once again, we emphasize that in this section, we are dealing with the ideal cases where the singularities are omitted at any order. Section 4 is devoted to derive the sector profile problems to account for the singularities, that appear in the general case.

Geometrical setting We consider the problems obtained after change of variables (solved by \mathring{u}^δ and \mathring{u}^{δ} , respectively). We denote by Π the domain where the problem is set, it can be split into

$$\Pi = \Pi^{\text{ext}} \cup \Gamma^{\text{trans}} \cup \Pi^{\text{lay}},$$

where $\Gamma^{\text{trans}} = \partial \Pi^{\text{ext}} \cap \partial \Pi^{\text{lay}}$. The layer Π^{lay} has the form $(\eta_-, \eta_+) \times I$. Finally, we set $\Gamma = \partial \Pi \setminus \Gamma^{\text{trans}}$.

- In the translation case, $\eta_\pm = \pm 1$, $I = \mathbb{R}$, and $\Gamma^{\text{trans}} = \partial \Pi^{\text{lay}} = \{-1, 1\} \times I$.
- In the contraction case, $\eta_- = 0$, $\eta_+ = 1$, $I = \partial \Pi_0$, and $\Gamma^{\text{trans}} = \partial \Pi^{\text{ext}} = \{0\} \times I$.

For the sake of conciseness, for any function u defined in Π we naturally denote by u^{ext} and u^{lay} its restrictions to Π^{ext} and Π^{lay} respectively.

Layer problem We consider the following equations

$$\left\{ \begin{array}{ll} \partial_\eta^2 u_{n+1}^{\text{lay}} = j_n & \text{for } \eta \in (\eta_-, \eta_+) \\ u_{n+1}^{\text{lay}} = u_{n+1}^{\text{ext}} & \text{for } \eta = \eta_-, \\ \partial_\eta u_{n+1}^{\text{lay}} = k_n & \text{for } \eta = \eta_+, \end{array} \right. \quad (16)$$

where the term

$$\mathbf{j}_n = - \sum_{k+\ell=n+1, k \geq 1} \mathbb{T}_k \mathbf{u}_\ell^{\text{lay}} \quad (17)$$

is given explicitly by $\mathbf{u}_\ell^{\text{lay}}$ for $0 \leq \ell \leq n$. The expression of \mathbf{k}_n depends on the geometrical framework:

- In the translation case,

$$\mathbf{k}_n = \partial_\nu \mathbf{u}_n^{\text{ext}}|_{\eta=\eta_+}.$$

- In the contraction case,

$$\mathbf{k}_n = f \mathbf{g}_n^* - \sum_{k=1}^n \mathbb{B}_k^* \mathbf{u}_{n-k}^{\text{lay}}.$$

Problem (16) is exactly obtained from (15) in the contraction case whereas the two following equations are missing in the translation case:

$$\begin{cases} \mathbf{u}_{n+1}^{\text{ext}} = \mathbf{u}_{n+1}^{\text{lay}} & \text{for } \eta = \eta_+, \\ \partial_\nu \mathbf{u}_n^{\text{ext}} = \partial_\eta \mathbf{u}_{n+1}^{\text{lay}} & \text{for } \eta = \eta_-. \end{cases} \quad (18)$$

They will be taken into account in (22). In order to solve Problem (16), let us now define the operator \mathbb{R} for any function \mathbf{j} as

$$\mathbb{R}[\mathbf{j}](\eta, \tau) = - \int_\eta^{\eta^+} \mathbf{j}(s, \tau) \, ds. \quad (19)$$

Thanks to (16), one easily obtains the formal expressions

$$\partial_\eta \mathbf{u}_{n+1}^{\text{lay}}(\eta, \tau) = \mathbf{k}_n(\tau) + \mathbb{R}[\mathbf{j}_n](\eta, \tau), \quad (20)$$

$$\mathbf{u}_{n+1}^{\text{lay}}(\eta, \tau) = \mathbf{u}_{n+1}^{\text{ext}}(\eta_-, \tau) + (\eta - \eta_-) \mathbf{k}_n(\tau) + \int_{\eta_-}^{\eta} \mathbb{R}[\mathbf{j}_n](t, \tau) \, dt. \quad (21)$$

Exterior problem In the outer domain Π^{ext} the problem at the order n has the following form:

$$\text{Find } \mathbf{u}_n^{\text{ext}} \text{ such that } \begin{cases} -\Delta \mathbf{u}_n^{\text{ext}} = 0 & \text{in } \Pi^{\text{ext}}, \\ \bar{\gamma}(\mathbf{u}_n^{\text{ext}}) = \phi_n & \text{on } \Gamma^{\text{trans}}, \\ \partial_\nu \mathbf{u}_n^{\text{ext}} = \mathbf{g}_n & \text{on } \partial \Pi^{\text{ext}} \setminus \Gamma^{\text{trans}}, \\ \mathbf{u}_n^{\text{ext}} \rightarrow 0 & \text{at infinity,} \end{cases} \quad (22)$$

where $\bar{\gamma}$, ϕ_n and \mathbf{g}_n are defined below.

- In the translation case, equations (18) together with (20)–(21) make appear the vector operator $\bar{\gamma}$ defined by

$$\bar{\gamma}(\mathbf{u}) = (\mathbf{u}|_{\eta=\eta_+} - \mathbf{u}|_{\eta=\eta_-}, \partial_\nu \mathbf{u}|_{\eta=\eta_+} - \partial_\nu \mathbf{u}|_{\eta=\eta_-}),$$

and the data given by

$$\phi_n(\tau) = (\mathbf{u}_n^{\text{lay}}(\eta^+, \tau) - \mathbf{u}_n^{\text{lay}}(\eta^-, \tau), \partial_\eta \mathbf{u}_{n+1}^{\text{lay}}(\eta^+, \tau) - \partial_\eta \mathbf{u}_{n+1}^{\text{lay}}(\eta^-, \tau)).$$

On the other hand, the Neumann datum \mathbf{g}_n is given by the Taylor expansion of g – see (9) – $\mathbf{g}_n = \dot{\mathbf{g}}_n$.

- In the contraction case $\partial\Pi^{\text{ext}} \setminus \Gamma^{\text{trans}} = \emptyset$ thus no need to define g_n . The operator $\bar{\gamma}$ is scalar and given by

$$\bar{\gamma}(\mathbf{u}) = \partial_\nu \mathbf{u}|_{\Gamma^{\text{trans}}},$$

and the datum ϕ_n is given by

$$\phi_n(\tau) = \frac{1}{f(\tau)} \partial_\eta \mathbf{u}_{n+1}^{\text{lay}}(\eta^-, \tau).$$

Sequential derivation of the terms At order $n = 0$, the solution to Problem (16) is formally given by formulae (20)–(21):

$$\mathbf{u}_0^{\text{lay}}(\eta, \tau) = \mathbf{u}_0^{\text{ext}}(\eta_-, \tau), \quad \partial_\eta \mathbf{u}_1^{\text{lay}}(\eta, \tau) = \mathbf{k}_0(\tau),$$

where $\mathbf{k}_0 = \partial_\nu \mathbf{u}_0^{\text{ext}}|_{\eta=\eta_+}$ in the translation case and $\mathbf{k}_0 = f \dot{\mathbf{g}}_0$ in the contraction case. The exterior problem (22) is thus completely determined since

- In the translation case, $\phi_0 = (0, 0)$, and no $\dot{\mathbf{g}}_0$ is needed,
- In the contraction case, $\phi_0 = \frac{1}{f} \partial_\eta \mathbf{u}_1^{\text{lay}}|_{\eta_-} = \dot{\mathbf{g}}_0$,

which thus entirely determines *a posteriori* the terms $\mathbf{u}_0^{\text{lay}}$ and $\partial_\eta \mathbf{u}_1^{\text{lay}}$.

The general procedure works as follows: if $(\mathbf{u}_\ell^{\text{ext}}, \mathbf{u}_\ell^{\text{lay}})$ are known for any $\ell \leq n-1$, then

$$\mathbf{u}_n^{\text{lay}}(\eta, \tau) = \mathbf{u}_n^{\text{lay}}(\eta_-, \tau) + (\eta - \eta_-) \mathbf{k}_{n-1}(\tau) + \int_{\eta_-}^{\eta} \mathbf{R}[\mathbf{j}_{n-1}](t, \tau) dt, \quad (23a)$$

$$\partial_\eta \mathbf{u}_{n+1}^{\text{lay}}(\eta, \tau) = \mathbf{k}_n(\tau) + \mathbf{R}[\mathbf{j}_n](\eta, \tau), \quad (23b)$$

where the terms \mathbf{j}_{n-1} , \mathbf{j}_n , \mathbf{k}_{n-1} , and \mathbf{k}_n are explicitly given. Note that neither $\mathbf{u}_n^{\text{lay}}$ nor $\mathbf{u}_n^{\text{ext}}$ are yet determined. However Problem (22) can be solved since ϕ_n is completely determined by

- In the translation case,

$$\phi_n(\tau) = \left((\eta_+ - \eta_-) \mathbf{k}_{n-1}(\tau) + \int_{\eta_-}^{\eta_+} \mathbf{R}[\mathbf{j}_{n-1}](t, \tau) dt, -\mathbf{R}[\mathbf{j}_n](\eta_-, \tau) \right),$$

- In the contraction case,

$$\phi_n(\tau) = \frac{1}{f(\tau)} \partial_\eta \mathbf{u}_{n+1}^{\text{lay}}(\eta^-, \tau) = \frac{1}{f(\tau)} (\mathbf{k}_n(\tau) + \mathbf{R}[\mathbf{j}_n](0, \tau)),$$

which thus determines *a posteriori* $\mathbf{u}_n^{\text{lay}}$ and $\partial_\eta \mathbf{u}_{n+1}^{\text{ext}}$.

The above derivation process can be summarized as follows

$$\mathbf{u}_n^{\text{ext}} = \mathbf{J}_n^{\text{ext}} \left((\mathbf{u}_\ell^{\text{ext}})_{0 \leq \ell \leq n-1}, (\mathbf{u}_\ell^{\text{lay}})_{0 \leq \ell \leq n-1} \right), \quad (24a)$$

$$\mathbf{u}_n^{\text{lay}} = \mathbf{J}_n^{\text{lay}} \left((\mathbf{u}_\ell^{\text{ext}})_{0 \leq \ell \leq n}, (\mathbf{u}_\ell^{\text{lay}})_{0 \leq \ell \leq n-1} \right), \quad (24b)$$

where $\mathbf{J}_n^{\text{ext}}$ and $\mathbf{J}_n^{\text{lay}}$ are the operators defined by the above problems (22) and (16) respectively. We will make use of $\mathbf{J}_0^{\text{lay}}$ as well, setting $\mathbf{u}_{-1}^{\text{lay}} = 0$ by convention.

Remark 3.3 *In the layer, the problem defining the terms $\mathbf{u}_n^{\text{lay}}$ is one-dimensional, and the tangential variable τ only acts as a parameter. The integration is only performed in the normal variable η . It is straightforward that the dependence on $\mathbf{u}_n^{\text{lay}}$ is polynomial in η , its degree being increased by 1 at each step. The dependence with respect to τ is more complex since it comes from traces on $\eta = \eta^\pm$ of the terms $\mathbf{u}_\ell^{\text{ext}}$ for $\ell < n$.*

3.4 Back to the physical domain

Remember that the above procedure is based on the assumption that all the coefficients c_λ^\pm and $a_n^{\pm,\sigma}$ vanish in the expansions of type (5). This implies that at any order, all the terms u_n^{ext} and thus u_n^{lay} are flat near the origin, see Section 2. It is however important to transfer the obtained approximate solution in the original domain.

- *Limit domain $\delta = 0$.*

In the contraction case, the first term u_0^{ext} previously determined coincides with the limit term u_0 defined by (2).

In the translation case, the domain $\dot{\Pi}^{\text{ext}}$, where u_0^{ext} is defined, is not connected. However the function \tilde{u}_0^{ext} defined in Π_0 by

$$\forall (x_1, x_2) \in \Pi_0, \quad \tilde{u}_0^{\text{ext}}(x_1, x_2) = \begin{cases} u_0^{\text{ext}}(x_1 + 1, x_2), & \text{if } x_1 > 0, \\ u_0^{\text{ext}}(x_1 - 1, x_2), & \text{if } x_1 < 0, \end{cases}$$

coincides with u_0 .

- *Approximation of u_δ in Π_δ .*

In the translation case, an approximation at order N is given by

$$u_\delta^{[N]}(x_1, x_2) = \begin{cases} \sum_{n=0}^N \delta^n u_n^{\text{ext}}(x_1 - 1 + \delta, x_2) & \text{if } x_1 < -\delta, \\ \sum_{n=0}^N \delta^n u_n^{\text{lay}}\left(\frac{x_1}{\delta}, x_2\right) & \text{if } -\delta < x_1 < \delta, \\ \sum_{n=0}^N \delta^n u_n^{\text{ext}}(x_1 + 1 - \delta, x_2) & \text{if } x_1 > \delta. \end{cases}$$

In the contraction case, approximations far from the corner have been derived thanks to the parameterization. In the vicinity of the corner, singularities should be dominant. However since we have assumed throughout Section 3 that no singularity appears, this means that at any order the functions are flat enough. Precisely, our hypothesis states that u_δ is $\mathcal{O}(|\mathbf{x}|^p)$ for any $p \in \mathbb{N}$. This implies that 0 is a good approximation of u_δ near the corner. Therefore an approximation of u_δ at the order δ^N is given by

$$u_\delta^{[N]}(\mathbf{x}) = \begin{cases} \sum_{n=0}^N \delta^n u_n^{\text{ext}}(\mathbf{x}) & \text{in } \Pi_0 \setminus \mathcal{B}_{\delta\rho_\star} \\ 0 & \text{in } \mathcal{B}_{\delta\rho_\star} \text{ (i.e. for } |x| < \delta\rho_\star), \\ \sum_{n=0}^N \delta^n u_n^{\text{lay}}\left(\Phi_\delta^{\star-1}(\mathbf{x})\right) & \text{elsewhere,} \end{cases}$$

where the mapping Φ_δ^\star is defined in (11).

4 Handling singularities

4.1 Need for a specific procedure

As mentioned before, the procedure described in Section 3.3 fails to provide an asymptotic expansion of u_δ when singularities appear. We detail here the reasons why.

First obstruction: lack of regularity for the sequential derivation

The formal derivation described in Section 3.3 fails since singularities appear in the solutions to (22). To explain this point, we distinguish the two configurations.

- *Translation case.*

In order to solve Problem (22) in a variational framework, it is necessary that ϕ_n has regularity $H^{1/2}(\dot{\Gamma}^{\text{trans}}) \times H^{-1/2}(\dot{\Gamma}^{\text{trans}})$. Using Equations (23), the condition over the first component needs generically $k_{n-1} \in H^{1/2}(\dot{\Gamma}^{\text{trans}})$ which requires $\dot{u}_{n-1}^{\text{ext}} \in H^2(\dot{\Pi}^{\text{ext}})$.

Thus we need that $\dot{u}_{n-1}^{\text{ext}}$ belongs to $H^2(\dot{\Pi}^{\text{ext}})$ to ensure the existence of $\dot{u}_n^{\text{ext}} \in H^1(\dot{\Pi}^{\text{ext}})$. Actually one order of regularity is lost at each step. Since \dot{u}_0^{ext} has limited regularity (due to the presence of singularities \mathfrak{s}_λ^\pm), an expansion at any order is hopeless (if the angles α^\pm are close to π , the maximal order for the construction may be $n = 1$).

- *Contraction case.*

Here, ϕ_n has to belong to $H^{-1/2}(\Gamma^{\text{trans}})$, which requires, thanks to Equations (23) and Definition (19), that $\tau \mapsto j_n(\eta, \tau)$ has $H^{-1/2}$ -regularity. But the expression of j_n , see (17), involves $\overset{\star}{\mathbb{T}}_2 \overset{\star}{u}_{n-1}^{\text{lay}}$ and with $\overset{\star}{u}_{n-1}^{\text{lay}}|_{\eta=\eta_-} = \overset{\star}{u}_{n-1}^{\text{ext}}|_{\Gamma^{\text{trans}}}$, and the condition $\overset{\star}{u}_{n-1}^{\text{ext}} \in H^2(\overset{\star}{\Pi}^{\text{ext}})$ is needed. Once again, we loose 2 orders of regularity every 2 steps.

Second obstruction: terms are not necessarily flat near the origin

We assume here that the first obstruction occurs for the construction of u_2^{ext} . This arises if $u_1^{\text{ext}} \notin H^2(\Pi^{\text{ext}})$. Precisely, according to (5) for $p = 1$, the following splitting holds for the exterior part

$$u_1^{\text{ext}}(\Phi_\delta^{-1}(\mathbf{x})) = u_{\text{flat},1}^{1,\text{ext}}(\Phi_\delta^{-1}(\mathbf{x})) + \chi(\mathbf{x}) \sum_{\pm} \sum_{\lambda \in \Lambda_2^\pm} c_{\lambda,1}^\pm \mathfrak{s}_\lambda^\pm(\mathbf{x}) + \chi(\mathbf{x}) \sum_{\pm} \sum_{\sigma=1,2} a_{1,1}^{\pm,\sigma} \mathfrak{a}_1^{\pm,\sigma}(\mathbf{x}), \quad (25)$$

with $u_{\text{flat},1}^{1,\text{ext}} \in H^2(\Pi^{\text{ext}})$, $\mathfrak{s}_\lambda^\pm \notin H^2(\Pi_0)$, $\mathfrak{a}_1^{\pm,\sigma} \in C^\infty(\overline{\Pi_0})$. Therefore, if the coefficients $c_{\lambda,1}^\pm$ vanish (again, this is not the general situation), we have $u_1^{\text{ext}} \in H^2(\Pi^{\text{ext}})$, which seems sufficient for the procedure. However the terms $a_{1,1}^{\pm,\sigma}$, which come from the Neumann traces, add new difficulties.

- *Translation case.*

In order to build u_2^{lay} , it is necessary to derive $u_0^{\text{lay}}(\eta, \cdot) = u_0^{\text{ext}}|_{\Gamma^{\text{trans}}}$ twice with respect to τ , which is impossible in general, since $a_{1,1}^{+,\sigma} \neq a_{1,1}^{-,\sigma}$. Introducing a truncation near the origin makes it possible to prevent this case, similarly to the contraction case. However this makes appear a non negligible error near $\mathbf{0}$.

- *Contraction case.*

Here, u_2^{ext} can be built. Nevertheless, we do not end up with a suitable asymptotic expansion. Indeed, coming back to the physical domain (see Section 3.4), we get a correct approximation only away from the origin because of the extension by 0 near $\mathbf{0}$.

Third obstruction: error estimates fail

Should the above obstructions be overcome to define u_1 , a third issue would appear for the error estimates. Actually, the terms $u_\delta - u_\delta^{[1]} \psi(|\mathbf{x}|/\delta)$ cannot be used to obtain error estimates by computing the residual, since Δu_1 is not well-defined in Π^{lay} as a function.

How to overcome these obstructions ?

We have seen that the problem comes from the limited regularity of u_0^{ext} . In the domain Π_0 , let us use the splitting into regular and singular parts, see (5),

$$\begin{aligned} u_0^{\text{ext}}(\mathbf{x}) &:= u_0^{\text{ext}}(\Phi_\delta^{-1}(\mathbf{x})) \\ &= u_{\text{flat},0}^{p,\text{ext}}(\mathbf{x}) + \chi(\mathbf{x}) \sum_{\pm} \sum_{\lambda \in \Lambda_p^\pm} c_{\lambda,0}^\pm \mathfrak{s}_\lambda^\pm(\mathbf{x}) + \chi(\mathbf{x}) \sum_{\pm} \sum_{\sigma=1,2} \sum_{1 \leq n \leq p-1} a_{n,0}^{\pm,\sigma} \mathfrak{a}_n^{\pm,\sigma}(\mathbf{x}), \end{aligned} \quad (26)$$

with $u_{\text{flat},0}^{p,\text{ext}} := u_{\text{flat},0}^{p,\text{ext}} \circ \Phi_\delta^{-1} \in H_{\text{loc}}^p(\Pi_0)$, flat near the origin. In the layer Π^{lay} , we set

$$\begin{aligned} u_0^{\text{lay}}(\mathbf{x}) &:= u_0^{\text{lay}}(\Phi_\delta^{-1}(\mathbf{x})) \\ &= u_{\text{flat},0}^{p,\text{lay}}(\Phi_\delta^{-1}(\mathbf{x})) + \chi(\mathbf{x}) \sum_{\pm} \sum_{\lambda \in \Lambda_p^\pm} c_{\lambda,0}^\pm \mathfrak{s}_\lambda^\pm(\mathbf{x}) + \chi(\mathbf{x}) \sum_{\pm} \sum_{\sigma=1,2} \sum_{1 \leq n \leq p-1} a_{n,0}^{\pm,\sigma} \mathfrak{a}_n^{\pm,\sigma}(\mathbf{x}), \end{aligned} \quad (27)$$

with $u_{\text{flat},0}^{p,\text{lay}} = J_0^{\text{lay}} \left(\left(u_{\text{flat},0}^{p,\text{ext}}, (0) \right) \right)$.

It is possible to build further the expansion, with $u_{\text{flat},0}^p$ instead of u_0 . The term u_1 is replaced with \check{u}_1 defined by

$$\check{u}_1^{\text{ext}} = J_1^{\text{ext}} \left(\left(u_{\text{flat},0}^{p,\text{ext}}, u_{\text{flat},0}^{p,\text{lay}} \right) \right), \quad (28a)$$

$$\check{u}_1^{\text{lay}} = J_1^{\text{lay}} \left(\left(u_{\text{flat},0}^{p,\text{ext}}, \check{u}_1^{\text{ext}} \right) \right). \quad (28b)$$

Of course, we do not have taken the singularities \mathfrak{s}_λ^\pm and $\mathfrak{a}_n^{\pm,\sigma}$ into account. This requires to introduce *corner profiles* in the expansion.

4.2 Heuristics for the introduction of the first profile

According to the above remarks, $u_\delta^{[1]}$ is not an approximation of u_δ at the order δ . It is not the case either for the function $\check{u}_\delta^{[1]}$, defined similarly to $u_\delta^{[1]}$ in Section 3.4 with \check{u}_1 instead of u_1 .

Let us go back to the approximation at the order 0. We set $r_\delta = u_\delta - \psi(\cdot/\delta)u_\delta^{[0]}$ where $u_\delta^{[0]}$ is given in Section 3.4. Here ψ is a smooth radial¹ cutoff function independent of δ and equal to 1 far from $\mathbf{0}$ and which vanishes near $\mathbf{0}$.

The remainder term r_δ is well-defined in Π_δ and satisfies the following transmission problem:

$$\left\{ \begin{array}{ll} \Delta r_\delta = - [\Delta; \psi(\frac{\cdot}{\delta})] u_0 + \frac{1}{\delta^2} \psi(\frac{\cdot}{\delta}) \left(\mathbb{T}_1 u_0 + \sum_{k \geq 2} \delta^k \mathbb{T}_k u_0 \right) \circ \Phi_\delta^{-1} & \text{in } \Pi_\delta^{\text{lay}}, \\ \Delta r_\delta = - [\Delta; \psi(\frac{\cdot}{\delta})] u_0 & \text{in } \Pi_\delta^{\text{ext}}, \\ [r_\delta] = 0 & \text{on } \Gamma_\delta^{\text{trans}}, \\ [\partial_\nu r_\delta] = -\psi(\frac{\cdot}{\delta}) \partial_\nu u_0 & \text{on } \Gamma_\delta^{\text{trans}}, \\ \partial_\nu r_\delta = (1 - \psi(\frac{\cdot}{\delta})) \sum_{n \geq 0} \delta^n g_n - \frac{\psi(\frac{\cdot}{\delta})}{f} \sum_{k \geq 2} \delta^k (\mathbb{B}_k u_0) \circ \Phi_\delta^{-1} & \partial \Pi_\delta^{\text{ext}}, \\ r_\delta \rightarrow 0 & \text{at infinity,} \end{array} \right. \quad (29)$$

¹By abuse of notation, we use ‘‘radial’’ in the layer as well, but in the translation case, this corresponds to a dependence in η .

where u_0 is the solution to (2). Note that in the translation case, the operators B_k are zero. The term $[\Delta; \Psi]$ generically refers to the commutator operator given by

$$[\Delta; \Psi] V = 2\nabla\Psi \cdot \nabla V + V\Delta\Psi,$$

for smooth enough Ψ and V , and thus the term $[\Delta; \psi(\cdot/\delta)] u_0$ is of order δ^{-2} . In Problem (29), the main source terms involve the contribution of u_0 and g_0 both taken near the origin $\mathbf{0}$, since $\psi(\cdot/\delta)$ equals 1 far from it.




Figure 4: The infinite domain \mathfrak{P} (corresponding to the non-symmetric case, see Fig. 1).

Splitting u_0 near the origin into regular and singular parts, see (26)–(27), we see that the principal part of the error is given by s_0^\pm . It is natural to introduce the profile in the infinite domain $\mathfrak{P} = \lim_{\delta \rightarrow 0} \delta^{-1} \Pi_\delta$, see Fig. 4, defined as

$$\begin{cases} \Delta \mathfrak{R}_0^\pm &= [\Delta; \psi] s_0^\pm & \text{in } \mathfrak{P}, \\ \partial_\nu \mathfrak{R}_0^\pm &= 0 & \text{on } \partial\mathfrak{P}, \\ \mathfrak{R}_0^\pm &\rightarrow 0 & \text{at infinity,} \end{cases} \quad (30)$$

where s_0^\pm have been extended by a constant on the support of $\nabla\psi$. Moreover, the Neumann boundary condition also generates a profile denoted by \mathfrak{M} :

$$\begin{cases} \Delta \mathfrak{M} &= 0 & \text{in } \mathfrak{P}, \\ \partial_\nu \mathfrak{M} &= (1 - \psi) & \text{on } \partial\mathfrak{P}, \\ \mathfrak{M} &\rightarrow 0 & \text{at infinity.} \end{cases} \quad (31)$$

The well-posedness and the behavior at infinity of \mathfrak{R}_0^\pm and \mathfrak{M} are discussed in Section 4.4. These profiles make it possible to correct the expansion as follows

$$u_\delta(\mathbf{x}) = \psi\left(\frac{\mathbf{x}}{\delta}\right) u_0(\mathbf{x}) - \chi(\mathbf{x}) \left(g_0(\mathbf{0}) \mathfrak{M}\left(\frac{\mathbf{x}}{\delta}\right) + \sum_{\pm} c_{0,0}^\pm \mathfrak{R}_0^\pm\left(\frac{\mathbf{x}}{\delta}\right) \right) + \mathfrak{R}_\delta(\mathbf{x}).$$

Thanks to (30), the remainder \mathfrak{R}_δ satisfies the following problem:

$$\begin{cases} \Delta \mathfrak{R}_\delta &= [\Delta; \psi(\cdot/\delta)] (\sum_{\pm} c_{0,0}^\pm s_0^\pm - u_0) \\ &\quad + [\Delta; \chi] \left(g_0(\mathbf{0}) \mathfrak{M}(\cdot/\delta) + \sum_{\pm} c_{0,0}^\pm \mathfrak{R}_0^\pm(\cdot/\delta) \right) & \text{in } \Pi_\delta^{\text{lay}} \cup \Pi_\delta^{\text{ext}}, \\ [\mathfrak{R}_\delta] &= 0 & \text{on } \Gamma_\delta^{\text{trans}}, \\ [\partial_\nu \mathfrak{R}_\delta] &= \mathcal{O}(1) & \text{on } \Gamma_\delta^{\text{trans}}, \\ \partial_\nu \mathfrak{R}_\delta &= \mathcal{O}(\delta) & \text{on } \partial\Pi_\delta^{\text{ext}}, \\ \mathfrak{R}_\delta &\rightarrow 0 & \text{at infinity,} \end{cases} \quad (32)$$

The leading term of $c_{0,0}^+ s_0^+ + c_{0,0}^- s_0^- - u_0$ near the origin is supported by $(\mathfrak{a}_{1,i}^\pm)_{i=1,2}$ according to the splitting (26) of u_0 , and it is thus of order δ . The term $[\Delta; \chi] \left(g_0(\mathbf{0}) \mathfrak{M}(\cdot/\delta) + \sum_{\pm} c_{0,0}^\pm \mathfrak{R}_0^\pm(\cdot/\delta) \right)$

vanishes inside a ball of radius $\mathcal{O}(1)$ centered at the origin, and thus its leading term is given by the behavior of the profiles \mathfrak{K}_0^\pm and \mathfrak{M} at infinity, that is $\mathcal{O}(\delta^{\min(\lambda_1^+, \lambda_1^-)})$, where $\lambda_1^\pm = \pi/\alpha^\pm > 1$. Therefore \mathfrak{R}_δ has a smaller residual than r_δ , and the expansion is improved.

In order to build the terms of order δ , it is necessary to account for three contributions

- The next term in the splitting of u_0^{ext} which involves the singular functions $a_1^{\pm, \sigma}$, see Section 2.
- The Neumann source term $(1 - \psi(\cdot/\delta))g_0$ near the origin.
- The term \check{u}_1 defined in (28), and in particular its first singular terms, involving the singular functions s_0^\pm .

The third contribution is addressed by the profile \mathfrak{K}_0^\pm defined by (30), while the first two contributions will be handled by new specific profiles $\mathfrak{L}_1^{\pm, \sigma}$ defined as:

$$\begin{cases} \Delta \mathfrak{L}_1^{\pm, \sigma} &= [\Delta; \psi] a_1^{\pm, \sigma} & \text{in } \mathfrak{B}, \\ \partial_\nu \mathfrak{L}_1^{\pm, \sigma} &= 0 & \text{on } \partial \mathfrak{B}, \\ \mathfrak{L}_1^{\pm, \sigma} &\rightarrow 0 & \text{at infinity.} \end{cases} \quad (33)$$

The expansion at the next order reads then

$$\begin{aligned} u_\delta(\mathbf{x}) &= \psi\left(\frac{\mathbf{x}}{\delta}\right) u_0(\Phi_\delta^{-1}(\mathbf{x})) - \chi(\mathbf{x}) \left(\sum_{\pm} c_{0,0}^\pm \mathfrak{K}_0^\pm\left(\frac{\mathbf{x}}{\delta}\right) + g_0(\mathbf{0}) \mathfrak{M}\left(\frac{\mathbf{x}}{\delta}\right) \right) \\ &\quad + \delta \psi\left(\frac{\mathbf{x}}{\delta}\right) \check{u}_1(\Phi_\delta^{-1}(\mathbf{x})) - \delta \chi(\mathbf{x}) \left(\sum_{\pm} \left(c_{1,0}^\pm \mathfrak{K}_0^\pm\left(\frac{\mathbf{x}}{\delta}\right) + \sum_{\sigma=1,2} a_1^{\pm, \sigma} \mathfrak{L}_1^{\pm, \sigma}\left(\frac{\mathbf{x}}{\delta}\right) \right) + g_1(\mathbf{0}) \mathfrak{M}\left(\frac{\mathbf{x}}{\delta}\right) \right) \\ &\quad + \mathcal{O}(\delta^{\min(\lambda_1^+, \lambda_1^-)}), \end{aligned} \quad (34)$$

where $c_{1,0}^\pm$ are the first singular coefficients of \check{u}_1^{ext} in the splitting near the origin.

The above expansion has *a priori* an order of accuracy in $\mathcal{O}(\delta^{\min(\lambda_1^+, \lambda_1^-)})$. To go one step further, it is necessary to change the whole expansion, and to account for one more singular term in the splitting of u_0^{ext} , which modifies $u_{\text{flat},0}^{0,\text{ext}}$ and thus \check{u}_1 . So it is necessary to determine *a priori* the desired order of accuracy, so that u_0^{ext} be split with the appropriate number of singular functions, and then the expansion can be performed.

4.3 Structure of the complete asymptotic expansion

In the previous section, we have described the construction of the first terms in the expansion, taking into account the singularities coming from the corners of the domain. It is possible to go further and build a complete asymptotic expansion at any order. To do so, we need to fix a target order P , and expand every term defined in Π^{ext} into singular and regular parts according to this choice. The contribution of all singular functions arising in such splittings will be handled through profiles (of type \mathfrak{K} , \mathfrak{L} , \mathfrak{M}) defined in the infinite domain \mathfrak{B} . The expansion of these profiles at infinity will give birth to other terms defined in Π^{ext} , which in turn will be split into regular and singular parts.

In order to obtain an expansion at order P , the overall procedure leads to

$$u_\delta^{\text{ext}}(\mathbf{x}) = \psi\left(\frac{\mathbf{x}}{\delta}\right) \sum_{\mu \in M_P} \delta^\mu u_\mu^{P,\text{ext}}(\mathbf{x}) - \chi(\mathbf{x}) \sum_{\mu \in M_P} \sum_{\pm} \sum_{\sigma=1,2} \delta^\mu \mathfrak{W}_\mu^{\pm, \sigma, P}\left(\frac{\mathbf{x}}{\delta}\right) + R_{\delta, P}^{\text{ext}}(\mathbf{x}),$$

where

- the terms $u_\mu^{P,\text{ext}}$ are defined in the standard variable \mathbf{x} inside Π^{ext} ,
- the terms $\mathfrak{W}_\mu^{\pm,\sigma,P}$ are defined on the infinite domain \mathfrak{F} , and are involved in the asymptotic expansion as functions of the scaled variable $\frac{\mathbf{x}}{\delta}$,
- the set of indices M_P is given by

$$M_P = \left\{ \mu = p + \frac{q^- \pi}{\alpha_-} + \frac{q^+ \pi}{\alpha_+} ; p, q^\pm \in \mathbb{N}_0, \text{ and } \mu \leq P \right\},$$

- the remainder R_P^{ext} satisfies on any compact subset $\omega \subset \Pi_\delta$:

$$\|R_{\delta,P}^{\text{ext}}\|_{\text{H}^1(\omega)} = o(\delta^P).$$

In the layer, the expansion has a similar structure, but involves the normal dilation used to build the terms :

$$u_\delta^{\text{lay}}(\mathbf{x}) = \psi\left(\frac{\mathbf{x}}{\delta}\right) \sum_{\mu \in M_P} \delta^\mu u_\mu^{P,\text{lay}}\left(\frac{\eta}{\delta}, \tau\right) - \chi(\mathbf{x}) \sum_{\mu \in M_P} \sum_{\pm} \sum_{\sigma=1,2} \delta^\mu \mathfrak{W}_\mu^{\pm,\sigma,P}\left(\frac{\mathbf{x}}{\delta}\right) + R_{\delta,P}^{\text{lay}}(\mathbf{x}).$$

The remainder R_P^{ext} satisfies on any compact subset $\omega \subset \Pi_\delta$:

$$\sqrt{\delta} \|R_{\delta,P}^{\text{lay}}\|_{\text{H}^1(\omega)} = o(\delta^P).$$

Remark 4.1 *If P is changed, the terms involved in the expansion will change as well, since the order of every splitting needs to be adapted accordingly.*

4.4 Construction of profiles: existence and behavior at infinity

In the last two sections, we have made use of various *profiles*, which account for the singularities appearing at each order in the construction of the expansion. All the profiles $(\mathfrak{R}_\lambda^\pm, \mathfrak{L}_\lambda^{\pm,\sigma}, \mathfrak{M})$ satisfy a problem such as

$$\begin{cases} \Delta \mathfrak{Z} = \mathfrak{f} & \text{in } \mathfrak{F}, \\ \partial_\nu \mathfrak{Z} = \mathfrak{g} & \text{on } \partial \mathfrak{F}, \\ \mathfrak{Z} \rightarrow 0 & \text{at infinity.} \end{cases} \quad (35)$$

The existence for Problem (35) relies on the space

$$\mathfrak{V} = \left\{ \mathfrak{Z} \in \text{H}_{\text{loc}}^1(\mathfrak{F}) ; \nabla \mathfrak{Z} \in \text{L}^2(\mathfrak{F}) \quad \text{and} \quad \frac{\mathfrak{Z}}{(1 + |\mathbf{X}|) \log(2 + |\mathbf{X}|)} \in \text{L}^2(\mathfrak{F}) \right\},$$

and the bilinear form

$$a(\mathfrak{Z}_1, \mathfrak{Z}_2) = \int_{\mathfrak{F}} \nabla \mathfrak{Z}_1 \cdot \nabla \mathfrak{Z}_2$$

which is coercive on the quotient space \mathfrak{V}/\mathbb{R} , see [2, 14]. Within this variational framework, we can build the profiles and make precise their behavior at infinity:

Theorem 4.2 *If \mathfrak{f} and \mathfrak{g} are smooth functions with compact support in \mathfrak{B} , and satisfy*

$$\int_{\mathfrak{B}} \mathfrak{f} + \int_{\partial\mathfrak{B}} \mathfrak{g} = 0,$$

then problem (35) admits a unique solution $\mathfrak{Z} \in \mathfrak{B}/\mathbb{R}$. The behavior of the solution \mathfrak{Z} at infinity is given by

$$\mathfrak{Z} = \sum_{\pm} \sum_{\lambda \in \Lambda_p^{\pm}} \mathfrak{Z}_{-\lambda}^{\pm} + \mathfrak{R}_p, \quad (36)$$

with $\partial^{\beta} \mathfrak{Z}_{-\lambda}^{\pm} = \mathcal{O}(|\mathbf{X}|^{-\lambda-|\beta|})$, and $\partial^{\beta} \mathfrak{R}_p = \mathcal{O}(|\mathbf{X}|^{-p-|\beta|})$ for any multi-index β (as $|\mathbf{X}| \rightarrow \infty$).

Relation (36) show in particular that $\mathfrak{Z} \rightarrow 0$ at infinity, which is crucial in the construction of the asymptotic expansion, see § 4.2. Note that Assumption (H4) allows us to avoid the use of the logarithmic singularity, which naturally appears for the twodimensional Laplace-Neumann problem, see [5, Th. 3.25].

5 Discussion and concluding remarks

5.1 Representation formula

From the asymptotic expansion, one can easily infer a representation formula far from the singularity. For any \mathbf{y} far from the inclusions, let define the Neumann function $N_{\mathbf{y}}$ as the solution to

$$\begin{cases} -\Delta N_{\mathbf{y}} &= \delta_{\mathbf{y}} & \text{in } \Pi_0, \\ \partial_{\nu} N_{\mathbf{y}} &= 0 & \text{on } \partial\Pi_0, \\ N_{\mathbf{y}} &\rightarrow 0 & \text{at infinity.} \end{cases} \quad (37)$$

Then, simple integration by parts leads to the following equality

$$u_{\delta}(\mathbf{y}) - u_0(\mathbf{y}) = -\delta \int_{\partial\Pi_0} \partial_{\nu} \check{u}_1^{\text{ext}} N_{\mathbf{y}} d\mathbf{x} + \mathcal{O}(\delta),$$

and by definition of \check{u}_1^{ext} and thanks to (26), one infers the following representation formula:

$$u_{\delta}(\mathbf{y}) - u_0(\mathbf{y}) = \delta \int_{\partial\Pi_0} \partial_{\tau} u_{\text{flat},0}^{1,\text{ext}} \partial_{\tau} N_{\mathbf{y}} d\mathbf{x} + \mathcal{O}(\delta).$$

Such a representation formula is useful to characterise (or to detect) the detachment of the two inclusions tied up by the corners.

5.2 Other possible geometrical frameworks

Let us now discuss some possible extensions. We have detailed the two asymptotic models (translation and contraction) in the special situation of assumptions (H1)–(H2)–(H3). Other geometric situations may be considered, for which the two above methods have to be slightly modified. We give in Figure 5 two examples for which assumption (H3) is violated.

In Figure 5a, the domains Ω_{δ}^L and Ω_{δ}^R do not lie in separated half planes orthogonal to the horizontal direction. The contraction method can be applied directly but the translation method needs to be adapted. Indeed, a vertical strip cannot be inserted between the two inclusions and we need to use two slanted half-strips. The analysis remains very similar.




Figure 5: Geometrical situations which do not fulfill assumption (H3).

In Figure 5b, the domains Ω_δ^L and Ω_δ^R lie on the upper half plane. The translation method applies directly but the contraction needs adaptation in this case. Actually, the contraction method cannot be applied if the direction \mathbf{d} remains to be horizontal. It is necessary to assume that the inclusions converge to each other at a point which lies under the origin $\mathbf{0}$.

In Figure 6, we present the situation of two nearly touching squares where both methods need more important adaptations. The limit situation shows a contact on a segment and not at a single point. Other profiles must be introduced to account for the geometry near its extremity points.




Figure 6: Geometrical situations which do not fulfill assumption (H1).

Let us emphasize that hypothesis (H2) is crucial to our two asymptotic expansions. In particular regarding the case of two *kissing balls*, the limit problem is no more a corner problem, but a problem with a cusp. The structure of the singularities is then rather different in this case as presented in see [7]. Our methods cannot be applied.




Figure 7: Geometrical situations which do not fulfill assumption (H2).

References

- [1] H. Ammari, H. Kang, and K. Touibi. An asymptotic formula for the voltage potential in the case of a near-surface conductivity inclusion. *Z. Angew. Math. Phys.*, 57(2):234–243, 2006.
- [2] C. Amrouche, V. Girault, and J. Giroire. Dirichlet and Neumann exterior problems for the n -dimensional Laplace operator: an approach in weighted Sobolev spaces. *J. Math. Pures Appl. (9)*, 76(1):55–81, 1997.
- [3] E. Bonnetier and F. Triki. On the spectrum of the Poincaré variational problem for two close-to-touching inclusions in 2D. *Arch. Ration. Mech. Anal.*, 209(2):541–567, 2013.
- [4] E. Bonnetier, F. Triki, and C.-H. Tsou. Eigenvalues of the Neumann-Poincaré operator for two inclusions with contact of order m : a numerical study. *J. Comput. Math.*, 36(1):17–28, 2018.
- [5] G. Caloz, M. Costabel, M. Dauge, and G. Vial. Asymptotic expansion of the solution of an interface problem in a polygonal domain with thin layer. *Asymptot. Anal.*, 50(1-2):121–173, 2006.
- [6] M. Dauge. *Elliptic boundary value problems on corner domains*, volume 1341 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 1988. Smoothness and asymptotics of solutions.
- [7] M. Dauge. Strongly elliptic problems near cuspidal points and edges. In *Partial differential equations and functional analysis*, volume 22 of *Progr. Nonlinear Differential Equations Appl.*, pages 93–110. Birkhäuser Boston, Boston, MA, 1996.
- [8] P. Grisvard. *Elliptic problems in nonsmooth domains*, volume 24 of *Monographs and Studies in Mathematics*. Pitman (Advanced Publishing Program), Boston, MA, 1985.
- [9] V. A. Kondratiev. Boundary value problems for elliptic equations in domains with conical or angular points. *Trudy Moskov. Mat. Obsc.*, 16:209–292, 1967.
- [10] V. Maz’ya, S. A. Nazarov, and B. A. Plamenevskij. *Asymptotic theory of elliptic boundary value problems in singularly perturbed domains*. Birkhäuser, Berlin, 2000.
- [11] A. B. Movchan and S. A. Nazarov. Asymptotic behavior of the stress-strained state near sharp inclusions. *Dokl. Akad. Nauk SSSR*, 290(1):48–51, 1986.
- [12] S. A. Nazarov and M. E. Pérez. On multi-scale asymptotic structure of eigenfunctions in a boundary value problem with concentrated masses near the boundary. *Rev. Mat. Complut.*, 31(1):1–62, 2018.
- [13] S. A. Nazarov and J. Taskinen. Singularities at the contact point of two kissing Neumann balls. *J. Differential Equations*, 264(3):1521–1549, 2018.
- [14] J.-C. Nédélec. *Acoustic and electromagnetic equations*, volume 144 of *Applied Mathematical Sciences*. Springer-Verlag, New York, 2001. Integral representations for harmonic problems.