



# **Density-dependent increase in superpredation linked to food limitation in a recovering population of northern goshawks *Accipiter gentilis***

S. R Hoy, S. J Petty, Alexandre Millon, D. P Whitfield, M. Marquiss, D. I. K. Anderson, M. Davison, X. Lambin

## **► To cite this version:**

S. R Hoy, S. J Petty, Alexandre Millon, D. P Whitfield, M. Marquiss, et al.. Density-dependent increase in superpredation linked to food limitation in a recovering population of northern goshawks *Accipiter gentilis*. *Journal of Avian Biology*, 2017, 48 (9), pp.1205-1215. <10.1111/jav.01387>. <hal-01768454>

**HAL Id: hal-01768454**

**<https://hal.science/hal-01768454v1>**

Submitted on 2 May 2018

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

1       **Density-dependent increase in superpredation linked to food**  
2       **limitation in a recovering population of northern goshawks,**  
3       *Accipiter gentilis*

4       S.R. Hoy <sup>a,b</sup>, S.J. Petty <sup>a</sup>, A. Millon <sup>c</sup>, D.P. Whitfield <sup>d</sup>, M. Marquiss <sup>a</sup>, D.I.K. Anderson <sup>e</sup>, M.  
5       Davison <sup>f</sup>, & X. Lambin <sup>a\*</sup>

6       <sup>a</sup> School of Biological Sciences, University of Aberdeen, Aberdeen AB24 2TZ, UK

7       <sup>b</sup> Michigan Technological University, Houghton, Michigan 49931

8       <sup>c</sup> Aix Marseille Universite, Universite Avignon, CNRS, IRD, IMBE, Technopôle Arbois-  
9       Méditerranée, 13545 Aix-en-Provence, France

10       <sup>d</sup> Natural Research Limited, Banchory, AB31 4BY, UK

11       <sup>e</sup> Forest Enterprise Scotland, Aberfoyle, Stirling, FK8 3UX

12       <sup>f</sup> Forestry Commission, Bellingham, NE48 2HP, UK

13       \* Corresponding author: x.lambin@abdn.ac.uk

14  
15       **Keywords:** food-stress hypothesis, intraguild predation, mesopredator suppression

## ABSTRACT

A better understanding of the mechanisms driving superpredation, the killing of smaller mesopredators by larger apex predators, is important because of the crucial role superpredation can play in structuring communities and because it often involves species of conservation concern. Here we document how the extent of superpredation changed over time, and assessed the impact of such temporal variation on local mesopredator populations using 40 years of dietary data collected from a recovering population of northern goshawks (*Accipiter gentilis*), an archetypical avian superpredator. We then assessed which mechanisms were driving variation in superpredation, e.g., was it opportunistic, a response to food becoming limited (due to declines in preferred prey) or to reduce competition. Raptors comprised 8% of goshawk diet on average in years when goshawk abundance was high, which is higher than reported elsewhere. Additionally, there was a per capita increase in superpredation as goshawks recovered, with the proportion of goshawk diet comprising raptors increasing from 2% to 8% as the number of goshawk home-ranges increased from  $\leq 14$  to  $\geq 25$ . This increase in superpredation coincided with a population decline in the most commonly killed mesopredator, the Eurasian kestrel (*Falco tinnunculus*), which may represent the reversal of the “mesopredator release” process (i.e., mesopredator suppression) which occurred after goshawks and other large raptors declined or were extirpated. Food limitation was the most likely driver of superpredation in this system given: 1) the substantial decline of two main prey groups in goshawk diet, the increase in diet diversity and decrease in goshawk reproductive success are all consistent with the goshawk population becoming food-limited; 2) it’s unlikely to be purely opportunistic as the increase in superpredation did not reflect changes in the availability of mesopredator species; and 3) the majority of mesopredators killed by goshawks do not compete with goshawks for food or nest sites.

## INTRODUCTION

Understanding the mechanisms driving variation in superpredation, the killing of smaller mesopredators by larger apex predators, is an important issue in ecology. This is partly because superpredation can directly impact mesopredator population dynamics which may then cascade to affect lower trophic levels (Paine 1980), but also because many of the superpredator and mesopredator species involved are of conservation concern (Palomares and Caro 1999, Caro and Stoner 2003, Ripple and Beschta 2004, Ritchie and Johnson 2009). However, despite this, and the crucial role superpredation can play in structuring communities, it is still not clear what mechanism (or combination of mechanisms) drives one predator to kill another.

Optimal foraging theory suggests that predators should attempt to kill prey when the energy gained outweighs the energetic cost and potential risk of injury involved (Berger-Tal et al. 2009). However, mesopredators are unlikely to represent a profitable prey source, even when they fall within the preferred size range of the superpredator, because their densities are often relatively low compared to that of other prey species. Furthermore, the risk of injury associated with attacking mesopredators is presumably higher than for other prey types, because mesopredators have evolved to kill other species (Lourenço et al. 2011a). Consequently, several alternative (but not mutually exclusive) hypotheses have been put forward to explain superpredation. The competitor-removal hypothesis suggests that superpredators kill mesopredators to free up shared resources (Serrano 2000). This leads to the prediction that superpredation will largely involve mesopredator species which compete with the superpredator for food or other resources, such as nest sites (e.g. intraguild predation). In contrast, the predator-removal hypothesis suggests that superpredation is a pre-emptive tactic to decrease the probability of the superpredator or their offspring being killed (Lourenço et al.

2011b). Under this scenario, the mesopredator species expected to be killed the most frequently are those which pose a threat to the superpredator or their offspring.

Alternatively, rather than being a response to the presence of other predators, the food-limitation hypothesis, also known as the food-stress hypothesis, suggests that mesopredators are killed to make up the shortfall in the superpredators diet when preferred prey species decline (Polis et al. 1989, Serrano 2000, Rutz and Bijlsma 2006, Lourenço et al. 2011a, b). Food limitation may also occur if there is an increase in the number of individuals (conspecifics or other species) exploiting preferred prey species, particularly if increasing predator densities elicit anti-predator behaviours in their prey (such as spatial or temporal avoidance of risky areas) which make prey more difficult to find and/or catch. Many populations of large predator species are currently increasing in abundance and recovering their former ranges across both North America and Europe (Maehr et al. 2001, Deinet et al. 2013, Chapron et al. 2014). Consequently, if superpredation is a response to density dependent food limitation, then the extent of superpredation occurring might be expected to increase during the recolonisation process, even if mesopredators are not a preferred prey species. However, whether such a per capita increase in superpredation has actually occurred or whether it coincides with or follows the colonisation process is as yet unknown.

Here we evaluate support for the food-limitation hypothesis, and other proposed determinants of superpredation in a recovering population of northern goshawks (*Accipiter gentilis*), using data collected between 1973 and 2014, over a 964km<sup>2</sup> area of Kielder Forest, United Kingdom (55°13'N, 2°33'W). The northern goshawk (hereafter goshawk) is an archetypical avian superpredator known to prey upon a large diversity of both avian and mammalian prey, including other raptors (Rutz et al. 2006, Sergio and Hiraldo 2008, Lourenço et al. 2011a).

Goshawks are the apex predator in this study system as Kielder Forest lacks other predator species known to prey on goshawks, such as Eurasian eagle owls (*Bubo bubo*) (Chakarov and Krüger 2010).

Goshawks were extirpated from the UK in the late 19<sup>th</sup> century. However, scattered populations were subsequently re-established in the 1960s and 70s after birds escaped or were released by falconers (Marquiss and Newton 1982, Petty and Anderson 1995, Petty 1996). In Kielder Forest the goshawk population recovered rapidly after the first recorded breeding attempt in 1973, and 25-33 goshawk home-ranges are now occupied (see Appendix S1; Petty & Anderson 1995). Such a large increase in goshawk abundance, was presumably concomitant with an increase in intraspecific competition for food and nest sites. However, goshawks may also have become food-limited because of a long-term decline in the abundance of red grouse (*Lagopus lagopus*) and a substantial decline of feral pigeon (*Columba livia*) in recent years in England (Robinson et al. 2015), as both species are important prey for goshawks in our study area (Petty et al. 2003a). The term feral pigeon includes both racing and homing pigeons.

The first aim of this study was to quantify the extent of superpredation and then to test the prediction that there has been a per capita increase in superpredation during the colonisation process by examining goshawk dietary data. Our second aim was to determine whether superpredation was impacting local populations of the most commonly preyed upon mesopredator species, for which local population trends are well characterised. We then assessed whether the goshawk population had become food-limited as the population recovered. It is difficult to directly assess food limitation for generalist predators, such as goshawks, without comprehensive prey abundance surveys (Rutz and Bijlsma 2006). Consequently, we examined two different lines of evidence to proximately assess food

limitation. First, we examined temporal variation in goshawk diet to determine whether there had been any decline in the contribution of certain prey species/groups known to be important for goshawks. We then examined changes in the reproductive success of the goshawk population, because reproductive success is closely associated with food availability in goshawks and other raptor species (Newton 1979, 1998, Rutz and Bijlsma 2006, Millon et al. 2008). Lastly, we evaluated evidence supporting the alternative hypotheses of superpredation by examining which mesopredator species were being killed by goshawks (e.g. were they known to compete with goshawks for food or nesting sites).

## **METHODS**

Kielder Forest is situated in Northumberland, in the north of England, adjacent to the border with Scotland. For a map of the study area see (Petty et al. 2003b). Each year active goshawk home-ranges were located by searching suitable nesting habitat within the forest (between the end of February and end of the breeding season). The locations of active nests were recorded and these sites were then visited at least four times to establish whether a breeding attempt took place, to record the number of chicks that fledged and collect dietary data.

### **Quantifying superpredation**

To quantify superpredation and to determine whether there had been a per capita increase in superpredation as the goshawk population expanded, we used goshawk dietary data. Specifically, we quantified the proportion of goshawk diet comprised of other raptor species each year. Here we use the term raptor to refer to all diurnal and nocturnal birds of prey. Goshawk diet was characterised by searching for the remains of prey (feathers, bones and fur) in the area surrounding active nest sites during nest visits between March-August, 1975-2014 (except between 1999-2001), in the same way as described in Petty *et al.* (2003). When possible,

at the end of the breeding season the top layer of nesting material was removed from active nests and searched for additional prey remains. Prey remains were removed or buried to avoid double counting in subsequent searches. We were able to identify 7763 prey items to species level by comparison with reference collections. It was not always possible to differentiate carrion crow (*Corvus corone*) from rook (*C. frugilegus*) remains. Therefore crow/rook refers to the abundance of both species in the diet, although rooks were scarce in the study area. We identified and quantified the minimum number of individuals of medium to large prey species by counting skeletal remains, whereas small avian prey (less than 100g) were identified and quantified by plucked feathers. Collecting and quantifying dietary data in this way is likely to underestimate the contribution of small prey items (Ziesemer 1983). However, this should not influence the results of our analyses as such items are relatively unimportant to goshawk diet in terms of biomass.

Once the proportion of goshawk diet comprising raptors had been calculated for each nest site/year, we examined how it varied in relation to the number of occupied goshawk home-ranges (measured as a continuous variable) using generalised linear mixed effect models (GLMM) with a binomial error structure fitted using the *lme4* package (Bates et al. 2015). Goshawk diet has previously been shown to change with altitude, presumably reflecting changes in abundance and diversity of prey species at different altitudes (Marquiss and Newton 1982, Toyne 1998). Consequently, we also examined whether the contribution of raptors to diet varied with the altitude of the goshawks nest site. Goshawk home-ranges were grouped into three altitudinal bands as follows: low, if the nest site was 225m or below, medium if between 226-354m, and high if 355m or above. We used these cut-offs because goshawk home-ranges above 355m were generally surrounded by open moorland habitat, whereas home-ranges below 225m were surrounded by forest, pasture and water (streams, rivers and a large



reservoir). The identity of home-ranges and year were both fitted as random effects to account for variation in diet between years and between home-ranges. Model selection was based on Akaike's information criterion corrected for small sample size (AICc) and AICc weights ( $W$ ; Burnham and Anderson 2002). The best performing model will have a  $\Delta AIC$  of zero, because  $\Delta AICc$  is the AICc for the model of interest minus the smallest AICc for the set of models being considered. Models are generally considered inferior if they have a  $\Delta AICc > 2$  units. AICc weights ( $w$ ) are an estimate of the relative likelihood of a particular model for the set of models being considered. Model assumptions were validated by visually inspecting residual plots; but they did not reveal any obvious nonlinear relationships, unless otherwise mentioned. Correlograms (with a lag distance up to 10km) were used to check for spatial-autocorrelation in the residuals of the best performing model. However, we found no evidence of spatial-autocorrelation in this, nor any other analyses of goshawk diet.

### **Impact on local mesopredator populations**

To assess the impact of changes in goshawk predation on local populations of the three most frequently preyed upon raptor species Eurasian kestrels (*Falco tinnunculus*), tawny owls (*Strix aluco*) and Eurasian sparrowhawks (*Accipiter nisus*; Petty *et al.* 2003), we first examined whether the proportion of goshawk diet comprising these three raptor species changed with goshawk abundance (measured as a continuous variable) using the same GLMM approach described above. We then used dietary data to estimate the minimum number of each mesopredator species killed each year by goshawks when  $\leq 14$ , 15-24 and 25+ goshawk home-ranges were occupied (for methods see Appendix 2). We used these three goshawk abundance categories to keep broadly similar sample sizes despite large variation in the number of prey remains recovered each year (range 10- 678). Temporal variation in predation rates on kestrel, tawny owl and sparrowhawk were then compared to changes in the local population dynamics

of these species. Annual counts of territorial kestrel pairs in and around the forest have been recorded since 1975 as part of a larger study on merlins (*Falco columbarius*; Newton, Meek & Little 1986; Little, Davison & Jardine 1995). Breeding tawny owls have been monitored continuously in a subsection of the forest since 1979 (Petty 1992, Petty et al. 1994, Hoy et al. 2015). The number of occupied sparrowhawk territories in a subsection of the study area has been recorded since 1974 (Petty 1979; Petty *et al.* 1995).

## **Assessing food limitation**

### *Declines in important prey*

To indirectly infer if the goshawk population had become food-limited we assessed whether there had been any decline in the contribution of important prey species/groups in the diet. We first examined how the dominance of main prey species in the diet changed as the goshawk population expanded. This was done by ranking species from most to least important, firstly in terms of their frequency contribution to diet and then in terms of their biomass when the number of occupied goshawk home-ranges was  $\leq 14$ , 15-24 and 25+. A full list of species killed by goshawk and their mean body mass values used in biomass calculations can be found in Appendix 3. Certain taxonomic groups are known to be important to goshawk diet. For example, *Columbiformes* are an important prey for goshawks across most of Europe, comprising up to 69% of all prey items (reviewed in Rutz *et al.* 2006). Whereas, *Tetraonidae* comprised almost 80% of goshawk diet in some years at northerly latitudes, and in southern Europe *Lagomorphs* were an important prey source (reviewed in Kenward 2006). We therefore categorised prey into taxonomic groups as follows: raptors, pigeons (*Columbiformes*), corvids (*Corvidae*), game birds (*Tetraonidae* and *Phasianidae*) mammals (mainly *Lagomorpha*, and *Sciuridae*), and ‘other’. This ‘other’ group largely consists of passerines but also includes other prey species, which are only occasionally taken. We then estimated both the frequency and

biomass contribution of these groups to goshawk diet and examined how the frequency contribution varied in relation to goshawk abundance (measured as a continuous variable) using the same GLMM approach. We were unable to assess whether variation in the proportion of goshawk diet comprised of these different prey species/groups were related to changes in the abundance of these prey species/groups as local population trends were not available. Lastly, because diet diversity had generally been observed to increase in other raptor populations as they became food-limited (Rutz and Bijlsma 2006, Lourenço et al. 2011a), we also examined how prey diversity changed with goshawk abundance using estimates of the Shannon-Wiener diversity index when  $\leq 14$ ; 15-24; 25+ goshawk home-ranges were occupied.

#### *Goshawk reproductive success*

We also indirectly assessed whether the goshawk population had become food-limited by examining how goshawk reproductive success varied in relation to the number of occupied home-ranges (measured as a continuous variable). In this analysis, we used two different measures of reproductive success: the average number of chicks fledged per successful breeding attempt and the proportion of breeding attempts which failed. We did not analyse variation in goshawk reproductive success prior to 1977 because goshawks did not reproduce successfully until a few years after the first home-ranges became established. Because the relationship between goshawk abundance and the number of chicks fledged per successful breeding attempt appeared to be non-linear we used generalised additive models (GAM) to characterise this relationship, fitted using the *mgcv* package (Wood 2015). In contrast, the relationship between the proportion of failed goshawk breeding attempts and goshawk abundance could be adequately characterised by generalised linear models (GLM) with a binomial error structure. All analyses were carried out in R version 3.0.3 (R Core Development Team 2015). Descriptive statistics are presented as the mean  $\pm$  1SD.

## RESULTS

### Superpredation increased during the colonisation process

Overall, raptors comprised 6% of all identifiable prey killed by goshawks ( $N=7763$ ) and represented 4% of goshawk prey in terms of biomass (Appendix 4). There was a per capita increase in superpredation as goshawks recovered, with the proportion of goshawk diet comprised of raptors increasing from 2% to 8% as the number of goshawk home-ranges increased from  $\leq 14$  to  $\geq 25$ . However, the proportion of raptors in goshawk diet was best modelled by an interaction between goshawk abundance and the altitude of the goshawk home-range (Table 1). The contribution of raptors to goshawk diet increased with goshawk abundance in home-ranges at the two lower elevations bands ( $\leq 225\text{m}$  and  $226\text{--}354\text{m}$ ). However, there was no significant change in the proportion of raptors in the diet at higher altitudes (e.g., above  $350\text{m}$ ), where the contribution of raptors to goshawk diet was highest (Fig. 1a).

### Impact on local mesopredator populations

Kestrels and tawny owls were both ranked within the 10 most important prey species, both in terms of their frequency and biomass contribution to diet (Table 2). Kestrels were the most commonly preyed raptor species, representing almost half (49%) of all raptors killed by goshawks ( $N=465$ ; Appendix 5). However, this proportion declined from 55% to 39% as the number of occupied goshawk home-ranges increased from  $<15$  to  $>24$  (Appendix 5). Kestrels contributed most to goshawk diet in high altitude home-ranges (Fig. 2a). The number of kestrels estimated to be killed each year by the goshawk population initially increased with goshawk abundance, from 14 [11–18 95% CI] when fewer than 15 goshawk home-ranges were occupied to 223 [197–248 95% CI] when 15–24 home-ranges were occupied. However, it then declined to 176 [154–198 95% CI] when more than 24 goshawk home-ranges were occupied

(Table 3). At the same time, there has been a substantial decline in the number of kestrel pairs recorded breeding in the study site. For example, there were 29 breeding pairs in 1981 compared to only five pairs in 2014.

Tawny owls then sparrowhawks were the next most commonly preyed upon raptor species, representing 23% and 10% respectively of all raptors killed by goshawks (Appendix 5). The contribution of both tawny owls and sparrowhawks in goshawk diet increased as the number of goshawk home-ranges increased; however, there was no evidence to suggest that it varied with altitude (Table 1; Fig.2b-c). The rank order importance of tawny owls to goshawk diet also increased from 9 to 7 as the number of occupied goshawk home-ranges increased from 15-24 to  $\geq 25$  (Table 2). Our estimates suggested there was huge increase in the number of tawny owls killed by goshawks each year, from an average of 5 [3-8 95% CI] to 159 owls [141-176 95% CI] as the number of occupied goshawk home-ranges increased from <15 to >24 (Table 3). The number of sparrowhawks killed by goshawks was also estimated to increase, from 1 [1-2 95% CI] to 53 [44-61 95% CI] as the number of occupied goshawk home-ranges increased from <15 to >24 (Table 3). Despite the estimated increase in predation on both tawny owls and sparrowhawks, there was no evidence to suggest that local populations had declined. That is, there was little interannual variation in the number of occupied tawny owl territories, which averaged  $56 \pm 4.07$  between 1985-2014 (Hoy et al. 2015), and sparrowhawks were known to occupy 7-14 home-ranges between 1974-1979 (Petty 1979) and 7-16 home-ranges between 2002-2014 (unpublished data).

## **Assessing food limitation**

### *Declines in important prey species/groups*

Almost half (48%) of all identifiable prey items were pigeons (Appendix 4). Wood pigeon (*Columba palumbus*) then feral pigeon were the two commonest prey species, irrespective of the number of goshawk home-ranges occupied (Table 2). The proportion of pigeons in goshawk diet declined over the study period as goshawk abundance increased, irrespective of home-range altitude (Fig. 1b; Table 4). For example, the biomass contribution of pigeons to diet declined from 52% to 40% as the number of home-ranges increased from <15 to >24. This decline in pigeon in goshawk diet appeared to be driven by a decrease in feral rather than wood pigeons. The contribution of pigeons to goshawk diet was lowest in higher altitude home-ranges, where moorland habitat predominated (Fig 1b; Table 4).

Crow/rook, red grouse and rabbit consistently ranked within the top-5 most important prey species, both in terms of biomass and frequency contribution to diet, irrespective of the number of goshawk home-ranges occupied (Table 2). The proportion of corvids and mammals in the diet increased with goshawk abundance (Fig 2c-d). For example, corvids and mammals comprised 11% and 4% respectively of diet (in terms of frequency) when <15 home-ranges were occupied, but 19% and 8% of diet respectively when >24 home-ranges were occupied (Appendix 4). In contrast, the contribution of game birds (including red grouse) declined as goshawk abundance increased, in all three altitudinal categories (Table 4; Fig. 2e). Although the dietary contribution of corvids and mammals did not vary with altitude, the proportion of game birds (especially red grouse) was noticeably higher for high altitude home-ranges (e.g., above 350m; where moorland habitat is more common). Goshawk diet also became more diverse as goshawk abundance increased, with the Shannon-Wiener diversity index increasing by 24% from 2.1 to 2.6 when the number of occupied goshawk territories increased from being <15 to >24.

### *Declines in reproductive success*

Overall the reproductive success of goshawks declined as the number of occupied home-ranges increased. This decline appeared to be driven by both a decline in the average number of chicks fledging per successful breeding attempt and an increase in the number of nesting attempts failing (Fig. 3). The number of chicks fledging per successful breeding attempt decreased from an average of  $2.90 \pm 0.24$  chicks to  $2.29 \pm 0.36$  chicks as the number of occupied goshawk home-ranges increased from  $<15$  to  $>24$  (Fig. 3a). The proportion of successful breeding attempts declined from an average of  $0.81 \pm 0.18$  to  $0.58 \pm 0.14$  when the number of occupied home-ranges increased from  $< 15$  to  $> 24$  (Fig. 3b).

## **DISCUSSION**

### **Superpredation has increased during the recolonization process**

The amount of superpredation in our study site, particularly in recent years, is noticeably higher than recorded elsewhere. For example, raptors comprised up to 8% of goshawk diet in Kielder Forest when goshawk abundance was high (Appendix 4) which is higher than the average of 2% estimated in a review of goshawk diet in Europe (Rutz et al. 2006, Lourenço et al. 2011a). Whilst many other studies provide a snapshot indication of the frequency of superpredation in a given system, relatively few have documented temporal variation in the frequency of superpredators killing other predators (but see Serrano 2000, Rutz and Bijlsma 2006), particularly in a recovering superpredator population. That there was a per capita increase in superpredation as the goshawk population recovered (Fig. 1a) has potentially important implications for conservation and management, because similar increases in superpredation may be expected in other superpredator populations currently recolonising former ranges in both North America and Europe (Maehr et al. 2001, Deinet et al. 2013, Chapron et al. 2014). For example, if increases in superpredation negatively affect the dynamics of mesopredator

species which are also of conservation concern it could lead to management conundrums for conservation projects aimed at restoring apex predator populations. However, it is important to note that when apex predator populations were reduced or extirpated, many previously suppressed mesopredator populations increase dramatically (Soulé et al. 1988, Crooks and Soulé 1999, Ritchie and Johnson 2009). Consequently, any declines in mesopredator populations which accompany the restoration of large predators may represent the reversal of this “mesopredator release” process (i.e., mesopredator suppression), rather than a shift to a new state.

#### **Impact on local mesopredator populations**

It’s important to note that our calculations of how many kestrels, tawny owls and sparrowhawks were killed by goshawks each year is not only likely to include breeding birds (i.e. the ones on which local population counts were based) but also non-breeders (e.g., “floaters”), individuals migrating through the study site (in the case of kestrels) and immigrants from neighbouring populations. Nevertheless, the large increase in the number of kestrels being killed each year (from an estimated 14 to 176; Table 3) coincided with a decline in the local kestrel population, which is consistent with an increase in goshawk predation having a negative impact upon the local kestrel population. However, the decline in kestrels could also be partly related to other factors, such as habitat changes or a decline in the amplitude of field vole (*Microtus agrestis*) population cycles in the study area (Cornulier et al. 2013), as they are the main prey for kestrels in our study site. Nevertheless, our results suggest that goshawks were killing a progressively greater proportion of a declining kestrel population, which may have contributed to the study area becoming a sink habitat, as previously suggested by Petty et al. (2003a).



In contrast, local breeding population of tawny owls and sparrowhawks did not decline over the study period, despite the substantial increase in the number killed each year by goshawks (Table 3; Fig. 2). This suggests that goshawk predation on tawny owls and sparrowhawks is compensatory rather than additive. Indeed, the impact of goshawk predation on the local tawny owl population is likely to be mitigated by goshawks selectively killing individuals with low reproductive values (e.g. juveniles and old owls, which have a lower probability of surviving and reproducing than prime-aged adults Millon et al. 2010, 2011), thus reducing the overall impact of predation at the population level (Hoy et al. 2015). Another factor which may be compensating for the increase in goshawk predation on tawny owls is the increase in immigrants entering the local population in recent years (Millon et al. 2014). Hence, goshawk predation may also have led to Kielder becoming a sink habitat for tawny owls. Unfortunately, we do not have equivalent data for sparrowhawks to be able to evaluate whether changes in immigration and/or selective predation of individuals with low reproductive values was also mitigating the impact of increased goshawk predation.

#### **Goshawks have become food-limited**

The substantial decline of two main prey groups (pigeon and game birds) in goshawk diet, the increase in diet diversity and decrease in goshawk reproductive success are all consistent with the goshawk population becoming increasingly food-limited as the population increased. The decline of pigeon and game birds in goshawk diet over the study period (Figs. 2b and 2c), presumably reflected a decline in the availability of two important prey species, namely feral pigeon and red grouse. Although, we cannot directly compare observed changes in the prevalence of feral pigeon and red grouse in diet to changes in their abundance (because regional population trends are not available), there is indirect evidence to suggest that the availability of these two prey species has declined. At a national level there has been a long-

term decline in red grouse populations in England, and feral pigeon populations are also thought to have declined by 26% since 1995 (Robinson et al. 2015). The decline in feral pigeons in goshawk diet may also be related to a decline in the number of stray racing pigeons entering the forest, because of a sustained decrease in the number of people participating in pigeon racing since the late 1980's (RPRA 2012). Furthermore, there is also anecdotal evidence of a local decrease in the abundance of red grouse and their main habitat over the study period (M.D., personal observation).

The increase in diet diversity we observed also indirectly supports the notion that there has been a decline in the availability of important prey, because when such food become scarce, predators are forced to switch to alternative species to make up the shortfall. Indeed, diet diversity was found to be negatively related to the abundance of important prey species for both goshawks (Rutz and Bijlsma 2006), sparrowhawks (Millon et al. 2009) and Eagle owls (*Bubo bubo*; Serrano 2000). Thus, together our results are consistent with goshawks switching to alternative prey species (raptors, corvids and mammals) as the availability of preferred prey species (e.g. feral pigeons and grouse) declined and they become food limited.

The decline in goshawk reproductive success as goshawk numbers increased (Fig. 3) provided additional and independent evidence that goshawks became food-limited given that goshawk reproductive success is known to be positively related to food availability (Rutz and Bijlsma 2006). A decline in reproductive success could also arise if goshawks had smaller home-ranges in high density years. However, this seems unlikely given that the average distance between goshawk nest sites has varied little since the mid-80's (mean distance between nest sites = 3.97 km  $\pm$  0.43; coefficient of variation = 0.11). A decline in reproductive success with increasing density could also arise if individuals establishing home-ranges in later years were forced to

settle in “poor-quality sites” because all the “good-quality sites” were already occupied (Rodenhuse et al. 1997). Although the biggest decline in reproductive success was observed in territories established towards the end of the study period, reproductive success also declined in territories established in the early and mid-part of the study period (Appendix 6). One likely reason for the decline in reproductive success, hence food availability being population wide (rather than restricted to certain home-ranges), may be because goshawk hunting ranges overlap (Kenward 2006), such that individuals nesting in the later established “poor-quality sites” may still forage and deplete prey in areas used for hunting by birds nesting in “good-quality site”.

### **Mechanisms underlying superpredation**

The increase in the proportion of raptors in the diet as goshawk abundance increased (Fig. 1), viewed in combination with the results of other analyses is consistent with predictions for the food-limitation hypothesis of superpredation. That is, as the availability of preferred prey (pigeon and grouse) declined goshawks appear to have switched to alternative, less profitable prey species, such as raptors. Furthermore, predictions of alternative hypotheses were not supported by our data. For example, if superpredation was purely opportunistic then changes in the frequency of mesopredator species in the diet are expected to reflect changes in mesopredator abundance (Polis et al. 1989). However, the contribution of kestrels to goshawk diet was higher in the later part of the study, despite kestrels declining. Furthermore, only two buzzards were known to have been killed by goshawks, despite a substantial increase in the abundance of buzzards in the forest since 1995 (over 80+ home-ranges now occupied). Our results also do not provide support for the predator-removal hypothesis given that the raptor species killed by goshawks were of no or little threat either to adult or juvenile goshawks (Appendix 5). Support for the competitor removal-hypothesis is also lacking because the

majority (83%) of raptors killed by goshawks are unlikely to compete with goshawks for food as they are largely dependent on field voles (e.g., kestrels and tawny owls; Appendix 5), yet voles only make up 0.06% of goshawk diet in terms of biomass. Furthermore, buzzards were seldom preyed upon by goshawks, yet they are known to compete with goshawks for nest sites and kill some of the same species as goshawks (Bijlsma 1994, Krüger 2002a, b). We therefore conclude that food limitation is the most likely driver of superpredation in this system given:

- 1) the decline in two main prey groups, the increase in diet diversity and the decrease in goshawk reproductive success suggest that the goshawk population has become food-limited;
- 2) superpredation does not appear to be purely opportunistic, given that variation in goshawk predation on different raptor species did not mirror local mesopredator population trends; and
- 3) the species of mesopredator killed offer little support for either the predator- or competitor-removal hypotheses of superpredation.

## CONCLUSIONS

Here we have provided evidence to show how superpredation varied over time in a recovering population of an apex predator, the northern goshawk. Our results suggest that increasing rates of superpredation were a response to declining food availability (pigeon and grouse) linked to increasing goshawk numbers. Thus, this study offers insights into the mechanisms driving variation in superpredation. We found evidence suggesting that an increase in goshawk predation may be contributing to a decline in the most frequently preyed mesopredator, Eurasian kestrels, a species which is also of conservation concern nationally. Thus, our results indicate that superpredation is likely to be an important factor to consider when developing conservation and management strategies for mesopredator species in the future. However, rather than a shift to a new alternative state, we suggest that the decline in kestrel numbers (and their likely persistence in refuges/areas with lower superpredator abundance) possibly

represents a reversal of a mesopredator release process (i.e. mesopredator suppression) following the extirpation of goshawks, and decline in other larger raptor species in the UK. Thus the results presented here may also offer insights into how other raptor communities will change in areas where goshawks are starting to recover. Lastly, in addition to the direct effect that an increase in predation can have on mesopredator population dynamics by increasing mortality rates, it is also important to consider that recovering superpredator populations may also be influencing mesopredator dynamics by negatively affecting mesopredator reproduction success. For example, mesopredators are more likely to abandon breeding attempts when superpredator densities are high (Mueller et al. 2016, Hoy et al. 2016).

## ACKNOWLEDGEMENTS

We are grateful to R. Lourenço and A.K. Mueller for their helpful comments. We thank Forest Research for funding all fieldwork on goshawks during 1973-1996, Forest Enterprise for funding fieldwork after 1998 and T. Dearnley and N. Geddes for allowing and facilitating work in Kielder Forest. This work was also partly funded by a Natural Environment Research Council studentship NE/J500148/1 to SH and a grant NE/F021402/1 to XL and by Natural Research. We thank I. Yoxall and B. Little for the data they collected and their contributions to this study. Lastly, we thank English Nature and the British Trust for Ornithology for kindly issuing licences to monitor goshawk nest sites.

## REFERENCES

- Bates, D., Maechler, M., Bolker, B. and Walker, S. 2015. Fitting Linear Mixed-Effects Models Using lme4. - J. Statistical Softw. in press.
- Berger-Tal, O., Mukherjee, S., Kotler, B. P. and Brown, J. S. 2009. Look before you leap: Is risk of injury a foraging cost? - Behav. Ecol. Sociobiol. 63: 1821–1827.
- Bijlsma, R. G. 1994. Voedselkeus van Havik *Accipiter gentilis*, Sperwer *A. nisus* en Buizerd *Buteo buteo* in de Flevopolders. - Tak. 2: 22–35.
- Burnham, K. P. and Anderson, D. R. 2002. Model Selection and Multimodel Inference: a

- 490 Practical Information-theoretic Approach. - Springer-Verlag.
- 491 Caro, T. M. and Stoner, C. J. 2003. The potential for interspecific competition among African  
492 carnivores. - Biol. Conserv. 110: 67–75.
- 493 Chakarov, N. and Krüger, O. 2010. Mesopredator release by an emergent superpredator: a  
494 natural experiment of predation in a three level guild. - PLoS One 5: e15229.
- 495 Chapron, G., Kaczensky, P., Linnell, J. D. C., von Arx, M., Huber, D., Andrén, H., López-  
496 Bao, J. V., Adamec, M., Álvares, F., Anders, O., Balčiauskas, L., Balys, V., Bedő, P.,  
497 Bego, F., Blanco, J. C., Breitenmoser, U., Brøseth, H., Bufka, L., Bunikyte, R., Ciucci,  
498 P., Dutsov, A., Engleder, T., Fuxjäger, C., Groff, C., Holmala, K., Hoxha, B., Iliopoulos,  
499 Y., Ionescu, O., Jeremić, J., Jerina, K., Kluth, G., Knauer, F., Kojola, I., Kos, I., Krofel,  
500 M., Kubala, J., Kunovac, S., Kusak, J., Kutal, M., Liberg, O., Majić, A., Männil, P.,  
501 Manz, R., Marboutin, E., Marucco, F., Melovski, D., Mersini, K., Mertzanis, Y.,  
502 Mysłajek, R. W., Nowak, S., Odden, J., Ozolins, J., Palomero, G., Paunović, M.,  
503 Persson, J., Potočník, H., Quenette, P.-Y., Rauer, G., Reinhardt, I., Rigg, R., Ryser, A.,  
504 Salvatori, V., Skrbinšek, T., Stojanov, A., Swenson, J. E., Szemethy, L., Trajçe, A.,  
505 Tsingarska-Sedefcheva, E., Váňa, M., Veeroja, R., Wabakken, P., Wölfl, M., Wölfl, S.,  
506 Zimmermann, F., Zlatanova, D. and Boitani, L. 2014. Recovery of large carnivores in  
507 Europe's modern human-dominated landscapes. - Science (80-. ). 346: 1517–1519.
- 508 Cornulier, T., Yoccoz, N. G., Bretagnolle, V., Brommer, J. E., Butet, A., Ecke, F., Elston, D.  
509 a, Framstad, E., Henttonen, H., Hörnfeldt, B., Huitu, O., Imholt, C., Ims, R. a, Jacob, J.,  
510 Jędrzejewska, B., Millon, A., Petty, S. J., Pietiäinen, H., Tkadlec, E., Zub, K. and  
511 Lambin, X. 2013. Europe-wide dampening of population cycles in keystone herbivores.  
512 - Science (80-. ). 340: 63–6.
- 513 Crooks, K. R. and Soulé, M. E. 1999. Mesopredator release and avifaunal extinctions in a  
514 fragmented system. - Nature 400: 563–566.
- 515 Deinet, S., Ieronymidou, C., McRae, L., Burfield, I. J., Foppen, R. P., Collen, B. and Böhm,  
516 M. 2013. Wildlife comeback in Europe: The recovery of selected mammal and bird  
517 species.
- 518 Hoy, S. R., Petty, S. J., Millon, A., Whitfield, D. P., Marquiss, M., Davison, M. and Lambin,  
519 X. 2015. Age and sex-selective predation as moderators of the overall impact of  
520 predation. - J. Anim. Ecol. 84: 692–701.
- 521 Hoy, S. R., Millon, A., Petty, S. J., Whitfield, D. P. and Lambin, X. 2016. Food availability  
522 and predation risk, rather than intrinsic attributes, are the main factors shaping the  
523 reproductive decisions of a long-lived predator. - J. Anim. Ecol. 85: 892–902.
- 524 Kenward, R. E. 2006. The Goshawk. - T & A D Poyser.
- 525 Krüger, O. 2002a. Interactions between common buzzard *Buteo buteo* and goshawk *Accipiter*  
526 *gentilis*: trade-offs revealed by a field experiment. - Oikos 96: 441–452.
- 527 Krüger, O. 2002b. Analysis of nest occupancy and nest reproduction in two sympatric  
528 raptors: common buzzard *Buteo buteo* and goshawk *Accipiter gentilis*. - Ecography  
529 (Cop.). 25: 523–532.
- 530 Little, B., Davison, M. and Jardine, D. 1995. Merlins *Falco columbarius* in Kielder Forest:  
531 influences of habitat on breeding performance. - For. Ecol. Manage. 79: 147–152.
- 532 Lourenço, R., Santos, S. M., Rabaça, J. E. and Penteriani, V. 2011a. Superpredation patterns  
533 in four large European raptors. - Popul. Ecol. 53: 175–185.
- 534 Lourenço, R., Penteriani, V., Delgado, M. D. M., Marchi-Bartolozzi, M. and Rabaça, J. E.

535 2011b. Kill before being killed: an experimental approach supports the predator-removal  
536 hypothesis as a determinant of intraguild predation in top predators. - Behav. Ecol.  
537 Sociobiol.: 1709–1714.

538 Maehr, D. S., Noss, R. F. and Larkin, J. L. 2001. Large Mammal Restoration: Ecological and  
539 Sociological Challenges in the 21st Century. - Island Press.

540 Marquiss, M. and Newton, I. 1982. The goshawk in Britain. - Br. Birds 75: 243–260.

541 Millon, A., Arroyo, B. E. and Bretagnolle, V. 2008. Variable but predictable prey availability  
542 affects predator breeding success: natural versus experimental evidence. - J. Zool. 275:  
543 349–358.

544 Millon, A., Nielsen, J. T., Bretagnolle, V. and Møller, A. P. 2009. Predator-prey relationships  
545 in a changing environment: The case of the sparrowhawk and its avian prey community  
546 in a rural area. - J. Anim. Ecol. 78: 1086–1095.

547 Millon, A., Petty, S. J. and Lambin, X. 2010. Pulsed resources affect the timing of first  
548 breeding and lifetime reproductive success of tawny owls. - J. Anim. Ecol. 79: 426–435.

549 Millon, A., Petty, S. J., Little, B. and Lambin, X. 2011. Natal conditions alter age-specific  
550 reproduction but not survival or senescence in a long-lived bird of prey. - J. Anim. Ecol.  
551 80: 968–975.

552 Millon, A., Petty, S. J., Little, B., Gimenez, O., Cornulier, T. and Lambin, X. 2014.  
553 Dampening prey cycle overrides the impact of climate change on predator population  
554 dynamics: A long-term demographic study on tawny owls. - Glob. Chang. Biol. 20:  
555 1770–1781.

556 Mueller, A.-K., Chakarov, N., Heseker, H. and Krüger, O. 2016. Intraguild predation leads to  
557 cascading effects on habitat choice, behaviour and reproductive performance (J Quinn,  
558 Ed.). - J. Anim. Ecol. 85: 774–784.

559 Newton, I. 1979. Population ecology of raptors. - T. and A.D. Poyser.

560 Newton, I. 1998. Population limitation in birds. - Academic Press.

561 Newton, I., Meek, E. and Little, B. 1986. Population and breeding of Northumbrian Merlins.  
562 - Br. Birds 79: 155–170.

563 Paine, R. T. 1980. Food webs: linkage, interaction strength, and community infrastructure. -  
564 J. Anim. Ecol. 49: 667–685.

565 Palomares, F. and Caro, T. M. 1999. Interspecific killing among mammalian carnivores. -  
566 Am. Nat. 153: 492–508.

567 Petty, S. J. 1979. Breeding biology of the sparrowhawk in Kielder Forest 1975-78. - In:  
568 Galloway, B. and Hodgson, M. S. (eds), Birds in Northumbria: 1978 Northumberland  
569 bird report. Tyneside bird club, pp. 92–109.

570 Petty, S. J. 1992. Ecology of the Tawny owl *Strix aluco* in the spruce forests of  
571 Northumberland and Argyll.

572 Petty, S. J. 1996. History of Northern Goshawk *Accipiter gentilis* in Britain. - In: Holmes, J.  
573 S. and Simons, J. R. (eds), The Introduction and Naturalisation of Birds. The Stationery  
574 Office, pp. 95–102.

575 Petty, S. J. and Anderson, D. I. K. 1995. Goshawks *Accipiter gentilis*. - In: Day, J. C. et al.  
576 (eds), The Atlas of Breeding Birds in Northumbria. Newcastle upon Tyne:  
577 Northumberland and Tyneside Bird Club, pp. 44–45.

578 Petty, S. J., Shaw, G. and Anderson, D. I. K. 1994. Value of nest boxes for population studies

579 and conservation of owls in coniferous forests in Britain. - J. Raptor Res. 28: 134–142.

580 Petty, S. J., Patterson, I. J., Anderson, D. I. K., Little, B. and Davison, M. 1995. Numbers,  
581 breeding performance, and diet of the sparrowhawk *Accipiter nisus* and merlin *Falco*  
582 *columbarius* in relation to cone crops and seed-eating finches. - For. Ecol. Manage. 79:  
583 133–146.

584 Petty, S. J., Anderson, D. I. K., Davison, M., Little, B., Sherratt, T. N., Thomas, C. J. and  
585 Lambin, X. 2003a. The decline of common kestrels *Falco tinnunculus* in a forested area  
586 of northern England: the role of predation by Northern Goshawks *Accipiter gentilis*. -  
587 Ibis (Lond. 1859). 145: 472–483.

588 Petty, S. J., Lurz, P. W. W. and Rushton, S. P. 2003b. Predation of red squirrels by northern  
589 goshawks in a conifer forest in northern England : can this limit squirrel numbers and  
590 create a conservation dilemma ? - Biol. Conserv. 111: 105–114.

591 Polis, G. A., Myers, C. A. and Holt, R. D. 1989. The ecology and evolution of intraguild  
592 predation: Potential Competitors That Eat Each Other. - Annu. Rev. Ecol. Syst. 20: 297–  
593 330.

594 R Core Development Team 2015. R Development Core Team. - R A Lang. Environ. Stat.  
595 Comput. . R Found. Stat. Comput. , Vienna, Austria. URL <http://www.R-project.org/>. in  
596 press.

597 Ripple, W. J. and Beschta, R. L. 2004. Wolves and the ecology of fear : can predation risk  
598 structure ecosystems ? - Bioscience 54: 755–766.

599 Ritchie, E. G. and Johnson, C. N. 2009. Predator interactions, mesopredator release and  
600 biodiversity conservation. - Ecol. Lett. 12: 982–998.

601 Robinson, R. A., Marchant, J. H., Leech, D. I., Massimino, D., Sullivan, M. J. P., Eglington,  
602 S. M., Barimore, C., Dadam, D., Downie, I. S., Hammond, M. J., Harris, S. J., Noble, D.  
603 G., Walker, R. H. and Baillie, S. R. 2015. BirdTrends 2015: trends in numbers, breeding  
604 success and survival for UK breeding birds. Research Report 678.

605 Rodenhouse, N. L., Sherry, T. W. and Holmes, R. T. 1997. Site-dependent regulation of  
606 population size: a new synthesis. - Ecology 78: 2025–2042.

607 RPRA 2012. Royal Pigeon Racing Association .: <http://www.rpra.org>.

608 Rutz, C. and Bijlsma, R. G. 2006. Food-limitation in a generalist predator. - Proc. R. Soc.  
609 Biol. Sci. 273: 2069–76.

610 Rutz, C., Bijlsma, R. G., Marquiss, M. and Kenward, R. E. 2006. Population limitation in the  
611 northern goshawk in Europe: a review with case studies. - Stud. Avian Biol.: 158–197.

612 Sergio, F. and Hiraldo, F. 2008. Intraguild predation in raptor assemblages: a review. - Ibis  
613 (Lond. 1859). 150: 132–145.

614 Serrano, D. 2000. Relationship between raptors and rabbits in the diet of eagle owls in  
615 southwestern Europe: competition removal or food stress? - J. Raptor Res. 34: 305–310.

616 Soulé, M. ., Bolger, D. ., Alberts, A. ., Wright, J., Sorice, M. and Hill, S. 1988. Reconstructed  
617 of rapid extinctions of dynamics birds in urban habitat islands. - Conserv. Biol. 2: 75–  
618 92.

619 Toyne, E. P. 1998. Breeding season diet of the Goshawk *Accipiter gentilis* in Wales. - Ibis  
620 (Lond. 1859). 140: 569–579.

621 Wood, S. 2015. Package: mgcv 1.8-9 Mixed GAM Computation Vehicle with  
622 GCV/AIC/REML Smoothness Estimation. - R 3.2.3



623 Ziesemer, F. 1983. Untersuchungen zum Einfluss des Habichts (*Accipiter gentilis*) auf  
624 Populationen seiner Beutetiere.  
625

Table 1. Model selection and parameters for the analysis of variation in the proportion of goshawk diet comprised of all raptor species and also kestrels, tawny owls and sparrowhawks separately. We examined whether diet varied in relation to goshawk abundance (number of home-ranges occupied) and the altitude of the goshawk home-range (e.g., below 225m, between 226-354m and above 355m). The most parsimonious model will have a  $\Delta\text{AICc} = 0$  and is highlighted in bold. The number of parameters estimated in each model is designated in the *np* column. AICc weights (*W*) are an estimate of the relative likelihood of a model.

Model	<i>np</i>	All raptors				Kestrel				Tawny owl				Sparrowhawk			
		Estimate	SE	$\Delta\text{AICc}$	<i>W</i>	Estimate	SE	$\Delta\text{AICc}$	<i>W</i>	Estimate	SE	$\Delta\text{AICc}$	<i>W</i>	Estimate	SE	$\Delta\text{AICc}$	<i>W</i>
1. Null	2			26.74	<0.01			11.87	<0.01			13.51	<0.01			9.36	0.01
2. Goshawk abundance (GA)	4	0.06	0.02	14.46	<0.01	0.02	0.02	12.53	<0.01	<b>0.11</b>	<b>0.03</b>	<b>0</b>	<b>0.84</b>	<b>0.12</b>	<b>0.04</b>	<b>0</b>	<b>0.61</b>
3. Altitude (226-354m)	5	-0.01	0.19	13.2	<0.01	-0.18	0.26	2.7	0.18	0.15	0.35	17.09	<0.01	-0.2	0.49	9.89	<0.01
Altitude (above 355m)		0.95	0.28			0.96	0.38			0.36	0.52			0.92	0.76		
4. Altitude (226-354m)	6	0.03	0.18	4.89	0.08	-0.15	0.25	3.94	0.1	0.22	0.34	3.66	0.13	-0.12	0.47	2.66	0.16
Altitude (above 355m)		0.84	0.27			0.95	0.38			0.14	0.5			0.58	0.72		
+ Goshawk abundance		0.05	0.02			0.02	0.02			0.11	0.03			0.11	0.04		
5. Altitude (226-354m)	8	<b>-0.87</b>	<b>0.68</b>	<b>0</b>	<b>0.92</b>	<b>-0.98</b>	<b>0.93</b>	<b>0</b>	<b>0.71</b>	-0.72	1.62	6.94	0.03	-4.18	1.94	2.07	0.22
Altitude (above 355m)		<b>2.57</b>	<b>1.01</b>			<b>3.18</b>	<b>1.26</b>			-2.98	3.73			-1.22	3.97		
Goshawk abundance		<b>0.03</b>	<b>0.03</b>			<b>0.01</b>	<b>0.04</b>			0.08	0.06			0.01	0.06		
Altitude (226-354m) x GA		<b>0.04</b>	<b>0.03</b>			<b>0.04</b>	<b>0.04</b>			0.04	0.06			0.16	0.08		
Altitude (above 355m) x GA		<b>-0.07</b>	<b>0.04</b>			<b>-0.09</b>	<b>0.05</b>			0.11	0.13			0.07	0.15		

Table 2. The 10 most important prey species in northern goshawk breeding season diet, ranked in order of decreasing importance in terms of both their frequency in the diet and biomass contribution to diet when the number of occupied goshawk home-ranges was estimated to be less than 14, between 15-24 and 25 or more.

Species rank	Frequency						Species rank	Biomass		
	<14 occupied goshawk home-ranges	<i>n</i>	15-24 occupied goshawk home-ranges	<i>n</i>	>25 occupied goshawk home-ranges	<i>n</i>		<14 occupied goshawk home-ranges	15-24 occupied goshawk home-ranges	>25 occupied goshawk home-ranges
1.	Feral pigeon	531	Wood pigeon	848	Wood pigeon	625	1.	Wood pigeon	Wood pigeon	Wood pigeon
2.	Wood pigeon	488	Feral pigeon	752	Feral pigeon	479	2.	Red grouse	Crow/rook	Crow/rook
3.	Red grouse	277	Crow/rook	597	Crow/rook	383	3.	Feral pigeon	Feral pigeon	Rabbit
4.	Crow/rook	149	Red grouse	221	Red grouse	133	4.	Crow/rook	Rabbit	Feral pigeon
5.	Jay	35	Kestrel	128	Rabbit	108	5.	Rabbit	Red grouse	Red grouse
6.	Song thrush	30	Rabbit	126	Red squirrel	89	6.	Pheasant	Pheasant	Pheasant
7.	Mistle thrush	27	Mistle thrush	85	Mistle thrush	88	7.	European hare	Kestrel	Tawny owl
8.	Field vole	25	Red squirrel	84	Jay	78	8.	Jay	Red squirrel	Red squirrel
9.	Kestrel	24	Song thrush	78	Kestrel	76	9.	Mallard	Tawny owl	Kestrel
10.	Red squirrel	22	Jay	61	Tawny owl	69	10.	Kestrel	Mistle thrush	Jay

Table 3. Estimated number of kestrels, tawny owls and sparrowhawks killed during the breeding season (March-August) each year by the Kielder Forest goshawk population when the number of occupied goshawk home-ranges was estimated to be less than 14, between 15-24 and 25 or more.

Species	Occupied goshawk home-ranges	Estimated % biomass of goshawk diet	Average number killed per pair	Mean number killed each year the by entire goshawk population	95% CI lower bound	95% CI upper bound
Kestrel	< 14	0.47	2.20	14	11	18
	15-24	2.29	10.23	223	197	248
	> 25	1.63	6.44	176	154	198
Tawny owl	< 14	0.34	0.70	5	3	6
	15-24	1.69	3.33	72	62	83
	> 25	3.32	5.79	159	141	176
Sparrowhawk	< 14	0.04	0.21	1	1	2
	15-24	0.37	1.66	36	28	45
	> 25	0.48	1.92	53	44	61

Table 4. Model selection and parameters for the analysis of variation in the proportion of goshawk diet comprised of different prey groups (pigeons, corvids, game birds, mammals and other). We examined whether diet varied in relation to goshawk abundance (number of home-ranges occupied) and the altitude of the goshawk home-range (e.g., below 225m, between 226-354m and above 355m). The most parsimonious model will have a  $\Delta AICc = 0$  and is highlighted in bold. The number of parameters estimated in each model is designated in the *np* column. *AICc* weights (*W*) are an estimate of the relative likelihood of a model.

Model	<i>np</i>	Pigeons				Corvids				Game birds				Mammals				Other			
		Estimate	SE	$\Delta AICc$	<i>W</i>	Estimate	SE	$\Delta AICc$	<i>W</i>	Estimate	SE	$\Delta AICc$	<i>W</i>	Estimate	SE	$\Delta AICc$	<i>W</i>	Estimate	SE	$\Delta AICc$	<i>W</i>
Null	2			16.82	<0.01			7.94	0.02			23.01	<0.01			2.60	0.19			6.77	0.03
Goshawk abundance (GA)	4	-0.02	0.01	10.49	<0.01	<b>0.04</b>	<b>0.01</b>	<b>0</b>	<b>0.8</b>	-0.03	0.01	9.66	0.01	<b>0.03</b>	<b>0.01</b>	<b>0</b>	<b>0.68</b>	<b>0.03</b>	<b>0.01</b>	<b>0</b>	0.77
Altitude (226-354m)	5	0.04	0.10	4.04	0.08	-0.11	0.12	10.96	<0.01	0.32	0.22	22.75	<0.01	-0.05	0.17	6.55	0.03	-0.05	0.13	9.21	0.01
Altitude (above 355m)		-0.42	0.15			-0.02	0.19			0.55	0.26			0.001	0.26			0.16	0.20		
Altitude (226-354m)	6	<b>0.03</b>	<b>0.10</b>	<b>0</b>	<b>0.62</b>	-0.09	0.12	3.55	0.14	<b>0.29</b>	<b>0.21</b>	<b>0</b>	<b>0.64</b>	-0.03	0.17	4.04	0.09	-0.04	0.13	3.20	0.16
Altitude (above 355m)		<b>-0.40</b>	<b>0.15</b>			-0.04	0.19			<b>0.85</b>	<b>0.26</b>			-0.06	0.26			0.12	0.20		
+ Goshawk abundance		<b>-0.02</b>	<b>0.01</b>			0.04	0.01			<b>-0.04</b>	<b>0.01</b>			0.03	0.01			0.03	0.01		
Altitude (226-354m)	8	-0.29	0.34	1.54	0.29	-0.12	0.42	5.96	0.04	0.30	0.60	1.15	0.36	0.01	0.64	7.76	0.01	0.32	0.49	6.10	0.04
Altitude (above 355m)		-1.29	0.60			0.77	0.71			-0.78	1.15			-0.83	1.39			1.03	0.89		
Goshawk abundance		-0.03	0.01			0.04	0.02			-0.04	0.02			0.03	0.03			0.05	0.02		
Altitude (226-354m) x GA		0.01	0.01			0.002	0.02			-0.001	0.03			-0.002	0.03			-0.01	0.02		
Altitude (above 355m) x GA		0.04	0.02			-0.03	0.03			0.06	0.04			0.03	0.05			-0.04	0.03		


## Figure Legends

Figure 1. Changes in the proportion of northern goshawk breeding season diet comprised of: a) raptors; b) pigeons; c) corvids; d) mammals; e) game birds and f) other species as the number of occupied goshawk home-ranges increased in Kielder Forest, UK. Error bars are the 95% confidence intervals.


Figure 2. Changes in the proportion of northern goshawk breeding season diet comprised of: a) kestrels; b) tawny owls; c) sparrowhawks as the number of occupied goshawk home-ranges increased in Kielder Forest, UK. Error bars are the 95% confidence intervals.

Figure 3. Inter-annual variation in goshawk reproductive success measured as: a) the number of chicks fledged per successful breeding attempt (line represents predicted values generated from a GAM) and b) the proportion of breeding attempts which were successful (line represents predicted values generated from a GLM) as the number of occupied goshawk home-ranges increased in Kielder Forest, UK.


a)




b)




c)




d)




e)




f)




a)





b)




c)







**Appendix 1:** The number of occupied northern goshawk home-ranges in Kielder Forest, UK



## **Appendix 2:** The average number of kestrels, tawny owls and sparrowhawks killed by the goshawk population each year

To estimate the average number of each species killed by the goshawk population each year, we first calculated the average number of each species killed per pair of goshawks, each year when 1-14, 15-24 and 25+ goshawk home-ranges were occupied, using the following equation taken from Petty *et al.* (2003).

$$IK = (CF + CM + CY) * (PT) / M$$

IK is the estimated number of individuals killed by a pair of goshawks between March and August (184 days). CF = estimated total food consumption of a female goshawk during the breeding season (189g of food per day \* 184 days). CM = total food consumption of a male goshawk during the breeding season (133g of food per day \* 184 days). The daily food consumption values used for male and female goshawk are the same as those used by Petty *et al.* (2003), originally calculated by Kenward *et al.* (1981). CY = total food consumption of young goshawks (i.e. offspring) during the breeding season (161g of food per day (CF+CM/2) \* 108 days \* mean fledged brood size of breeding pairs). The mean fledged brood size of goshawks was 2.19 in years when fewer than 15 home-ranges were occupied, 1.93 when 15-24 home-ranges were occupied and 1.31 when 25 or more home-ranges were occupied. The CY estimate assumes that young goshawks: 1) hatch around mid-May; 2) do not leave their natal territory until August; and 3) that juveniles have the same overall food intake as adults. Although young nestlings require less food than adults, older nestlings require more, such that when averaged over the entire period nestling food intake can be assumed to be equivalent to that of adults. M = average mass of the prey species. We used an average mass of 208g for kestrel (Ratcliffe 1993); 470g for tawny owl and 205g for sparrowhawk (Robinson 2005). PT = proportion biomass of the prey species in the diet. We used the dietary data to estimate of the proportion biomass of each of the three mesopredator species in goshawk diet for each of the three goshawk abundance categories (i.e. using pooled annual diet data collected when the number of occupied goshawk home-ranges was 1-14, 15-24 and 25+). This average proportion was then used in the above equation to calculate the number of individuals of each species killed during the breeding season by a goshawk pair. To get an estimate of the total number of each species killed each year by the entire goshawk population and how that has changed as the goshawk population increased in abundance, we multiplied our estimate of the number of individuals killed by a pair of goshawks (IK) by the average number of home-ranges occupied by goshawks for each of the goshawk abundance categories. The average number of home-ranges occupied in each goshawk abundance category was estimated to be 6.5, 21.75 and 27.38 when 1-14, 15-24 and 25+ goshawk home-ranges were occupied respectively.

**Appendix 3:** List of the species killed by northern goshawks in Kielder Forest, and the taxonomic prey group they were assigned to, along with the body mass used for each species to estimate their percentage biomass contribution to goshawk diet. We were not always able to differentiate between male and female prey remains, consequently we used the midpoint between the average mass for males and females in our biomass estimates. Body mass estimates for birds were obtained from the British Trust for Ornithology's website ([www.bto.org/birdfacts](http://www.bto.org/birdfacts)) and mass estimates for mammals were obtained from the British Mammal Societies website (<http://www.mammal.org.uk>).

Prey group	Common name	Mass (g)
Corvid	Carrion crow/rook ( <i>Corvus corone/C. frugilegus</i> )	510
Corvid	Eurasian jay ( <i>Garrulus glandarius</i> )	170
Corvid	Jackdaw ( <i>Corvus monedula</i> )	220
Corvid	Magpie ( <i>Pica pica</i> )	220
Corvid	Raven ( <i>Corvus corax</i> )	1200
Game	Black grouse ( <i>Tetrao tetrix</i> )	1065
Game	Pheasant ( <i>Phasianus colchicus</i> )	1190
Game	Red grouse ( <i>Lagopus lagopus scotica</i> )	600
Game	Red-legged partridge ( <i>Alectoris rufa</i> )	490
Mammal	Common shrew ( <i>Sorex araneus</i> )	9.5
Mammal	European hare ( <i>Lepus europaeus</i> )	3500
Mammal	European rabbit ( <i>Oryctolagus cuniculus</i> )	1600
Mammal	Field vole ( <i>Microtus agrestis</i> )	30
Mammal	Grey squirrel ( <i>Sciurus carolinensis</i> )	552.5
Mammal	Mole ( <i>Talpa europaea</i> )	100
Mammal	Pygmy shrew ( <i>Sorex minutus</i> )	4
Mammal	Rat ( <i>Rattus norvegicus</i> )	360
Mammal	Red squirrel ( <i>Sciurus vulgaris</i> )	200
Mammal	Stoat ( <i>Mustela erminea</i> )	266.25
Mammal	Weasel ( <i>Mustela nivalis</i> )	90.25
Other	Blackbird ( <i>Turdus merula</i> )	100
Other	Black-headed gull ( <i>Chroicocephalus ridibundus</i> )	290
Other	Blue tit ( <i>Cyanistes caeruleus</i> )	10.5
Other	Budgerigar ( <i>Melopsittacus undulatus</i> )	35
Other	Chaffinch ( <i>Fringilla coelebs</i> )	24
Other	Coal tit ( <i>Periparus ater</i> )	9
Other	Common frog ( <i>Rana temporaria</i> )	22.7
Other	Common gull ( <i>Larus canus</i> )	400
Other	Common lizard ( <i>Zootoca vivipara</i> )	4
Other	Common toad ( <i>Bufo bufo</i> )	55
Other	Crossbill ( <i>Loxia curvirostra</i> )	43
Other	Cuckoo ( <i>Cuculus canorus</i> )	120
Other	Curlew ( <i>Numenius arquata</i> )	985
Other	Domestic chicken ( <i>Gallus gallus domesticus</i> )	1900
Other	Eurasian bullfinch ( <i>Pyrrhula pyrrhula</i> )	21
Other	Fieldfare ( <i>Turdus pilaris</i> )	100

Prey group	Common name	Mass (g)
Other	Goldcrest ( <i>Regulus regulus</i> )	6
Other	Great spotted woodpecker ( <i>Dendrocopos major</i> )	85
Other	Great tit ( <i>Parus major</i> )	18.5
Other	Green woodpecker ( <i>Picus viridis</i> )	190
Other	Kittiwake ( <i>Rissa tridactyla</i> )	410
Other	Lapwing ( <i>Vanellus vanellus</i> )	230
Other	Lesser black-backed gull ( <i>Larus fuscus</i> )	830
Other	Lesser redpoll ( <i>Acanthis cabaret</i> )	11
Other	Mallard ( <i>Anas platyrhynchos</i> )	1090
Other	Meadow pipit/tree pipit ( <i>Anthus pratensis/A. trivialis</i> )	19
Other	Mistle thrush ( <i>Turdus viscivorus</i> )	130
Other	Moorhen ( <i>Gallinula chloropus</i> )	320
Other	Newt ( <i>Triturus vulgaris</i> )	30
Other	Oyster catcher ( <i>Haematopus ostralegus</i> )	540
Other	Pied wagtail ( <i>Motacilla alba</i> )	21
Other	Redshank ( <i>Tringa totanus</i> )	120
Other	Robin ( <i>Erithacus rubecula</i> )	18
Other	Siskin ( <i>Spinus spinus</i> )	15
Other	Skylark ( <i>Alauda arvensis</i> )	38.5
Other	Snipe ( <i>Gallinago gallinago</i> )	110
Other	Song thrush ( <i>Turdus philomelos</i> )	83
Other	Starling ( <i>Sturnus vulgaris</i> )	78
Other	Swallow ( <i>Hirundo rustica</i> )	18.5
Other	Teal ( <i>Anas crecca</i> )	330
Other	Tree creeper ( <i>Certhia familiaris</i> )	10
Other	Whinchat ( <i>Saxicola rubetra</i> )	17
Other	Willow warbler ( <i>Phylloscopus trochilus</i> )	10
Other	Woodcock ( <i>Scolopax rusticola</i> )	280
Pigeon	Collared dove ( <i>Streptopelia decaocto</i> )	200
Pigeon	Feral pigeon ( <i>Columba livia</i> )	300
Pigeon	Wood pigeon ( <i>Columba palumbus</i> )	450
Raptor	Barn owl ( <i>Tyto alba</i> )	300
Raptor	Common buzzard ( <i>Buteo buteo</i> )	890
Raptor	Common kestrel ( <i>Falco tinnunculus</i> )	208
Raptor	Long-eared owl ( <i>Asio otus</i> )	290
Raptor	Merlin ( <i>Falco columbarius</i> )	205
Raptor	Northern goshawk ( <i>Accipiter gentilis</i> )†	1000
Raptor	Short-eared owl ( <i>Asio flammeus</i> )	330
Raptor	Sparrowhawk ( <i>Accipiter nisus</i> )	205
Raptor	Tawny owl ( <i>Strix aluco</i> )	470

† Goshawk chicks were only included in the diet if there was evidence to suggest that it was a case of cannibalism rather than fledglings dying in the nest.

**Appendix 4:** Variation in the proportion of northern goshawk diet made up different prey groups as goshawk abundance increased. Total refers to all dietary data collected during the breeding season between 1973 and 2014, the other columns show estimates from dietary data collected in years when 1-14, 15-24 and 25 or more goshawk home-ranges were occupied.

Prey group	<i>n</i>				% Biomass				% Frequency			
	Total	1-14	15-24	25+	Total	1-14	15-24	25+	Total	1-14	15-24	25+
Pigeon (Columbidae)	3724	1019	1601	1104	43.37	51.80	41.75	39.82	47.97	57.18	46.61	43.36
Corvid (Corvidae)	1379	190	694	495	19.15	11.52	22.18	20.25	17.76	10.66	20.20	19.44
Game (Phasianidae, Tetraonidae)	748	296	287	165	15.55	25.46	13.97	10.91	9.64	16.61	8.36	6.48
Mammal	541	74	255	212	14.08	6.13	15.16	18.05	6.97	4.15	7.42	8.33
Raptor (Accipitridae, Falconidae, Strigidae)	465	44	225	196	4.35	2.05	4.16	6.17	5.99	2.47	6.55	7.70
Other	906	159	373	374	3.50	3.04	2.77	4.80	11.67	8.92	10.86	14.69
Total	7763	1782	3435	2546	100	100	100	100	100	100	100	100

**Appendix 5:** Occurrence of raptor species in the breeding season diet of a northern goshawk population in Kielder Forest, UK when the number of goshawk home-ranges occupied each year was estimated to be 1-14, 15-24 and 25 or more.

Species	<i>n</i>				% Biomass				% Frequency				% of raptors			
	Total	1-14	15-24	25+	Total	1-14	15-24	25+	Total	1-14	15-24	25+	Total	1-14	15-24	>25
Common kestrel *	228	24	128	76	1.46	0.68	1.83	1.48	2.94	1.35	3.73	2.99	49.03	54.55	56.89	38.78
Tawny owl *	106	7	30	69	1.53	0.45	0.97	3.04	1.37	0.39	0.87	2.71	22.80	15.91	13.33	35.20
Sparrowhawk	48	2	22	24	0.30	0.06	0.31	0.46	0.62	0.11	0.64	0.94	10.32	4.55	9.78	12.24
Short-eared owl *	23	7	14	2	0.23	0.32	0.32	0.06	0.30	0.39	0.41	0.08	4.95	15.91	6.22	1.02
Barn owl *	14	0	3	11	0.13	0	0.06	0.31	0.18	0	0.09	0.43	3.01	0	1.33	5.61
Long-eared owl *	17	0	12	5	0.15	0	0.24	0.14	0.22	0	0.35	0.20	3.66	0	5.33	2.55
Merlin	14	0	12	2	0.09	0	0.17	0.04	0.18	0	0.35	0.08	3.01	0	5.33	1.02
Northern goshawk	13	4	3	6	0.40	0.55	0.21	0.56	0.17	0.22	0.09	0.24	2.80	9.09	1.33	3.06
Common buzzard	2	0	1	1	0.05	0	0.06	0.08	0.03	0	0.03	0.04	0.43	0	0.44	0.51

\* Denotes raptor species which are dependent on field voles.

**Appendix 6:** The proportion of goshawk breeding attempts which were successful (i.e. fledged at least one chick) shown in relation to goshawk abundance (number of occupied goshawk territories) and according to when the goshawk home-range first became established. Numbers in parentheses are the total number of breeding attempts for each category.

Home-ranges established	Number of occupied goshawk home ranges		
	$\leq 14$	15-24	$\geq 25$
Early (1973-1986)	0.73 (60)	0.76 (114)	0.51 (112)
Middle (1987-2001)	-	0.77 (96)	0.61 (125)
Late (2002-2014)	-	0.75 (20)	0.25 (44)