



HAL
open science

Taylor's experiment in a periodically sheared particulate suspension

Mathieu Souzy, Phong Pham, Bloen Metzger

► **To cite this version:**

Mathieu Souzy, Phong Pham, Bloen Metzger. Taylor's experiment in a periodically sheared particulate suspension. *Physical Review Fluids*, 2016, 1 (4), pp.42001 - 42001. 10.1103/PhysRevFluids.1.042001 . hal-01768667

HAL Id: hal-01768667

<https://hal.science/hal-01768667>

Submitted on 17 Apr 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Taylor's experiment in a periodically sheared particulate suspension

Mathieu Souzy,¹ Phong Pham,^{1,2} and Bloen Metzger¹

¹*Aix-Marseille Université, IUSTI-CNRS UMR 7343, 13453 Marseille Cedex 13, France*

²*Department of Chemical Engineering, University of Florida, Gainesville, Florida 32611, USA*

(Received 15 April 2016; published 5 August 2016)

We revisit Taylor's experiment investigating the evolution of a blob of dye in a periodically sheared suspension of non-Brownian particles. Above a critical strain amplitude, particulate suspensions are subject to phase transition where reversibility is lost and particles fail to return to their original positions. We investigate the effect of this transition on the dispersion of a blob of dye. Beyond the critical strain, the dispersion of the blob is found to increase significantly. The dispersion coefficient of the blob of dye is measured and compared to the self-diffusivity coefficient of the particles.

DOI: [10.1103/PhysRevFluids.1.042001](https://doi.org/10.1103/PhysRevFluids.1.042001)

In 1966, Taylor illustrated in a now famous experiment the reversibility of Stokes flow [1]. A blob of dye is initially injected in a Couette cell filled with a very viscous and Newtonian fluid. When rotating one of the cylinder, the fluid is linearly sheared and the blob of dye stretches into a thin filament until almost disappearing. Then, upon reversing the rotation for the same number of turns, the drop astoundingly recovers its initial shape (except for slight blurring due to molecular diffusion).

Inspired by Taylor's work, we performed the exact same experiment but in a suspension of non-Brownian particles to investigate how the presence of particles within the fluid breaks the reversibility of the flow and influences the dispersion process of the blob of dye. Under periodic shear, Pine *et al.* [2] demonstrated that non-Brownian suspensions undergo a dynamical phase transition. For a given particulate volume fraction and a small enough strain amplitude, the suspension relaxes to an absorbing reversible state in which particles return to their original position after every cycle of shear. However, when the strain amplitude exceeds a critical value γ_c , which depends on the volume fraction and the particle roughness [3], the suspension transitions to a fluctuating state: the particles do not return to their original positions, and, when tracked stroboscopically at the end of each cycle of shear, the particles exhibit large fluctuations analogous to a random walk. In that fluctuating state, the loss of reversibility of the particles motion should also induce a loss of reversibility in the fluid motion. The goal in the present Rapid Communication is to perform Taylor's experiment in a particulate suspension to study the effect of this transition on the dispersion of the blob of dye.

Under continuous shear, the phenomenon called "shear-induced diffusion", which was widely investigated [3–10], was shown to significantly enhance the transfer of heat [9,11] or mass [12,13] across suspensions of non-Brownian particles. This enhancement occurs since under shear, particles within the fluid constantly collide with one another and change streamlines. The particles thus generate disturbances within the fluid which promote the dispersion of the scalar (temperature or concentration), prelude to its subsequent mixing. Particles also significantly affect transfers close to boundaries: the particle rotation was shown to disrupt the diffusive boundary layer by a "rolling-coating effect" (Souzy *et al.* [13]) which convects the scalar at a constant rate from the wall, where it is injected, to the bulk of the suspension.

The dispersion enhancement of a scalar field has already been extensively studied in suspensions undergoing a continuous shear flow, but it has not yet been studied under periodic shear. Our motivation is that this latter flow configuration is particularly appealing since, after each cycle of shear, the affine part of the flow (the reversible part) cancels out. Therefore, by measuring the evolution of the blob of ink stroboscopically, i.e., at the end of each cycle of shear, one can investigate the blob's dispersion solely focusing on the effect of the fluctuations induced by the particles, not having to account for the background linear stretching as in a continuous shear.

MATHIEU SOUZY, PHONG PHAM, AND BLOEN METZGER




FIG. 1. (a) Schematic of the setup, top and side views. (b) Image of the suspension seen from the top (particles appear as white discs) and stroboscopic particle trajectories for cycles of shear of strain amplitude $\gamma_0 = 2.8$. (c) Corresponding particle mean square displacement versus total accumulated strain $\gamma = 2N_c \gamma_0$.

An original experimental setup was developed to perform direct visualizations of the dispersion of a blob of dye in a periodically sheared suspension of non-Brownian particles. By measuring the diffusion coefficients of both the particle and the fluid phases, we find that the blob's dispersion follows the same dynamical phase transition as the particles.

The experimental setup, shown in Fig. 1(a), consists of a transparent cell within which a transparent mylar belt, mounted tightly on two vertical rotating cylinders, generates a linear shear flow. A precision rotation stage drives the rotation of one of the vertical cylinders. The fluid is a Newtonian mixture of Triton X-100 (73.86 wt%), zinc chloride (14.24 wt%), and water (11.90 wt%), identical to that used in Souzy *et al.* [13]. Its viscosity $\eta = 3$ Pa s is large enough to ensure a viscous flow ($Re \approx 10^{-4}$), and its composition was chosen to match both the particle density ($\rho_f = \rho_p = 1.19$ g cm $^{-3}$) to avoid sedimentation and their optical index of refraction. A small amount of hydrochloric acid (≈ 0.05 wt%) is added to the solution to prevent the formation of zinc hypochlorite precipitate,

TAYLOR'S EXPERIMENT IN A PERIODICALLY SHEARED ...

thereby significantly improving the optical transparency of the solution. The temperature of the setup is also adjusted (± 0.05 °C) with a water-bath jacket connected to a cryo-thermostat in order to finely tune index matching between the fluid and the particles. We use monodisperse spherical PMMA beads from Engineering Laboratories Inc. with a diameter $d = 2$ μm , which were chosen for their high surface quality and good transparency. Note that, above the region of interest, a small plexiglass window is placed on the free surface of the suspension to avoid its deflection during shear which would deteriorate the visualization. A small hole, through the window, is used to inject the blob of dye in the middle of the suspension with a thin needle. The molecular diffusion coefficient of the dye in the fluid at rest is $D_0 \approx 10^{-13}$ $\text{m}^2 \text{s}^{-1}$ [14]. All experiments were done at a volume fraction of $\phi = 35\%$.

Two types of experiments were performed: (1) to follow the evolution of the blob of dye, the suspension is perfectly transparent and the blob consists in a small injection of fluid doped with a fluorescent tracer (rhodamine 6G), and (2) to track the particle motion, the whole fluid is homogeneously doped with the fluorescent dye. In both cases, visualization is performed from the top of the cell by imaging the rhodamine, which fluoresces under illumination provided by a thin horizontal laser sheet (≈ 60 μm thick). Images are acquired using a high-resolution camera (Nikon D-300S). A high-pass filter ($\lambda > 590$ nm) is used to eliminate direct light reflections and collect the light solely emitted by the fluorescent rhodamine.

As presented in Fig. 1(b), when the whole fluid is doped with rhodamine, the particles within the light sheet appear as white discs, while the fluid is dark. Note that all the particles have the same size; the apparent size differences arises from their different vertical positions relative to the light plane. Images are first thresholded, then an algorithm of particle detection based on the circular Hough transform [9,15] is applied to find the center location of each particle within ± 1 pixel. Particles trajectories are then reconstructed by applying a tracking algorithm. A very important point is that for an experiment where periodic oscillations are performed at the strain amplitude γ_0 , prior to injecting the blob or measuring the particle trajectories, the suspension is first presheared at that strain amplitude for 40 cycles in order to let the suspension self-organize into its steady-state configuration [16]. To track the particles, images are taken continuously every strain increment of 0.1, while for blob experiments, images are acquired only at the end of each cycle of shear. A cycle of shear consists in slowly shearing (at 0.1 s^{-1}) the suspension forward for a strain γ_0 , then shearing it backward to the initial position. For each experiment, N_c cycles of shear are performed, and the total accumulated strain is thus $\gamma = 2N_c\gamma_0$.

We first present results characterizing the dynamics of the particles. Trajectories of the particles are recorded continuously to be able to track particles from one cycle to the next. However, to quantify the amount of irreversibility, we compute the mean square displacement of the particles using only their positions at the end of each cycle. An example of such trajectories is represented in Fig. 1(b) for an experiment performed at large strain amplitude. The mean square displacement in the gradient (y) direction $\langle \Delta y \Delta y \rangle$, where $\Delta y = y(t) - y(t=0)$ and the angle brackets denote an average over all particles, is close to zero for small applied strain amplitudes. This means that the particle motion is reversible. However, for experiments performed at larger strains, the mean square displacement increases linearly with time, indicating that the particles do not return to their original position and follow a diffusive behavior; see Fig. 1(c). Following Pine *et al.*, the slope of the particle mean square displacements is used to define a dimensionless particle diffusion coefficient $D_p^* = \langle \Delta y \Delta y \rangle / (2d^2 \dot{\gamma} t)$, which quantifies the irreversible motion of the particles. We recover the results of Pine *et al.* that particles, beyond a critical strain amplitude of $\gamma_c \approx 2$ for the present volume fraction $\phi = 35\%$ (equivalent of $\gamma_c \approx 1$ with the definition of the strain amplitude used by Pine [2]), transit to a fluctuating state; see Fig. 4.

We now investigate the effect of this transition on the dispersion of a blob of dye. After pouring the suspension (without rhodamine) within the cell and preshearing it, a cylindrical blob of dyed fluid is injected at $t = 0$ and with the fluid at rest in the middle of the shear cell. The blob has an initial diameter $s_0 \simeq d$ and is aligned with the vorticity direction of the shear flow and centered on the neutral velocity plane. This results in a macroscopically two-dimensional initial configuration

MATHIEU SOUZY, PHONG PHAM, AND BLOEN METZGER




FIG. 2. Image of the blob after 40 cycles of shear performed at a strain amplitude: $\gamma_0 = 0.5$, $\gamma_0 = 2.5$, $\gamma_0 = 3$. Each image corresponds to the superposition of 10 pictures taken from 10 independent experiments performed at the same strain amplitude; see also Movie 1 and Movie 2 [18].

and ensures that the blob does not drift with the flow. The concentration of rhodamine within the blob was adjusted so that absorption of the light intensity can be neglected. The suspension is then slowly sheared back and forth over $N_c = 40$ cycles at the strain amplitude γ_0 .

Figures 2(a)–2(c) show how the blob has evolved after 40 cycles of shear for experiments performed at different strain amplitudes. Note that each image is the superposition of pictures taken from 10 independent experiments performed with the same imposed strain amplitude. At small strain amplitude $\gamma_0 = 0.5$, as in Taylor’s experiment, the blob after each cycle of shear recovers its initial shape; see Fig. 2(a). Conversely, for larger applied strain amplitudes, the blob rapidly spreads in the surrounding medium, evolving towards a topology constituted of filament-like structures [17] (to visualize the continuous evolution of the blob; see Movie 1 and Movie 2 [18]). Larger strain amplitudes result, after the same number of cycles, into a larger dispersion of the blob.

More quantitatively, we can characterize the blob effective dispersion by examining its spreading. Figure 3(a) shows successive concentration profiles obtained by averaging the light-intensity level along the x direction. The low concentration of rhodamine used for the experiments ensures a linear relation between light intensity and dye concentration. Two important observations can be made. First, the concentration distribution is normally distributed. The dashed curves are the best Gaussian fit where, once the amplitude of the initial distribution (for $N_c = 0$) is fixed, the sole remaining adjustable parameter is the standard deviation σ . Second, the relative variance of the concentration distribution $\Delta\sigma^2(t) = \sigma^2(t) - \sigma^2(t = 0)$ increases linearly with the total accumulated




FIG. 3. (a) Successive concentration profiles for a blob submitted to a strain amplitude $\gamma_0 = 2.9$. The profiles are obtained by averaging over 10 independent runs. The dashed curves correspond to the best Gaussian fit. (b) Evolution of the corresponding dimensionless relative variance $\Delta\sigma^2/d^2$ as a function of the total accumulated strain $\gamma = 2N_c\gamma_0$.

TAYLOR'S EXPERIMENT IN A PERIODICALLY SHEARED ...




FIG. 4. Dimensionless self-diffusivity coefficient of the particle D_p^* and dimensionless dispersion coefficient of the fluid D_f^* , as a function of the applied strain amplitude γ_0 . Reversibility is lost above a critical strain amplitude $\gamma_c \approx 2$.

strain $\gamma = 2N_c\gamma_0$ (equivalent to time); see Fig. 3(b). These two features evidence that the dispersion process of the blob, and thus of the interstitial fluid phase between the particles, is diffusive. The effective dispersion coefficient of the fluid phase is provided by the classical definition

$$\Delta\sigma^2 = 2D_f t. \quad (1)$$

The dimensionless effective dispersion coefficient of the fluid phase, D_f^* , can be defined by the dimensionless form of Eq. (1) ($\Delta\sigma^2/d^2 = 2\frac{D_f}{\dot{\gamma}d^2}\gamma$) as $D_f^* = D_f/\dot{\gamma}d^2$, which corresponds to one half the slope of the linear fits presented on Fig. 3(b). Error bars are also estimated from the typical standard deviation between runs; see Fig. 4.

Similarly to the behavior of the particles, for the volume fraction $\phi = 35\%$ investigated, above a critical strain $\gamma_c \approx 2$, the dispersion coefficient of the blob goes through an abrupt transition; see Fig. 4 where D_p^* and D_f^* are plotted versus the applied strain amplitude γ_0 . The loss of reversibility induced by the presence of the particles significantly accelerates the dispersion process. Indeed, if one was only considering the effect of molecular diffusion, i.e., if the blob was sheared in a pure fluid without particle, the transverse dispersion coefficient of the dye would be $D_f = D_0$, leading to a dimensionless effective dispersion coefficient of the fluid phase, $D_f^* = D_0/\dot{\gamma}d^2 = 10^{-6}$. In the fluctuating state (beyond γ_c), the dimensionless dispersion coefficient of the fluid is $\approx 10^{-2}$, so about four orders of magnitude larger than that of molecular diffusion alone. Note that, as one can guess from Fig. 2, the dispersion coefficients in the flow direction (not shown here) follow very similar trends as those in the gradient direction.

For $\gamma_0 < \gamma_c$, in the so-called reversible state, we measure effective diffusion coefficients of $\approx 10^{-5}$ both for the particles and for the blob. For the particles, which are non-Brownian, D_p^* should vanish when $\gamma_0 \rightarrow 0$. This small discrepancy arises from experimental artifacts which are noise from the Hough detection algorithm (detection of the particle center locations within ± 1 pixel) and unavoidable slight sedimentation or natural convection currents within the fluid. In that limit, the latter two effects probably also bias the measurement of the blob dispersion coefficient. It is noteworthy that the measured self-diffusivity coefficient of the particles obtained at large strain amplitudes is in good agreement with previous measurements done in continuously sheared particulate suspensions [7,9]. Similarly, the enhancement of the dye diffusion coefficient at large applied strain amplitude is of the same order $\approx 10^4$ than in a continuously sheared suspension [13].

This result could be useful in certain mixing applications where oscillatory shear might be easier to implement than continuous shear [19].

The molecular diffusion coefficient of rhodamine 6G in the present suspending liquid is small ($D_0 \approx 10^{-13} \text{ m}^2 \text{ s}^{-1}$). In such a condition of large Péclet number, $Pe = \dot{\gamma} d^2 / D_0 = 10^6$, (with $s_0 \simeq d$) where the effect of molecular diffusion is negligible relative to the fluctuations induced by the presence of the particles [5,20], the measurement of the dye-effective dispersion coefficient provides a good estimate of the fluid dispersion coefficient. Conversely to what was reported by Breedveld *et al.* [21], we find that the fluid dispersion coefficient is always slightly larger than that of the particles; see Fig. 4. One can explain this trend by the fact that the self-diffusivity coefficient of the particles accounts only for their translational diffusivity, while the fluid dispersion coefficient arises from the perturbations induced by both the translational and the rotational motion of the particles.

We performed Taylor's experiment in a suspension of non-Brownian particles to investigate how the loss of reversibility induced by the presence of the particles affects the dispersion of the blob of dye. By measuring both the particles and the blob of dye transverse effective diffusion coefficients, we found that the dispersion of the blob undergoes the same transition as the one observed by the particles [2]. At small strain amplitude, as in Taylor's experiment, the system is reversible and the blob thus recovers its initial shape after each cycle of shear. For a volume fraction $\phi = 35\%$, beyond a critical strain amplitude $\gamma_c \approx 2$, the system transits to a fluctuating state where reversibility is lost. As a result, the blob spreads very rapidly. The blob dispersion is found to be diffusive with an effective diffusive coefficient of the same order as the self-diffusivity coefficient of the particles.

We would like to thank Paul Cervetti, Sady Noel, and Stephane Martinez for building the experimental setup. This work was supported by the National Science Foundation (Grants No. 0968313 and No. 1362060), ANR JCJC SIMI 9, the Labex MEC (ANR-10-LABX-0092) and the A*MIDEX project (ANR-11-IDEX-0001-02), funded by the "Investissements d'Avenir" French Government program managed by the French National Research Agency (ANR).

-
- [1] G. I. Taylor, Kinematic Reversibility, in *Multi-Media Fluid Mechanics CD-ROM*, edited by G. M. Homsy (Cambridge University Press, Cambridge, 2000).
 - [2] D. J. Pine, J. P. Gollub, J. F. Brady, and A. M. Leshansky, Chaos and threshold for irreversibility in sheared suspensions, *Nature (London)* **438**, 997 (2005).
 - [3] P. Pham, J. E. Butler, and B. Metzger, Origin of critical strain amplitude in periodically sheared suspensions, *Phys. Rev. Fluids* **1**, 022201(R) (2016).
 - [4] P. A. Arp, and S. G. Mason, The kinetics of flowing dispersions IX. Doublets of rigid spheres (experimental), *J. Colloid Interface Sci.* **61**, 44 (1976).
 - [5] E. C. Eckstein, D. G. Bailey, and A. H. Shapiro, Self-diffusion of particles in shear flow of a suspension, *J. Fluid Mech.* **79**, 191 (1977).
 - [6] F. R. Da Cunha and E. J. Hinch, Shear-induced dispersion in a dilute suspension of rough spheres, *J. Fluid Mech.* **309**, 211 (1996).
 - [7] A. Sierou and J. F. Brady, Shear-induced self-diffusion in non-colloidal suspensions, *J. Fluid Mech.* **506**, 285 (2004).
 - [8] F. Blanc, F. Peters, and E. Lemaire, Experimental Signature of the Pair Trajectories of Rough Spheres in the Shear-Induced Microstructure in Noncolloidal Suspensions, *Phys. Rev. Lett.* **107**, 208302 (2011).
 - [9] B. Metzger, O. Rahli, and X. Yin, Heat transfer across sheared suspensions: role of the shear-induced diffusion, *J. Fluid Mech.* **724**, 527 (2013).
 - [10] P. Pham, B. Metzger, and J. E. Butler, Particle dispersion in sheared suspensions: Crucial role of solid-solid contacts, *Phys. Fluids* **27**, 051701 (2015).
 - [11] C. W. Sohn and M. M. Chen, Heat transfer enhancement in laminar slurry pipe flows with power law thermal conductivities, *J. Heat Transfer* **106**, 539 (1984).

TAYLOR'S EXPERIMENT IN A PERIODICALLY SHEARED . . .

- [12] N. L. Wang and K. H. Keller, Augmented transport of extracellular solutes in concentrated erythrocyte suspensions in Couette flow, *J. Colloid Interface Sci.* **103**, 210 (1985).
- [13] M. Souzy, X. Yin, E. Villiermaux, C. Abid, and B. Metzger, Super-diffusion in sheared suspensions, *Phys. Fluids* **27**, 041705 (2015).
- [14] C. T. Culbertson, S. C. Jacobson, and J. M. Ramsey, Diffusion coefficient measurements in microfluidic devices, *Talanta* **56**, 365 (2002).
- [15] T. Peng, Detect circles with various radii in grayscale image via Hough Transform, <http://www.mathworks.com/matlabcentral/fileexchange/9168-detect-circleswith-various-radii-in-greyscale-image-via-hough-transform>.
- [16] L. Cort e, P. Chaikin, J. P. Gollub, and D. J. Pine, Random organization in periodically driven systems, *Nature Phys.* **4**, 420 (2008).
- [17] J. M. Ottino, *The Kinematics of Mixing: Stretching, Chaos, and Transport* (Cambridge University Press, Cambridge, 1989).
- [18] See Supplemental Material at <http://link.aps.org/supplemental/10.1103/PhysRevFluids.1.042001> for two movies that show the evolution of a drop of dye in a particulate suspension (volume fraction $\phi = 35\%$) periodically sheared for a strain amplitude of 1.5 (Movie 1) and a strain amplitude of 4 (Movie 2).
- [19] M. Workamp, V. Saggiomo, and J. A. Dijksman, A simple low pressure drop suspension-based microfluidic mixer, *J. Micromech. Microeng.* **25**, 094003 (2015).
- [20] D. Leighton and A. Acrivos, Measurement of shear-induced self-diffusion in concentrated suspensions of spheres, *J. Fluid Mech.* **177**, 109 (1987).
- [21] V. Breedveld, D. van den Ende, A. Tripathi, and A. Acrivos, The measurement of the shear-induced particle and fluid tracer diffusivities in concentrated suspensions by a novel method, *J. Fluid Mech.* **375**, 297 (1998).