



HAL
open science

On complete monotonicity of linear combination of finite psi functions

Bai-Ni Guo, Feng Qi

► **To cite this version:**

Bai-Ni Guo, Feng Qi. On complete monotonicity of linear combination of finite psi functions. Communications of the Korean Mathematical Society, 2019, 34 (4), pp.1223–1228. 10.4134/CKMS.c180430 . hal-01773131v2

HAL Id: hal-01773131

<https://hal.science/hal-01773131v2>

Submitted on 18 Jan 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Commun. Korean Math. Soc. **34** (2019), No. 4, pp. 1223–1228
<https://doi.org/10.4134/CKMS.c180430>
pISSN: 1225-1763 / eISSN: 2234-3024

**ON COMPLETE MONOTONICITY OF LINEAR
COMBINATION OF FINITE PSI FUNCTIONS**

BAI-NI GUO AND FENG QI

Reprinted from the
Communications of the Korean Mathematical Society
Vol. 34, No. 4, October 2019

©2019 Korean Mathematical Society

ON COMPLETE MONOTONICITY OF LINEAR COMBINATION OF FINITE PSI FUNCTIONS

BAI-NI GUO AND FENG QI

ABSTRACT. In the paper, the authors supply complete monotonicity of linear combination of finite psi functions and extend some known results.

1. Preliminaries

It is well-known [1, 12, 14] that the classical gamma function $\Gamma(z)$ can be defined by

$$\Gamma(z) = \int_0^{\infty} t^{z-1} e^{-t} dt, \quad \Re(z) > 0$$

or by

$$\Gamma(z) = \lim_{n \rightarrow \infty} \frac{n! n^z}{\prod_{k=0}^n (z+k)}, \quad z \in \mathbb{C} \setminus \{0, -1, -2, \dots\}$$

and that the logarithmic derivative

$$\psi(z) = [\ln \Gamma(z)]' = \frac{\Gamma'(z)}{\Gamma(z)}$$

is called the psi or digamma function.

From [11, Chapter XIII], [30, Chapter 1], and [31, Chapter IV], we recall that an infinitely differentiable and nonnegative function $f(x)$ is said to be completely monotonic on an interval I if and only if

$$(-1)^{m-1} f^{(m-1)}(x) \geq 0$$

for all $m \in \mathbb{N}$ and $x \in I$. The Bernstein–Widder theorem [31, p. 161, Theorem 12b] states that a necessary and sufficient condition for $f(x)$ to be completely monotonic on $(0, \infty)$ is that

$$f(x) = \int_0^{\infty} e^{-xt} d\mu(t)$$

Received October 15, 2018; Accepted February 27, 2019.

2010 *Mathematics Subject Classification.* Primary 26A48; Secondary 33B15, 44A10.

Key words and phrases. complete monotonicity, psi function, linear combination, alternative proof.

for $x \in (0, \infty)$, where μ is non-decreasing and the above integral converges. In other words or simply speaking, a function is completely monotonic on $(0, \infty)$ if and only if it is a Laplace transform.

2. Motivations

In [2, Theorem 4.1], it was obtained that, if a_k and b_k for $1 \leq k \leq m$ satisfy $a_1 \geq a_2 \geq \cdots \geq a_m$ and $b_1 \geq b_2 \geq \cdots \geq b_m > 0$, then

$$\phi_0(x) = \sum_{k=1}^m a_k \psi(b_k x)$$

is completely monotonic on $(0, \infty)$ if and only if

$$\sum_{k=1}^m a_k = 0 \quad \text{and} \quad \sum_{k=1}^m a_k \ln b_k \geq 0.$$

In [8, Lemma 2.1], the function $\phi_0(x)$ was extended as

$$\phi_\delta(x) = \sum_{k=1}^m a_k \psi(b_k x + \delta)$$

for $x > 0$ and $\delta \geq 0$ and it was acquired that, if $a_1 \geq a_2 \geq \cdots \geq a_m$ and $b_1 \geq b_2 \geq \cdots \geq b_m \geq 0$ such that $\sum_{k=1}^m a_k \geq 0$, then, when $\delta \geq \frac{1}{2}$, the first derivative $\phi'_\delta(x)$ is completely monotonic and, consequently, the function $\phi_\delta(x)$ is increasing and concave, on $(0, \infty)$. These results were applied in [8] to discuss a problem arising in the context of statistical density estimation based on Bernstein polynomials.

In the proof of [29, Theorem 2.2], it was obtained that the function

$$\left(\sum_{i=1}^m a_i \right)^2 \psi' \left(1 + x \sum_{i=1}^m a_i \right) - \sum_{i=1}^m a_i^2 \psi'(1 + a_i x), \quad a_i > 0$$

is completely monotonic and, consequently,

$$(1) \quad \left(\sum_{i=1}^m a_i \right) \psi \left(1 + x \sum_{i=1}^m a_i \right) - \sum_{i=1}^m a_i \psi(1 + a_i x), \quad a_i > 0$$

is positive, increasing, and concave, with respect to $x \in (0, \infty)$. See also the proof of [13, Theorem 2.1]. In other words, the function defined by (1) is a Bernstein function of $x \in (0, \infty)$. For detailed information on the Bernstein functions, please refer to the monograph [30], the papers [21–23, 25, 26] and closely related references.

In this paper, we supply an alternative proof for the above complete monotonicity of the functions $\phi_0(x)$ and $\phi_\delta(x)$ for $\delta \geq \frac{1}{2}$ on $(0, \infty)$ and obtain slightly extended conclusions.

3. Main results and their proofs

We now state and prove our main results alternatively.

Theorem 1. *If $\delta \geq \frac{1}{2}$, $(a_i - a_j)(b_i - b_j) \geq 0$ for all $1 \leq i, j \leq m$, and $\sum_{k=1}^m a_k \geq 0$, then the first derivative $\pm\phi'_\delta(x)$ is completely monotonic and, consequently, the function $\pm\phi_\delta(x)$ is increasing and concave, on $(0, \infty)$.*

If $(a_i - a_j)(b_i - b_j) \geq 0$ for all $1 \leq i, j \leq m$, then the function $\pm\phi_0(x)$ is completely monotonic on $(0, \infty)$ if and only if $\sum_{k=1}^m a_k = 0$ and $\sum_{k=1}^m a_k \ln b_k \geq 0$.

Proof. Using the formula

$$\psi'(z) = \int_0^\infty \frac{t}{1 - e^{-t}} e^{-zt} dt, \quad \Re(z) > 0$$

in [1, p. 260, 6.4.1] gives

$$\begin{aligned} \psi'(\tau x + \delta) &= \int_0^\infty \frac{t}{1 - e^{-t}} e^{-(\tau x + \delta)t} dt \\ &= \int_0^\infty \frac{te^{-\delta t}}{1 - e^{-t}} e^{-\tau x t} dt \\ &= \frac{1}{\tau} \int_0^\infty h\left(\frac{v}{\tau}\right) e^{-vx} dv, \end{aligned}$$

where $\tau > 0$ and $h_\delta(t) = \frac{te^{-\delta t}}{1 - e^{-t}}$. Hence

$$\phi'_\delta(x) = \sum_{k=1}^m a_k b_k \psi'(b_k x + \delta) = \int_0^\infty \left[\sum_{k=1}^m a_k h_\delta\left(\frac{v}{b_k}\right) \right] e^{-vx} dv.$$

In [9, 32], it was established that the positive function

$$h_\delta(t) = \begin{cases} \frac{t}{e^{\delta t} - e^{(\delta-1)t}}, & t \neq 0, \\ 1, & t = 0, \end{cases}$$

is decreasing on \mathbb{R} if $\delta \geq 1$, increasing on \mathbb{R} if $\delta \leq 0$, increasing in $(-\infty, 0)$ if $\delta \leq \frac{1}{2}$, and decreasing in $(0, \infty)$ if $\delta \geq \frac{1}{2}$. For more information on properties and applications of $h_\delta(t)$, please refer to the papers and review articles [3, 4, 6, 15–19, 24, 28] and closely related references therein. Therefore, by virtue of the Čebyšev inequality in [10, p. 36, Section 2.5, Theorem 1], we acquire

$$\frac{1}{m} \sum_{k=1}^m a_k h_\delta\left(\frac{v}{b_k}\right) \geq \left(\frac{1}{m} \sum_{k=1}^m a_k\right) \left[\frac{1}{m} \sum_{k=1}^m h_\delta\left(\frac{v}{b_k}\right)\right] \geq 0.$$

This means that the first derivative $\pm\phi'_\delta(x)$ is completely monotonic on $(0, \infty)$. Consequently, the function $\pm\phi_\delta(x)$ is increasing and concave on $(0, \infty)$.

When $\delta = 0$, the function $h_0(t)$ is positive and increasing on $[0, \infty)$ and

$$\frac{1}{m} \sum_{k=1}^m a_k h_0\left(\frac{v}{b_k}\right) \geq \left(\frac{1}{m} \sum_{k=1}^m a_k\right) \left[\frac{1}{m} \sum_{k=1}^m h_0\left(\frac{v}{b_k}\right)\right].$$

Accordingly, when $\sum_{k=1}^m a_k \leq 0$, the first derivative $\mp\phi'_0(x)$ is completely monotonic and the function $\mp\phi_0(x)$ is increasing on $(0, \infty)$. Furthermore, utilizing

$$\lim_{x \rightarrow \infty} [\ln x - \psi(x)] = 0$$

in [5, Theorem 1] and [7, 27] yields that the limit

$$\phi_0(x) = \sum_{k=1}^m a_k [\psi(b_k x) - \ln(b_k x)] + \sum_{k=1}^m a_k \ln b_k + (\ln x) \sum_{k=1}^m a_k \rightarrow \sum_{k=1}^m a_k \ln b_k$$

as $x \rightarrow \infty$ is valid and

$$\mp\phi_0(x) \leq \mp \sum_{k=1}^m a_k \ln b_k$$

if and only if $\sum_{k=1}^m a_k = 0$. When and only when $\sum_{k=1}^m a_k \ln b_k \geq 0$, the function $\pm\phi_0(x)$ is positive and, consequently, completely monotonic, on $(0, \infty)$. The proof of Theorem 1 is complete. \square

Remark 1. This paper is a slight revision of the preprint [20].

References

- [1] M. Abramowitz and I. A. Stegun, *Handbook of mathematical functions with formulas, graphs, and mathematical tables*, National Bureau of Standards Applied Mathematics Series, **55**, For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, DC, 1964.
- [2] H. Alzer and C. Berg, *Some classes of completely monotonic functions. II*, Ramanujan J. **11** (2006), no. 2, 225–248. <https://doi.org/10.1007/s11139-006-6510-5>
- [3] W.-S. Cheung and F. Qi, *Logarithmic convexity of the one-parameter mean values*, Taiwanese J. Math. **11** (2007), no. 1, 231–237. <https://doi.org/10.11650/twjmath/1500404648>
- [4] B.-N. Guo and F. Qi, *A simple proof of logarithmic convexity of extended mean values*, Numer. Algorithms **52** (2009), no. 1, 89–92. <https://doi.org/10.1007/s11075-008-9259-7>
- [5] ———, *Two new proofs of the complete monotonicity of a function involving the PSI function*, Bull. Korean Math. Soc. **47** (2010), no. 1, 103–111. <https://doi.org/10.4134/BKMS.2010.47.1.103>
- [6] ———, *The function $(b^x - a^x)/x$: logarithmic convexity and applications to extended mean values*, Filomat **25** (2011), no. 4, 63–73. <https://doi.org/10.2298/FIL1104063G>
- [7] B.-N. Guo, F. Qi, J.-L. Zhao, and Q.-M. Luo, *Sharp inequalities for polygamma functions*, Math. Slovaca **65** (2015), no. 1, 103–120. <https://doi.org/10.1515/ms-2015-0010>
- [8] A. Leblanc and B. C. Johnson, *On a uniformly integrable family of polynomials defined on the unit interval*, JIPAM. J. Inequal. Pure Appl. Math. **8** (2007), no. 3, Article 67, 5 pp.

- [9] A.-Q. Liu, G.-F. Li, B.-N. Guo, and F. Qi, *Monotonicity and logarithmic concavity of two functions involving exponential function*, *Internat. J. Math. Ed. Sci. Tech.* **39** (2008), no. 5, 686–691. <https://doi.org/10.1080/00207390801986841>
- [10] D. S. Mitrinović, *Analytic Inequalities*, Springer-Verlag, New York, 1970.
- [11] D. S. Mitrinović, J. E. Pečarić, and A. M. Fink, *Classical and New Inequalities in Analysis*, Mathematics and its Applications (East European Series), **61**, Kluwer Academic Publishers Group, Dordrecht, 1993. <https://doi.org/10.1007/978-94-017-1043-5>
- [12] F. W. J. Olver, D. W. Lozier, R. F. Boisvert, and C. W. Clark (eds.), *NIST Handbook of Mathematical Functions*, Cambridge University Press, New York, 2010; Available online at <http://dlmf.nist.gov/>.
- [13] F. Ouimet, *Complete monotonicity of multinomial probabilities and its application to Bernstein estimators on the simplex*, *J. Math. Anal. Appl.* **466** (2018), no. 2, 1609–1617. <https://doi.org/10.1016/j.jmaa.2018.06.049>
- [14] F. Qi, *Limit formulas for ratios between derivatives of the gamma and digamma functions at their singularities*, *Filomat* **27** (2013), no. 4, 601–604. <https://doi.org/10.2298/FIL1304601Q>
- [15] F. Qi, P. Cerone, S. S. Dragomir, and H. M. Srivastava, *Alternative proofs for monotonic and logarithmically convex properties of one-parameter mean values*, *Appl. Math. Comput.* **208** (2009), no. 1, 129–133. <https://doi.org/10.1016/j.amc.2008.11.023>
- [16] F. Qi and J.-X. Cheng, *Some new Steffensen pairs*, *Anal. Math.* **29** (2003), no. 3, 219–226. <https://doi.org/10.1023/A:1025467221664>
- [17] F. Qi, J.-X. Cheng, and G. Wang, *New Steffensen pairs*, in *Inequality theory and applications*. Vol. I, 273–279, Nova Sci. Publ., Huntington, NY, 2001.
- [18] F. Qi and B.-N. Guo, *On Steffensen pairs*, *J. Math. Anal. Appl.* **271** (2002), no. 2, 534–541. [https://doi.org/10.1016/S0022-247X\(02\)00120-8](https://doi.org/10.1016/S0022-247X(02)00120-8)
- [19] ———, *Some properties of extended remainder of Binet’s first formula for logarithm of gamma function*, *Math. Slovaca* **60** (2010), no. 4, 461–470. <https://doi.org/10.2478/s12175-010-0025-7>
- [20] ———, *An alternative proof for complete monotonicity of linear combinations of many psi functions*, HAL archives (2018), available online at <https://hal.archives-ouvertes.fr/hal-01773131>.
- [21] ———, *The reciprocal of the weighted geometric mean is a Stieltjes function*, *Bol. Soc. Mat. Mex. (3)* **24** (2018), no. 1, 181–202. <https://doi.org/10.1007/s40590-016-0151-5>
- [22] ———, *The reciprocal of the weighted geometric mean of many positive numbers is a Stieltjes function*, *Quaest. Math.* **41** (2018), no. 5, 653–664. <https://doi.org/10.2989/16073606.2017.1396508>
- [23] ———, *Lévy-Khintchine representation of Toader-Qi mean*, *Math. Inequal. Appl.* **21** (2018), no. 2, 421–431. <https://doi.org/10.7153/mia-2018-21-29>
- [24] F. Qi, B.-N. Guo, and C.-P. Chen, *The best bounds in Gautschi-Kershaw inequalities*, *Math. Inequal. Appl.* **9** (2006), no. 3, 427–436. <https://doi.org/10.7153/mia-09-41>
- [25] F. Qi and W.-H. Li, *Integral representations and properties of some functions involving the logarithmic function*, *Filomat* **30** (2016), no. 7, 1659–1674. <https://doi.org/10.2298/FIL1607659Q>
- [26] F. Qi and D. Lim, *Integral representations of bivariate complex geometric mean and their applications*, *J. Comput. Appl. Math.* **330** (2018), 41–58. <https://doi.org/10.1016/j.cam.2017.08.005>
- [27] F. Qi and A.-Q. Liu, *Completely monotonic degrees for a difference between the logarithmic and psi functions*, *J. Comput. Appl. Math.* **361** (2019), 366–371. <https://doi.org/10.1016/j.cam.2019.05.001>

- [28] F. Qi, Q.-M. Luo, and B.-N. Guo, *The function $(b^x - a^x)/x$: ratio's properties*, in Analytic number theory, approximation theory, and special functions, 485–494, Springer, New York, 2014. https://doi.org/10.1007/978-1-4939-0258-3_16
- [29] F. Qi, D.-W. Niu, D. Lim, and B.-N. Guo, *Some logarithmically completely monotonic functions and inequalities for multinomial coefficients and multivariate beta functions*, HAL archives (2018), available online at <https://hal.archives-ouvertes.fr/hal-01769288>.
- [30] R. L. Schilling, R. Song, and Z. Vondraček, *Bernstein Functions*, second edition, De Gruyter Studies in Mathematics, **37**, Walter de Gruyter & Co., Berlin, 2012. <https://doi.org/10.1515/9783110269338>
- [31] D. V. Widder, *The Laplace Transform*, Princeton Mathematical Series, **6**, Princeton University Press, Princeton, NJ, 1941.
- [32] S.-Q. Zhang, B.-N. Guo, and F. Qi, *A concise proof for properties of three functions involving the exponential function*, Appl. Math. E-Notes **9** (2009), 177–183.

BAI-NI GUO
SCHOOL OF MATHEMATICS AND INFORMATICS
HENAN POLYTECHNIC UNIVERSITY
JIAOZUO 454010, HENAN, P. R. CHINA
Email address: bai.ni.guo@gmail.com

FENG QI
SCHOOL OF MATHEMATICAL SCIENCES
TIANJIN POLYTECHNIC UNIVERSITY
TIANJIN 300387, P. R. CHINA
Email address: qifeng618@gmail.com