



**HAL**  
open science

# QUASI-STATIC LARGE DEVIATIONS

Anna de Masi, Stefano Olla

► **To cite this version:**

| Anna de Masi, Stefano Olla. QUASI-STATIC LARGE DEVIATIONS. 2018. hal-01777505v1

**HAL Id: hal-01777505**

**<https://hal.science/hal-01777505v1>**

Preprint submitted on 24 Apr 2018 (v1), last revised 10 Feb 2019 (v4)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# QUASI-STATIC LARGE DEVIATIONS

ANNA DE MASI AND STEFANO OLLA

ABSTRACT. We consider the symmetric simple exclusion with open boundaries that are incontact with particle reservoirs at different densities. The reservoir densities changes at a slower time scale with respect the natural time scale the system reach the stationary state. This gives rise to a quasi static hydrodynamic limit proven in [6]. We study here the large deviations with respect to this limit for the particle density field and the total current. We identify explicitly the large deviation functional and prove that it satisfies a fluctuation relation.

## 1. INTRODUCTION

In usual hydrodynamic limits it is studied the macroscopic evolution of the conserved quantities of a *large* microscopic system, under a space-time scaling such that the time scale is the typical one where the system reaches equilibrium. When the system is open and connected to thermal or particle reservoirs, the parameters of these reservoirs give the boundary conditions for the partial differential equations which describe the macroscopic evolution of the conserved quantities.

When the parameters of the boundary reservoirs (for example the particle density or the temperature) are not constant but change in time at a scale slower than the time to relaxation at equilibrium, rescaling the time at this slower scale give a *quasi static* macroscopic evolution for the conserved quantities: at each *macroscopic* time the profile of the conserved quantity is equal to the stationary one corresponding to the given boundary condition at that time. When these stationary profiles are of equilibrium, this limit models the thermodynamic quasi static transformations, where the Clausius equality holds, i.e. the *work* done by the boundaries.

---

*Date:* April 24, 2018. **File:** `dmoLD-9.tex`.

*2000 Mathematics Subject Classification.* 60F10, 82C22, 82C70.

*Key words and phrases.* Quasi-Static thermodynamic, hydrodynamic limits, Large Deviations.

We thank Errico Presutti for stimulating discussions. We thank the kind hospitality of GSSI.

SO's research is supported by ANR-15-CE40-0020-01 grant LSD.

In [1], [11] and [10], quasi static transformations are obtained from a time rescaling of the macroscopic diffusive equation.

In [6] we have started to analyse, for various stochastic particle systems whose macroscopic evolution is described by a diffusive equation, the direct hydrodynamic quasi static limit by rescaling properly space and time in the microscopic dynamics,

We study here the large deviation for one of these models, the symmetric simple exclusion process.

The system we consider is composed by  $2N + 1$  sites (denoted by  $-N, \dots, N$ ) where particles move like symmetric random walks with exclusion. At the boundaries the system is connected to given reservoirs with time changing densities  $\rho_-(t)$  on the left and  $\rho_+(t)$  on the right. We refer at *macroscopic time scale* the scale at which boundary conditions change. The particles jumps or are created and annihilated at the boundaries with rate of the order  $N^{2+\alpha}$ . When  $\alpha = 0$  the empirical density converges, as  $N \rightarrow \infty$ , to the solution of the linear heat equation with Dirichlet slowly varying boundary conditions  $\rho_{\pm}(t)$ . In the case  $\rho_{\pm}$  constants the result is known since the '80's, [7]. As proved in [6], when  $\alpha > 0$ , at each time  $t$  the empirical density converges, as  $N \rightarrow \infty$ , to the solution of the stationary heat equation with boundary conditions  $\rho_{\pm}(t)$ :

$$\bar{\rho}(t, y) = \frac{1}{2}[\rho_+(t) - \rho_-(t)]y + \frac{1}{2}[\rho_+(t) + \rho_-(t)], \quad y \in [-1, 1] \quad (1.1)$$

and a space homogeneous (time integrated) current

$$\bar{J}(t, y) = - \int_0^t \frac{1}{2}[\rho_+(s) - \rho_-(s)] ds. \quad (1.2)$$

Thus the quasi-static limit gives the evolution of macroscopic profile, completely driven by the boundary conditions.

In [6] it is also proven that the distribution of the process is close to a product Bernoulli measure with parameter  $\bar{\rho}(t, y)$ . Stronger results are proved by controlling the correlation functions with the same techniques used in [5]. First order corrections have been studied in [4].

Observe that if  $\rho_+(t) = \rho_-(t)$  then the above result says that the stochastic process at any macroscopic time is close to equilibrium but the order parameter changes in time thus performing a quasi static transformation.

We study in this article the the joint large deviations of the density and the current with respect to this quasi-static limit. We prove that the probability to observe a profile of density  $\rho(t, y)$  that satisfies the

boundary conditions  $\rho_{\pm}(t)$  and a total current  $J(t)$  is asymptotically  $\mathbb{P}(\rho, J) \sim e^{-N^{1+\alpha}I(J, \rho)}$  where

$$I(J, \rho) = \frac{1}{4} \int_0^T \int_{-1}^1 \frac{(J'(t) + \partial_y \rho(t, y))^2}{\phi(\rho(t, y))} dy dt. \quad (1.3)$$

See section 4 for a precise statement. Notice that  $J(t)$  is space constant: in the quasi static limit the total current must be homogeneous in space.

Clearly  $I(\bar{J}, \bar{\rho}) = 0$ , and  $I$  satisfy the fluctuation relation

$$I(J, \rho) - I(-J, \rho) = \int_0^T J'(t)(z_+(t) - z_-(t))dt, \quad (1.4)$$

where  $z_{\pm}(t) = \log \frac{\rho_{\pm}(t)}{1-\rho_{\pm}(t)}$ . Notice that  $I(-\bar{J}, \rho)$  does not depend on  $\rho$  (as long as  $\rho$  satisfy the boundary conditions). We also prove that the large deviations do not depend on the initial configuration.

From our analysis it follows that when imposing a smooth density profile  $\rho$  that satisfies the boundary conditions  $\rho_{\pm}(t)$  and a differentiable current  $J(t)$  with bounded derivative, the “typical path” for the process is obtained as follows. First find  $H(t, y)$  so that

$$\partial_x \rho(t, x) - \rho(t, x)(1 - \rho(t, x))\partial_y H(t, x) = -J'(t) \quad (1.5)$$

Then the process follows a weakly asymmetric exclusion process with asymmetry given by  $\frac{1}{N}\partial_y H(\frac{x}{N}, t)$ , see Section 8.

The main scheme of the proof goes along the lines of [9] and [2], but with the further feature of controlling also the deviations of the current. Using a variational characterization of  $I$ , for deviations such that  $I(J, \rho) < +\infty$ , it is possible to find suitable regular approximations  $J_{\varepsilon}$  and  $\rho_{\varepsilon}$  such that  $I(\rho_{\varepsilon}, J_{\varepsilon}, \rho_{\varepsilon}) \rightarrow I(J, \rho)$ . For regular  $J, \rho$ , it is possible to construct the weakly asymmetric exclusion dynamics as described above such that the corresponding quasi-static limit is given by  $J$  and  $\rho$  and whose relative entropy with respect to the original process converges to  $I(J, \rho)$ . This takes care of the lower bound. For the upper bound we use suitable exponential martingales that control also the current, and a superexponential estimate adapted from the original idea in [9].

## 2. SIMPLE EXCLUSION WITH BOUNDARIES

We consider the exclusion process in  $\{0, 1\}^{\Lambda_N}$ ,  $\Lambda_N := \{-N, \dots, N\}$  with reservoirs at the boundaries with density  $\rho_{\pm}(t) \in [a, 1 - a]$  for some  $a > 0$ . We assume that  $\rho_{\pm}(t) \in C^1$ .

Denoting by  $\eta(x) \in \{0, 1\}$  the occupation number at  $x \in \Lambda_N$  we define the dynamics via the generator

$$L_{N,t} = N^{2+\alpha}[L_{\text{exc}} + L_{b,t}], \quad t \geq 0, \quad \alpha > 0 \quad (2.1)$$

where for a given  $\gamma > 0$ ,

$$L_{\text{exc}}f(\eta) = \gamma \sum_{x=-N}^{N-1} \left( f(\eta^{(x,x+1)}) - f(\eta) \right) =: \sum_{x=-N}^{N-1} \nabla_{x,x+1} f(\eta) \quad (2.2)$$

$\eta^{(x,y)}$  is the configuration obtained from  $\eta$  by exchanging the occupation numbers at  $x$  and  $y$ , and

$$L_{b,t}f(\eta) = \sum_{j=\pm} \rho_j(t)^{1-\eta(jN)} (1 - \rho_j(t))^{\eta(jN)} [f(\eta^{jN}) - f(\eta)] \quad (2.3)$$

where  $\eta^x(x) = 1 - \eta(x)$ , and  $\eta^x(y) = \eta(y)$  for  $x \neq y$ .

We recall the quasi-static hydrodynamic limit proven in [6]:

**Theorem 2.1.** *For any  $\alpha > 0$  and for any macroscopic time  $t > 0$  the following holds. For any initial configuration  $\eta_0$ , for any  $r \in [-1, 1]$  and for any local function  $\varphi$*

$$\lim_{N \rightarrow \infty} \mathbb{E}_{\eta_0} \left( \theta_{[Nr]} \varphi(\eta_t) \right) = \langle \varphi(\eta) \rangle_{\bar{\rho}(r,t)} =: \hat{\varphi}(\rho(r,t)) \quad (2.4)$$

where  $[\cdot]$  denotes integer part,  $\theta$  is the shift operator,  $\langle \cdot \rangle_{\rho}$  is the expectation with respect to the product Bernoulli measure of density  $\rho$ , and

$$\bar{\rho}(y,t) = \frac{1}{2}[\rho_+(t) - \rho_-(t)]r + \frac{1}{2}[\rho_+(t) + \rho_-(t)], \quad y \in [-1, 1] \quad (2.5)$$

is the quasi-static profile of density at time  $t$ . In particular for any  $t > 0$  and any  $y \in [-1, 1]$

$$\lim_{N \rightarrow \infty} \mathbb{E}_{\eta_0} [\eta_t([Ny])] = \bar{\rho}(y,t) \quad (2.6)$$

In the following we will use the notation

$$\phi(\rho) = \rho(1 - \rho). \quad (2.7)$$

Define the following counting processes: for  $x = -N - 1, \dots, N$

$$\begin{aligned} \mathbf{h}_+(t, x) &= \{\text{number of jumps } x \rightarrow x + 1 \text{ up to time } t\}, \\ \mathbf{h}_-(t, x) &= \{\text{number of jumps } x + 1 \rightarrow x \text{ up to time } t\}, \\ \mathbf{h}(t, x) &= \mathbf{h}_+(t, x) - \mathbf{h}_-(t, x) \end{aligned} \quad (2.8)$$

When  $x = -N - 1$  the corresponding  $\mathbf{h}_+(t, -N - 1)$  is the number of particles that enters on the left boundary, and  $\mathbf{h}_+(t, N)$  is the number of particles that exit at the right boundary.

The conservation law is microscopically given by the relation

$$\eta_t(x) - \eta_0(x) = \mathbf{h}(t, x-1) - \mathbf{h}(t, x), \quad x = -N, \dots, N \quad (2.9)$$

For  $y \in [-1, 1]$ , define

$$\mathbf{h}_N(t, y) = \frac{1}{N^{1+\alpha}} \mathbf{h}(t, [Ny]).$$

Notice that, for any  $x, x' \in \{-N, \dots, N\}$ , by (2.9) we must have  $|\mathbf{h}(t, x) - \mathbf{h}(t, x')| \leq 2N$ , that implies

$$|\mathbf{h}_N(t, y) - \mathbf{h}_N(t, y')| \leq (2N)^{-\alpha}, \quad \forall y, y' \in [-1, 1]. \quad (2.10)$$

We have then

$$\begin{aligned} \frac{1}{N^{1+\alpha}} \mathbb{E}_{\eta_0}(\mathbf{h}(t, x)) &= \frac{1}{N^{1+\alpha}} \frac{1}{2N} \mathbb{E}_{\eta_0} \left( \sum_{x'=-N}^{N-1} \mathbf{h}(t, x') \right) + O(N^{-\alpha}) \\ &= \frac{1}{N^{1+\alpha}} \frac{1}{2N} \mathbb{E}_{\eta_0} \left( \gamma N^{2+\alpha} \int_0^t \sum_{x'=-N}^{N-1} (\eta_s(x') - \eta_s(x'+1)) \right) + O(N^{-\alpha}) \\ &= \frac{\gamma}{2} \mathbb{E}_{\eta_0} \left( \int_0^t (\eta_s(-N) - \eta_s(N)) ds \right) + O(N^{-\alpha}) \end{aligned}$$

Thus from (2.6)

$$\lim_{N \rightarrow \infty} \mathbb{E}_{\eta_0}(\mathbf{h}_N(t, y)) = \bar{h}(t) := -\frac{\gamma}{2} \int_0^t [\rho_+(s) - \rho_-(s)] ds. \quad (2.11)$$

Observe that the total current depends on  $\gamma$  while the quasi-static profile does not. In the sequel we will set  $\gamma = 1$ .

### 3. THE RATE FUNCTION

We denote by  $\mathcal{M} = \{\rho(t, y) \text{ measurable, } t \in [0, T], y \in [-1, 1], 0 \leq \rho(t, y) \leq 1\}$ . We endow  $\mathcal{M}$  of the weak topology, i.e. for any continuous function  $G \in \mathcal{C}([0, T] \times [-1, 1])$ ,  $\rho \mapsto \int_0^T dt \int_{-1}^1 dy \rho(t, y) G(t, y)$  is continuous on  $\mathcal{M}$ . Note that  $\mathcal{M}$  is compact under this topology.

Let  $\rho(t, y) \in \mathcal{M}$  and  $J(t) \in \mathcal{D}([0, T], \mathbb{R})$ , with  $J(0) = 0$ . The rate function is defined by:

$$I(J, \rho) = \sup_{\substack{H \in \mathcal{C}^{1,2}([0, T] \times [-1, 1]), \\ H(\cdot, -1) = 0}} \left\{ \mathcal{L}(H; J, \rho) - \int_0^T dt \int_{-1}^1 dy (\partial_y H(t, y))^2 \phi(\rho(t, y)) \right\} \quad (3.1)$$

where

$$\begin{aligned} \mathcal{L}(H; J, \rho) := & H(1, T)J(T) + \int_0^T dt \left( -\partial_t H(t, 1)J(t) - \int_{-1}^1 \partial_{yy} H(t, y)\rho(t, y)dy \right. \\ & \left. + (\partial_y H(t, 1)\rho_+(t) - \partial_y H(t, -1)\rho_-(t)) \right) \end{aligned} \quad (3.2)$$

Observe that if  $\partial_y \rho$  and  $J'$  exist and are regular enough and  $\rho(t, \pm 1) = \rho_{\pm}(t)$ , then  $I(J, \rho)$  is given by

$$I(J, \rho) = \frac{1}{4} \int_0^T \int_{-1}^1 \frac{(J'(t) + \partial_y \rho(t, y))^2}{\phi(\rho(t, y))} dy dt, \quad (3.3)$$

and the maximum is reached on the function

$$\bar{H}(t, y) = \frac{1}{2} \int_{-1}^y \frac{J'(t) + \partial_y \rho(t, y')}{\phi(\rho(t, y'))} dy'. \quad (3.4)$$

Define

$$\begin{aligned} \mathcal{H}_+ = \text{clos} \left\{ H(t, y) \in \mathcal{C}^{1,2}([0, T] \times [-1, 1]), H(\cdot, -1) = 0 : \right. \\ \left. \|H\|_{\rho,+}^2 = \int_0^T dt \int_{-1}^1 dy (\partial_y H(t, y))^2 \phi(\rho(t, y)) < +\infty \right\}. \end{aligned}$$

and the dual space

$$\begin{aligned} \mathcal{H}_- = \text{clos} \left\{ H(t, y) \in \mathcal{C}^{1,2}([0, T] \times [-1, 1]) : \right. \\ \left. \|H\|_{\rho,-}^2 = \int_0^T dt \int_{-1}^1 dy (\partial_y H(t, y))^2 \phi(\rho(t, y))^{-1} < +\infty \right\}. \end{aligned}$$

**Proposition 3.1.** *If  $I(J, \rho) < \infty$  then the weak derivatives  $J'(t)$  and  $\partial_y \rho(t, y)$  exists in  $\mathcal{H}_-$  and  $\rho(t, \pm 1) = \rho_{\pm}(t)$ , furthermore*

$$I(J, \rho) = \frac{1}{4} \int_0^T \int_{-1}^1 \frac{(J'(t) + \partial_y \rho(t, y))^2}{\phi(\rho(t, y))} dy dt. \quad (3.5)$$

*Proof.* Choose  $H(t, y) = z(t)(1 + y)$  for a given smooth function  $z(t)$  on  $[0, T]$  in the variational formula (3.1). Then defining

$$\begin{aligned} \mathcal{Q}(J, \rho) = \sup_{z \in \mathcal{C}^1([0, T])} \left\{ 2z(T)J(T) - \int_0^T (2z'(t)J(t)dt + z(t)(\rho_+(t) - \rho_-(t))) \right. \\ \left. - \int_0^T z(t)^2 \int_{-1}^1 \phi(\rho(t, y))dy \right\} \\ := \sup_z \left\{ \tilde{L}(J, z) - \int_0^T dt z(t)^2 \int_{-1}^1 \phi(\rho(t, y))dy \right\} \end{aligned} \quad (3.6)$$

we have that  $\mathcal{Q}(J, \rho) \leq I(J, \rho) < +\infty$ . It follows that

$$|\tilde{L}(J, z)|^2 \leq 2\mathcal{Q}(J, \rho) \int_0^T dt z(t)^2 \int_{-1}^1 \phi(\rho(t, y)) dy.$$

This means that  $z \rightarrow \tilde{L}(J, z)$  is a bounded linear functional on the Hilbert space  $\mathcal{N}_+$ , where

$$\mathcal{N}_\pm = \text{clos} \left\{ z \in \mathcal{C}^1([0, T]) : \|z\|^2 = \int_0^T z(t)^2 \bar{\phi}(t)^{\pm 1} dt \right\}$$

where  $\bar{\phi}(t) = \int_{-1}^1 \phi(\rho(t, y)) dy$ . Applying Riesz representation theorem, there exists a function  $g(t) \in \mathcal{N}_+$  such that

$$\tilde{L}(J, z) = \int_0^T g(t) z(t) \bar{\phi}(t) dt.$$

This implies the existence of the weak derivative  $J'(t)$  such that  $2J'(t) + (\rho_+(t) - \rho_-(t)) \in \mathcal{N}_-$ . Furthermore  $g(t)\bar{\phi}(t) = 2J'(t) + (\rho_+(t) - \rho_-(t))$  and

$$\mathcal{Q}(J, \rho) = \frac{1}{4} \int_0^T [2J'(t) + (\rho_+(t) - \rho_-(t))]^2 \bar{\phi}(t)^{-1} dt. \quad (3.7)$$

The linear functional  $\mathcal{L}(H; J, \rho)$  defined in (3.2) is bounded by

$$|\mathcal{L}(H; J, \rho)|^2 \leq 2I(J, \rho) \int_0^T dt \int_{-1}^1 dy (\partial_y H(t, y))^2 \phi(\rho(t, y)), \quad (3.8)$$

and it can be extended to a bounded linear functional on  $\mathcal{H}_+$ . By Riesz representation there exists a function  $G \in \mathcal{H}_+$  such that

$$\mathcal{L}(H; J, \rho) = \int_0^T dt \int_{-1}^1 dy \partial_y H(t, y) \partial_y G(t, y) \phi(\rho(t, y)). \quad (3.9)$$

Since we have already proven that  $I(J, \rho) < +\infty$  implies the existence of the weak derivative  $J'(t) \in \mathcal{N}_-$ , we can rewrite the variational



formula (3.1) as

$$\begin{aligned}
I(J, \rho) &= +\infty \quad \text{if } J(t) \text{ is not differentiable.} \\
I(J, \rho) &= \sup_{\substack{H \in \mathcal{C}^{1,2}([0,T] \times [-1,1]), \\ H(\cdot, -1) = 0}} \left\{ \int_0^T dt H(t, 1) J'(t) \right. \\
&\quad + \int_0^T dt (\partial_y H(t, 1) \rho_+(t) - \partial_y H(t, -1) \rho_-(t)) \\
&\quad \left. - \int_0^T dt \int_{-1}^1 dy [\partial_{yy} H(t, y) \rho(t, y) + (\partial_y H(t, y))^2 \phi(\rho(t, y))] \right\} \\
&= \sup_H \left\{ \mathcal{L}(H; J, \rho) - \int_0^T dt \int_{-1}^1 dy (\partial_y H(t, y))^2 \phi(\rho(t, y)) \right\}.
\end{aligned} \tag{3.10}$$

Notice that we can rewrite

$$\begin{aligned}
\mathcal{L}(H; J, \rho) &= - \int_0^T dt \left[ \int_{-1}^1 \partial_{yy} H(t, y) (J'(t)y + \rho(t, y)) dy \right. \\
&\quad \left. - \partial_y H(t, 1) (\rho_+(t) + J'(t)) - \partial_y H(t, -1) (\rho_-(t) - J'(t)) \right]
\end{aligned}$$

Let us first show that  $I(J, \rho) < \infty$  implies that  $\rho(t, \pm 1) = \rho_{\pm}(t)$ , in the sense that, for any continuous function  $g(t)$  on  $[0, T]$ ,

$$\begin{aligned}
\lim_{\delta \rightarrow 0} \frac{1}{\delta} \int_0^T dt \int_{-1}^{-1+\delta} dy \rho(t, y) g(t) &= \int_0^T \rho_-(t) g(t) dt, \\
\lim_{\delta \rightarrow 0} \frac{1}{\delta} \int_0^T dt \int_{1-\delta}^1 dy \rho(t, y) g(t) &= \int_0^T \rho_+(t) g(t) dt.
\end{aligned}$$

In fact assume that these boundary conditions are not satisfied, and choose the functions  $H(s, y)$  such that

$$\partial_y H(t, y) = A \left( 1 - \frac{1-y}{\delta} \right) 1_{[1-\delta, 1]}(y) g(t)$$

This function is not smooth, but it can be smoothed up by some convolution without changing the argument. On this function we have

$$\begin{aligned}
\mathcal{L}(H; J, \rho) &= -\frac{A}{\delta} \int_0^T dt g(t) \int_{1-\delta}^1 dy (J'(t)y + \rho(t, y)) + A \int_0^T g(t) (\rho_+(t) + J'(t)) dt \\
&= -\frac{1}{2} A \delta \int_0^T g(t) J'(t) dt - A \int_0^T dt g(t) \left( \delta^{-1} \int_{1-\delta}^1 dy \rho(t, y) - \rho_+(t) \right)
\end{aligned}$$

while we have

$$\begin{aligned} & \int_0^T dt \int_{-1}^1 dy (\partial_y H(t, y))^2 \phi(\rho(t, y)) \\ &= A^2 \int_0^T dt g(t)^2 \int_{1-\delta}^1 dy \left(1 - \frac{1-y}{\delta}\right)^2 \phi(\rho(t, y)) \\ & \leq CA^2 T \delta. \end{aligned} \quad (3.11)$$

Then if the boundary condition in  $y = 1$  is not satisfied, one can construct a sequence of functions  $H$  such that  $I(J, \rho) = +\infty$ . The other side is treated in a similar way.

Now that the boundary conditions must be satisfied, it follows from (3.9) that  $I(J, \rho) < \infty$  implies the existence of the weak derivative  $\partial_x \rho(t, x)$  such that we can identify  $\partial_x G(t, x) \phi(\rho(t, x)) = J'(t) + \partial_x \rho(t, x)$ .

Notice that, from this identification we can rewrite  $I$  as

$$\begin{aligned} I(J, \rho) &= \frac{1}{4} \int_0^T \int_{-1}^1 \frac{J'(t)^2 + \partial_y \rho(t, y)^2}{\phi(\rho(t, y))} dy dt \\ &+ \frac{1}{2} \int_0^T J'(t) \log \left( \frac{\rho_+(t)(1 - \rho_-(t))}{\rho_-(t)(1 - \rho_+(t))} \right) dt \end{aligned} \quad (3.12)$$

that proves that  $J$  and  $\rho$  are in  $\mathcal{H}_-$ , since  $\rho_{\pm}(t)$  are assumed bounded away from 0 and 1.  $\square$

**Proposition 3.2.** *If  $I(J, \rho) < \infty$ , there exists an approximation by bounded functions  $J_{\varepsilon}(t)$  and  $x$ -smooth  $\rho_{\varepsilon}(t, x)$  of  $J$  and  $\rho$ , such that  $\rho_{\varepsilon}(t, \pm 1) = \rho_{\pm}(t)$  and*

$$\lim_{\varepsilon \rightarrow 0} I(J_{\varepsilon}, \rho_{\varepsilon}) = I(J, \rho). \quad (3.13)$$

*Proof.* We follow a similar argument used in [2].

We proceed in two steps. We first define an approximation  $\tilde{\rho}_{\varepsilon}(t, x)$  bounded away from 0 and 1 such that  $\phi(\tilde{\rho}_{\varepsilon}(t, x)) \geq C\varepsilon^2$  and  $I(J, \tilde{\rho}_{\varepsilon}) \rightarrow I(J, \rho)$ . Then assuming that  $\phi(\rho(t, x)) \geq a > 0$ , we construct a smooth approximation  $\rho_{\varepsilon}(t, x)$  such that  $I(J, \rho_{\varepsilon}) \rightarrow I(J, \rho)$ . The conclusion follows then by a diagonal argument.

For the first step, define (recall (2.5))

$$\tilde{\rho}_{\varepsilon}(t, x) = (1 - \varepsilon)\rho(t, x) + \varepsilon\bar{\rho}(t, x). \quad (3.14)$$

Notice that  $\tilde{\rho}_{\varepsilon}(t, \pm 1) = \rho_{\pm}(t)$  and

$$\phi(\tilde{\rho}_{\varepsilon}(t, x)) \geq \varepsilon^2 (\rho_-(t) \wedge \rho_+(t)) (1 - (\rho_-(t) \vee \rho_+(t))).$$

Since  $I(J, \rho)$  is convex in  $\rho$ , we have

$$I(J, \tilde{\rho}_{\varepsilon}) \leq (1 - \varepsilon)I(J, \rho) + \varepsilon I(J, \bar{\rho}) \xrightarrow{\varepsilon \rightarrow 0} I(J, \rho), \quad (3.15)$$

while by lower semicontinuity

$$\liminf_{\varepsilon \rightarrow 0} I(J, \tilde{\rho}_\varepsilon) \geq I(J, \rho), \quad (3.16)$$

that concludes the first approximation step.

Now assuming that  $\phi(\rho(t, x)) \geq \phi(a) > 0$ , i.e.  $a < \rho(t, x) < 1 - a$ , we construct a smooth approximation  $\rho_\varepsilon(t, x)$  such that

$$\rho_\varepsilon(t, x) \xrightarrow{\varepsilon \rightarrow 0} \rho(t, y), \quad \rho_\varepsilon(t, \pm 1) = \rho_\pm(t).$$

and such that  $I(J, \rho_\varepsilon) \rightarrow I(J, \rho)$ .

Let be  $\Delta_D$  the laplacian on  $[-1, 1]$  with Dirichlet boundary conditions, and

$$R_\varepsilon^D(x, y) = (I - \varepsilon \Delta_D)^{-1}(x, y).$$

Then define

$$\rho_\varepsilon(t, x) = \bar{\rho}(t, x) + \int_{-1}^1 R_\varepsilon^D(x, y) (\rho(t, y) - \bar{\rho}(t, y)) dy. \quad (3.17)$$

We next prove that there exist  $a_* > 0$  such that

$$\phi(\rho_\varepsilon(t, x)) \geq \phi(a_*). \quad (3.18)$$

In fact notice that  $0 \leq \bar{\rho}(t, x) - \int_{-1}^1 R_\varepsilon^D(x, y) \bar{\rho}(t, y) dy \leq 1$ , that implies

$$\rho_\varepsilon(t, x) \geq \int_{-1}^1 R_\varepsilon^D(x, y) \rho(t, y) dy \geq a \int_{-1}^1 R_\varepsilon^D(x, y) dy \geq a',$$

for some positive  $a' < a$ .

Similarly we have

$$\begin{aligned} 1 - \rho_\varepsilon(t, x) &= 1 - \bar{\rho}(t, x) - \int_{-1}^1 R_\varepsilon^D(x, y) (1 - \bar{\rho}(t, y)) dy + \int_{-1}^1 R_\varepsilon^D(x, y) (1 - \rho(t, y)) dy \\ &\geq \int_{-1}^1 R_\varepsilon^D(x, y) (1 - \rho(t, y)) dy \geq (1 - a) \int_{-1}^1 R_\varepsilon^D(x, y) dy \geq (1 - a''). \end{aligned}$$

Then choosing  $a_* = a' \wedge a''$  we obtain (3.18).

Since  $\rho_\varepsilon$  is smooth, by (3.10), we have:

$$I(J, \rho_\varepsilon) = \frac{1}{4} \int_0^T dt \int_{-1}^1 dy (J'(t) + \partial_y \rho_\varepsilon(t, y))^2 \phi(\rho_\varepsilon(t, y))^{-1} \quad (3.19)$$

Let  $R_\varepsilon^N(x, y) = (I - \varepsilon \Delta_N)^{-1}(x, y)$  the resolvent for the laplacian with Neumann boundary conditions, then we have the property that

$$\partial_x R_\varepsilon^D(x, y) = -\partial_y R_\varepsilon^N(x, y), \quad (3.20)$$

that implies

$$\partial_x \rho_\varepsilon(t, x) = \int_{-1}^1 R_\varepsilon^N(x, y) \partial_y \rho(t, y) dy := (R_\varepsilon^N \partial_y \rho)(t, x). \quad (3.21)$$

Furthermore  $R_\varepsilon^N(x, y)$  is a probability kernel:  $\int_{-1}^1 R_\varepsilon^N(x, y) dy = 1$ .

Since  $R_\varepsilon^N(x, y) \leq 1$ , by convexity we have

$$\begin{aligned} (J'(t) + \partial_y \rho_\varepsilon(t, y))^2 &\leq \int_{-1}^1 R_\varepsilon^N(x, y) (J'(t) + \partial_y \rho(t, x))^2 dx \\ &\leq \int_{-1}^1 (J'(t) + \partial_y \rho(t, x))^2 dx \leq \frac{1}{4} I(J, \rho) \end{aligned} \quad (3.22)$$

Then we have

$$\frac{(J'(t) + \partial_y \rho_\varepsilon(t, y))^2}{\phi(\rho_\varepsilon(t, y))} \leq \frac{1}{4\phi(a_*)} I(J, \rho) \quad (3.23)$$

Then by dominated convergence we have

$$\lim_{\varepsilon \rightarrow 0} \mathcal{I}(J, \rho_\varepsilon) = \mathcal{I}(J, \rho).$$

It remains to prove the  $J'$  approximation by bounded functions. Let us define

$$J_k(t) = \int_0^t J'(s) \wedge k \vee (-k) ds \quad (3.24)$$

and let us assume that  $\rho(t, x)$  is smooth in  $x$  and bounded away from 0 and 1. Then it is clear that

$$\frac{(J'(t) \wedge k \vee (-k) + \partial_y \rho(t, y))^2}{\phi(\rho(t, y))}$$

is not decreasing in  $k$  for  $k$  large enough. Then by monotone convergence  $I(J_k, \rho) \rightarrow I(J, \rho)$ .  $\square$

Notice that the minimum of  $I(J, \rho)$  is correctly achieved for  $J'(t) = \bar{J}'(t) = -\frac{1}{2} [\rho_+(t) - \rho_-(t)]$  and  $\rho(t, y) = \bar{\rho}(t, y)$ .

Furthermore, from (3.12), we have that for any given  $\rho$ ,  $I(J, \rho)$  is a quadratic function of  $J'$ . In particular we have the following Gallavotti-Cohen type of symmetry:

$$I(-J, \rho) = I(J, \rho) - \int_0^T J'(t) \log \frac{\rho_+(t)(1 - \rho_-(t))}{\rho_-(t)(1 - \rho_+(t))} dt, \quad (3.25)$$

The rate function for  $\mathfrak{h}_N$  alone is obtained, by contraction principle (cf. [12]), minimizing over all possible  $\rho$  satisfying the given boundary conditions:

$$I(J) = \inf_{\rho} I(J, \rho), \quad (3.26)$$

and also satisfy the symmetry relation

$$I(-J) = I(J) - \int_0^T J'(t) \log \frac{\rho_+(t)(1 - \rho_-(t))}{\rho_-(t)(1 - \rho_+(t))} dt. \quad (3.27)$$

For  $\rho_+, \rho_-$  and current  $J'$  constant in time, and the stationary profile  $\bar{\rho}(\cdot)$ , we have

$$I(J, \bar{\rho}) = \frac{T}{4} \left( J' + \frac{1}{2} [\rho_+ - \rho_-] \right)^2 \int_{-1}^1 (\bar{\rho}(y))(1 - \bar{\rho}(y))^{-1} dy \quad (3.28)$$

in particular

$$I(-\bar{J}, \bar{\rho}) = \frac{T}{4} (\rho_+ - \rho_-)^2 \int_{-1}^1 (\bar{\rho}(y))(1 - \bar{\rho}(y))^{-1} dy, \quad (3.29)$$

while from (3.28)

$$I(-\bar{J}) = \frac{T}{2} (\rho_+ - \rho_-) \log \frac{\rho_+(1 - \rho_-)}{\rho_-(1 - \rho_+)}. \quad (3.30)$$

**3.1. Relation with the Bodineau-Derrida functional.** Bodineau and Derrida [3] write the LD rate for the density field  $\rho(t, y)$  and a current field  $j(t, y)$  such that  $\partial_t \rho(t, y) = -\partial_y j(t, y)$  (the conservation law), as

$$I_{bd}(j, \rho) = \frac{1}{4} \int_0^T \int_{-1}^1 \frac{(j(t, y) + \partial_y \rho(t, y))^2}{\phi(\rho(t, y))} dy dt. \quad (3.31)$$

The formula (3.3) looks similar, but differs in an important point. In the quasistatic case the current field (*instantaneous currents*)  $j(t, y)$  has to be constant in  $y$  and it is not related to the time derivative of  $\rho$ : we cannot have large deviations where the intensity of the current is not constant in space, see (2.10).

A direct connection with the Bodineau-Derrida functional can be found when the boundary conditions  $\rho_{\pm}$  do not change in time, minimizing over  $\rho(t, y)$  in order to obtain the rate function  $I(J)$  for the large deviation of the only current  $h_N$ . If one looks at deviation such that  $J'$  is constant in time, the minimizing  $\rho$  will also time independent and the rate  $I(J)$  will coincide, up to a factor  $T$ , to the expression given in [3] formula (3.2) for the large deviations of the total current averaged in time.

## 4. THE LARGE DEVIATION THEOREM

Let us define the empirical density profile

$$\pi_N(t, y) = \sum_{x=-N}^{N-1} \eta_x(t) 1_{[x, x+1)}(Ny), \quad y \in [-1, 1] \quad (4.1)$$

that has values on  $\mathcal{M}$ , so that the couple  $(\mathbf{h}_N, \pi_N)$  has values on  $\Xi = \mathcal{D}([0, T], \mathbb{R}) \times \mathcal{M}$ .

**Theorem 4.1.** *Under the dynamics generated by (2.1), starting with an arbitrary configuration  $\eta$ , the couple  $(\mathbf{h}_N, \pi_N)$  satisfy a large deviation principle with rate function  $I(J, \rho)$ , i.e.*

- For any closed set  $C \subset \Xi$

$$\limsup_{N \rightarrow \infty} \frac{1}{N^{1+\alpha}} \log P_\eta((\mathbf{h}_N, \pi_N) \in C) \leq - \inf_{(J, \rho) \in C} I(J, \rho), \quad (4.2)$$

- For any open set  $\mathcal{O} \subset \Xi$

$$\liminf_{N \rightarrow \infty} \frac{1}{N^{1+\alpha}} \log P_\eta((\mathbf{h}_N, \pi_N) \in \mathcal{O}) \geq - \inf_{(J, \rho) \in \mathcal{O}} I(J, \rho), \quad (4.3)$$

## 5. THE SUPEREXPONENTIAL ESTIMATE

One of the main steps in the proof of Theorem 4.1 is a super-exponential estimate which allows the replacement of local functions by functionals of the empirical density. Define

$$V_{N, \varepsilon}(t, \eta) = \sum_{x=-N}^{N-1} G(t, x/N) (\eta(x)(1 - \eta(x+1)) - \phi(\bar{\eta}_{N, \varepsilon}(x/N))) \quad (5.1)$$

The local averages are defined in the bulk as

$$\bar{\eta}_{N, \varepsilon}(x/N) = \frac{1}{2N\varepsilon + 1} \sum_{|x' - x| \leq \varepsilon N} \eta(x'), \quad |x| < N(1 - \varepsilon), \quad (5.2)$$

and at the boundaries as

$$\begin{aligned} \bar{\eta}_{N, \varepsilon}(x/N) &= \rho_+(t) & x &\geq N(1 - \varepsilon), \\ \bar{\eta}_{N, \varepsilon}(x/N) &= \rho_-(t) & x &\leq -N(1 - \varepsilon). \end{aligned} \quad (5.3)$$

Notice that  $V_{N, \varepsilon}(t, \eta)$  depends on  $t$  not only by the function  $G(t, y)$  but also by the special definition (5.3).

In the following we use as reference measures the inhomogeneous product measures

$$\mu_t(\eta) = \prod_{x=-N}^N \bar{\rho}\left(\frac{x}{N}, t\right)^{\eta(x)} \left[1 - \bar{\rho}\left(\frac{x}{N}, t\right)\right]^{1-\eta(x)}. \quad (5.4)$$

Observe that, since  $\rho_{\pm}(t)$  are uniformly away from 0 and 1, there is  $C > 0$  so that

$$\sup_t \sup_{\eta} \mu_t(\eta) \geq e^{-CN}. \quad (5.5)$$

**Proposition 5.1.** *For any  $\delta > 0$  and initial configuration  $\eta$*

$$\lim_{\varepsilon \rightarrow 0} \limsup_{N \rightarrow \infty} \frac{1}{N^{1+\alpha}} \log \mathbb{P}_{\eta} \left( \int_0^T V_{N,\varepsilon}(t, \eta_t) dt \geq N\delta \right) = -\infty. \quad (5.6)$$

*Proof.* By (5.5), it is enough to prove (5.6) for the dynamics with initial distribution given by  $\mu_0$ . By exponential Chebichef inequality we have for any  $a > 0$ :

$$\mathbb{P}_{\mu_0} \left( \int_0^T V_{N,\varepsilon}(t, \eta_t) dt \geq N\delta \right) \leq e^{-N^{1+\alpha} \delta a} \mathbb{E}_{\mu_0} \left( e^{N^{\alpha} \int_0^T a V_{N,\varepsilon}(t, \eta_t) dt} \right)$$

All we need to prove is that there exists  $K < +\infty$  such that for all  $a > 0$

$$\limsup_{\varepsilon \rightarrow 0} \limsup_{N \rightarrow \infty} \frac{1}{N^{1+\alpha}} \log \mathbb{E}_{\mu_0} \left( e^{N^{\alpha} \int_0^T a V_{N,\varepsilon}(t, \eta_t) dt} \right) \leq KT$$

To simplify notations, denote  $\tilde{V}(t, \eta) = N^{\alpha} a V_{N,\varepsilon}(t, \eta)$ . Consider the equation

$$\partial_s u(\eta, s) = L_{N,T-s} u(\eta, s) + \tilde{V}(T-s, \eta) u(\eta, s), \quad u(\eta, 0) = 1, \quad 0 \leq s \leq T. \quad (5.7)$$

By Feynman-Kac formula

$$u(\eta, T) = \mathbb{E}_{\eta} \left( e^{\int_0^T \tilde{V}(s, \eta_s) ds} \right) \quad (5.8)$$

Then

$$\begin{aligned} \frac{d}{ds} \frac{1}{2} \sum_{\eta} u(\eta, s)^2 \mu_{T-s}(\eta) &= \frac{1}{2} \sum_{\eta} u(\eta, s)^2 \frac{d\mu_{T-s}(\eta)}{ds} \\ &+ \sum_{\eta} \left( u(\eta, s) L_{N,T-s} u(\eta, s) + \tilde{V}(T-s, \eta) u^2(\eta, s) \right) \mu_{T-s}(\eta) \end{aligned} \quad (5.9)$$

and, since  $\left| \frac{d\mu_{T-s}(\eta)}{ds} \right| \leq CN\mu_{T-s}(\eta)$ , this is bounded by

$$\leq (CN + \Gamma(s)) \sum_{\eta} u(\eta, s)^2 \mu_{T-s}(\eta),$$

where, setting  $\|f\|_s^2 = \sum_{\eta} f(\eta)^2 \mu_{T-s}(\eta)$ ,

$$\Gamma(s) = \sup_{f, \|f\|_s=1} \left\{ \sum_{\eta} \tilde{V}(T-s, \eta) f^2(\eta) \mu_{T-s}(\eta) + \sum_{\eta} f(\eta) L_{N, T-s} f(\eta) \mu_{T-s}(\eta) \right\}. \quad (5.10)$$

By Gronwall inequality and (5.9) we have

$$\sum_{\eta} u(\eta, T)^2 \mu_0(\eta) \leq e^{2 \int_0^T \Gamma(s) ds + TCN} \sum_{\eta} u(\eta, 0)^2 \mu_T(\eta) = e^{2 \int_0^T \Gamma(t) dt + TCN}. \quad (5.11)$$

Then using Schwarz inequality we get

$$\mathbb{E}_{\mu_0} \left( e^{\int_0^T \tilde{V}(\eta(t)) dt} \right) \leq e^{\int_0^T \Gamma(t) dt + TCN}. \quad (5.12)$$

The Dirichlet forms associated to the generator are

$$\begin{aligned} \mathfrak{D}_{x,t,\rho}(f) &= \frac{1}{2} \sum_{\eta} \rho^{1-\eta(x)} (1-\rho)^{\eta(x)} [f(\eta^x) - f(\eta)]^2 \mu_t(\eta), \quad x = \pm N \\ \mathfrak{D}_{ex,t}(f) &= \frac{1}{2} \sum_{\eta} \sum_{x=-N}^{N-1} (\nabla_{x,x+1} f(\eta))^2 \mu_t(\eta) \end{aligned} \quad (5.13)$$

and we define

$$\mathfrak{D}_t(f) = - \sum_{\eta} f(\eta) L_{N,t} f(\eta) \mu_t(\eta) = \mathfrak{D}_{-N,t,\rho_-(t)}(f) + \mathfrak{D}_{N,t,\rho_+(t)}(f) + \mathfrak{D}_{ex,t}(f).$$

Observe that

$$\begin{aligned} \sum_{\eta} f(\eta) (L_{N,t} f)(\eta) \mu_t(\eta) &= -N^{2+\alpha} \sum_{\eta} \sum_{x=-N}^{N-1} \nabla_{x,x+1} f(\eta) \nabla_{x,x+1} (f \mu_t)(\eta) \\ &\quad - N^{2+\alpha} (\mathfrak{D}_{N,t,\rho_+(t)}(f) + \mathfrak{D}_{-N,t,\rho_-(t)}(f)) \end{aligned} \quad (5.14)$$



and

$$\begin{aligned}
\sum_{\eta} \sum_{x=-N}^{N-1} \nabla_{x,x+1} f(\eta) \nabla_{x,x+1} (f \mu_t)(\eta) &= \sum_{\eta} \sum_{x=-N}^{N-1} (\nabla_{x,x+1} f)^2 \mu_t(\eta) \\
&\quad + \sum_{\eta} \sum_{x=-N}^{N-1} f(\eta^{x,x+1}) \nabla_{x,x+1} f(\eta) \nabla_{x,x+1} \mu_t(\eta) \\
&= 2\mathfrak{D}_{ex,t}(f) + \frac{1}{N} \tilde{B}_N(t) + O(N^{-1}) \tag{5.15}
\end{aligned}$$

where

$$\tilde{B}_N(t) = \sum_{\eta} \sum_{x=-N}^{N-1} f(\eta^{x,x+1}) \nabla_{x,x+1} f(\eta) (\eta(x) - \eta(x+1)) B\left(\frac{x}{N}, t\right) \mu_t(\eta)$$

and

$$B\left(\frac{x}{N}, t\right) = \frac{\bar{\rho}'\left(\frac{x}{N}, t\right)}{\bar{\rho}\left(\frac{x}{N}, t\right)(1 - \bar{\rho}\left(\frac{x}{N}, t\right))}, \quad \|B\|_{\infty} \leq c \|\rho'\|_{\infty} \tag{5.16}$$

where  $c > 0$  since  $\rho_{\pm}(t)$  are uniformly away from 0 and 1. By an elementary inequality we have:

$$\begin{aligned}
\left| \tilde{B}_N(t) \right| &\leq \frac{N}{2} \mathfrak{D}_{ex,t}(f) + \frac{2}{2N} \sum_{\eta} \sum_{x=-N}^{N-1} f(\eta^{x,x+1})^2 (\eta(x) - \eta(x+1))^2 B\left(\frac{x}{N}, t\right)^2 \mu_t(\eta) \\
&\leq \frac{N}{2} \mathfrak{D}_{ex,t}(f) + \frac{1}{N} \sum_{x=-N}^{N-1} B\left(\frac{x}{N}, t\right)^2 \sum_{\eta} f^2(\eta) \mu_t(\eta^{x,x+1})
\end{aligned}$$

Since  $\mu_t(\eta^{x,x+1}) = F(x, t) \mu_t(\eta)$  with  $F(x, t)$  a uniformly bounded function of  $\bar{\rho}\left(\frac{x}{N}, t\right)$  and  $\bar{\rho}\left(\frac{x+1}{N}, t\right)$  and  $\eta$ , using (5.16) we get

$$\left| \tilde{B}_N(t) \right| \leq \frac{N}{2} \mathfrak{D}_{ex}(f) + C \leq \frac{N}{2} \mathfrak{D}_t(f) + C \tag{5.17}$$

From (5.14), (5.15) and (5.17) we get

$$\begin{aligned}
\Gamma(s) &\leq \sup_f \left\{ N^{\alpha} a \sum_{\eta} V_{N,\varepsilon}(T-s, \eta) f^2(\eta) \mu_{T-s}(\eta) - N^{2+\alpha} \frac{1}{2} \mathfrak{D}_{T-s}(f) \right\} + N^{1+\alpha} C \\
&\leq N^{1+\alpha} \sup_f \left\{ \frac{a}{N} \sum_{\eta} V_{N,\varepsilon}(T-s, \eta) f^2(\eta) \mu_{T-s}(\eta) - N \frac{1}{2} \mathfrak{D}_{T-s}(f) \right\} + N^{1+\alpha} C \tag{5.18}
\end{aligned}$$

Then we are left to prove that for any  $a > 0$  and any  $t > 0$ :

$$\limsup_{\varepsilon \rightarrow 0} \limsup_{N \rightarrow \infty} \sup_f \left\{ \frac{1}{N} \sum_{\eta} a V_{N,\varepsilon}(t, \eta) f^2(\eta) \mu_t(\eta) - \frac{N}{2} \mathfrak{D}_t(f) \right\} \leq C' \quad (5.19)$$

This follows by proving that, for a suitable constant  $C$

$$\limsup_{\varepsilon \rightarrow 0} \limsup_{N \rightarrow \infty} \sup_{\mathfrak{D}_t(f) \leq CN^{-1}} \frac{1}{N} \sum_{\eta} V_{N,\varepsilon}(t, \eta) f^2(\eta) \mu_t(\eta) \leq 0 \quad (5.20)$$

The rest of the proof is identical as in theorem 3.1 in [6] (see pages 1051-2).  $\square$

With a similar argument follows also the super-exponential control of the densities at the boundaries:

**Proposition 5.2.** *For any  $\delta > 0$  we have*

$$\limsup_{\varepsilon \rightarrow 0} \limsup_{N \rightarrow \infty} \frac{1}{N^{1+\alpha}} \log \mathbb{P}_{\eta} \left( \int_0^T \left| \frac{1}{\varepsilon N} \sum_{\pm N}^{\pm N(1 \mp \varepsilon)} \eta_t(x) - \rho_{\pm}(t) \right| \geq \delta \right) = -\infty \quad (5.21)$$

## 6. THE EXPONENTIAL MARTINGALES

We use the following notations: for  $x = -N, \dots, N-1$

$$\phi_{-}(\eta, x) = \eta(x+1)(1-\eta(x)), \quad \phi_{+}(\eta, x) = \eta(x)(1-\eta(x+1)), \quad (6.1)$$

and at the boundaries

$$\begin{aligned} \phi_{-}(\eta, -N-1, t) &:= (1-\rho_{-}(t))\eta(-N) \\ \phi_{+}(\eta, -N-1, t) &:= \rho_{-}(t)(1-\eta(-N)) \\ \phi_{-}(\eta, N, t) &:= \rho_{+}(t)(1-\eta(N)) \\ \phi_{+}(\eta, N, t) &:= (1-\rho_{+}(t))\eta(N). \end{aligned} \quad (6.2)$$

Given two function  $z_{\pm}(t, y)$ , we associate the following exponential martingales, for  $x = -N-1, \dots, N$

$$\begin{aligned} \mathcal{E}_{\pm}(z_{\pm}, x, T) &= \\ \exp \left\{ \int_0^T z_{\pm}(t, x/N) dh_{\pm}(t, x) - \int_0^T N^{2+\alpha} (e^{z_{\pm}(t, x/N)} - 1) \phi_{\pm}(\eta_t, x) dt \right\} \\ &= \exp \left\{ z_{\pm}(T, x/N) h_{\pm}(T, x) \right. \\ &\quad \left. - \int_0^T \left[ \partial_t z_{\pm}(t, x/N) h_{\pm}(t, x) + N^{2+\alpha} (e^{z_{\pm}(t, x/N)} - 1) \phi_{\pm}(\eta_t, x) \right] dt \right\} \end{aligned} \quad (6.3)$$

We now choose a smooth function  $H(t, y)$ ,  $y \in [-1, 1]$  such that  $H(t, -1) = 0$  and we set

$$z_+(t, y) = H(t, y + 1/N) - H(t, y) = -z_-(t, y),$$

and furthermore

$$z_+(t, -1-y) := -y\partial_y H(t, -1), \quad z_+(t, 1+y) := y\partial_y H(t, 1) - H(t, 1), \quad y \geq 0.$$

The martingales defined by (6.3) are orthogonal, consequently taking the product we still have an exponential martingale equal to

$$\begin{aligned} & \prod_{x=-N-1}^N \prod_{\sigma=\pm} \mathcal{E}_\sigma(z_\sigma, x, T) = \\ & \exp \left\{ \sum_{x=-N-1}^N \left[ z_+(T, x/N) \mathbf{h}(T, x) - \int_0^T \partial_t z_+(t, x/N) \mathbf{h}(t, x) dt \right. \right. \\ & \left. \left. - N^{2+\alpha} \int_0^T \left( (e^{z_+(t, x/N)} - 1) \phi_+(\eta_t, x) + (e^{-z_+(t, x/N)} - 1) \phi_-(\eta_t, x) \right) dt \right] \right\} \end{aligned} \quad (6.4)$$

For large  $N$  we use Taylor approximation, for  $x = -N, \dots, N-1$ ,

$$\begin{aligned} & (e^{z_+(t, x/N)} - 1) \phi_+(\eta_t, x) + (e^{-z_+(t, x/N)} - 1) \phi_-(\eta_t, x) \\ & = -z_+(t, x/N) (\eta_t(x+1) - \eta_t(x)) + \frac{1}{2} z_+(t, x/N)^2 (\phi_+(\eta_t, x) + \phi_-(\eta_t, x)) + O(z_+(t, x/N)^3). \end{aligned}$$

and on the boundaries

$$\begin{aligned} & (e^{z_+(t, -1-1/N)} - 1) \phi_+(\eta_t, -N-1, t) + (e^{-z_+(t, -1-1/N)} - 1) \phi_-(\eta_t, -N-1, t) \\ & = -z_+(t, -1-1/N) (\eta_t(-N) - \rho_-(t)) \\ & + \frac{1}{2} z_+(t, -1-1/N)^2 (\phi_+(\eta_t, -N-1, t) + \phi_-(\eta_t, -N-1, t)) + O(z_+(t, -1-1/N)^3). \end{aligned}$$

$$\begin{aligned} & (e^{z_+(t, 1)} - 1) \phi_+(\eta_t, N) + (e^{-z_+(t, 1)} - 1) \phi_-(\eta_t, N) \\ & = -z_+(t, 1) (\rho_+(t) - \eta(t, N)) + \frac{1}{2} z_+(t, 1)^2 (\phi_+(\eta_t, N) + \phi_-(\eta_t, N)) + O(z_+(t, 1)^3). \end{aligned}$$

We can rewrite the exponential martingale (6.4) as

$$\begin{aligned}
& \prod_{x=-N-1}^N \prod_{\sigma=\pm} \mathcal{E}_\sigma(z_\sigma, x, T) = \\
& \exp \left\{ \sum_{x=-N-1}^N \left[ z_+(T, x/N) \mathbf{h}(T, x) - \int_0^T \partial_t z_+(t, x/N) \mathbf{h}(t, x) dt \right] \right. \\
& \left. + N^{2+\alpha} \int_0^T \sum_{x=-N-1}^N \left[ z_+(t, x/N) (\eta_t(x+1) - \eta_t(x)) \right. \right. \\
& \quad \left. \left. - \frac{1}{2} z_+(t, x/N)^2 (\phi_+(\eta_t, x) + \phi_-(\eta_t, x)) \right] dt + O(N^\alpha) \right\} \tag{6.5}
\end{aligned}$$

where in the above expression we set the convention  $\eta_t(N+1) := \rho_+(t)$  and  $\eta_t(-N-1) := \rho_-(t)$ .

After a summation by parts we have

$$\begin{aligned}
& \exp \left\{ \sum_{x=-N-1}^N \left[ z_+(T, x/N) \mathbf{h}(T, x) - \int_0^T \partial_t z_+(t, x/N) \mathbf{h}(t, x) dt \right] \right. \\
& \left. + N^{2+\alpha} \int_0^T \sum_{x=-N}^N \left[ \left( z_+(t, \frac{x-1}{N}) - z_+(t, \frac{x}{N}) \right) \eta_t(x) \right. \right. \\
& \quad \left. \left. + z_+(t, 1) \rho_+(t) - z_+(t, -1 - 1/N) \rho_-(t) \right. \right. \\
& \quad \left. \left. - \frac{1}{2} z_+(t, x/N)^2 (\phi_+(\eta_t, x) + \phi_-(\eta_t, x)) \right] dt + O(N^\alpha) \right\} \tag{6.6}
\end{aligned}$$

Using (2.9) and the special expression for  $z_+$ , we have for the logarithm of (6.6)

$$\begin{aligned}
& - \sum_{x=-N}^N H(T, x/N) [\eta_T(x) - \eta_0(x)] \\
& \quad + [H(T, 1 + 1/N) \mathbf{h}(T, N) - H(T, -1 - 1/N) \mathbf{h}(T, -N - 1)] \\
& + \int_0^T \sum_{x=-N}^N \partial_t H(t, x/N) [\eta_t(x) - \eta_0(x)] dt \\
& - \int_0^T [\partial_t H(t, 1 + 1/N) \mathbf{h}(t, N) - \partial_t H(t, -1 - 1/N) \mathbf{h}(t, -N - 1)] dt \\
& - N^\alpha \int_0^T \left( \sum_{x=-N-1}^N \left[ \partial_{yy} H(t, \frac{x}{N}) \eta_t(x) + \frac{1}{2} \left( \partial_y H(t, \frac{x}{N}) \right)^2 (\phi_+(\eta_t, x) + \phi_-(\eta_t, x)) \right] \right. \\
& \quad \left. + N (\partial_y H(t, 1) \rho_+(t) - \partial_y H(t, -1) \rho_-(t)) \right) dt + O(N^\alpha)
\end{aligned}$$

$$\begin{aligned}
&= - \sum_{x=-N}^N H(T, x/N) [\eta_T(x) - \eta_0(x)] \\
&\quad + \left[ H(T, 1 + 1/N) \mathbf{h}(T, N) + \frac{1}{N} \partial_y H(T, -1) \mathbf{h}(T, -N - 1) \right] \\
&\quad + \int_0^T \sum_{x=-N}^N \partial_t H(t, x/N) [\eta_t(x) - \eta_0(x)] dt \\
&\quad - \int_0^T \left[ \partial_t H(t, 1 + 1/N) \mathbf{h}(t, N) + \frac{1}{N} \partial_t \partial_y H(t, -1) \mathbf{h}(t, -N - 1) \right] dt \\
&\quad - N^\alpha \int_0^T \left( \sum_{x=-N-1}^N \left[ \partial_{yy} H(t, \frac{x}{N}) \eta_t(x) + \frac{1}{2} \left( \partial_y H(t, \frac{x}{N}) \right)^2 (\phi_+(\eta_t, x) + \phi_-(\eta_t, x)) \right] \right. \\
&\quad \quad \left. + N (\partial_y H(t, 1) \rho_+(t) - \partial_y H(t, -1) \rho_-(t)) \right) dt + O(N^\alpha)
\end{aligned}$$

Since we are interested only in the terms that have order  $N^{1+\alpha}$ , we can forget all terms of order  $N$  in the above expression, and the exponential martingale has the form:

$$\begin{aligned}
&\exp N^{1+\alpha} \left\{ H(T, 1) \mathbf{h}_N(T, N) - \int_0^T \partial_t H(t, 1) \mathbf{h}_N(t, N) dt \right. \\
&\quad - \int_0^T \left( \frac{1}{N} \sum_{x=-N-1}^N \partial_{yy} H(t, \frac{x}{N}) \eta_t(x) + \partial_y H(t, 1) \rho_+(t) - \partial_y H(t, -1) \rho_-(t) \right. \\
&\quad \left. \left. + \frac{1}{2N} \sum_{x=-N-1}^N \left( \partial_y H(t, \frac{x}{N}) \right)^2 (\phi_+(\eta_t, x) + \phi_-(\eta_t, x)) \right) dt + O(N^{-(\alpha \wedge 1)}) \right\}.
\end{aligned} \tag{6.7}$$

After the superexponential estimate proved in the previous section, that also fix the densities at the boundaries, it follows the variational formula for the rate function given by (3.1).

## 7. THE UPPER BOUND

**7.1. Exponential tightness.** The following proposition uses standard arguments and we give a proof for completeness in Appendix A:

**Proposition 7.1.** *There exist a sequence of compact sets  $K_L$  in  $\mathcal{D}([0, T], \mathbb{R})$  such that*

$$\lim_{L \rightarrow \infty} \lim_{N \rightarrow \infty} \frac{1}{N^{1+\alpha}} \log \mathbb{P}_N (\mathbf{h}_N \in K_L^c) = -\infty. \tag{7.1}$$

**7.2. Proof of the upper bound.** By (6.7) we can rewrite the exponential martingale defined by (6.5) as

$$\begin{aligned}
M_N^H &= \exp \left[ N^{1+\alpha} \left\{ H(T, 1) \mathbf{h}_N(T, N) - \int_0^T \partial_t H(t, 1) \mathbf{h}_N(t, N) dt \right. \right. \\
&\quad - \int_0^T \left( \int_{-1}^1 \mu_N(t, y) \partial_{yy} H(t, y) dy + \partial_y H(t, 1) \rho_+(t) - \partial_y H(t, -1) \rho_-(t) \right. \\
&\quad \left. \left. + \frac{1}{2N} \sum_{x=-N-1}^N \left( \partial_y H(t, \frac{x}{N}) \right)^2 (\phi_+(\eta_t, x) + \phi_-(\eta_t, x)) \right) dt + O(N^{-(\alpha \wedge 1)}) \right\} \right].
\end{aligned} \tag{7.2}$$

where

$$\lim_{N \rightarrow \infty} \mathbb{E}^{\mathbb{P}_N} (O(N^{-(\alpha \wedge 1)})) = 0.$$

After applying the superexponential estimates of section 5, we have

$$\begin{aligned}
M_N^H &= \exp \left[ N^{1+\alpha} \left\{ H(T, 1) \mathbf{h}_N(T, N) - \int_0^T \partial_t H(t, 1) \mathbf{h}_N(t, N) dt \right. \right. \\
&\quad - \int_0^T \left( \int_{-1}^1 \pi_N(t, y) \partial_{yy} H(t, y) dy + \partial_y H(t, 1) \rho_+(t) - \partial_y H(t, -1) \rho_-(t) \right. \\
&\quad \left. \left. + \int_{-1}^1 (\partial_y H(t, y))^2 \phi(\bar{\eta}_{N, \varepsilon}(t, y)) dy \right) dt + O_{\varepsilon, N} + O(N^{-(\alpha \wedge 1)}) \right\} \right].
\end{aligned}$$

Then for every set  $A \subset \Xi$  we have

$$1 = \mathbb{E}^{\mathbb{P}_N} (M_N^H) \geq e^{N^{1+\alpha} \sup_H \inf_{(J, \rho) \in A} I(H, J, \rho)} \mathbb{P}((\mathbf{h}_N, \pi_N) \in A)$$

where

$$\begin{aligned}
I(H, J, \rho) &= H(T, 1) J(T) - \int_0^T \partial_t H(t, 1) J(t) dt \\
&\quad - \int_0^T \left( \int_{-1}^1 \rho(t, y) \partial_{yy} H(t, y) dy + \partial_y H(t, 1) \rho_+(t) - \partial_y H(t, -1) \rho_-(t) \right. \\
&\quad \left. + \int_{-1}^1 (\partial_y H(t, y))^2 \phi(\rho(t, y)) dy \right) dt.
\end{aligned}$$

Using the lower semicontinuity of  $I(J, \rho)$  and a standard argument (see [12], lemma 11.3 or [8] lemma A2.3.3) we have for a compact set  $C \subset \Xi$ :

$$\limsup_{N \rightarrow \infty} \frac{1}{N^{1+\alpha}} \log \mathbb{P}((\mathbf{h}_N, \pi_N) \in C) \leq - \inf_{(J, \rho) \in C} I(J, \rho).$$

The extension to closed set follows from the exponential compactness proved in (7.1), see [8] pag.271.

## 8. THE LOWER BOUND

The proof of the lower bound follows a standard argument, consequently we will only sketch it here, since all the ingredients are already proven. It is enough to prove that given  $(J, \rho) \in \Xi$  such that  $I(J, \rho) < \infty$ , then for any open neighbor  $\mathcal{O}$  of it we have

$$\liminf_{N \rightarrow \infty} \frac{1}{N^{1+\alpha}} \log P((\mathbf{h}_N, \mu_N) \in \mathcal{O}) \geq -I(J, \rho). \quad (8.1)$$

By proposition 3.2, we can assume  $J$  and  $\rho$  such that  $J'(t)$  exists and is bounded,  $\rho$  bounded away from 0 and 1, and  $\partial_y \rho(t, y)$  exists and is bounded. Then we consider the weakly asymmetric excusion dynamics with drift given by

$$\partial_y \bar{H}(t, y) = \frac{1}{2} \frac{J'(t) + \partial_y \rho(t, y)}{\phi(\rho(t, y))}, \quad (8.2)$$

more precisely the jump rate from  $x$  to  $x + 1$  at time  $t$  is taken to be  $N^{2+\alpha} e^{z(t, x/N)}$ , where

$$z(t, x/N) = \bar{H}(t, (x+1)/N) - \bar{H}(t, x/N) \quad (8.3)$$

and from  $x + 1$  to  $x$  is given by  $N^{2+\alpha} e^{-z(t, x/N)}$ . We call  $Q_N$  the law of this weakly asymmetric process.

The Radon-Nykodyn derivative  $\frac{dQ_N}{dP_N}$  is given by (6.4).

The quasi static limit for this process is the following:

**Proposition 8.1.** *Let  $\tilde{Q}_N$  the law on  $\Xi$  of  $(\mathbf{h}_N, \pi_N)$  under  $Q_N$ , then*

$$\tilde{Q}_N \longrightarrow \delta_{(J, \rho)} \quad (8.4)$$

*Proof of proposition 8.1.* By (9.1) we can extend the superexponential estimates contained in section 5 to  $Q_N$ . [In fact we have that](#)

$$\mathbb{E}^{P_N} \left( \left( \frac{dQ_N}{dP_N} \right)^2 \right)^{1/2} \leq e^{cN^{1+\alpha}}$$

and by Schwarz inequality

$$Q_N(A_{N, \varepsilon}) \leq P_N(A_{N, \varepsilon}) e^{cN^{1+\alpha}}$$

where  $A_{N, \varepsilon} = \{\int_0^T V_{N, \varepsilon}(t, \eta_t) dt \geq N\delta\}$ , and (5.6) extends immediately to  $Q_N$ . At this point the proof of the quasi-static hydrodynamic limit follows similar to the one in [6].  $\square$

We then write

$$P((\mathbf{h}_N, \pi_N) \in \mathcal{O}) = \mathbb{E}^{Q_N} \left( \frac{dQ_N}{dP_N} \mathbf{1}_{(\mathbf{h}_N, \pi_N) \in \mathcal{O}} \right)$$

Since  $\mathcal{O}$  contains  $(J, \rho)$ , by Proposition 8.1, under  $Q_N$  the probability of the event  $(\mathbf{h}_N, \pi_N) \in \mathcal{O}$  is close to one. By Jensen inequality

$$\frac{1}{N^{1+\alpha}} \log P((\mathbf{h}_N, \pi_N) \in \mathcal{O}) \geq \mathbb{E}^{Q_N} \left( \frac{1}{N^{1+\alpha}} \log \frac{dQ_N}{dP_N} \right) = -\frac{1}{N^{1+\alpha}} H(Q_N|P_N)$$

where  $H(Q_N|P_N)$  is the relative entropy of  $Q_N$  with respect to  $P_N$ .

The lower bound is then a consequence of the following Proposition.

**Proposition 8.2.** *Let  $H(Q_N|P_N) = \mathbb{E}^{Q_N} \left( \log \frac{dQ_N}{dP_N} \right)$  the relative entropy of  $Q_N$  with respect to  $P_N$ . Then*

$$\lim_{N \rightarrow \infty} H(Q_N|P_N) = I(J, \rho), \quad (8.5)$$

The proof of proposition 8.2 is a direct consequence of (6.4) and of Proposition 8.1.

## 9. APPENDIX A: THE EXPONENTIAL TIGHTNESS

We prove here proposition 7.1. The arguments used here are just slight variations of the standard ones (e.g. section 10.4 in [8]). Since  $\mathcal{M}$  is compact, we have only to control that the distribution of  $\mathbf{h}_N$  is exponentially tight. This is consequence of the following 2 propositions.

**Proposition 9.1.**

$$\lim_{L \rightarrow \infty} \lim_{N \rightarrow \infty} \frac{1}{N^{1+\alpha}} \log \mathbb{P}_\eta \left( \sup_{0 \leq t \leq T} |\mathbf{h}_N(t)| \geq L \right) = -\infty. \quad (9.1)$$

**Proposition 9.2.** *For any  $\varepsilon > 0$ :*

$$\lim_{\delta \rightarrow 0} \lim_{N \rightarrow \infty} \frac{1}{N^{1+\alpha}} \log \mathbb{P}_\eta \left( \sup_{|t-s| \leq \delta} |\mathbf{h}_N(t) - \mathbf{h}_N(s)| \geq \varepsilon \right) = -\infty. \quad (9.2)$$

*Proof of Proposition 9.1 .* Since the difference between  $\mathbf{h}_N(t)$  and  $\bar{\mathbf{h}}_N(t) := \frac{1}{2N} \sum_x \mathbf{h}_N(t, x)$  is uniformly small, we have just to prove it for  $\bar{\mathbf{h}}_N(t)$ . For  $\beta \in \mathbb{R}$ , consider the exponential martingale (6.4) with  $z_+(t, x) = \frac{\beta}{N}$ ,  $x = -N, \dots, N-1$  and  $z_+(t, -N-1) = 0$ . This is given by

$$M_t = \exp \left\{ N^{1+\alpha} (\beta \bar{\mathbf{h}}_N(t) - A_N(\beta, t)) \right\}$$

$$A_N(\beta, t) = N \int_0^t \sum_{x=-N}^{N-1} ((e^{\beta/N} - 1) \phi_+(\eta_s, x) + (e^{-\beta/N} - 1) \phi_-(\eta_s, x)) ds$$



Notice that  $0 \leq A_N(\beta, t) \leq CT(|\beta| + \beta^2)$  for some constant  $C$ . Then for any  $\beta > 0$  we have, by Doob's inequality,

$$\begin{aligned}
\mathbb{P}_\eta \left( \sup_{0 \leq t \leq T} |\mathbf{h}_N(t)| \geq L \right) &\leq \mathbb{P}_\eta \left( \sup_{0 \leq t \leq T} |\log M_t| \geq N^{1+\alpha} (\beta L - CT(\beta + \beta^2)) \right) \\
&\leq \mathbb{P}_\eta \left( \sup_{0 \leq t \leq T} \log M_t \geq N^{1+\alpha} (\beta L - CT(\beta + \beta^2)) \right) \\
&= \mathbb{P}_\eta \left( \sup_{0 \leq t \leq T} M_t \geq e^{N^{1+\alpha}(\beta L - CT(\beta + \beta^2))} \right) \\
&\leq e^{-N^{1+\alpha}(\beta L - CT(\beta + \beta^2))},
\end{aligned} \tag{9.3}$$

that concludes the proof.  $\square$

*Proof of Proposition 9.2 .* Since

$$\left\{ \sup_{|t-s| \leq \delta} |\mathbf{h}_N(t) - \mathbf{h}_N(s)| \geq \varepsilon \right\} \subset \bigcup_{k=0}^{\lceil T\delta^{-1} \rceil} \left\{ \sup_{k\delta \leq t \leq (k+1)\delta} |\mathbf{h}_N(t) - \mathbf{h}_N(k\delta)| \geq \varepsilon/4 \right\}$$

Since

$$\log \mathbb{P}_\eta \left( \sup_{|t-s| \leq \delta} |\mathbf{h}_N(t) - \mathbf{h}_N(s)| \geq \varepsilon \right) \leq \max_k \log \mathbb{P}_\eta \left( \sup_{k\delta \leq t \leq (k+1)\delta} |\mathbf{h}_N(t) - \mathbf{h}_N(k\delta)| \geq \varepsilon/4 \right) + \log(\lceil T\delta^{-1} \rceil)$$

By the same estimate made in (9.3) we have

$$\log \mathbb{P}_\eta \left( \sup_{k\delta \leq t \leq (k+1)\delta} |\mathbf{h}_N(t) - \mathbf{h}_N(k\delta)| \geq \varepsilon/4 \right) \leq -N^{1+\alpha} (\beta\varepsilon/4 - C\delta(\beta + \beta^2))$$

and with a proper choice of  $\beta$  we get the following bound with a constant  $C'$  independent of  $k$ :

$$\frac{1}{N^{1+\alpha}} \log \mathbb{P}_\eta \left( \sup_{k\delta \leq t \leq (k+1)\delta} |\mathbf{h}_N(t) - \mathbf{h}_N(k\delta)| \geq \varepsilon/4 \right) \leq -\frac{C'\varepsilon^2}{\delta}.$$

$\square$

## REFERENCES

- [1] Bertini L., Gabrielli D., Jona-Lasinio G., and Landim C. *Clausius inequality and optimality of quasistatic transformations for nonequilibrium stationary states*. Physical review letters, **110**(2):020601 (2013).
- [2] Bertini L., Landim C., Mourragui M., *Dynamical large deviations for the boundary driven weakly asymmetric exclusion process*, Annals of Probability, **37**, 6, 2357-2403, (2009).

- [3] Bodineau T., Derrida B., *Current Large Deviations for Asymmetric Exclusion Processes with Open Boundaries*, Journal of Statistical Physics, **123**, n.2, 277–300 (2006).
- [4] Chavez E., Landim C., *A Correction to the Hydrodynamic Limit of Boundary Driven Weakly Asymmetric Exclusion Processes in a Quasi-Static Time Scale*, Journal of Statistical Physics, **163**, Issue 5, pp 10791107, (2016).
- [5] De Masi A., Presutti E., Tsagkarogiannis D., and Vares M.E.. *Truncated correlations in the stirring process with births and deaths*, Electronic Journal of Probability **17** 1-35 (2012)
- [6] De Masi A., Olla S. *Quasi Static Hydrodynamic Limits*, Journal of Statistical Physics, **161**, n.5, 1037–1058 (2015).
- [7] Galves A., Kipnis C., Marchioro C. and Presutti E. *Non equilibrium measures which exhibit a temperature gradient: study of a model*. Commun. Math. Phys., **819**, 127-1474 (1981).
- [8] Kipnis C., Landim C.. *Scaling Limits of Interacting Particle Systems*, Grundlehren der mathematischen Wissenschaften 320, Springer-Verlag, Berlin, New York, (1999).
- [9] Kipnis C., Olla S., and Varadhan S.R.. *Hydrodynamics and large deviation for simple exclusion processes*, Commun. Pure Appl. Math. **42** 115 –137(1989). *Stationary and non-equilibrium fluctuations in boundary driven exclusion process* Markov Proces. Related Fields, **14**, 165–184 (2008).
- [10] Letizia V., and Olla S.. *Non-equilibrium isothermal transformations in a temperature gradient from a macroscopic dynamics*, Annals of Probability, **45**, N. 6A (2017), 3987–4018. doi:10.1214/16-AOP1156.
- [11] Olla S.. *Microscopic derivation of an isothermal thermodynamic transformation*, In From Particle Systems to Partial Differential Equations, pages 225–238. Springer, 2014.
- [12] Varadhan, S.R.S., *Large Deviations and Applications*, CBMSNSF Regional Conference Series in Applied Mathematics 46, Philadelphia, Society for Industrial and Applied Mathematics, 1984.

ANNA DE MASI, UNIVERSITÀ DELL’AQUILA,, 67100 L’AQUILA, ITALY  
*E-mail address:* demasi@univaq.it

STEFANO OLLA, CEREMADE, UMR CNRS, UNIVERSITÉ PARIS-DAUPHINE,  
PSL RESEARCH UNIVERSITY, 75016 PARIS, FRANCE  
*E-mail address:* olla@ceremade.dauphine.fr