



**HAL**  
open science

## Defining and measuring asymmetry

Michel Petitjean

► **To cite this version:**

Michel Petitjean. Defining and measuring asymmetry. First European Asymmetry Symposium (FEAS), Mar 2018, Nice, France. , 2018. hal-01778387

**HAL Id: hal-01778387**

**<https://hal.science/hal-01778387v1>**

Submitted on 25 Apr 2018

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

## DEFINING AND MEASURING ASYMMETRY

Michel Petitjean

(member of the executive board of the International Symmetry Association)

MTi, INSERM UMR-S 973, University Paris 7  
 petitjean.chiral@gmail.com <http://petitjeanmichel.free.fr/itoweb.petitjean.html>

### AN UNIFYING DEFINITION OF SYMMETRY [1]

*It is crucial to understand that mathematical constructs are models of real physical situations, and that a mathematical model of symmetry is a simplified image in our mind of some physical situation in which we would like to see symmetry.*

An object  $Y$  is a **function** having its input argument in a metric space  $E$ .  $Y$  is transformed via distance-preserving bijections of  $E$  onto  $E$ .

$F$  is the set of all these distance-preserving bijections.  $F$  is a group and  $I_F$  is its neutral element.

An object  $Y$  is symmetric if there is a bijection  $U$  of  $F$ , with  $U \neq I_F$ , such that for all element  $x$  of  $E$ ,  $Y(Ux) = Y(x)$ .

Five groups appear during the building process of this symmetry definition, including the one above.

These groups are not imposed a priori during this process: they appear naturally as consequences of several trivial properties that a symmetric object is expected to have.

*The definition applies to: geometric figures in Euclidean spaces (with or without constraints of colors), functions, probability distributions, matrices, strings, graphs, etc.*

### CHIRALITY : LACK OF INDIRECT SYMMETRY [2]

Owing to the unifying definition of symmetry, we define  $F^+$  as a subset of  $F$  such that the elements of  $F^+$  are writable as products of squared elements of  $F$ . It is a subgroup of  $F$  and its elements are called direct isometries. The elements of  $F \setminus F^+$ , when existing, are called indirect isometries.

An object having symmetry due to an indirect isometry has indirect symmetry and is called achiral. An object having no indirect symmetry is called chiral.

*We defined chirality in metric spaces: no need of an Euclidean space, no need of an orientable space. However, the definition generalizes the usual one of Lord Kelvin (1904)*

### MEASURING ASYMMETRY IN EUCLIDEAN SPACES

Many measures of asymmetry or chirality are unsafe because they are not based on a distance between objects [3]. A suitable choice is the colored Wasserstein distance [4], a generalization of the  $L^2$ -Wasserstein distance taking colors in account. The Direct Symmetry Index  $DSI$  and the chiral index  $\chi$  measure respectively the degree of direct asymmetry and the degree of chirality [5]. Then we are able to characterize several extreme asymmetric figures and distributions [6, 7, 8, 9]. We present below the examples of the most asymmetric triangles.

$DSI$  takes values in the interval  $[0..1]$

Direct symmetry  $\Leftrightarrow DSI = 0$

The most direct-asymmetric triangles are below [5]

*The unequivalence of all vertices precludes the existence of any direct symmetry: at least two vertices should be equivalent*



This degenerate triangle with only two equivalent vertices is maximally direct-asymmetric

in the  $d$ -dimensional Euclidean space,  $d > 1$ :  $DSI = 1$

Abscissas:  $(-1 - \sqrt{3})/2$ ,  $(-1 + \sqrt{3})/2$ , 1

The most direct-asymmetric triangle with 3 equivalent vertices


$DSI = 1 - \sqrt{2}/2$  Angles:  $\pi/4$ ,  $\pi/8$ ,  $5\pi/8$



$\chi$  takes values in the interval  $[0..1]$


Achirality  $\Leftrightarrow \chi = 0$

The most chiral triangles are below [6]



The regular  $d$ -simplex with all  $d + 1$  non equivalent vertices is maximally chiral in the  $d$ -dimensional Euclidean space

$$\chi = 1$$



The most chiral triangle with only two equivalent vertices

$$\chi = 1 - \sqrt{2}/2$$

Distances ratios:  $\sqrt{1 - \sqrt{6}/4} : 1 : \sqrt{1 + \sqrt{6}/4}$



The most chiral triangle with three equivalent vertices

$$\chi = 1 - 2\sqrt{5}/5$$

Distances ratios:  $1 : \sqrt{4 + \sqrt{15}} : \sqrt{(5 + \sqrt{15})/2}$

### REFERENCES

- [1] Petitjean, M. A Definition of symmetry. *Symmetry Cult. Sci.* **2007**, 18[2-3], 99–119. <https://hal.archives-ouvertes.fr/hal-01552499>
- [2] Petitjean, M. Chirality in metric spaces. In memoriam Michel Deza. *Optim. Letters* **2018**, under press. DOI 10.1007/s11590-017-1189-7
- [3] Petitjean, M. Chirality and symmetry measures: a transdisciplinary review. *Entropy* **2003**, 5[3], 271–312.
- [4] Petitjean, M. Chiral mixtures. *J. Math. Phys.* **2002**, 43[8], 4147–4157.
- [5] Petitjean, M. On the root mean square quantitative chirality and quantitative symmetry measures. *J. Math. Phys.* **1999**, 40[9], 4587–4595.
- [6] Petitjean, M. About second kind continuous chirality measures. 1. Planar sets. *J. Math. Chem.* **1997**, 22[2-4], 185–201.
- [7] Coppersmith, D.; Petitjean, M. About the optimal density associated to the chiral index of a sample from a bivariate distribution. *Compt. Rend. Acad. Sci. Paris (Sér. I)* **2005**, 340[8], 599–604.
- [8] Schwartz, A.; Petitjean, M. [6.6]Chiralane: a remarkably symmetric chiral molecule. *Symmetry Cult. Sci.* **2008**, 19[4], 307–316.
- [9] Petitjean, M. The most chiral disphenoid. *MATCH Commun. Math. Comput. Chem.* **2015**, 73[2], 375–384.