



HAL
open science

Fungi diversity in PM_{2.5} and PM₁ at the summit of Mt. Tai: abundance, size distribution, and seasonal variation

Caihong Xu, Min Wei, Jianmin Chen, Chao Zhu, Jiarong Li, Ganglin Lv, Xianmang Xu, Lulu Zheng, Guodong Sui, Weijun Li, et al.

► **To cite this version:**

Caihong Xu, Min Wei, Jianmin Chen, Chao Zhu, Jiarong Li, et al.. Fungi diversity in PM_{2.5} and PM₁ at the summit of Mt. Tai: abundance, size distribution, and seasonal variation. *Atmospheric Chemistry and Physics*, 2017, 17 (18), pp.11247-11260. 10.5194/acp-17-11247-2017 . hal-01785118

HAL Id: hal-01785118

<https://hal.science/hal-01785118v1>

Submitted on 4 May 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Fungi diversity in PM_{2.5} and PM₁ at the summit of Mt. Tai: abundance, size distribution, and seasonal variation

Caihong Xu¹, Min Wei^{1,a}, Jianmin Chen^{1,2,3}, Chao Zhu¹, Jiarong Li¹, Ganglin Lv¹, Xianmang Xu¹, Lulu Zheng², Guodong Sui², Weijun Li¹, Bing Chen¹, Wenxing Wang¹, Qingzhu Zhang¹, Aijun Ding³, and Abdelwahid Mellouki^{1,4}

¹Environment Research Institute, School of Environmental Science and Engineering, Shandong University, Jinan 250100, China

²Shanghai Key Laboratory of Atmospheric Particle Pollution and Prevention (LAP3), Fudan Tyndall Centre, Department of Environmental Science & Engineering, Fudan University, Shanghai 200433, China

³Institute for Climate and Global Change Research, School of Atmospheric Sciences, Nanjing University, Nanjing 210023, Jiangsu, China

⁴Institut de Combustion, Aérothermique, Réactivité et Environnement, CNRS, 45071 Orléans CEDEX 02, France

^anow at: College of Geography and Environment, Shandong Normal University, Jinan 250100, China

Correspondence to: Jianmin Chen (jmchen@fudan.edu.cn, jmchen@sdu.edu.cn)

Received: 5 March 2017 – Discussion started: 29 March 2017

Revised: 28 July 2017 – Accepted: 16 August 2017 – Published: 22 September 2017

Abstract. Fungi are ubiquitous throughout the near-surface atmosphere, where they represent an important component of primary biological aerosol particles. This study combined internal transcribed spacer region sequencing and quantitative real-time polymerase chain reaction (qPCR) to investigate the ambient fungi in fine (PM_{2.5}, 50% cutoff aerodynamic diameter $D_{a50} = 2.5 \mu\text{m}$, geometric standard deviation of collection efficiency $\sigma_g = 1.2$) and submicron (PM₁, $D_{a50} = 1 \mu\text{m}$, $\sigma_g = 1.2$) particles at the summit of Mt. Tai located in the North China Plain, China. Fungal abundance values were 9.4×10^4 and 1.3×10^5 copies m^{-3} in PM_{2.5} and PM₁, respectively. Most of the fungal sequences were from Ascomycota and Basidiomycota, which are known to actively discharge spores into the atmosphere. The fungal community showed a significant seasonal shift across different size fractions according to Metastats analysis and the Kruskal–Wallis rank sum test. The abundance of *Glomerella* and *Zasmidium* increased in larger particles in autumn, whereas *Penicillium*, *Bullera*, and *Phaeosphaeria* increased in smaller particles in winter. Environmental factors, namely Ca^{2+} , humidity, and temperature, were found to be crucial for the seasonal variation in the fungal community. This study might serve as an important reference for fungal contribution to primary biological aerosol particles.

1 Introduction

Inhaled particulate matter (PM), categorized as PM_{2.5} and PM₁ (aerodynamic equivalent diameters of ≤ 2.5 and $\leq 1 \mu\text{m}$, respectively), has proven to be associated with the increasing morbidity and mortality from cardiovascular and respiratory diseases (Brauer et al., 2013; Wang et al., 2014). Primary biological aerosol particles (PBAPs; about 10^4 – 10^8 cells cm^{-2}) constitute an important component of PM. They can actively metabolize in the atmosphere with their mass concentrations ranging from 5.49 to 102 ng m^{-3} (Zhong et al., 2016). Furthermore, they play an important role in agriculture, the biosphere, cloud formation, global climate, and atmospheric dynamics (Brodie et al., 2007; Despres et al., 2012; Christner et al., 2008; Zhou et al., 2014; Jaenicke et al., 2005). Fungi, the primary group of PBAPs, include 1.5 million unique species, distributed across rural and urban environments (Hawksworth et al., 2001). They actively eject their spores with aqueous jets or droplets into the atmosphere. The global emissions of fungal spores are estimated as the largest source of bioaerosols (Elbert et al., 2007). Pioneering studies have reported global fungal emissions to reach 28 Tg per year and contribute to about 4–13% of the mass concentration of PM_{2.5} (Heald et al., 2009; Womiloju et al., 2003). More recently, some specific fungal species have been verified to be linked with the occurrence of public health problems (Mor-

ris et al., 2002; Yadav et al., 2004; Bowers et al., 2012, 2013; Cao et al., 2014; Ryan et al., 2009). Despite their importance, the abundance, diversity, and community structure of fungi associated with PM have received limited attention in terms of research.

Earlier studies on airborne fungal communities, primarily based on culturing methods, found the dominant phyla to be Ascomycota (AMC) and Basidiomycota (BMC). Some of the species are considered major pathogens and allergens of plants, animals, and humans, e.g., *Hemileia vastatrix*, *Aspergillus*, *Cryptococcus*, and *Pneumocystis* spp. (Despres et al., 2012; Smets et al., 2016). While most of the fungal species remain unknown because cultivable species (typically less than 100) occupy only a tiny minority of all existing species, advances in nucleic acid sequencing allow the accurate determination of both cultured and uncultured microbial communities in environmental samples. For bacterial community composition, Xu et al. (2017a) investigated the abundance and community of bacteria in submicron particles during severe haze episodes in Jinan, China. Later, they discussed the diurnal variation of diverse bacterial communities in cloud water at Mt. Tai, China (Xu et al., 2017b). For diverse fungi in Mainz, Germany, Frohlich-Nowoisky et al. (2009) described the fungal community in coarse ($> 3 \mu\text{m}$) and fine ($\leq 3 \mu\text{m}$) PM using internal transcribed spacer (ITS) region sequencing. Yamamoto et al. (2012) reported the crucial influence of aerodynamic diameter and season on the fungal taxonomic composition in the northeastern United States by 454 pyrosequencing. The fungal allergens clustered in the largest size ranges ($> 9 \mu\text{m}$) in the fall season, whereas the pathogens were most abundant in the spring season and were typically observed in particles with aerodynamic diameters of $< 4.7 \mu\text{m}$. Subsequently, DeLeon-Rodriguez et al. (2013) discussed the effect of tropical storm or hurricane periods on the shift of airborne fungal species over the upper troposphere. Gou et al. (2016) described the fungal abundance and taxonomic composition of fungi in PM₁ and PM₁₀ in winter in China by 18S rRNA gene sequencing. However, that study focused on the ambient fungi in total suspended particles (TSP), PM₁₀, and PM_{2.5}, and was primarily conducted over the ground's surface; therefore, fungal populations in PM₁ at high-elevation sites were not well accounted for. Diverse microbes at high altitudes (such as in cloud water and precipitation) can act as nucleating agents for cloud and ice condensation, influence precipitation patterns (Xu et al., 2017b; Pratt et al., 2009; Creamean et al., 2013; Bower et al., 2013), and drive the biogeochemical cycling of elements in ecosystem processes. Hence, it is essential to advance the knowledge of microbes in PM, especially across the East Asian regions which are frequently ravished by dust, haze or other weather phenomenon. During 2013, 2014, and 2015, serious air pollution events associated with the inadequate use of clean energy in the transport, domestic, and industrial sectors affected northern China, which includes several areas with severe air pollution, namely Beijing, Tian-

jin, Shijiazhuang, Jinan, and Qingdao. Most researchers focus their attention on the case study of bacterial abundance and diversity (Gao et al., 2014, 2017a; Xu, et al., 2017a; Wei et al., 2017). The various physical, chemical, and biological factors caused by the severe haze or dust episodes may cause shifts in the bacterial community structure. Moreover, the airborne microbial abundance and diversity are also effected by seasonal and meteorological factors; however, the investigations into the seasonal variation of fungal characteristics in aerosol particles have been very limited.

Mt. Tai ($36^{\circ}15' \text{N}$, $117^{\circ}06' \text{E}$; 1534 m a.s.l.), the highest site in the North China Plain, is a tilted fault block mountain, its height increasing from the north to the south, facing the Japanese islands, Korean Peninsula, East China Sea, and the Yellow Sea. The vegetation cover is 80%, with nearly 1000 kinds of plants growing in the area. The number of tourists, from both China and abroad, visiting this mountain increased from 5.5 million in 2014 to 5.9 million in 2015. Past investigations in this region mainly concentrated on the physicochemical characteristics of aerosol particles and cloud water and their influence on air quality and human health. Thus far, there have been no studies addressing the diverse fungal community in aerosol particles at Mt. Tai, necessitating the development of a reliable knowledge base on the atmospheric aerosols in such scenic destinations.

The objectives of the present study were: (i) to fill the knowledge gaps regarding the ambient fungi of PM_{2.5} and PM₁ at a high-elevation site of East Asia, (ii) to elucidate the size-based differences between the data of ambient fungal concentration and viable fungal community structure across different seasons, and (iii) to estimate whether environmental factors play a role in the variation of fungal characteristics at Mt. Tai.

2 Materials and methods

2.1 Sample collection

At Mt Tai, spring occurs from March to May; summer, June to August; fall, September to November; and winter, December to February, according to the environmental temperature. Two middle-volume (100 L min^{-1}) samplers (TH-150A; Wuhan Tianhong Instruments Co. Ltd., Wuhan, China) were deployed with particles larger than 2.5 and $1 \mu\text{m}$ trapped by the impactors and particles smaller than 2.5 and $1 \mu\text{m}$ collected on the quartz filters. The 50% cutoff aerodynamic diameters are 2.5 and $1 \mu\text{m}$. The smaller the aerosol particles, the higher the collection efficiency. Sixty quartz membrane filters (Pall, NY, USA; 88 mm) were obtained for 23 h (09:00 to 08:00 the next day) over 8–13 days during each season from 2014 to 2015 at the summit of Mt. Tai (Table 1). The blank filters were obtained by placing sterilized quartz microfibre filters inside the sampler without any operation. Before sampling, all the filters were baked in a muffle fur-

Table 1. Sample descriptions and the associated meteorological characteristics of the atmosphere, including the temperature (*T*), relative humidity (RH), PM_{2.5} mass concentration (MC), PM₁ mass concentration, and fungal cell concentrations on the basis of qPCR analysis of rRNA copy numbers in PM_{2.5} and PM₁.

Season	Date of collection	<i>T</i>	RH	PM _{2.5}				PM ₁			
				MC	Fungal SSU rRNA gene copy number	Fungal spore OC MC	Fungal spore MC	MC	Fungal SSU rRNA gene copy number	Fungal spore OC MC	Fungal spore MC
Unit		°C	%	µg m ⁻³	10 ⁴ copy m ⁻³	ng C m ⁻³	µg m ⁻³	µg m ⁻³	10 ⁴ m ⁻³	ng C m ⁻³	µg m ⁻³
Summer	06/25/15	12.6	98	5.5	11.00	7.15	0.02	BDL	18.00	11.69	0.03
	06/26/15	14	97	2.8	3.58	2.33	0.01	BDL	6.51	4.23	0.01
	06/27/15	15.1	94.6	52.7	1.01	0.66	0.00	18.1	0.40	0.26	0.00
	06/28/15	16.9	84.4	91.1	6.85	4.45	0.01	40.0	3.10	2.02	0.01
	06/29/15	17.3	62.6	16.8	4.71	3.06	0.01	13.3	2.79	1.81	0.00
	07/03/15	17.7	31.0	15.3	12.20	7.92	0.02	12.7	11.40	7.42	0.02
	07/07/15	16.9	84.4	94.0	47.70	31.00	0.08	39.9	4.20	2.73	0.01
	07/08/15	17.3	62.6	110.9	22.60	14.71	0.04	42.0	5.31	3.45	0.01
08/07/15	17.4	97.6	13.5	5.64	3.67	0.01	11.4	6.94	4.51	0.01	
Autumn	10/22/14	6.7	60.7	40.1	9.34	6.07	0.02	28.1	0.37	0.24	0.00
	10/25/14	10.3	80.7	48.1	7.28	4.73	0.01	34.6	45.70	29.73	0.08
	10/26/14	11.4	73.6	50.8	6.78	4.41	0.01	31.9	8.26	5.37	0.01
	11/03/14	0.3	21.6	4.9	1.57	1.02	0.00	BDL	22.20	14.44	0.04
	11/04/14	2.6	33.7	31.6	7.95	5.17	0.01	24.5	6.18	4.01	0.01
	11/05/14	4.1	30.9	33.1	3.70	2.41	0.01	25.6	103.00	66.75	0.17
	11/06/14	5.1	19.3	22.7	12.70	8.24	0.02	18.0	8.86	5.76	0.01
	11/07/14	2.6	34.0	19.8	7.89	5.13	0.01	16.0	6.39	4.15	0.01
	11/08/14	2.4	45.7	22.4	14.70	9.54	0.02	17.8	2.92	1.90	0.00
	11/09/14	1.1	73.1	77.1	5.56	3.61	0.01	33.5	3.61	2.34	0.01
	11/10/14	3.0	49.0	49.2	9.38	6.10	0.02	37.2	16.70	10.87	0.03
	11/11/14	2.7	65.4	32.7	27.50	17.85	0.05	25.3	26.30	17.07	0.04
11/12/14	1.0	50.1	51.7	7.50	4.87	0.01	25.7	18.30	11.87	0.03	
Winter	12/03/14	-8.9	24.4	13.7	5.03	3.27	0.01	9.7	6.84	4.45	0.01
	12/04/14	-11	39.1	35.0	8.68	5.64	0.01	30.6	2.78	1.81	0.00
	12/05/14	-10.6	23.4	14.5	1.09	0.71	0.00	13.3	16.20	10.52	0.03
	12/06/14	-5.7	11.0	9.1	6.32	4.11	0.01	8.3	4.15	2.70	0.01
	12/07/14	-5.4	45.7	38.8	7.90	5.14	0.01	30.9	9.36	6.08	0.02
	12/08/14	-7.9	35.7	36.5	1.33	0.86	0.00	29.0	7.17	4.66	0.01
	12/09/14	-5.3	16.1	16.5	10.10	6.55	0.02	13.5	5.73	3.72	0.01
	12/10/14	-5.6	58.3	9.3	3.24	2.10	0.01	8.1	7.10	4.62	0.01

C – carbon, MC – mass concentration, *T* – temperature, RH – relative humidity, BDL – below the detection line.

nance at 500 °C for 5 h, placed into sterilized aluminum foil, and then deposited into a sealed bag. To avoid contamination, the sampling filter holder and materials used for changing filters were treated with 75 % ethanol every day. After sampling, the samples were stored at -80 °C until the next analysis. PM_{2.5} and PM₁ mass concentrations were monitored by a synchronized hybrid ambient real-time particulate monitor (Model 5030; Thermo Fisher Scientific, Wilmington, DE, USA). Half of the PM_{2.5} and PM₁ filters were used to analyze water-soluble inorganic ions (NO₃⁻, SO₄²⁻, NH₄⁺, K⁺, Ca²⁺, Na⁺, and Mg²⁺) by an ambient ion monitor (URG-9000; URG Corporation, Chapel Hill, NC, USA). The remaining filters were analyzed in the same batch of laboratory experiments, including DNA extraction, PCR amplification, quantitative real-time PCR (qPCR), and Illumina sequencing, except for sample A29 on 9 December 2014 (ac-

identally omitted in the first batch of Illumina sequencing). Considering that a part of the sequences in the two batches of experiments differed, we removed this sample before quality control. Meteorological data, including relative humidity, wind speed, wind direction, and temperature, were obtained from <http://www.underground.com> at a resolution of 3 h during the sampling period. The visibility was monitored online by a visibility sensor (Model PWD22; Vaisala, Finland) with a maximum limit of 20 km.

2.2 DNA extraction and PCR amplification

The sample pretreatment and DNA extraction experiments were performed following a protocol optimized by Jiang et al. (2015). This protocol can extract sufficient DNA from low-biomass environmental samples (e.g., aerosol particles) and boosted the DNA extraction efficiency by more than

twice as compared to the non-optimized extraction method. Besides, it has been applied for studying airborne microbial diversity in different environments (Cao et al., 2014; Deng et al., 2016; Tong et al., 2017; Gao et al., 2017b). Half of the filters (about 121.64 cm² in area) were cut into small pieces, inserted into 50 mL Falcon tubes that were filled with sterilized 1 × PBS buffer, and centrifuged at 200 × *g* for 3 h at 4 °C. The resuspension was collected into a 0.2 μm Supor 200 PES membrane disc filter. We cut the PES membrane disc filter into small pieces, heated the pieces to 65 °C in PowerBead tubes for 15 min and then vortexed them for 15 min. DNA was extracted according to the standard PowerSoil DNA isolation protocol (Judd et al., 2016) and purified by AMPure XP bead purification. A parallel extraction procedure was performed with the blank filter to check for sample contamination. DNA concentrations were quantified by a NanoDrop 2000 spectrophotometer (Thermo Fisher Scientific). The fragments of ITS1 regions were amplified from genomic DNA by PCR using the forward primer ITS1F (5'-CTTGGTCATTTAGAGGAAGTAA-3') and the reverse primer ITS4 (5'-TCCTCCGCTTATTGATATGC-3'), which target the fungal ITS region of the rRNA gene (Manter et al., 2007). The experiment was conducted using the Gene Amp[®] PCR System 9700 (Applied Biosystems, CA, USA) in a total volume of 50 μL PCR mix containing PCR buffer (1 ×), 1.5 μM MgSO₄, 0.4 μM of each deoxynucleotide triphosphate, 0.3 μM each of the forward and reverse primers, 0.5 U Ex Taq (TaKaRa, Dalian, China), 100 ng template DNA, and double distilled H₂O. The thermal cycling profile was 94 °C for 1 min; 35 cycles of denaturation at 98 °C for 20 s, annealing at 68 °C for 30 s, and elongation at 72 °C for 45 s; and final extension at 72 °C for 5 min. Three replicates of PCR for each sample were combined together. The final products were separated by 1.5 % agarose gel electrophoresis and purified using the Qiaquick PCR purification kit (Qiagen, Valencia, CA, USA). Purified amplicons were quantified by a Qubit 2.0 fluorometer (Thermo Scientific) and pooled with equal molar amounts. Sequencing libraries were generated using the Truseq DNA PCR-Free Sample Prep Kit following manufacturer's instructions. Sequencing was performed on an Illumina MiSeq instrument (Illumina, San Diego, CA, USA) with the MiSeq reagent kit V3 (Illumina) according to the standard protocols.

2.3 Sequence analyses

After high-throughput sequencing, we removed the chimeric and low-quality sequences using the FASTX-ToolKit (http://hannonlab.cshl.edu/fastx_toolkit) and UCHIME algorithm (Edge et al., 2011) before statistical analysis. The remaining high-quality sequences were normalized to 7973 reads to compare the different samples effectively. They were then clustered into operational taxonomic units (OTUs) at a 97 % similarity cutoff using USEARCH software (Version 7.1, <http://drive5.com/uparse/>). We used the OTUs as the basis

for estimating the alpha diversity and beta diversity. Alpha diversity estimators, including Chao1, Simpson's index, and Shannon's index, were performed by the Quantitative Insights into Microbial Ecology software (Version 1.8.0, http://qiime.org/scripts/assign_taxonomy.html; Kuczynski et al., 2011). The taxonomy of ITS sequences was analyzed by RDP Classifier against the UNITE database (release 7.0, <http://unite.ut.ee/index.php>; Koljalg et al., 2013) using a confidence threshold of 70 %. RDP Classifier was used to determine the taxonomic composition at the phylum, class, order, family, genus, and species levels (Koiv et al., 2015; Miettinen et al., 2015). The raw reads were deposited into the NCBI Sequence Read Archive database under accession number SRR5146156.

2.4 qPCR for ITS regions

To determine the fungal biomass, we performed qPCR (Gao et al., 2017a; Yamaguchi et al., 2016; Lee et al., 2010) using a CFX96 real-time PCR detection system (Bio-Rad, Hercules, CA, USA) in 25 μL reaction mixtures containing 12.5 μL TransStart Green qPCR SuperMix, 1 μL ITS3-KYO2 (5'-GATGAAGAACGYAGYRAA-3'), 1 μL ITS4 (5'-TCCTCCGCTTATTGATATGC-3'), 5 μL sample DNA, and 5.5 μL double-distilled H₂O. The amplification followed a three-step PCR for fungal ITS regions: 40 cycles of denaturation at 95 °C for 30 s, primer annealing at 52 °C for 30 s, and extension at 72 °C for 30 s. A standard curve was created using tenfold dilution series of fungal ITS region plasmids. Assuming that the average fungal genome has about 30–200 rRNA copies, the fungal concentrations were calculated using the methods described by Lee et al. (2010) and van Doorn et al. (2007).

2.5 Fungal contribution to atmospheric organic carbon

The contributions of fungal spores to organic carbon (OC) were calculated using mannitol as a biotracer. We assumed 1.7 pg mannitol and 13 pg OC per spore. To assess the contribution of fungal spores to the OC and to the mass balance of atmospheric aerosol particles quantitatively, we used the weighted-average carbon (C) conversion factor of 13 pg C per spore and of 33 pg fresh weight per spore, which had been obtained earlier as the average carbon content of spores from airborne fungal species (Bauer et al., 2008; Zhu et al., 2016; Liang et al., 2017).

2.6 Statistical analyses

To determine the differences in the fungal community variations among different size fractions, meta-analyses based on the permutation *t*-test were conducted using Mothur software (version 1.35.1). The program Metastats can produce a tab-delimited table to display the mean relative abundance of the mean, variance, and standard error, together with the *p* values and *q* values. Values were considered significant




Figure 1. Relationships between fungal number concentrations of PM_{2.5} and PM₁ with wind speed and wind direction.

if $p \leq 0.05$ and $q \leq 0.05$. The Kruskal–Wallis rank sum test was used to evaluate the seasonal variation of the microbial community. Boxplots and q values have been provided for illustration. The relationship between the ambient microbial community and environmental factors, including PM concentrations and chemical compositions, was assessed with nonparametric Spearman's rank correlation coefficients by SPSS 16.0.

3 Results and discussion

3.1 Concentration of fungal spores in PM_{2.5} and PM₁

PM_{2.5} and PM₁ samples were collected during summer, autumn, and winter at the summit of Mt. Tai. Temporal variations of the mass concentration and corresponding fungal spore numbers of PM_{2.5} and PM₁ are summarized in Table 1. PM₁ mass concentration was stable over different seasons, while PM_{2.5} demonstrated a high seasonal variation, with higher average concentrations in summer ($44.7 \mu\text{g m}^{-3}$) than in autumn ($37.2 \mu\text{g m}^{-3}$) and winter ($21.7 \mu\text{g m}^{-3}$). The values were much lower than that in the summer of 2006 ($123.1 \mu\text{g m}^{-3}$; Deng et al., 2011) and comparable with that in the summer of 2007 ($59.3 \mu\text{g m}^{-3}$; Zhou et al., 2009). The average PM₁ / PM_{2.5} ratios were 0.45 in summer, 0.65 in autumn, and 0.84 in winter, implying that fine particles dominated in summer, while submicron particles dominated in autumn and winter.

qPCR revealed an average fungal gene copy number of 9.4×10^4 copies m^{-3} (ranging from 1.0×10^4 to 4.8×10^5 copies m^{-3}) and 1.3×10^5 copies m^{-3} (ranging from 3.7×10^3 to 1.0×10^6 copies m^{-3}) in PM_{2.5} and PM₁, respectively. There is no significant differences between PM_{2.5} and PM₁ based on the uncertainty estimate (95 % confidence intervals). Assuming an average rRNA gene copy

number of 200 per fungal genome (van Doorn et al., 2007; Lee et al., 2010), we obtained an average fungal concentration of 467 and 644 spores m^{-3} in PM_{2.5} and PM₁, respectively. The concentrations at Mt. Tai were lower than those at surface ground sites, including those in South Korea (ranging from 9.56×10^1 to 4.2×10^4 cells m^{-3} ; Lee et al., 2010), Austria (1.8×10^4 cells m^{-3} in urban sites and 2.3×10^4 cells m^{-3} in suburban sites; Bauer et al., 2008), Portugal (ranging from 891 to 964 spores m^{-3} ; Oliveira et al., 2009), and the United States (6450 spores m^{-3} ; Tsai et al., 2007). Our lower values might be ascribed to an underestimation of the fungal numbers. We used a higher gene copy number of 200 for each microbe studied, whereas DeLeon-Rodriguez et al. (2013) employed a lower number of rRNA copies of fungal genomes (30–100 copies per genome). The discrepancy between our results and those of Lee et al. (2010) might be because of the differences in sample type, sampling time, and altitude. Lee et al. (2010) focused on the fungal concentration in TSP by a high-volume TSP sampler ($0.225 \text{ m}^3 \text{ min}^{-1}$) 15 m above the ground in autumn and winter, whereas we obtained the PM_{2.5} and PM₁ by middle-volume samplers ($0.1 \text{ m}^3 \text{ min}^{-1}$) 1534 m above the ground in summer, autumn, and winter. It is difficult to explain the disparity between different studies without uniform guidelines for the sampling and quantitative assessment of bioaerosols.

Fungal abundance varied seasonally with different size particles in the near-surface atmosphere. Saari et al. (2015) found that coarse fluorescent bioaerosol particles ($1.5\text{--}5 \mu\text{m}$) increased in summer, whereas in winter, these particles primarily existed in smaller particles ($0.5\text{--}1.5 \mu\text{m}$). The snow cover and decreased biological activity in winter resulted in the disappearance of microbes from the coarse fluorescent bioaerosol particles. In this study, the highest fungal concentration in PM_{2.5} was observed in summer (641 spores m^{-3}), whereas the highest value in PM₁ was found in autumn

(1033 spores m⁻³), indicating different origins of fungal spores. Huffman et al. (2010) found that long-range transport of aerosols and anthropogenic sources such as combustion influence the fluorescent biological aerosol particles having diameters less than 1 μm. During the autumn sampling, no obvious straw combustion phenomena occurred, and we detected some long-range transportation events in November 2014. Long-range transported airborne PM were mainly derived from the outer Mongolia regions, well-known to be one of the dustiest places in East Asia (6 November), Siberia (3 and 12 November), and the Taklimakan and Gobi desert regions (5 November). Influenced by the air movements from the desert region, the corresponding fungal abundance increased from 6.18×10^4 to 10.3×10^5 copies m⁻³ (about 16.7-fold). Similarly, the corresponding fungal abundance influenced by air parcels from the Siberian, and Taklimakan and Gobi desert regions increased to 22.2×10^4 copies m⁻³ and 18.3×10^4 copies m⁻³, respectively. Hence, we hypothesized that the long-range transport of air parcels from north China might have contributed to the fungal enrichment of PM₁. In addition, the increased fungal abundance might be explained by meteorological diversity (Abdel Hameed et al., 2012). Low wind speed hinders fungal dispersal owing to the accumulation effect. According to Almaguer et al. (2014), in Cuba, the calm winds coming from the southwest direction induce the accumulation of fungal spores over the northern coast of the island. Lin et al. (2000) observed a strongly negative correlation between wind speeds of 4 m s^{-1} and fungal concentration; the fungal concentration increased as the wind speed became higher than 5 m s^{-1} in the Taipei area. In our present study, the fungal abundance in PM₁ showed no obvious increase under breezy conditions (wind speed 2 m s^{-1}) mainly from the southern direction (Fig. 1). When the wind speed was higher than 2 m s^{-1} , the fungal abundance increased markedly under the influence of westerly winds. As the westerly wind velocity increased, the fungal concentration increased slowly. Meanwhile, in PM_{2.5}, the fungal abundance increased with wind velocities higher than 2 m s^{-1} , mainly from the northwest direction of the continental areas, where diverse vegetation grows. The phenomenon implies that westerly and northwesterly winds might highly induce fungal growth and abundance in PM at Mt. Tai.

3.2 Contribution of spores to OC concentrations and PM mass

OC, accounting for 7–80 % of PM mass, constitutes a significant fraction of atmospheric aerosols (Yu et al., 2004; Ram et al., 2012; Ho et al., 2012). Ambient fungi are considered a possible source of OC in PMs. Cheng et al. (2009) estimated the mean fungal OC concentrations in Hong Kong to be 3.7, 6.0, and 9.7 ng m^{-3} , corresponding to 0.1, 1.2, and 0.2 % of the total OC in PM_{2.5}, PM_{2.5–10}, and PM₁₀, respectively. In the present study, the range and average concentrations of fungal contribution to atmospheric OC and mass




Figure 2. Statistical comparisons of OTUs, and Chao1 and Shannon indices among summer, autumn, and winter in PM_{2.5} and PM₁.

concentration for PM_{2.5} and PM₁ are listed in Table 1. The daily averaged concentrations of fungal OC in PM_{2.5} and PM₁ were 6.1 and 8.3 ng C m^{-3} , respectively, with the respective contributions to PM being 0.067 and 0.096 %, indicating that airborne fungal spores as a minor source of carbonaceous aerosols cannot be ignored at Mt. Tai. The fungal contribution to OC obtained at Mt. Tai was comparable with that observed at an urban site in Hong Kong (3.7 ng C m^{-3} ; Cheng et al., 2009) but lower than that obtained at an urban site in Austria ($117.9 \text{ ng C m}^{-3}$; Bauer et al., 2008) and a forest site on Hainan Island ($147\text{--}923 \text{ ng C m}^{-3}$; Zhang et al., 2015). The discrepancy between the abovementioned studies can be justified by the difference in particle type studied (TSP, PM₁₀, PM_{2.5}, and PM₁), fungal concentration, spore carbon content, and assessment method (e.g., sugar alcohol, cultivation, mannitol, and light microscopy). On the basis of the same conversion factor of $13 \text{ pg C spore}^{-1}$ by mannitol, the results were much lower than that obtained at an urban site in Beijing ($0.3 \pm 0.2 \mu \text{ C m}^{-3}$; Liang et al., 2017), implying a lower fungal concentration at Mt. Tai than that in Beijing. More studies are needed to better understand the spatial, temporal, and size distributions of fungal OC contributions to atmospheric particles in urban areas in the North China Plain.

3.3 Taxonomic diversity and composition of ambient fungi

On average, 509 and 475 OTUs were obtained in PM_{2.5} and PM₁, respectively, which were higher than those obtained in earlier airborne fungal studies at the ground level in Beijing, China (34–285; Yan et al., 2016) and Rehovot, Israel (121–178; Dannemiller et al., 2014). The OTUs associated with

Table 2. The relative abundance of the top five orders and two genera in TSP, PM₁₀, PM_{2.5}, and PM₁.

Taxonomic level	Common fungi	RAS ^a	RAF ^b	References	Samplers	Sample type	Concentration or abundance
Genera	Alternaria	11.7	6.2	Adhikari et al. (2004)	Andersen sampler (Thermo Andersen, Smyrna, 300082-5211, USA)	TSP	2.6 %
				Dannemiller et al. (2014)	High volume PM10 samplers (Ecotech, Knoxfield, VIC, Australia)	PM ₁₀	> 1 %
				Alghamdi et al. (2014)	PM _{2.5} samplers (Staplex Air Sampler Division, USA)	PM _{2.5}	2.6 %
	Aspergillus	2.3	1.9	Gou et al. (2016)	Low volume air sampler (BGI, USA)	PM ₁	> 1 %
				Cao et al. (2014)	Air samplers (Thermo Electron Corp., MA, USA)	PM ₁₀ and PM _{2.5}	abundant
				Gou et al. (2016)	Low volume air sampler (BGI, USA)	PM ₁₀ and PM ₁	abundant
Order	Pleosporales	18.4	45.4	Rittenour et al. (2014)	Buck Bioaire sampler (A.P. Buck, Inc, Orlando, FL, USA)	TSP	46 %
				Yan et al. (2016)	Air samplers (Air Metrics, USA, 5 L min ⁻¹)	PM ₁₀ and PM _{2.5}	29.4 %
				Gou et al. (2016)	Low volume air sampler (BGI, USA)	PM ₁₀ and PM ₁	10–15 %
	Xylariales	5.0	14.4	Womack et al. (2015)	SKC Biosamplers (BioSampler SKC Inc.)	TSP	abundant
				Gou et al. (2016)	Low volume air sampler (BGI, USA)	PM ₁₀ and PM ₁	0–5 %
	Eurotiales	4.8	13.3	Yan et al. (2016)	Air samplers (Air Metrics, USA, 5 L min ⁻¹)	PM ₁₀ and PM _{2.5}	10.6 %
				Gou et al. (2016)	Low volume air sampler (BGI, USA)	PM ₁₀ and PM ₁	10–15 %
	Capnodiales	4.4	12.5	Yan et al. (2016)	Air samplers (Air Metrics, USA, 5 L min ⁻¹)	PM ₁₀ and PM _{2.5}	27.96 %
				Gou et al. (2016)	Low volume air sampler (BGI, USA)	PM ₁₀ and PM ₁	~ 25 %
	Polyporales	2.5	6.4	Womack et al. (2015)	SKC Biosamplers (BioSampler SKC Inc.)	TSP	abundant
				Yan et al. (2016)	Air samplers (Air Metrics, USA, 5 L min ⁻¹)	PM ₁₀ and PM _{2.5}	3.6 %
				Yamamoto et al. (2012)	Eight-stage Andersen sampler (New Star Environmental, Roswell, GA, USA)	PM with aerodynamic diameter is 2.1–3.3, 3.3–4.7, 4.7–5.8, 5.8–9.0 and > 9.0 μm	abundant

RAS^a indicates relative abundance in submicron particles and RAF^b indicates relative abundance in fine particles.

PM_{2.5} in summer, autumn, and winter were higher than those associated with PM₁, implying more diverse fungal spores in PM_{2.5}. However, the Shannon and Chao1 indices showed different trends in PM_{2.5} and PM₁ (Fig. 2). The ambient fungi showed the highest richness and diversity in winter, followed by autumn and summer. Although PM₁ mass concentration dominated in autumn and winter, the corresponding fungal diversity was lower than that in PM_{2.5}. Similarly, the dominant PM_{2.5} mass concentration in summer presented lower diversity than that in PM₁.

In the fungal community, AMC (89.7 %) and BMC (7.0 %) were the predominant phyla, and they are known to actively discharge spores into the atmosphere (Fig. 3a). The remaining phyla were Zygomycota (ZMC) and Glomeromycota. AMC and BMC present a global pattern across continental (Austria, Arizona, Brazil, and Germany), coastal (Taiwan, Puerto Rico, and UK), and marine sites (Pacific, Indian, Atlantic, and Southern Ocean) (Frohlich-Nowoisky et al., 2012). In continental samples, BMC (64 %) seems to be more abundant than AMC (34 %), whereas in marine sites, AMC (72 %) is about 2.6 times more abundant than BMC. Herein, the abundance of AMC was approximately 12.8 times higher than that of BMC. Members of AMC have single-celled or filamentous vegetative growth forms that are easily aerosolized, unlike BMC (Womack et al., 2015). Furthermore, 10 classes belonging to AMC, 10 to BMC, and 1 to ZMC were observed (Fig. 3b). The preponderant classes belonging to AMC were Dothideomycetes (37.3 %),

Sordariomycetes (15.0 %), and Eurotiomycetes (6.1 %). The dominant orders in Dothideomycetes included Pleosporales (14.9 %), Capnodiales (5.3 %), and Botryosphaerales (1.6 %) (Fig. 3c). Pleosporales has been reported to include fungi allergenic to local residents (Rittenour et al., 2014). The values were lower than those reported in Beijing's PM (Pleosporales: 29.39 % and Capnodiales: 27.96 %) (Yan et al., 2016). Likewise, the dominant classes in BMC were Agaricomycetes (4.4 %) and Tremellomycetes (1.5 %), including the orders Polyporales (2.5 %), Agaricales (1.6 %), and Tremellales (1.2 %). About 291 taxa from the genus level were determined, including *Alternaria*, *Glomerella*, *Zasmidium*, *Pestalotiopsis*, *Aspergillus*, and *Phyllosticta*. The distribution was discrepant with that at the ground level, wherein *Cladosporium* occupied more than 50 % of total fungi, followed by *Alternaria*, *Didymella*, and *Khuskia* (Oh et al., 2014). The top five orders (Pleosporales, Xylariales, Eurotiales, Capnodiales, Polyporales) and genera (*Alternaria* and *Aspergillus*) were commonly observed in suspended aerosol particles (including TSP, PM₁₀, PM_{2.5}, and PM₁) but showed variable relative abundances, as shown in Table 2. We attribute this disparity to the different sampling approaches, instruments, and analysis methods. This aspect needs to be probed and studied in depth in the future.




Figure 3. Relative abundances of fungal communities of PM_{2.5} and PM₁ at phylum (a), class (b), and order level (c).

3.4 Implication of the allergenic and pathogenic fungi

To date, about 123 fungal genera (mainly belonging to the phylum AMC) have been identified to be human allergens (Simon-Nobbe et al., 2008). Of the 11 potentially allergy-inducing AMC species and 1 potentially allergy-inducing BMC species found at Mt. Tai, the 3 most common species were *Aspergillus flavus*, *Blumeria graminis*, and *Saccharomyces cerevisiae*. *Aspergillus flavus* is a common human pathogen found in air, and it is also a human allergen and mycotoxin producer (Adhikari et al., 2004). It is associated with invasive aspergillosis and superficial infections (Hedayati et al., 2007). *Blumeria graminis*, found on the surface of plant leaves, causes powdery mildew on cereal plants (Belanger et al., 2003). Such pathogens and allergens are expected to be widely spread around the atmospheric environment in temperate and tropical zones (Vermani et al., 2010). Our results also revealed that the abundance of potential allergenic and pathogenic fungal spores in summer were the highest compared to those in autumn and winter. Clinicians should consider the fungal spores described herein as a possible cause of human and plant disease under long exposure to airborne particles throughout the year, especially in the summer season. Furthermore, the abundance of the above-mentioned allergenic and pathogenic fungal spores in PM₁ was about 3.8 times higher than that in PM_{2.5} in summer, implying relatively higher health risks for smaller particles.

Residents and even visitors at Mt. Tai should be warned about this phenomenon.

3.5 Size distribution and seasonal variation of fungal communities

Both fungal abundance and fungal community show a seasonal trend across different size fractions (Awad et al., 2013). Yamamoto et al. (2012) observed that the pathogenic fungi were mainly detected at PM_{4.7} (PM with aerodynamic diameter < 4.7 μm), while the allergenic fungi existed primarily at PM sizes with aerodynamic diameter > 9 μm. In the present study, a discrepant size distribution of the fungal community was observed according to the Metastat analysis by permutation *t*-tests (Table 3). *Glomerella*, *Zasmidium*, and *Phyllosticta* were abundantly enriched in PM_{2.5}, while the abundance of *Preussia*, *Truncatella*, *Umbelopsis*, *Sebacina*, and *Cordyceps* increased in PM₁. The Kruskal–Wallis rank sum test showed that 6 fungal genera had apparent seasonal variation (Fig. 4). *Glomerella* and *Zasmidium* increased in autumn and decreased as the particle size increased. *Glomerella* was widely found on the surface of leaves, suggesting that leaf senescence is an important source of fungi in PM_{2.5} in autumn (Wang et al., 2015). Some crucial environmental factors having a potential influence on fungal release and growth, such as temperature; NO₂; PM₁₀; SO₂; CO; relative humidity (Yan et al., 2016); radiation; vegetation




Figure 4. Variance analysis of fungal genera based on the Kruskal–Wallis rank sum test.

Table 3. Metastats analysis showing the fungal genera that are significantly different among PM_{2.5} and PM₁.

Taxa	PM ₁			PM _{2.5}			<i>p</i> value	<i>q</i> value
	mean	SE	variance	mean	SE	variance		
Glomerella	10.51984	0.021813	0.013798	22.49025	0.01807	0.009796	0.000999	0.025543
Zasmidium	6.523201	0.011769	0.004017	12.71881	0.012239	0.004494	0.000999	0.025543
Phyllosticta	2.507228	0.004038	0.000473	5.948659	0.004366	0.000572	0.000999	0.025543
Trussia	0.039161	0.000195	1.10E-06	0.009109	6.81E-05	1.39E-07	0.002322	0.042885
Truncatella	0.030579	0.00024	1.67E-06	0.005152	2.44E-05	1.79E-08	0.002784	0.046274
Umbelopsis	0.027549	0.000252	1.84E-06	0.005369	2.55E-05	1.95E-08	0.001669	0.034675
Sebacina	0.021306	0.000196	1.11E-06	0.001261	1.26E-05	4.77E-09	0.000550	0.022857
Cordyceps	0.020939	0.000137	5.45E-07	0.002518	1.75E-05	9.18E-09	0.001392	0.030848

(Moreau et al., 2016); urbanization; and accidental events, e.g., dust storms (Prospero et al., 2005), rainfall (Zhang et al., 2015), hurricanes (DeLeon-Rodriguez et al., 2013), and haze (Yan et al., 2016), have been identified. Herein Spearman's rank coefficient analysis indicated that Ca²⁺, a typical water-soluble inorganic ion from dust, was negatively related to the prevalence of *Glomerella* and *Zasmidium* in autumn (Fig. 5). The increase of *Penicillium*, *Bullera*, and *Geosmithia* in winter is ascribed to their sensitivity to low temperature (Sousa et al., 2008; Abdel Hameed et al., 2012). The results based on Spearman's rank correlation test analy-

sis support this notion (Fig. 5, $p < 0.01$). Humidity, another important factor for fungal release into the atmosphere either by active or passive modes, is a crucial factor for the variation in fungal spores such as *Lophium* ($p < 0.01$), *Cenococcum* ($p < 0.05$), *Tricholoma* ($p < 0.05$), and *Candida* ($p < 0.05$). In summer, no distinct difference was observed based on the top 40 fungal genera (Fig. 4). However, some trace fungal genera presented an inverse correlation with temperature (*Coccomyces*, $p < 0.01$; and *Dictyosporium*, $p < 0.01$), humidity (*Botryosphaeria*, $p < 0.001$; *Coccomyces*, $p < 0.01$; and *Dictyosporium*, $p < 0.01$), PM_{2.5} (*Acremonium*, $p < 0.01$;




Figure 5. Heatmap analysis of the top 64 fungal genera based on Spearman's rank correlations (***) $p < 0.001$; ** $p < 0.01$; * $p < 0.05$). Red arrows indicate that the specific fungi varied significantly in different seasons.

Phoma, $p < 0.01$), and Ca^{2+} (*Talaromyces*, $p < 0.01$; *Acaromyces*, $p < 0.01$). The crucial environmental factors we identified contributed to the variation in the fungal community. Due to the limited culture studies on the mechanism for the effects of environmental factors on specific fungal spores, the relationship between bioaerosols and environmental factors still needs to be surveyed over a longer duration.

4 Conclusions

Diverse airborne fungal spores are relevant for studies on the atmosphere, biogeoscience, climate and ecology, environmental hygiene, agriculture, and bioengineering. As the details of fungal spores present at high-elevation sites re-

main unknown, the detection and characterization of ambient fungi can help elucidate the regional and global distribution of diverse fungi. To that end, we provide a comprehensive framework of the fungal abundance and communities associated with different seasons across PM_{2.5} and PM₁ at Mt. Tai. The results revealed that the concentration and fungal community structure at Mt. Tai differs considerably from those reported for surface ground sites. Over the sampling period, average fungal concentrations of 467 and 644 spores m^{-3} in PM_{2.5} and PM₁, respectively, were calculated. In addition to long-distance air-mass movement events, westerly and northwesterly winds also favor the increase in fungal abundance. Diverse fungal communities presented significant seasonal variation across different size particles. The prevalence of *Glomerella* and *Zasmidium* increased in autumn and de-

creased as the particle size increased. In winter, the prevalence of *Penicillium*, *Bullera*, and *Geosmithia* increased with the decrease in particle size. No distinct disparity was observed in summer. The variation in fungal profile can be influenced by environmental factors including humidity, temperature, wind speed, PM_{2.5}, and some chemical components in PMs (including Ca²⁺). Nevertheless, the detailed specific effects of environmental factors on the ambient fungal community remain poorly explained. In further studies, a combination of traditional culture-based methods and metagenomics may help answer the various unresolved questions.

Data availability. The meteorological data are accessible from <http://www.underground.com> at a resolution of 3 h. Mass concentration and chemical properties of PM_{2.5} and PM₁ are available from the authors upon request (caihongxu@mail.sdu.edu.cn). The back trajectory data from the Hybrid Single Particle Lagrangian Integrated Trajectory (HYSPPLIT) model are available at http://ready.arl.noaa.gov/HYSPLIT_traj.php. The raw 18S rRNA gene sequences are available at the NCBI Sequence Read Archive database under accession number SRR5146156.

Competing interests. The authors declare that they have no conflict of interest.

Special issue statement. This article is part of the special issue “Regional transport and transformation of air pollution in eastern China”. It is not associated with a conference.

Acknowledgements. This work was supported by the National Natural Science Foundation of China (no. 41375126, 21527814), Taishan Scholar Grant (grant number ts20120552), Cyrus Tang Foundation (no. CTF-FD2014001), the Ministry of Science and Technology of China (no. 2016YFC0202701, 2014BAC22B01), and the European Union’s Horizon 2020 research and innovation programme under grant agreement no. 690958 (MARSU project).

Edited by: Tong Zhu

Reviewed by: four anonymous referees

References

- Abdel Hameed, A. A., Khoder, M. I., Ibrahim, Y. H., Saeed, Y., Osman, M. E., and Ghanem, S.: Study on some factors affecting survivability of airborne fungi, *Sci. Total Environ.*, 414, 696–700, 2012.
- Adhikari, A., Sen, M. M., Gupta-Bhattacharya, S., and Chanda, S.: Airborne viable, non-viable, and allergenic fungi in a rural agricultural area of India: a 2-year study at five outdoor sampling stations, *Sci. Total Environ.*, 326, 123–141, 2004.
- Almaguer, M., Aira, M. J., Rodriguez-Rajo, F. J., and Rojas, T. I.: Temporal dynamics of airborne fungi in Havana (Cuba) during

- dry and rainy seasons: influence of meteorological parameters, *Int. J. Biometeorol.*, 58, 1459–1470, 2014.
- Awad, A. H., Gibbs, S. G., Tarwater, P. M., and Green, C. F.: Seasonal evaluation of fine and coarse culturable bacterial aerosols from residences within a rural and an urban city in Egypt, *Int. J. Environ. Heal. R.*, 23, 269–280, 2013.
- Bauer, H., Schueller, E., Weinke, G., Berger, A., Hitzemberger, R., Marr, I. L., and Puxbaum, H.: Significant contributions of fungal spores to the organic carbon and to the aerosol mass balance of the urban atmospheric aerosol, *Atmos. Environ.*, 42, 5542–5549, 2008.
- Belanger, R. R., Benhamou, N., and Menzies, J. G.: Cytological evidence of an active role of silicon in wheat resistance to powdery mildew (*Blumeria graminis f. sp. tritici*), *Phytopathology*, 93, 402–412, 2003.
- Bowers, R. M., Lauber, C. L., Wiedinmyer, C., Hamady, M., Hallar, A. G., Fall, R., Knight, R., and Fierer, N.: Characterization of airborne microbial communities at a high-elevation site and their potential to act as atmospheric ice nuclei, *Appl. Environ. Microbiol.*, 75, 5121–5130, 2009.
- Bowers, R. M., McCubbin, I. B., Hallar, A. G., and Fierer, N.: Seasonal variability in airborne bacterial communities at a high-elevation site, *Atmos. Environ.*, 50, 41–49, 2012.
- Bowers, R. M., Clements, N., Emerson, J. B., Wiedinmyer, C., Hannigan, M. P., and Fierer, N.: Seasonal variability in bacterial and fungal diversity of the near-surface atmosphere, *Environ. Sci. Technol.*, 47, 12097–12106, 2013.
- Brauer, M., Freedman, G., Frostad, J., van Donkelaar, A., Martin, R. V., Dentener, F., van Dingenen, R., Estep, K., Amini, H., Apte, J. S., Balakrishnan, K., Barregard, L., Broday, D., Feigin, V., Ghosh, S., Hopke, P. K., Knibbs, L. D., Kokubo, Y., Liu, Y., Ma, S., Morawska, L., Sangrador, J. L., Shaddick, G., Anderson, H. R., Vos, T., Forouzanfar, M. H., Burnett, R. T., and Cohen, A.: Ambient air pollution exposure estimation for the global burden of disease 2013, *Environ. Sci. Technol.*, 50, 79–88, 2013.
- Brodie, E. L., DeSantis, T. Z., Parker, J. P., Zubietta, I. X., Piceno, Y. M., and Andersen, G. L.: Urban aerosols harbor diverse and dynamic bacterial populations, *P. Natl. Acad. Sci. USA*, 104, 299–304, 2007.
- Cao, C., Jiang, W., Wang, B., Fang, J., Lang, J., Tian, G., Jiang, J., and Zhu, T. F.: Inhalable microorganisms in Beijing’s PM_{2.5} and PM₁₀ pollutants during a severe smog event, *Environ. Sci. Technol.*, 48, 1499–1507, 2014.
- Cheng, J. Y. W., Chan, C. K., Lee, C. T., and Lau, A. P. S.: Carbon content of common airborne fungal species and fungal contribution to aerosol organic carbon in a subtropical city, *Atmos. Environ.*, 43, 2781–2787, 2009.
- Christner, B. C., Morris, C. E., Foreman, C. M., Cai, R., and Sands, D. C.: Ubiquity of biological ice nucleators in snowfall, *Science*, 319, 1214, <https://doi.org/10.1126/science.1149757>, 2008.
- Creamean, J. M., Suski, K. J., Rosenfeld, D., Cazorla, A., DeMott, P. J., Sullivan, R. C., White, A. B., Ralph, F. M., Minnis, P., Comstock, J. M., Tomlinson, J. M., and Prather, K. A.: Dust and biological aerosols from the Sahara and Asia influence precipitation in the western US, *Science*, 339, 1572–1578, 2013.
- Dannemiller, K. C., Lang-Yona, N., Yamamoto, N., Rudich, Y., and Peccia, J.: Combining real-time PCR and next-generation DNA sequencing to provide quantitative comparisons of fungal aerosol populations, *Atmos. Environ.*, 84, 113–121, 2014.

- DeLeon-Rodriguez, N., Latham, T. L., Rodriguez, R. L., Barazesh, J. M., Anderson, B. E., Beyersdorf, A. J., Ziemba, L. D., Bergin, M., Nenes, A., and Konstantinidis, K. T.: Microbiome of the upper troposphere: species composition and prevalence, effects of tropical storms, and atmospheric implications, *P. Natl. Acad. Sci. USA*, 110, 2575–2580, 2013.
- Deng, W. J., Chai, Y. M., Lin, H. Y., So, W. W. M., Ho, K. W. K., Tsui, A. K. Y., and Wong, P. K. S.: Distribution of bacteria in inhalable particles and its implications for health risks in kindergarten children in Hong Kong, *Atmos. Environ.*, 128, 268–275, 2016, .
- Després, V. R., Nowoisky, J. F., Klose, M., Conrad, R., Andreae, M. O., and Pöschl, U.: Characterization of primary biogenic aerosol particles in urban, rural, and high-alpine air by DNA sequence and restriction fragment analysis of ribosomal RNA genes, *Biogeosciences*, 4, 1127–1141, <https://doi.org/10.5194/bg-4-1127-2007>, 2007.
- Despres, V. R., Huffman, J. A., Burrows, S. M., Hoose, C., Safatov, A. S., Buryak, G., Fröhlich-Nowoisky, J., Elbert, W., Andreae, M. O., and Pöschl, U.: Primary biological aerosols in the atmosphere: A review of observations and relevance, *Tellus B*, 64, 1–58, 2012.
- Edgar, R. C., Haas, B. J., Clemente, J. C., Quince, C., and Knight, R.: UCHIME improves sensitivity and speed of chimera detection, *Bioinformatics*, 27, 2194–2200, 2011.
- Elbert, W., Taylor, P. E., Andreae, M. O., and Pöschl, U.: Contribution of fungi to primary biogenic aerosols in the atmosphere: wet and dry discharged spores, carbohydrates, and inorganic ions, *Atmos. Chem. Phys.*, 7, 4569–4588, <https://doi.org/10.5194/acp-7-4569-2007>, 2007.
- Fröhlich-Nowoisky, J., Pickersgill, D. A., Despres, V. R., and Pöschl, U.: High diversity of fungi in air particulate matter, *P. Natl. Acad. Sci. USA*, 106, 12814–12819, 2009.
- Fröhlich-Nowoisky, J., Burrows, S. M., Xie, Z., Engling, G., Solomon, P. A., Fraser, M. P., Mayol-Bracero, O. L., Artaxo, P., Begerow, D., Conrad, R., Andreae, M. O., Després, V. R., and Pöschl, U.: Biogeography in the air: fungal diversity over land and oceans, *Biogeosciences*, 9, 1125–1136, <https://doi.org/10.5194/bg-9-1125-2012>, 2012.
- Gandolfi, I., Bertolini, V., Bestetti, G., Ambrosini, R., Innocente, E., Rampazzo, G., Papacchini, M., and Franzetti, A.: Spatio-temporal variability of airborne bacterial communities and their correlation with particulate matter chemical composition across two urban areas, *Appl. Microbiol. Biotechnol.*, 99, 4867–4877, 2015.
- Gao, J. F., Fan, X. Y., Li, H. Y., and Pan, K. L.: Airborne bacterial communities of PM_{2.5} in Beijing-Tianjin-Hebei megalopolis, China as revealed by Illumina MiSeq sequencing: a case study, *Aerosol Air Qual. Res.*, 17, 778–798, 2017a.
- Gao, M., Jia, R. Z., Qiu, T. L., Han, M. L., and Wang, X. M.: Size-related bacterial diversity and tetracycline resistance gene abundance in the air of concentrated poultry feeding operations, *Environ. Pollut.*, 220, 1342–1348, 2017b.
- Gou, H., Lu, J. J., Li, S., Tong, Y. B., Xie, C. B., and Zheng, X.: Assessment of microbial communities in PM₁ and PM₁₀ of Urumqi during winter, *Environ. Pollut.*, 214, 202–210, 2016.
- Hawksworth, D. L.: The magnitude of fungal diversity: the 1.5 million species estimate revisited, *Mycol. Res.*, 105, 1422–1432, 2001.
- Heald, C. L. and Spracklen, D. V.: Atmospheric budget of primary biological aerosol particles from fungal spores, *Geophys. Res. Lett.*, 36, 269–277, 2009.
- Hedayati, M. T., Pasqualotto, A. C., Warn, P. A., Bowyer, P., and Denning, D. W.: *Aspergillus flavus*: human pathogen, allergen and mycotoxin producer, *Microbiology*, 153, 1677–1692, 2007.
- Ho, K. F., Lee, S. C., Yu, J. C., Zou, S. C., and Fung, K.: Carbonaceous characteristics of atmospheric particulate matter in Hong Kong, *Sci. Total Environ.*, 300, 59–67, 2002.
- Huffman, J. A., Treutlein, B., and Pöschl, U.: Fluorescent biological aerosol particle concentrations and size distributions measured with an Ultraviolet Aerodynamic Particle Sizer (UV-APS) in Central Europe, *Atmos. Chem. Phys.*, 10, 3215–3233, <https://doi.org/10.5194/acp-10-3215-2010>, 2010.
- Jiang, W. J., Liang, P., Wang, B. Y., Fang, J. H., Lang, J. D., Tian, G., Jiang, J. K., and Zhu, T. F.: Optimized DNA extraction and metagenomic sequencing of airborne microbial communities, *Nat. Protoc.*, 10, 768–779, 2015.
- Jaenicke, R.: Abundance of cellular material and proteins in the atmosphere, *Science*, 308, p. 73, 2005.
- Judd, C. R., Koyama, A., Simmons, M. P., Brewer, P. E., and von Fischer, J. C.: Co-variation in methanotroph community composition and activity in three temperate grassland soils, *Soil Biol. Biochem.*, 95, 78–86, 2016.
- Koljalg, U., Nilsson, R. H., and Abarenkov, K.: Towards a unified paradigm for sequence-based identification of fungi, *Mol. Ecol.*, 22, 5271–5277, 2013.
- Koiv, V., Roosaare, M., Vedler, E., Kivistik, P. A., Toppi, K., Schryer, D. W., Remm, M., Tenson, T., and Mae, A.: Microbial population dynamics in response to *Pectobacterium atrosepticum* infection in potato tubers, *Sci. Rep-UK*, 5, 11606, <https://doi.org/10.1038/srep11606>, 2015.
- Kuczynski, J., Stombaugh, J., Walters, W. A., Gonzalez, A., Gregory Caporaso, J., and Knight, R.: Using QIIME to analyze 16S rRNA gene sequences from microbial communities, *Curr. Protoc. Bioinformatics*, Chapter 10, Unit 10.7, <https://doi.org/10.1002/0471250953.bi1007s36>, 2011.
- Lee, S. H., Lee, H. J., Kim, S. J., Lee, H. M., Kang, H., and Kim, Y. P.: Identification of airborne bacterial and fungal community structures in an urban area by T-RFLP analysis and quantitative real-time PCR, *Sci. Total Environ.*, 408, 1349–1357, 2010.
- Liang, L., Engling, G., Du, Z., Duan, F., Cheng, Y., Liu, X., and He, K.: Contribution of fungal spores to organic carbon in ambient aerosols in Beijing, China, *Atmos. Pollut. Res.*, 8, 351–358, 2017.
- Lin, W. H. and Li, C. S.: Associations of fungal aerosols, air pollutants, and meteorological factors, *Aerosol Sci. Technol.*, 32, 359–368, 2000.
- Manter, D. K. and Vivanco, J. M.: Use of the ITS primers, ITS1F and ITS4, to characterize fungal abundance and diversity in mixed-template samples by qPCR and length heterogeneity analysis, *J. Microbiol. Meth.*, 71, 7–14, 2007.
- Miettinen, H., Kietavainen, R., Sohlberg, E., Numminen, M., Ahonen, L., and Itavaara, M.: Microbiome composition and geochemical characteristics of deep subsurface high-pressure environment, Pyhasalmi mine Finland, *Front. Microbiol.*, 6, 1203, <https://doi.org/10.3389/fmicb.2015.01203>, 2015.
- Moreau, C. S., Weigl, F., Tischer, C., Probst, A. J., Heinrich, J., Markevych, I., Jochner, S., and Pritsch, K.: Fun-

- gal and bacterial communities in indoor dust follow different environmental determinants, *Plos One*, 11, e0154131, <https://doi.org/10.1371/journal.pone.0154131>, 2016.
- Morris, C. E., Kinkel, L.L., Lindow, S. E., Hechtponar, E. I., and Elliott, V. J.: Fifty years of phyllosphere microbiology: significant contributions to research in related fields, *Phyllosphere Microbiology*, St. Paul, MN, USA, APS Press, 365–375, 2002.
- Oh, S. Y., Fong, J. J., Park, M. S., Chang, L., and Lim, Y. W.: Identifying airborne fungi in Seoul, Korea using metagenomics, *J. Microbiol.*, 52, 465–472, 2014.
- Oliveira, M., Ribeiro, H., Delgado, J. L., and Abreu, I.: The effects of meteorological factors on airborne fungal spore concentration in two areas differing in urbanisation level, *Int. J. Biometeorol.*, 53, 61–73, 2009.
- Pratt, K. A., DeMott, P. J., French, J. R., Wang, Z., Westphal, D. L., Heymsfield, A. J., Twohy, C. H., Prenni, A. J., and Prather, K. A.: In situ detection of biological particles in cloud ice-crystals, *Nat. Geosci.*, 2, 398–401, 2009.
- Prospero, J. M., Blades, E., Mathison, G., and Naidu, R.: Interhemispheric transport of viable fungi and bacteria from Africa to the Caribbean with soil dust, *Aerobiologia*, 21, 1–19, 2005.
- Ram, K., Sarin, M. M., and Tripathi, S. N.: Temporal trends in atmospheric PM_{2.5}, PM₁₀, elemental carbon, organic carbon, water-soluble organic carbon, and optical properties: impact of biomass burning emissions in the Indo Gangetic Plain, *Environ. Sci. Technol.*, 46, 686–695, 2012.
- Rittenour, W. R., Ciaccio, C. E., Barnes, C. S., Kashon, M. L., Lemons, A. R., Beezhold, D. H., and Green, B. J.: Internal transcribed spacer rRNA gene sequencing analysis of fungal diversity in Kansas City indoor environments, *Environ. Sci. Proc. Impacts*, 16, 33–43, 2014.
- Ryan, P. H., Bernstein, D. I., Lockey, J., Reponen, T., Levin, L., Grinshpun, S., Villareal, M., Hershey, G. K. K., Burkle, J., and Lemasters, G.: Exposure to traffic-related particles and endotoxin during infancy is associated with wheezing at age 3 years, *Am. J. Resp. Crit. Care*, 180, 1068–1075, 2009.
- Saari, S.: Seasonal and diurnal variations of fluorescent bioaerosol concentration and size distribution in the urban environment, *Aerosol Air Qual. Res.*, 15, 572–581, 2015.
- Simon-Nobbe, B., Denk, U., Poll, V., Rid, R., and Breitenbach, M.: The spectrum of fungal allergy, *Int. Arch. Allergy Imm.*, 145, 58–86, 2008.
- Smets, W., Moretti, S., Denys, S., and Lebeer, S.: Airborne bacteria in the atmosphere: presence, purpose, and potential, *Atmos. Environ.*, 139, 214–221, 2016.
- Sousa, S. I. V., Martins, F. G., Pereira, M. C., Alvim-Ferraz, M. C. M., Ribeiro, H., Oliveira, M., and Abreu, I.: Influence of atmospheric ozone, PM₁₀ and meteorological factors on the concentration of airborne pollen and fungal spores, *Atmos. Environ.*, 42, 7452–7464, 2008.
- Tong, X. L., Xu, H. T., Zou, L. H., Cai, M., Xu, X. F., Zhao, Z. T., Xiao, F., and Li, Y. M.: High diversity of airborne fungi in the hospital environment as revealed by meta-sequencing-based microbiome analysis, *Sci. Rep.-UK*, 7, 39606, <https://doi.org/10.1038/srep39606>, 2017.
- Tsai, F. C., Macherb, J. M., and Hung, Y. Y.: Biodiversity and concentrations of airborne fungi in large US office buildings from the BASE study, *Atmos. Environ.*, 41, 5181–5191, 2007.
- van Doorn, R., Szemes, M., Bonants, P., Kowalchuk, G. A., Salles, J. F., Ortenberg, E., and Schoen, C. D.: Quantitative multiplex detection of plant pathogens using a novel ligation probe-based system coupled with universal, high-throughput real-time PCR on OpenArrays, *BMC genomics*, 8, 1–4, 2007.
- Vermani, M., Vijayan, V. K., Kausar, M. A., and Agarwal, M. K.: Quantification of airborne *Aspergillus* allergens: redefining the approach, *J. Aasthma.*, 47, 754–761, 2010.
- Wang, L., Song Z. W., Xu, A. L., Wu, D. D., and Yan, X.: Phylogenetic diversity of airborne microbes in Qingdao downtown in autumn, *J. Appl. Ecol.*, 26, 1121–1129, 2015 (in Chinese).
- Wang, X., Chen, J., Sun, J., Li, W., Yang, L., Wen, L., Wang, W., Wang, X., Collett Jr, J. L., Shi, Y., Zhang, Q., Hu, J., Yao, L., Zhu, Y., Sui, X., Sun, X., and Mellouki, A.: Severe haze episodes and seriously polluted fog water in Ji'nan, China, *Sci. Total Environ.*, 493, 133–137, 2014.
- Wei, M., Xu, C., Chen, J., Zhu, C., Li, J., and Lv, G.: Characteristics of bacterial community in cloud water at Mt Tai: similarity and disparity under polluted and non-polluted cloud episodes, *Atmos. Chem. Phys.*, 17, 5253–5270, doi:10.5194/acp-17-5253-2017, 2017.
- Womack, A. M., Artaxo, P. E., Ishida, F. Y., Mueller, R. C., Saleska, S. R., Wiedemann, K. T., Bohannon, B. J. M., and Green, J. L.: Characterization of active and total fungal communities in the atmosphere over the Amazon rainforest, *Biogeosciences*, 12, 6337–6349, <https://doi.org/10.5194/bg-12-6337-2015>, 2015.
- Womilolu, T. O., Miller, J. D., Mayer, P. M., and Brook, J. R.: Methods to determine the biological composition of particulate matter collected from outdoor air, *Atmos. Environ.*, 37, 4335–4344, 2003.
- Xu, C. H., Wei, M., Chen, J. M., Wang, X. F., Sui, X., Zhu, C., Li, J. R., Zheng, L. L., Sui, G. D., Li, W. J., Wang, W. X., Zhang, Q. Z., and Mellouki, A.: Bacterial characterization in ambient submicron particles during severe haze episodes at Ji'nan, China, *Sci. Total Environ.*, 580, 188–1196, 2017a.
- Xu, C. H., Wei, M., Chen, J. M., Sui, X., Zhu, C., Li, J. R., Zheng, L. L., Sui, G. D., Li, W. J., Wang, W. X., Zhang, Q. Z., and Mellouki, A.: Investigation of diverse bacteria in cloud water at Mt. Tai, China, *Sci. Total Environ.*, 580, 258–265, 2017b.
- Yadav, R. K. P., Halley, J. M., Karamanoli, K., Constantinidou, H. I., and Vokou, D.: Bacterial populations on the leaves of Mediterranean plants: quantitative features and testing of distribution models, *Environ. Exp. Bot.*, 52, 63–77, 2004.
- Yamaguchi, N., Baba, T., Ichijo, T., Himezawa, Y., Enoki, K., Saraya, M., Li, P. F., and Nasu, M.: Abundance and community structure of bacteria on Asian dust particles collected in Beijing, China, during the Asian dust season, *Biological and Pharmaceutical Bulletin*, 39, 68–77, 2016.
- Yamamoto, N., Bibby, K., Qian, J., Hospodsky, D., Rismani-Yazdi, H., Nazaroff, W. W., and Peccia, J.: Particle-size distributions and seasonal diversity of allergenic and pathogenic fungi in outdoor air, *ISME J.*, 6, 1801–1811, 2012.
- Yan, D., Zhang, T., Su, J., Zhao, L. L., Wang, H., Fang, X. M., Zhang, Y. Q., Liu, H. Y., and Yu, L. Y.: Diversity and composition of airborne fungal community associated with particulate matters in Beijing during haze and non-haze days, *Front. Microbiol.*, 7, 1–12, 2016.
- Yu, J. Z., Tung, J. W. T., Wu, A. W. M., Lau, A. K. H., Louie, P. K. K., and Fung, J. C. F.: Abundance and seasonal characteristics

- of elemental and organic carbon in Hong Kong PM₁₀, *Atmos. Environ.*, 38, 1511–1521, 2004.
- Zhang, Z., Engling, G., Zhang, L., Kawamura, K., Yang, Y., Tao, J., Zhang, R., Chan, C. Y., and Li, Y.: Significant influence of fungi on coarse carbonaceous and potassium aerosols in a tropical rainforest, *Environ. Res. Lett.*, 10, 034015, <https://doi.org/10.1088/1748-9326/10/3/034015>, 2015.
- Zhong, X., Qi, J., Li, H., Dong, L., and Gao, D.: Seasonal distribution of microbial activity in bioaerosols in the outdoor environment of the Qingdao coastal region, *Atmos. Environ.*, 140, 506–513, 2016.
- Zhou, M., Liu, Y., Wang, L., Kuang, X., Xu, X., and Kan, H.: Particulate air pollution and mortality in a cohort of Chinese men, *Environ. Pollut.*, 186, 1–6, 2014.
- Zhu, C., Kawamura, K., Fukuda, Y., Mochida, M., and Iwamoto, Y.: Fungal spores overwhelm biogenic organic aerosols in a midlatitudinal forest, *Atmos. Chem. Phys.*, 16, 7497–7506, <https://doi.org/10.5194/acp-16-7497-2016>, 2016.