



HAL
open science

Intersection of moving convex sets in a normed space

Jean Jacques Moreau

► **To cite this version:**

Jean Jacques Moreau. Intersection of moving convex sets in a normed space. *Mathematica Scandinavica*, 1975, 36, pp.159-173. hal-01788536

HAL Id: hal-01788536

<https://hal.science/hal-01788536>

Submitted on 9 May 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

INTERSECTION OF MOVING CONVEX SETS IN A NORMED SPACE

J. J. MOREAU

To Werner Fenchel on his 70th birthday

1. Introduction.

Let I be an interval of \mathbb{R} ; let $t \mapsto A(t)$ or $t \mapsto A_t$ denote a *multifunction* (i.e. a set-valued mapping) from I into a metric space (E, d) . When t is interpreted as the time the language of kinematics may be used and A is called a *moving set* in E .

Several recent papers concerning evolution processes or selection properties (cf. [3], [4], [8], [9]) have drawn the attention to the concept of the *variation* of such a multifunction in the sense of the *Hausdorff distance* between subsets of (E, d) .

A classical preliminary in the study of Hausdorff distance consists in defining, for every two subsets C and D of (E, d) , the non symmetric *écart*

$$(1) \quad \begin{aligned} e(C, D) &= \sup_{x \in C} \inf_{y \in D} d(x, y) \\ &= \sup \{d(x, D) : x \in C\} \end{aligned}$$

which we propose to call *the metric excess of C over D* . The considered sets may be empty; let us agree that “sup” and “inf” above are understood in the sense of the ordered set $\bar{\mathbb{R}}_+ = [0, +\infty]$. The supremum of an empty collection of elements of this ordered set is 0 and the infimum is $+\infty$. Thus $e(\emptyset, D) = 0$ for any D , and $e(C, \emptyset) = +\infty$ for any $C \neq \emptyset$. One easily proves that the *écart* e satisfies the triangle inequality; clearly $e(C, D) = 0$ if and only if C is contained in the closure $\text{cl}D$ of D .

The Hausdorff distance between C and D is then

$$h(C, D) = \max \{e(C, D), e(D, C)\},$$

possibly infinite; it is zero if and only if C and D have the same closure.

Let $[s, t]$ be a compact subinterval of I ; for every finite subdivision of $[s, t]$, namely

$$S: \quad s = \tau_0 \leq \tau_1 \leq \dots \leq \tau_n = t,$$

put

$$V(S) = \sum_{i=1}^n h(A(\tau_{i-1}), A(\tau_i)) \in \bar{\mathbb{R}}_+.$$

In accordance with classical terminology, the supremum of $V(S)$ for S ranging over all the finite subdivisions of $[s, t]$ is called *the variation of the multifunction A over the interval $[s, t]$* ; it is denoted $\text{var}(A; s, t)$.

If, for every $[s, t] \subset I$, this variation is finite, let us say that A is of *finite variation in I* ; in this case there exists a non decreasing function $v: I \rightarrow \mathbb{R}$ (unique up to an additive constant), called *the indefinite variation of A in I* , such that for every $[s, t] \subset I$

$$\text{var}(A; s, t) = v(t) - v(s).$$

Let us say that the multifunction A is of *continuous finite variation in I* if v is continuous in I .

The *absolute continuity* of A over a compact subinterval K of I is defined, by means of Hausdorff distance, in the conventional way, i.e. for every $\varepsilon > 0$, there exists $\eta > 0$, such that the implication

$$\sum_i |\tau_i - \sigma_i| < \eta \Rightarrow \sum_i h(A(\sigma_i), A(\tau_i)) < \varepsilon$$

holds for any finite family $]\sigma_i, \tau_i[$ of non overlapping subintervals of K . This is equivalent to A being of finite variation in K , with the numerical function v absolutely continuous. In this case v is almost everywhere differentiable in K ; its derivative, denoted by \dot{v} , is a nonnegative element of $L^1(K)$ which, kinematically speaking, may be called *the speed function of the moving set A* .

The multifunction A is said to be *Lipschitzian over I* if there exists $\lambda \geq 0$ such that, for every σ and τ in I ,

$$h(A(\sigma), A(\tau)) \leq \lambda |\sigma - \tau|.$$

This holds if and only if A is absolutely continuous on every compact subinterval of I , with $\dot{v} \leq \lambda$ almost everywhere.

Having recalled this we now turn to the main purpose of the present paper: Suppose that the metric space E is actually a *normed real linear space*, that K is a compact interval of \mathbb{R} and that $t \mapsto A(t)$ and $t \mapsto B(t)$ are two multifunctions from K into E with *convex* values. If these two multifunctions are of continuous finite variation, respectively are absolutely continuous, respectively are Lipschitzian, does the same hold for the multifunction $t \mapsto A(t) \cap B(t)$? It will be shown that the answer is yes under the additional assumptions that $A(t)$ has, for every t , a non empty intersection with the interior of $B(t)$ and that $\text{diam } A(t) \cap B(t) < +\infty$. In this connection, for x fixed in E , the numerical function $t \mapsto d(x, A(t) \cap B(t))$ is also studied.

The proofs rest on various inequalities which may be of more general interest. For instance they could be used in studying the *retraction* of the multifunction $t \mapsto A(t) \cap B(t)$, a concept similar to that of variation, in the definition of which h is replaced by e (cf. [13]).

All this was motivated by the author's theory of the evolution of *elastoplastic* mechanical systems [7], [9], [12].

2. Summary.

Throughout the paper, the distance function d on E is supposed finite-valued; but, as E is not necessarily bounded, the diameters of subsets or the expressions e and h associated with two subsets may take the value $+\infty$.

The suprema or infima of expressions involving d , e or h , will always be understood in the sense of the ordering of $\bar{\mathbb{R}}_+ = [0, +\infty]$: In this ordered set $\sup \emptyset = 0$ and $\inf \emptyset = +\infty$.

Sections 3 to 5 establish some technical inequalities, mainly concerning the case where E is a real normed linear space. In particular, $e(A, E \setminus B)$ is studied; clearly A meets the interior of B if and only if this is > 0 .

The elementary facts about convex functions and their conjugates, a theory initiated by W. Fenchel [2], used in these sections, may today be considered as classical; the reader could refer to [5], [6] or [14]. (The latter treatise, though restricted to finite dimensional spaces, supplies much of the fundamental information.)

Sections 6 and 7 introduce what can be called the two *metric semi-continuities* of multifunctions from an arbitrary topological space T into a metric space E (here actually a normed space), i.e. continuity-like properties defined by means of the non-symmetric écart e . Other situations involving these concepts can be found in [1], [13], [15]. Considering two multifunctions $t \mapsto A_t$ and $t \mapsto B_t$ from T into the normed space E , with *convex* values and possessing such a semi-continuity at the point $s \in T$, it is proved that the assumption $A_s \cap \text{int} B_s \neq \emptyset$, with $\text{diam} A_s \cap B_s < \infty$, ensures the same semi-continuity for $t \mapsto A_t \cap B_t$. This yields also semi-continuity properties, in the classical sense, for the numerical functions $t \mapsto \text{diam} A_t \cap B_t$ and $t \mapsto e(A_t, E \setminus B_t)$.

Section 8 makes use of all what precedes to answer the question formulated in the introduction. The final section 9 shows that, in what concerns the study of the numerical function $t \mapsto d(x, A_t \cap B_t)$, with x fixed in E , the boundedness assumption on $A_t \cap B_t$ may be dropped.

All this reproduces, with some improvements, most of the material formerly presented in two multigraph seminar reports [10]; short sum-

maries appeared in [11] and [12] (to which a correction must be brought: In the formulation of what corresponds to paragraph 4 below, the assumption $\text{int}B_2 \neq \emptyset$ had been forgotten).

3. Diameters.

Let A be a subset of a metric space (E, d) ; the *diameter* of A , i.e.

$$\text{diam } A = \sup \{d(x, y) : x \in A, y \in A\}$$

takes its value in $\bar{\mathbb{R}}_+ = [0, +\infty]$. As already stipulated such a supremum is understood in the sense of the ordering of $\bar{\mathbb{R}}_+$, so that $\text{diam } \emptyset = 0$.

If A' denotes another subset of E , one has, with the definition (1) of the *excess* e ,

$$(2) \quad \text{diam } A' \leq \text{diam } A + 2e(A', A).$$

In fact, this inequality is trivial when A or A' is empty. Otherwise let x' and y' be elements of A' and let $\varepsilon > 0$; there exist x and y in A such that

$$d(x', x) \leq d(x', A) + \varepsilon, \quad d(y', y) \leq d(y', A) + \varepsilon.$$

Using the distance inequality for the chain of points x', x, y, y' , then taking suprema for x' and y' ranging over A' one finally obtains

$$\text{diam } A' \leq \text{diam } A + 2e(A', A) + 2\varepsilon.$$

Since ε is arbitrary this proves (2).

In a similar way the following inequality can be established:

For every $a \in A$ and $b \in E$

$$(3) \quad d(a, b) \leq \text{diam } A + d(b, A).$$

4. Balls contained in convex sets.

If (E, d) denotes an arbitrary metric space and B a subset of E , the open ball with center a and radius ϱ is contained in B if and only if $\varrho \leq d(a, E \setminus B)$. One concludes that, if A denotes another subset of E , the following equivalence holds

$$(4) \quad A \cap \text{int}B \neq \emptyset \Leftrightarrow e(A, E \setminus B) > 0.$$

From now on let us suppose that E is a real normed linear space. Let F be its dual; in both spaces the norm will be denoted by $\|\cdot\|$.

If B_1 and B_2 are two nonempty *convex*, not necessarily closed, subsets of E , their respective *support functions* γ_1 and γ_2 are sublinear functions from F into $]-\infty, +\infty]$, taking the value zero at the origin. An expression

of $e(B_1, B_2)$ in terms of γ_1 and γ_2 is easily found (cf. [13]) and, in particular, denoting by σ the *unit sphere* of F , one obtains

$$(5) \quad \forall y \in \sigma: \gamma_1(y) \leq \gamma_2(y) + e(B_1, B_2) .$$

LEMMA. *Let B_1 and B_2 denote two convex subsets of the normed space E , with $\text{int} B_2 \neq \emptyset$ and let $a \in E$; then*

$$(6) \quad d(a, E \setminus B_1) \leq d(a, E \setminus B_2) + e(B_1, B_2) .$$

PROOF. In order to simplify the calculations, assume that $a = 0$. Suppose B_1 nonempty, since otherwise the inequality is trivial.

Let ρ be a real number such that

$$(7) \quad 0 < \rho \leq d(0, E \setminus B_1);$$

this means that the open ball with center 0 and radius ρ is contained in B_1 . As the support function of the ball is $y \mapsto \rho\|y\|$, this inclusion implies

$$\forall y \in F: \rho\|y\| \leq \gamma_1(y)$$

or equivalently, as γ_1 is positively homogeneous,

$$(8) \quad \forall y \in \sigma: \rho \leq \gamma_1(y) .$$

Let us make use now of inequality (5) and suppose $e(B_1, B_2) < +\infty$ (otherwise the lemma is trivial); then (8) implies

$$\forall y \in \sigma: \rho - e(B_1, B_2) \leq \gamma_2(y) .$$

Again, let us interpret this as an inequality between two support functions, therefore equivalent to the relation of inclusion between the corresponding *closed* convex sets. Suppose first $\rho - e(B_1, B_2) > 0$; the preceding inequality means that the closed ball centered at the origin with this radius is contained in $\text{cl} B_2$. By an elementary property of convex sets in topological linear spaces, the hypothesis $\text{int} B_2 \neq \emptyset$ ensures $\text{cl} B_2 = \text{clint} B_2$; thus the corresponding open ball is contained in B_2 , i.e.

$$(9) \quad \rho - e(B_1, B_2) \leq d(0, E \setminus B_2) .$$

This, on the other hand, is trivial in the case $\rho - e(B_1, B_2) \leq 0$. The fact that (7) implies (9) proves the lemma.

PROPOSITION. *Let B_1 and B_2 denote two convex subsets of the normed space E , with $\text{int} B_2 \neq \emptyset$, and let A_1 and A_2 denote arbitrary subsets of E . Then*

$$(10) \quad e(A_1, E \setminus B_1) \leq e(A_2, E \setminus B_2) + e(A_1, A_2) + e(B_1, B_2) .$$

PROOF. Taking the suprema of both members of (6) for a ranging over A_1 yields

$$e(A_1, E \setminus B_1) \leq e(A_1, E \setminus B_2) + e(B_1, B_2).$$

Now, as e satisfies the triangle inequality, one has

$$e(A_1, E \setminus B_2) \leq e(A_1, A_2) + e(A_2, E \setminus B_2).$$

5. Main inequality.

LEMMA. Let X be a real topological linear space and Y its topological dual. Let f and g denote two convex functions defined on X , with values in $]-\infty, +\infty]$; suppose there exists a point $a \in X$ at which both functions take finite values and that one of them is continuous at this point. Then, denoting by f^* their conjugate functions (i.e. the polar functions in the terminology of [6]) defined on Y , the function $(f+g)^*$ equals the infimal convolute of f^* and g^* .

If A and B are two convex subsets of X such that $A \cap \text{int} B \neq \emptyset$, the support function of $A \cap B$ equals the infimal convolute of the support functions of A and B .

PROOF. By the definition of polar functions, the continuous affine function $x \mapsto \langle x, y \rangle - r$, with $y \in Y$ and $r \in \mathbb{R}$ is a minorant, for instance of f if and only if $r \geq f^*(y)$. By the definition of the infimal convolute $f^* \nabla g^*$

$$(f^* \nabla g^*)(y) = \inf \{f^*(u) + g^*(v) : u + v = y\}$$

the assumption

$$r > (f^* \nabla g^*)(y)$$

implies the existence of s and t in \mathbb{R} , and of u and v in Y , such that

$$r = s + t, \quad y = u + v, \quad s > f^*(u), \quad t > g^*(v).$$

Then the affine functions $\langle \cdot, u \rangle - s$ and $\langle \cdot, v \rangle - t$ are minorants of f and g , respectively; consequently $\langle \cdot, y \rangle - r$ is a minorant of $f+g$, thus

$$(11) \quad r \geq (f+g)^*(y).$$

This proves $f^* \nabla g^* \geq (f+g)^*$.

To establish the reverse inequality, we are going to prove that (11) implies

$$r \geq (f^* \nabla g^*)(y).$$

This will be done by showing that every continuous affine minorant m

of $f+g$ equals the sum of continuous affine minorants of f and g . In fact, for such an m one has

$$\forall x \in X: f(x) \geq m(x) - g(x).$$

Therefore, in the topological linear space $X \times \mathbb{R}$ the two convex sets

$$\{(x, r) \in X \times \mathbb{R} : f(x) < r\}$$

and

$$\{(x, r) \in X \times \mathbb{R} : r < m(x) - g(x)\}$$

have an empty intersection. As f possesses a point of continuity the first of these sets possesses a nonempty interior; by a standard separation theorem it follows that there exists a closed hyperplane separating these two subsets of $X \times \mathbb{R}$. The fact that both sets meet the line $\{(x, r) : x = a\}$, and again the continuity of f at the point a , ensures that this hyperplane is “nonvertical”, i.e. it is the graph of an affine continuous functions $n: X \rightarrow \mathbb{R}$ such that

$$m - g \leq n \leq f.$$

Thus n and $m - n$ are continuous affine minorants of f and g , respectively; their sum equals m .

The last part of the lemma is a special case of what precedes: Take as f and g the respective indicator functions of A and B (i.e. the functions taking the value zero on the sets and $+\infty$ outside).

PROPOSITION. *Let A and B denote two convex subsets of the normed space E . If there exists an open ball with radius $\rho > 0$ contained in B , whose center a belongs to A , one has*

$$(12) \quad \forall x \in E: d(x, A \cap B) \leq (1 + \rho^{-1}\|x - a\|)(d(x, A) + d(x, B)).$$

PROOF. Let φ and γ denote the support functions of A and B , defined on the dual F of E . Let $x \in E$; standard arguments from the theory of conjugate convex functions yield

$$d(x, A) = \sup \{ \langle x, u \rangle - \varphi(u) : u \in F, \|u\| \leq 1 \}$$

and a similar expression for $d(x, B)$, hence

$$d(x, A) + d(x, B) = \sup \{ \langle x, u + v \rangle - \varphi(u) - \gamma(v) : \|u\| \leq 1, \|v\| \leq 1 \}.$$

As the function θ defined on $F \times F$ by

$$\theta(u, v) = \langle x, u + v \rangle - \varphi(u) - \gamma(v)$$

is positively homogeneous, one has equivalently, for every $k > 0$,

$$(13) \quad k(d(x, A) + d(x, B)) = \sup \{ \theta(u, v) : \|u\| \leq k, \|v\| \leq k \}.$$

Let us make calculation easier by supposing a translation performed in E such that $a=0$; in that case the hypothesis $a \in A$ implies

$$(14) \quad \forall u \in F: \varphi(u) \geq 0$$

and the hypothesis that the open ball with center $a=0$ and radius ϱ is contained in B implies

$$(15) \quad \forall v \in F: \gamma(v) \geq \varrho \|v\|.$$

By the lemma, the support function of $A \cap B$ is the inf-convolute $\varphi \nabla \gamma$, thus

$$\begin{aligned} d(x, A \cap B) &= \sup \{ \langle x, w \rangle - (\varphi \nabla \gamma)(w) : \|w\| \leq 1 \} \\ &= \sup \{ \sup \{ \langle x, w \rangle - \varphi(u) - \gamma(v) : u + v = w \} : \|w\| \leq 1 \} \\ &= \sup \{ \theta(u, v) : \|u + v\| \leq 1 \}. \end{aligned}$$

Now, in view of (14) and (15) one has the implication

$$\|u + v\| \leq 1 \Rightarrow \theta(u, v) \leq \|x\| - \varrho \|v\|.$$

Therefore, when $\|v\| > \|x\|/\varrho$, the value of $\theta(u, v)$ is less than $\theta(0, 0) = 0$; hence,

$$d(x, A \cap B) = \sup \{ \theta(u, v) : \|u + v\| \leq 1, \|v\| \leq \|x\|/\varrho \}.$$

But

$$\|u + v\| \leq 1, \|v\| \leq \|x\|/\varrho \Rightarrow \|u\| \leq 1 + \|x\|/\varrho.$$

After putting $k = 1 + \|x\|/\varrho$ in (13), the comparison of the sets over which the suprema of $\theta(u, v)$ are taken yields

$$d(x, A \cap B) \leq (1 + \|x\|/\varrho)(d(x, A) + d(x, B));$$

since $a=0$, this is (12).

COROLLARY. *Let A and B denote two convex subsets of the normed space E ; let α and ϱ be real numbers such that*

$$(16) \quad 0 < \alpha < \varrho < e(A, E \setminus B).$$

Then, for every $x \in E$ such that

$$(17) \quad d(x, A) + d(x, B) \leq \alpha,$$

one has

$$(18) \quad d(x, A \cap B) \leq \frac{\varrho + \text{diam } A \cap B}{\varrho - \alpha} (d(x, A) + d(x, B)).$$

PROOF. In view of (16) there exists $a \in A$ satisfying with ϱ the hypothesis of the proposition. On the other hand, similarly to inequality (3),

$$\|x - a\| \leq \text{diam } A \cap B + d(x, A \cap B).$$

Therefore inequality (12) ensures

$$d(x, A \cap B) \leq (1 + \varrho^{-1}(\text{diam } A \cap B + d(x, A \cap B)))(d(x, A) + d(x, B)).$$

Denoting by s the left member of (17) and by r the left member of (18), this is equivalent to

$$\varrho r \leq (\varrho + \text{diam } A \cap B + r)s.$$

If x satisfies (17) one has $\varrho - s \geq \varrho - \alpha > 0$, hence

$$r \leq \frac{\varrho + \text{diam } A \cap B}{\varrho - s} s \leq \frac{\varrho + \text{diam } A \cap B}{\varrho - \alpha} s,$$

which is (18).

REMARK. When E is a Hilbert or pre-Hilbert space some more precise inequalities may be obtained by using trigonometrical arguments (see [10]); they may be of use, for instance, in the study of multifunctions with discontinuous finite variation.

6. First semi-continuity.

PROPOSITION. Let T denote a topological space and let $t \mapsto A_t$ and $t \mapsto B_t$ be two multifunctions from T into the normed space E , with convex values. Let $s \in T$ be such that

$$\begin{aligned} \text{diam } A_s \cap B_s &< +\infty, \\ (19) \quad A_s \cap \text{int } B_s &\neq \emptyset, \end{aligned}$$

$$(20) \quad \lim_{t \rightarrow s} e(A_t, A_s) = 0,$$

$$(21) \quad \lim_{t \rightarrow s} e(B_t, B_s) = 0.$$

Then

$$(22) \quad \lim_{t \rightarrow s} e(A_t \cap B_t, A_s \cap B_s) = 0$$

and the two functions $t \mapsto \text{diam } A_t \cap B_t$ and $t \mapsto e(A_t, E \setminus B_t)$ are upper-semi-continuous at the point s .

PROOF. In view of assumption (19) there exist two real numbers α and ϱ such that $0 < \alpha < \varrho < e(A_s, E \setminus B_s)$. By assumptions (20) and (21) there exists a neighborhood V of s in T such that, for every t in V one

has $e(A_t, A_s) \leq \alpha/2$ and $e(B_t, B_s) \leq \alpha/2$. Hence, if $t \in V$ and $x \in A_t \cap B_t$ one has $d(x, A_s) + d(x, B_s) \leq \alpha$. Therefore, using (18) we get

$$d(x, A_s \cap B_s) \leq \frac{\varrho + \text{diam } A_s \cap B_s}{\varrho - \alpha} (e(A_t, A_s) + e(B_t, B_s)) .$$

Taking the supremum for x ranging over $A_t \cap B_t$, one obtains an upper bound for $e(A_t \cap B_t, A_s \cap B_s)$ which shows that (22) follows from (20) and (21).

The upper semi-continuity of the function $t \mapsto \text{diam } A_t \cap B_t$ is a consequence of the inequality (2), namely

$$\text{diam } A_t \cap B_t \leq \text{diam } A_s \cap B_s + 2e(A_t \cap B_t, A_s \cap B_s) .$$

The upper semi-continuity of the function $t \mapsto e(A_t, E \setminus B_t)$ (trivial in the case where $e(A_s, E \setminus B_s) = +\infty$) follows from the proposition of section 4, which implies

$$e(A_t, E \setminus B_t) \leq e(A_s, E \setminus B_s) + e(A_t, A_s) + e(B_t, B_s)$$

REMARK. Concerning, for instance the multifunction $t \mapsto A_t$, observe that assumption (20) holds in particular when this multifunction is *upper semi-continuous* at the point s in the following classical sense: For every open set Ω containing A_s , there exists a neighborhood V of s in T such that $t \in V$ implies $A_t \subset \Omega$. Furthermore, under the additional assumption that A_s is compact this upper semi-continuity is equivalent to (20).

7. Second semi-continuity.

PROPOSITION. *In the same framework as in the preceding section, let the hypotheses (20) and (21) be replaced by*

$$(23) \quad \lim_{t \rightarrow s} e(A_s, A_t) = 0 ,$$

$$(24) \quad \lim_{t \rightarrow s} e(B_s, B_t) = 0 .$$

Then

$$(25) \quad \lim_{t \rightarrow s} e(A_s \cap B_s, A_t \cap B_t) = 0$$

and the two functions $t \mapsto \text{diam } A_t \cap B_t$ and $t \mapsto e(A_t, E \setminus B_t)$ are lower semi-continuous at the point s .

PROOF. As the assumption $A_s \cap \text{int } B_s \neq \emptyset$ still holds, there exist $r > 0$ and $a \in A_s$ such that $d(a, E \setminus B_s) > r$. The assumptions (23) and (24) yield a neighborhood V of s in T , such that, for every $t \in V$,

$$e(A_s, A_t) < \frac{1}{4}r, \quad e(B_s, B_t) < \frac{1}{4}r .$$

Thus, for each $t \in V$, one has $d(a, A_t) < \frac{1}{4}r$ and this ensures the existence of a point a_t in A_t such that $\|a - a_t\| < \frac{1}{4}r$. On the other hand, inequality (6), rewritten as

$$d(a, E \setminus B_s) \leq d(a, E \setminus B_t) + e(B_s, B_t)$$

implies that, if $t \in V$,

$$d(a, E \setminus B_t) > r - e(B_s, B_t) > \frac{3}{4}r.$$

On the other hand

$$d(a, E \setminus B_t) \leq \|a - a_t\| + d(a_t, E \setminus B_t)$$

thus $d(a_t, E \setminus B_t) \geq \frac{1}{2}r$. Therefore, inequality (12) yields, for every $t \in V$,

$$\forall x \in E: d(x, A_t \cap B_t) \leq (1 + 2r^{-1}\|x - a_t\|)(d(x, A_t) + d(x, B_t)).$$

If $x \in A_s \cap B_s$ one has

$$\|x - a_t\| \leq \|x - a\| + \|a - a_t\| \leq \text{diam } A_s \cap B_s + \frac{1}{4}r$$

hence finally, for every $t \in V$,

$$e(A_s \cap B_s, A_t \cap B_t) \leq (\frac{3}{2} + 2r^{-1} \text{diam } A_s \cap B_s)(e(A_s, A_t) + e(B_s, B_t)).$$

This shows that (25) follows from (23) and (24).

The lower semi-continuity of $t \mapsto \text{diam } A_t \cap B_t$ results from inequality (2) rewritten as

$$\text{diam } A_s \cap B_s \leq \text{diam } A_t \cap B_t + 2e(A_s \cap B_s, A_t \cap B_t).$$

The lower semi-continuity of $t \mapsto e(A_t, E \setminus B_t)$ results from inequality (10) rewritten as

$$e(A_s, E \setminus B_s) \leq e(A_t, E \setminus B_t) + e(A_s, A_t) + e(B_s, B_t).$$

REMARK. Concerning, for instance, the multifunction $t \mapsto A_t$, observe that assumption (23) implies that this multifunction is *lower semi-continuous* in the following classical sense: For every open set Ω meeting A_s , there exists a neighborhood V of s in T such that $t \in V$ implies $A_t \cap \Omega \neq \emptyset$.

8. Variation over a compact interval.

PROPOSITION. *Let K be a compact interval of R ; let $t \mapsto A_t$ and $t \mapsto B_t$ be two multifunctions from K into the normed space E , with convex values; it is supposed that, for every $t \in K$,*

$$\text{diam } A_t \cap B_t < +\infty, \quad A_t \cap \text{int } B_t \neq \emptyset.$$

Then, if both multifunctions are of continuous finite variation in K (resp. are absolutely continuous, resp. are Lipschitzian), so is also the multifunction $t \mapsto A_t \cap B_t$.

PROOF. The assumption involves that both multifunctions are continuous in the sense of Hausdorff distance h which majorizes the écart e ; therefore the propositions of sections 6 and 7 can be applied. This shows that the finite-valued function $t \mapsto \text{diam } A_t \cap B_t$ is upper semi-continuous at every point of the compact interval K , thus majorized by some finite constant D . Similarly the function $t \mapsto e(A_t, E \setminus B_t)$, with strictly positive values, is lower semi-continuous on K , thus minorized by some constant $\varrho > 0$; let us choose α in $]0, \varrho[$.

The indefinite variations of the two multifunctions are continuous finite numerical functions on K , hence uniformly continuous. Consequently there exists $p > 0$ such that for any two points σ and τ of K , the condition $|\sigma - \tau| \leq p$ ensures that $h(A_\sigma, A_\tau)$ and $h(B_\sigma, B_\tau)$ are majorized by $\frac{1}{2}\alpha < \frac{1}{2}\varrho$. When these majorations hold, by putting $k = \varrho + D/\varrho - \alpha$, inequality (18) yields

$$\forall x \in A_\sigma \cap B_\sigma: d(x, A_\tau \cap B_\tau) \leq k(d(x, A_\tau) + d(x, B_\tau)).$$

After taking the suprema for x ranging over $A_\sigma \cap B_\sigma$, one obtains an inequality concerning the écart e ; the same holds when σ and τ are exchanged; hence finally

$$h(A_\sigma \cap B_\sigma, A_\tau \cap B_\tau) \leq k(h(A_\sigma, A_\tau) + h(B_\sigma, B_\tau)).$$

In view of this, the definition of the variation of multifunctions over a compact subinterval $[s, t]$ of K will be applied, under the precaution of considering only sufficiently fine subdivisions of this subinterval, in order that the distance between successive points be less than p . This yields

$$\text{var}(A \cap B; s, t) \leq k(\text{var}(A; s, t) + \text{var}(B; s, t))$$

which establishes the proposition.

A useful special case consists in taking as B_t a fixed open ball β :

COROLLARY. *Let K be a compact interval of R ; let $t \mapsto A_t$ be a multifunction from K into the normed space E , with convex values and let β denote a fixed open ball such that, for every t in K , one has $A_t \cap \beta \neq \emptyset$. Then, if $t \mapsto A_t$ is of finite continuous variation (resp. is absolutely continuous, resp. is Lipschitzian) so is also the multifunction $t \mapsto A_t \cap \beta$.*

REMARK. For an interval I possibly non-compact, the above proposition may be used from a local standpoint: If the multifunctions $t \mapsto A_t$ and $t \mapsto B_t$ are of finite continuous variation in I and if $s \in I$ is such that

$$\text{diam } A_s \cap B_s < +\infty \quad \text{and} \quad A_s \cap \text{int } B_s \neq \emptyset,$$

sections 6 and 7 yield a compact interval, neighborhood of s in I , to which the proposition can be applied.

9. Distance function.

Let x be a fixed point of E . By elementary inequalities, the semi-continuity properties of the multifunction $t \mapsto A_t \cap B_t$ possibly established by applying the foregoing propositions, as well as the properties concerning the variation of this multifunction, imply similar properties for the numerical function $t \mapsto d(x, A_t \cap B_t)$. The purpose of the present section is to show that the conclusions concerning this numerical function may actually be obtained in a slightly more general framework, free from the boundedness assumption precedingly made about the intersection $A_t \cap B_t$.

PROPOSITION. *Let T denote a topological space and let $t \mapsto A_t$ and $t \mapsto B_t$ be two multifunctions from T into the normed space E , with convex values. Let $s \in T$ such that $A_s \cap \text{int} B_s \neq \emptyset$ and*

$$\begin{aligned} \lim_{t \rightarrow s} e(A_t, A_s) &= 0 && \text{(respectively } \lim_{t \rightarrow s} e(A_s, A_t) = 0) \\ \lim_{t \rightarrow s} e(B_t, B_s) &= 0 && \text{(respectively } \lim_{t \rightarrow s} e(B_s, B_t) = 0). \end{aligned}$$

Then, for every $x \in E$, the function $t \mapsto d(x, A_t \cap B_t)$ is lower semi-continuous (resp. upper semi-continuous) at the point s .

PROOF OF THE LOWER SEMI-CONTINUITY. As $A_s \cap B_s$ is nonempty, $d(x, A_s \cap B_s)$ is finite. We have to prove that, for every $\varepsilon > 0$, there exists a neighborhood V of s in T ensuring the implication

$$(26) \quad t \in V \Rightarrow d(x, A_t \cap B_t) \geq d(x, A_s \cap B_s) - \varepsilon.$$

Let us choose an open ball β , with center x and large enough to meet $A_s \cap \text{int} B_s$. To establish (26) it suffices to prove the similar implication where the considered sets are replaced by their intersections with β . In fact, if $A_t \cap B_t \cap \beta \neq \emptyset$, one has

$$d(x, A_t \cap B_t \cap \beta) = d(x, A_t \cap B_t)$$

and if $A_t \cap B_t \cap \beta = \emptyset$, the distance $d(x, A_t \cap B_t)$ is greater than or equal to the radius of β which itself is greater than $d(x, A_s \cap B_s)$. Put $B'_t = B_t \cap \beta$ and $B'_s = B_s \cap \beta$; the triangle inequality concerning the écart e , applied to the three sets $\{x\}$, $A_t \cap B'_t$ and $A_s \cap B'_s$ yields

$$(27) \quad d(x, A_s \cap B'_s) \leq d(x, A_t \cap B'_t) + e(A_t \cap B'_t, A_s \cap B'_s).$$

The results of section 6 hold for the pair of multifunctions $t \mapsto B_t$ and $t \mapsto \beta$, thus

$$\lim_{t \rightarrow s} e(B_t', B_s') = 0.$$

As these results hold again for the pair of multifunctions $t \mapsto A_t$ and $t \mapsto B_t'$, one has

$$\lim_{t \rightarrow s} e(A_t \cap B_t', A_s \cap B_s') = 0.$$

In view of (27) this proves the expected implication.

PROOF OF THE UPPER SEMI-CONTINUITY. The arguments of section 6, at their beginning, do not make use of the boundedness of $A_s \cap B_s$: In the present case they still yield a real number $r > 0$, a point $a \in A_s$, a neighborhood V of s in T such that for every $t \in V$ there exists $a_t \in A_t$ with $\|a - a_t\| < \frac{1}{2}r$ and $d(a_t, E \setminus B_t) \geq \frac{1}{2}r$. The point a_t belongs to $A_t \cap B_t$, thus

$$d(x, A_t \cap B_t) \leq \|x - a_t\| \leq \|x - a\| + \|a - a_t\| < \|x - a\| + \frac{1}{4}r$$

which means that the open ball β with center x and radius $\|x - a\| + \frac{1}{4}r$ meets $A_t \cap B_t$. Consequently, for every $t \in V$,

$$(28) \quad d(x, A_t \cap B_t) = d(x, A_t \cap B_t \cap \beta).$$

The results of section 6 apply to the pair of multifunctions $t \mapsto B_t$ and $t \mapsto \beta$, thus by putting $B_t' = B_t \cap \beta$

$$\lim_{t \rightarrow s} e(B_s', B_t') = 0.$$

The same results apply also to the pair $t \mapsto A_t$ and $t \mapsto B_t'$, with the conclusion

$$\lim_{t \rightarrow s} e(A_s \cap B_s', A_t \cap B_t') = 0.$$

Then, in view of (28), the upper semi-continuity of $t \mapsto d(x, A_t \cap B_t)$ follows from the triangle inequality

$$d(x, A_t \cap B_t') \leq d(x, A_s \cap B_s') + e(A_s \cap B_s', A_t \cap B_t').$$

As an application of this proposition, let us indicate:

COROLLARY. *Let K be a compact interval of \mathbf{R} ; let $t \mapsto A_t$ and $t \mapsto B_t$ be two multifunctions from K into the normed space E , with convex values; let x be a point of E . It is supposed that for every $t \in K$ one has $A_t \cap \text{int} B_t \neq \emptyset$. Then, if both multifunctions are of finite continuous variation on K (resp. are absolutely continuous, respectively are Lipschitzian) so is also the function $t \mapsto d(x, A_t \cap B_t)$.*

PROOF. The above proposition ensures that this function, with finite values, is upper semi-continuous on K , thus strictly majorized by some constant $D > 0$. For every t in K , the open ball β with center x and radius D meets $A_t \cap B_t$, thus meets the interior of B_t . The results of section 8 apply to the pair of multifunctions $t \mapsto A_t$ and $t \mapsto B_t' = B_t \cap \beta$; therefore the multifunction $t \mapsto A_t \cap B_t'$ is of continuous finite variation (resp. is absolutely continuous, resp. is Lipschitzian). Then the conclusions concerning the function $t \mapsto d(x, A_t \cap B_t) = d(x, A_t \cap B_t')$ follow from the inequality

$$|d(x, A_\tau \cap B_\tau') - d(x, A_\sigma \cap B_\sigma')| \leq h(A_\tau \cap B_\tau', A_\sigma \cap B_\sigma')$$

for every σ and τ in K .

REFERENCES

1. T. F. Bridgland, *Extreme limits of compacta valued functions*, Trans. Amer. Math. Soc. 170 (1972), 149–163.
2. W. Fenchel, *On conjugate convex functions*, Canad. J. Math. 1 (1949), 73–77.
3. H. Hermes, *On continuous and measurable selections and the existence of solutions of generalized differential equations*, Proc. Amer. Math. Soc., 29 (1971), 535–542.
4. N. Kikuchi, and Y. Tomita, *On the absolute continuity of multifunctions and orientor fields*, Funkcial. Ekvac. 14 (1971), 161–170.
5. P. J. Laurent, *Approximation et optimisation*, Hermann, Paris, 1972.
6. J. J. Moreau, *Fonctionnelles convexes*, Séminaire sur les Equations aux Dérivées partielles, Collège de France, (multigraphié 108 p.) Paris, 1967.
7. J. J. Moreau, *Sur l'évolution d'un système élasto-visco-plastique*, C.R. Acad. Sci. Paris, Sér. A, 273 (1971), 118–121.
8. J. J. Moreau, *Sélections de multiapplications à rétraction finie*, C.R. Acad. Sci. Paris, Sér. A, 276 (1973), 265–268.
9. J. J. Moreau, *Problème d'évolution associé à un convexe mobile d'un espace hilbertien*, C.R. Acad. Sci. Paris, Sér. A, 276 (1973), 791–794.
10. J. J. Moreau, *Intersection de deux convexes mobiles*, Séminaire d'Analyse Convexe, Montpellier, 3 (1973), exposé n° 1 (multigraphié 26 p.); *Note complémentaire sur les intersections de convexes mobiles*, ibid. exposé n° 16 (multigraphié 4 p.)
11. J. J. Moreau, *Intersection de deux convexes mobiles dans un espace normé*, C.R. Acad. Sci. Paris, Sér. A, 276 (1973), 1505–1508.
12. J. J. Moreau, *On unilateral constraints, friction and plasticity*, in: *New Variational Techniques in Mathematical Physics*, p. 173–322, Centro Internazionale Matematico Estivo, II Ciclo 1973, Edizioni Cremonese, Roma, 1974.
13. J. J. Moreau, *Multiplications à rétraction finie*, Ann. Scuola Norm. Sup. Pisa. (To appear.)
14. R. T. Rockafellar, *Convex analysis*, (Princeton Mathematical Series 28), Princeton University Press, 1970.
15. B. van Cutsem, *Problems of convergence in stochastic linear programming*, in: *Techniques of Optimization* (ed. A. V. Balakrishnan), p. 445–454, Academic Press, 1972.